The genetic architecture of recombination rates is sexually-dimorphic and

polygenic in wild house sparrows (*Passer domesticus*).

John B. McAuley[1], Bertrand Servin[2], Hamish A. Burnett[3], Cathrine Brekke[1], Lucy Peters[2],

Ingerid J. Hagen[3,4], Alina K. Niskanen[3,5], Thor Harald Ringsby[3], Arild Husby[6], Henrik Jensen[3],

Susan E. Johnston[1,*

[1] Institute of Ecology and Evolution, School of Biology, University of Edinburgh, Edinburgh, EH9 3FL,

United Kingdom.

[2] GenPhySE, Université de Toulouse, INRAE, ENVT, 31326 Castanet Tolosan, France.

[3] Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and

Technology, 7491 Trondheim, Norway.

[4] Norwegian Institute for Nature Research, Høgskoleringen 9, 7034 Trondheim, Norway

[5] Ecology and Genetics Research Unit, University of Oulu, 90014 Oulu, Finland.

[6] Evolutionary Biology, Department of Ecology and Genetics, Uppsala University, 75236 Uppsala,

Sweden.

* Corresponding author: Susan.Johnston@ed.ac.uk

Abstract

Meiotic recombination through chromosomal crossing-over is a fundamental feature of sex and an

important driver of genomic diversity. It ensures proper disjunction, allows increased selection responses,

and prevents mutation accumulation; however, it is also mutagenic and can break up favourable

haplotypes. This cost/benefit dynamic is likely to vary depending on mechanistic and evolutionary

contexts, and indeed, recombination rates show huge variation in nature. Identifying the genetic

architecture of this variation is key to understanding its causes and consequences. Here, we investigate

individual recombination rate variation in wild house sparrows (*Passer domesticus*). We integrate

genomic and pedigree data to identify autosomal crossover counts (ACC) and intra-chromosomal allelic

shuffling ($\overline{r}_{intra}$) in 13,056 gametes. Females had 1.37 times higher ACC, and 1.55 times higher $\overline{r}_{intra}$

than males. ACC and $\overline{r}_{intra}$ were heritable in females and males (ACC h^2 = 0.23 and 0.11; $\overline{r}_{intra}$ h^2 =

0.12 and 0.14), but cross-sex additive genetic correlations were low (r_A = 0.29 and 0.32 for ACC and

$\overline{r}_{intra}$). Conditional bivariate analyses showed that all measures remained heritable after accounting for

genetic values in the opposite sex, indicating that sex-specific ACC and $\overline{r}_{intra}$ can evolve independently.

Genome-wide models showed that ACC and $\overline{r}_{intra}$ are polygenic and driven by many small-effect loci,

many of which are likely to act in *trans* as global recombination modifiers. Our findings show that

recombination rates of females and males can have different evolutionary potential in wild birds, providing

a compelling mechanism for the evolution of sexual dimorphism in recombination.

Introduction

Meiotic recombination via chromosomal crossing-over is an essential process in sexual reproduction and has a key role in generating diversity in eukaryotic genomes (Coop and Przeworski 2007). Crossing-over can be beneficial; it ensures proper segregation of chromosomes (Koehler et al. 1996), prevents the accumulation of deleterious alleles, and increases the speed at which populations can respond to selection (Hill and Robertson 1966; Felsenstein 1974; Sarah P. Otto and Lenormand 2002). On the other hand, recombination can increase the risk of mutations at DNA double-strand break sites (Halldorsson et al. 2019; Hinch, Donnelly, and Hinch 2023), and can break down favourable allele combinations previously built up by selection (Charlesworth and Barton 1996). Despite these trade-offs, there is extensive variation in recombination rate both within and between chromosomes, individuals, populations, sexes, and species (Myers et al. 2005; Coop and Przeworski 2007; Jessica Stapley et al. 2017; Ritz, Noor, and Singh 2017). The cost-benefit dynamic of recombination rate is likely to vary depending on mechanistic and evolutionary contexts, and if rates are heritable (i.e. there is underlying additive genetic variation), then they have the potential to respond to selection. Therefore, understanding the genetic basis of recombination rate variation - that is, the amount of additive genetic variation, and the effect sizes and distributions of gene variants that contribute to such variation - is a critical first step in determining if and how they are contributing to adaptation, and how they themselves are evolving (Ritz, Noor, and Singh 2017; Jessica Stapley et al. 2017).

To date, studies investigating the genetic basis of recombination rate variation have mostly been limited to model systems and livestock, and have demonstrated that crossover rates can be heritable and associated with particular genetic variants (Kong et al. 2014; Smukowski and Noor 2011; Petit et al. 2017; Johnsson et al. 2021; C. Brekke, Johnston, et al. 2022; Kadri et al. 2016; Johnston et al. 2016; Johnston, Huisman, and Pemberton 2018; Weng et al. 2019; Cathrine Brekke et al. 2023; Cirulli, Kliman, and Noor 2007; Cattani, Kingan, and Presgraves 2012; Chan, Jenkins, and Song 2012; Hunter et al. 2016; Samuk et al. 2020). In some systems, such as mammals, the genetic architecture can be oligogenic, with loci with meiotic functions being repeatedly implicated (e.g. *RNF212*, *RNF212B*, *MEI1*, *MSH4*); whereas in other systems, such as Atlantic salmon, variation can be polygenic (i.e., controlled by many loci of small effect. Nevertheless, the patterns observed in these studies may not be generalisable to other systems due to a number of specific factors, such as: relatively low individual variation in recombination rates (e.g. in *Drosophila spp.*); long-periods of female meiotic arrest (e.g. in humans) (MacLennan et al. 2015); dynamic variation in karyotype that can affect crossover distributions (e.g. in mice) (Capilla et al. 2014; Garagna et al. 2014); and a history of strong and sustained directional selection, which theoretically imposes indirect selection for increased recombination in small populations (e.g. in crops and livestock) (S. P. Otto and Barton 2001; Ross-Ibarra 2004) (but see (Muñoz-Fuentes et al. 2015)). Another consideration is that in many vertebrates, the positioning of recombination hotspots is determined by *PRDM9*, a gene coding for a zinc finger protein that binds to particular sequence motifs in the genome to

Figure 1: Phylogeny of birds where within-sex recombination rates have been estimated using linkage mapping or cytogenetic analysis of chiasma counts (indicated by *). The underlying data is provided in Table S1 using data compiled in (L. P. Malinovskaya, Tishakova, et al. 2020) and other sources. The phylogeny was obtained from TimeTree v5 (Kumar et al. 2022). This figure should be taken as illustrative only - some of the linkage map data is likely to have had poor marker densities in telomeric regions and have less resolution to detect double crossovers. As recombination landscape and crossover rates are sex-dependent, this could underestimate recombination in one sex, hence why we have not made a formal comparison. **References:** (Jaari, Li, and Merilä 2009; J. Stapley et al. 2008; Calderón and Pigozzi 2006; B. Hansson et al. 2010; Bengt Hansson et al. 2005; Backström et al. 2008; Kawakami et al. 2014; van Oers et al. 2014; Robledo-Ruiz et al. 2022; Peñalba et al. 2020; L. P. Malinovskaya, Tishakova, et al. 2020; Pigozzi 2001; Groenen et al. 2009; Weng et al. 2019; Torgasheva and Borodin 2017; Pigozzi and Solari 1998; Aslam et al. 2010).

initiate double strand breaks (Baudat et al. 2010; Baker et al. 2017). As double strand break repair is mutagenic (Hinch, Donnelly, and Hinch 2023), this can erode the recognised motifs and reduce the number of binding sites, leading to rapid evolution of *PRDM9* alleles and a corresponding rapid turnover of hotspots (Grey, Baudat, and de Massy 2018; Myers et al. 2010). Indeed, *PRDM9* is one of the fastest evolving genes in species where it is functional, but it has also been lost from clades such as birds, canids and amphibians, where recombination hotspots appear to be stable and enriched at functional elements (Baker et al. 2017; Singhal et al. 2015). Therefore, research in both non-model, non-*PRDM9*, and non-mammalian systems is required to better understand the genetic architecture of recombination rate more broadly, and how it is affected by other intrinsic and extrinsic variation in natural environments.

A near-universal feature of recombination is that rates and landscapes differ in their degree and magnitude between male and female gametes, a phenomenon known as "heterochiasmy" (Lenormand and Dutheil 2005; Sardell and Kirkpatrick 2020). Numerous hypotheses have been proposed to explain how it has evolved, including: reducing or increasing crossing-over in the sex under stronger selection (Trivers 1988; Burt, Bell, and Harvey 1991); gametic selection reducing crossovers in the sex with

stronger haploid selection (Nei 1969; Lenormand 2003; Lenormand and Dutheil 2005); counteracting the
effects of meiotic drive (Brandvain and Coop 2012); or that it has arisen through drift (Burt, Bell, and
Harvey 1991). Yet despite several comparative investigations of overall rate (Lenormand and Dutheil
2005; Mank 2009; Cooney, Mank, and Wright 2021), there remains little empirical support for these
hypotheses. Interestingly, some of the empirical studies above show that the genetic architecture of
recombination rate can differ between the sexes, in terms of the large effect loci underpinning them (C.
Brekke, Berg, et al. 2022; Kong et al. 2014; Johnston et al. 2016; Johnston, Huisman, and Pemberton
2018; C. Brekke, Johnston, et al. 2022), and/or their cross-sex additive genetic correlations being
significantly less than one (Cathrine Brekke et al. 2023; Johnston et al. 2016). Therefore, studies that
specifically dissect within and between-sex genetic architectures in diverse species are imperative to gain
a full understanding on the molecular mechanisms and evolutionary drivers/constraints contributing to
both recombination rate variation and heterochiasmy from the chromosomal to species levels.
113
Avian systems present a unique opportunity for further developing our understanding of the genetic basis
of recombination rate variation and heterochiasmy in the wild. Linkage mapping and cytogenetic studies
in birds have shown that despite relatively conserved karyotypes, there is substantial variation in
recombination rates, broad-scale landscapes, and heterochiasmy (Figure 1; see Table S1 for a full list of
references). In addition, birds lack *PRDM9* and have highly conserved, stable recombination hotspots that
are enriched at transcription start sites (TSS) and gene promoter regions, likely due to increased
chromatin accessibility (Singhal et al. 2015; Bascón-Cardozo et al. 2022; Kawakami et al. 2017; Pan et al.
2011). A number of wild bird populations have been subject to long-term, individual based studies of their
ecology and evolution. As genomic and pedigree data for these populations proliferates, this presents an
opportunity to investigate recombination in wild, non-model systems, and ultimately, the relationships
between recombination rates, their genetic architectures, and individual fitness components such as
reproduction and offspring viability.
126
Here, we examine sex-specific variation in individual recombination rate and its genetic architecture in a
wild meta-population of house sparrows (*Passer domesticus*). House sparrows are small passerine birds
native to Eurasia, and are now distributed across most human-habited areas of the world. They are
human commensals, often found in cities and farmland, and their ubiquity makes them a model system
for ecology, evolution and adaptation (Anderson 2006). House sparrows in the Helgeland archipelago
have relatively low dispersal from their natal island (Saatoglu et al. 2021), allowing for detailed data on
survival and reproduction to be collected since 1993 (Jensen et al. 2004; Stubberud et al. 2017; Araya-
Ajoy et al. 2021). This population has extensive genomic resources, including an annotated reference
genome (Elgvin et al. 2017), a 6.5K SNP linkage map (Hagen et al. 2020), high-density SNP arrays
(Lundregan et al. 2018) and a dense genetic pedigree (Niskanen et al. 2020). In this study, we
characterise individual recombination rates using SNP and pedigree data to determine: (a) how

recombination varies at the broad scale across chromosomes; (b) the influence of intrinsic (individual)

and extrinsic (environment) effects on recombination rates; (c) the additive genetic variance and the

genetic architecture of individual recombination rates; and (d) how these factors vary between the sexes,,

including the potential for male and female recombination rates to evolve independently.

Results

Linkage mapping.

Autosomal linkage maps were constructed using data from 1,320 full-sib families, incorporating linkage

information from 9,777 gametes transmitted by 1,805 individuals to their offspring. We mapped 56,765

autosomal SNPs to 45,022 unique centiMorgan (cM) positions on 28 autosomes, with heterogeneity in

the landscape of recombination within and between the sexes (Figure S1, Tables S2 and S3). The female

and male autosomal maps were 1994.4 cM (2.18 cM/Mb) and 1627.2 cM (1.78 cM/Mb), respectively. The

female map length was 1.23 times longer than the male map length. There was a strong correlation

between chromosome length and genetic map length in each sex (r^2 = 0.97 and 0.96 for females and

males, respectively; P < 0.001) and recombination rates were female-biased for most chromosomes

(Figure 2A). Chromosome-wide recombination rate (cM/Mb) was higher in smaller autosomes (r^2 =

0.749, P < 0. 0001, fitted as a logarithmic function), with micro-chromosomes demonstrating

recombination rates 3 to 5 times that of the genome-wide average (Figure 2B). Despite their high

recombination rates, chromosomes 21, 22, 25, 27 and 28 still had linkage map lengths much shorter than

the expectation of 50 cM, indicating that the obligate crossover was not always sampled on those

chromosomes with the current genomic dataset. Summary statistics for map lengths are provided in Table

S2, and the full sex-specific linkage maps are provided in Table S3.

Identification of meiotic crossovers.

We used the full pedigree of 12,959 genotyped individuals to phase chromosomes and identify crossover

(CO) positions in gametes transmitted from focal individuals (FIDs) to their offspring. In total, we identified

236,671 COs in 13,054 phased gametes, including 6,409 gametes transmitted from 1,354 unique female

FIDs and 6,647 gametes transmitted from 1,299 unique male FIDs. Larger chromosomes had more COs

per gamete than small chromosomes, with chromosomes 1 and 2 having up to 8 COs per gamete in

some rare cases (Figure 3). Chromosomes 10 to 20 generally had 50% non-recombinant chromosomes,

likely reflective of crossover interference over these short chromosomes leading to a single obligate CO,

which has a 50% chance of segregation into the gamete. Chromosomes 21 to 28 had more than 50% of

chromosomes with no COs, meaning that we had low power to pick up COs on these chromosomes;

these chromosomes were discarded from estimates of genome-wide recombination rates.

173
Figure 2. Variation in recombination rates between chromosomes, showing correlations between (A) the
sex-specific linkage map lengths (cM) and chromosome length (Mb); and (B) sex-specific chromosomal
recombination rate (cM/Mb) and chromosome length. Numbers are individual chromosomes, and lines
and the grey-shaded areas indicate the regression slopes and standard errors, respectively.
Chromosome 25 has been omitted from Figure 2B due to its exceptionally high recombination rate (30
cM/Mb and 26.9 cM/Mb in females and males, respectively).
180

181 **Figure 3:** Distribution of crossover (CO) counts per chromosome as the proportion of total number of
182 gametes (N = 13,054). The white dashed line is the minimum expected proportion of gametes with 0 COs
183 per chromosome due to obligate crossing-over and Mendelian segregation of COs into gametes.

185 **Figure 4.** Distributions of female and male recombination rates in gametes transmitted from FIDs to their

186 offspring for A) ACC (autosomal crossover count); and B) $\overline{r}_{intra}$, (rate of intra-chromosomal shuffling).

Individual recombination rate variation

The crossover dataset (for chromosomes 1 to 20 and 1A) was used to calculate recombination rates in

each gamete using two measures: the total autosomal crossover count (ACC), and the rate of intra-

chromosomal allelic shuffling ($\overline{r}_{intra}$), which is the probability that two randomly chosen SNP loci on the

same chromosome are uncoupled during meiosis (Veller, Kleckner, and Nowak 2019). The mean ACC

was 1.37 times higher in females than in males, with mean ACCs of 18.9 and 13.8, respectively (Figure

4). The mean rate of intra-chromosomal shuffling $\overline{r}_{intra}$ was 1.55 times higher in females (Figure 4). ACC

and $\overline{r}_{intra}$ were positively correlated (r^2 = 0.684, P < 0.0001; Figure S2A).

Heritability and genetic correlations of individual recombination rates.

The proportion of phenotypic variance in ACC and $\overline{r}_{intra}$ explained by additive genetic effects (the

narrow-sense heritability, h^2) was determined within each sex using an animal model approach fitting a

genomic relatedness matrix as a random effect. For all models of ACC and $\overline{r}_{intra}$, only fixed effects of

total phase coverage, and total phase coverage^2 were significant (Table S4), as were the additive genetic,

permanent environment (repeated measures) and residual random effects (Table 1).

ACC was heritable and evolvable in females (h^2 = 0.232, I_A = 0.016) and males (h^2 = 0.112, I_A = 0.006, P

< 0.001, Table 1). The cross-sex additive genetic correlation was positive (r_A = 0.292, se = 0.154), but

was not significantly different from zero (χ^2_1 LRT = 3.48, P = 0.062) and was significantly lower than one

(χ^2_1 LRT = 19.06, P < 0.001), indicating that the genetic basis of ACC has a significant unshared

component between males and females. This was confirmed when conditioning the within sex ACC on

the genetic value in the opposite sex, where female and male ACC remained independently heritable (h^2

= 0.126 and 0.077, respectively, P < 0.001, Table 1). The permanent environment effect was lower but

significant in both females and males (pe^2 = 0.057 and 0.077 respectively, P < 0.05, Table 1), indicating

that ACC is also partially explained by individual differences not attributed to additive genetic effects.

$\bar{r}_{intra}$ was heritable and evolvable in females and males (h^2 = 0.116 & 0.140 and I_A = 0.009 & 0.027,

respectively, P < 0.001, Table 1); when corrected for ACC by adding it as an additional fixed covariate,

heritabilities were reduced but remained significant (h^2 = 0.042 & 0.084 respectively, P < 0.001). The

cross-sex additive genetic correlation of ACC was positive (r_A = 0.319, se = 0.151) and was significantly

different from zero (χ_1^2 LRT = 4.54, P = 0.033) and one (χ_1^2 LRT = 23.21, P < 0.001), again indicating that

the genetic basis of $\bar{r}_{intra}$ has an unshared component between males and females. This was confirmed

when conditioning the within-sex $\bar{r}_{intra}$ on the genetic value in the opposite sex, where female and male

$\bar{r}_{intra}$ were independently heritable (h^2 = 0.082 and 0.077, respectively, P < 0.001, Table 1). There was a

significant permanent environment effect for $\bar{r}_{intra}$ in males (pe^2 = 0.049), but not females (Table 1).

For both ACC and $\bar{r}_{intra}$, we investigated the potential contribution of female-restricted chromosomes

(i.e. the germline-restricted chromosome (Borodin et al. 2022) and the W chromosome) to phenotypic

variation by fitting individual matriline (i.e. the path of inheritance from mothers to offspring) as an

additional random effect. Matriline did not explain any of the phenotypic variance, indicating that neither

female-restricted chromosome significantly contributed to variation in recombination rates in this

population. A full explanation of why this analysis was carried out is provided in the Discussion and

Methods sections.

Genome-wide association studies of individual recombination rates.

Genome-wide association studies were carried out using a larger genomic dataset of 65,840 SNPs, which

included Z-linked SNPs and SNPs of unknown position. These models accounted for repeated measures

within the same individuals. We did not identify any loci that were significantly associated with variation in

ACC or $\bar{r}_{intra}$ in females or males (Figure 5, Figure S3), with a power analysis indicating that we had

95% power to identify any loci explaining >3.2% of the phenotypic variance. We carried out Empirical

Bayes analyses of the GWAS summary statistics to identify loci with significant non-zero effects. For

female ACC, 4,696 SNPs (7.13%) and 1,178 SNPs (1.79%) had non-zero effects at the significance

thresholds of α = 0.05 and 0.01, respectively (Figure S4). For male ACC, 179 SNPs (0.27%) and 7 SNPs

(0.01%) had non-zero effects on ACC at the levels of α = 0.05 and 0.01, respectively (Figure S4). For

female $\bar{r}_{intra}$, 1213 SNPs (1.84%) and 143 SNPs (0.22%) had non-zero effects at the significance

thresholds of α = 0.05 and 0.01, respectively (Figure S4). For male $\bar{r}_{intra}$, 1254 SNPs (1.9%) and 151

SNPs (0.23%) had non-zero effects on ACC at the levels of α = 0.05 and 0.01, respectively (Figure S4).

Table 1: Proportions of phenotypic variance explained by additive genetic (h^2), permanent environment
effect (pe^2) and residual variance (r^2). V_P and V_P (obs) are the phenotypic variances as estimated from the
animal model and from the raw data, respectively. The mean value is from the raw data. I_A is the
evolvability. h^2_{indep} is the trait heritability independent of the genetic value of the other sex, and r_A is the
additive genetic correlation. Values in parentheses are the standard errors. Significances of h^2, pe^2 and
h^2_{indep} estimates are indicated by * (P < 0.05), ** (P < 0.001) and *** (P < 0.001). All results are from 6,409
gametes from 1,354 females and 6,647 gametes from 1,299 males on all crossovers occurring on
chromosomes 1A and 1 to 20.

Rate Measure	Sex	V_P	V_P (Obs)	Mean	V_A	h^2	pe^2	I_A	h^2_{indep}	r_A
ACC	F	23.93	24.41	18.92	5.561	0.232	0.057	0.016	0.126	0.292
		(0.56)			(0.761)	(0.029)***	(0.023)*	(2.13e^{-3})	(0.023)***	(0.154)
	M	10.76	10.56	13.76	1.202	0.112	0.077	6.35e^{-3}	0.077	
		(0.221)			(0.273)	(0.025)***	(0.023)*	(1.44e^{-3})	(0.019)***	
$\bar{r}_{intra}$	F	5.50e^{-5}	5.56e^{-5}	0.027	6.40e^{-6}	0.116	1.28e^{-5}	8.94e^{-3}	0.077	0.319
		(1.04e^{-6})			(7.24e^{-7})	(0.012)***	(2.43e^{-7})	(1.01e^{-3})	(0.01)***	(0.151)
	M	5.77e^{-5}	5.75e^{-5}	0.017	8.07e^{-6}	0.14	0.049	0.027	0.082	
		(1.18e^{-6})			(1.43e^{-6})	(0.027)***	(0.020)*	(4.82e^{-3})	(0.019)***	

Figure 5: Genome-wide association plots for autosomal crossover count (ACC) and intra-chromosomal
shuffling ($\bar{r}_{intra}$) in females and males. The dashed line is the genome-wide Bonferroni-corrected
significance level (equivalent to α = 0.05). Chromosome 0 indicates SNPs of unknown genomic position.
Association statistics have been corrected with the genomic control parameter λ. P-P plots of the null
expectations of each plot is provided in Figure S3. Sample sizes are given in Table 1. Note that visually,
P values on the Z chromosome appear to be lower than on other chromosomes. This is an artefact of
over-plotting of much higher marker densities on the other chromosomes, and association statistics are
not significantly lower on the Z chromosome.

Figure 6: Chromosome partitioning of additive genetic effects on autosomal crossover count (ACC) and
intra-chromosomal shuffling ($\bar{r}_{intra}$) in females and males. Each point indicates the proportion of
phenotypic variance explained by a chromosome-specific genomic relatedness matrix modelled as a
function of chromosome length. The top row partitions variance for recombination rates estimated on all
chromosomes (1-20 & 1A), which accounts for *cis* and *trans* effects combined. The bottom row partitions
variance for recombination rates excluding that chromosome, accounting for *trans* effects only.
Correlations are analysed with a linear regression (full results in Table S6). Full chromosome partitioning
results are provided in Table S7.

Chromosome partitioning of additive genetic variance.

To confirm the hypothesis of a polygenic architecture of recombination rates, we used a chromosome

partitioning approach estimating the contribution of each chromosome to the phenotypic variance (Yang,

Manolio, et al. 2011). We computed separate genomic relatedness matrices for (a) chromosome *i* and (b)

all chromosomes excluding *i*, and fit both as random effects in an animal model. Larger chromosomes

contributed more to the total additive genetic variance for female ACC, female $\bar{r}_{intra}$, and male $\bar{r}_{intra}$

(linear regression P < 0.01, Figure 6, Table S6, Table S7), supporting the hypothesis of a polygenic

architecture underpinning these traits. The absence of an effect for male ACC is likely due to a lack of

model convergence leading to zero estimates for two of the largest chromosomes, 2 and 3; when

removing these two chromosomes, the linear regression was significant (t = 4.120, P < 0.001, Adjusted

R^2 = 0.5).

We then modified this approach to determine if the polygenic components underpinning recombination

rate are commonly acting in *trans* (i.e. they affect the global recombination rate) rather than *cis* (i.e. they

affect the recombination rate on the chromosome on which they are situated, and/or are in linkage

disequilibrium with heritable aspects of chromatin structure that affect local recombination rates). For

each chromosome i, we calculated ACC and $\overline{r}_{intra}$ for chromosomes 1A and 1 to 20 excluding i, and

then reran the animal models and correlations as above. Similarly, larger chromosomes contributed more

to the total additive genetic variance for female ACC, female $\overline{r}_{intra}$, and male $\overline{r}_{intra}$ (linear regression P

< 0.05, Figure 6, Table S6, Table S7). These effects were weaker for female $\overline{r}_{intra}$ and male $\overline{r}_{intra}$ when

compared to the full model above, suggesting that a significant component of variation in intra-

chromosomal shuffling may still act in *cis*. However, for female ACC, this effect was highly similar in both

models, indicating that polygenic variation in female autosomal crossover count is likely to be more

commonly acting in *trans* within this population.

DISCUSSION

In this study, we have shown that female house sparrows have 1.37 times higher autosomal crossover

counts (ACC) and 1.55 times more intra-chromosomal shuffling ($\overline{r}_{intra}$) than males. ACC was moderately

heritable in both females and males (h^2 = 0.23 and 0.11, respectively), as was $\overline{r}_{intra}$ (h^2 = 0.12 and 0.14,

respectively). A genome-wide association study found no regions of the genome with a significant effect

on recombination rate, with chromosome partitioning and Empirical Bayes approaches supporting a

polygenic genetic architecture. Larger chromosomes contributed more additive genetic variation to ACC

and $\overline{r}_{intra}$ in the rest of the genome, with further analysis implying that polygenic effects in female ACC

are mostly acting in *trans*, i.e. affecting the global recombination rate. For both ACC and $\overline{r}_{intra}$, the inter-

sex additive genetic correlation was low (r_A ~0.3), and both traits remained heritable within each sex after

conditioning on the genetic values of the opposite sex, indicating that the genetic architecture can evolve

independently within each sex. Here, we discuss in more detail the mechanisms by which polygenic

architecture is likely to contribute to variation in recombination rate, how this differs between the sexes,

and how the broad scale recombination landscape compares with other studies.

Sex differences in recombination and its genetic architecture.

This study provides a compelling example of heterochiasmy (i.e. sexual dimorphism) in recombination

rate and broad-scale landscapes at the phenotypic level. More importantly, our study shows that these

sex differences are also underpinned by different genetic architectures. Male and female recombination

rates are likely to have some degree of shared genetic architecture, as shown by additive genetic

correlations of $r_A \approx 0.3$; whilst there were moderate standard errors around these estimates, this value

was significantly lower than 1 for both ACC and $\overline{r}_{intra}$. Both traits remained heritable within each sex,

even after conditioning on genetic values in the opposite sex, with values ranging from h^2 = 0.077 to 0.13.

This indicates that there is potential for evolution of recombination rates within each sex independent of

that in the other sex, which could affect the degree and potentially the direction of heterochiasmy within

the population. With the exception of domestic chickens (Weng et al. 2019), our study is the largest study

of individual variation in recombination in birds, and our study is the largest to investigate sex-specific

effects at the individual and population levels. As the number of gametes from males and females are

similar in our study, and coverage is high across the chromosomes we analysed here, we are confident

that we capture the vast majority of biologically meaningful COs within each sex. While we cannot directly

relate heterochiasmy in a single population to a particular hypothesis outlined in the introduction, our

results add to a growing body of data that there is sexual dimorphism in not only the rate of crossing over,

but also in its positioning and in its genetic architecture. More broadly, our results highlight that

recombination in female and male meiosis are distinct biological processes. Future studies investigating

the evolutionary causes and consequences of recombination should endeavour to consider how sex-

differences in meiotic processes contribute to observed variation.

Variation in crossover count and intra-chromosomal shuffling.

Much of the theory proposed to explain the advantages and disadvantages of recombination rate

variation centres around the generation and preservation of beneficial haplotypes through shuffling of

alleles at linked sites (Hill and Robertson 1966; Felsenstein 1974; Kondrashov 1988). The efficacy and

extent to which crossovers shuffle linked alleles within chromosomes is not only a function of CO count,

but also of the CO position (Veller, Kleckner, and Nowak 2019). For example, a crossover in close

proximity to a chromosome end will lead to much less allele shuffling than one in the centre of a

chromosome (Veller, Kleckner, and Nowak 2019). However, almost all previous studies investigating the

genetic architecture of recombination rates have focussed solely on autosomal crossover counts (e.g.

(Kong et al. 2014; Ma et al. 2015; Petit et al. 2017; Johnston et al. 2016); but see (Cathrine Brekke et al.

2023). In our study, we modelled both ACC and $\overline{r}_{intra}$ to consider two different (but not necessarily

independent) phenomena, that respectively represent more the variation in the mechanistic process of

CO formation (ACC) and the variation in crossover positioning and potentially the evolutionary

consequences of recombination ($\overline{r}_{intra}$). We identified a substantial correlation between ACC and $\overline{r}_{intra}$

(r^2 = 0.684), reflecting that much of the variation in $\overline{r}_{intra}$ is driven by ACC, whereas the remainder may

be due to other factors, such as individual differences in chromatin landscape, differences in the genetic

architecture, and/or random processes that affect CO positioning. Females demonstrate higher ACC and

$\overline{r}_{intra}$ than males, but this difference is stronger in terms of allele shuffling rather than the number of COs

(~1.5× vs 1.37×, respectively). Therefore, females may exhibit a stronger capacity to drive responses to

selection than males in this population, although in reality, it may be that the sex-averaged rate is more

meaningful in terms of longer term responses to selection (Burt, Bell, and Harvey 1991).

One limitation of our method to characterise COs using pedigree information is that we only use data from

gametes that resulted in an offspring. This leads to a "missing fraction" of recombination measures in the

population, meaning that our measured rates may not reflect the true rate of crossing over during meiosis;

for example, problems in meiosis and lower rates of crossing over can translate into lower fertility and

increased rates of aneuploidy in humans (Hassold and Hunt 2001; Kong et al. 2004; MacLennan et al.

2015; Handel and Schimenti 2010). A full understanding of any missing fraction would aim to characterise

variation at the pre-zygotic stage e.g. by verifying rate variation using chiasma count data (L. P.

Malinovskaya, Tishakova, et al. 2020) or gamete sequencing approaches (Dréau et al. 2019) to compare

with our pedigree estimated measures. Nevertheless, these methods also have limitations: cytogenetic in

birds requires sacrificing individuals to obtain reproductive tissues, and at present, gamete sequencing

approaches would be prohibitively expensive to generate large amounts of data (Dréau et al. 2019).

Therefore, our pedigree approach is a powerful and useful method to generate large numbers of sex-

specific recombination measures (often from pre-existing data in long-term ecological studies).

What is the nature of polygenic variation underpinning recombination rates?

In our study, we show that recombination rates are heritable and present compelling evidence that this

variation is polygenic and driven by many loci of small effect. Indeed, the estimate of $h^2 = 0.232$ for

female ACC is relatively high compared to other studies of recombination in other vertebrate species

(Kong et al. 2014; Petit et al. 2017; Johnsson et al. 2021; C. Brekke, Johnston, et al. 2022; Kadri et al.

2016; Johnston et al. 2016; Johnston, Huisman, and Pemberton 2018; Weng et al. 2019; Cathrine Brekke

et al. 2023). Our results are similar to mammal studies in that females had higher phenotypic variance

and heritability of autosomal crossover counts, but are in contrast in that these studies often identify a

conserved suite of loci (e.g. *RNF212*, *RNF212B*, *REC8*, *MSH4*, *HEI10*, among others) that explain a

moderate to large effect of heritable variation (e.g. (Kong et al. 2014; Petit et al. 2017; Kadri et al. 2016;

Johnston, Huisman, and Pemberton 2018; C. Brekke, Berg, et al. 2022)). A lack of significant GWAS in

wild populations is often attributed to reduced power to detect trait loci, due to relatively small sample

sizes compared to human and livestock studies (Johnston, Chen, and Josephs 2022; Santure and Garant

2018). Our power analysis indicated that a locus would have to contribute >3.2% of the phenotypic

variance to be detected, meaning that we would only detect moderate to large effect loci. However, we

were able to leverage chromosome partitioning and Empirical Bayes approaches to show strong evidence

that the additive genetic variance in recombination rates can be attributed to small effect loci throughout

the genome, indicating a polygenic architecture.

The question remains - how does polygenic variation contribute variation in recombination rate? This trait

ultimately is a phenotype of the genome, and polygenic effects will likely operate through two main routes.

First, SNPs may be in LD with genomic features that contribute to local regulation of recombination rate

variation in *cis*, e.g. via polymorphic hotspots and heritable aspects of chromatin accessibility that are

linked to nearby SNPs. Second, SNPs may be in LD with regions of the genome that affect crossover

formation in *trans*, e.g. through modifying the cell environment, chromatin structure, and/or the expression

and structure of meiotic proteins. It is difficult to directly test *cis* effects in isolation; this could feasibly test

if local variation is associated with local recombination rate. However, as crossovers are inherently rare at
a localised level, and with a large number of tests to be conducted, this analysis becomes challenging
and underpowered. Nevertheless, we were able to test *trans* effects at a global level by adapting the
genome partitioning approach to account for recombination rates on chromosomes excluding the focal
chromosome. This indicated that male and female $\overline{r}_{intra}$ are likely to be affected by both *cis-* and *trans-*
acting polygenic variation, whereas female ACC is largely driven by *trans*-acting polygenic variation.
Future work will investigate the correlation of non-zero effect SNPs with functional variation, as well as
individual and population fine-scale variation of recombination rate to identify common genomic features
within the population.

Our study has also shown that the female-restricted chromosomes, namely the W and germline-restricted
chromosome (GRC), are likely to make a negligible contribution to heritable variation in recombination
rates. The GRC is present in all passerines and can comprise around 10% of the genome, but is not
present in somatic cells (Warren et al. 2010; Pigozzi and Solari 1998; Borodin et al. 2022; Torgasheva et
al. 2019). In males, it is generally ejected before meiosis, whereas in females it duplicates and forms
crossovers with itself (Pei et al. 2022; L. P. Malinovskaya, Zadesenets, et al. 2020). The function and
evolution of the GRC is still poorly understood, but recent work in blue tits (*Cyanistes caeruleus*) has
shown that it is enriched for meiotic genes associated with the synaptonemal complex, although its gene
content may differ between species (Kinsella et al. 2019; Mueller et al. 2023). Whilst genetic variation on
these chromosomes could not be captured by the SNP array (as this only targets somatic genomes), the
dense sparrow pedigree allowed us to solve this question by fitting matriline as a random effect. Whilst a
negligible effect is likely true in our study population, we cannot rule out that the GRC contributes to
variation in recombination rate in other systems.

Trends and comparisons in linkage mapping and broad-scale recombination landscapes.
Our motivation for constructing a linkage map was to ensure that the SNPs were correctly positioned
relative to one another, as any wrongly positioned SNPs could lead to false calling of crossovers. Our
higher-density linkage maps showed strong concordance with a previous 6.5K map in the same
population (Hagen et al. 2020), although both male and female maps were shorter in the current study
(~90% of the previous estimates; Table S1). It should be noted that both studies use different mapping
functions and softwares (Morgan function in Lep-MAP (Rastas 2017) in the current study, Kosambi
function in Cri-MAP in the previous study (Green, Falls, and Crooks 1990)). Nevertheless, differences in
map distances between functions are negligible in high-density maps (Kivikoski et al. 2023), and Lep-
MAP is less sensitive to phasing or genotyping errors through design (Rastas 2017). It is likely that our
higher-density map has improved resolution at telomeric regions and micro-chromosomes to pick up more
COs where recombination rates tend to be higher, and more power to detect double crossovers.

However, it should also be noted that as marker densities increase, recombination frequencies can be

underestimated relative to map distance using these mapping approaches (Kivikoski et al. 2023).

Our map confirmed heterogeneity in the recombination landscape, particularly between the sexes, where

some regions showed strong divergence in rates (e.g. at ~45Mb on chromosome 1A, or 15-17.5Mb on

chromosome 10, among others; Figure S1). These sex differences indicate that sudden changes in

landscape are not necessarily indicative of rearrangements (i.e. when the other sex does not show the

same trend), but they could indicate the positions of broader landscape features where males and

females can have pronounced heterochiasmy (i.e. differences in male and female rates). Future work will

investigate how recombination rate variation, particularly between the sexes, is associated with genomic

features. Our overall sex-averaged recombination rate of ~1.81 cM/Mb was concordant with

recombination rates estimated based on cytogenetic analysis of chiasma counts and recombination

nodules in birds, which range from 1.6 cM/Mb in female common terns (*Sterna hirundo*, (Lisachov et al.

2017) to 2.9 cM/Mb in female domestic geese (*Anser anser*; (Torgasheva and Borodin 2017); see data

compiled in (L. Malinovskaya et al. 2018) and Table S1). There is less concordance with previous linkage

maps in other species, particularly with those carried out with low marker densities in the early days of

linkage mapping, where cM map lengths could be as low as 694 cM for e.g. male Siberian jays (Jaari, Li,

and Merilä 2009). It is likely that marker densities in these cases have led to underestimation of

recombination by having low sub-telomeric coverage of markers, low sample sizes, and/or markers being

too widely spaced to quantify double crossovers. Therefore, understanding how landscapes of

recombination vary relative to other avian species requires generation and standardisation of higher

density linkage maps in a wider range of systems (Kivikoski et al. 2023).

CONCLUSIONS

In this study, we revealed sex differences in the rate and the genetic architecture of recombination in wild

house sparrows. This study is a first step to investigating the selective and evolutionary importance of

recombination in wild birds, with future analyses investigating fine-scale variation in recombination rate,

and association between recombination rates and fitness effects. Our results expand our understanding

of individual variation in recombination in a non-mammal system, and our approach has the potential to

be extended to other long-term studies with genomic, pedigree, and fitness information.

Material and Methods

Study system.

All data was collected from the meta-population of house sparrows inhabiting an 18-island archipelago

covering 1600 km^2 off the Helgeland coast in Northern Norway (midpoint of 66°32'N, 12°32'E), which has

been subject to an individual-based long-term study since 1993 (Jensen et al. 2004). Birds are routinely

captured and individually marked from the beginning of May to the middle of August and for

approximately one month in the autumn using mist nets (adults and fledged juveniles), or as fledglings in

accessible nests during the breeding season. A small (25 µl) blood sample is collected from the brachial

vein for DNA from every captured bird. Individual hatch year was determined as either (a) the first year

recorded for nestlings or fledged juveniles captured in the summer and autumn, or (b) the year prior to

first year recorded for birds first captured as female adults before June 1 or as males before August 1, or

(c) a range including the year first recorded and the year prior for birds first captured as adult females

after June 1 or adult males after August 1; hatch island is also recorded alongside hatch year (Saatoglu et

al. 2021; Ranke et al. 2021).

SNP genotyping and Quality Control.

All SNPs used in our analysis were taken from two custom house sparrow Axiom SNP arrays (200K and

70K) based on the resequencing of 33 individual house sparrows (Lundregan et al. 2018). SNPs on both

arrays are distributed across 29 autosomes in the house sparrow genome (chromosome 16 was excluded

as sequences on this chromosome are difficult to assemble due to containing the highly repetitive major

histocompatibility complex). All SNPs on the 70K array are present on the 200K array. All SNP positions

are given relative to the house sparrow genome assembly Passer_domesticus-1.0 (GenBank Assembly

GCA_001700915.1; (Elgvin et al. 2017; Lundregan et al. 2018)). A total of 3,886 recruited adults were

successfully genotyped on the 200K array (Niskanen et al. 2020), and an additional 9,079 recruited adults

and non-recruited fledglings and juveniles were successfully genotyped on the 70K array. For our

analyses, we merged the two datasets using *--bmerge* in PLINK v1.90b7 (Chang et al. 2015), and

removed individuals and SNPs with call rates of < 0.1 (using *--mind* and *--geno*) and SNPs with minor

allele frequencies (MAF) of < 0.01 in founder individuals (*--maf*). The merged dataset contained 65,840

SNPs in 12,965 individuals. For all autosomal SNPs used in the estimation of recombination rates below,

Mendelian errors were identified with the *--mendel* function in PLINK, and SNPs with more than 100

Mendelian mismatches were discarded. We generated summary statistics of SNP loci using the function

summary.snp.data in GenABEL v1.8-0 (Aulchenko et al. 2007) in R v3.6.3. To ensure strong

concordance between 200K and 70K data, we removed SNPs where there was a difference between the

datasets of more than 3 standard deviations between (i) the minor allele frequency, (ii) deviation from

Hardy-Weinberg equilibrium, and/or (iii) the proportion of heterozygote individuals. After these steps, we

retained 56,767 autosomal SNPs and 617 Z-linked SNPs in 12,959 individuals. The mean and median

autosomal inter-marker distances in this final dataset were 16.1 kb and 9.6 kb, respectively, and the

mean and median Z-chromosome inter-marker distances were 111.2 kb and 77.9 kb, respectively. We

investigated the linkage disequilibrium (LD) exponential decay profile for all autosomal loci occurring

within 500 kb windows using the flag *--ld-window-kb 500* in PLINK. LD decayed to $r^2 = 0.01$ at a distance

of ~100 kb (Figure S5).

Genetic pedigree construction.

508
A metapopulation-level pedigree was constructed using a subset of 873 SNP markers in the software
Sequoia (Huisman 2017). This subset of SNPs was selected for the pedigree construction by filtering the
70K SNP set using PLINK 1.9 (Chang et al. 2015) using the *--indep* command, specifying a window size
of 1000 KB, step size of 10Kb and an r^2 threshold of 0.01, whilst filtering for MAF > 0.35 and excluding Z-
chromosome mapped SNPs. The SNP set was further refined by constructing a preliminary parentage-
only pedigree, and removing SNPs with Mendelian error rates > 0.03. After these steps, 873 SNPs
remained. For genetic sex assignment, we calculated the *Fhat2* inbreeding coefficients (F_Z) using the --
ibc command using 306 high-quality Z-chromosome SNPs. Individuals with F_Z > 0.95 were assigned as
female and those with F_Z < 0.5 assigned as male. Individuals with F_Z > 0.5 < 0.95 and were left as
"unknown" sex, as were individuals assigned as genetic females that had autosomal F_{ROH} > 0.1, due to
inbreeding biasing F_Z estimates. We used the R-package Sequoia (Huisman 2017) to construct a
pedigree including 11,073 individuals sampled from Helgeland and several other nearby populations. We
used the genetic sex for all individuals and assigned known hatch years as the year of first capture for
individuals sampled as nestlings or first captured as juveniles, and assumed sparrows first captured as
female adults before June 1 or as males before August 1 were hatched the previous year. We set a
possible hatch year range of year *t*-1 or *t* for sparrows first captured as adult females after June 1 or adult
males after August 1 in year *t*, because by this stage some juveniles are difficult to distinguish from
adults. After the initial parentage-only pedigree, a final pedigree was constructed with sib-ship clustering
and LLR calculation enabled, with a maximum of eight sib-ship iterations and a genotyping error rate of
0.002 (Niskanen *et al.* 2020).
529

Linkage map construction.

530
Autosomal linkage map construction was conducted using Lep-MAP v3 (Rastas 2017). The pedigree was
ordered into full-sib families as follows: for each unique male-female mating pairing (hereafter referred to
as the focal individuals, or FIDs, in which meiosis took place), we constructed a three-generation family
including all genotyped parents and offspring. Whilst an FID can be present in several families (i.e. as an
offspring, parent, or when mating with a different individual), this design meant that each meiosis was
only counted once. A total of 4,534 full-sib families were constructed, with 1 to 20 offspring per family. We
assumed the same marker order as the house sparrow genome above, and treated each chromosome as
a separate linkage group. The module *filtering2* was run with the parameter *datatolerance* = 0.01 to filter
based on segregation distortion, although all markers passed this step. The module
separatechromosomes2 was run for each linkage group, and markers that were not assigned to the main
group (i.e. LOD score < 5) were excluded (N = 2 SNPs removed). Finally, the module *ordermarkers2* was
run to calculate the centiMorgan (cM) positions using the Morgan mapping function for each sex
separately. Relationships between (a) chromosome length (in megabases) and linkage map length (in

cM) and (b) male and female linkage map lengths (in cM) were analysed using linear regressions in R

v4.2.2.

Calculating individual recombination rates.

Chromosome phasing and crossover estimation:

The software YAPP v0.2a0 (https://yapp.readthedocs.io/) (Servin 2021) was used to phase chromosomes

and identify crossover (CO) positions in gametes transmitted from FIDs to their offspring. This approach

uses the whole pedigree rather than the smaller sub-pedigrees above, and is more robust to missing

individuals, allowing us to characterise crossovers in more individual gametes. YAPP identified 14,769

parent-offspring pairs, representing 14,769 gametes in which COs could potentially be inferred. First, the

mendel command was run with default parameters, removing 143 pairs with higher rates of Mendelian

errors using the default parameters. Next, the *phase* command was used to infer the gametic phase of

chromosomes, by implementing a Weighted Constraints Satisfaction Problem solver in the software

ToulBar2 (Favier et al. 2010). Then, the *recomb* command was run to identify COs. For each

chromosome per meiosis, YAPP outputs the start and stop positions of the informative length of the

chromosome (i.e. the total region where phase can be inferred for a particular individual from the

pedigree, or "coverage") and the start and stop positions of each crossover interval as determined by a

Hidden Markov Model (Figure S2A); this was included to account for CO detection being potentially

reduced in individuals with lower coverage. This process was run in three iterations of the *phase* and

recomb commands to allow conservative quality control and minimise the risks of calling false COs:

Iteration 1: After the first iteration, we removed parent-offspring links with > 60 COs per gamete and/or >

9 COs on any chromosomes, and removed parents or offspring with an autosomal heterozygosity of <

0.36, as this threshold was associated in an uptick in estimated CO counts, suggesting a reduced sample

quality and/or increased phasing difficulty (N = 14,127 gametes remaining).

Iteration 2: After the second iteration, we investigated the genomic locations of double crossovers

(DCOs) in close proximity (< 3Mb between adjacent CO mid-points). Chromosome 26 had a high number

of close DCOs despite its short length (~6.9 Mb), and so all COs on this chromosome were discarded.

We then identified genomic regions enriched for close DCOs by splitting the genome into 100 kb bins and

tallying the number of CO window mid-points in each bin. Two regions of the genome showed elevated

close DCOs at chromosome ends: these were from 0-3 Mb of chromosome 4; and from 68-69.9 Mb of

chromosome 1A (Figure S8). We speculate that these regions have large structural variants (e.g.

inversions) and/or rearrangements relative to the reference genome that lead to false calling of DCOs in

these regions. In this case, we removed all COs overlapping these two regions from all further analyses.

We then removed all parent-offspring links with > 45 COs per gamete and/or > 9 COs on any

chromosome, and removed parents and offspring with a SNP call rate of < 0.98 and/or a heterozygosity
value outwith 3 standard deviations of the mean (N = 13,159 gametes remaining).

Iteration 3: After running for a third iteration, we re-investigated all DCOs in the data. We assumed that
all COs are subject to interference, and that any short DCOs were indicative of phasing and/or genotyping
errors. After visual examination of the distribution of distances between DCOs to determine an
appropriate threshold, we removed all COs that were less than 2Mb apart (from the right-hand boundary
of the first CO interval to the left-hand boundary of the second CO interval; Figure S7B). Any gametes
with more than 5 short DCOs were removed. In cases of clustered short COs (i.e. 3 or more adjacent
COs that are separated by distances of 2 Mb or less), 1 CO was called in the case of odd numbers of
COs (i.e. a phase change occurred at either side of the cluster), or 0 COs in the case of even numbers of
COs (i.e. there was no phase change on either side of the cluster).

It should be noted that pedigree-based methods to estimate COs can only identify those present in one of
the four cells resulting from meiosis. For each CO, there will be two recombinant and two non-
recombinant chromatids at that position. Therefore, our CO counts represent a sample of the crossovers
that happened in meiosis I. We assume that each CO is sampled with a 50:50 probability, but we cannot
rule out that meiosis with two or more COs on the same chromosome may be more likely to be co-
inherited on the same chromatid. Therefore, there is an expectation that at least 50% of gametes will
have at least one CO per chromosome due to obligate crossing-over, as each CO per meiosis has a 50%
chance of being observed in the gamete due to Mendelian segregation. We observed that the micro-
chromosomes 21 to 28 had a higher-than-expected number of gametes with no observed COs (>50%;
Figure 3), meaning that not all COs can be detected. Therefore, all COs occurring on these chromosomes
were discarded from downstream analyses. In total, we identified 212,711 COs in 13,056 phased
gametes, including 6,409 gametes from 1,354 unique females and 6,647 gametes from 1,299 unique
males.

Recombination rate calculation:
The CO dataset was used to calculate recombination rates in FIDs using two approaches. First, we
determined the autosomal crossover count (ACC) by summing the number of COs per gamete. Second,
we calculated the rate of intra-chromosomal allelic shuffling, $\bar{r}_{intra}$, which is the probability that two
randomly chosen SNP loci on the same chromosome are uncoupled in meiosis (Veller, Kleckner, and
Nowak 2019). This was defined as:

613
$$\bar{r}_{intra} = \sum_{k=1}^{n} 2p_k(1-p_k)L_k^2$$

where for chromosome k, p_k is the proportion of the SNPs inherited from one parent, L_k is its fraction of

the genome length (or gene count), and n is the number of autosomes. In our initial data exploration, we

also quantified $\overline{r}_{intra}$ as a function of individual genes rather than SNP loci (defined as $\overline{r}_{gene}$). However,

$\overline{r}_{intra}$ and $\overline{r}_{gene}$ were highly correlated (Pearson's correlation r = 0.948, P < 0.001; Figure S2B) and so

only the $\overline{r}_{intra}$ measure was used in downstream analyses.

Determining Heritability of Recombination Rate.

Univariate models: Variance components and the proportion of phenotypic variance in recombination

measures attributed to additive genetic effects (the narrow-sense heritability, h^2) were determined using

an "animal model" fitted by restricted maximum-likelihood in the package ASReml-R v4 (Butler et al.

2009) in R v4.2.2. A genomic relatedness matrix (GRM) based on all autosomal markers was constructed

with GCTA v1.94.1 (Yang, Lee, et al. 2011). The GRM was adjusted for sampling error using the --*grm-*

adj 0 argument, which assumes the frequency spectra of genotyped and causal loci are similar. Models

were run for ACC and $\overline{r}_{intra}$ in males and females separately. The fixed effect structure included the total

phase coverage from YAPP (the length of the genome that can be phased and therefore within which

crossovers can be detected) and the total phase coverage squared. For models of $\overline{r}_{intra}$, we fit with and

without ACC as an additional continuous fixed effect, as intra-locus shuffling is a function of the crossover

count, with both measures highly correlated (Pearson's correlation r = 0.684, P < 0.001; Figure S4A).

Random effects included the additive genetic effect (GRM) and permanent environment effect (repeated

measures in each FID). Models were also run with a pedigree-based relatedness matrix calculated using

the *ainv* function in ASReml-R, but variance estimates were highly similar to those from the GRM. Initial

models were run with age of the FID in year of gamete formation (defined as the difference between

offspring hatch year and parent hatch year) as a continuous fixed covariate, and random effects of FID

hatch year, FID natal island and offspring hatch year (to investigate cohort effects and parse apart

environment effects) and FID's mother identity (to estimate maternal effects). However, these effects

were estimated as bounded at zero and were not significant in any models, and so were discarded from

further analyses.

The heritability of each measure (h^2) was determined as the ratio of the additive genetic variance V_A to

the total phenotypic variance V_P, defined as the sum of random effect variances and the residual variance

as estimated by the animal model, using the equation $h^2 = V_A/V_P$. We also calculated the mean-

standardised additive genetic variance, defined as the evolvability (I_A) using the equation $I_A = V_A/(\bar{x}^2)$,

where $\bar{x}$ is the trait mean. This measure quantifies the expected proportional change per one unit of

selection (Hansen, Pélabon, and Houle 2011). Standard errors of these estimates were derived using the

delta method implemented in the ASReml-R function *vpredict*.

Bivariate models: Bivariate models were run to determine the additive genetic covariances and

correlations between male and female ACC and $\bar{r}_{intra}$ with total phase coverage as a continuous fixed

effect. The additive genetic correlation, r_A, was calculated without constraint using the CORGH function

(i.e. correlation with heterogenous variances) in ASReml-R v4. This function also allowed us to run

additional models where r_A was constrained to zero or 0.999 (as a correlation of one cannot be fit by the

software). Significant differences between the observed value and constrained models were tested using

likelihood ratio tests, calculated as 2 times the difference between the model log-likelihoods, distributed

as χ^2 with 1 degree of freedom.

The CORGH function reports the heritabilities of traits when unconditional on the genetic values of the

other sex. To confirm this, we also estimated how much of V_A in females was conditional on genetic

values in males (i.e. $V_{A(f|m)}$), using the following equation:

663
$$V_{A(f|m)} = V_{Af} - \left(cov_{A(fm)}\right)^2 / V_{Am}$$

where V_{Af} and V_{Am} are the additive genetic variance estimates in females and males, respectively, and

$cov_{A(fm)}$ is the additive genetic covariance of the trait between the sexes (Hansen et al. 2003). The

conditional heritability was then calculated as $h^2_{(f|m)} = V_{A(f|m)}/V_P$. The $cov_{A(fm)}$ estimate was obtained

from the same bivariate animal model as above, specifying the US function (i.e. the covariance matrix is

unstructured). This was then repeated for males, conditional on genetic values in females (i.e. $V_{A(m|f)}$).

Potential Contribution of Female-Restricted Chromosomes to Variation: House sparrows have two

female-restricted chromosomes present in the cell during meiosis I - a haploid germline-restricted

chromosome (GRC) (Warren et al. 2010; Pigozzi and Solari 1998; Borodin et al. 2022; Torgasheva et al.

2019; Pei et al. 2022; L. P. Malinovskaya, Zadesenets, et al. 2020) and the W chromosome. The function

and evolution of the GRC is poorly understood, but recent work in blue tits (*Cyanistes caeruleus*) has

shown that it is enriched for meiotic genes associated with the synaptonemal complex, although its gene

content may differ between species (Kinsella et al. 2019; Mueller et al. 2023). If there is between-

individual genetic variation for recombination rate on the GRC or W chromosomes, this may be partially

captured by the GRM (as related females will carry more genetically similar GRCs and W chromosomes),

but *not* by the SNP array variation above. To determine their potential contribution to additive genetic

variation in recombination, we used the pedigree to determine the matriline identity of each bird (i.e. the

path of inheritance from mother to offspring). Mother identity was known for 83% of birds through direct

observation or inference of unsampled parents from previous pedigree construction in Sequoia (Huisman

2017; Niskanen et al. 2020). We identified 351 unique matrilines for the 1,354 female birds with

recombination rate estimates, with 90% of birds assigned to the 150 most common matrilines. We then fit

individual matriline (representing maternal inheritance of GRC and W, as well as mitochondria) as an

additional random effect in the animal models above to partition the proportion of variance explained by

the line of maternal inheritance independent of the additive genetic effect. If significant, this implies that

variation on the GRC, W chromosome, or mitochondria may also contribute to heritable variation in

recombination rate measures.

Determining Genomic Variants Associated with Recombination Rate.

***Genome-wide association studies (GWAS)*:** GWAS of recombination measures were conducted with

all FIDs using the merged SNP dataset (N = 65,840) implemented in RepeatABEL v1.1 (Rönnegård et al.

2016) in R v3.6.3. Models were run for ACC and $\overline{r}_{intra}$ in males and females separately. The package

models the additive effect of SNPs, where genotypes (AA, AB and BB) correspond to 0, 1 and 2, and

slopes and standard errors of associations are calculated. The total phase coverage was fit as a fixed

effect, and the GRM was fit to account for inflation of test statistics due to population structure. To

account for any further inflation, we divided association statistics using the genomic control parameter λ,

which was calculated as the observed median χ^2 statistic, divided by the null expectation median χ^2

statistic (Devlin and Roeder 1999). The significance threshold was calculated using a Bonferroni

correction, with the threshold set at α = 0.05 of P = 2.765 ×10^{-7}. We performed a power analysis to

evaluate the capacity of our GWAS to detect biologically meaningful quantitative trait loci using the

method outlined in (Visscher et al. 2017) implemented in an R function provided by Kaustubh Adhikari

(https://github.com/kaustubhad/gwas-power) in R v4.2.2. When specifying a minimum sample size of N =

1299 unique males in our dataset (i.e. the most conservative threshold), we determined that we had 95%

power to identify a locus explaining 3.2% of the phenotypic variance.

Distribution of polygenic effects: We determined the distribution of allele effect sizes and estimated

false discovery rates and false sign rates to identify loci with non-zero effects on recombination rate using

the *ash* function in the R package ashR v2.2-32 (Stephens 2017). This package models the slopes and

standard errors of the additive SNP effects from the GWAS in an Empirical Bayes framework to compute

a posterior distribution of SNP effect sizes across all loci. For SNPs estimated to have non-zero effect on

the trait, the significance of a SNP effect is determined by a local false sign rate, defined as probability of

error when classifying the slope of the effect as positive or negative, with cut-off thresholds at α = 0.05

and α = 0.01. The prior distribution was specified to be unimodal and symmetric around 0 when applying

the false discovery rate estimation, i.e. effect sizes are most likely to be 0 and equally likely to be positive

or negative; this was specified using the arguments *mixcompdist = "uniform"* and *method= "fdr"*.

Chromosome partitioning of additive genetic variance: We estimated the contribution of each

chromosome to the additive genetic variation in recombination rate, to determine if larger chromosomes

(i.e. those with more genes) contribute more to the total additive genetic variance, and thus supporting

the hypothesis of a polygenic genetic architecture. For each chromosome *i*, we calculated two GRMs: one

for chromosome *i* and one for all autosomes excluding *i*. GRMs were determined using GCTA v1.94.1

with the same parameters above. We then fit both GRMs in place of the single GRM in the animal model

structure above. This allowed us to determine the proportion of variance of the global recombination rate

explained by each chromosome. We then investigated the correlation between chromosome i size and

the proportion of additive genetic variance explained by chromosome i using a linear regression in R

v4.3.1. It should be noted that chromosome partitioning analyses can be biased as a result of

heteroscedasticity and censoring (Kemppainen and Husby 2018), as the trait heritability, SNP effect

sizes, and their physical location will impact inferences of polygenic architecture (Kemppainen and Husby

2018). To quantify the effect of this, we repeated the chromosome partitioning analysis above, but

permuting the phenotype values across all individuals within the model. This was done twenty times, due

to computational constraints. We compared the permuted data linear regression with the true regression

above, identifying little impact of censoring and heteroscedasticity on our current analysis (Figure S9).

Finally, we adapted the chromosome partitioning approach to determine if genetic variants underpinning

recombination rate are more likely to commonly act in *trans* (i.e. they affect the global recombination rate)

or *cis* (i.e. they affect the recombination rate on the chromosome on which they are situated). More

simply, this investigates the contribution of each chromosome to recombination on the remaining

chromosomes. We repeated the analysis above, except for each chromosome i, we calculated the ACC

and $\overline{r}_{intra}$ response variables excluding measures from chromosome i. We then investigated the

correlation between chromosome i size and the proportion of additive genetic variance explained by

chromosome i using a linear regression in R v4.3.1, as above.

DATA STATEMENT

Data will be publicly archived on publication. All scripts are available at the following link for review:

https://github.com/susjoh/2024_Sparrow_Recomb_GWAS.

ACKNOWLEDGEMENTS

We thank students and fieldworkers for help with fieldwork, and the hospitality of inhabitants at the study

area in Helgeland who made this study possible. Sampling was conducted in accordance with permits

from the Norwegian Food Safety Authority and the Ringing Centre at Stavanger Museum, Norway. We

thank Katie Abson, Jarrod Hadfield, Anna Hewett, Mark Ravinet, Martin Stoffel, Alexander Suh, Anna

Torgasheva, and Carl Veller for helpful discussions on many aspects of the study. This work made

extensive use of the Ashworth Computing Cluster Co-operative (AC3) hosted at the Institute of Ecology

and Evolution, with assistance from Dominik Laetsch, Cei Abreu-Goodger, and Jobran Chebib. The

house sparrow field study and genomic resource development was funded by the Research Council of

Norway (RCN grant nos. 221956, 274930 and 302619), the RCN's Centres of Excellence funding scheme

(grant no. 223257) and The Royal Society (RGF/R1/180090). S.E.J. was funded by a Royal Society

University Research Fellowship (UF150448 and URF/R/211008). J.B.M. was funded by a Darwin Trust of
Edinburgh PhD studentship.
763

764 **AUTHOR CONTRIBUTIONS**
765 SEJ & HJ conceived and designed the study. HJ and THR organised and collected field data together
766 with AH, HAB and IJH. HAB, HJ, AH, IJH & AKN generated and curated the genomic dataset. HAB &
767 AKN constructed the pedigree. BS developed and adapted YAPP software for the analysis. JBM, CB, LP
768 & SEJ analysed the data. SEJ & JBM wrote the manuscript with input from all authors.

769
770

Referenced Literature

Anderson, Ted R. 2006. *Biology of the Ubiquitous House Sparrow: From Genes to Populations*. Oxford,
UK: Oxford University Press.
Araya-Ajoy, Yimen G., Alina K. Niskanen, Hannah Froy, Peter Sjolte Ranke, Thomas Kvalnes, Bernt
Rønning, Michael Le Pepke, et al. 2021. "Variation in Generation Time Reveals Density
Regulation as an Important Driver of Pace of Life in a Bird Metapopulation." *Ecology Letters* 24
(10): 2077–87.
Aslam, Muhammad L., John W. M. Bastiaansen, Richard P. M. A. Crooijmans, Addie Vereijken, Hendrik-
Jan Megens, and Martien A. M. Groenen. 2010. "A SNP Based Linkage Map of the Turkey
Genome Reveals Multiple Intrachromosomal Rearrangements between the Turkey and Chicken
Genomes." *BMC Genomics* 11 (November): 647.
Aulchenko, Yurii S., Stephan Ripke, Aaron Isaacs, and Cornelia M. van Duijn. 2007. "GenABEL: An R
Library for Genome-Wide Association Analysis." *Bioinformatics (Oxford, England)* 23 (10): 1294–
96.
Backström, Niclas, Nikoletta Karaiskou, Erica H. Leder, Lars Gustafsson, Craig R. Primmer, Anna
Qvarnström, and Hans Ellegren. 2008. "A Gene-Based Genetic Linkage Map of the Collared
Flycatcher (Ficedula Albicollis) Reveals Extensive Synteny and Gene-Order Conservation during
100 Million Years of Avian Evolution." *Genetics* 179 (3): 1479–95.
Baker, Zachary, Molly Schumer, Yuki Haba, Lisa Bashkirova, Chris Holland, Gil G. Rosenthal, and Molly
Przeworski. 2017. "Repeated Losses of PRDM9-Directed Recombination despite the
Conservation of PRDM9 across Vertebrates." *ELife* 6 (June). https://doi.org/10.7554/elife.24133.
Bascón-Cardozo, Karen, Andrea Bours, Georg Manthey, Peter Pruisscher, Gillian Durieux, Julien Dutheil,
Linda Odenthal-Hesse, and Miriam Liedvogel. 2022. "Fine-Scale Map Reveals Highly Variable
Recombination Rates Associated with Genomic Features in the European Blackcap."
https://doi.org/10.22541/au.165423614.49331155/v1.
Baudat, F., J. Buard, C. Grey, A. Fledel-Alon, C. Ober, M. Przeworski, G. Coop, and B. de Massy. 2010.
"PRDM9 Is a Major Determinant of Meiotic Recombination Hotspots in Humans and Mice."
Science (New York, N.Y.) 327 (5967): 836–40.
Borodin, Pavel, Augustin Chen, Wolfgang Forstmeier, Simone Fouché, Lyubov Malinovskaya, Yifan Pei,
Radka Reifová, et al. 2022. "Mendelian Nightmares: The Germline-Restricted Chromosome of
Songbirds." *Chromosome Research: An International Journal on the Molecular, Supramolecular
and Evolutionary Aspects of Chromosome Biology* 30 (2–3): 255–72.
Brandvain, Yaniv, and Graham Coop. 2012. "Scrambling Eggs: Meiotic Drive and the Evolution of Female
Recombination Rates." *Genetics* 190 (2): 709–23.
Brekke, C., P. Berg, A. B. Gjuvsland, and S. E. Johnston. 2022. "Recombination Rates in Pigs Differ
between Breeds, Sexes and Individuals, and Are Associated with the RNF212, SYCP2, PRDM7,
MEI1 and MSH4 Loci." *Genetics, Selection, Evolution: GSE* 54 (1): 33.
Brekke, C., S. E. Johnston, A. B. Gjuvsland, and P. Berg. 2022. "Variation and Genetic Control of
Individual Recombination Rates in Norwegian Red Dairy Cattle." *Journal of Dairy Science*,
December. https://doi.org/10.3168/jds.2022-22368.

Brekke, Cathrine, Susan E. Johnston, Tim Knutsen, and Peer Berg. 2023. "Genetic Architecture of Individual Meiotic Crossover Rate and Distribution in a Large Atlantic Salmon (Salmo Salar) Breeding Population." *BioRxiv*. https://doi.org/10.1101/2023.06.07.543993.

Burt, Austin, Graham Bell, and Paul H. Harvey. 1991. "Sex Differences in Recombination." *Journal of Evolutionary Biology* 4 (2): 259–77.

Butler, D. G., B. R. Cullis, A. R. Gilmour, and B. J. Gogel. 2009. "Mixed Models for S Language Environments: ASReml-R Reference Manual."

Calderón, P. L., and M. I. Pigozzi. 2006. "MLH1-Focus Mapping in Birds Shows Equal Recombination between Sexes and Diversity of Crossover Patterns." *Chromosome Research: An International Journal on the Molecular, Supramolecular and Evolutionary Aspects of Chromosome Biology* 14 (6): 605–12.

Capilla, Laia, Nuria Medarde, Alexandra Alemany-Schmidt, Maria Oliver-Bonet, Jacint Ventura, and Aurora Ruiz-Herrera. 2014. "Genetic Recombination Variation in Wild Robertsonian Mice: On the Role of Chromosomal Fusions and Prdm9 Allelic Background." *Proceedings. Biological Sciences* 281 (1786): 20140297.

Cattani, M. V., S. B. Kingan, and D. C. Presgraves. 2012. "Cis- and Trans-Acting Genetic Factors Contribute to Heterogeneity in the Rate of Crossing over between the Drosophila Simulans Clade Species." *Journal of Evolutionary Biology* 25 (10): 2014–22.

Chan, Andrew H., Paul A. Jenkins, and Yun S. Song. 2012. "Genome-Wide Fine-Scale Recombination Rate Variation in Drosophila Melanogaster." *PLoS Genetics* 8 (12): e1003090.

Chang, Christopher C., Carson C. Chow, Laurent Cam Tellier, Shashaank Vattikuti, Shaun M. Purcell, and James J. Lee. 2015. "Second-Generation PLINK: Rising to the Challenge of Larger and Richer Datasets." *GigaScience* 4 (1): 7.

Charlesworth, B., and N. H. Barton. 1996. "Recombination Load Associated with Selection for Increased Recombination." *Genetical Research* 67 (1): 27–41.

Cirulli, Elizabeth T., Richard M. Kliman, and Mohamed A. F. Noor. 2007. "Fine-Scale Crossover Rate Heterogeneity in Drosophila Pseudoobscura." *Journal of Molecular Evolution* 64 (1): 129–35.

Cooney, Christopher R., Judith E. Mank, and Alison E. Wright. 2021. "Constraint and Divergence in the Evolution of Male and Female Recombination Rates in Fishes." *Evolution; International Journal of Organic Evolution* 75 (11): 2857–66.

Coop, Graham, and Molly Przeworski. 2007. "An Evolutionary View of Human Recombination." *Nature Reviews Genetics*. https://doi.org/10.1038/nrg1947.

Devlin, B., and K. Roeder. 1999. "Genomic Control for Association Studies." *Biometrics* 55 (4): 997–1004.

Dréau, Andreea, Vrinda Venu, Elena Avdievich, Ludmila Gaspar, and Felicity C. Jones. 2019. "Genome-Wide Recombination Map Construction from Single Individuals Using Linked-Read Sequencing." *Nature Communications* 10 (1): 4309.

Elgvin, Tore O., Cassandra N. Trier, Ole K. Tørresen, Ingerid J. Hagen, Sigbjørn Lien, Alexander J. Nederbragt, Mark Ravinet, Henrik Jensen, and Glenn-Peter Sætre. 2017. "The Genomic Mosaicism of Hybrid Speciation." *Science Advances* 3 (6): e1602996.

Favier, Aurélie, Jean-Michel Elsen, Simon de Givry, and Andrés Legarra. 2010. "Optimal Haplotype Reconstruction in Half-Sib Families." In *ICLP-10 Workshop on Constraint Based Methods for Bioinformatics*. https://doi.org/10.29007/rpn1.

Felsenstein, J. 1974. "The Evolutionary Advantage of Recombination." *Genetics* 78 (2): 737–56.

Garagna, Silvia, Jesus Page, Raul Fernandez-Donoso, Maurizio Zuccotti, and Jeremy B. Searle. 2014. "The Robertsonian Phenomenon in the House Mouse: Mutation, Meiosis and Speciation." *Chromosoma* 123 (6): 529–44.

Green, P., K. Falls, and S. Crooks. 1990. *Documentation for CRIMAP, Version 2.4* (version 2.4). Washington University School of Medicine, St. Louis, MO.

Grey, Corinne, Frédéric Baudat, and Bernard de Massy. 2018. "PRDM9, a Driver of the Genetic Map." *PLoS Genetics* 14 (8): e1007479.

Groenen, Martien A. M., Per Wahlberg, Mario Foglio, Hans H. Cheng, Hendrik-Jan Megens, Richard P. M. A. Crooijmans, Francois Besnier, et al. 2009. "A High-Density SNP-Based Linkage Map of the Chicken Genome Reveals Sequence Features Correlated with Recombination Rate." *Genome Research* 19 (3): 510–19.

Hagen, Ingerid J., Sigbjørn Lien, Anna M. Billing, Tore O. Elgvin, Cassandra Trier, Alina K. Niskanen, Maja Tarka, Jon Slate, Glenn-Peter Saetre, and Henrik Jensen. 2020. "A Genome-Wide Linkage

Map for the House Sparrow (Passer Domesticus) Provides Insights into the Evolutionary History of the Avian Genome." *Molecular Ecology Resources* 20 (2): 544–59.

Halldorsson, Bjarni V., Gunnar Palsson, Olafur A. Stefansson, Hakon Jonsson, Marteinn T. Hardarson, Hannes P. Eggertsson, Bjarni Gunnarsson, et al. 2019. "Characterizing Mutagenic Effects of Recombination through a Sequence-Level Genetic Map." *Science (New York, N.Y.)* 363 (6425): eaau1043.

Handel, Mary Ann, and John C. Schimenti. 2010. "Genetics of Mammalian Meiosis: Regulation, Dynamics and Impact on Fertility." *Nature Reviews. Genetics* 11 (2): 124–36.

Hansen, Thomas F., Christophe Pélabon, W. Scott Armbruster, and Matthew L. Carlson. 2003. "Evolvability and Genetic Constraint in Dalechampia Blossoms: Components of Variance and Measures of Evolvability." *Journal of Evolutionary Biology* 16 (4): 754–66.

Hansen, Thomas F., Christophe Pélabon, and David Houle. 2011. "Heritability Is Not Evolvability." *Evolutionary Biology* 38 (3): 258–77.

Hansson, B., M. Ljungqvist, D. A. Dawson, J. C. Mueller, J. Olano-Marin, H. Ellegren, and J-A Nilsson. 2010. "Avian Genome Evolution: Insights from a Linkage Map of the Blue Tit (Cyanistes Caeruleus)." *Heredity* 104 (1): 67–78.

Hansson, Bengt, Mikael Akesson, Jon Slate, and Josephine M. Pemberton. 2005. "Linkage Mapping Reveals Sex-Dimorphic Map Distances in a Passerine Bird." *Proceedings. Biological Sciences* 272 (1578): 2289–98.

Hassold, Terry, and Patricia Hunt. 2001. "To Err (Meiotically) Is Human: The Genesis of Human Aneuploidy." *Nature Reviews Genetics*. https://doi.org/10.1038/35066065.

Hill, W. G., and Alan Robertson. 1966. "The Effect of Linkage on Limits to Artificial Selection." *Genetical Research*. https://doi.org/10.1017/s0016672300010156.

Hinch, Robert, Peter Donnelly, and Anjali Gupta Hinch. 2023. "Meiotic DNA Breaks Drive Multifaceted Mutagenesis in the Human Germ Line." *Science (New York, N.Y.)* 382 (6674): eadh2531.

Huisman, Jisca. 2017. "Pedigree Reconstruction from SNP Data: Parentage Assignment, Sibship Clustering and Beyond." *Molecular Ecology Resources* 17 (5): 1009–24.

Hunter, Chad M., Wen Huang, Trudy F. C. Mackay, and Nadia D. Singh. 2016. "The Genetic Architecture of Natural Variation in Recombination Rate in Drosophila Melanogaster." *PLoS Genetics* 12 (4): e1005951.

Jaari, Sonja, Meng-Hua Li, and Juha Merilä. 2009. "A First-Generation Microsatellite-Based Genetic Linkage Map of the Siberian Jay (Perisoreus Infaustus): Insights into Avian Genome Evolution." *BMC Genomics* 10 (1): 1.

Jensen, Henrik, Bernt-Erik SAEther, Thor Harald Ringsby, Jarle Tufto, Simon C. Griffith, and Hans Ellegren. 2004. "Lifetime Reproductive Success in Relation to Morphology in the House Sparrow Passer Domesticus." *The Journal of Animal Ecology* 73 (4): 599–611.

Johnsson, Martin, Andrew Whalen, Roger Ros-Freixedes, Gregor Gorjanc, Ching-Yi Chen, William O. Herring, Dirk-Jan de Koning, and John M. Hickey. 2021. "Genetic Variation in Recombination Rate in the Pig." *Genetics, Selection, Evolution: GSE* 53 (1): 54.

Johnston, Susan E., Camillo Bérénos, Jon Slate, and Josephine M. Pemberton. 2016. "Conserved Genetic Architecture Underlying Individual Recombination Rate Variation in a Wild Population of Soay Sheep (Ovis Aries)." *Genetics* 203 (1): 583–98.

Johnston, Susan E., Nancy Chen, and Emily B. Josephs. 2022. "Taking Quantitative Genomics into the Wild." *Proceedings of the Royal Society B: Biological Sciences* 289 (September): 20221930.

Johnston, Susan E., Jisca Huisman, and Josephine M. Pemberton. 2018. "A Genomic Region Containing REC8 and RNF212B Is Associated with Individual Recombination Rate Variation in a Wild Population of Red Deer (Cervus Elaphus)." *G3* 8 (7): 2265–76.

Kadri, Naveen Kumar, Chad Harland, Pierre Faux, Nadine Cambisano, Latifa Karim, Wouter Coppieters, Sébastien Fritz, et al. 2016. "Coding and Noncoding Variants in HFM1, MLH3, MSH4, MSH5, RNF212, and RNF212B Affect Recombination Rate in Cattle." *Genome Research* 26 (10): 1323–32.

Kawakami, Takeshi, Carina F. Mugal, Alexander Suh, Alexander Nater, Reto Burri, Linnéa Smeds, and Hans Ellegren. 2017. "Whole-Genome Patterns of Linkage Disequilibrium across Flycatcher Populations Clarify the Causes and Consequences of Fine-Scale Recombination Rate Variation in Birds." *Molecular Ecology* 26 (16): 4158–72.

Kawakami, Takeshi, Linnéa Smeds, Niclas Backström, Arild Husby, Anna Qvarnström, Carina F. Mugal,
Pall Olason, and Hans Ellegren. 2014. "A High-Density Linkage Map Enables a Second-
Generation Collared Flycatcher Genome Assembly and Reveals the Patterns of Avian
Recombination Rate Variation and Chromosomal Evolution." *Molecular Ecology* 23 (16): 4035–
58.
Kemppainen, Petri, and Arild Husby. 2018. "Inference of Genetic Architecture from Chromosome
Partitioning Analyses Is Sensitive to Genome Variation, Sample Size, Heritability and Effect Size
Distribution." *Molecular Ecology Resources* 18 (4): 767–77.
Kinsella, Cormac M., Francisco J. Ruiz-Ruano, Anne-Marie Dion-Côté, Alexander J. Charles, Toni I.
Gossmann, Josefa Cabrero, Dennis Kappei, et al. 2019. "Programmed DNA Elimination of
Germline Development Genes in Songbirds." *Nature Communications* 10 (1): 5468.
Kivikoski, Mikko, Pasi Rastas, Ari Löytynoja, and Juha Merilä. 2023. "Predicting Recombination
Frequency from Map Distance." *Heredity* 130 (3): 114–21.
Koehler, K. E., R. S. Hawley, S. Sherman, and T. Hassold. 1996. "Recombination and Nondisjunction in
Humans and Flies." *Human Molecular Genetics* 5 Spec No: 1495–1504.
Kondrashov, A. S. 1988. "Deleterious Mutations and the Evolution of Sexual Reproduction." *Nature* 336
(6198): 435–40.
Kong, Augustine, John Barnard, Daniel F. Gudbjartsson, Gudmar Thorleifsson, Gudrun Jonsdottir, Sigrun
Sigurdardottir, Bjorgvin Richardsson, et al. 2004. "Recombination Rate and Reproductive
Success in Humans." *Nature Genetics* 36 (11): 1203–6.
Kong, Augustine, Gudmar Thorleifsson, Michael L. Frigge, Gisli Masson, Daniel F. Gudbjartsson, Rasmus
Villemoes, Erna Magnusdottir, Stefania B. Olafsdottir, Unnur Thorsteinsdottir, and Kari
Stefansson. 2014. "Common and Low-Frequency Variants Associated with Genome-Wide
Recombination Rate." *Nature Genetics* 46 (1): 11–16.
Kumar, Sudhir, Michael Suleski, Jack M. Craig, Adrienne E. Kasprowicz, Maxwell Sanderford, Michael Li,
Glen Stecher, and S. Blair Hedges. 2022. "TimeTree 5: An Expanded Resource for Species
Divergence Times." *Molecular Biology and Evolution* 39 (8).
https://doi.org/10.1093/molbev/msac174.
Lenormand, Thomas. 2003. "The Evolution of Sex Dimorphism in Recombination." *Genetics* 163 (2):
811–22.
Lenormand, Thomas, and Julien Dutheil. 2005. "Recombination Difference between Sexes: A Role for
Haploid Selection." *PLoS Biology* 3 (3): e63.
Lisachov, A. P., L. P. Malinovskaya, A. V. Druzyaka, P. M. Borodin, and A. A. Torgasheva. 2017.
"Synapsis and Recombination of Autosomes and Sex Chromosomes in Two Terns (Sternidae,
Charadriiformes, Aves)." *Vavilovskii Zhurnal Genetiki i Selektsii* 21 (2): 259–68.
Lundregan, Sarah L., Ingerid J. Hagen, Jostein Gohli, Alina K. Niskanen, Petri Kemppainen, Thor Harald
Ringsby, Thomas Kvalnes, et al. 2018. "Inferences of Genetic Architecture of Bill Morphology in
House Sparrow Using a High-Density SNP Array Point to a Polygenic Basis." *Molecular Ecology*
27 (17): 3498–3514.
Ma, Li, Jeffrey R. O'Connell, Paul M. VanRaden, Botong Shen, Abinash Padhi, Chuanyu Sun, Derek M.
Bickhart, et al. 2015. "Cattle Sex-Specific Recombination and Genetic Control from a Large
Pedigree Analysis." *PLoS Genetics* 11 (11): e1005387.
MacLennan, Marie, James H. Crichton, Christopher J. Playfoot, and Ian R. Adams. 2015. "Oocyte
Development, Meiosis and Aneuploidy." *Seminars in Cell & Developmental Biology* 45
(September): 68–76.
Malinovskaya, Lyubov P., Katerina Tishakova, Elena P. Shnaider, Pavel M. Borodin, and Anna A.
Torgasheva. 2020. "Heterochiasmy and Sexual Dimorphism: The Case of the Barn Swallow
(Hirundo Rustica, Hirundinidae, Aves)." *Genes* 11 (10): 1119.
Malinovskaya, Lyubov P., Kira S. Zadesenets, Tatyana V. Karamysheva, Ekaterina A. Akberdina, Elena
A. Kizilova, Margarita V. Romanenko, Elena P. Shnaider, et al. 2020. "Germline-Restricted
Chromosome (GRC) in the Sand Martin and the Pale Martin (Hirundinidae, Aves): Synapsis,
Recombination and Copy Number Variation." *Scientific Reports* 10 (1): 1058.
Malinovskaya, Lyubov, Elena Shnaider, Pavel Borodin, and Anna Torgasheva. 2018. "Karyotypes and
Recombination Patterns of the Common Swift (Apus Apus Linnaeus, 1758) and Eurasian Hobby
(Falco Subbuteo Linnaeus, 1758)." *Avian Research* 9 (1). https://doi.org/10.1186/s40657-018-
0096-7.

Mank, Judith E. 2009. "The Evolution of Heterochiasmy: The Role of Sexual Selection and Sperm
Competition in Determining Sex-Specific Recombination Rates in Eutherian Mammals." *Genetics*
Research 91 (5): 355–63.
Mueller, Jakob C., Stephen A. Schlebusch, Yifan Pei, Manon Poignet, Niki Vontzou, Francisco J. Ruiz-
Ruano, Tomáš Albrecht, et al. 2023. "Micro Germline-Restricted Chromosome in Blue Tits:
Evidence for Meiotic Functions." *Molecular Biology and Evolution* 40 (5).
https://doi.org/10.1093/molbev/msad096.
Muñoz-Fuentes, Violeta, Marina Marcet-Ortega, Gorka Alkorta-Aranburu, Catharina Linde Forsberg, Jane
M. Morrell, Esperanza Manzano-Piedras, Arne Söderberg, et al. 2015. "Strong Artificial Selection
in Domestic Mammals Did Not Result in an Increased Recombination Rate." *Molecular Biology*
and Evolution 32 (2): 510–23.
Myers, Simon, Leonardo Bottolo, Colin Freeman, Gil McVean, and Peter Donnelly. 2005. "A Fine-Scale
Map of Recombination Rates and Hotspots across the Human Genome." *Science* 310 (5746):
321–24.
Myers, Simon, Rory Bowden, Afidalina Tumian, Ronald E. Bontrop, Colin Freeman, Tammie S. MacFie,
Gil McVean, and Peter Donnelly. 2010. "Drive against Hotspot Motifs in Primates Implicates the
PRDM9 Gene in Meiotic Recombination." *Science (New York, N.Y.)* 327 (5967): 876–79.
Nei, M. 1969. "Linkage Modifications and Sex Difference in Recombination." *Genetics* 63 (3): 681–99.
Niskanen, Alina K., Anna M. Billing, Håkon Holand, Ingerid J. Hagen, Yimen G. Araya-Ajoy, Arild Husby,
Bernt Rønning, et al. 2020. "Consistent Scaling of Inbreeding Depression in Space and Time in a
House Sparrow Metapopulation." *Proceedings of the National Academy of Sciences of the United*
States of America 117 (25): 14584–92.
Oers, K. van, A. W. Santure, I. De Cauwer, N. E. M. van Bers, R. P. M. A. Crooijmans, B. C. Sheldon, M.
E. Visser, J. Slate, and M. A. M. Groenen. 2014. "Replicated High-Density Genetic Maps of Two
Great Tit Populations Reveal Fine-Scale Genomic Departures from Sex-Equal Recombination
Rates." *Heredity* 112 (3): 307–16.
Otto, S. P., and N. H. Barton. 2001. "Selection for Recombination in Small Populations." *Evolution;*
International Journal of Organic Evolution 55 (10): 1921–31.
Otto, Sarah P., and Thomas Lenormand. 2002. "Resolving the Paradox of Sex and Recombination."
Nature Reviews. Genetics 3 (4): 252–61.
Pan, Jing, Mariko Sasaki, Ryan Kniewel, Hajime Murakami, Hannah G. Blitzblau, Sam E. Tischfield, Xuan
Zhu, et al. 2011. "A Hierarchical Combination of Factors Shapes the Genome-Wide Topography
of Yeast Meiotic Recombination Initiation." *Cell* 144 (5): 719–31.
Pei, Yifan, Wolfgang Forstmeier, Francisco J. Ruiz-Ruano, Jakob C. Mueller, Josefa Cabrero, Juan Pedro
M. Camacho, Juan D. Alché, et al. 2022. "Occasional Paternal Inheritance of the Germline-
Restricted Chromosome in Songbirds." *Proceedings of the National Academy of Sciences of the*
United States of America 119 (4): e2103960119.
Peñalba, Joshua V., Yuan Deng, Qi Fang, Leo Joseph, Craig Moritz, and Andrew Cockburn. 2020.
"Genome of an Iconic Australian Bird: High-Quality Assembly and Linkage Map of the Superb
Fairy-Wren (Malurus Cyaneus)." *Molecular Ecology Resources* 20 (2): 560–78.
Petit, Morgane, Jean-Michel Astruc, Julien Sarry, Laurence Drouilhet, Stéphane Fabre, Carole Moreno,
and Bertrand Servin. 2017. "Variation in Recombination Rate and Its Genetic Determinism in
Sheep Populations." *Genetics* 207 (2): 767–84.
Pigozzi, M. I. 2001. "Distribution of MLH1 Foci on the Synaptonemal Complexes of Chicken Oocytes."
Cytogenetics and Cell Genetics 95 (3–4): 129–33.
Pigozzi, M. I., and A. J. Solari. 1998. "Germ Cell Restriction and Regular Transmission of an Accessory
Chromosome That Mimics a Sex Body in the Zebra Finch, Taeniopygia Guttata." *Chromosome*
Research: An International Journal on the Molecular, Supramolecular and Evolutionary Aspects
of Chromosome Biology 6 (2): 105–13.
Ranke, Peter S., Yimen G. Araya-Ajoy, Thor Harald Ringsby, Henrik Pärn, Bernt Rønning, Henrik Jensen,
Jonathan Wright, and Bernt-Erik Saether. 2021. "Spatial Structure and Dispersal Dynamics in a
House Sparrow Metapopulation." *The Journal of Animal Ecology* 90 (12): 2767–81.
Rastas, Pasi. 2017. "Lep-MAP3: Robust Linkage Mapping Even for Low-Coverage Whole Genome
Sequencing Data." *Bioinformatics* 33 (23): 3726–32.
Ritz, Kathryn R., Mohamed A. F. Noor, and Nadia D. Singh. 2017. "Variation in Recombination Rate:
Adaptive or Not?" *Trends in Genetics: TIG* 33 (5): 364–74.

Robledo-Ruiz, Diana A., Han Ming Gan, Parwinder Kaur, Olga Dudchenko, David Weisz, Ruqayya Khan,
Erez Lieberman Aiden, et al. 2022. "Chromosome-Length Genome Assembly and Linkage Map of
a Critically Endangered Australian Bird: The Helmeted Honeyeater." *GigaScience* 11 (March).
https://doi.org/10.1093/gigascience/giac025.
Rönnegård, Lars, S. Eryn McFarlane, Arild Husby, Takeshi Kawakami, Hans Ellegren, and Anna
Qvarnström. 2016. "Increasing the Power of Genome Wide Association Studies in Natural
Populations Using Repeated Measures - Evaluation and Implementation." *Methods in Ecology
and Evolution / British Ecological Society* 7 (7): 792–99.
Ross-Ibarra, Jeffrey. 2004. "The Evolution of Recombination under Domestication: A Test of Two
Hypotheses." *The American Naturalist* 163 (1): 105–12.
Saatoglu, Dilan, Alina K. Niskanen, Markku Kuismin, Peter S. Ranke, Ingerid J. Hagen, Yimen G. Araya-
Ajoy, Thomas Kvalnes, et al. 2021. "Dispersal in a House Sparrow Metapopulation: An Integrative
Case Study of Genetic Assignment Calibrated with Ecological Data and Pedigree Information."
Molecular Ecology 30 (19): 4740–56.
Samuk, Kieran, Brenda Manzano-Winkler, Kathryn R. Ritz, and Mohamed A. F. Noor. 2020. "Natural
Selection Shapes Variation in Genome-Wide Recombination Rate in Drosophila Pseudoobscura."
Current Biology: CB 30 (8): 1517-1528.e6.
Santure, Anna W., and Dany Garant. 2018. "Wild GWAS—Association Mapping in Natural Populations."
Molecular Ecology Resources 18 (4): 729–38.
Sardell, Jason M., and Mark Kirkpatrick. 2020. "Sex Differences in the Recombination Landscape." *The
American Naturalist* 195 (2): 361–79.
Servin, Bertrand. 2021. *YAPP: Yet Another Phasing Program*. https://yapp-doc.netlify.app/.
Singhal, Sonal, Ellen M. Leffler, Keerthi Sannareddy, Isaac Turner, Oliver Venn, Daniel M. Hooper, Alva I.
Strand, et al. 2015. "Stable Recombination Hotspots in Birds." *Science (New York, N.Y.)* 350
(6263): 928–32.
Smukowski, C. S., and M. A. F. Noor. 2011. "Recombination Rate Variation in Closely Related Species."
Heredity 107 (6): 496–508.
Stapley, J., T. R. Birkhead, T. Burke, and J. Slate. 2008. "A Linkage Map of the Zebra Finch Taeniopygia
Guttata Provides New Insights into Avian Genome Evolution." *Genetics* 179 (1): 651–67.
Stapley, Jessica, Philine G. D. Feulner, Susan E. Johnston, Anna W. Santure, and Carole M. Smadja.
2017. "Variation in Recombination Frequency and Distribution across Eukaryotes: Patterns and
Processes." *Philosophical Transactions of the Royal Society of London. Series B, Biological
Sciences* 372 (1736): 20160455.
Stephens, Matthew. 2017. "False Discovery Rates: A New Deal." *Biostatistics* 18 (2): 275–94.
Stubberud, Marlene Waege, Ane Marlene Myhre, Håkon Holand, Thomas Kvalnes, Thor Harald Ringsby,
Bernt-Erik Saether, and Henrik Jensen. 2017. "Sensitivity Analysis of Effective Population Size to
Demographic Parameters in House Sparrow Populations." *Molecular Ecology* 26 (9): 2449–65.
Torgasheva, Anna A., and Pavel M. Borodin. 2017. "Immunocytological Analysis of Meiotic
Recombination in the Gray Goose (Anser Anser)." *Cytogenetic and Genome Research* 151 (1):
27–35.
Torgasheva, Anna A., Lyubov P. Malinovskaya, Kira S. Zadesenets, Tatyana V. Karamysheva, Elena A.
Kizilova, Ekaterina A. Akberdina, Inna E. Pristyazhnyuk, et al. 2019. "Germline-Restricted
Chromosome (GRC) Is Widespread among Songbirds." *Proceedings of the National Academy of
Sciences of the United States of America* 116 (24): 11845–50.
Trivers, Robert. 1988. "Sex Differences in Rates of Recombination and Sexual Selection." In *Evolution of
Sex: An Examination of Current Ideas*, edited by Bruce R. Levin Richard E. Michod, 270–86.
Sunderland, Massachusetts: Sinauer Associates Inc.
Veller, Carl, Nancy Kleckner, and Martin A. Nowak. 2019. "A Rigorous Measure of Genome-Wide Genetic
Shuffling That Takes into Account Crossover Positions and Mendel's Second Law." *Proceedings
of the National Academy of Sciences of the United States of America* 116 (5): 1659–68.
Visscher, Peter M., Naomi R. Wray, Qian Zhang, Pamela Sklar, Mark I. McCarthy, Matthew A. Brown,
and Jian Yang. 2017. "10 Years of GWAS Discovery: Biology, Function, and Translation." *The
American Journal of Human Genetics* 101 (1): 5–22.
Warren, Wesley C., David F. Clayton, Hans Ellegren, Arthur P. Arnold, Ladeana W. Hillier, Axel Künstner,
Steve Searle, et al. 2010. "The Genome of a Songbird." *Nature* 464 (7289): 757–62.

Weng, Ziqing, Anna Wolc, Hailin Su, Rohan L. Fernando, Jack C. M. Dekkers, Jesus Arango, Petek
Settar, Janet E. Fulton, Neil P. O'Sullivan, and Dorian J. Garrick. 2019. "Identification of
Recombination Hotspots and Quantitative Trait Loci for Recombination Rate in Layer Chickens."
Journal of Animal Science and Biotechnology 10 (1): 20.
Yang, Jian, S. Hong Lee, Michael E. Goddard, and Peter M. Visscher. 2011. "GCTA: A Tool for Genome-
Wide Complex Trait Analysis." *The American Journal of Human Genetics* 88 (1): 76–82.
Yang, Jian, Teri A. Manolio, Louis R. Pasquale, Eric Boerwinkle, Neil Caporaso, Julie M. Cunningham,
Mariza de Andrade, et al. 2011. "Genome Partitioning of Genetic Variation for Complex Traits
Using Common SNPs." *Nature Genetics* 43 (6): 519–25.6666