Single cell RNA sequencing unravels the transcriptional network underlying zebrafish retina regeneration

Laura Celotto[1], Fabian Rost[2], Anja Machate[1], Juliane Bläsche[2], Andreas Dahl[2], Anke Weber[1], Stefan Hans[1], and Michael Brand[*1]

[1] Technische Universität Dresden, CRTD - Center for Regenerative Therapies Dresden, Center for Molecular and Cellular Bioengineering (CMCB), Fetscherstraße 105, 01307 Dresden, Germany

[2] Technische Universität Dresden, DRESDEN-Concept Genome Center, Center for Molecular and Cellular Bioengineering (CMCB), Fetscherstraße 105, 01307 Dresden, Germany

*Corresponding author

ORCID: Laura Celotto: 0000-0002-9191-4056; Fabian Rost: 0000-0001-6466-2589; Andreas Dahl: 0000-0002-2668-8371; Anke Weber: 0000-0003-2220-658X; Stefan Hans: 0000-0003-0283-0211; Michael Brand: 0000-0001-5711-6512

Keywords: Retina, regeneration, zebrafish, Müller glia, progenitors, light lesion, photoreceptors, scRNAseq

SUMMARY

In the lesioned zebrafish retina, Müller glia produce multipotent retinal progenitors that generate all retinal neurons, replacing lost cell types. To study the molecular mechanisms linking Müller glia reactivity to progenitor production and neuronal differentiation, we used single cell RNA sequencing of Müller glia, progenitors and regenerated progeny from uninjured and light-lesioned retinae. We discover an injury-induced Müller glia differentiation trajectory that leads into a cell population with a hybrid identity expressing marker genes of Müller glia and progenitors. A glial self-renewal and a neurogenic trajectory depart from the hybrid cell population. We further observe that neurogenic progenitors progressively differentiate to generate retinal ganglion cells first and bipolar cells last, similar to the events observed during retinal development. Our work provides a comprehensive description of Müller glia and progenitor transcriptional changes and fate decisions in the regenerating retina, which are key to tailor cell differentiation and replacement therapies for retinal dystrophies in humans.

INTRODUCTION

The vertebrate retina is the neural tissue in the back of the eyes that detects and processes visual stimuli prior to transmission to the brain (Dowling, 1987). The retina has a layered structure, with outer, inner and ganglion cell nuclear layers alternating with the outer and inner plexiform layers (Amini et al., 2018; Hoon et al., 2014; Stenkamp, 2015). The nuclear layers comprise the nuclei of retinal neurons, whereas the plexiform layers contain neuronal synapses. There are six distinct classes of retinal neurons and one main type of glia: retinal ganglion, cone photoreceptor, horizontal, amacrine, rod photoreceptor and bipolar cells, as well as Müller glia. This

heterogeneity arises during development in the indicated order from a common pool

of multipotent retinal progenitor cells (RPCs), which progressively restrict their

competence to generate early and mid-late born cell fates (Agathocleous and Harris,

2009; Alexiades and Cepko, 1997; Amini et al., 2018; Cepko, 2014; Cepko et al.,

1996; Livesey and Cepko, 2001; Stenkamp, 2015, 2007; Turner and Cepko, 1987;

Weber et al., 2014; Young, 1985). Retinal development, structure and function are

highly conserved among vertebrates, including the teleost zebrafish (*Danio rerio*) with

some variations. For instance, humans and zebrafish possess a cone-dominated

retina adapted for diurnal visual tasks, whereas nocturnal animals, like mice and rats,

have rod-dominated retinae adapted for a dim light lifestyle (Fleisch et al., 2011;

Saade et al., 2013; Stenkamp, 2007). Intriguingly, and in contrast to warm-blooded

vertebrates like birds and mammals, adult zebrafish can fully regenerate their retinae

upon lesions affecting different neuronal cell types, depending on the injury

paradigm. For instance, intense light ablates preferentially light-sensing rod and cone

photoreceptors in the central and dorsal retina, whereas application of neurotoxins,

like ouabain or NMDA, affects mainly retinal ganglion and amacrine cells (Bernardos

et al., 2007; Fimbel et al., 2007; Powell et al., 2016; Sherpa et al., 2008; Vihtelic et

al., 2006; Vihtelic and Hyde, 2000; Weber et al., 2013). Independently of the lesion

type, Müller glia (MG), acting as resident stem cells, are the primary cellular origin of

regenerated cells. Specifically, zebrafish MG react to the lesion by partial

dedifferentiation that takes place between 6 and 24 hours post lesion, followed by

cell cycle re-entry, including up-regulation of proliferating cell nuclear antigen

(PCNA), a marker of the S-G2/M cell cycle phases (Bernardos et al., 2007; Fausett et

al., 2008; Nagashima et al., 2013; Raymond et al., 2006). PCNA expression in MG is

initiated at 24-31 hours post lesion (Lahne et al., 2015; Nelson et al., 2013; Thummel

et al., 2008) and can still be detected up to 2 days post lesion (Nagashima et al.,

2013; Nelson et al., 2013). PCNA upregulation is followed by asymmetric cell division of the MG that produces renewed MG and multipotent RPCs, which are detected starting from 51 hours post lesion (Gorsuch and Hyde, 2014; Lahne et al., 2020; Lenkowski and Raymond, 2014; Thummel et al., 2008; Wan and Goldman, 2016). Dedifferentiation of MG to a progenitor-like state is not complete, as MG upregulate progenitor-associated genes like *pax6* or *vsx2*, but retain their glial morphology. In this context, glial markers like Glial Fibrillary Acidic Protein (*gfap*) as well as glutamine synthase (*glula*) have been reported to be either retained or downregulated, depending on the lesion paradigm, the regeneration time point, the antibody and the transgenic line used to label MG (Fimbel et al., 2007; Lenkowski and Raymond, 2014; Nagashima et al., 2013; Thummel et al., 2008). However, in all reported cases, MG-derived RPCs do not express *gfap* (Lenkowski and Raymond, 2014; Thummel et al., 2008). Asymmetric cell division of MG occurs between 31 hours and 2 days post-lesion, followed by further proliferation of MG-derived RPCs that peaks at 4-5 days post lesion, depending on the injury paradigm (Bernardos et al., 2007; Fausett and Goldman, 2006; Fimbel et al., 2007; Lahne et al., 2015; Nagashima et al., 2013; Nelson et al., 2013; Thummel et al., 2008). Studies on zebrafish retina regeneration have so far mainly focused on very early phases of MG dedifferentiation and RPC production (Fausett et al., 2008; Nelson et al., 2013; Ramachandran et al., 2010, 2011; Thummel et al., 2008, 2010; Wan et al., 2012; Zhao et al., 2014). Recent work showed that MG initially assume a reactive state, characterized by the expression of *hmga1a* and *yap1*, prior to giving rise to proliferative, neurogenic RPCs (Hoang et al., 2020). Consequently, impaired Yap1 or Hmga1a function impedes MG reprogramming and neurogenesis in the regenerating retina (Hoang et al., 2020). In contrast, the molecular and cellular identities of self-renewed MG and MG-derived RPCs remain elusive. RPCs have fusiform nuclei that

closely associate with GFAP-positive glial fibers across the inner nuclear layer. They also express markers of multipotency, like Pax6 and Sox2, as well as markers of neurogenesis, like Atoh7 (Fausett and Goldman, 2006; Gorsuch et al., 2017; Nagashima et al., 2013; Raymond et al., 2006; Thummel et al., 2010). Previous studies have investigated RPCs molecular identity mainly at the population level (Cameron et al., 2005; Hoang et al., 2020; Kassen et al., 2007; Kramer et al., 2021; Lahne et al., 2021; Morris et al., 2011; Ng Chi Kei et al., 2017; Powell et al., 2016; Qin et al., 2009; Ramachandran et al., 2012; Thummel et al., 2008). However, a detailed characterization of the transcriptional network underlying RPC production and RPC differentiation, as well as the lineage relationships between MG, RPCs and differentiating progeny is currently missing. Here, we profiled the transcriptome of MG and MG-derived RPCs in response to an intense light lesion, and followed their differentiation trajectories towards regenerated retinal neurons at the single cell level. In particular, we addressed the transcriptional state of reactive MG prior to and after their asymmetric cell division. Moreover, we addressed whether MG-derived RPCs are heterogeneous and to what extent they express molecular markers that are observed during embryonic retinogenesis (Xu et al., 2020). To this aim, we employed a short-term lineage tracing strategy that allowed the isolation of individual MG and MG-derived cells of the regenerating zebrafish retina, which were subsequently subjected to droplet-based, 10x Genomics single cell RNA sequencing (scRNAseq). We find that MG can be non-reactive in uninjured and lesioned conditions, as well as reactive upon lesion. Reactive MG give rise to a hybrid cell population, which replenishes the pool of MG as well as feeds into the pool of neurogenic RPCs. Moreover, MG-derived RPCs are highly heterogeneous and display a progressive restriction in their competence during subsequent differentiation. They upregulate developmental markers of retinogenesis and produce regenerated neurons partially

according to the developmental birth order, with retinal ganglion cells born first and

bipolar cells born last.

RESULTS

Isolation of individual MG, RPCs and regenerated progeny using fluorescent

reporters

Previous work on the regenerating zebrafish retina described the early phases of

MG de-differentiation and RPC production (Cameron et al., 2005; Fausett et al.,

2008; Fausett and Goldman, 2006; Lahne et al., 2015; Nagashima et al., 2013;

Nelson et al., 2013; Thummel et al., 2010). However, knowledge on the molecular

identity and cell fate choices of RPCs is scarce and based on analyses at a

population level (Cameron et al., 2005; Hoang et al., 2020; Kramer et al., 2021;

Lahne et al., 2021; Ng Chi Kei et al., 2017; Powell et al., 2016; Qin et al., 2009;

Ramachandran et al., 2012; Thummel et al., 2008). To study MG activation, RPC

identity and their relationship with MG in more detail, we profiled the transcriptome of

MG and MG-derived RPCs and followed their differentiation trajectories towards

regenerated retinal neurons at the single cell level. To do so, we devised a strategy

to isolate individual MG, RPCs and regenerated progeny, using retinae of uninjured

and light-lesioned fish carrying the two fluorescent reporters *pcna:EGFP* and

gfap:mCherry. The former drives the expression of EGFP under the regulatory

elements of *proliferating cell nuclear antigen* (*pcna*), a known marker of proliferating

cells, and thereby labels actively dividing cells, and, due to the persistence of EGFP,

their short-term progeny. The other reporter drives expression of mCherry under the

regulatory elements of *glial fibrillary acidic protein* (*gfap*), a known marker of

zebrafish MG. Both transgenes recapitulate the endogenous expression in the adult

zebrafish retina faithfully (Figure1-figure supplement 1). Immunohistochemistry

against the proliferation marker PCNA co-labeled *pcna:EGFP*-positive cells mainly in

the ciliary marginal zone, one of the stem cells niches in the adult zebrafish retina

(Bernardos et al., 2007; Raymond et al., 2006). Occasionally, *pcna:EGFP*/PCNA-

double positive cells were observed in the central retina as well (Figure1-figure

supplement 1A). The MG-specific antibody Zrf1 labeled virtually all *gfap:mCherry*-

positive cells in the central retina as well as in the ciliary marginal zone (Bernardos et

al., 2007) (Figure1-figure supplement 1B). We also addressed the expression of the

two fluorescent reporter lines following light lesion, at three different time points upon

injury. First, 44 hours post light lesion (hpl), when MG are about to undergo or have

already completed their asymmetric cell division giving rise to the first RPCs and self-

renewed MG (Lahne et al., 2015; Nagashima et al., 2013). Second, 4 days post light

lesion (dpl), representing the peak of RPC proliferation (Bernardos et al., 2007;

Fausett and Goldman, 2006) and third, 6 dpl, when the first differentiating,

regenerated progeny arise (Bernardos et al., 2007). Immunohistochemistry against

EGFP and mCherry in double transgenic *Tg(pcna:EGFP);Tg(gfap:mCherry)* animals

corroborated the expected expression pattern of the transgenes in the central retina

at the chosen time points (Figure1-figure supplement 2). Uninjured, central retinae

displayed no EGFP, but strong mCherry signal consistent with non-proliferating MG

(white arrowheads). In contrast, numerous EGFP/mCherry double-positive cells were

present at 44 hpl, indicating potential cell-cycle re-entry of MG (yellow arrowheads).

At 4 dpl, overall mCherry expression was reduced, consistent with the previously

described downregulation of the *gfap* promoter by MG-derived RPCs (Fimbel et al.,

2007; Lenkowski and Raymond, 2014; Thummel et al., 2008). In contrast, EGFP was

strongly found throughout all retinal layers, matching the reported peak in

proliferation of MG-derived RPCs (red arrowheads) (Bernardos et al., 2007; Fausett

and Goldman, 2006; Fimbel et al., 2007; Lenkowski and Raymond, 2014; Thummel

et al., 2008). Finally, in comparison to 4 dpl, an increase in mCherry accompanies a

decline in EGFP signal at 6 dpl (yellow arrowhead). In order to obtain single cells of

MG, RPCs and regenerated progeny, we utilized the persistence of EGFP and

mCherry fluorescence, providing a short-term label. Initially, only MG express

mCherry in uninjured, central retinae, but activate EGFP in preparation of the

asymmetric cell division upon light lesion at 44 hpl (Figure 1A). Firstborn RPCs inherit

high levels of persisting fluorescent mCherry protein from dividing MG and slowly

degrade/dilute the protein during subsequent rounds of cell division. Hence, MG-

derived, but further dividing, RPCs are EGFP-positive, but low mCherry-positive at 4

dpl. At 6 dpl, RPCs have further diluted the mCherry reporter, thus displaying only a

faint mCherry signal, but remain EGFP-positive. Similarly, RPC-derived, regenerated

progeny is characterized by even lower levels of intrinsic mCherry fluorescence, and

low EGFP signal during subsequent cell differentiation at 6 dpl. Following this

rationale, we dissected retinae of uninjured and light-lesioned samples, dissociated

them into single cell suspensions that were subjected to flow cytometry followed by

droplet-based, 10x Genomics scRNAseq (Figure 1B). Flow cytometry plots confirmed

the rationale of the above-mentioned strategy (Figure 1C and Figure1-figure

supplement 3). Figure 1C depicts contour plots, which in flow cytometry represent the

relative frequency of events, i.e. cells, in a sample. Each contour line in the plot

encloses an equal percentage of cells, and contour lines that are closely packed

indicate a high concentration of cells. In all samples, we detected non-fluorescent

and EGFP-positive only cells, consistent with non-proliferating and proliferating cells

in the adult zebrafish retina. We also detected cell populations expressing only

mCherry at all sampled time points, most likely representing non-reactive MG. In

addition to cells expressing either EGFP or mCherry, we find EGFP/mCherry-double

positive cells in all samples consistent with proliferating MG originating from the

ciliary marginal zone and the central retina (see Figure1-figure supplement 1 and 2).

The EGFP/mCherry-double positive population increased at 44 hpl, presumably due

to an increase in the cell number of reactive, proliferating MG. A second, EGFP-

positive, lower mCherry-positive cell population was detected at 44 hpl, presumably

including the very first RPCs resulting from MG cell division at this time point. At 4

dpl, the large EGFP/mCherry-double positive cell population was still present, but

displayed a lower mCherry, yet an increased EGFP signal, in line with MG-derived

RPCs, which exhibit maximal proliferation at 4 dpl (Bernardos et al., 2007; Fausett

and Goldman, 2006). At 6 dpl, a further decreased mCherry intensity was observed

for the EGFP/mCherry-double positive cell population. In order to profile the

transcriptome of individual MG, RPCs and progeny of the uninjured and light lesioned

retinae, we dissected retinae of the before-mentioned time points (uninjured, 44 hpl,

4 dpl, 6 dpl) and eventually sorted 15.000 cells per time point. We applied a sorting

gate that included both mCherry-positive only and EGFP/mCherry-double positive

cells at the sampled time points (Figure 1C and Figure 1-figure supplement 3). Non-

fluorescent as well as EGFP- only positive cells were not included. Taken together,

we devised a short-term lineage tracing strategy that allows the isolation of individual

MG and MG-derived cells of the regenerating zebrafish retina, which can be

subjected to droplet-based,10x Genomics.

scRNAseq identifies 15 distinct cell populations in the light-lesioned retina

Sorted cells underwent 10x Genomics scRNAseq and 13.139 cells passed the

sequencing quality control. We detected a median of 1.407 genes and 4.061 mean

read counts per cell. Initially, we identified three clusters for the full dataset (Figure 2-

figure supplement 1A): a retinal core dataset and two additional clusters identified as

microglia and ribosomal gene-enriched cells (Figure 2-figure supplement 1B). Both

populations were excluded from the subsequent analysis. Re-clustering of the core

dataset, which eventually included 11.690 cells, revealed 15 potential, different cell

populations (Figure 2A). From left to right on the UMAP, which visualizes the cell

lineage trajectory, we observed the progressive emergence of MG, RPC and progeny

clusters. We assigned the cellular identity by inspecting the top 100, upregulated

marker genes per cluster (supplementary file 1). In detail, we identified four MG

clusters (clusters 1 to 4) that locate to the left and middle portion of the map, followed

by five RPC clusters (clusters 5 to 9) locating to the right side of the map.

Regenerated progeny branch from the RPCs and include retinal ganglion (cluster

10), amacrine (cluster 13) and bipolar cells (cluster 15) at the bottom-right, as well as

red and blue cones (clusters 11 and 12) at the top-right of the map. Additionally,

there is a second cluster of amacrine cells (cluster 14) that separates from the first

amacrine cell cluster.

To investigate how the composition of cells changed during regeneration, we

computed embedding densities for each time point (Figure 2B). Similar to the

progression of cellular identities, cells progressed over time from left to right, from the

uninjured, to the 44 hpl, the 4 dpl, as well as the 6 dpl enriched samples. As

expected, cells belonging to non-reactive MG were found at all sampled time points.

In particular, the vast majority of cells of the uninjured sample belonged to cluster 1

as well as cluster 2. However, we also observed a small proportion of cells (Figure

2B) aligning in a trajectory from non-reactive MG (left side of the map) through

reactive MG (middle part) to RPCs and differentiating progeny (right side) in the

uninjured sample. At 44 hpl, cells mapping to non-reactive MG were still abundantly

present, although mostly shifted right towards a cluster representing reactive MG

(cluster 3). Cells belonging to this cluster of reactive MG were also highly enriched in

this sample. In contrast, cells mapping to the second cluster of reactive MG (cluster

4) as well as to RPCs (clusters 5-9) were overrepresented at 4 dpl, in line with the

reported peak in proliferation at this time point (Bernardos et al., 2007; Fausett and

Goldman, 2006). Cells mapping to RPCs were still frequently present at 6 dpl.

Moreover, cells belonging to regenerated progeny (clusters 10-15), which first appear

at 4 dpl, became predominantly enriched at 6 dpl. In addition to the overall

progression from left to right, we noted that cells mapping to reactive MG of the

second cluster (cluster 4) and the abutting portion of non-reactive MG (cluster 1)

were also enriched at this time point, indicating a reversed direction.

Analysis of the cell cycle state further supported our cluster annotation. Specifically,

known cell cycle marker genes (given in materials and methods) were used to

calculate the predicted cell cycle state (Macosko et al., 2015; Tirosh et al., 2016;

Whitfield et al., 2002), and the cell cycle phase (G1, S, G2/M) was projected onto the

UMAP of the single cell transcriptome (Figure 2C). Differentiating or differentiated

cells were considered as cells in G1 (Wu et al., 2021). Consistently, we observed that

most cells of the left side and lower middle part as well as cells branching from the

right side of the map were in G1, which comprised most non-reactive MG (clusters 1

and 2), one cluster of reactive MG (cluster 4), one RPC population (cluster 7) and the

regenerated progeny (clusters 10-15). Cells in S phase were also present on the left

side of the map, which represent non-reactive MG (clusters 1 and 2), but were in

particular enriched in two domains of the middle part corresponding to reactive MG

(cluster 3) and RPCs (clusters 5 and 7). In addition, two domains on the right side of

the map showed an accumulation of cells in S phase, which include photoreceptor

and horizontal cell precursors (clusters 8 and 9). The pattern of cells in G2/M phase

complemented the pattern of cells in G1 and S phase. Cell proliferation pattern was

further confirmed by *pcna* expression (Figure 2D). When plotting *pcna* expression

onto the main UMAP of the single cell transcriptome, we observed high counts in one

population of reactive MG (cluster 3) and in RPCs (clusters 5, 6, 8 and 9). In

summary, scRNAseq analysis enabled the establishment of a transcriptome map,

which reveals the progressive emergence of non-reactive MG, reactive MG and MG-

derived progenitors and regenerated progeny.

287   scRNAseq identifies two populations of non-reactive and two of reactive MG in

288   the light-lesioned retina

MG in the homeostatic zebrafish retina are characterized by the expression of GFAP,

as well as other canonical glial markers, like *apoeb*, *glula*, *cahz*, *rlbp1a* (Bernardos

and Raymond, 2006; Peterson et al., 2001; Raymond et al., 2006; Thummel et al.,

2008; Yurco and Cameron, 2005). In our scRNAseq dataset, we found clusters 1 and

2 expressing *apoeb*, *glula*, *cahz*, *rlbp1a* as marker genes (Figure 2-figure supplement

2 and supplementary file 1). When plotting *apoeb* expression onto the UMAP, we

observed high counts in clusters 1 and 2, as well as in clusters 3 and 4 (Figure 3A).

The glial marker *gfap*, found among the top 100 upregulated marker genes for cluster

4 (supplementary file 1), showed a similar pattern, with high counts in clusters 1 to 4,

but displayed high counts in cluster 5 as well (Figure 3A). Given the expression

pattern of *apoeb* and *gfap*, we identified clusters 1 to 4 as MG populations. Further

analysis indicated clusters 1 and 2 as non-reactive MG and clusters 3 and 4 as

reactive MG (Figure 3B, C). Non-reactive MG, which were mostly in G1 and did not

express any injury-specific markers (Figure 2-figure supplement 2) showed counts for

mstnb (Figure 3B). In addition, members of the retinoic acid signalling pathway, like

rdh10a, *aldh1a3*, *dhrs3a* (D'Aniello et al., 2015; Feng et al., 2010; Holmes and

Hempel, 2011) were enriched in these clusters (Figure 3-figure supplement 1A,
supplementary file 2).

We next looked into differentially expressed genes (DEG) between the non-reactive
MG clusters 1 and 2. For instance, the UMAP plot of *inhbaa*, an evolutionarily
conserved vertebrate glia marker (Hoang et al., 2020) showed strong expression only
in clusters 1, 3 and 4, but not in cluster 2 (Figure 3B, Figure 3-figure supplement 1A,
and supplementary file 2). Reactive MG on the other hand downregulated some glial
markers, like *glula* and *rlbp1a*, and upregulated inflammatory cytokines, like *crlf1a*,
matrix metalloproteases, like *mmp9*, and growth factors like *hbegfa* in the
regenerating retina (Figure 2-figure supplement 2, Figure 3-figure supplement 1B,
and supplementary file 1). Consistent with that, plotting of *crlf1a* expression onto the
UMAP showed high read counts in clusters 3 and 4, as well as in cluster 5 (Figure
3C). Again, DEG analysis showed that reactive MG can be subdivided further,
because only cluster 3, but not cluster 4 expressed Hairy-related downstream targets
of active Notch signalling including *her15.1-1*, *her9* and *her4.1* (Figure 2-figure
supplement 2, Figure 3-figure supplement 1B, and supplementary file 3). Indeed, the
UMAP plot of *her4.1* showed strong expression in cluster 3 as well as in cluster 5
(Figure 3C). To validate the expression of *her4.1* in the regenerating retina we used
in situ hybridization. In agreement with our scRNAseq dataset, *her4.1* signal was not
present in the uninjured retina, but appeared at 44 hpl in the inner nuclear layer of
the central retina (Figure 3D). At 4 and 6 dpl, *her4.1* transcripts remained in the inner
nuclear layer and appeared in cells of the outer nuclear layer too (Figure 3D). DEG
analysis revealed *starmaker* (*stm*) as a highly specific marker for reactive MG in
cluster 3, but not for reactive MG in cluster 4 (supplementary files 1 and 3). We used
fluorescent *in situ* hybridization to validate *stm* expression, combined with
immunohistochemistry against GFAP (located mainly to MG fibers) as well as the

proliferation marker PCNA in the uninjured as well as light-lesioned retina (Figure 3-

figure supplement 2). In agreement with our scRNAseq results, we observed *stm*

expression only at 44 hpl, but did not detect any *stm* expression in the uninjured as

well as in the 4 and 6 dpl retinal samples (Figure 3-figure supplement 2). Virtually all

stm-positive cells at 44 hpl displayed a MG morphology (Figure 3-figure supplement

2, yellow arrowhead) and were positive for Zrf1 (detecting GFAP) as well as PCNA.

We continued our validation by further analysing the expression of *crlf1a*, a marker of

reactive   MG,   with   fluorescent   *in   situ*   hybridization   combined   with

immunohistochemistry against GFAP and PCNA (Figure 4). Consistent with our

scRNAseq dataset, *crlf1a* was not detectable in the uninjured, central retina, but was

upregulated in Zrf1/PCNA-double positive cells at 44 hpl (Figure 4A, B). At 4 dpl,

crlf1a signal was still present in Zrf1/PCNA-double positive cells and returned back to

undetectable levels at 6 dpl (Figure 4A, B). Taken together, we observed cellular

heterogeneity in MG during regeneration, and identified two non-reactive and two

reactive MG populations that express distinct marker genes.

scRNAseq identifies a transient cell population with hybrid characteristics of

reactive MG and early RPCs in the light-lesioned retina

As mentioned above, several MG-related genes, like *gfap* or *crlf1a*, were also

expressed in cluster 5, which locates next to the reactive MG (clusters 3 and 4), is

highly proliferative and only becomes eminent at 4 dpl. Further analysis of this cluster

showed expression of several members of the High Mobility Group (Hmg) protein

family among the top 100 upregulated marker genes (supplementary file 1).

Consistent with that, the UMAP expression profile of *hmgb2b* showed presence of

high counts already in cluster 3 (reactive MG), as well as in other, further

downstream RPC clusters and regenerating progeny (Figure 5A). This indicated a

transition from MG to RPC fate in the course of regeneration and that cells of cluster

5 might have hybrid characteristics of reactive MG and downstream RPCs. Hence,

we conducted a DEG analysis of cluster 5 in comparison to reactive MG (clusters 3

and 4), as well as to the neighbouring, downstream RPCs (cluster 6). When

compared to reactive MG, cells of cluster 5 showed expression of several genes,

including *stmn1a*, *nr2e1* and *sox9b* (Figure 3-figure supplement 1C, supplementary

files 4 and 5). Consistently, UMAP expression profile of *stmn1a* showed counts

prevalently in cluster 5 and neighbouring, downstream RPCs on the right side of the

map, but not in the reactive or non-reactive MG on the left side (Figure 5A). The

UMAP profiles of *sox9b* and *nr2e1* showed strong expression almost exclusively

restricted to cluster 5. DEG analysis between cluster 5 and the neighbouring

downstream cluster 6 revealed also expression of *nr2e1* and *sox9b*, but additionally

transcripts like *id1* (Figure 3-figure supplement 1D and supplementary file 6).

Consistently, high *id1* counts were found in cluster 5 and in other reactive or non-

reactive MG clusters, but not in downstream RPCs on the right side of the map

(Figure 5A). In order to validate the RNAseq data, we analysed the expression of

hmgb2b and *id1* transcripts in the regenerating retina using *in situ* hybridization and

fluorescent *in situ* hybridization combined with immunohistochemistry, respectively.

Consistent with our scRNAseq data, *hmgb2b* signal was not detectable in the

uninjured, central retina, but was initiated in individual cells residing in the inner

nuclear layer at 44 hpl (Figure 5B). Subsequently, *hmgb2b* transcripts could be

detected in clusters of cells in the inner and outer nuclear layers at 4 and 6 dpl

(Figure 5B). The expression of *id1* in the regenerating tissue matched the scRNAseq

data. Expression of *id1* was already detectable in Zrf1-positive but PCNA-negative

cells in the inner nuclear layer of the uninjured, central retina (Figure 6A, B). At 44

hpl, *id1* was strongly upregulated and co-localized with Zrf1/PCNA-double positive

cells in the inner nuclear layer. At 4 and 6 dpl, *id1* expression was still associated

with Zrf1/PCNA-double positive cells and, occasionally, with PCNA-only positive cells

(Figure 6A, B). In summary, marker analysis reveals that cells in cluster 5 display a

transcriptional signature of reactive MG, as well as RPCs, which is why we assigned

it a hybrid identity.

scRNAseq identifies two neurogenic progenitor populations in the light-

lesioned retina

The before-mentioned DEG-analysis between cluster 5 (hybrid cells) and cluster 6

(neighbouring, downstream RPCs) also revealed that the latter upregulated *insm1a*

(Figure 3-figure supplement 1D and supplementary file 6), which has been described

to promote cell cycle exit of RPCs in the injured fish retina at 4 dpl (Ramachandran et

al., 2012). In addition, our DEG analysis showed upregulation of *atoh7* and *onecut2*

in cluster 6 (Figure 3-figure supplement 1D and supplementary file 6). We found that

the highly redundant gene *onecut1* was also expressed in cluster 6, albeit this was

not found in the DEG analysis. As expected, the UMAP expression profile of *insm1a*

confirmed high read counts in cluster 6 but also in clusters 7, 8 and 10 (Figure 7A).

The UMAP expression profile of *atoh7* also confirmed expression in cluster 6, as well

as in clusters 7 and 10 (Figure 7A). Moreover, the UMAP expression profile of

onecut1 and *onecut2* confirmed that both genes were expressed in clusters 6 and 7

and in addition in cluster 9 (Figure 7A). Cluster 7 expressed additional neurogenic

marker genes like *tubb5* and *sox11b* (Baraban et al., 2013; Lange et al., 2020; Mu et

al., 2016) (Figure 3-figure supplement 1D and supplementary file 1). Finally, the cell

cycle status and the UMAP expression of *mki67,* a mitotic marker, showed read

counts in cluster 6, but not in cluster 7 (Figure 7A). Hence, we concluded that cluster

6 and cluster 7 represented neurogenic progenitors, and were named progenitors 2

and progenitors 3, respectively. To validate *atoh7*, *onecut1* and *onecut2* expression

in the regenerating retina, we employed again *in situ* hybridization. In agreement with

our scRNAseq data, *atoh7* signal was absent in the uninjured and 44 hpl retina, but

became detectable in the inner and outer nuclear layers at 4 and 6 dpl (Figure 7B).

Similarly, *onecut1* and *onecut2* were prominently expressed in the inner and outer

nuclear layer at 4 and 6 dpl, in addition to their homeostatic expression in the

presumptive horizontal cell layer in the uninjured and 44 hpl retina (Figure 7B, black

arrowheads). In summary, scRNAseq showed heterogeneity of neurogenic

progenitors, which upregulate known markers of retinogenesis in the regenerating

retina.

scRNAseq identifies fate-restricted progenitors in the light-lesioned retina

Gene expression analysis of cluster 8, which locates next to the neurogenic

progenitors (cluster 6 and 7), showed significant upregulation of several well-known

photoreceptor precursor markers, including *crx*, and *thrb* (supplementary file 1).

Consistently, UMAP plots for *crx* and *thrb* revealed high counts in cluster 8 (Figure

8A). Moreover, the former was also highly expressed in clusters 11 and 12, whereas

the latter was restricted to the additional cluster 1. Subsequent further gene

expression analysis identified clusters 11 and 12 as differentiating cone

photoreceptors arising from cluster 8. Precisely, clusters 11 and 12 represent

differentiating photoreceptors giving rise to blue and red cones as shown by the

UMAP plots of *opn1sw2* and *opn1lw2*, respectively (Figure 8B). Cluster 9, which

abuts the neurogenic progenitor 3 (cluster 7), expressed known developmental

markers of horizontal cell precursors like *prox1a* and *rem1* (supplementary file 1).

Consistently, the UMAP plot for *rem1* showed high expression almost restricted to

this cluster specifically. Gene expression analysis also revealed *ptf1a* expression in

cluster 9, which has been shown to be required for lineage specification of horizontal

and amacrine cells in the developing retina (Jusuf et al., 2011; Jusuf and Harris,

2009). In line with these findings, the UMAP expression profile of *ptf1a* confirms high

counts in the left part of cluster 9, but is even more pronounced in the upstream part

of cluster 7 feeding into cluster 9 (Figure 8C). Gene expression analysis of the

remaining clusters 10, 13, 14 and 15 identified them as differentiating progeny giving

rise to retinal ganglion cells (cluster 10), amacrine cells (clusters 13 and 14), as well

as bipolar cells (cluster 15), respectively (supplementary file 1). In this context, UMAP

plots of *gap43* and the more mature retinal ganglion cell marker *isl2b* showed a

strong upregulation in cluster 10 (Figure 8D). UMAP plots of *stx1b* and *pvalb6*,

markers of differentiating and mature amacrine cells, revealed strong expression in

clusters 13 and 14, whereas UMAP plots of *vsx1* and *samsn1a*, known and novel

markers of bipolar cells, respectively, showed high counts in cluster 15 (Figure 8E,

F).

With this respect, it was recently shown that the regenerating adult zebrafish retina

recapitulates developmental fate specification programs (Lahne et al., 2021). Hence,

we plotted the fraction of cells that differentiate into a specific cell fate during the time

course of regeneration (Figure 8-figure supplement 1). We found that the earliest

committed progeny emerged at 4 dpl giving rise to retinal ganglion cells or to

photoreceptors, in particular red cones. At 6 dpl, the number of retinal ganglion cells

and red cones continued to increase. In addition, we detected a significant number of

blue cones and amacrine cells, as well as a small number of bipolar cells at this time

point. We did not detect any differentiating progeny giving rise to horizontal cells or

rods at this stage. To address this issue, we further investigated the expression of

more mature neuronal markers in our dataset (Figure 8-figure supplement 2).

Expression of *rho* and *gnat1*, two markers of rod photoreceptors, was very low and

sparse, and did not define any specific rod cluster (Figure 8-figure supplement 2A).

Expression of *gad1b*, which encodes for glutamate decarboxylase synthetizing the

neurotransmitter GABA, was mostly restricted to amacrine cells (clusters 13 and 14)

(Figure 8-figure supplement 2B). To address the expression of more mature markers

of amacrine and horizontal cells, we investigated *chata* (*choline O-acetyltransferase*

a, an amacrine cell-specific marker), as well as *calb2a* and *calb2b* (*calbindin 2a* and

calbindin 2b, markers of amacrine and horizontal cells) (Figure 8-figure supplement

2B, C). As expected, high counts of *chata* were specifically found in amacrine cells

(clusters 13 and 14). In particular, *chata* expression was detected only in cells

located at the tip of cluster 13 and in all cells of cluster 14. *calb2a* was expressed in

some reactive MG (cluster 3) and quite strongly in some amacrine cells (cluster 13).

Importantly, we did not detect any *calb2a* and *calb2b* in horizontal cell precursors

(cluster 9).

We also checked the expression of transcripts for variants of protein kinase C using

prkcab and *prkcbb* (*protein kinase alpha b*, *protein kinase beta b*) to identify mature

bipolar cells (Figure 8-figure supplement 2D). Counts for *prkcab* were sparsely found

in some non-reactive and reactive MG (clusters 1-4). *prkcbb*-positive cells were found

more frequently in some non-reactive and reactive MG (clusters 1-4), progenitor 1

(cluster 5), as well as in amacrine cells (clusters 13, 14). Importantly, we did not

detect *prkcab* or *prkcbb* in our bipolar cells. When plotting the expression of general

synaptic markers like *snap25a* and *snap25b*, we observed mainly transcripts in the

differentiating progeny (Figure 8-figure supplement 2E). Specifically, high counts of

snap25a were found in retinal ganglion and amacrine cells (clusters 10, 13 and 14).

High counts of *snap25b* were observed in the same clusters as well as in red and

blue cones (clusters 11 and 12) and the right part of the horizontal cell precursors

(cluster 9). Taken together, our scRNAseq data reveal the presence of fate-restricted

progenitors and the sequential appearance of differentiating progeny in the light-

lesioned, adult zebrafish retina. Moreover, the analysis of more mature markers

shows that the differentiation of retinal ganglion cells and red cones is preceding the

appearance of blue cones, amacrine and bipolar cells.

scRNAseq of MG and MG-derived cells of the regenerating zebrafish retina

corroborates and complements previously published scRNAseq data

In order to compare our newly obtained scRNAseq dataset of MG and MG-derived

cells of the regenerating zebrafish retina to a previously published dataset of the

light-lesioned retina (Hoang et al., 2020), we employed the ingestion method

(Scanpy, https://scanpy-tutorials.readthedocs.io/en/latest/integrating-data-using-

ingest.html) and mapped the clusters identified by Hoang and colleagues to our

clusters (Figure 9). While we applied a short-term lineage tracing strategy and only

sequenced the enriched population of FAC-sorted MG and MG-derived cells of the

regenerating zebrafish retina, Hoang and colleagues sequenced all retinal cells in the

light-lesioned retina. Consequently, comparison between the two datasets unravelled

similarities, but also significant differences due to the different experimental set-ups

(Figure 9A). Consistently, the cluster annotated as resting MG in Hoang et al.

mapped to clusters annotated as non-reactive MG 1 and 2 in our dataset (Figure 9B).

The cluster annotated as activated MG in Hoang et al. mapped to clusters annotated

as reactive MG 1 and 2, as well as to the cluster with hybrid identity of

MG/progenitors in our dataset. Interestingly, some cells annotated as activated MG in

Hoang et al. mapped also to neurogenic progenitor 2 and 3 clusters in our dataset

(Figure 9B). The cluster annotated as progenitors in Hoang et al. mapped to the

progenitor area in our dataset, which included neurogenic progenitors 2, 3 as well as

photoreceptor and horizontal cell precursors (Figure 9B). Finally, retinal ganglion

cells, cones, GABAergic amacrine cells and bipolar cells annotated in Hoang et al.

perfectly mapped to retinal ganglion cells, cone, amacrine and bipolar cells in our

dataset (Figure 9B). While we did not detect a mature horizontal cell cluster, Hoang

and colleagues annotated a horizontal cell cluster, which cells mapped to reactive

MG 2, MG/progenitors 1 and part of progenitors 3 in our dataset (Figure 9B).

Moreover, Hoang and colleagues annotated rod photoreceptors that mapped to

progenitors 3, photoreceptor precursors, red and blue cones, horizontal cell

precursors and bipolar cells in our dataset (Figure 9B). Finally, Hoang and colleagues

annotated additional cell clusters that did not map to any cluster in our dataset, and

included oligodendrocytes, pericytes, retinal pigmented epithelial cells as well as

vascular/endothelial cells (Figure 9B).

Next, we selected representative marker genes per cluster from our scRNAseq

dataset and checked their expression in the dataset by Hoang and colleagues

(Figure 9-figure supplement 1). The dot plot showing the expression of selected gene

candidates per cluster further corroborated the large overlap between clusters

annotated in the present study with those annotated in the study by Hoang and

colleagues.

DISCUSSION

In this study, we employed a short-term lineage tracing strategy, which allowed the

selective enrichment of MG and their respective progeny for up to six days post light

lesion. Subsequent scRNAseq resolved the transcriptome of individual MG, RPCs as well as regenerated progeny, and allowed to reconstruct their lineage relationship in the homeostatic and regenerating zebrafish retina (Figure 1). A previous study combined bulk RNA sequencing of sorted MG with scRNAseq of whole retinae from zebrafish upon NMDA as well as light lesions (Hoang et al., 2020). This work focussed specifically on the early reprogramming of MG and progenitor production, covering up to 36 hours post light lesion. The authors report that MG go through an activated, non-proliferative state characterized by upregulation of gliosis markers. However, neither did they find any evidence of MG heterogeneity in uninjured conditions nor did they provide any thorough characterization of MG-derived RPCs (Hoang et al., 2020). Additional RNA sequencing studies of the regenerating zebrafish retina have used microarray as well as RT-PCR and qRT-PCR to evaluate transcriptional changes at a population level, without providing any MG/RPC single cell resolution (Cameron et al., 2005; Craig et al., 2008; Kassen et al., 2007; Kramer et al., 2021; Morris et al., 2011; Ng Chi Kei et al., 2017; Qin et al., 2009). Previous immunohistochemistry and *in situ* hybridization experiments provided a broad description of MG-derived RPCs. Indeed, progenitors are reported as fusiform cells that proliferate and express markers of multipotency like Pax6 and Sox2, as well as some neurogenic markers like Atoh7, Ptf1a, Vsx1 and others (Bernardos et al., 2007; Fausett and Goldman, 2006; Fimbel et al., 2007; Gorsuch et al., 2017; Lahne et al., 2021; Ramachandran et al., 2012). In contrast to these previous results, our study is the first providing a comprehensive overview of MG, MG-derived cells and their lineage relationship up to 6 days post lesion in the zebrafish retina. Due to our selective enrichment of MG and their respective progeny, we gain an adequate single cell resolution to observe substantial heterogeneity in the molecular identity of MG in homeostatic as well as regenerating conditions. We also provide an in depth

description of MG as well as RPC molecular identity, cell fate decisions and

production of regenerated progeny that cover all key aspects during zebrafish retina

regeneration. Our strategy allowed us to define two main trajectories in the

regenerating zebrafish retina (Figure 10). The first trajectory (solid and dashed red

lines) comprises clusters 1 to 5 and constitutes a closed cycle that both maintains the

MG pool and generates the first-born neurogenic RPCs. The second trajectory (solid

black lines) involves clusters 6 to 15 and gives rise to all retinal neurons via the

progressive restriction in cell fate competence of neurogenic RPCs. We discuss the

two trajectories separately.

Müller glia self-renew and give rise to early, multipotent progenitors

The glial trajectory starts with two populations of non-reactive MG (clusters 1 and 2),

which express canonical glial markers like *gfap*, *apoeb* and additional genes like

glula, *rlbp1a* and *cahz* that ensure the housekeeping function of MG during

homeostasis (Figure 3A, Figure 2-figure supplement 2, supplementary file 1). The

majority of cells in the non-reactive MG clusters of our scRNAseq dataset are

quiescent based on the analysis of the cell cycle, and are frequently found in the

uninjured, but also in the lesioned retina at all sampled time points (Figure 2).

However, some non-reactive MG are in S or G2/M phases of the cell cycle, and

express *pcna*, albeit at a lower level as compared to the expression in injury-induced,

proliferating cell clusters (Figure 2C, D). This is consistent with the proposed model

that zebrafish MG proliferate occasionally also in the uninjured, central retina and

generate rod progenitors that feed the rod precursor pool at the base of the outer

nuclear layer (Bernardos et al., 2007; Stenkamp, 2007). Accordingly, we detect

PCNA-positive cells in the uninjured, central retina (Figure 1-figure supplement 1) as

well as a small fraction of cells that align in a trajectory from non-reactive MG on the

left through reactive MG in the middle, to photoreceptor precursors on the right side

of the UMAP in the uninjured retina (Figure 2B). However, some cells end up in the

amacrine as well as retinal ganglion cell clusters. This suggests that MG might have

the potential to generate not only rods, but also all other retinal neurons in the

homeostatic adult retina. Indeed, MG in the uninjured retina express, albeit at a low

level, multipotency markers like *pax6*, and share many marker genes with RPCs

(Bernardos et al., 2007; Blackshaw et al., 2004; Raymond et al., 2006). Still, MG and

injury-induced RPCs are considered distinct cell populations. Upon injury, MG divide

only once between 31 and 48 hpl, whereas MG-derived RPCs continue to divide,

reaching a peak of proliferation at 4-5 dpl (discussed in Fausett and Goldman, 2006;

Lahne et al., 2020; Lenkowski and Raymond, 2014; Thummel et al., 2008). We

isolated cells at 44 hpl, a time point when MG are about to undergo, or already

underwent, their asymmetric cell division (Bernardos et al., 2007; Lahne et al., 2015;

Lenkowski and Raymond, 2014; Nagashima et al., 2013) and identify cells in cluster

3 as early reactive MG. Consistent with that, we find that cells in cluster 3 do not only

express *gfap* and *apoeb*, but also *pcna* (Figure 2D, 3A). Moreover, the predicted cell

cycle status supports this notion. Whereas cells in the left part of cluster 3 are mostly

in G1, cells in G2/M phase are found in the right part, with cells in S phase

overlapping with both domains (Figure 2C). Cells in cluster 3 also upregulate

inflammation-related markers like *crlf1a*, as well as *mmp9* and *hbegfa* (Figure 3C, 4,

Figure 3-figure supplement 1B and supplementary file 1). The upregulation of

inflammatory genes makes MG similar to immune cells (Silva et al., 2020; Wan et al.,

2012; Zhao et al., 2014). In this context, our population of early reactive MG (cluster

3), identified mostly at 44 hpl, resembles the reactive MG pool found in the

regenerating zebrafish retina at 36 hpl (Hoang et al., 2020). However, our MG cluster

is proliferative, whereas the one discussed by the mentioned study is not. The

presence of a distinct pool of non-proliferative, reactive MG as early as 36 hpl in the

earlier study might be explained by the different time points of sample collection,

and/or by the lower number of cells sequenced in our experiment (11.690) in

comparison to the previous one (45.153) (Hoang et al., 2020). In summary, MG in the

homeostatic zebrafish retina behave like multipotent RPCs. Upon injury, reactive MG

express genes associated with inflammation, like *crlf1a*, and undergo asymmetric cell

division. Daughter cells derived from this cell division most likely map to cluster 5,

which is discussed below.

A crossroad in the glial trajectory: hybrid cells with MG and RPC

characteristics

The close similarity between zebrafish MG and MG-derived RPCs at the level of

expressed markers as well as cellular and subcellular morphology has been

described extensively in the literature (Fausett and Goldman, 2006; Lahne et al.,

2020; Lenkowski and Raymond, 2014). Thus, we suggest that cells in cluster 5,

which displays a transcriptional hybrid identity between MG and RPCs, are the

product of the asymmetric cell division from the early reactive MG (cluster 3). Cells in

cluster 5 are still *gfap* positive, and share marker expression with early reactive MG,

like *her4.1* and *crlf1a* (Figure 3A, C). However, cluster 5 is highly proliferative and

becomes only prominent at 4 dpl, a time point when predominantly RPCs are

proliferating (Bernardos et al., 2007; Fausett and Goldman, 2006; Lenkowski and

Raymond, 2014; Nagashima et al., 2013). Consistently, *pcna* is highly expressed in

cluster 5 and the predicted cell cycle status indicates that most cells are in S or G2/M

phase (Figure 2C, D). Moreover, previous studies have found *her4.1* and *crlf1a* in

MG-derived RPCs in the light- as well as stab-lesioned retina, where they control

RPC proliferation at 4 dpl (Wan et al., 2012; Wilson et al., 2016; Zhao et al., 2014).

We found that *her4.1* at 4 and 6 dpl locates to the inner nuclear layer as well as to

the outer nuclear layer, where presumptive RPCs reside at this time point (Figure 3D)

(Campbell et al., 2022; Wilson et al., 2016). In addition, cells in cluster 5 upregulate

sox9b and *nr2e1*, which have been found in early, multipotent RPCs of the

developing fish, avian and mammalian retinae, as well as in late retinogenesis, where

they regulate MG genesis (Buenaventura et al., 2018; Corso-Díaz and Simpson,

2015; Kitambi and Hauptmann, 2007; Muto et al., 2009; Poché et al., 2008). Hence,

we propose that cells in cluster 5 represent a crossroad of the glial and neurogenic

trajectory, containing self-renewed MG as well as first-born, MG-derived RPCs as a

direct result of the earlier (44 hpl) asymmetric MG cell division. We further propose

that self-renewed MG close the cycle via late reactive MG (cluster 4), which is

evident at 4 dpl, and return into non-reactive MG (clusters 1 and 2, red, dashed line

in Figure 10). In this respect, we note that reactive MG 2 (cluster 4) share many

marker genes with early reactive MG 1 in cluster 3 (Figure 2-figure supplement 2,

supplementary files 1 and 3). Both reactive MG cell populations express marker

genes that are involved in inflammation, like *crlf1a* (Figure 3C, supplementary file 1)

encoding for an inflammatory cytokine (Zhao et al., 2014). However, proliferation

markers like *pcna* are expressed only in reactive MG in cluster 3 but not in cluster 4

(Figure 3-figure supplement 1). We corroborated *crlf1a* expression in the

regenerating tissue using fluorescent *in situ* hybridization combined with

immunohistochemistry for PCNA and GFAP (Figure 4). The staining confirmed co-

localization of the transcript with PCNA-, GFAP-double positive cells at 44 hpl as well

as at 4 dpl. The co-localization of *crlf1a* with PCNA at 4 dpl does not seem to match

the sequencing data for reactive MG 2, which is eminent at 4 dpl and should not be

proliferative. However, it is possible that *crlf1a*/PCNA-double positive cells detected

at 4 dpl are still positive for the PCNA protein, but no longer express the *pcna*

transcript. Double *in situ* hybridization for *pcna* and *crlf1a* would be needed to fully

address whether *crlf1a*-positive cells are still *pcna*-positive at 4 dpl. It is also possible

that *crlf1a*/GFAP-double positive but PCNA-negative Müller glia are fewer and only

masked in the crowd of *crlf1a*-, PCNA-double positive, GFAP-negative progenitors at

4 dpl (Raymond et al., 2006). Nevertheless, we would like to mention that we cannot

exclude the possibility that non-proliferative, reactive MG in cluster 4 represent MG

that are reactive to the tissue injury, but never re-enter the cell cycle upon light-

lesion.

In contrast to MG, MG-derived RPCs embark into the neurogenic pathway (cluster 6,

Figure 10, solid, black line). In an alternative model, cells in cluster 5 might represent

only MG cells that have upregulated early RPC markers like *sox9b* and *nr2e1* and

are dividing a second time at 4 dpl to generate renewed MG (cluster 4) and

neurogenic RPCs (cluster 6). However, MG have been described to divide only once

between 31 and 48 hpl and additional and/or asynchronous MG divisions later than 2

dpl have not been investigated (Lahne et al., 2015; Lenkowski and Raymond, 2014;

Nagashima et al., 2013; Thummel et al.,2008). BrdU/EdU lineage tracing

experiments will be required to address this issue, with the caveat that MG and RPCs

are hardly distinguishable from each other in the 4 dpl retina, especially when using

the Zrf1 antibody to label MG (Fausett and Goldman, 2006; Raymond et al., 2006).

Live imaging of the regenerating retina would be the best tool to follow MG and MG-

derived progenitors *in vivo*, but the technique is not optimized yet for long recordings

from the intact tissue in the living animal (Lahne et al., 2015). Either way, our

proposed model of self-renewing MG is further supported by the change of cell

density over time (Figure 2B). Following the cells across the four time points on the

UMAP, we observe that initially most cells of the uninjured sample map to non-

reactive MG (clusters 1 and 2) located at the far left. At 44 hpl, the cell density shifts

towards the upper right portion of cluster 1 and towards the early reactive MG (cluster

3) in the upper, middle of the map. At 4 dpl, sampled cells mostly map to RPCs

(clusters 5-9) on the right, as well as to late reactive MG (cluster 4) in the lower,

middle part. Additionally, cells mapping to the lower right part of non-reactive MG

(clusters 1 and 2) are enriched in this sample, indicating a reverse direction from right

to left. This population is significantly further increased at 6 dpl, when the number of

RPCs is less prominent and the differentiating neural types at the top and bottom

right are predominantly present in the sample. According to this model, sample

collection at even later time points, e.g. 10 or 14 dpl, might show that the initial status

has been restored and that non-reactive MG are mostly enriched. However, it is

unlikely that this experiment could be performed with our short-term labelling

strategy.

The neurogenic trajectory produces regenerated progeny that partially follow

the developmental order of neurogenesis

Neurogenic RPCs map to the right of cluster 5, and include proliferating, neurogenic

RPCs (cluster 6) and the downstream, non-proliferating, neurogenic RPCs (cluster

7). Cells in these clusters express *insm1a*, *atoh7* and additional neurogenic genes

that are required for the generation of several neuronal lineages (Figure 7, Figure 3-

figure supplement 1D and supplementary files 1 and 6). Indeed, *atoh7* expression

has been described to be initiated in RPCs on their way to the last cell division giving

rise to a retinal ganglion cell and another cell that can be either an amacrine cell, or a

precursor of a horizontal or of a cone cell in the developing zebrafish retina (Amini et

al., 2018; Nerli et al., 2022; Poggi et al., 2005). Cluster 6 and 7 appear also highly

similar to the transient, neurogenic RPC state that has been described recently in the

developing mouse retina (Wu et al., 2021). We propose that cluster 6 and 7 are the

roots of the neurogenic trajectory from which fate-restricted RPCs and regenerated

progeny arise at 4 and 6 dpl (solid, black line in Figure 10). Cells in cluster 6 are

mki67-positive, but additionally express *insm1a*, which promotes RPC cell cycle exit

at 4 dpl in the injured retina (Ramachandran et al., 2012). Cells in cluster 6 might

undergo their last cell division, as suggested by the concomitant expression of *atoh7*.

Consistently, in our scRNAseq high counts of *atoh7* are found in cells of cluster 6, to

cluster 7 and eventually cluster 10. Cluster 10 represents differentiating retinal

ganglion cells that are the first regenerated progeny, evident as early as 4 dpl (Figure

7, 10 and Figure 8- figure supplement 1).

Clusters 14 and 15 highly express amacrine cell genes (Figure 8- figure supplement

2B). Interestingly, our fluorescent-reporter sorting strategy allows the discrimination

of distinct cell subsets belonging to the same cell type. In this respect, cluster 14

includes GABAergic amacrine cells, as shown by the expression of *gad1b*, which

encodes for the GABA-synthesizing enzyme GAD67 (Bosma et al., 1999). In

contrast, cluster 15 includes a class of amacrine cells known as starburst amacrine

cells, as indicated by the expression of both *gad1b* and *chata* (O'Malley et al., 1992).

While amacrine cells arise directly from the core of neurogenic RPCs, fate restricted

precursors for regenerated cones and for horizontal cells are identified (Figure 8 and

10), as it occurs during development (Godinho et al., 2007; Weber et al., 2014).

Consistently with the events during development, red cones (cluster 11) branch from

thrb-positive cells of the photoreceptor RPC cluster (cluster 8). Interestingly, *thrb*

transcripts are found in neurogenic RPCs of cluster 7 as well as at the beginning of

cluster 9, which contains horizontal cell precursors becoming evident at 4 dpl (Figure

8B). This is in agreement with observations in the developing zebrafish retina, where

horizontal and cone cells share a common Thrb-positive progenitor (Suzuki et al.,

2013). Mature markers of horizontal cells, like calbindins, are not found in cluster 9

(Figure 8B, Figure 8-figure supplement 2C and supplementary file 1), in contrast to

cone photoreceptor mature markers, like red light- and blue light-sensitive opsins, at

4 and 6 dpl. We did not detect UV-sensitive opsin marker gene expression labelling

UV cones, which might arise later than 6 dpl and were hence not included in our

sorted samples. While it might be that mature horizontal cells arise later than 6 dpl,

their absence from our dataset could also be due to our experimental set-up. As

explained before, we isolated mCherry-only positive and EGFP/mCherry-double

positive cells to obtain MG and MG-derived cells. Whereas mCherry, driven by a *gfap*

promoter fragment, is actively transcribed in MG, the label of MG-derived cells is

completely dependent on the half-life of the fluorophore. Protein turnover and further

cell divisions would result in a constant depletion of mCherry in MG-derived cells over

time. In this context, we find a strong expression of *pcna* in cluster 9, indicating that

additional cell divisions might have diluted the mCherry label to undetectable levels

(Figure 2D). Inclusion of EGFP-positive only cells in our sample collection may reveal

the pool of mature horizontal cells, which inherit the fluorophore from their

pcna:EGFP-positive precursors. Accordingly, we find the presence of horizontal cell

precursor markers in our dataset. For instance, cells in cluster 9 express *onecut1* and

onecut2, which encode transcription factors and are upregulated in horizontal cell

RPCs in the developing mouse retina, and which regulate horizontal cell

maintenance in adult animals (Emerson et al., 2013; Klimova et al., 2015; Sapkota et

al., 2014). Counts for *onecut1* and *onecut2* are found also in earlier neurogenic

RPCs of clusters 6 and 7, where they most likely contribute to the transient state of

RPCs together with additional neurogenic markers, like *atoh7* (Wu et al., 2021). This

is again consistent with observations in the developing mouse retina, where Onecut1

and Onecut2 regulate the development of retinal ganglion cells, as well as the

development of cones, horizontal cells, and of a subset of amacrine cells, while

inhibiting rod specification genes (Emerson et al., 2013; Sapkota et al., 2014).

In addition to mature horizontal cells, we were also unable to detect any rod-specific

cluster expressing mature rod markers like *rho* and *gnat1* (Figure 8-figure

supplement 2A). We assume that mature rods might arise as a third branch from the

photoreceptor RPC cluster (Figure 10), which indeed expresses rod specification

markers like *nr2e3* (supplementary file 1). We speculate that rods either become

evident only later than 6 dpl, or that they do not survive the isolation procedure, since

rods possess a complex morphology with long outer segments. A third possibility

might be that rod photoreceptors are replenished by their dedicated rod precursors

faster than the triggering of the MG-dependent injury response. In this context, rod

precursors form a pool of cells at the base of the outer nuclear layer and proliferate

as early as 16 hours post-light lesion (Morris et al., 2005; Stenkamp, 2011; Thummel

et al., 2008). Since the already available pool of rod precursors would be

gfap:mCherry-negative, we might have missed the derived rod progeny, given our

sorting strategy. In summary, we show that in the light-lesioned retina, zebrafish MG

go through an inflammatory-like state before generating hybrid MG/RPCs that

eventually cycle-back to non-reactive MG as well generate neurogenic RPCs. The

neurogenic RPCs progressively restrict their competence to produce neurons that

include retinal ganglion cells first and bipolar cell last up to 6 dpl. A previously

published study used scRNAseq to profile the transcriptome of cells in the zebrafish

retina upon several lesion paradigms, including the light injury, up to 72 hpl (Hoang et

al., 2020). In their study, the authors dissociated uninjured and lesioned retinae to

single cells that were all subsequently subjected to scRNAseq. Conversely, in our

study we used a short-term lineage tracing strategy to enrich for and sequence specifically MG and MG-derived cells in uninjured and light-lesioned retinae up to 6 dpl. Bioinformatic comparison between the two datasets revealed a high degree of overlap between the respective cell clusters, as it would be expected when the same tissue is analysed. Specifically, Hoang and colleagues annotated resting MG and activated MG, as well as progenitor clusters that all map to non-reactive MG, reactive MG and progenitor clusters in our scRNAseq dataset, respectively (Figure 9 and Figure 9-figure supplement 1). In contrast to Hoang and colleagues, we were able to capture a certain degree of heterogeneity in MG and progenitor populations, and thus identify two clusters of non-reactive MG, two clusters of reactive MG and five clusters of progenitors and neuronal committed precursors. Consistently with the heterogeneity of MG described in our work, a recent study found MG heterogeneity in both quiescent and lesioned zebrafish retina using scRNAseq as well (Krylov et al., 2023). Specifically, this study reports six clusters of quiescent Müller glia that display differential spatial distribution along the dorsal/ventral retinal axis. The authors describe a ventral, quiescent Müller glia population that shares some marker genes (*aldh1a3*, *rdh10a*, *smoc1*) with our non-reactive Müller glia 2 (cluster 2, supplementary files 1 and 2). Furthermore, Krylov et al., 2023 show that Müller glia that locate to different spatial position show differential activation and proliferation response upon photoreceptor ablation (Krylov et al., 2023). Due to the half-life of mCherry and EGFP, we were also able to sample MG-derived cells to capture large parts of the regenerated neurons and birthdate their order specifically. This is in contrast to Hoang and colleagues, who on the one hand sequenced all retinal cells, independently of their contribution to regeneration, and only used samples up to 72 hpl, thus lacking sufficient MG lineage resolution to detect lesion-specific, regenerated progeny (Hoang et al., 2020). While they were able to annotate clusters

for almost all mature retinal cell types, the applied scRNAseq strategy is unable to distinguish between neurons that were already present in the retina and those that were regenerated specifically upon lesion. Given that we only sequenced MG and MG-derived cells, this could explain why Hoang and colleagues detected mature horizontal and rod cells, while we could not. Interestingly, and despite this discrepancy, horizontal cells detected by Hoang and colleagues share some marker genes with the horizontal cell precursors we annotated in our dataset (Figure 9-figure supplement 1). Likewise, rod photoreceptors from the Hoang dataset map largely, although not exclusively, to photoreceptor precursors in our dataset (Figure 9 and Figure 9-figure supplement 1). Importantly, there is a consistent overlap between retinal ganglion, cone, amacrine and bipolar cells from Hoang and colleagues' dataset and the corresponding cell clusters in our dataset, which further strengthens our results.

In summary, our study is the first comprehensive description of the molecular network underlying MG self-renewal and RPC differentiation to regenerating neurons which will inform future regenerative therapies to treat retinal dystrophies in humans.(Javed and Cayouette, 2017).

MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
antibody	Zrf1 (Mouse, monoclonal, IgG1)	ZIRC	RRID:AB_10013806	IHC, 1:250

antibody	anti-mCherry (Rabbit, polyclonal)	Takara Bio	Catalog #632475; RRID:AB_2737298	IHC, 1:500
antibody	anti-PCNA (Mouse, monoclonal, IgG2a)	Dako	RRID:AB_2160651	IHC, 1:500
antibody	anti-GFP (Chicken, polyclonal)	Abcam	RRID:AB_300798	IHC, 1:2000
antibody	Sheep anti-Digoxigenin-AP Fab fragments	Roche	Catalog #11093274910; RRID: AB_514497	ISH, 1:2000
antibody	Sheep anti-Digoxigenin-POD Fab fragments	Roche	Catalog #11207733910; RRID:AB_514500	ISH, 1:2000
antibody	Alexa Fluor® 633 Goat anti-Mouse IgG1	Thermo Fisher Scientific	Catalog #A21126; RRID: AB_2535768	IHC, 1:500
antibody	Alexa Fluor® 555 Goat anti-Rabbit	Thermo Fisher Scientific	Catalog #A21429; RRID: AB_2535850	IHC, 1:500
antibody	Alexa Fluor® 488 Goat anti-mouse IgG2a	Thermo Fisher Scientific	Catalog #A21131; RRID: AB_2535771	IHC, 1:500
chemical compound, drug	NBT/BCIP stock solution	Roche	Catalog #11681451001	ISH, 20 µl/ml
chemical compound, drug	Blocking reagent	Roche	Catalog # 11096176001	ISH

chemical compound, drug	Calcein blue, AM	Invitrogen	Catalog #C1429	Living cell dye, 1 µM
chemical compound, drug	Neutral protease (Dispase®)	Worthington Biochemical Corporation	Catalog #LS02100	0.2 U/ml
commercial assay or kit	Papain dissociation System	Worthington Biochemical Corporation	Catalog #LK003150	
commercial assay or kit	10x Chromium single - cell kit	10x Genomics		V3 Chemistry
commercial assay or kit	Total RNA purification kit	Norgen Biotek Coporation	Catalog #17200	
commercial assay or kit	cDNA synthesis	Roche	Catalog #4379012001	
commercial assay or kit	TOPO TA® cloning dual promoter	Thermo Fisher Scientific	Catalog #450640	
commercial assay or kit	Zymoclean Gel DNA Recovery Kit	Zymo Research	Catalog #D4002	
commercial assay or kit	Gene Jet Plasmid Miniprep kit	Thermo Fisher Scientific	Catalog #k0503	
commercial assay or kit	Nucleo Bond® Xtra	Macherey-Nagel	Catalog #740410.5	

	Midi			
commercial assay or kit	TSA™ Plus Cyanine 3 System,	AKOYA Biosciences®	Catalog #NEL744A001KT	
other	20x SSC buffer			175.3 g NaCl, 88.2 g HOC(COONa)(CH2COONa)2 · 2H2O, adjust pH to 6 with acetic acid in DEPC water
other	Washing buffer			50% deionized formamide, 1x SSC in DEPC water
other	MABT buffer			100 mM maleic acid, 150 mM NaCl, adjust pH to 7.5 with NaOH, 0.1% Tween 20, in DEPC water sterile filter
other	Staining buffer for Alkaline Phosphatase			100 mM NaCl, 50 mM MgCl2, 100 mM Tris base, pH = 9.5, 0.1% Tween 20 in DEPC water
other	TNT buffer			100 mM Tris HCl, pH = 7.5, 150 mM NaCl, 0.1% Tween 20 in DEPC water

Animals

Zebrafish (*Danio rerio*) were raised as previously described (Brand et al., 2002).

Adult fish (6-12 months old) of either sex were used for all experiments. WT animals

were in the AB genetic background.

Transgenic lines

To generate the *Tg(pcna:EGFP)* (tud119), the BAC CH73-140L11, containing more

than 100 kb of the genomic *pcna* locus, was obtained from the BACPAC resource

centre. A cassette containing EGFP and a polyadenylation signal was recombined at

the starting ATG of exon 1. First, FRTgb- neo-FRT and EGFP cassettes where

amplified by PCR. Then, the cassettes where amplified and merged by fusion PCR

with primers carrying 50 nucleotides of homology of the targeting sequence.

Recombineering of the BAC was performed as previously described (Zhang et al.,

1998). BAC DNA was purified and about 50 pg of BAC DNA were injected into

fertilized eggs at the one cell stage. F0 were raised and incrossed. F1 were identified

by visual screening for GFP expression and raised. The *Tg(gfap:nls-mCherry)* line

was generated as previously described (Lange et al., 2020). In addition, to the line

described in Lange et al. (tud117tg), a second independent insertion was isolated

(tud120). This insertion displays a stronger expression in MG and was hence used

for this study.

Light injury

WT AB and *Tg(pcna:EGFP);Tg(gfap:nls-mCherry)* fish were dark adapted for 3 days.

Afterwards, diffuse light lesion was performed as previously described (Weber et al.,

2013). Briefly, up to 4 fish were placed in a beaker filled with system water (300 ml),

positioned 3 cm away from the light bulb of a metal halide lamp (EXFO X-Cite 120W,

EXFO Photonic Solutions, Mississauga, Ontario, Canada). Fish were exposed for 30

min to a bright light (≥ 100.000 lux), and then put back to system tanks in normal light

conditions. Uninjured fish controls were dark adapted and processed in parallel to

light injured animals.

Dissociation of retinae to a single cell suspension

Uninjured and light-lesioned *Tg(pcna:EGFP);Tg(gfap:nls-mCherry)* fish were exposed

to an overdose (0.2%) of MS-222 (Sigma) (Brand et al., 2002) until cessation of the

operculum movement. Subsequently, each cornea was poked with a needle, and

tweezers were gently pressured on each side of the eye to remove the lens. Then,

tweezers were used to carefully isolate the retinae, which were put in a dissociation

solution of papain (16 U/ml), dispase (0.2 U/ml) and DNAse (168 U/ml) dissolved in

Earle's Balanced Salt Solution (EBSS, Worthington Biochemical Corporation). Up to

6 retinae were put in 1.7 ml of dissociation solution. Retinae were digested for 30

minutes at 28°C, and then calcein blue (final concentration: 1µM, Invitrogen) was

added for 10 minutes to the digestion solution at 28°C to stain for living cells. At the

end of the enzymatic digestion, mechanical trituration was applied using fire-polished

glass pipettes of decreasing tip diameter to help tissue dissociation. Single cells were

centrifuged (300 g) at 4°C, for 6 minutes. The cell pellet was re-suspended in EBSS

supplemented with DNAse (100 U/ml) to eliminate residual, ambient DNA, and

albumin ovomucoid inhibitor (1:10 of stock solution, Worthington Biochemical

Corporation) to stop papain action. The single cell solution was eventually filtered (20

µm filters) to remove clumps and debris, and put on ice, ready for Fluorescent

Activated Cell Sorting (FACS). Each single cell suspension was obtained by pooling

8 retinae (from 4 fish, 2 males and 2 females) of uninjured and light-lesioned

Tg(pcna:EGFP);Tg(gfap:mCherry) animals. Uninjured WT AB, *Tg(gfap:mCherry)* and

Tg(pcna:EGFP) fish were used to prepare single cell suspensions of dissected

retinae for setting the single fluorophore gates in flow cytometry. Additionally, a

calcein only stained single cell suspension was prepared from uninjured WT AB fish

retinae for the setting of the calcein gate in the flow cytometry.

Flow cytometry and FACS

A BD FACSAria™ III cell sorter was used for flow cytometry and FACS. mCherry fluorophore was detected using a 561 nm excitation laser and a 610/20 nm band pass filter, EGFP using a 488 nm laser and a 530/30 nm band pass filter and calcein blue using a 405 nm laser and a 450/40 nm band pass filter. The gating strategy is shown in Figure 1-supplemental figure 3 for uninjured samples and was the same for all the light-lesioned samples. Forward and side scatters were used to gate cells among all the events in the single cell suspension and to gate singlets among those cells (Figure 1-figure supplement 3A). mCherry-positive only as well as EGFP-positive only gates were set in comparison to the unstained, WT control (Figure1-figure supplement 3B). Calcein blue positive, single cells were eventually gated as living cells (60.000 living cells from uninjured, 44 hpl and 4 dpl samples, 41.389 living cells from 6 dpl sample). 15.000 cells were sorted from each samples, put on ice and sent to the CMCB Deep Sequencing Facility for single cell RNA sequencing (scRNAseq).

(Droplet based) single cell transcriptomics

Single cell transcriptome sequencing was performed based on the 10x Genomics Single cell transcriptome workflow (Zheng et al., 2017). Specifically, 15.000 FACS-sorted cells were recovered in BSA-coated tubes containing 5 ul PBS with 0.04% BSA. Cell samples were carefully mixed with Reverse Transcription Reagent (Chromium Single Cell 3' Library Kit v3) and loaded onto a Chromium Single Cell B Chip to reach a recovery of 10.000 cells per sample. The samples were processed further following the guidelines of the 10x Genomics user manual for single cell 3' RNA-seq v3. In short, the droplets were directly subjected to reverse transcription,

the emulsion was broken and cDNA was purified using silane beads (Chromium

Single Cell 3' Gel Bead Kit v2). After amplification of cDNA with 11 cycles, samples

were purified with 0.6x volume of SPRI select beads to deplete DNA fragments

smaller than 400 base pairs and cDNA quality was monitored using the Agilent

FragmentAnalyzer 5200 (NGS Fragment Kit). 10 ul of the resulting cDNA were used

to prepare single cell RNA seq libraries - involving fragmentation, dA-Tailing, adapter

ligation and 11 cycles of indexing PCR following manufacturer's guidelines. After

quantification, both libraries were sequenced on an Illumina NextSeq500 in 75 base

pair paired-end mode, generating 45-109 million fragments per transcriptome library.

The raw sequencing data was processed with the 'count' command of the Cell

Ranger software (v2.1.0) provided by 10X Genomics. To build the reference, the

zebrafish genome (GRCz11) as well as gene annotation (Ensembl 90) were

downloaded from Ensembl and the annotation was filtered with the 'mkgtf' command

of Cell Ranger (options: '—e attribute=gene_biotype:protein_coding --

attribute=gene_biotype:lincRNA –attribute=gene_biotype:antisense'). Genome

sequence and filtered annotation were then used as input to the 'mkref' command of

Cell Ranger to build the appropriate Cell Ranger.

Bioinformatic analysis

The complete software stack for the analysis is available as a Singularity container

(https://gitlab.hrz.tu-chemnitz.de/dcgc-bfx/singularity/singularity-single-cell, tag

1f5da0b). The count matrices generated with Cell Ranger were further analysed

using scanpy 1.8.2

(https://genomebiology.biomedcentral.com/articles/10.1186/s13059-017-1382-0). For

quality control, we removed cells with less than 2.000 counts, less than 500 detected

genes, more than 15% mitochondrial reads or more than 60% of the counts in the top

50 highest expressed genes. The counts were normalised with the size factors

computed with calculateSumFactors from scran 1.18.5 (min.mean=0.1)(Lun et al.,

2016). The normalised counts were log-transformed using the numpy function log1p.

The top 4000 highly variable genes were detected using the scanpy function

scanpy.pp.highly_variable_genes (n_top_genes= 4000). Cell cycle scores were

computed using scanpy.tl.score_genes_cell_cycle on each sample separately. As

genes associated to S-phase, the genes *mcm5*, *pcna*, *tyms*, *fen1*, *mcm2*, *mcm4*,

rrm1, *unga*, *gins2*, *mcm6*, *cdca7a*, *dtl*, *prim1*, *uhrf1*, *si:dkey-185e18.7*, *hells*, *rfc2*,

rpa2, *nasp*, *rad51ap1*, *gmnn*, *wdr76*, *slbp*, *ccne2*, *ubr7*, *pold3*, *msh2*, *atad2*, *rad51*,

rrm2, *cdc45*, *cdc6*, *exo1*, *tipin*, *dscc1*, *blm*, *casp8ap2*, *usp1*, *pola1*, *chaf1b*, *brip1* and

e2f8 were used. As genes associated to G2/M phase, the genes *hmgb2a*, *cdk1*,

nusap1, *ube2c*, *birc5a*, *tpx2*, *top2a*, *ndc80*, *cks2*, *nuf2*, *cks1b*, *mki67*, *tmpoa*, *cenpf*,

tacc3, *smc4*, *ccnb2*, *ckap2l*, *aurkb*, *bub1*, *kif11*, *anp32e*, *tubb4b*, *gtse1*, *kif20ba*,

si:ch211-69g19.2, *jpt1a*, *cdc20*, *ttk*, *kif2c*, *rangap1a*, *ncapd2*, *dlgap5*, *si:ch211-*

244o22.2, *cdca8*, *ect2*, *kif23*, *hmmr*, *aurka*, *anln*, *lbr*, *ckap5*, *cenpe*, *ctcf*, *nek2*, *g2e3*,

gas2l3, *cbx5* and *selenoh* were used. A principal component analysis (PCA) was

performed on the highly variable genes (function scanpy.pp.pca using

svd_solver='arpack'). A k-nearest-neighbour (knn) graph was computed using

scanpy.pp.neighbors and a UMAP was computed using scanpy.tl.umap. Clustering

was performed with scnapy.tl.leiden (resolution= 0.01). For the three resulting

clusters, marker genes were computed using scanpy.tl.rank_genes_groups. Two

smaller clusters identified as microglia and low-quality cells were removed from the

downstream analysis. Detection of highly variable genes PCA were performed as

above on the remaining cells. A new knn graph was computed (this time with

n_neighbors= 50) and a new UMAP was computed. Embedding densities were

computed using scanpy.tl.embedding_density. Clustering was performed in multiple

steps, always using scanpy.tl.leiden: First, the data were clustered with the resolution

parameter set to 0.45. Next, cluster 10 was subclustered with resolution 0.1 and

cluster 0 was subclustered with resolution 0.2. Marker genes of the resulting 15

clusters were computed with scanpy.tl.rank_genes_groups. Differential expression

analysis of pairs of clusters was performed by subsetting the data to cells from those

two clusters and computing differentially expressed genes with

scanpy.tl.rank_genes_groups. The source code for the analysis is available on

GitHub (https://github.com/fbnrst/celotto_2023).

Cluster annotation

Cluster annotation was performed in several steps. Initially, we identified clusters by

considering the top 100 upregulated genes per cluster, the time point at which they

appeared and their cell cycle profile. Genes were manually annotated by consulting

the ZFIN database (http://zfin.org/) and Pubmed (https://pubmed.ncbi.nlm.nih.gov/)

indexed literature about retina development and regeneration. In particular, we took

as a reference the single cell RNA sequencing dataset of the regenerating zebrafish

retina from Hoang et al., 2020, for marker genes lists. Following the initial

identification of cell clusters, we proceeded with a finer characterization by

considering differential gene expression analysis for clusters that shared a similar

identity. Occasionally (e.g., in the case of horizontal cell precursor marker *ptf1a*), we

plotted the UMAP profile of known candidates from retina development or

regeneration. Selected marker genes were then checked for expression in the retina

tissue via *in situ* hybridization, fluorescent *in situ* hybridization and

immunohistochemistry on cryosections.

Comparison with Light Damage data in Hoang et al., 2020

The count matrix, for the light damage single-cell data was downloaded from https://github.com/jiewwwang/Single-cell-retinal-regeneration (Hoang et al., 2020). The data was used as a reference data for Scanpy's ingestion method (https://genomebiology.biomedcentral.com/articles/10.1186/s13059-017-1382-0). The light-damage data was normalized with scanpy's pp.normalize_total and then log-transformed pp.log1p. This log-normalised count-matrix was used to create the dotplot in Figure 9-figure supplement 1. Next, our data and the light-damage data were subsetted to the set of genes that was present for both count matrices. 3.000 highly variable genes were computed using pp.highly_variable_genes, and then 50 principal components were computed using pp.pca. Then, the cell types from the light-damage reference data were ingested onto our data using tl.ingest.

Tissue preparation and cutting

Fish heads were fixed with 4% paraformaldehyde in 0.1 M phosphate buffer (PB, pH: 7.4) overnight, at 4°C. Subsequently, fixed heads where decalcified and cryoprotected in 20% EDTA and 20% sucrose in 0.1M PB (pH: 7.5) overnight, at 4°C. Heads were eventually embedded in 7.5% gelatin, 20% sucrose in 0.1M PB and sectioned (12 or 14 μm) at the cryostat (CryoStar Nx70, Thermo Scientific). Cryosections were stored at -20°C.

in situ **hybridization probe synthesis**

For the synthesis of *in situ* hybridization RNA probes, 4 dpl retinae were dissected and sonicated at 4°C to homogenate the tissue and to extract total RNA. We chose 4

dpl samples, because transcripts of interest were enriched at 4 dpl. RNA was extracted using the total RNA purification kit and following manufacturer's instructions (Norgen Biotek Corporation). cDNA was then transcribed using the first strand cDNA synthesis kit (Roche), followed by PCR to amplify the fragments of interest using primer pairs listed in the key resource table. The amplicon was subsequently purified (gel and DNA recovery kit, Zymo Research) and cloned in a TOPO TA® vector (Thermo Fisher), with which competent *E coli* bacteria were transformed and cultured on Petri dishes coated with agar and ampicillin, overnight at 37°C. The next day, colonies were picked up and cultured in LB medium supplemented with ampicillin (1:2000) overnight at 37°C, shaking, to enrich cultures for mini-prep (Gene Jet Plasmid Miniprep kit k0503, Thermo Fisher). Purified plasmids were sent for sequencing (Eurofins) to verify the sequence. Insert-positive colonies were further enriched in LB/ampicillin cultures for midi-prep (Nucleo Bond® Xtra Midi, Macherey – Nagel) and purified plasmids were linearized to transcribe anti-sense, digoxigenin labelled (DIG) RNA probes, which were stored in hybridization buffer at -20°C. Probes for *her4.1* (Takke et al., 1999), *stm* (Söllner et al., 2003), *crlf1a*, *hmgb2b*, *atoh7*, *onecut1* (Matthews et al., 2004), *onecut2* were transcribed from a TOPO TA® cloned insert. Instead, the *id1* probe was transcribed directly from the PCR – amplified cDNA using SP6-conjugated forward and T7-conjugated reverse primers (key resource table).

in situ hybridization

in situ hybridization and fluorescent i*n situ* hybridization were performed on three animals per injured time point and on three uninjured controls processed in parallel. Cryosections were dried for 30 minutes at room temperature, then washed two times

for 10 minutes in phosphate buffer saline supplemented with 0.3% Triton X-100

(PBSTx) for permeabilization. *in situ* hybridization probes were denatured by warming

up at 70°C for 10 minutes and brief vortexing. Permeabilized sections were incubated

overnight in denatured probes at 62°C. The next day, sections were rinsed three

times (once for 15 minutes, followed by two washes of 30 minutes each) in washing

buffer (refer to key resource table) at 62°C. Then, in the case of chromogenic *in situ*

hybridization, sections were washed twice in MABT (key resource table) for 30

minutes each and incubated with 2% digoxigenin blocking reagent (key resource

table) in MABT for 1 hour at room temperature. In the case of fluorescent *in situ*

hybridization, sections were washed in TNT buffer (key resource table), and then

incubated with 2% digoxigenin blocking reagent dissolved in TNT buffer. After,

sections were incubated with the AP-conjugated, anti-digoxigenin antibody

(chromogenic *in situ* hybridization, 1:2000, key resource table) or with the POD-

conjugated, anti-digoxigenin antibody (fluorescent *in situ* hybridization, 1:2000, key

resource table) overnight, at 4°C. The next day, sections were washed four times for

20 minutes each in MABT (chromogenic *in situ* hybridization) or three times for 10

minutes each in TNT buffer (fluorescent *in situ* hybridization) at room temperature.

Then, sections for chromogenic *in situ* hybridization were washed for 5 minutes in

staining buffer (key resource table) and eventually incubated with 20 µl/ml NBT/BCIP

stock solution in staining buffer, at room temperature, to develop the colour reaction.

Time of incubation varied between three hours and overnight, depending on the

target mRNA and probe concentration. In the case of fluorescent *in situ* hybridization,

after TNT washing steps, sections were incubated for 30 minutes with 1:200 Cy3-

Tyramide in TSA amplification buffer (Akoya Biosciences®), at room temperature.

Finally, sections from chromogenic *in situ* hybridization were washed three times with

PBS, for 10 minutes each and mounted in 70% glycerol in PBS, ready for bright field

microscopy. Fluorescent *in situ* hybridization sections were rinsed three times in

PBS, for 10 minutes each, and then incubation with primary antibody for

immunohistochemistry was performed as described below.

1072

Immunohistochemistry

Immunohistochemistry was conducted as previously described (Grandel et al., 2006;

Kroehne et al., 2011). Immunohistochemistry was performed on three light-lesioned

animals and three, parallel processed, uninjured control per condition. Sections were

dried at room temperature for 30 minutes and washed 3 times in PBSTx for 10

minutes each. Then, primary antibodies diluted in PBSTx were applied overnight at

4°C (refer to the key resource table for primary antibody host, antigen specificity and

dilution). For proliferating cell nuclear antigen (PCNA) retrieval, sections were

immersed in 10 mM sodium citrate (pH: 6) for 6 minutes at a temperature higher than

80°C, then washed, and finally primary anti-PCNA antibody was applied. After

primary antibody incubation, sections were washed three times in PBSTx for 10

minutes each, then Alexa® fluorophore conjugated, secondary antibodies diluted In

PBSTx were applied accordingly (refer to key resource table for secondary antibody

host, primary antibody specificity and dilution) for 2 hours at room temperature.

Sections were eventually washed three times in PBS, mounted in 70% glycerol in

PBS, and stored at 4°C. Each staining was performed on three fish per time point

upon injury and on three uninjured fish controls processed in parallel. When

combined with fluorescent *in situ* hybridization, immunohistochemistry followed

fluorescent *in situ* hybridization processed sections.

1092

Microscopy

Bright field images of chromogenic *in situ* hybridizations and fluorescent images in Fiure1-figure supplement 2 were acquired with the Widefield ApoTome microscope (Zeiss Axio Imager.Z1) using the 20x/0.8 Plan-Apochromat, Air, DIC objective. Fluorescent images in Figure 1-figure supplement 1, Figure 3-figure supplement 2 and in Figure 4 and 6 were acquired with the laser scanning confocal microscope 780 (Zeiss Axio Imager.Z1) using the 40x/1.2 C-Apochromat, Water, DIC, Zeiss (Figure1-figure supplement 1 and Figure 4) and the 20x/0.8 Plan-Apochromat, Air, DIC, Zeiss (Figure 6). Fluorescent channels were acquired sequentially to minimize crosstalk in multicolour specimens.

Image acquisition and processing

Bright field microscope images were processed with the Zeiss ZEN blue, V 2012 software. The extended depth of focus function (contrast method) was applied to raw, bright field images of *her4.1*, *hmgb2b*, *atoh7* and *onecut1 in situ* hybridizations to project the chromogenic signal of the optical stack to a single plane. Fluorescent microscope images were processed using Fiji (Schindelin et al., 2012) for image contrast adjustments, crop and rotation. Adobe Illustrator CC 2015.3 was used to assemble figures. Excel 2016 (Microsoft) was used to generate the plot in Figure 8-figure supplement 1.

STATISTICS

Marker genes were ranked by their test statistic computed with a t-test for each gene, comparing cells within the cluster to the cells outside the cluster, as implemented in

the scanpy function tl.rank_genes_groups. The same was done to identify

differentially expressed genes between pairs of clusters.

ETHICAL STATEMENT

Fish were kept according to FELASA guidelines (Aleström et al., 2020; Brand et al.,

2002). All animal experiments were conducted according to the guidelines and under

supervision of the Regierungspräsidium Dresden (permit: TVV 44/2017). All efforts

were made to minimize animal suffering and the number of animals used.

DATA AVAILABILITY

All sequencing data that support the findings of this study have been deposited at

https://singlecell.broadinstitute.org/single_cell/reviewer_access/5a29ec20-d572-

4461-bfd3-cb8973c05f33 and are accessible through BioProject ID SCP1973.

Additionally, the data have been deposited at NCBI GEO

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE226373) with the GEO

accession number GSE226373.

Acknowledgements

We thank Dr. Judith Konantz, M. Fischer, S. Kunadt and D. Mögel for excellent

zebrafish care. We thank Katja Bernhardt for the excellent technical assistance in

flow cytometry and cell sorting. We would like to thank also Susanne Reinhardt and

Dr. Andreas Petzold from the DRESDEN-concept Genome Center for support with

initial planning and bioinformatic processing of the scRNAseq experiment,

respectively. We thank Prof. Marius Ader, Prof. Mike O. Karl, Dr. Volker Kroehne,

Juliane Hammer and Elisa Nerli for critically reading the manuscript. In addition, we

thank the members of the Brand laboratory for continued support and discussions as

well as helpful comments on the manuscript. We thank also Dr. Christian Lange, Dr.

Robert Münch and Dr. Steffen Rulands for excellent scientific discussion. This work

was supported by grants to M.B. from the Deutsche Forschungsgemeinschaft (BR

1746/11-1), the European Union (European Research Council AdG Zf-BrainReg) and

the German Excellence Initiative (Wissenschaftsrat and Deutsche

Forschungsgemeinschaft) (EXC 168). Work was supported by the DRESDEN-

concept Genome Center, the Light Microscopy Facility and Flow Cytometry Facility,

core facilities of the CMCB at the Technische Universität Dresden. The funders had

no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

1153

REFERENCES

Agathocleous M, Harris WA. 2009. From progenitors to differentiated cells in the
vertebrate retina. *Annu Rev Cell Dev Biol* **25**:45–69.
doi:10.1146/annurev.cellbio.042308.113259
Aleström P, D'Angelo L, Midtlyng PJ, Schorderet DF, Schulte-Merker S, Sohm F,
Warner S. 2020. Zebrafish: Housing and husbandry recommendations. *Lab
Anim* **54**:213–224. doi:10.1177/0023677219869037
Alexiades MR, Cepko CL. 1997. Subsets of retinal progenitors display temporally
regulated and distinct biases in the fates of their progeny. *Development*
124:1119–1131. doi:10.1242/dev.124.6.1119
Amini R, Rocha-Martins M, Norden C. 2018. Neuronal Migration and Lamination in
the Vertebrate Retina. *Front Neurosci* **11**:742. doi:10.3389/fnins.2017.00742
Baraban M, Anselme I, Schneider-Maunoury S, Giudicelli F. 2013. Zebrafish
Embryonic Neurons Transport Messenger RNA to Axons and Growth Cones
In Vivo. *J Neurosci* **33**:15726–15734. doi:10.1523/JNEUROSCI.1510-13.2013
Bernardos RL, Barthel LK, Meyers JR, Raymond PA. 2007. Late-Stage Neuronal
Progenitors in the Retina Are Radial Muller Glia That Function as Retinal Stem
Cells. *Journal of Neuroscience* **27**:7028–7040.
doi:10.1523/JNEUROSCI.1624-07.2007
Bernardos RL, Raymond PA. 2006. GFAP transgenic zebrafish. *Gene Expr Patterns*
6:1007–1013. doi:10.1016/j.modgep.2006.04.006

Blackshaw S, Harpavat S, Trimarchi J, Cai L, Huang H, Kuo WP, Weber G, Lee K, Fraioli RE, Cho S-H, Yung R, Asch E, Ohno-Machado L, Wong WH, Cepko CL. 2004. Genomic analysis of mouse retinal development. *PLoS Biol* **2**:E247. doi:10.1371/journal.pbio.0020247

Bosma PT, Blázquez M, Collins MA, Bishop JD, Drouin G, Priede IG, Docherty K, Trudeau VL. 1999. Multiplicity of glutamic acid decarboxylases (GAD) in vertebrates: molecular phylogeny and evidence for a new GAD paralog. *Molecular Biology and Evolution* **16**:397–404. doi:10.1093/oxfordjournals.molbev.a026120

Brand: Zebrafish: A practical approach - Google Scholar. n.d. https://scholar.google.com/scholar_lookup?title=Zebrafish:+A+Practical+Appro ach&author=M+Brand&author=M+Granato&author=C+N%C3%BCsslein-Volhard&publication_year=2002&

Buenaventura DF, Ghinia-Tegla MG, Emerson MM. 2018. Fate-restricted retinal progenitor cells adopt a molecular profile and spatial position distinct from multipotent progenitor cells. *Dev Biol* **443**:35–49. doi:10.1016/j.ydbio.2018.06.023

Cameron DA, Gentile KL, Middleton FA, Yurco P. 2005. Gene expression profiles of intact and regenerating zebrafish retina. *Mol Vis* **11**:775–791.

Campbell LJ, Levendusky JL, Steines SA, Hyde DR. 2022. Retinal regeneration requires dynamic Notch signaling. *Neural Regen Res* **17**:1199–1209. doi:10.4103/1673-5374.327326

Cepko C. 2014. Intrinsically different retinal progenitor cells produce specific types of progeny. *Nat Rev Neurosci* **15**:615–627. doi:10.1038/nrn3767

Cepko CL, Austin CP, Yang X, Alexiades M, Ezzeddine D. 1996. Cell fate determination in the vertebrate retina. *PNAS* **93**:589–595. doi:10.1073/pnas.93.2.589

Corso-Díaz X, Simpson EM. 2015. Nr2e1 regulates retinal lamination and the development of Müller glia, S-cones, and glycineric amacrine cells during retinogenesis. *Molecular Brain* **8**:37. doi:10.1186/s13041-015-0126-x

Craig SEL, Calinescu A-A, Hitchcock PF. 2008. Identification of the molecular signatures integral to regenerating photoreceptors in the retina of the zebra fish. *j ocul biol dis inform* **1**:73–84. doi:10.1007/s12177-008-9011-5

D'Aniello E, Ravisankar P, Waxman JS. 2015. Rdh10a Provides a Conserved Critical Step in the Synthesis of Retinoic Acid during Zebrafish Embryogenesis. *PLOS ONE* **10**:e0138588. doi:10.1371/journal.pone.0138588

Dowling JE. 1987. The Retina: An Approachable Part of the Brain. Harvard University Press.

Emerson MM, Surzenko N, Goetz JJ, Trimarchi J, Cepko CL. 2013. Otx2 and Onecut1 promote the fates of cone photoreceptors and horizontal cells and repress rod photoreceptors. *Dev Cell* **26**:59–72. doi:10.1016/j.devcel.2013.06.005

Fausett BV, Goldman D. 2006. A Role for 1 Tubulin-Expressing Muller Glia in Regeneration of the Injured Zebrafish Retina. *Journal of Neuroscience* **26**:6303–6313. doi:10.1523/JNEUROSCI.0332-06.2006

Fausett BV, Gumerson JD, Goldman D. 2008. The Proneural Basic Helix-Loop-Helix Gene Ascl1a Is Required for Retina Regeneration. *Journal of Neuroscience* **28**:1109–1117. doi:10.1523/JNEUROSCI.4853-07.2008

Feng L, Hernandez RE, Waxman JS, Yelon D, Moens CB. 2010. Dhrs3a regulates retinoic acid biosynthesis through a feedback inhibition mechanism. *Developmental Biology* **338**:1–14. doi:10.1016/j.ydbio.2009.10.029

Fimbel SM, Montgomery JE, Burket CT, Hyde DR. 2007. Regeneration of Inner Retinal Neurons after Intravitreal Injection of Ouabain in Zebrafish. *Journal of Neuroscience* **27**:1712–1724. doi:10.1523/JNEUROSCI.5317-06.2007

Fleisch VC, Fraser B, Allison WT. 2011. Investigating regeneration and functional integration of CNS neurons: Lessons from zebrafish genetics and other fish species. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1812**:364–380. doi:10.1016/j.bbadis.2010.10.012

Godinho L, Williams PR, Claassen Y, Provost E, Leach SD, Kamermans M, Wong ROL. 2007. Nonapical Symmetric Divisions Underlie Horizontal Cell Layer Formation in the Developing Retina In Vivo. *Neuron* **56**:597–603. doi:10.1016/j.neuron.2007.09.036

Gorsuch RA, Hyde DR. 2014. Regulation of Müller glial dependent neuronal regeneration in the damaged adult zebrafish retina. *Experimental Eye Research* **123**:131–140. doi:10.1016/j.exer.2013.07.012

Gorsuch RA, Lahne M, Yarka CE, Petravick ME, Li J, Hyde DR. 2017. Sox2 regulates Müller glia reprogramming and proliferation in the regenerating zebrafish retina via Lin28 and Ascl1a. *Exp Eye Res* **161**:174–192. doi:10.1016/j.exer.2017.05.012

Grandel H, Kaslin J, Ganz J, Wenzel I, Brand M. 2006. Neural stem cells and neurogenesis in the adult zebrafish brain: Origin, proliferation dynamics, migration and cell fate. *Developmental Biology* **295**:263–277. doi:10.1016/j.ydbio.2006.03.040

Hoang T, Wang J, Boyd P, Wang F, Santiago C, Jiang L, Yoo S, Lahne M, Todd LJ, Jia M, Saez C, Keuthan C, Palazzo I, Squires N, Campbell WA, Rajaii F, Parayil T, Trinh V, Kim DW, Wang G, Campbell LJ, Ash J, Fischer AJ, Hyde DR, Qian J, Blackshaw S. 2020. Gene regulatory networks controlling vertebrate retinal regeneration. *Science* **370**:eabb8598. doi:10.1126/science.abb8598

Holmes RS, Hempel J. 2011. Comparative studies of vertebrate aldehyde dehydrogenase 3: Sequences, structures, phylogeny and evolution. Evidence for a mammalian origin for the ALDH3A1 gene. *Chemico-Biological Interactions*, Enzymology and Molecular Biology of Carbonyl Metabolism **191**:113–121. doi:10.1016/j.cbi.2011.01.014

Hoon M, Okawa H, Santina LD, Wong ROL. 2014. Functional Architecture of the Retina: Development and Disease. *Prog Retin Eye Res* **42**:44–84. doi:10.1016/j.preteyeres.2014.06.003

Javed A, Cayouette M. 2017. Temporal Progression of Retinal Progenitor Cell Identity: Implications in Cell Replacement Therapies. *Frontiers in Neural Circuits* **11**.

Jusuf PR, Almeida AD, Randlett O, Joubin K, Poggi L, Harris WA. 2011. Origin and Determination of Inhibitory Cell Lineages in the Vertebrate Retina. *J Neurosci* **31**:2549–2562. doi:10.1523/JNEUROSCI.4713-10.2011

Jusuf PR, Harris WA. 2009. Ptf1a is expressed transiently in all types of amacrine cells in the embryonic zebrafish retina. *Neural Dev* **4**:34. doi:10.1186/1749-8104-4-34

Kassen SC, Ramanan V, Montgomery JE, T. Burket C, Liu C-G, Vihtelic TS, Hyde DR. 2007. Time course analysis of gene expression during light-induced photoreceptor cell death and regeneration in *albino* zebrafish. *Devel Neurobio* **67**:1009–1031. doi:10.1002/dneu.20362

Kitambi SS, Hauptmann G. 2007. The zebrafish orphan nuclear receptor genes
nr2e1 and nr2e3 are expressed in developing eye and forebrain. *Gene*
Expression Patterns **7**:521–528. doi:10.1016/j.modgep.2006.10.006
Klimova L, Antosova B, Kuzelova A, Strnad H, Kozmik Z. 2015. Onecut1 and
Onecut2 transcription factors operate downstream of Pax6 to regulate
horizontal cell development. *Developmental Biology* **402**:48–60.
doi:10.1016/j.ydbio.2015.02.023
Kramer AC, Gurdziel K, Thummel R. 2021. A Comparative Analysis of Gene and
Protein Expression Throughout a Full 28-Day Retinal Regeneration Time-
Course in Adult Zebrafish. *Front Cell Dev Biol* **9**:741514.
doi:10.3389/fcell.2021.741514
Kroehne V, Freudenreich D, Hans S, Kaslin J, Brand M. 2011. Regeneration of the
adult zebrafish brain from neurogenic radial glia-type progenitors.
Development **138**:4831–4841. doi:10.1242/dev.072587
Krylov A, Yu S, Veen K, Newton A, Ye A, Qin H, He J, Jusuf PR. 2023. Heterogeneity
in quiescent Müller glia in the uninjured zebrafish retina drive differential
responses following photoreceptor ablation. *Frontiers in Molecular*
Neuroscience **16**.
Lahne M, Brecker M, Jones SE, Hyde DR. 2021. The Regenerating Adult Zebrafish
Retina Recapitulates Developmental Fate Specification Programs. *Front Cell*
Dev Biol **8**:617923. doi:10.3389/fcell.2020.617923
Lahne M, Li J, Marton RM, Hyde DR. 2015. Actin-Cytoskeleton- and Rock-Mediated
INM Are Required for Photoreceptor Regeneration in the Adult Zebrafish
Retina. *J Neurosci* **35**:15612–15634. doi:10.1523/JNEUROSCI.5005-14.2015
Lahne M, Nagashima M, Hyde DR, Hitchcock PF. 2020. Reprogramming Müller Glia
to Regenerate Retinal Neurons. *Annu Rev Vis Sci* **6**:171–193.
doi:10.1146/annurev-vision-121219-081808
Lange C, Rost F, Machate A, Reinhardt S, Lesche M, Weber A, Kuscha V, Dahl A,
Rulands S, Brand M. 2020. Single cell sequencing of radial glia progeny
reveals diversity of newborn neurons in the adult zebrafish brain. *Development*
1855951. doi:10.1242/dev.185595
Lenkowski JR, Raymond PA. 2014. Müller glia: Stem cells for generation and
regeneration of retinal neurons in teleost fish. *Progress in Retinal and Eye*
Research **40**:94–123. doi:10.1016/j.preteyeres.2013.12.007
Livesey FJ, Cepko CL. 2001. Vertebrate neural cell-fate determination: lessons from
the retina. *Nat Rev Neurosci* **2**:109–118. doi:10.1038/35053522
Lun ATL, McCarthy DJ, Marioni JC. 2016. A step-by-step workflow for low-level
analysis of single-cell RNA-seq data with Bioconductor.
doi:10.12688/f1000research.9501.2
Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas
AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek
AK, Regev A, McCarroll SA. 2015. Highly Parallel Genome-wide Expression
Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**:1202–1214.
doi:10.1016/j.cell.2015.05.002
Matthews RP, Lorent K, Russo P, Pack M. 2004. The zebrafish onecut gene hnf-6
functions in an evolutionarily conserved genetic pathway that regulates
vertebrate biliary development. *Developmental Biology* **274**:245–259.
doi:10.1016/j.ydbio.2004.06.016
Morris AC, Forbes-Osborne MA, Pillai LS, Fadool JM. 2011. Microarray Analysis of
XOPS-mCFP Zebrafish Retina Identifies Genes Associated with Rod

Photoreceptor Degeneration and Regeneration. *Investigative Ophthalmology & Visual Science* **52**:2255–2266. doi:10.1167/iovs.10-6022

Morris AC, Schroeter EH, Bilotta J, Wong ROL, Fadool JM. 2005. Cone survival despite rod degeneration in XOPS-mCFP transgenic zebrafish. *Invest Ophthalmol Vis Sci* **46**:4762–4771. doi:10.1167/iovs.05-0797

Mu Z, Zhang S, He C, Hou H, Liu D, Hu N, Xu H. 2016. Expression of SoxC Transcription Factors during Zebrafish Retinal and Optic Nerve Regeneration. *Neurosci Bull* **33**:53–61. doi:10.1007/s12264-016-0073-2

Muto A, Iida A, Satoh S, Watanabe S. 2009. The group E Sox genes Sox8 and Sox9 are regulated by Notch signaling and are required for Müller glial cell development in mouse retina. *Exp Eye Res* **89**:549–558. doi:10.1016/j.exer.2009.05.006

Nagashima M, Barthel LK, Raymond PA. 2013. A self-renewing division of zebrafish Müller glial cells generates neuronal progenitors that require N-cadherin to regenerate retinal neurons. *Development* **140**:4510–4521. doi:10.1242/dev.090738

Nelson CM, Ackerman KM, O'Hayer P, Bailey TJ, Gorsuch RA, Hyde DR. 2013. Tumor Necrosis Factor-Alpha Is Produced by Dying Retinal Neurons and Is Required for Muller Glia Proliferation during Zebrafish Retinal Regeneration. *Journal of Neuroscience* **33**:6524–6539. doi:10.1523/JNEUROSCI.3838-12.2013

Nerli E, Kretzschmar J, Bianucci T, Rocha-Martins M, Zechner C, Norden C. 2022. Deterministic and probabilistic fate decisions co-exist in a single retinal lineage. doi:10.1101/2022.08.11.503564

Ng Chi Kei J, Currie PD, Jusuf PR. 2017. Fate bias during neural regeneration adjusts dynamically without recapitulating developmental fate progression. *Neural Dev* **12**:12. doi:10.1186/s13064-017-0089-y

O'Malley DM, Sandell JH, Masland RH. 1992. Co-release of acetylcholine and GABA by the starburst amacrine cells. *J Neurosci* **12**:1394–1408. doi:10.1523/JNEUROSCI.12-04-01394.1992

Peterson RE, Fadool JM, Mcclintock J, Linser PJ. 2001. Müller cell differentiation in the zebrafish neural retina: Evidence of distinct early and late stages in cell maturation. *Journal of Comparative Neurology* **429**:530–540. doi:10.1002/1096-9861(20010122)429:4<530::AID-CNE2>3.0.CO;2-C

Poché RA, Furuta Y, Chaboissier M-C, Schedl A, Behringer RR. 2008. Sox9 is expressed in mouse multipotent retinal progenitor cells and functions in Müller Glial cell development. *Journal of Comparative Neurology* **510**:237–250. doi:10.1002/cne.21746

Poggi L, Vitorino M, Masai I, Harris WA. 2005. Influences on neural lineage and mode of division in the zebrafish retina in vivo. *J Cell Biol* **171**:991–999. doi:10.1083/jcb.200509098

Powell C, Cornblath E, Elsaeidi F, Wan J, Goldman D. 2016. Zebrafish Müller glia-derived progenitors are multipotent, exhibit proliferative biases and regenerate excess neurons. *Sci Rep* **6**:24851. doi:10.1038/srep24851

Qin Z, Barthel LK, Raymond PA. 2009. Genetic evidence for shared mechanisms of epimorphic regeneration in zebrafish. *Proc Natl Acad Sci U S A* **106**:9310–9315. doi:10.1073/pnas.0811186106

Ramachandran R, Reifler A, Parent JM, Goldman D. 2010. Conditional gene expression and lineage tracing of tuba1a expressing cells during zebrafish development and retina regeneration. *J Comp Neurol* **518**:4196–4212. doi:10.1002/cne.22448

Ramachandran R, Zhao X-F, Goldman D. 2012. Insm1a-mediated gene repression is essential for the formation and differentiation of Müller glia-derived progenitors in the injured retina. *Nat Cell Biol* **14**:1013–1023. doi:10.1038/ncb2586

Ramachandran R, Zhao X-F, Goldman D. 2011. Ascl1a/Dkk/ -catenin signaling pathway is necessary and glycogen synthase kinase-3 inhibition is sufficient for zebrafish retina regeneration. *Proceedings of the National Academy of Sciences* **108**:15858–15863. doi:10.1073/pnas.1107220108

Raymond PA, Barthel LK, Bernardos RL, Perkowski JJ. 2006. Molecular characterization of retinal stem cells and their niches in adult zebrafish. *BMC Dev Biol* **6**:36. doi:10.1186/1471-213X-6-36

Saade CJ, Alvarez-Delfin K, Fadool JM. 2013. Rod photoreceptors protect from cone degeneration-induced retinal remodeling and restore visual responses in zebrafish. *J Neurosci* **33**:1804–1814. doi:10.1523/JNEUROSCI.2910-12.2013

Sapkota D, Chintala H, Wu F, Fliesler SJ, Hu Z, Mu X. 2014. Onecut1 and Onecut2 redundantly regulate early retinal cell fates during development. *PNAS* **111**:E4086–E4095. doi:10.1073/pnas.1405354111

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Sherpa T, Fimbel SM, Mallory DE, Maaswinkel H, Spritzer SD, Sand JA, Li L, Hyde DR, Stenkamp DL. 2008. Ganglion cell regeneration following whole-retina destruction in zebrafish. *Devel Neurobio* **68**:166–181. doi:10.1002/dneu.20568

Silva NJ, Nagashima M, Li J, Kakuk-Atkins L, Ashrafzadeh M, Hyde DR, Hitchcock PF. 2020. Inflammation and matrix metalloproteinase 9 (Mmp-9) regulate photoreceptor regeneration in adult zebrafish. *Glia* **68**:1445–1465. doi:10.1002/glia.23792

Söllner C, Burghammer M, Busch-Nentwich E, Berger J, Schwarz H, Riekel C, Nicolson T. 2003. Control of crystal size and lattice formation by starmaker in otolith biomineralization. *Science* **302**:282–286. doi:10.1126/science.1088443

Stenkamp DL. 2015. Development of the Vertebrate Eye and Retina. *Prog Mol Biol Transl Sci* **134**:397–414. doi:10.1016/bs.pmbts.2015.06.006

Stenkamp DL. 2011. The rod photoreceptor lineage of teleost fish. *Progress in Retinal and Eye Research* **30**:395–404. doi:10.1016/j.preteyeres.2011.06.004

Stenkamp DL. 2007. Neurogenesis in the fish retina. *Int Rev Cytol* **259**:173–224. doi:10.1016/S0074-7696(06)59005-9

Suzuki SC, Bleckert A, Williams PR, Takechi M, Kawamura S, Wong ROL. 2013. Cone photoreceptor types in zebrafish are generated by symmetric terminal divisions of dedicated precursors. *PNAS* **110**:15109–15114. doi:10.1073/pnas.1303551110

Takke C, Dornseifer P, Weizsäcker E v., Campos-Ortega JA. 1999. her4, a zebrafish homologue of the Drosophila neurogenic gene E(spl), is a target of NOTCH signalling. *Development* **126**:1811–1821. doi:10.1242/dev.126.9.1811

Thummel R, Enright JM, Kassen SC, Montgomery JE, Bailey TJ, Hyde DR. 2010. Pax6a and Pax6b are required at different points in neuronal progenitor cell proliferation during zebrafish photoreceptor regeneration. *Experimental Eye Research* **90**:572–582. doi:10.1016/j.exer.2010.02.001

Thummel R, Kassen SC, Enright JM, Nelson CM, Montgomery JE, Hyde DR. 2008. Characterization of Müller glia and neuronal progenitors during adult zebrafish retinal regeneration. *Experimental Eye Research* **87**:433–444. doi:10.1016/j.exer.2008.07.009

Tirosh I, Izar B, Prakadan SM, Wadsworth MH, Treacy D, Trombetta JJ, Rotem A, Rodman C, Lian C, Murphy G, Fallahi-Sichani M, Dutton-Regester K, Lin J-R, Cohen O, Shah P, Lu D, Genshaft AS, Hughes TK, Ziegler CGK, Kazer SW, Gaillard A, Kolb KE, Villani A-C, Johannessen CM, Andreev AY, Van Allen EM, Bertagnolli M, Sorger PK, Sullivan RJ, Flaherty KT, Frederick DT, Jané-Valbuena J, Yoon CH, Rozenblatt-Rosen O, Shalek AK, Regev A, Garraway LA. 2016. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**:189–196. doi:10.1126/science.aad0501

Turner DL, Cepko CL. 1987. A common progenitor for neurons and glia persists in rat retina late in development. *Nature* **328**:131–136. doi:10.1038/328131a0

Vihtelic TS, Hyde DR. 2000. Light-induced rod and cone cell death and regeneration in the adult albino zebrafish (Danio rerio) retina. *J Neurobiol* **44**:289–307. doi:10.1002/1097-4695(20000905)44:3<289::aid-neu1>3.0.co;2-h

Vihtelic TS, Soverly JE, Kassen SC, Hyde DR. 2006. Retinal regional differences in photoreceptor cell death and regeneration in light-lesioned albino zebrafish. *Experimental Eye Research* **82**:558–575. doi:10.1016/j.exer.2005.08.015

Wan J, Goldman D. 2016. Retina regeneration in zebrafish. *Curr Opin Genet Dev* **40**:41–47. doi:10.1016/j.gde.2016.05.009

Wan J, Ramachandran R, Goldman D. 2012. HB-EGF Is Necessary and Sufficient for Müller Glia Dedifferentiation and Retina Regeneration. *Developmental Cell* **22**:334–347. doi:10.1016/j.devcel.2011.11.020

Weber A, Hochmann S, Cimalla P, Gärtner M, Kuscha V, Hans S, Geffarth M, Kaslin J, Koch E, Brand M. 2013. Characterization of light lesion paradigms and optical coherence tomography as tools to study adult retina regeneration in zebrafish. *PLoS One* **8**:e80483. doi:10.1371/journal.pone.0080483

Weber IP, Ramos AP, Strzyz PJ, Leung LC, Young S, Norden C. 2014. Mitotic Position and Morphology of Committed Precursor Cells in the Zebrafish Retina Adapt to Architectural Changes upon Tissue Maturation. *Cell Reports* **7**:386–397. doi:10.1016/j.celrep.2014.03.014

Whitfield ML, Sherlock G, Saldanha AJ, Murray JI, Ball CA, Alexander KE, Matese JC, Perou CM, Hurt MM, Brown PO, Botstein D. 2002. Identification of genes periodically expressed in the human cell cycle and their expression in tumors. *Mol Biol Cell* **13**:1977–2000. doi:10.1091/mbc.02-02-0030

Wilson SG, Wen W, Pillai-Kastoori L, Morris AC. 2016. Tracking the fate of her4 expressing cells in the regenerating retina using her4:Kaede zebrafish. *Exp Eye Res* **145**:75–87. doi:10.1016/j.exer.2015.11.002

Wu F, Bard JE, Kann J, Yergeau D, Sapkota D, Ge Y, Hu Z, Wang J, Liu T, Mu X. 2021. Single cell transcriptomics reveals lineage trajectory of retinal ganglion cells in wild-type and Atoh7-null retinas. *Nat Commun* **12**:1465. doi:10.1038/s41467-021-21704-4

Xu B, Tang X, Jin M, Zhang H, Du L, Yu S, He J. 2020. Unifying developmental programs for embryonic and postembryonic neurogenesis in the zebrafish retina. *Development* **147**:dev185660. doi:10.1242/dev.185660

Young RW. 1985. Cell differentiation in the retina of the mouse. *Anat Rec* **212**:199–205. doi:10.1002/ar.1092120215

Yurco P, Cameron DA. 2005. Responses of Müller glia to retinal injury in adult zebrafish. *Vision Research* **45**:991–1002. doi:10.1016/j.visres.2004.10.022

Zhang Y, Buchholz F, Muyrers JPP, Stewart AF. 1998. A new logic for DNA engineering using recombination in Escherichia coli. *Nat Genet* **20**:123–128. doi:10.1038/2417

Zhao X-F, Wan J, Powell C, Ramachandran R, Myers MG, Goldman D. 2014. Leptin
and IL-6 family cytokines synergize to stimulate Müller glia reprogramming and
retina regeneration. *Cell Rep* **9**:272–284. doi:10.1016/j.celrep.2014.08.047
Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB,
Wheeler TD, McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L,
Underwood JG, Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW,
Hindson CM, Bharadwaj R, Wong A, Ness KD, Beppu LW, Deeg HJ,
McFarland C, Loeb KR, Valente WJ, Ericson NG, Stevens EA, Radich JP,
Mikkelsen TS, Hindson BJ, Bielas JH. 2017. Massively parallel digital
transcriptional profiling of single cells. *Nat Commun* **8**:14049.
doi:10.1038/ncomms14049

1488

1489

FIGURE CAPTIONS

Figure 1: Isolation of individual MG, RPCs and regenerated progeny using

fluorescent reporters. (A) Schematic illustration of the rationale to isolate individual

cells from regenerating retinae of double transgenic

Tg(pcna:EGFP);Tg(gfap:mCherry) animals. In the uninjured retina, MG express

strong levels of mCherry but no EGFP. Upon light lesion, which results in dying

photoreceptors, MG turn on EGFP resulting in the presence of both fluorophores at

44 hours post lesion (hpl). At 4 days post lesion (dpl), MG remain to contain strong

mCherry and EGFP levels. In contrast, further proliferating RPCs display weak

mCherry but strong EGFP fluorescence. Finally, MG have lost the EGFP signal at 6

dpl, expressing mCherry only. RPCs, which start to migrate and differentiate, are still

detectable but show diluted levels of mCherry and EGFP. **(B)** Schematic illustration

of the workflow for the scRNAseq experiment. Uninjured control and light lesioned

retinae (44 hpl, 4 dpl, 6 dpl) of double transgenic *Tg(pcna:EGFP);Tg(gfap:mCherry)*

animals were dissociated to obtain a single cell suspension. Following fluorescent

activated cell sorting of mCherry-positive only and EGFP/mCherry-double positive

cells, individual cells were eventually subjected to droplet-based, 10x Genomics

scRNAseq. **(C)** Flow cytometry plots depicting EGFP and mCherry fluorescence of

dissociated retinal cells at the indicated time points. Cells in the sorting gate include

mCherry-positive only as well as mCherry/EGFP-double positive cells. Number of

cells sorted from the sorting gate per time point: 15.000.

Associated: figure supplements 1, 2 and 3.

1512

Figure 2: scRNAseq identifies 15 different cell populations in the light-lesioned

retina. (A) 11.690 cells depicted on the uniform manifold approximation and

projection (UMAP) map. From left to right, the map shows a progressive emergence

of MG, RPCs and progeny clusters. See figure legend for the detailed information

about cluster identity. **(B)** Normalized cell density on the UMAP for the 4 different

samples (uninjured, 44 hpl, 4 and 6 dpl). Dark red shows strong cell enrichment and

no color absence of cells. **(C)** Cells colored by cell cycle phases (G1, S, and G2/M

phase in blue, green and orange, respectively). **(D)** Expression of *pcna*. Legend:

logarithm of normalized counts per million. **Associated: figure supplements 1 and**

2.

1523

Figure 3: scRNAseq reveals heterogeneity in Müller glia and identifies two

populations of non-reactive and two of reactive Müller glia in the light-lesioned

retina. (A) Four Müller glia clusters (cluster 1-4) and the expression levels of *apoeb*

and *gfap*. **(B)** The two non-reactive Müller glia clusters (cluster 1 and 2) and the

expression levels of *mstnb* and *inhbaa*. Cells of both clusters express *mstnb*, but only

cells in cluster 1 expresses *inhbaa*. **(C)** Visualization of the two reactive Müller glia

clusters (cluster 3 and 4) and the relative expression levels of *crlf1a* and *her4.1*

projected onto the main transcriptional UMAP. Cells of both clusters express *crlf1a*,

but only cells in cluster 3 expresses *her4.1*. **(D)** Bright field microscopy images of

chromogenic *in situ* hybridization detecting *her4.1* on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 days post lesion (dpl) and 6 dpl. N= 3 fish per time point and uninjured controls. The position of the outer nuclear layer (ONL), inner nuclear layer (INL) and ganglion cell layer (GCL) is indicated in the uninjured sample. Scale bar: 30 μm. **Associated: figure supplements 1 and 2.**

Figure 4: expression profile of *crlf1a* during retina regeneration. (A) Confocal images of fluorescent *in situ* hybridization for *crlf1a* (grey) and immunohistochemistry detecting Zrf1-positive MG fibers (magenta) and the proliferation marker PCNA (green) on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 days post lesion (dpl) and 6 dpl. The position of the outer nuclear layer (ONL), inner nuclear laser (INL) and ganglion cell layer (GCL) is indicated in the uninjured sample. N= 3 fish per time point and uninjured controls. Scale bar: 30 μm. **(B)** Magnification of the areas indicated by the squares 1 and 2 in panel A. Scale bar: 10 μm.

Figure 5: scRNAseq identifies a transient cell population with hybrid characteristics of reactive Müller glia and early progenitors in the light-lesioned retina. (A) The progenitor 1 cluster (cluster 5) and the expression levels of *hmgb2b*, *stmn1a*, *nr2e1 sox9b* and *id1*. **(B)** Bright field microscopy images of chromogenic *in situ* hybridization detecting *hmgb2b* on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 days post lesion (dpl) and 6 dpl. N= 3 fish per time point and uninjured controls. The position of the outer nuclear layer (ONL), inner nuclear layer (INL) and ganglion cell layer (GCL) is indicated in the uninjured sample. Scale bar: 30 μm.

1558

Figure 6: expression profile of *id1* during retina regeneration. (A) Confocal images of fluorescent *in situ* hybridization for *id1* (grey) and immunohistochemistry detecting Zrf1-positive MG fibers (magenta) and the proliferation marker PCNA (green) on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 and 6 days post lesion (dpl). The position of the outer nuclear layer (ONL), inner nuclear layer (INL) and ganglion cell layer (GCL) is indicated in the uninjured sample. N= 3 fish per time point and uninjured controls. Scale bar: 30 µm. **(B)** Magnification of the areas indicated by the squares 1 and 2 in panel A. Scale bar: 10 µm.

1568

Figure 7: scRNAseq identifies two neurogenic progenitor populations in the light-lesioned retina. (A) The two progenitor clusters (cluster 6 and 7) and the relative expression levels of *insm1a*, *atoh7*, *onecut1*, *onecut2* and *mki67*. **(B)** Bright field microscopy images of chromogenic *in situ* hybridization detecting *atoh7*, *onecut1* or *onecut2* on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 days post lesion (dpl) and 6 dpl. Each ISH was carried on N= 3 fish per time point and uninjured controls.The position of the outer nuclear layer (ONL), inner nuclear layer (INL) and ganglion cell layer (GCL) is indicated in the uninjured sample. Scale bar: 30 µm.

1578

Figure 8: scRNAseq identifies fate-restricted progenitors and differentiating progeny in the light-lesioned retina. (A) The photoreceptor precursor cluster (cluster 8) and the expression levels of *crx* and *thrb*. **(B)** The red and blue cone

cluster (cluster 11 and 12) and the expression levels of the blue cone marker *opn1sw2* and red cone marker *opn1lw2*. **(C)** The horizontal cell precursor cluster (cluster 9) and the expression levels of *ptf1a* and *rem1*. **(D)** The ganglion cell cluster (cluster 10) and the expression levels of *gap43* and *isl2b*. **(E)** The two amacrine cell clusters (cluster 14 and 15) and the expression levels of *stx1b* and *pvalb6*. **(D)** The bipolar cell cluster (cluster 15) and the expression levels of the known bipolar cell marker *vsx1* and the novel marker *samsn1a*. **Associated: figure supplements 1 and 2.**

Figure 9: comparison between the present scRNAseq dataset and the scRNAseq dataset by Hoang et al., 2020. (A) UMAP showing all cell clusters annotated in Hoang et al., 2020 and mapped to the scRNAseq dataset described in the present study. **(B)** UMAP showing the individual cell clusters annotated in Hoang et al., 2020 and mapped to the scRNAseq dataset described in the present study.

Figure 10: model summarizing the cellular events underlying retina regeneration in zebrafish. Upon lesion, non-reactive MG on the left side of the main transcriptional UMAP feed into a hybrid Müller glia/progenitor population via reactive Müller glia (solid red line). Based from this hybrid cell population, we find a Müller glia trajectory reverting to the initial starting point indicating Müller glia self-renewal (dashed red line). Moreover, the hybrid cell population represents the tip of the neurogenic trajectory, resulting in the formation of neuronal precursors and differentiating neurons (solid black lines). Differentiating rods and horizontal cells, which have not been found in the scRNAseq, are indicated with dashed lines and positioned in the predicted location on the main UMAP.

1607

Figure 1-supplemental figure 3: fluorescence activated cell sorting (FACS) and

gating strategies. (A) Flow cytometry plots of dissociated, uninjured retinae from

Tg(pcna:EGFP);Tg(gfap:mCherry) fish and gating strategy for sorting calcein-positive

cells. Plot 1 displays all the events in a single cell suspension according to the

forward (x axis) and side scatter (y axis) areas. Among cells gated in 1, singlets (P2)

were chosen by considering the area (x axis) and the height (y axis) of the forward

scatter in 2. Height (x axis) and width (y axis) of the side scatter were plotted in 3 to

further gate singlets among P2. Finally, only calcein-positive (i.e. living) cells were

gated among singlets in 4. The gating strategy was the same for the uninjured as well

as all lesioned samples. Samples consisted of single cell suspensions derived from 8

retinae of 4 fish (2 males and 2 females) each. **(B)** Flow cytometry plots showing

EGFP and mCherry fluorescence of cells from WT, *Tg(gfap:mCherry)* and

Tg(pcna:EGFP) dissociated retinae.

1621

Figure 1-figure supplement 1: characterization of the fluorescent reporter lines

***Tg(pcna:EGFP)* and *Tg(gfap:mCherry)* in double transgenic animals. (A)**

Confocal images of immunohistochemistry against the proliferation marker PCNA

(grey) and EGFP (blue) on retinal sections of *Tg(pcna:EGFP);Tg(gfap:mCherry)*

animals. Schemes in the thumbnails on the top-right and top-left of the PCNA panels

indicate the region of interest, either the central retina or the ciliary marginal zone

(CMZ). N= 3 fish. **(B)** Confocal images of immunohistochemistry detecting Zrf1-

positive MG fibers (magenta) and mCherry on retinal sections of

Tg(pcna:EGFP);Tg(gfap:mCherry) animals. N= 3 fish. Scale bar: 30 μm for main

panels, 10 μm for inserts.

Figure 1-figure supplement 2: expression profile of *Tg(pcna:EGFP);Tg(gfap:mCherry)* during retina regeneration. Apotome images showing endogenous immunofluorescence of EGFP and mCherry driven by *Tg(pcna:EGFP)* and *Tg(gfap:mCherry)* on retinal sections in uninjured as well as lesioned conditions at different time points. White arrowheads indicate mCherry-positive, EGFP-negative MG. Yellow arrowheads indicate mCherry-/EGFP-double positive MG. Red arrowheads indicate low-mCherry-/EGFP-double positive, putative RPCs. N= 3 fish per time point and uninjured controls. Scale bar: 30 µm.

Figure 2-figure supplement 1: original scRNAseq dataset. (A) UMAP of the original scRNAseq dataset containing 13.139 cells and consisting of a core cluster (circled in red) and two clusters of microglia and ribosomal gene-enriched cells (circled in grey). **(B)** Heat map depicting the expression of the top 10 marker genes per cluster of the original dataset.

Figure 2-figure supplement 2: dot plot of top 5 marker genes per cluster. Dot plot depicting the expression of top 5 marker genes per cluster (see Figure 2A). Dot size scales with a fraction of cells expressing the gene, and the dot colour indicates the scaled mean gene expression in the cluster.

Figure 3-figure supplement 1: expression of selected genes per cluster. (A) Heat map showing differentially expressed genes between non-reactive MG 1 and non-reactive MG 2. **(B)** Heat map showing differentially expressed genes (*gfap, pcna, her15.1-1, her9, her4.1*) as well as shared (*hbegfa*) genes between reactive MG 1

and reactive MG 2. **(C)** Heat map showing differentially expressed genes in reactive MG1, reactive MG2 and P1. **(D)** Heat map showing differentially expressed genes in P1, P2 and P3.

Figure 3-figure supplement 2: expression profile of *stm* during retina regeneration. (A) Confocal images of fluorescent *in situ* hybridization for *stm* (grey) and immunohistochemistry detecting Zrf1-positive MG fibers (magenta) and the proliferation marker PCNA (green) on retinal sections of uninjured control animals and at 44 hours post lesion (hpl), 4 and 6 days post lesion (dpl). The yellow arrowhead at 44 hpl indicates a *stm*-, Zrf1- and PCNA-triple positive Müller glia. N= 3 fish per time point and uninjured controls. Scale bar: 30 µm.

Figure 8-figure supplement 1: birthdate order of regenerated progeny. Graph plotting the fraction of cells (y-axis) embarking on a specific retinal cell fate (coloured curves) in relation to a given time point (x-axis). Total number of cells per condition: uninjured= 2243, 44 hpl= 1723, 4 dpl= 4899, 6 dpl= 2825.

Figure 8-figure supplement 2: expression of mature markers. (A) Expression levels of the known rod markers *rho* and *gnat1*. **(B)** Expression levels of the known mature amacrine and horizontal cell markers *gad1b* and *chata*. **(C)** Expression levels of the known mature amacrine cell and horizontal cell markers *calb2a* and *calb2b*. **(D)** Expression levels of the known mature bipolar cell markers *prkcab* and *prkcbb*. **(E)** Expression levels of the general synaptic markers *snap25a* and *snap25b*.

Figure 9-figure supplement 1: dot plot of selected marker genes per cluster in the present scRNAseq dataset and the scRNAseq dataset by Hoang et al., 2020. Dot plot depicting the expression of our selected marker genes (x-axis) in the clusters annotated by Hoang et al., 2020 (y-axis). Marker genes were chosen from the scRNAseq dataset described in the present study

SUPPLEMENTAL FIGURES

Figure 1-figure supplement 1: characterization of the fluorescent reporter lines *Tg(pcna:EGFP)* and *Tg(gfap:mCherry)* in double transgenic animals.

Figure 1-figure supplement 2: expression profile of *Tg(pcna:EGFP);Tg(gfap:mCherry)* during retina regeneration.

Figure 1-figure supplement 3: fluorescence activated cell sorting (FACS) and gating strategies.

Figure 2-figure supplement 1: original scRNAseq dataset

Figure 2-figure supplement 2: dot plot of top 5 marker genes per cluster.

Figure 3-figure supplement 1: expression of selected genes per cluster.

Figure 3-figure supplement 2: expression profile of *stm* during retina regeneration.

Figure 8-figure supplement 1: birthdate order of regenerated progeny.

Figure 8-figure supplement 2: expression of mature markers.

Figure 9-figure supplement 1: dot plot of selected marker genes per cluster in the present scRNAseq dataset and the scRNAseq dataset by Hoang et al., 2020.

SUPPLEMENTARY FILES

Appendix: sequence based reagents

Supplementary file 1: Top 100 marker genes per cluster

Supplementary file 2: Top 100 differentially expressed genes between Non-reactive

Müller glia (1) and Non-reactive Müller glia (2)

Supplementary file 3: Top 100 differentially expressed genes between Reactive

Müller glia (1) and Reactive Müller glia (2)

Supplementary file 4: Top 100 differentially expressed genes between Reactive

Müller glia (1) and Müller glia/Progenitors 1

Supplementary file 5: Top 100 differentially expressed genes between Reactive

Müller glia (2) and Müller glia/Progenitors 1

Supplementary file 6: Top 100 differentially expressed genes between Müller

glia/Progenitors 1 and Progenitors 2

Figure 1

A Uninjured   44 hpl   4 dpl   6 dpl

LEGEND

Retinal neurons

Non-reactive MG
Strong mCherry
No EGFP

Reactive MG
Strong mCherry
Strong EGFP

Dying photoreceptors

Proliferating RPC
Strong mCherry
Strong EGFP

Proliferating RPCs
Weak mCherry
Strong EGFP

Proliferating RPCs
Faint mCherry
Strong EGFP

B *Tg(pcna:EGFP);Tg(gfap:mCherry)*

> 100000 lux, 30'

Uninjured   44 hpl
4 dpl   6 dpl

mCherry

EGFP

10x Genomics
scRNAseq

ATTCATC
GAATACT

C

mCherry

Uninjured   44 hpl   4 dpl   6 dpl

sorting gate

EGFP

A

1. Non-reactive Müller glia 1
2. Non-reactive Müller glia 2
3. Reactive Müller glia 1
4. Reactive Müller glia 2
5. Müller glia/Progenitors 1
6. Progenitors 2
7. Progenitors 3
8. Photoreceptor precursors
9. Horizontal cell precursors
10. Retinal ganglion cells
11. Red cones
12. Blue cones
13. Amacrine cells 1
14. Amacrine cells 2
15. Bipolar cells

B

Uninjured   44 hpl   4 dpl   6 dpl

C

G1   S   G2/M

D

pcna

Figure 3

A

Müller glia

apoeb

gfap

B

Non-reactive Müller glia

mstnb

inhbaa

C

Reactive Müller glia

crlf1a

her4.1

D

| Uninjured | 44 hpl | 4 dpl | 6 dpl |

ONL

INL

GCL

her4.1

Figure 4

A

	Uninjured	44 hpl	4 dpl	6 dpl
Zrf1	ONL INL GCL			
crlf1a	ONL INL GCL			
PCNA	ONL INL GCL			
Merge	ONL INL GCL	1	2	

B

1 | **Zrf1 PCNA** | **Zrf1 PCNA** *crlf1a*

44 hpl

2 | **Zrf1 PCNA** | **Zrf1 PCNA** *crlf1a*

4 dpl

Figure 5

A

Müller glia/Progenitors 1

B

Figure 6

A

B

Figure 7

A

Progenitors 2 - 3

insm1a

atoh7

onecut1

onecut2

mki67

B

Uninjured	44 hpl	4 dpl	6 dpl

ONL
INL
GCL
atoh7

ONL
INL
GCL
onecut1

ONL
INL
GCL
onecut2

Figure 8

Figure 9

A

Cluster annotation according to Hoang et al., 2020

● Resting Müller glia ● Retinal ganglion cells ● Rods ● Bipolar cells ● Pericytes
● Activated Müller glia ● Horizontal cells ● GABAergic amacrine cells ● Microglia ● Retinal pigmented epithelium
● Progenitors ● Cones ● Glycinergic amacrine cells ● Oligodendrocytes ● Vascular/Endothelial cells

B

Resting Müller glia   Activated Müller glia   Progenitors

Retinal ganglion cells   Cones   GABAergic amacrine cells

Bipolar cells   Horizontal cells   Rods

Figure 10

LEGEND

Non-reactive Müller glia | Reactive Müller glia 1 | Reactive Müller glia 2 | Progenitor 1 | Progenitor 2 | Progenitor 3 | Horizontal cell precursor | Photoreceptor precursor

Retinal ganglion cell | Red cone | Blue cone | Amacrine cell | Bipolar cell | Predicted rod | Predicted horizontal cell

Injury-induced MG trajectory | Injury-induced MG self-renewal trajectory | Injury-induced neurogenic trajectory

Figure 1-figure supplement 1

A

B

Figure 1-figure supplement 2

Figure 1-figure supplement 3

Figure 2-figure supplement 1

A

Figure 2-figure supplement 2

LEGEND

MG 1 = Non-reactive Müller glia 1
MG 2 = Non-reactive Müller glia 2
rMG 1 = Reactive Müller glia 1
rMG 2 = Reactive Müller glia 2
P1 = Müller glia/Progenitors 1
P2 = Progenitors 2
P3 = Progenitors 3
PHP = Photoreceptor precursors
HCP = Horizontal cell precursors
RGC = Retinal ganglion cells
Red C = Red cones
Blue C = Blue cones
AC 1 = Amacrine cells 1
AC 2 = Amacrine cells 2
BP = Bipolar cells

Figure 3-figure supplement 1

LEGEND
- 1. MG 1 = Non-reactive Müller glia 1
- 2. MG 2 = Non-reactive Müller glia 2
- 3. rMG 1 = Reactive Müller glia 1
- 4. rMG 2 = Reactive Müller glia 2
- 5. P1 = Müller glia/Progenitors 1
- 6. P2 = Progenitors 2
- 7. P3 = Progenitors 3

Figure 3-figure supplement 2

Figure 8-figure supplement 1

A

Figure 8-figure supplement 2

A Rod markers

B Amacrine and horizontal cell markers

C Amacrine and horizontal cell markers

D Bipolar cell markers

E General synaptic markers

Figure 9-figure supplement 1