Early whole-body mutant huntingtin lowering averts changes in proteins and lipids important for synapse function and white matter maintenance in the LacQ140 mouse model

Kai Shing[1*], Ellen Sapp[1*], Adel Boudi[1], Sophia Liu[1], Connor Seeley[1], Deanna Marchionini[2], Marian DiFiglia[1], and Kimberly B. Kegel-Gleason[1]

[1]Department of Neurology, Massachusetts General Hospital, Charlestown, MA 02129

[2]CHDI Management/CHDI Foundation, New York, NY 10001

*These authors contributed equally.

Corresponding Author: Kimberly B. Kegel-Gleason

Address: 114 16th Street, Room 2001

MassGeneral Institute for Neurodegeneration (MIND)

Charlestown, MA 02129

Phone: 617-724-8754

Email: kkegelgleason@mgh.harvard.edu

Keywords: Huntington's disease, striatum, lipidomics, transcriptomics, myelin, sphingolipid

ABSTRACT

Background: Expansion of a triplet repeat tract in exon1 of the HTT gene causes Huntington's disease (HD). The mutant HTT protein (mHTT) has numerous aberrant interactions with diverse, pleiomorphic effects. No disease modifying treatments exist but lowering mutant huntingtin (mHTT) by gene therapy is a promising approach to treat Huntington's disease (HD). It is not clear when lowering should be initiated, how much lowering is necessary and for what duration lowering should occur to achieve benefits. Furthermore, the effects of mHTT lowering on brain lipids have not been assessed.

Methods: Using a mHtt-inducible mouse model we analyzed whole body mHtt lowering initiated at different ages and sustained for different time-periods. Subcellular fractionation (density gradient ultracentrifugation), protein chemistry (gel filtration, western blot, and capillary electrophoresis immunoassay), liquid chromatography and mass spectrometry of lipids, and bioinformatic approaches were used to test effects of mHTT transcriptional lowering.

Results: mHTT protein in cytoplasmic and synaptic compartments of the caudate putamen, which is most affected in HD, was reduced 38-52%. Little or no lowering of mHTT occurred in nuclear and perinuclear regions where aggregates formed at 12 months of age. mHtt transcript repression partially or fully preserved select striatal proteins (SCN4B, PDE10A). Total lipids in striatum were reduced in LacQ140 mice at 9 months and preserved by early partial mHtt lowering. The reduction in total lipids was due in part to reductions in subclasses of ceramide (Cer), sphingomyelin (SM), and monogalactosyldiacylglycerol (MGDG), which are known to be important for white matter structure and function. Lipid subclasses phosphatidylinositol (PI), phosphatidylserine (PS), and bismethyl phosphatidic acid (BisMePA) were also changed in LacQ140 mice. Levels of all subclasses other than ceramide were preserved by early mHtt lowering. Pathway enrichment analysis of RNAseq data imply a transcriptional mechanism is responsible in part for changes in myelin lipids, and some but not all changes can be rescued by mHTT lowering.

Conclusions: Our findings suggest that early and sustained reduction in mHtt can prevent changes in levels of select striatal proteins and most lipids but a misfolded, degradation-resistant form of mHTT hampers some benefits in the long term.

BACKGROUND

Huntington's disease (HD) is a heritable neurodegenerative disease caused by a CAG expansion in the

huntingtin (*HTT*) gene. The protein product, huntingtin (HTT), is ubiquitously expressed but enriched in neurons [1,2]

and has been implicated in numerous molecular functions including vesicle trafficking, autophagy, transcription and

DNA repair (Reviewed by [3–5]). HTT has also been shown to have essential functions during development [6–11]. The CAG-

repeat expansion encodes a polyglutamine expansion in HTT, which causes protein accumulation and aggregation

and has pleiomorphic effects that contribute to HD pathology ranging from mitochondrial dysfunction,

transcriptional defects, cholesterol mishandling, altered palmitoylation, and metabolic changes from altered signaling

in the hypothalamus [12,13].

HTT transcription lowering strategies have become central to HD translational studies (reviewed by [14]).

Interventional trials have targeted total *HTT* (that is, normal and mutant *HTT*) using antisense oligonucleotides (ASOs)

(reviewed by [15]) or *HTT* pre-mRNA using an oral drug, branaplam (LMI070) [16]. In animal models (mice, sheep, and

nonhuman primates), additional strategies of gene therapy using transcriptional repressors to target expression from

the mutant allele [17] and modified interfering RNAs (RNAi) [18] or AAV expressing microRNAs (miRNAs) or short hairpin

RNAs (shRNAs) [19,20] to target RNA levels are actively being pursued [17,21–28].

Although lowering total *HTT* in humans was hoped to be generally safe [29], insufficient evidence currently

exists to make this conclusion. There is support for selective lowering of m*HTT* based on data in mice that show loss

of wild type (WT) *HTT* may affect neuronal function including synaptic connectivity [6,8,30,31].

Proteins known to change with HD progression may serve as useful readouts for investigating the efficacy of

HTT lowering. For instance, DARPP32 is enriched in striatal projection neurons and is progressively reduced in HD

patient postmortem brain and in mouse HD models [32]. The cAMP phosphodiesterase PDE10A is reduced early and

sustainably in HD striatum measured both by western blot and mass spectrometry [17,33,34] or ligand binding [17,35].

Microarray analysis in neurons derived from human stem cells [36], mass spectrometry and western blot analyses in

striatal synaptosomes [33] and immunofluorescence (IF) studies in mice [37] showed SCN4B is lowered in HD models.

ATP5A protein levels are altered in numerous mass spectrometry studies [33,38–40].

The intracellular location of HTT positions it to affect lipids in membranes. HTT normally associates with

membranes where it interacts directly with lipid bilayers [41]. Lipids comprise ~50% of the total dry weight in brain and

have diverse cellular roles as membrane structural components, signaling molecules, and sources of energy [42].

Disruptions to lipid composition/metabolism have been linked to disease [43]; thus, their study may provide insight into

disease mechanism or lead to discovery of new biomarkers. We and others have described alterations to lipid levels

in HD mouse models [44–48] and post-mortem HD brain tissue [49–51]. Specific glycerophospholipids showed differential

associations with mHTT [52] and mHTT can alter the biophysical properties of lipid bilayers [41,53] underscoring the

potential for direct consequences of the presence of mHTT on cellular lipids.

The effects of lowering total *Htt* or m*Htt* alone on striatal proteins and behavioral and psychiatric measures

have been investigated in HD mouse models after delivery of reagents to the striatum or lateral ventricle [17,54].

Biomarkers (mHTT levels and aggregation) that are responsive to total or m*Htt* lowering have been investigated in HD

mouse models [35,55,56]. However, the impact of lowering m*Htt* on lipids has not been examined. Here we used the

inducible HD knock-in mouse model, LacQ140, in which the expression of m*Htt* alone is regulated by adding or

withdrawing the lactose analog isopropyl b-D-1-thiogalactopyranoside (IPTG) in their drinking water [57–60], to study the

effects of m*Htt* lowering for different time periods on proteins known to be affected in HD models and using a survey

of lipids. We also tracked changes in levels of soluble and aggregated mHTT protein in different subcellular

compartments. The results show that early and sustained reduction in m*Htt* in this HD mouse model can delay select

protein changes and prevent numerous lipid derangements. Using a systems biology approach, we integrated our

lipidomic findings with a previously generated transcriptomic dataset, identifying dysregulation of pathways relating

to myelin and white matter including genes directly involved in biosynthetic pathways for lipids we found changed in

LacQ140 mice. An SDS-soluble misfolded form of mHTT persists, however, and benefits eventually dwindled.

MATERIALS AND METHODS

Animals

The *LacO/LacIR*-regulatable HD mouse model (LacQ140) was generated by crossing the *Htt^LacQ140/+* mouse to the

*Tg^ACTB-lacI*Scrb* mouse as previously described [58,59]. The default state of the LacQ140 mouse is global repression of m*Htt*

due to *Lac* Repressor binding to the *Lac* operators. The continuous administration of IPTG starting from E5 interrupts

the binding between the *Lac* repressor and operators, resulting in a de-repressed state, and maximal expression of

m*Htt* in LacQ140. All WT mice were Htt^LacO+/+; b-actin-LacI^R tg. Mice were provided with enrichment (envirodry, play

100 tunnels, Bed-o'cobs and plastic bones) and housed uniform for genotype, gender, and treatment. Mice were fed *ad*

101 *libitum*. The lactose analog IPTG was provided in drinking water (at 10mM) which de-represses the *LacQ140* allele

102 and keeps normal m*Htt* expression. During embryonic development, m*Htt* expression levels were maintained at

103 normal levels by administering IPTG to pregnant dams starting at embryonic day 5 (E5). IPTG was administered never

104 (*mHtt* repressed), always (*mHtt* always expressed) or withdrawn at 2 or 8 months (*mHtt* expressed normally then

105 repressed at 2 or 8 months). The CAG repeat length range in $Htt^{LacO-Q140/+}$ mice was 143-157 with average of 148 and

106 median of 148 CAG.

107 **Sample preparation**

108 The striatum from one hemisphere for each mouse was homogenized in 750 µl 10mM HEPES pH7.2, 250mM sucrose,

109 1uM EDTA + protease inhibitor tablet (Roche Diagnostics GmbH, Mannheim, Germany) + 1mM NaF + 1mM Na3VO4.

110 A 150 µl aliquot of this crude homogenate was removed and protein concentration was determined using the

111 Bradford method (BioRad, Hercules, CA). Subcellular fractionation by density gradient ultracentrifugation using

112 Optiprep was performed on remaining 600 µl sample for the 6- and 12-month-old mice as previously described [47].

113 **Capillary immunoassay**

114 Equal amounts of protein from the crude homogenates were analyzed using the automated simple western system,

115 Wes (ProteinSimple, Bio-Techne, San Jose, CA), which utilizes a capillary-based immunoassay. The protocol described

116 in the manual was followed to detect HTT, GFAP and DARPP32 using 0.6 µg of sample. Quantitative analysis of

117 protein levels is done automatically using the Compass for Simple Western Software (ProteinSimple) on

118 electropherogram data. The peak area (using automatic "dropped line" option in software) of each protein of interest

119 was normalized to the peak area of the vinculin loading control. Figures show protein bands similar to traditional

120 western blots using "lane view" option in the Compass software to create a blot-like image from the

121 electropherogram data.

122 **Western blot analysis**

123 Equal amounts of protein from the crude homogenates were analyzed by western blot for levels of HTT and other

124 proteins of interest as previously described [33]. Briefly, 10 µg of protein were separated by SDS-PAGE, transferred to

125 nitrocellulose, and probed with primary antibody overnight. Nitrocellulose membranes were cut into horizontal strips

to maximize the number of antibodies that could be probed on one blot. Peroxidase labeled secondary antibodies

were used with the SuperSignal West Pico Chemiluminescent substrate (ThermoScientific, Rockford, IL, #34580) and

digital images were captured with a CCD camera (AlphaInnotech, Bayern, Germany). For western blot analysis of

subcellular fractions, equal volumes of each sample (15 µl) were separated by SDS-PAGE. Pixel intensity

quantification of the western blot signals on the digital images was determined using ImageJ software (NIH) by

manually circling each band and multiplying the area by the average signal intensity. The total signal intensity was

normalized to vinculin or GAPDH loading controls.

Antibodies

The following antibodies and dilutions were used in this study: Anti-HTT Ab1 (aa1-17,[1]) 1:50 for capillary

immunoassay and 1:2000 for western blot; anti-HTT EPR5526 (Abcam, Waltham, MA, ab109115, 1:2000 for western

blot); anti-polyQ MW1 (MilliporeSigma, Burlington, MA, MABN2427, 1:50 for capillary immunoassay); anti-polyQ

PHP3 (generous gift from Dr. Ali Khoshnan, 1:2000 for western blot); Anti-PDE10A (Abcam, Waltham, MA,

#ab177933, 1:2000 for western blot); Anti-DARPP32 (Abcam, #ab40801, 1:2000 for capillary immunoassay); Anti-

GFAP (MilliporeSigma, Burlington, MA, AB5804, 1:3000 for capillary immunoassay); Anti-GAPDH (MilliporeSigma,

Burlington, MA, #MAB374, 1:10000 for western blot); Anti-Sodium channel subunit beta-4 (Abcam, Waltham, MA,

#ab80539, 1:500 for western blot); Anti-vinculin (Sigma, St. Louis, MO, #V9131, 1:5000 for capillary immunoassay,

1:2000 for western blot); Anti-ATP5A (Abcam, Waltham, MA, #ab14748, 1:2000 for western blot); Anti-HTT MW8

(University of Iowa Developmental Studies Hybridoma Bank, 1:1000 for filter trap); Anti-HTT S830 (generous gift from

Dr. Gillian Bates, [61] 1:8000), HDAC1 (Abcam, Waltham, MA, ab32369-7, 1:4000).

Filter trap assay

Based on protocol described in [62,63], equal protein amounts for each sample (40 µg) were brought up to 50 µl volume

with PBS and 50 µl 4% SDS in PBS was added to each sample to make final concentration 2% SDS. A cellulose acetate

membrane was wet in 2% SDS/PBS and placed in dot blot apparatus. The 100 µl samples were added to each well

and pulled through the membrane with a vacuum then washed 3 times with 200 µl 2% SDS/PBS. The membrane was

removed from the apparatus, washed in Tris buffered saline + 0.1% Tween-20 (TBST) then processed for western blot

using MW8 or S830 antibodies. The total signal intensity of each dot was measured in ImageJ by circling the entire

dot and multiplying the area by the average signal intensity minus the background signal from an empty dot.

Statistical analysis

One-way ANOVA with Tukey's multiple comparison test was performed to determine significance between groups.

Asterisks on graphs show p values and are described in the figure legends.

Lipid extraction

Lipids were extracted using methyl tert-butyl ether (MTBE) as previously described and analyzed using ion switching

and molecular assignment as previously described [47,64,65]. Each age group was processed together. Crude

homogenates (100 µl) of dissected mouse striatum were transferred into 20 ml glass scintillation vials. 750 µl of HPLC

grade methanol was added to each sample, then vials were vortexed. 2.5 ml of MTBE was then added to each sample

and incubated on a platform shaker for 1 hour. After incubation, 625 µl of water was added to induce polar and non-

polar phase separation. The non-polar lipid containing (upper) phase was collected into a new vial, and the polar

(lower) phase was subsequently re-extracted with 1 ml of MTBE/methanol/water (10/3/2.5, v/v/v). Following re-

extraction, the lipid containing phases were combined and allowed to dry on a platform shaker, then further dried

with nitrogen gas. Extracted lipids were hand delivered to the Beth Israel Deaconess Medical Center Mass

Spectrometry Core Facility.

Lipid Annotation

Data for each timepoint was classified by LIPID MAPS category[66]: glycerophospholipids, glycerolipids, sphingolipids,

sterol lipids, fatty acyls, and prenol lipids were detected. Each category contains distinct subclasses as annotated

below. _Glycerophospholipids:_ Phosphatidylcholine (PC), Phosphatidylethanolamine (PE), Phosphatidylserine (PS),

Phosphatidylinositol (PI), Methylphosphocholine (MePC), Phosphatidic acid (PA), Bis-methyl phosphatidic acid

(BisMePA), Dimethyl phosphatidylethanolamine (dMePE), Phosphatidylgylcerol (PG), Bis-methylphosphatidylserine

(BisMePS), Bis-methyl phosphatidyl ethanolamine (BisMePE), Cardiolipin (CL), Phosphatidylethanol (PEt), Biotinyl-

phosphoethanolamine (BiotinylPE), Phosphatidylmethanol (PMe), Phosphatidylinositol-bisphosphate (PIP2),

Phosphatidylinositol-monophosphate (PIP), Lysophosphatidylcholine (LPC), Lysophosphatidylethanolamine (LPE),

Lysophosphatidylserine (LPS), Lysophosphatidylinositol (LPI), Lysophosphosphatidylgylcerol (LPG), Lysodimethyl

phosphatidyl ethanolamine (LdMePE). *Glycerolipids:* Triglyceride (TG), Monogalactosyldiacylglycerol (MGDG),

Monogalactosylmonoacylglycerol (MGMG), Diglyceride (DG), Sulfoquinovosylmonoacylglycerol (SQMG),

Sulfoquinovosyldiacylglycerol (SQDG). *Sphingolipids:* Hexosylceramides (Hex1Cer), Simple Glc series (CerG1),

Sphingomyelin (SM), Ceramide (Cer), Ceramide phosphate (CerP), Sulfatide (ST), Sphingoid base (So), Sphingomyelin

phytosphingosine (phSM), Simple Glc series (CerG2GNAc1), Ceramide phosphorylethanolamine (CerPE), Sphingosine

(SPH), Dihexosylceramides (Hex2Cer). *Sterol lipids*: Cholesterol ester (ChE), Zymosterol (ZyE). *Fatty acyls:* Fatty acid

(FA), Acyl Carnitine (AcCa). *Prenol lipids:* Coenzyme (Co).

Individual lipid species were annotated according to sum composition of carbons and double bonds in the format

Lipid Subclass (total number of carbons: total number of double bonds). If distinct fatty acid chains could be

identified, they were annotated separated by an underscore (ex. PC 32:1, or PC (16:0_16:1). Using this approach, we

cannot determine the *sn*-1 or *sn*-2 positions of the fatty acid chains. Lipid species within the sphingolipid category

contain prefixes 'd' or 't' to denote di-hydroxy or tri-hydroxy bases. For example, SM(d18:1_23:0) contains 2 hydroxyl

groups. The Hex1Cer subclass is comprised of both glucosylceramide (GlcCer) and galactosylceramide (GalCer); the

orientation of one of the hydroxyl groups in Glc differs from in Gal, and thus cannot be resolved by these methods [67].

Plasmanyl lipid species (ether linked) are annotated by 'e' and plasmenyl/plasmalogen (vinyl ether linked) lipid

species are annotated by 'p' (ex. PC (36:5e) or PE (16:0p_20:4) [68].

Lipidomics statistics and data visualization

Heatmap and hierarchical clustering in Figure 4 were generated using Morpheus from the Broad Institute

(Cambridge, MA, https://software.broadinstitute.org/morpheus). Hierarchical clustering was performed across all

rows (lipid subclasses) using the one minus Pearson correlation distance metric. Rows determined to be the most

similar are merged first to produce clusters, working iteratively to merge rows into clusters. The dendrogram displays

the order of clustering with the most similar rows shown in closest proximity. Lipid expression values are assigned to

colors based on the minimum (blue, low relative expression) and maximum (red, high relative expression) values for

each independent row. Heatmap in Figures 4 & 5 was generated in R using the ComplexHeatmap package v2.16.0 [69].

Each column represents data from one animal. Statistical significance was determined by one-way analysis of

variance (ANOVA) with Tukey's multiple comparison test between treatment groups for lipid subclasses (6mo: N=36,

9mo: N=24, 12mo: N=29) or lipid species (6mo: N=800, 9mo: N=632, 12mo: N=735). The two-stage linear step-up

procedure of Benjamini, Krieger, and Yekutieli was applied to ANOVA p-values to control the false discovery rate

(FDR) with significance accepted at q<0.05 (Q=5%).

RNA-sequencing analysis

The raw counts matrix derived from striatal samples in the LacQ140 mouse model was accessed from Gene

Expression Omnibus (GEO GSE156236). The 6-month cohort included n=10 mice per group (WT, LacQ140,

LacQ140_2M, LacQ140_A) and the 12-month cohort included n=10 WT, n=9 LacQ140, n=9 LacQ140_8M, n=10

LacQ140_2M, and n=10 LacQ140_A. Analysis was performed using R v4.3.0 in the RStudio IDE (2023.03.0+386).

Transcripts with fewer than 10 total counts were removed from the analysis. Differential expression analysis was

performed using DESeq2 v1.40.1 [70]. P-values were computed with the Wald test and were corrected for multiple

testing using the Benjamini and Hochberg procedure [71]. Genes with a fold change greater than 20% in either direction

and with an adjusted p-value < 0.05 were considered differentially expressed. Functional enrichment analysis was

conducted using the ClusterProfiler package v4.8.1. Over-representation analysis was performed for upregulated or

downregulated DEGs with the enrichGO function (org.Mm.eg.db v3.17.0). Gene set enrichment analysis was

performed with the gseGO function using a ranked list of genes (Wald statistic) for each comparison (LacQ140 vs WT,

LacQ140_8M vs WT, LacQ140_2M vs WT, and LacQ140_A vs WT) at 6 and 12 months. Heatmaps were plotted using

ComplexHeatmap package v2.16.0 with rows representing individual animals and rows representing genes.

RESULTS

Time course of mHTT protein lowering with regulated transcriptional repression using LacQ140 mouse striatum

We used the inducible HD knock-in mouse model, LacQ140, in which the expression of m*Htt* throughout the

body was regulated by adding or withdrawing the lactose analog isopropyl b-D-1-thiogalactopyranoside (IPTG) in

drinking water using an established *Lac* operator/repressor system (**Fig. 1a**) [59,60]. We used the model to lower m*Htt*

from conception, starting at 2 or 8 months of age. To control for the effects of IPTG, WT mice also received IPTG

treatment over their lifetime. The striatum was examined at 6, 9, and 12 months of age (**Fig. 1b**). Antibodies against

varied epitopes within HTT detect different pools of normal and mutant HTT by immunofluorescence and

immunohistochemistry [1,72–75]. Numerous conformations (over 100 distinct classes) of purified HTT and mHTT have

been identified by electron microscopy in vitro [76]. Polyglutamine (polyQ) expansion induces confirmational changes

particular to mHTT detectible even after SDS-PAGE and/or western blot with several antibodies to regions in the

poly-Q region (MW1 [77,78] and 3B5H10 [79]) and in the adjacent polyproline region (PHP1-4 [80] and S830 [81]). To assess

lowering, full length HTT and mHTT proteins were detected by capillary immunoassay using anti-HTT antibody Ab1

targeting aa 1-17 and anti-polyQ antibody MW1 (**Fig. 2**), and by western blot using anti-HTT antibody EPR5526, and

anti-mHTT antibody PHP3, which recognizes an altered conformer of polyproline in mHTT [80] (**Fig. S1**).

Overall, results showed lowering of 35-52% of mHTT detected with Ab1, MW1 and PHP3 in 6- and 9-month

mice with transcriptional repression **(Fig. 2a, Fig. S1a)** and **(Fig. 2b, Fig. S1b)**. Although EPR5526 recognizes both WT

and mHTT, no significant lowering of mHTT protein was measured with mHtt gene repression using this antibody in

6-month mice. In the 12-month mice, mHTT was significantly reduced in LacQ140_A (51%) mice compared to

LacQ140 mice (**Fig. 2c**). However, this was observed only in the capillary immunoassay using antibody Ab1 but not

with antibodies MW1, EPR5526, or PHP3 **(Fig. S1c)**. m*Htt* repression in the LacQ140_2M and LacQ140_8M mice

yielded no significant reduction of mHTT protein levels compared to LacQ140 mice using any antibody. These results

show that systemic regulated repression of m*Htt* transcription in LacQ140 mice results in partial lowering of mHTT

protein levels in the striatum (38-52%), but a soluble form of full-length mHTT remains as mice age.

Effects of m*Htt* lowering on its distribution in different subcellular cytoplasmic compartments

We next looked at the effects of m*Htt* repression on its protein levels in different subcellular compartments.

Density gradient fractionation and ultracentrifugation for subcellular fractionation of cytoplasmic components was

performed as shown in **Fig. S2a**. The schematic in **Fig. S2b** indicates representative proteins that are enriched in

different cytoplasmic compartments. In 6-month-old mice, the mHTT/WT HTT ratio was significantly lower in

LacQ140_2M mice in fractions 13 and 14 compared to LacQ140 mice (**Fig. S2c and d**). Similar results were observed

in 12-month-old mice, where the mHTT/WT HTT ratio in fractions 1, 3, 13, and 14 in LacQ140 mice was significantly

lower with m*Htt* repression for different periods of time compared to LacQ140 mice (**Fig. S2e and f**). There was no

change in the distribution of HTT and mHTT in the fractions between groups in 6-month-old mice **(Fig. S3a and b)** or

in 12-month-old mice **(Fig. S3c and d)**. Altogether, these results show that mHTT is lowered in cytoplasmic fractions through 12 months.

Effects of m*Htt* lowering on its distribution in crude nuclear fractions

In 12-month-old LacQ140 mice, repressing m*Htt* transcription only partially reduced levels of mHTT protein in crude homogenates (**Fig. 2c**) even though mHTT was efficiently lowered in the sub cytoplasmic compartments contained in the S1 fraction (**Fig. S2f**). We speculated that mHTT may reside in other compartments where it is more resistant to removal by transcript repression. To address this idea, we examined P1 fractions which contain nuclei, ER, large perinuclear structures such as the recycling compartment, some mitochondria and autophagosomes [82]. HTT was detected with antibody Ab1 in WT mice (2 alleles worth) and LacQ140 mice (1 allele worth) and mHTT (1 allele worth) in LacQ140 mice at both 6 and 12 months (**Fig. 3a and b**). Significant lowering of mHTT protein was observed in 6-month-old LacQ140_A but not LacQ140_2M mice using antibody Ab1 (**Fig. 3a**). In 12-month mice, repression of the m*Htt* allele at any age or duration failed to lower mHTT protein levels in the P1 fraction (**Fig. 3b**).

We queried whether forms of mHTT with altered migration with SDS-PAGE could be detected in the P1 fractions using antibody S830, which has been reported by us and others to detect a smear using SDS-PAGE and western blot [61,83]. At 12-months, HD mice showed an S830-positive smear above the HTT/mHTT bands which was not lowered in the LacQ140_8M, LacQ140_2M, or LacQ140_A mice (**Fig. 3c**). Filter trap assay showed lowering of an SDS-insoluble aggregated species (recognized by S830) in both crude homogenate and P1 fractions; slightly less lowering of aggregated mHTT occurred in the P1 fraction when transcript repression was initiated at 2 or 8 months (**Fig. S4a and b**). Altogether, these results show that a soluble species of mHTT in the P1 fraction is more resistant to lowering after transcriptional repression.

Effects of m*Htt* lowering on levels of GFAP, DARPP32, SCN4B, PDE10A, and ATP5A

Prior studies in different mouse models of HD have shown that levels of some neuronal proteins are altered in the mouse striatum, namely DARPP32, PDE10A, SCN4B, and ATP5A [32–34,37–40,84]. To assess levels of these proteins and that of the astrocyte protein, glial acidic fibrillary protein (GFAP), in LacQ140 mice without and with m*Htt* gene repression, crude homogenates from 6, 9, and 12-month-old mice were analyzed by capillary immunoassay or western blot.

In agreement with previous studies in HD mouse models [17,33,34] striatum from the LacQ140 exhibited a

significant reduction in PDE10A at all ages examined. Early m*Htt* lowering starting at 2 months of age statistically

preserved PDE10A levels at 6 and 9 months (**Fig. S5a and b**), but the effect was lost by 12 months of age when

transcript repression was initiated at 2 or 8 months (**Fig. S5c**). Transcript repression initiated embryonically

(LacQ140_A) and examined at 6 and 12 months preserved PDE10A levels, although no protection was observed at 9

months. SCN4B expression was reduced in the striatum of LacQ140 at 6 and 9 months of age and levels were

preserved with early m*Htt* lowering (**Fig. S5a and b**). DARPP32 levels were significantly lower in LacQ140 compared

to WT mice at 9 months but not at 6 or 12 months, and there were no differences in levels of GFAP and ATP5A

between WT and LacQ140 at any age examined (**Fig. S6a-c**).

Effects of m*Htt* lowering on lipids detected by mass spectrometry

We surveyed for lipid changes in LacQ140 striatum compared to WT and the effects of lowering m*Htt* using

liquid chromatography and mass spectrometry (LC-MS) as previously described [47]. For each age, lipids were extracted

from crude homogenates of striatum from each treatment group/genotype and analyzed as a set. The total lipids per

group were compared. Our MS intensity measurements were relative measurements so only samples processed

together can be compared (i.e., by age group). The sum of lipids for each genotype and/or treatment group were

reported as a proportion of WT within each age group (**Fig. 4a-c**). No changes in total lipid were observed at 6

months or 12 months (**Fig. 4a and c**). However, at 9 months LacQ140 mice had significantly lower levels of total lipids

compared to WT or to LacQ140_A mice **(Fig. 4b)**. A heat map and hierarchical clustering of lipid changes by subclass

at 9 months revealed two major groups where-in subclasses moved in the same direction even if all weren't

statistically significant (**Fig. 4d**). The top cluster delimited in blue shows subclasses that decreased in LacQ140 mice

compared to WT and were corrected by lowering. In contrast, the cluster marked in red shows subclasses that were

increased in LacQ140 mice compared to WT and were improved by m*Htt* lowering**.** Other subclasses marked in black

did not change. Summaries of lipid subclasses and number of species detected for each age group are in **Table 1,** and

an overview of changes in individual species across ages is shown in **Table 2**. Data and graphs for 6-month-old mice

subclasses and species can be found in **Fig. S7 & 8** and **Additional File 1**; 9-month-old mice in **Fig. S9 & 10** and

Additional File 1; and 12-month-old mice in **Fig. S11 & 12** and **Additional File 1**.

We assessed if there were more subtle alterations at the subclass or individual species level that did not influence the overall lipid levels. In striatum of LacQ140 mice at 6-months-of-age, we detected a modest increase in the subclass phosphatidylinositol (PI) compared to WT (**Fig. S7**). Individual species of PI (18:0_20:4, 16:0_22:6), and phosphatidylserine PS (18:0_20:4, 22:6_22:6) were increased in LacQ140 mice compared to WT (**Fig. S8**). However, no subclass or species changes survived correction using the Benjamini, Krieger, and Yekutieli procedure with a 5% false discovery rate (FDR) of q <0.05, N=36 subclasses (**Table 1**) and N=800 species (**Additional File 1**). Consistent with our observations at 6 months, PS and PI were increased in 9-month-old LacQ140 mice compared to WT (**Fig. 5a and b**). A limited number of species are modestly increased at 6 months, which progresses to more robust increases in these glycerophospholipids at the subclass level at 9 months. A significant reduction in bismethyl phosphatidic acid (BisMePA) also occurred (**Fig. 5c**). Functionally, PI is the precursor for PIPs which are important for protein kinase c (PKC) signaling at the synapse [85] and can also act as important docking and activating molecules for membrane associated proteins, including HTT [52]. PS is an abundant anionic glycerophospholipid necessary for activation of several ion channels [86], fusion of neurotransmitter vesicles, regulation of AMPA signaling, and coordination of PKC, Raf-1, and AKT signaling [87].

In contrast to the increases observed in glycerophospholipids, 9-month-old LacQ140 mice exhibit significant reductions in the subclasses sphingomyelin (SM) and ceramide (Cer) (**Fig. 5d and e),** lipids central in sphingolipid metabolism and important for myelination. Cer is the backbone for SM as well as more complex glycosphingolipids, which are highly enriched in myelin [88]. Moreover, the low abundance glycerolipid monogalactosyldiacylglycerol (MGDG) was strikingly reduced at 9 months (**Figure 5f**). MGDG regulates oligodendrocyte differentiation and is considered a marker of myelination [89,90].

All subclasses (PI, PS, BisMePA, SM, Cer, MGDG) with changes between WT and LacQ140 had q values<0.05 (N=24 subclasses) (**Table 1; Fig. 4d**). Early lowering of m*Htt* (LacQ140_A) restored levels of each of these subclasses except for Cer and lowering m*Htt* starting at 2 months (LacQ140_2M) was sufficient to correct levels of SM and MGDG (**Fig. 5a-f).** To assess the effects of m*Htt* lowering on the distinct lipid species that comprise each subclass, lipid species were plotted in a heatmap **(Fig. 5g).** The broad changes at the subclass level can be attributed to large numbers of individual lipid species changing within each subclass even if each species is not significantly different (**Fig. 5g & Fig. S9**). Alternatively, subclasses can remain unchanged but still contain individual lipid species with

differences. For example, the subclass Hex1Cer was not significantly changed **(Fig. S9)**, but many individual species were significantly altered in LacQ140 mice **(Fig. S10).** These species included 3 that were significantly decreased in LacQ140 mice and normalized to WT levels with lowering of m*Htt*: Hex1Cer (d18:1_18:0), Hex1Cer (t18:0_18:0), Hex1Cer (d18:1_25:1) **(Fig. S10).** Combined, these 3 species represent approximately 1% of total Hex1Cer detected. Hex1Cer is comprised of galactosylceramide and glucosylceramide and cannot be distinguished by LC-MS/MS; however, previous studies have determined that the majority of Hex1Cer in the brain is galactosylceramide, which is highly enriched in myelin [67,91]. Galactosylceramides are major constituents of the myelin sheath and contribute to its stabilization and maintenance [92].

In total, we identified 14 lipid species significantly altered between LacQ140 and WT mice using a stringent FDR cutoff (FDR 5%/q<0.05, Benjamini, Krieger, and Yekutieli procedure, N=632 lipids) and all 14 showed improvement with m*Htt* lowering at 9 months (**Fig. S10**). This included increases in PI (16:0_20:4) & PI (18:1_20:4) and PS (18:0_20:4), PS (18:0_18:1) and PS (18:1_22:6). Lipid species decreased in LacQ140 mice and restored with m*Htt* lowering included Hex1Cer species as detailed above, SM (d40:2), PE (16:0p_20:4) and MGDG (16:0_18:1) (**Fig. 5g, Fig. S10)**. Taken together, these results indicate a profound disruption of myelin lipids in LacQ140 mice, which is attenuated with lowering of m*Htt.*

In the striatum of LacQ140 mice at 12 months, no significant differences at the subclass level were found compared to WT (**Table 1; Fig. S11**) although 29 subclasses were measured. At the individual lipid species level, we identified 26 lipid species significantly altered between LacQ140 and WT mice (FDR 5%/q<0.05, Benjamini, Krieger, and Yekutieli procedure, N=735 lipids). (**Fig. S12**). Contrary to the reversal of lipid dysregulation observed at 9 months, the only lipid species that was responsive to lowering m*Htt* at 12 months was CerP (t40:3), which was increased in LacQ140 and decreased with early m*Htt* lowering (LacQ140_A) (**Fig. S12**). Myelin enriched lipids, including 6 species of Hex1Cer (d18:1_18:0, d:40:2, d18:2_22:0, d41:2, t18:0_22:1 & d18:1_22:1) and ST (d18:1_22:1), decreased and were not improved with m*Htt* lowering (**Fig. S12**). Similarly, nine triacylglycerol (TG) species, eight of which contain oleic acid (C18:1) were decreased in LacQ140 mice and unchanged with m*Htt* lowering (**Fig. S12**). Overall, these results demonstrate that most individual lipid changes were no longer reversible by 12 months.

Bioinformatic analysis of transcriptional alterations to lipid metabolism and myelination

To determine if any of the lipid changes could be explained by altered transcription of genes regulating lipid-related metabolic pathways, we re-analyzed the previously generated RNAseq dataset in the LacQ140 striatum (GEO GSE156236). Differential expression analysis [70] detected 1360 upregulated and 1431 downregulated genes at 6 months and 1375 upregulated and 1654 downregulated genes at 12 months (FC>20%, adjusted p-value <0.05) **(Fig. S13a).** Over-representation analysis against GO Biological Processes (GO BP) revealed that downregulated genes were enriched in terms related to lipid metabolism and signaling as well as myelination. In 6-month LacQ140 downregulated genes, GO BP terms ensheathment of neurons (padj<0.01), axon ensheathment (padj<0.01), myelination (padj<0.05), cellular lipid metabolic process (padj<0.05), lipid metabolic process (padj<0.05), phospholipase C-activating G protein-coupled receptor signaling pathway (padj<0.05), response to lipid (padj<0.05) were significantly over-represented **(Fig. 6a)**. In LacQ140 upregulated genes, there was no over-representation of terms that could directly explain changes to lipid levels **(Additional File 2).**

Next, Gene Set Enrichment Analysis (GSEA) was conducted [93,94] to examine the impact of m*Htt* repression on the enrichment profiles of terms related to lipid metabolism and myelination. At 6 months, 3 of the top 20 enriched GO BP terms were related to myelination and all were negatively enriched: ensheathment of neurons, myelination, and axon ensheathment **(Additional File 2)**. Further, every significantly enriched myelin or lipid related GO BP term was negatively enriched in LacQ140 compared to WT (padj <0.05) (**Fig. 6b, Additional File 2)**. Repression of m*Htt* in LacQ140_2M and LacQ140_A groups was sufficient to reverse the negative enrichment of many GO BP terms including myelination/axon ensheathment/ensheathment of neurons, and sphingolipid/phospholipid metabolic processes suggesting some beneficial impact of m*Htt* lowering (**Fig. 6b**).

Specifically, 6-month LacQ140 mice had reduced expression of genes encoding proteins involved in oligodendrocyte development (*Akt2, Bcas1, Ptgds, Cldn11, Trf, Tcf7l2, Qki*), myelin structure and compaction (*Mbp, Plp1, Tspan2, Mal*), and myelin lipid biosynthesis (*Ugt8a, Fa2h, Aspa*) (**Fig. 6c**). LacQ140 mice also exhibited dysregulation of genes encoding enzymes including phospholipases (*Pla2g7, Pla2g4c, Pla2g4e, Plb1, Plcd3, Plcd4*), phosphatases & kinases (*Plpp1, Plpp3, Plppr2, Inpp4b, Pik3c2b*), and regulators of phospholipid biosynthesis (*Gpat2, Gpat3, Chpt1*) (**Fig. 6c**). Enzymes responsible for biosynthesis and metabolism of sphingolipids were altered in LacQ140 mice: *Sptssb, Ormdl2, Gba2, Asah,* were downregulated and *Smpdl3b* was upregulated. Genes encoding

enzymes responsible for fatty acid elongation (*Elovl1, Elovl5, Elovl6*) were all downregulated (**Fig. 6c**). Of the DEGs

related to lipid metabolism and myelination 23 were reversed in LacQ140_2M mice, 23 were reversed in LacQ140_A

mice and 15 were reversed in both (padj < 0.05, FC > 20% opposite direction of LacQ140) (**Fig. 6c & Fig. S13b).**

Three of the nine GO BP terms significantly over-represented in 6-month LacQ140 downregulated genes

were also over-represented in 12-month downregulated genes: ensheathment of neurons (padj <0.01), axon

ensheathment (padj <0.01), myelination (padj <0.05), and phospholipase C-activating G protein-coupled receptor

signaling pathway (padj<0.01) (**Fig. 6d**). Similarly, GSEA showed that only a subset (8/22) of the GO BP terms related

to lipid metabolism or myelination were enriched at both timepoints (**Fig. 6e**). Terms including glycerophospholipid

metabolic process and glycerolipid metabolic processes showed improvement with lowering of m*Htt*, whereas terms

related to myelination remained negatively enriched with m*Htt* repression (**Fig. 6e**).

DEGs associated with myelin and lipid biosynthesis were both downregulated and upregulated in 12-month

LacQ140 mice. LacQ140 mice had reduced (*Cntnap1, Mal, Tspan2, Plp1, Mobp*) and increased (*Omg, Opalin*)

expression of genes encoding proteins involved in myelin structure and compaction (**Fig. 6f**). Similarly,

oligodendrocyte development related genes were both decreased (*Akt2, Cldn11, Myt1l*) and increased (*Prkcq, Olig1,*

Sox11). Genes related to myelin lipid biosynthesis (down: *Ugt8a, Fa2h, Selenoi*), phospholipid synthesis/metabolism

(up: *Chkb, Agpat2, Mboat2,* down: *Agpat1, Lpcat4*), phosphatases (up: *Plpp1, Plpp2, Plppr2,* down: *Plpp3, Plpp6,*

Inpp5a), phospholipases (up: *Pla2g4e,* down: *Plcb1, Plcb3, Plce1, Plcl1*) and sphingolipid synthesis/metabolism (up:

Cers4, Smpdl3b, Acer3, down: *Gba2, St3gal2, Smpd3, Sgpp2, Sgms1, A4galt, Fads3*) were differentially expressed in

both directions (**Fig. 6f)**. In 12-month LacQ140 mice, glycerolipid synthesis/metabolism genes (up: *Dgat1,* down:

Dgat2, Dgat2l6, Lpin3, Dagla) and DAG kinase genes (up: *Dgka, Dgkg,* down: *Dgkb, Dgki*) were differentially

expressed (**Fig. 6f**). Consistent with the enrichment results at this timepoint, fewer genes were reversed with

repression of m*Htt* (5 genes LacQ140_2M, 4 genes LacQ140_A) (**Fig. 6f & Fig. S13c**).

Differentially expressed genes (DEGs) were also manually curated for genes that could directly impact levels

of PS, PI, sphingolipids, and glycerolipids. Phosphatidylserine synthase enzymes (*Ptdss1/Ptdss2*) [87] and rate limiting

enzymes for PI biosynthesis (CDP-Diacylglycerol Synthases: *Cds1/Cds2*) [95] were unchanged in 6- and 12-month

LacQ140 mice **(Additional Files 1 & 2)**. For glycerolipids, the only known enzymes that catalyze TG biosynthesis [96]

were increased (*Dgat1*) and decreased (*Dgat2*) at 12 months; mRNA levels remained unchanged with m*Htt*

repression **(Fig. 6f).**

Lipid changes might be explained by altered cellular composition of tissue. To address this possibility, we

surveyed for changes in transcript levels for cellular markers. No change in mRNA expression levels for the microglial

markers *Iba1* and *Cd68* or for the reactive astrocyte marker *Gfap* were observed that might indicate upregulation of

these cell types could account for the lipid changes. However, our transcriptional profiling did reveal an expression

signature consistent with altered oligodendrocyte development and alterations to myelin structure or compaction in

both 6- and 12-month-old mice **(Fig. 6c and f)**. UDP-galactose-ceramide galactosyltransferase (*Ugt8a*) is expressed in

oligodendrocytes where it catalyzes the formation of GalCer (Hex1Cer) from Cer; *Ugt8a* mRNA was decreased at both

6 and 12 months in the LacQ140 mice **(Fig. 6c & Fig. 7a).** Embryonic repression of m*Htt* and repression beginning at

2-months of age preserved *Ugt8a* mRNA measured at 6-months; however, the effect was lost by 12-months of age

(Fig. 6c and f). These mRNA results mirror our findings of Hex1Cer lipid levels measured using LC-MS in LacQ140

striatal lysates. GalCer and GlcCer are both members of the Hex1Cer subclass but cannot be distinguished by MS

since they are isomers with the same mass and charge [67]; the gene catalyzing the transfer of glucose to ceramide

(*Ugcg*) was not changed at the mRNA level, suggesting lower levels of Hex1Cer in LacQ140 mice are due to lower

levels of GalCer. Ugt8a can also transfer galactose to diacylglycerol to form the lipid MGDG [97] which we found was

decreased with m*Htt* expression and improved with m*Htt* lowering **(Fig. 5f and g).** In contrast, *Fa2h* mRNA was

reduced at 6 and 12-months but not improved with *mHtt* lowering **(Fig. 6c and f).** Fatty acid 2-hydroxylase, which is

encoded by *Fa2h,* modifies fatty acids substrates with a hydroxyl (-OH) group, producing 2-hydroxylated fatty acids

that are incorporated into myelin sphingolipids and provide stability[98] **(Fig. 7b and c)**. Overall, these data reveal

transcriptomic changes in key lipid metabolic and biosynthetic genes in LacQ140 striatum, which directly impact the

lipidomic profile. Critically, many of these lipidomic and transcriptomic alterations can be attenuated with lowering

of m*Htt*.

DISCUSSION

Here we used the LacQ140 inducible HD mouse model to initiate whole body m*Htt* reduction at different

ages and evaluate effects on proteins and lipids. Lowering mHTT protein by 38-52% in the LacQ140 caudate putamen,

starting from conception up to 12 months of age, was sufficient to prevent m*Htt* induced changes in the levels of

some proteins and some lipids. However, a resistant soluble species of the protein detected in older mice limited

long term benefit of m*Htt* lowering. Our lipid data show clear evidence of changes impacting myelin which

mechanistically is due in part to aberrant transcription as evidenced by differentially expressed genes regulating

some of the same lipids that were found altered. The correction of lipid changes we identified here correlates with

the behavioral changes measured previously in LacQ140 mice using a comprehensive set of unbiased, high-content

assays [59] in that behavior was closer to normal at 6 and 9 months with early mHTT lowering, but the protective effect

of lowering degraded by 12 months.

We identified forms of mHTT protein that were resistant to m*Htt* lowering detected by various HTT

antibodies. These resistant forms of mHTT identified by immunoblot may correspond to mHTT aggregates or foci

found to be resistant to lowering in LacQ140 mice using MSD and immunofluorescence methods [59]. Others have

described aggregated and soluble forms of mHTT that resist degradation in distinct cellular compartments, including

full length mHTT in the nucleus [17,99–101]. In the LacQ140 striatum, the SDS-soluble degradation-resistant form of full-

length mHTT, detected by us using antibodies S830, MW1 and EPR5526, resides in a perinuclear or nuclear

compartment which might correspond to the "juxtanuclear quality control compartment (JUNQ)" described by

Frydman and colleagues [102]. Accrual of misfolded mHTT in the nucleus likely contributes to transcriptional

interference and the eventual failure of prolonged benefits of modest mHTT lowering. Targeting these resistant

fractions of misfolded mHTT by a chaperone activity to aid in its degradation may be beneficial in combination with

gene therapy *HTT* lowering strategies.

We found that early, continuous partial lowering of mHTT protein for up to 12 months fully or partially

preserved PDE10A and SCN4B protein levels with initiation of lowering embryonically providing the highest benefits.

These data agree with previous findings showing preservation of *Pde10a* mRNA in the LacQ140 model after early

m*Htt* lowering [59] and preservation of the Pde10a PET signal in the Q175 model after striatal injection of AAV-HTT-ZFP

[17]. However, our data here indicate that the time points chosen for post-treatment analysis are important and that

changes do not follow a linear neurodegenerative trajectory in mice. The greatest number of protein and lipid

changes in LacQ140 mice occurred at 9 months, with few changes detected at 12 months. Similarly, Langfelder et al.

found a greater number of differentially expressed genes at 6 months compared to 10 months in zQ175 mice [39]. This

suggests that in the mouse brain, adverse responses to mHTT oscillate or go through waves of degeneration and

regeneration. Therefore, to appreciate any benefits afforded by m*Htt* lowering, frequent or continuous monitoring

should be conducted.

In this study, mass spectrometry of lipids identified numerous alterations in LacQ140 striatum, many of

which were prevented with modest m*Htt* lowering. To our surprise, lipidomic analysis showed that LacQ140 mice had

increased levels in species of glycerophospholipids PI and PS starting at 6 months and progressing to a significant

change in the PI and PS subclass level at 9 months. Proteomics in Q140 synaptosomes revealed changes in proteins

that regulate PI levels including PKC signaling and PIP2 hydrolysis, changes in two isoforms of DGKs, and alterations in

one of the rate-limiting enzymes in PI synthesis (CDS2) [33] all of which impact PI levels [95]. Transcriptomic profiling of

LacQ140 mice also showed a plethora of altered mRNA levels for enzymes that impact PI and PIPs **(Fig. 6c and f;**

Additional Files 2 and 3). Normally found on the inner leaflet of the plasma membrane, PS can be externalized by

apoptotic cells to signal for their demolition [103] and in neuronal synapses, externalized PS signals to microglia for

synaptic pruning [104].Microglial activation occurs in HD post-mortem brain [105] and increased pruning by microglia is

hypothesized to contribute to synaptic loss in R6/2 HD striatum [106] and in zQ175 and BACHD mice, as well as in HD

brain [107]. An overall increase in PS could inadvertently mark synapses or myelin [108] for engulfment by microglia. The

ratio of PS to PE impacts autophagy [109] which may in turn impair mHTT removal [110,111]. Both PI and PS are abundant in

astrocytes as well as neurons [112] so it is unclear which cell type(s) is producing the changes in these lipids. We cannot

rule out the possibility that lipid changes are due to altered cellular composition of the brain.

Changes in white matter detected through imaging are one of the first signs of disease in people with HD

(PwHD) [113–127]. Morphometric studies of postmortem HD brains showed reduced cross-sectional area of white matter

as well as gray matter atrophy [128,129]. In HD post-mortem brain tissue, a dramatic shift in the profile of various

sphingolipids including Cer, SM, hexosylceramides, and sulfatides occurred [51]. Here, we demonstrate in the striatum

of 9-month-old LacQ140 mice significant reductions (compared to WT mice) of relative levels of total lipids and the

lipid subclasses SM and Cer, and three individual species of Hex1Cer, all important for myelin [130]. Our data are in

alignment with data from R6/1 mice showing changes in cerebroside and sulfatide levels [131], from the R6/2 mouse

model showing reductions in components of the sphingolipid biosynthesis pathway [45], and findings in a transgenic

sheep model OVT73, similarly showing decreased levels of numerous species of SM [132]. A salient finding was a

profound reduction in LacQ140 striatum of the low abundance signaling lipid MGDG, which regulates

oligodendrocyte differentiation [89]. MGDG is considered a marker of myelination and stimulates PKC-alpha activity in

oligodendrocytes to support process formation [90]. We previously reported reduced levels of MGDG in subcellular

fractions of Q175/Q7 HD striatum at 2 and 6 months [47]. Crucially, in LacQ140 mice, lowering m*Htt* improved the loss

of SM and MGDG suggesting protection against white matter pathology.

Altered oligodendrocyte differentiation or survival due to direct effects of mHTT protein on transcription in

the nucleus may account for many of the lipid changes we observed. RNA transcripts of genes important for

oligodendrocyte differentiation and myelin sphingolipid biosynthesis were altered in LacQ140 mice. The transcription

factor *Tcf7l2*, which was lower at 6 months in LacQ140 mice, was recently implicated in altered myelin formation in

R6/2 and Q175 mice [133]. The basic helix-loop-helix transcription factor *Olig1*, which was increased at 12 months in

LacQ140 mice compared to WT, is important for commitment of cells to CNS oligodendrocyte identity [134]. Critically,

Lim et al. presented evidence showing abnormal oligodendrocyte maturation in multiple HD postmortem brain

regions, as well as R6/2 brain, with single-nucleus RNAseq showing changes in *OLIG1* and *OLIG2* [135]. Altered levels of

myelin transcripts were found in human embryonic stem cells differentiated along an oligodendrocyte pathway [136]

and an epigenic etiology for changes in myelin gene expression in human oligodendrocyte precursors that was

blocked by inactivation of m*HTT* allele was described [137].

Changes in particular enzyme levels that regulate lipid biosynthesis that were changed at the transcriptional

level in LacQ140 mice can have dire consequences and result directly in myelin defects. Both *Ugt8a* and *Fa2h* mRNA

were lower in the striatum of LacQ140 mice; *Ugt8a* mRNA levels were protected by m*Htt* lowering at 6 months, but

Fa2h mRNA was not. Work by others showed that mice deficient in the *Ugt8a* gene exhibited abnormal myelin

maturation and structure [138,139]. In humans, mutations in FA2H are associated with leukodystrophy and hereditary

spastic paraplegia type SPG35 [140,141] highlighting the importance of hydroxylated sphingolipids in myelin integrity.

Adult *Fa2h*-deficient mice have normal oligodendrocyte differentiation with normal appearing myelin that later

degenerates, showing "splitting of lammelae" by 18 months [142]. This is similar to the Ki140CAG mouse model [143]

where myelin appears to be quite normal into early adulthood, but then may start to degenerate with disease

progression.

Although transcriptional deregulation clearly impacts lipid modifying enzymes, levels of myelin-related lipids could be caused by an interaction of mHTT with oligodendrocyte membranes or a be a consequence of Wallerian degeneration of cortical-striatal axons of the neurons. Interestingly, the presence of SM increased permeabilization of monolayers by mHTT *in vitro* [53], suggesting mHTT could have particular effects on myelin lipids. Crucially, mHTT was localized within myelin sheaths using immunogold EM in 9-month-old Q175 striatum [80]. Moreover, mHTT can be secreted by neurons in culture and in the brain as a soluble free form [144]. We speculate that mHTT could insert directly into myelin bilayers to disrupt myelin architecture.

Observing white matter changes in animal models has been challenging. White matter loss was reported in R6/1 mice [145], and changes in myelination have been described in Yac128 [146,147] and HdhQ250 mice [148], but experiments designed to look for white matter changes in the Q150 HD mouse model showed brain atrophy but no white matter abnormalities [149]. A recent imaging study of OVT73 sheep brain reported changes in diffusivity in the internal capsule at 9-10 years, suggesting changes in white matter microstructure [150]. Our biochemical experiments here show that mHTT effects on striatal lipid homeostasis in HD mouse models are complex. We and others have reported lipidomic and metabolomic studies on knock-in Q140/Q140 HD mice at single time points [48] and Q111 HD mice [44] but did not observe loss of lipids important for white matter. Curiously, the lipid differences in LacQ140 mice measured at 9 months disappeared at 12 months suggesting that, even in the absence of m*Htt* lowering, the mouse brain insulted with mHTT attempts to heal itself and succeeds at some level. Consistent with our lipidomic findings, a longitudinal imaging study over 18 months showed transient changes in diffusivity/fractional anisotropy of corpus collosum in Q140 mouse brain [143]. These results echo imaging data from presymptomatic PwHD suggesting attempted remyelination [151]. Hence, if HD mouse models undergo a series of degeneration and regeneration cycles, observations at one or two time points may be misleading. By 12 months, although changes at the lipid subclass level were annulled in the LacQ140 mice, a detailed analysis of the individual lipid species comprising these subclasses shows a shift in species within each subclass. The altered composition of subclasses may alter function, weaken HD brains, or predispose them to further stress. In fact, the behavioral analyses of these mice which showed loss of protection by mHTT lowering at 12 months [59] indicates that although levels of lipids by subclasses are restored at 12 months (with or without mHTT lowering), the changes in individual lipid species which compose each subclasses correlate with behavioral degradation.

Not all lipid changes were reversed with m*Htt* lowering. Consistent with metabolic defects in HD, here we

report reduced levels of species of TG, glycerolipid molecules used for energy storage which can be metabolized by

mitochondria, in LacQ140 compared to WT. Of note, reductions in species of TG were not reversed by m*Htt* lowering.

Interestingly, the LacQ140 mice exhibited reciprocal changes in the two biosynthetic enzymes *Dgat1* and *Dgat2* at

the transcriptional level, suggesting the ability to store energy may arise in part from this variation.

CONCLUSIONS

Collectively, our studies advocate early lowering of m*HTT* for greatest benefit, and in this context, modest

lowering is sufficient to delay some protein and lipid changes. Furthermore, our work shows readily detectable but

transient changes in lipids highly enriched in myelin, consistent with possible white matter damage and regeneration

occurring in the LacQ140 mouse model.

LIST OF ABBREVIATIONS

HD: Huntington's disease

HTT: Huntingtin

ASOs: Antisense oligonucleotide

miRNA: MicroRNA

shRNA: Short hairpin RNA

DARPP32: Dopamine And CAMP-Regulated Neuronal Phosphoprotein 32

PDE10A: Phosphodiesterase 10A

SCN4B: Sodium Voltage-Gated Channel Beta Subunit 4

ATP5A: ATP Synthase F1 Subunit Alpha

PIP: Phosphatidylinositol phosphate

IPTG: Isopropyl b-D-1-thiogalactopyranoside

Poly-Q: Polyglutamine

GFAP: Glial fibrillary acidic protein

LC-MS: Liquid chromatography – mass spectrometry

PS: Phosphatidylserine

PI: Phosphatidylinositol

BisMePA: Bis-methyl phosphatidic acid

PKC: Protein kinase C

SM: Sphingomyelin

Cer: Ceramide

MGDG: Monogalactosyldiacylglycerol

Hex1Cer: Monohexosylceramide

GalCer: Galactosylceramide

GlcCer: Glucosylceramide

TG: Triacylglycerol

GO BP: Gene Ontology Biological Processes

GSEA: Gene Set Enrichment Analysis

Ugt8a: UDP-Galactose Ceramide Galactosyltransferase

Fa2h: Fatty acid 2-hydroxylase

DECLARATIONS

Ethics approval and consent to participate

Mice were housed at Psychogenics (Paramus, NJ) and all treatments and procedures were conducted with oversight

by Psychogenics Institutional Animal Care and Use Committee.

Consent for publication

Not applicable

Availability of data and materials

All datasets generated are included in this article and can be found in **Additional Files 1-5.** RNA-sequencing data

analyzed is available at Gene Expression Omnibus (GEO) accession number GSE156236.

Competing interests

KBK-G spouse owns less than 0.1% stock in the following companies: Advanced Microdevices, Aveo Pharmaceuticals,

Inc, Boston Scientific Corporation, Bristol-Myers Squibb Company, Cisco Systems, Inc., Fate Therapeutics, GE

Healthcare Life Sciences, Generex Biotechnology Corporation, Idera Pharmaceuticals, Inc., Nante Health,

Neurometrics, Inc., NuGenerex, Repligen Corporation, Sesen Bio, Inc., T2 Biosystems, and Vericel Corporation.

Other authors have no declarations of interest.

Funding

This work was funded by CHDI Foundation, Inc., a nonprofit biomedical research organization exclusively dedicated

to developing therapeutics that will substantially improve the lives of HD-affected individuals, the Dake family fund

to MD and KBK-G, and NIH 1S10RR023594S10 to MD.

Authors' contributions

KKG, MD, and DM conceived experimental plan and wrote manuscript. KS performed all lipidomics and

bioinformatics and wrote manuscript, ES performed protein chemistry analysis and wrote manuscript, AB aided in

the experimental plan and edited manuscript, CS and SL performed subcellular fractionations.

Acknowledgements

The authors would like to acknowledge the Dake family for their support.

REFERENCES

1. Difiglia, M. *et al.* Huntingtin Is a Cytoplasmic Protein Associated with Vesicles in Human and Rat Brain Neurons. *Neuron* **14**, 1075–1081 (1995).
2. Sharp, A. H. *et al.* Widespread expression of Huntington's disease gene (IT15) protein product. *Neuron* **14**, 1065–74 (1995).
3. Caviston, J. P. & Holzbaur, E. L. F. Huntingtin as an essential integrator of intracellular vesicular trafficking. *Trends in Cell Biology* **19**, 147–155 (2009).
4. Gao, R. *et al.* Mutant huntingtin impairs PNKP and ATXN3, disrupting DNA repair and transcription. *eLife* (2019) doi:10.7554/eLife.42988.001.
5. Saudou, F. & Humbert, S. The Biology of Huntingtin. *Neuron* **89**, 910–926 (2016).
6. Barron, J. C., Hurley, E. P. & Parsons, M. P. Huntingtin and the Synapse. *Front Cell Neurosci* **15**, 689332 (2021).
7. Duyao, M. P. *et al.* Inactivation of the mouse Huntington's disease gene homolog Hdh. *Science* **269**, 407–10 (1995).
8. McKinstry, S. U. *et al.* Huntingtin is required for normal excitatory synapse development in cortical and striatal circuits. *J Neurosci* **34**, 9455–72 (2014).
9. Nasir, J. *et al.* Targeted disruption of the Huntington's disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes. *Cell* **81**, 811–23 (1995).
10. O'Kusky, J. R., Nasir, J., Cicchetti, F., Parent, A. & Hayden, M. R. Neuronal degeneration in the basal ganglia and loss of pallido-subthalamic synapses in mice with targeted disruption of the Huntington's disease gene. *Brain Res* **818**, 468–79 (1999).
11. Zeitlin, S., Liu2, J.-P., Chapman3, D. L., Papaioannou, V. E. & Efstratiadis, A. Increased apoptosis and early embryonic lethality in mice nullizygou s for the Huntington 's disease gene homologu e. *Nature Genetics* **11**, 155–163 (1995).
12. Nopoulos, P. C. Huntington disease: a single-gene degenerative disorder of the striatum. *Dialogues Clin Neurosci* **18**, 91–8 (2016).
13. Petersen, A. & Bjorkqvist, M. Hypothalamic-endocrine aspects in Huntington's disease. *Eur J Neurosci* **24**, 961–7 (2006).
14. Tabrizi, S. J., Ghosh, R. & Leavitt, B. R. Huntingtin Lowering Strategies for Disease Modification in Huntington's Disease. *Neuron* **101**, 801–819 (2019).
15. Caron, N. S., Dorsey, E. R. & Hayden, M. R. Therapeutic approaches to huntington disease: From the bench to the clinic. *Nature Reviews Drug Discovery* **17**, 729–750 (2018).
16. Keller, C. G. *et al.* An orally available, brain penetrant, small molecule lowers huntingtin levels by enhancing pseudoexon inclusion. *Nature Communications* **13**, (2022).
17. Zeitler, B. *et al.* Allele-selective transcriptional repression of mutant HTT for the treatment of Huntington's disease. *Nature Medicine* **25**, 1131–1142 (2019).
18. Alterman, J. F. *et al.* A divalent siRNA chemical scaffold for potent and sustained modulation of gene expression throughout the central nervous system. *Nature Biotechnology* **37**, 884–894 (2019).
19. Keeler, A. M. *et al.* Cellular Analysis of Silencing the Huntington's Disease Gene Using AAV9 Mediated Delivery of Artificial Micro RNA into the Striatum of Q140/Q140 Mice. *J Huntingtons Dis* **5**, 239–248 (2016).
20. McBride, J. L. *et al.* Preclinical safety of RNAi-mediated HTT suppression in the rhesus macaque as a potential therapy for Huntington's disease. *Mol Ther* **19**, 2152–62 (2011).
21. Boudreau, R. L. *et al.* Nonallele-specific silencing of mutant and wild-type huntingtin demonstrates therapeutic efficacy in Huntington's disease mice. *Mol Ther* **17**, 1053–63 (2009).
22. Datson, N. A. *et al.* The expanded CAG repeat in the huntingtin gene as target for therapeutic RNA modulation throughout the HD mouse brain. *PLoS One* **12**, e0171127 (2017).
23. DiFiglia, M. *et al.* Therapeutic silencing of mutant huntingtin with siRNA attenuates striatal and cortical neuropathology and behavioral deficits. *Proc Natl Acad Sci U S A* **104**, 17204–9 (2007).
24. Drouet, V. *et al.* Sustained effects of nonallele-specific Huntingtin silencing. *Ann Neurol* **65**, 276–85 (2009).
25. Kordasiewicz, H. B. *et al.* Sustained Therapeutic Reversal of Huntington's Disease by Transient Repression of Huntingtin Synthesis. *Neuron* **74**, 1031–1044 (2012).
26. Rodriguez-Lebron, E., Denovan-Wright, E. M., Nash, K., Lewin, A. S. & Mandel, R. J. Intrastriatal rAAV-mediated delivery of anti-huntingtin shRNAs induces partial reversal of disease progression in R6/1 Huntington's disease transgenic mice. *Mol Ther* **12**, 618–33 (2005).

27. Stanek, L. M. *et al.* Antisense oligonucleotide-mediated correction of transcriptional dysregulation is correlated
with behavioral benefits in the YAC128 mouse model of Huntington's disease. *J Huntingtons Dis* **2**, 217–28
(2013).
28. Stanek, L. M. *et al.* Silencing mutant huntingtin by adeno-associated virus-mediated RNA interference
ameliorates disease manifestations in the YAC128 mouse model of Huntington's disease. *Hum Gene Ther* **25**,
461–74 (2014).
29. Leavitt, B. R., Kordasiewicz, H. B. & Schobel, S. A. Huntingtin-Lowering Therapies for Huntington Disease: A
Review of the Evidence of Potential Benefits and Risks. *JAMA Neurol* **77**, 764–772 (2020).
30. Dragatsis, I., Levine, M. S. & Zeitlin, S. Inactivation of Hdh in the brain and testis results in progressive
neurodegeneration and sterility in mice. *Nature Genetics* **26**, 300–306 (2000).
31. Mehler, M. F. *et al.* Loss-of-Huntingtin in Medial and Lateral Ganglionic Lineages Differentially Disrupts Regional
Interneuron and Projection Neuron Subtypes and Promotes Huntington's Disease-Associated Behavioral,
Cellular, and Pathological Hallmarks. *J Neurosci* **39**, 1892–1909 (2019).
32. Bibb, J. A. *et al.* Severe deficiencies in dopamine signaling in presymptomatic Huntington's disease mice. *Proc
Natl Acad Sci U S A* **97**, 6809–14 (2000).
33. Sapp, E. *et al.* Protein changes in synaptosomes of Huntington's disease knock-in mice are dependent on age
and brain region. *Neurobiology of Disease* **141**, (2020).
34. Skotte, N. H. *et al.* Integrative Characterization of the R6/2 Mouse Model of Huntington's Disease Reveals
Dysfunctional Astrocyte Metabolism. *Cell Reports* **23**, 2211–2224 (2018).
35. Bertoglio, D. *et al.* Development of a ligand for in vivo imaging of mutant huntingtin in Huntington's disease.
Science Translational Medicine **14**, 3682 (2022).
36. Mehta, S. R. *et al.* Human Huntington's Disease iPSC-Derived Cortical Neurons Display Altered Transcriptomics,
Morphology, and Maturation. *Cell Rep* **25**, 1081-1096 e6 (2018).
37. Oyama, F. *et al.* Sodium channel beta4 subunit: down-regulation and possible involvement in neuritic
degeneration in Huntington's disease transgenic mice. *J Neurochem* **98**, 518–29 (2006).
38. Agrawal, S. & Fox, J. H. Novel proteomic changes in brain mitochondria provide insights into mitochondrial
dysfunction in mouse models of Huntington's disease. *Mitochondrion* **47**, 318–329 (2019).
39. Langfelder, P. *et al.* Integrated genomics and proteomics define huntingtin CAG length-dependent networks in
mice. *Nature Neuroscience* **19**, 623–633 (2016).
40. McQuade, L. R. *et al.* Proteomics of Huntington's disease-affected human embryonic stem cells reveals an
evolving pathology involving mitochondrial dysfunction and metabolic disturbances. *J Proteome Res* **13**, 5648–
59 (2014).
41. Kegel, K. B. *et al.* Polyglutamine expansion in huntingtin increases its insertion into lipid bilayers. *Biochemical
and Biophysical Research Communications* **387**, 472–475 (2009).
42. Hyuk Yoon, J. *et al.* Brain lipidomics: From functional landscape to clinical significance. *Science Advances* **8**, 9317
(2022).
43. Harayama, T. & Riezman, H. Understanding the diversity of membrane lipid composition. *Nature Reviews
Molecular Cell Biology* **19**, 281–296 (2018).
44. Carroll, J. B. *et al.* HdhQ111 mice exhibit tissue specific metabolite profiles that include striatal lipid
accumulation. *PLoS ONE* **10**, (2015).
45. Di Pardo, A. *et al.* De novo synthesis of sphingolipids is defective in experimental models of Huntington's
disease. *Frontiers in Neuroscience* **11**, (2017).
46. Farzana, F. *et al.* Longitudinal spatial mapping of lipid metabolites reveals pre-symptomatic changes in the
hippocampi of Huntington's disease transgenic mice. *Neurobiology of Disease* **176**, 105933 (2023).
47. Iuliano, M. *et al.* Disposition of Proteins and Lipids in Synaptic Membrane Compartments Is Altered in Q175/Q7
Huntington's Disease Mouse Striatum. *Frontiers in Synaptic Neuroscience* **13**, (2021).
48. Vodicka, P. *et al.* Mass Spectrometry Analysis of Wild-Type and Knock-in Q140/Q140 Huntington's Disease
Mouse Brains Reveals Changes in Glycerophospholipids Including Alterations in Phosphatidic Acid and Lyso-
Phosphatidic Acid. *Journal of Huntington's Disease* **4**, 187–201 (2015).
49. Hunter, M., Demarais, N. J., Faull, R. L. M., Grey, A. C. & Curtis, M. A. Subventricular zone lipidomic architecture
loss in Huntington's disease. *Journal of Neurochemistry* **146**, 613–630 (2018).

50. Hunter, M., Demarais, N. J., Faull, R. L. M., Grey, A. C. & Curtis, M. A. An imaging mass spectrometry atlas of
lipids in the human neurologically normal and Huntington's disease caudate nucleus. *Journal of Neurochemistry*
157, 2158–2172 (2021).
51. Phillips, G. *et al.* The long and the short of Huntington's disease: how the sphingolipid profile is shifted in the
caudate of advanced clinical cases. *Brain Communications* **4**, (2022).
52. Kegel, K. B. *et al.* Polyglutamine expansion in huntingtin alters its interaction with phospholipids. *Journal of
Neurochemistry* **110**, 1585–1597 (2009).
53. Chaibva, M. *et al.* Sphingomyelin and GM1 Influence Huntingtin Binding to, Disruptionof, and Aggregation on
Lipid Membranes. *ACS Omega* **3**, 273–285 (2018).
54. Southwell, A. L. *et al.* Huntingtin suppression restores cognitive function in a mouse model of Huntington's
disease. *Science Translational Medicine* **10**, 3959 (2018).
55. Caron, N. S. *et al.* Cerebrospinal fluid mutant huntingtin is a biomarker for huntingtin lowering in the striatum of
Huntington disease mice. *Neurobiology of Disease* **166**, (2022).
56. Southwell, A. L. *et al.* Ultrasensitive measurement of huntingtin protein in cerebrospinal fluid demonstrates
increase with Huntington disease stage and decrease following brain huntingtin suppression. *Sci Rep* **5**, 12166
(2015).
57. Coffey, S. R. *et al.* Huntingtin lowering reduces somatic instability at CAG-expanded loci. *bioRxiv* (2020)
doi:10.1101/2020.07.23.218347.
58. Cronin, C. A., Gluba, W. & Scrable, H. The lac operator-repressor system is functional in the mouse. *Genes Dev*
15, 1506–17 (2001).
59. Marchionini, D. M. *et al.* Benefits of global mutant huntingtin lowering diminish over time in a Huntington's
disease mouse model. *JCI Insight* **7**, (2022).
60. Scrable, H. Say when: reversible control of gene expression in the mouse by lac. *Semin Cell Dev Biol* **13**, 109–19
(2002).
61. Landles, C. *et al.* Proteolysis of mutant huntingtin produces an exon 1 fragment that accumulates as an
aggregated protein in neuronal nuclei in Huntington disease. *J Biol Chem* **285**, 8808–23 (2010).
62. Scherzinger, E. *et al.* Huntingtin-encoded polyglutamine expansions form amyloid-like protein aggregates in
vitro and in vivo. *Cell* **90**, 549–58 (1997).
63. Weiss, A. *et al.* Sensitive biochemical aggregate detection reveals aggregation onset before symptom
development in cellular and murine models of Huntington's disease. *J Neurochem* **104**, 846–58 (2008).
64. Breitkopf, S. B. *et al.* A relative quantitative positive/negative ion switching method for untargeted lipidomics
via high resolution LC-MS/MS from any biological source. *Metabolomics* **13**, (2017).
65. Matyash, V., Liebisch, G., Kurzchalia, T. V., Shevchenko, A. & Schwudke, D. Lipid extraction by methyl-terf-butyl
ether for high-throughput lipidomics. *Journal of Lipid Research* **49**, 1137–1146 (2008).
66. Liebisch, G. *et al.* Update on LIPID MAPS classification, nomenclature, and shorthand notation for MS-derived
lipid structures. *Journal of Lipid Research* **61**, 1539–1555 (2020).
67. Reza, S., Ugorski, M. & Suchański, J. Glucosylceramide and galactosylceramide, small glycosphingolipids with
significant impact on health and disease. *Glycobiology* **31**, 1416–1434 (2021).
68. Koelmel, J. P., Ulmer, C. Z., Jones, C. M., Yost, R. A. & Bowden, J. A. Common cases of improper lipid annotation
using high-resolution tandem mass spectrometry data and corresponding limitations in biological
interpretation. *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids* **1862**, 766–770 (2017).
69. Gu, Z. Complex heatmap visualization. *iMeta* (2022) doi:10.1002/imt2.43.
70. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biology* **15**, 550 (2014).
71. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to
Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300 (1995).
72. Kegel, K. B. *et al.* Huntingtin is present in the nucleus, interacts with the transcriptional corepressor C-terminal
binding protein, and represses transcription. *J Biol Chem* **277**, 7466–76 (2002).
73. Sapp, E. *et al.* Huntingtin Localization in Brains of Normal and Huntington's Disease Patients. *Annals of
Neurology* **42**, 604–612 (1997).
74. Tousley, A. *et al.* Huntingtin associates with the actin cytoskeleton and alpha-actinin isoforms to influence
stimulus dependent morphology changes. *PLoS One* **14**, e0212337 (2019).

75. Trettel, F. Dominant phenotypes produced by the HD mutation in STHdhQ111 striatal cells. *Human Molecular Genetics* **9**, 2799–2809 (2000).

76. Seong, I. S. *et al.* Huntingtin facilitates polycomb repressive complex 2. *Human Molecular Genetics* **19**, 573–583 (2010).

77. Ko, J., Ou, S. & Patterson, P. H. New anti-huntingtin monoclonal antibodies: implications for huntingtin conformation and its binding proteins. *Brain Res Bull* **56**, 319–329 (2001).

78. Legleiter, J. *et al.* Monoclonal Antibodies Recognize Distinct Conformational Epitopes Formed by Polyglutamine in a Mutant Huntingtin Fragment *. *Journal of Biological Chemistry* **284**, 21647–21658 (2009).

79. Miller, J. *et al.* Identifying polyglutamine protein species in situ that best predict neurodegeneration. *Nature Chemical Biology* **7**, 925–934 (2011).

80. Ko, J. *et al.* Identification of distinct conformations associated with monomers and fibril assemblies of mutant huntingtin. *Human Molecular Genetics* **27**, 2330–2343 (2018).

81. Sathasivam, K. *et al.* Identical oligomeric and fibrillar structures captured from the brains of R6/2 and knock-in mouse models of Huntington's disease. *Human Molecular Genetics* **19**, 65–78 (2010).

82. Kegel, K. B. *et al.* Huntingtin Expression Stimulates Endosomal-Lysosomal Activity, Endosome Tubulation, and Autophagy. *Journal of Neuroscience* (2000).

83. Vodicka, P. *et al.* Assessment of chloroquine treatment for modulating autophagy flux in brain of WT and HD mice. *J Huntingtons Dis* **3**, 159–74 (2014).

84. Gu, M. *et al.* Mitochondrial defect in Huntington's disease caudate nucleus. *Ann Neurol* **39**, 385–9 (1996).

85. Berridge, M. J. The Inositol Trisphosphate/Calcium Signaling Pathway in Health and Disease. *Physiological Reviews* **96**, 1261–1296 (2016).

86. Hirt, U. A. & Leist, M. Rapid, noninflammatory and PS-dependent phagocytic clearance of necrotic cells. *Cell Death Differ* **10**, 1156–1164 (2003).

87. Kim, H.-Y., Huang, B. X. & Spector, A. A. Phosphatidylserine in the brain: Metabolism and function. *Progress in Lipid Research* **56**, 1–18 (2014).

88. Chrast, R., Saher, G., Nave, K. A. & Verheijen, M. H. G. Lipid metabolism in myelinating glial cells: Lessons from human inherited disorders and mouse models. *Journal of Lipid Research* **52**, 419–434 (2011).

89. Inoue, T., Deshmukh, D. S. & Pieringer, R. A. The Association of the Galactosyl Diglycerides of Brain with Myelination. *Journal of Biological Chemistry* **246**, 5688–5694 (1971).

90. Schmidt-Schultz, T. & Althaus, H. H. Monogalactosyl Diglyceride, a Marker for Myelination, Activates Oligodendroglial Protein Kinase C. *Journal of Neurochemistry* **62**, 1578–1585 (1994).

91. Vanier, M. T. & Svennerholm, L. Chemical pathology of Krabbe's disease. III. Ceramide-hexosides and gangliosides of brain. *Acta Paediatr Scand* **64**, 641–648 (1975).

92. Schmitt, S., Cantuti Castelvetri, L. & Simons, M. Metabolism and functions of lipids in myelin. *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids* **1851**, 999–1005 (2015).

93. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences* (2005).

94. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation* **2**, (2021).

95. Blunsom, N. J. & Cockcroft, S. CDP-Diacylglycerol Synthases (CDS): Gateway to Phosphatidylinositol and Cardiolipin Synthesis. *Frontiers in Cell and Developmental Biology* **8**, (2020).

96. Harris, C. A. *et al.* DGAT enzymes are required for triacylglycerol synthesis and lipid droplets in adipocytes [S]. *Journal of Lipid Research* **52**, 657–667 (2011).

97. van der Bijl, P., Strous, G. J., Lopes-Cardozo, M., Thomas-Oates, J. & van Meer, G. Synthesis of non-hydroxy-galactosylceramides and galactosyldiglycerides by hydroxy-ceramide galactosyltransferase. *Biochem J* **317**, 589–597 (1996).

98. Alderson, N. L., Maldonado, E. N., Kern, M. J., Bhat, N. R. & Hama, H. FA2H-dependent fatty acid 2-hydroxylation in postnatal mouse brain. *Journal of Lipid Research* **47**, 2772–2780 (2006).

99. Diaz-Hernandez, M. *et al.* Full motor recovery despite striatal neuron loss and formation of irreversible amyloid-like inclusions in a conditional mouse model of Huntington's disease. *J Neurosci* **25**, 9773–81 (2005).

100. Fox, L. M. *et al.* Huntington's Disease Pathogenesis Is Modified In Vivo by Alfy/Wdfy3 and Selective Macroautophagy. *Neuron* **105**, 813-821.e6 (2020).

101. Wheeler, V. C. Long glutamine tracts cause nuclear localization of a novel form of huntingtin in medium spiny
striatal neurons in HdhQ92 and HdhQ111 knock-in mice. *Human Molecular Genetics* **9**, 503–513 (2000).
102. Kaganovich, D., Kopito, R. & Frydman, J. Misfolded proteins partition between two distinct quality control
compartments. *Nature* **454**, 1088–95 (2008).
103. Williamson, P. & Schlegel, R. A. Transbilayer phospholipid movement and the clearance of apoptotic cells.
Biochim Biophys Acta **1585**, 53–63 (2002).
104. Scott-Hewitt, N. *et al.* Local externalization of phosphatidylserine mediates developmental synaptic pruning by
microglia. *The EMBO Journal* **39**, (2020).
105. Sapp, E. *et al.* Early and progressive accumulation of reactive microglia in the Huntington disease brain. *J
Neuropathol Exp Neurol* **60**, 161–72 (2001).
106. Savage, J. C. *et al.* Microglial physiological properties and interactions with synapses are altered at
presymptomatic stages in a mouse model of Huntington's disease pathology. *J Neuroinflammation* **17**, 98
(2020).
107. Wilton, D. *et al.* Microglia Mediate Early Corticostriatal Synapse Loss and Cognitive Dysfunction in Huntington's
Disease Through Complement-Dependent Mechanisms. *bioRxiv* (2021) doi:10.1101/2021.12.03.471180.
108. Djannatian, M. *et al.* Myelination generates aberrant ultrastructure that is resolved by microglia. *Journal of Cell
Biology* **222**, (2023).
109. Polyansky, A. *et al.* Phospholipid imbalance impairs autophagosome completion. *The EMBO Journal* (2022)
doi:10.15252/embj.2022110771.
110. Qin, Z. H. *et al.* Autophagy regulates the processing of amino terminal huntingtin fragments. *Hum Mol Genet* **12**,
3231–44 (2003).
111. Yamamoto, A., Cremona, M. L. & Rothman, J. E. Autophagy-mediated clearance of huntingtin aggregates
triggered by the insulin-signaling pathway. *Journal of Cell Biology* **172**, 719–731 (2006).
112. Fitzner, D. *et al.* Cell-Type- and Brain-Region-Resolved Mouse Brain Lipidome. *Cell Reports* **32**, (2020).
113. Bourbon-Teles, J. *et al.* Myelin Breakdown in Human Huntington's Disease: Multi-Modal Evidence from
Diffusion MRI and Quantitative Magnetization Transfer. *Neuroscience* **403**, 79–92 (2019).
114. Di Paola, M. *et al.* Multimodal MRI analysis of the corpus callosum reveals white matter differences in
presymptomatic and early Huntington's disease. *Cereb Cortex* **22**, 2858–66 (2012).
115. Faria, A. V. *et al.* Linking white matter and deep gray matter alterations in premanifest Huntington disease.
Neuroimage Clin **11**, 450–460 (2016).
116. Hobbs, N. Z. *et al.* The progression of regional atrophy in premanifest and early Huntington's disease: a
longitudinal voxel-based morphometry study. *J Neurol Neurosurg Psychiatry* **81**, 756–63 (2010).
117. Matsui, J. T. *et al.* Prefrontal cortex white matter tracts in prodromal Huntington disease. *Hum Brain Mapp* **36**,
3717–32 (2015).
118. Odish, O. F. *et al.* Microstructural brain abnormalities in Huntington's disease: A two-year follow-up. *Hum Brain
Mapp* **36**, 2061–74 (2015).
119. Oh, S. L. *et al.* Fixel-Based Analysis Effectively Identifies White Matter Tract Degeneration in Huntington's
Disease. *Front Neurosci* **15**, 711651 (2021).
120. Paulsen, J. S. Functional imaging in Huntington's disease. *Exp Neurol* **216**, 272–7 (2009).
121. Paulsen, J. S. *et al.* Detection of Huntington's disease decades before diagnosis: the Predict-HD study. *J Neurol
Neurosurg Psychiatry* **79**, 874–80 (2008).
122. Phillips, O. *et al.* Tractography of the corpus callosum in Huntington's disease. *PLoS One* **8**, e73280 (2013).
123. Poudel, G. R. *et al.* Longitudinal change in white matter microstructure in Huntington's disease: The IMAGE-HD
study. *Neurobiol Dis* **74**, 406–12 (2015).
124. Rosas, H. D. *et al.* Diffusion tensor imaging in presymptomatic and early Huntington's disease: Selective white
matter pathology and its relationship to clinical measures. *Movement Disorders* **21**, 1317–1325 (2006).
125. Singh, S., Mehta, H. & Fekete, R. Altered Fractional Anisotropy in Early Huntington's Disease. *Case Rep Neurol* **5**,
26–30 (2013).
126. Sweidan, W., Bao, F., Bozorgzad, N. S. & George, E. White and Gray Matter Abnormalities in Manifest
Huntington's Disease: Cross-Sectional and Longitudinal Analysis. *J Neuroimaging* **30**, 351–358 (2020).
127. Tereshchenko, A. *et al.* Brain structure in juvenile-onset Huntington disease. *Neurology* **92**, e1939–e1947
(2019).

128. de la Monte, S. M., Vonsattel, J. P. & Richardson, E. P., Jr. Morphometric demonstration of atrophic changes in the cerebral cortex, white matter, and neostriatum in Huntington's disease. *J Neuropathol Exp Neurol* **47**, 516–25 (1988).

129. Vonsattel, J. P. & DiFiglia, M. Huntington disease. *J Neuropathol Exp Neurol* **57**, 369–84 (1998).

130. O'Brien, J. S. & Sampson, E. L. Lipid composition of the normal human brain: gray matter, white matter, and myelin. *Journal of Lipid Research* **6**, 537–544 (1965).

131. Desplats, P. A. *et al.* Glycolipid and ganglioside metabolism imbalances in Huntington's disease. *Neurobiology of Disease* **27**, 265–277 (2007).

132. Skene, D. J. *et al.* Metabolic profiling of presymptomatic Huntington's disease sheep reveals novel biomarkers. *Scientific Reports* **7**, (2017).

133. Benraiss, A. *et al.* A TCF7L2-responsive suppression of both homeostatic and compensatory remyelination in Huntington disease mice. *Cell Reports* **40**, 111291 (2022).

134. Dai, J., Bercury, K. K., Ahrendsen, J. T. & Macklin, W. B. Olig1 function is required for oligodendrocyte differentiation in the mouse brain. *J Neurosci* **35**, 4386–402 (2015).

135. Lim, R. G. *et al.* Huntington disease oligodendrocyte maturation deficits revealed by single-nucleus RNAseq are rescued by thiamine-biotin supplementation. *Nature Communications* **13**, 7791 (2022).

136. Osipovitch, M. *et al.* Human ESC-Derived Chimeric Mouse Models of Huntington's Disease Reveal Cell-Intrinsic Defects in Glial Progenitor Cell Differentiation. *Cell Stem Cell* **24**, 107-122.e7 (2019).

137. Bardile, C. F. *et al.* Intrinsic mutant HTT-mediated defects in oligodendroglia cause myelination deficits and behavioral abnormalities in Huntington disease. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 9622–9627 (2019).

138. Bosio, A., Binczek, E. & Stoffel, W. Functional breakdown of the lipid bilayer of the myelin membrane in central and peripheral nervous system by disrupted galactocerebroside synthesis. *Proceedings of the National Academy of Sciences* **93**, 13280–13285 (1996).

139. Coetzee, T. *et al.* Myelination in the Absence of Galactocerebroside and Sulfatide: Normal Structure with Abnormal Function and Regional Instability. *Cell* **86**, 209–219 (1996).

140. Edvardson, S. *et al.* Mutations in the Fatty Acid 2-Hydroxylase Gene Are Associated with Leukodystrophy with Spastic Paraparesis and Dystonia. *American Journal of Human Genetics* **83**, 643–648 (2008).

141. Rattay, T. W. *et al.* FAHN/SPG35: a narrow phenotypic spectrum across disease classifications. *Brain* **142**, 1561–1572 (2019).

142. Zöller, I. *et al.* Absence of 2-hydroxylated sphingolipids is compatible with normal neural development but causes late-onset axon and myelin sheath degeneration. *Journal of Neuroscience* **28**, 9741–9754 (2008).

143. Pérot, J. B. *et al.* Longitudinal multimodal MRI characterization of a knock-in mouse model of Huntington's disease reveals early gray and white matter alterations. *Human molecular genetics* **31**, 3581–3596 (2022).

144. Caron, N. S. *et al.* Mutant Huntingtin Is Cleared from the Brain via Active Mechanisms in Huntington Disease. *J Neurosci* **41**, 780–796 (2021).

145. Rattray, I. *et al.* Correlations of behavioral deficits with brain pathology assessed through longitudinal MRI and histopathology in the R6/1 mouse model of Huntington's disease. *PLoS One* **8**, e84726 (2013).

146. Teo, R. T. Y. *et al.* Structural and molecular myelination deficits occur prior to neuronal loss in the YAC128 and BACHD models of Huntington disease. *Human Molecular Genetics* **25**, 2621–2632 (2016).

147. Teo, R. T. Y. *et al.* Impaired Remyelination in a Mouse Model of Huntington Disease. *Molecular Neurobiology* **56**, 6873–6882 (2019).

148. Jin, J. *et al.* Early white matter abnormalities, progressive brain pathology and motor deficits in a novel knock-in mouse model of Huntington's disease. *Human Molecular Genetics* **24**, 2508–2527 (2015).

149. Steventon, J. J. *et al.* Longitudinal in vivo MRI in a Huntington's disease mouse model: Global atrophy in the absence of white matter microstructural damage. *Scientific Reports* **6**, (2016).

150. Taghian, T. *et al.* Brain Alterations in Aged OVT73 Sheep Model of Huntington's Disease: An MRI Based Approach. *Journal of Huntington's Disease* 1–16 (2022) doi:10.3233/jhd-220526.

151. Phillips, O. *et al.* Deep white matter in Huntington's disease. *PLoS ONE* **9**, (2014).

Table 1. Overview of lipid subclass changes at 6, 9, and 12 months

Category	Subclass	6-month-old mice			9-month-old mice			12-month-old mice		
		Species detected	ANOVA p-val	Adj p-val (q), N =36	Species detected	ANOVA p-val	Adj p-val (q), N =24	Species detected	ANOVA p-val	Adj p-val (q), N =29
Glycerophos-pholipids	PC	134	0.7415	1	107	0.2928	0.3267	160	0.4243	0.4627
	PE	152	0.1975	1	154	0.0284*	0.0563	119	0.5653	0.5372
	PS	62	0.1171	1	58	**0.0014**	**0.0125**	25	0.46	0.4627
	PI	19	**0.0189**	0.7144	17	**0.0054**	**0.0168**	34	0.0086*	0.0338*
	MePC	17	0.9859	1	7	0.4901	0.486	15	0.7552	0.6329
	PA	13	0.9596	1	6	0.7008	0.5957	9	0.0025*	0.0184*
	BisMePA	4	0.1589	1	1	**0.0041**	**0.0168**	6	0.2812	0.3647
	dMePE	9	0.195	1	2	0.2041	0.269	0	N/A	N/A
	PG	19	0.6255	1	4	0.4685	0.486	4	0.859	0.6765
	BisMePE	1	0.8072	1	0	N/A	N/A	0	N/A	N/A
	BisMePS	0	N/A	N/A	0	N/A	N/A	2	0.0001*	0.0022*
	CL	22	0.5812	1	0	N/A	N/A	0	N/A	N/A
	PEt	7	0.3261	1	2	0.9792	0.7283	0	N/A	N/A
	PMe	2	0.8695	1	0	N/A	N/A	0	N/A	N/A
	PIP2	2	0.1393	1	0	N/A	N/A	3	0.0161*	0.0394*
	PIP	1	0.7034	1	0	N/A	N/A	0	N/A	N/A
	LPC	30	0.7943	1	10	0.6835	0.5957	7	0.2805	0.3647
	LPE	23	0.9947	1	16	0.6396	0.5957	4	0.0127*	0.035*
	LPS	7	0.9061	1	5	0.211	0.269	2	0.0817	0.1287
	LPI	7	0.7746	1	1	0.1715	0.2551	0	N/A	N/A
	LPG	4	0.8888	1	0	N/A	N/A	0	N/A	N/A
	LdMePE	1	0.4552	1	0	N/A	N/A	0	N/A	N/A
	BiotinylPE	0	N/A	N/A	1	0.8078	0.6554	1	0.0225*	0.0451*
Glycerolipids	TG	107	0.5998	1	44	0.0057*	0.0168*	72	0.0002*	0.0022*
	MGDG	3	0.8693	1	2	**0.0005**	**0.0089**	5	0.0214*	0.0451*
	MGMG	6	0.7537	1	0	N/A	N/A	6	0.0497*	0.0887
	DG	13	0.1381	1	70	0.1174	0.1905	59	0.6614	0.5834
	SQMG	2	0.5986	1	0	N/A	N/A	0	N/A	N/A
	SQDG	1	0.9316	1	0	N/A	N/A	0	N/A	N/A
Sphingolipids	Hex1Cer	0	N/A	N/A	85	0.0243*	0.0542	104	0.4617	0.4627
	CerG1	50	0.5466	1	0	N/A	N/A	0	N/A	N/A
	SM	47	0.6398	1	23	**0.0066**	**0.0168**	36	0.3243	0.3973
	Cer	14	0.9804	1	6	**0.0048**	**0.0168**	21	0.4374	0.4627
	CerP	0	N/A	N/A	0	N/A	N/A	17	0.0035*	0.0193*
	ST	10	0.3977	1	9	0.2283	0.2717	14	0.2176	0.3199
	So	3	0.4465	1	0	N/A	N/A	0	N/A	N/A
	CerPE	0	N/A	N/A	1	0.0766	0.1367	0	N/A	N/A
	phSM	4	0.3619	1	0	N/A	N/A	0	N/A	N/A
	CerG2GNAc1	1	0.5784	1	0	N/A	N/A	0	N/A	N/A
	SPH	0	N/A	N/A	0	N/A	N/A	1	0.775	0.6329
	Hex2Cer	0	N/A	N/A	0	N/A	N/A	1	0.0112	0.035
Sterol Lipids	ChE	1	0.5341	1	1	0.918	0.7124	2	0.9151	0.6958
	ZyE	0	N/A	N/A	0	N/A	N/A	3	0.0092*	0.0338*
Fatty Acyls	FA	2	0.5893	1	0	N/A	N/A	0	N/A	N/A
	AcCa	0	N/A	N/A	0	N/A	N/A	2	0.5847	0.5372
Prenol Lipids	Co	0	N/A	N/A	0	N/A	N/A	1	0.0523	0.0887
	Total	800			632			735		

Table 1. Overview of lipid subclass changes at 6, 9, and 12 months.

Bold text indicates lipid subclasses significantly different between WT and LacQ140 mice (p<0.05 or q<0.05).
Asterisk (*) indicates lipid subclasses significantly different among other groups (p<0.05 or q<0.05).
One way ANOVA was conducted followed by Tukey's multiple comparisons test, two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli, q<0.05.

Table 2. Overall comparison of lipidomic results across ages.

Age	# Species detected	# Species different at p<0.05*	# Species different for WT/LacQ140 at p<0.05	# Species different at q<0.05**	# Species different for WT/LacQ140 at q<0.05	# Species recovered with lowering (q<0.05)
6 months	800	9	4	0	0	0
9 months	632	192	72	17	14	14
12 months	735	162	36	72	26	1

*p value for ANOVA; **Adjusted p value determined using Benjamini, Krieger, and Yekutieli procedure

MAIN FIGURE LEGENDS

Figure 1. Generation of LacQ140 mice and treatment paradigm
(a) The *LacO/LacIR*-regulatable HD mouse model (LacQ140) was generated by crossing the *Htt*$^{LacQ140/+}$ mouse to the *Tg*$^{ACTB-lacI*Scrb}$ mouse [58] as previously described [59]. The default state of the LacQ140 mouse is global repression of m*Htt* due to *Lac* Repressor binding to the *Lac* operators. Administration of IPTG starting from embryonic day 5 (E5) interrupts the binding between the *Lac* repressor and operators, resulting in a de-repressed state, and maximal expression of m*Htt* in LacQ140. All WT mice were Htt$^{LacO+/+}$; b-actin-LacI^R tg. **(b)** Mice were fed *ad libitum*; the lactose analog IPTG was provided in drinking water (at 10mM) which de-represses the *LacQ140* allele and keeps normal m*Htt* expression. During embryonic development, m*Htt* expression levels were maintained at normal levels by administering IPTG to pregnant dams starting at embryonic day 5 (E5). IPTG was continuously administered to WT mice. IPTG was administered always (m*Htt* always expressed, LacQ140), withdrawn at 8 months (m*Htt* repressed beginning at 8 months, LacQ140_8M), withdrawn 2 at months (m*Htt* repressed beginning at 2 months, LacQ140_2M), or never administered (m*Htt* always repressed, LacQ140_A). Tissue for each group (except LacQ140_8M) was collected at 6, 9, and 12 months of age.

Figure 2. Analysis of mHTT protein levels in crude homogenates of 6-, 9- and 12-months old mice.
HTT levels were analyzed by capillary immunoassay on equal amounts of protein (0.6 µg) using anti-HTT antibody Ab1 and anti-polyQ antibody MW1 (**a**). Peak area analysis performed using Compass software in 6-month-old mice shows a significant decrease in WT HTT as detected with Ab1 in all LacQ140 mice compared to WT mice (F(3, 20) = 5.674, **P=0.0056, One-way ANOVA with Tukey's multiple comparison test, n=6). mHTT levels are significantly lower in LacQ140_2M and LacQ140_A as detected with both Ab1 and MW1 compared to LacQ140 (**a**, Ab1: F(2, 15) = 11.25, **P=0.0010, -38% and -43% respectively; MW1: F(2, 14) = 9.879, **P=0.0021, -40% and -47% respectively). Peak area analysis in 9-month-old mice shows a significant decrease in WT HTT as detected with Ab1 in all LacQ140 mice compared to WT mice (F(3, 20) = 34.67, ****P<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6). mHTT levels are significantly lower in LacQ140_2M and LacQ140_A, as detected with both Ab1 and MW1, compared to LacQ140 (**b**, Ab1: F(2, 15) = 10.82, **P=0.0012, -42% and -52% respectively; MW1: F(2, 15) = 20.82, ****P<0.0001, -44% and -43% respectively). Peak area analysis in 12-month-old mice shows a significant decrease in WT HTT as detected with Ab1 in all LacQ140 mice compared to WT mice (F(4, 25) = 15.81, ****P<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6). WT HTT was significantly lower in LacQ140_A compared to LacQ140_8M and LacQ140_2M mice. mHTT levels are significantly lower in LacQ140_A mice, as detected with Ab1, compared to LacQ140 (**c**, Ab1: F(3, 20) = 5.017, **P=0.0094, -51%, One-way ANOVA with Tukey's multiple comparison test, n=6). Asterisks on graphs represent Tukey's multiple comparison test, n=6 mice per group (*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001).

Figure 3. HTT protein levels in P1 fractions.
Equal protein (10 µg) from P1 fractions from 6-month-old (**a**) and 12-month-old (**b**) LacQ140 and WT mice were analyzed by western blot for HTT levels with anti-HTT Ab1. No aggregated protein was observed at the top of the gel (arrowhead). Total pixel intensity quantification for each band was measured using ImageJ software and normalized to HDAC1 signal. There was a significant decrease in WT HTT signal in all the treatment conditions for LacQ140 mice compared to WT mice in both **(a)** 6-month-old mice (F(3, 20) = 40.34, ****P<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6) and **(b)** 12-month-old mice (F(4, 25) = 11.01, ****P<0.0001 ,One-way ANOVA with Tukey's multiple comparison test, n=6). There were significantly lower levels of mHTT in the 6-month-old LacQ140_A mice compared to LacQ140 (**a,** F(2, 15) = 8.233, **P=0.0039, One-way ANOVA with Tukey's multiple comparison test, n=6) but no changes in mHTT levels were detected in the **(b)** 12-month-old LacQ140 mice (F(3, 20) = 1.137, P=0.3583, n.s., One-way ANOVA). Equal protein (10 µg) from P1 fractions from 12-month-old LacQ140 and WT mice were analyzed by western blot for HTT levels with anti-HTT S830 (**c**). The S830 antibody detected a smear of HTT signal (bracket) as well as full-length mHTT (arrow). There were significantly lower levels of full length mHTT in the 12-month-old LacQ140_A mice compared to LacQ140 (F(3, 20) = 3.548, *P=0.0330, One-way ANOVA with Tukey's multiple comparison test, n=6) and no changes detected in the HTT smear in all LacQ140 mice (F(3, 20) = 0.9281,

P=0.4453, n.s., One-way ANOVA). Asterisks on graphs represent Tukey's multiple comparison test, n=6 mice per group (*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001).

Figure 4. Analyses of lipids in crude homogenates LacQ140 caudate putamen by mass spectrometry.
(a) Total lipid intensity detected at 6 months normalized to WT; no significant difference between groups (one-way ANOVA: F(3, 20) = 0.4604, P=0.7130, n.s., n=6). **(b)** Total lipid intensity detected at 9 months normalized to WT; LacQ140 mice have decreased total lipid intensity which is reversed in LacQ140_A mice (one-way ANOVA & Tukey's multiple comparison test: F(3, 20) = 5.474, **P=0.0065, n=6). **(c)** Total lipid intensity detected at 12 months normalized to WT; no significant difference between groups (one-way ANOVA: F (4, 25) = 0.7504, P=0.5671, n.s., n=6). **(d)** Heatmap depicts the lipid subclass composition for WT, LacQ140, and treatment groups at 9 months. Hierarchical clustering was performed across lipid subclasses (rows) and columns (animals) using the one minus Pearson correlation distance metric. ANOVA p-value column indicates lipid subclasses significantly changed between LacQ140 and WT mice in green (p<0.05, One-way ANOVA with Tukey's multiple comparison test, n=6). Lipid subclasses with adjusted p-values (q) < 0.05 LacQ140 vs WT are indicated in purple (q<0.05, two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli, N=24 lipid subclasses, n=6 mice). Source data and full statistical details can be found in **Additional Files 1, 4, & 5**.

Figure 5. Restoration of dysregulated lipid subclasses with lowering of m*Htt.*
Graphs show relative intensities for lipid subclasses expressed as a percent of total lipid intensity detected (bars = mean, error bars = ± SD). **(a)** PS increased in LacQ140 mice and is reversed in LacQ140_A mice; F(3, 20) = 7.601, **P=0.0014, q=0.0125, **(b)** PI increased in LacQ140 mice and reversed in LacQ140_A mice; F(3, 20) = 5.707, **P=0.0054, q=0.0168, **(c)** BisMePA decreased in LacQ140 mice and is reversed in LacQ140 mice; F(3, 20) = 6.086, **P=0.0041, q=0.0168, **(d)** SM decreased in LacQ140 mice and is reversed in LacQ140_2M and LacQ140_A mice; F(3, 20) = 5.465, **P=0.0066, q=0.0168, **(e)** Cer decreased in LacQ140 mice; F(3, 20) = 5.883, **P=0.0048, q=0.0168, **(f)** MGDG decreased in LacQ140 mice and is reversed in LacQ140_2M and LacQ140_A mice; F(3, 20) = 9.350, ***P=0.0005, q=0.0089. Statistics are one-way ANOVA and asterisks on graphs represent Tukey's multiple comparison test, n=6 mice per group. **(g)** Heatmap shows individual lipid species that comprise each subclass. Hierarchical clustering was performed across individual lipid species (rows) and animals (columns) using the one minus Pearson correlation distance metric. Individual lipid species significantly changed between any group are indicated in green (p<0.05, one-way ANOVA), lipid species significantly changed between LacQ140 and WT are indicated in purple (p<0.05, one-way ANOVA, n=6) and red (q<0.05, one-way ANOVA & two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli, N=632 lipid species, n=6 mice). Source data and full statistical details can be found in **Additional Files 1, 4, & 5**.

Figure 6. Lipid metabolism and myelin associated transcriptional changes and reversal with m*Htt* lowering.
(a) Dotplot (clusterProfiler) of lipid or myelin related GO BP terms overrepresented (one sided hypergeometric test, padj<0.05) in 6-month LacQ140 downregulated genes (padj<0.05, FC>20%). GeneRatio represents the number of genes associated with each GO term/number of downregulated genes. Dots are sized by count of genes associated with respective terms. **(b)** Gene set enrichment analysis (clusterProfiler) of 6-month LacQ140, LacQ140_2M and LacQ140_A groups compared to WT. All lipid and myelin associated GO BP terms significantly enriched in LacQ140 compared to WT (padj<0.05) are displayed. X axis for each respective group represents the normalized enrichment score (NES) and bars are colored by significance (padj<0.05 = blue, padj>0.05 = tan). Adjusted p-values are displayed adjacent to bars. **(c)** Heatmap shows differentially expressed genes in 6-month-old LacQ140 mice compared to WT (padj<0.05, FC > ±20%). Gene expression is shown as median ratio normalized counts (DESeq2), scaled by respective gene (columns). Bars above the heatmap indicate DEGs reversed with m*Htt* lowering. LacQ140_2M = green, LacQ140_A = purple; padj<0.05, FC > 20% opposite of LacQ140**. (d)** Dotplot of lipid or myelin related GO BP terms overrepresented in 12-month downregulated genes. GO BP terms significantly overrepresented at 6-months (a) are displayed for comparison. 4/9 terms (ensheathment of neurons, axon ensheathment, myelination, and phospholipase C activating G-protein coupled receptor signaling pathway) are significantly overrepresented in 12-month LacQ140 downregulated genes (one sided hypergeometric test, padj < 0.05). **(e)** Gene set enrichment analysis

of LacQ140, LacQ140_8M, LacQ140_2M and LacQ140_A groups compared to WT. GO BP terms enriched at 6-months (b) are displayed for comparison. 8/22 GO BP terms are significantly negatively enriched (padj < 0.05) at 12-months, shown in blue. **(f)** Heatmap shows differentially expressed genes in 12-month-old LacQ140 mice compared to WT (padj<0.05, FC > ±20%). Gene expression is shown as median ratio normalized counts (DESeq2), scaled by respective gene (columns). Bars above the heatmap DEGs reversed with m*Htt* lowering (LacQ140_8M = bottom bar, no reversal, LacQ140_2M = green, LacQ140_A = purple; padj<0.05, FC >20% opposite of LacQ140). GO BP terms associated with genes, fold changes, and exact FDR values can be found in **Additional Files 2 & 3.**

Figure 7. Simplified de novo sphingolipid biosynthesis pathway.
(a) Serine and palmitoyl CoA are condensed by serine palmitoyltransferase (SPT) to generate 3-ketosphinganine. 3-ketophinganine is reduced to dihydrosphingosine by 3-ketodihydrosphingosine reductase (KDSR). Dihydrosphingosine is acetylated by ceramide synthases (CERS) and further desaturated by ceramide desaturase (DEGS) to generate ceramide. Ceramide is the substrate for generation of other sphingolipids (sphingomyelin, galactosylceramide, glucosylceramide, sulfatide, and lactosylceramide). Abbreviations: SGMS = sphingomyelin synthase, SMase = sphingomyelinase, UGT8A = UDP galactosyltransferase 8A, GALC = galactosylceramidase. CST = galactosylceramide sulfotransferase, ARSA = arylsulfatase, GBA = glucosylceramidase, UGCG = UDP-glucose ceramide glucosyltransferase, LacCer synthase = lactosylceramide synthase. **(b)** Biosynthesis of non-hydroxylated sphingolipids. **(c)** Biosynthesis of 2-hydroxy sphingolipids. Fatty acid 2-hydroxylase (FA2H) catalyzes hydroxylation of fatty acids in the C2 position, which can be incorporated into sphingolipid precursors (i.e., dihydroceramide) in the acylation step of de novo synthesis.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

Supplementary Figure 1. Analysis of mHTT protein levels with EPR5526 and PHP3 in crude homogenates of 6-, 9- and 12-months old mice.

HTT levels were analyzed by western blot on equal amounts of protein (10 µg) using anti-HTT antibody EPR5526 and anti-polyQ antibody PHP3. Total pixel intensity quantification for each band measured using ImageJ software in 6-month-old mice shows a significant decrease in WT HTT as detected with EPR5526 in all LacQ140 mice compared to WT mice but no change in mHTT levels (**a**). mHTT levels are significantly lower in LacQ140_2M and LacQ140_A mice as detected with PHP3 compared to LacQ140 (**a**, -42% and -35% respectively, $**p<0.01$, $***p<0.001$, One-way ANOVA with Tukey's multiple comparison test, n=6). Total pixel intensity quantification in 9-month-old mice shows a significant decrease in WT HTT as detected with EPR5526 in all LacQ140 mice compared to WT mice. (**b**). mHTT levels are significantly lower in LacQ140_2M and LacQ140_A mice as detected with EPR5526 and PHP3 compared to LacQ140 (**b**, EPR5526: -24% and -30%, respectively; PHP3: -46% and -49% respectively, $*p<0.05$, $**p<0.01$, $***p<0.001$, One-way ANOVA with Tukey's multiple comparison test, n=6).Total pixel intensity quantification in 12-month-old mice shows a significant decrease in WT HTT as detected with EPR5526 in LacQ140 and LacQ140_2M compared to WT mice. No changes in mHTT levels as detected with either EPR5526 or PHP3 were observed (**c**).

Supplementary Figure 2

Supplementary Figure 2. Effects of m*Htt* lowering on the subcellular distribution of WT and mHTT protein by density gradient ultracentrifugation.

Diagram depicts the centrifugation strategy for protein samples **(a)**. Schematic shows the approximate location in the fractions of protein markers and the organelles found in these compartments **(b)**. Representative western blot images for equal volumes of fractions 1-5 and 12-14 from 6-month-old mice probed with anti-HTT antibody Ab1 are shown in **(c)**. The remaining images are shown in **Supplementary Figure 3a**. Total pixel intensity quantification for each band measured using ImageJ software is graphed as average percent mutant/WT HTT ± SD for each fraction **(d)**. Since each fraction contains different levels of proteins normally used to control for protein loading, levels of mHTT were normalized to levels of WT HTT which was not repressed/lowered. The ratio mutant/WT HTT is significantly higher in LacQ140 mice compared to LacQ140_2M in fractions 13 and 14 (*$p<0.05$, One-way ANOVA with Tukey's multiple comparison test for each fraction, n=6). Representative western blot images for equal volumes of fractions 1-4 and 11-14 from 12-month-old mice probed with anti-HTT antibody Ab1 are shown in **(e)**. The remaining images are shown in **Supplementary Figure 3c**. Total pixel intensity quantification for each band is graphed as average percent mutant/WT HTT ± SD for each fraction **(f)**. The ratio mutant/WT HTT is significantly higher in LacQ140 compared to LacQ140_8M, LacQ140_2M and/or LacQ140_A mice in fractions 1, 3, 13, and 14 (*$p<0.05$, **$p<0.01$, One-way ANOVA with Tukey's multiple comparison test for each fraction, n=3). Graphs indicate data in fractions where mHTT was detected in at least 3 mice except LacQ140_A fractions 1 and 3 where only 2 mice had detectible mHTT.

Supplementary Figure 3

a

b

c

d

Supplementary Figure 3. Effects of m*Htt* lowering on the subcellular distribution of WT and mHTT.
Western blot images for equal volumes of fractions 1-14 (right blots, each strip is from one mouse) or fractions 1-5 and 12-14 (left blots) from 6-month-old mice probed with anti-HTT antibody Ab1 are shown in **(a)**. Each strip in left blots is a set of 1 mouse per treatment group with the groups labeled at the top of the blots. Total pixel intensity quantification for each band using ImageJ software is graphed as average percent of total HTT signal for each fraction ± SD **(b)**. Representative western blot images for equal volumes of fractions 1-14 (right blots, each strip is from one mouse) or 1-4 and 11-14 (left blots) from 12-month-old mice probed with anti-HTT antibody Ab1 are shown in **(c)**. Each strip in the left blots is from 2 mice per group with the groups labeled on the right. Total pixel intensity quantified for each band using ImageJ software are graphed as average percent of total HTT signal for each fraction ± SD **(d)**. There is no difference in WT or mHTT levels as a percent of total HTT in any fraction with any treatment of LacQ140 mice at 6 or 12 months (One-way ANOVA with Tukey's multiple comparison test for each fraction, n=6 for 6 months and n=3 for 12 months).

Supplementary Figure 4

Supplementary Figure 4. HTT levels in crude homogenates and P1 fractions from WT and LacQ140 mice by filter trap assay.

Filter trap assays of 12-month-old crude homogenates **(a)** and P1 fractions **(b)** were probed with S830 antibody. Each dot represents one animal and each of the 6 dots across equals one group which is labeled on the left and right sides. There are 2 dots for the lysates from R6/2 HD mice which have a highly expressing transgene for a small fragment of HTT containing a large CAG repeat (180CAGs) and that accumulate numerous aggregates which have been shown to be retained in the assay and were used as a positive control. There was significantly more signal for aggregated mHTT in the 12-month-old LacQ140 mice compared to WT, LacQ140_8M, LacQ140_2M and LacQ140_A mice (**a**, *p<0.05, **p<0.01, ****p<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6). In the P1 fractions, there was significantly more signal for aggregated mHTT detected with S830 antibody in the 12-month LacQ140 mice compared to WT, LacQ140_8M, LacQ140_2M and LacQ140_A mice and in LacQ140_8M compared to WT and LacQ140_A mice (**b**, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6).

Supplementary Figure 5

a

6 month old mice

WT | LacQ140 | LacQ140_2M | LacQ140_A

95- PDE10A
43- SCN4B
130- Vinculin

PDE10A
Signal intensity normalized to vinculin
**** ** ****
**
*
WT | LacQ140 | LacQ140_2M | LacQ140_A

SCN4B
** *
WT | LacQ140 | LacQ140_2M | LacQ140_A

b

9 month old mice

WT | LacQ140 | LacQ140_2M | LacQ140_A

95- PDE10A
43- SCN4B
130- Vinculin

PDE10A
Signal intensity normalized to vinculin
**** ***
**
**
WT | LacQ140 | LacQ140_2M | LacQ140_A

SCN4B
**** **
*
*
WT | LacQ140 | LacQ140_2M | LacQ140_A

c

12 month old mice

Females | Males

WT | LacQ140 | LacQ140_8M | LacQ140_2M | LacQ140_A | WT | LacQ140 | LacQ140_8M | LacQ140_2M | LacQ140_A

95- PDE10A
43- SCN4B
130- Vinculin

PDE10A
Signal intensity normalized to vinculin
*** *
* **
*
WT | LacQ140 | LacQ140_8M | LacQ140_2M | LacQ140_A

SCN4B
WT | LacQ140 | LacQ140_8M | LacQ140_2M | LacQ140_A

Supplementary Figure 5. Duration of m*Htt* lowering in 6, 9 and 12 months old LacQ140 mice affects levels of PDE10A and SCN4B.

PDE10A and SCN4B levels were analyzed by western blot on equal amounts of protein (10 µg). Total pixel intensity quantification for each band using ImageJ software in 6-month-old mice shows a significant decrease in PDE10A levels in LacQ140 compared to WT mice. There is an increase in PDE10A levels in LacQ140_2M and LacQ140_A mice compared to LacQ140 and no change from WT mice (**a,** *p<0.05, **p<0.01, ****p<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6). There is a significant decrease in SCN4B levels in LacQ140 compared to WT mice and a significant increase back to WT levels in LacQ140_2M mice. Total pixel intensity quantification in 9-month-old mice shows a significant decrease in PDE10A levels in LacQ140 compared to WT mice. There is an increase in PDE10A levels in LacQ140_2M compared to LacQ140 and no change from WT mice (**b,** **p<0.01, ***p<0.001, One-way ANOVA with Tukey's multiple comparison test, n=6). There is a significant decrease in SCN4B levels in LacQ140 compared to WT mice. There is a significant increase in SCN4B levels in LacQ140_2M and LacQ140_A mice compared to LacQ140 but significantly lower than in the WT mice (**b,** *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, One-way ANOVA with Tukey's multiple comparison test, n=6). Total pixel intensity quantification in 12-month-old mice shows a significant decrease in PDE10A levels in LacQ140, LacQ140_8M and LacQ140_2M compared to WT mice. There is an increase in PDE10A levels in LacQ140_A mice compared to LacQ140 and LacQ140_2M and no change from WT mice (**c,** *p<0.05, **p<0.01, ***p<0.001, One-way ANOVA with Tukey's multiple comparison test, n=6). There are no changes in SCN4B levels in any of the LacQ140 or WT mice.

Supplementary Figure 6

Supplementary Figure 6. Duration of m*Htt* lowering in 6, 9 and 12 months old LacQ140 mice has minimal effects on levels of GFAP, DARPP32 and ATP5A.

GFAP and DARPP32 levels were analyzed by capillary immunoassay on equal amounts of protein (0.6 µg) and ATP5A levels were analyzed by western blot on equal amounts of protein (10 µg). In 6-month-old mice, peak area analysis shows no significant change in GFAP or DARPP32 levels (**a**) and total pixel intensity quantification shows no changes in ATP5A levels in any of the LacQ140 or WT mice (**a**). Peak area analysis in 9-month-old mice shows a significant decrease in DARPP32 levels in all LacQ140 mice compared to WT mice (**b**, **$p<0.01$, ***$p<0.001$, One-way ANOVA with Tukey's multiple comparison test, n=6). Total pixel intensity quantification shows ATP5A levels are significantly higher in LacQ140_2M and LacQ140_A mice compared to WT or LacQ140 mice (**b,** *$p<0.05$, **$p<0.01$, ***$p<0.001$, One-way ANOVA with Tukey's multiple comparison test, n=6). Peak area analysis in 12-month-old mice shows no significant change in GFAP levels in LacQ140 or WT mice. There is a significant decrease in DARPP32 levels in LacQ140_8M and LacQ140_2M compared to LacQ140_A mice (**c,** *$p<0.05$, One-way ANOVA with Tukey's multiple comparison test, n=6). Total pixel intensity quantification for each band using ImageJ software shows no changes in ATP5A levels in any of the LacQ140 or WT mice.

Supplementary Figure 7. Lipid subclasses detected in caudate putamen of 6-month-old mice.
Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid subclasses expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent summed lipid subclass intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. PI was significantly increased in striatum of 6-month-old LacQ140 mice compared to WT mice (One-way ANOVA with Tukey's multiple comparisons test, $F_{(3, 20)} = 4.176$, * P=0.0189, n=6). This increase is not significant when p-values are adjusted for multiple testing using the Benjamini, Krieger and Yekutieli procedure with a false discovery rate of 5%, (q = 0.7144, N=36 subclasses). No changes between groups were found in any other subclasses.

Abbreviations:
Glycerophospholipids: PC = phosphatidylcholine, PE = phosphatidylethanolamine, PS = phosphatidylserine, PI = phosphatidylinositol, MePC = methylphosphocholine, PA = phosphatidic acid, BisMePA = bis-methyl phosphatidic acid, dMePE = dimethyl phosphatidylethanolamine, PG = phosphatidylgylcerol, BisMePE = bis-methyl phosphatidyl ethanolamine, CL = cardiolipin, PEt = phosphatidylethanol, PMe = phosphatidylmethanol, PIP2 = phosphatidylinositol-bisphosphate, PIP = phosphatidylinositol-monophosphate, LPC = lysophosphatidylcholine, LPE = lysophosphatidylethanolamine, LPS = lysophosphatidylserine, LPI = lysophosphatidylinositol LPG = lysophosphosphatidylgylcerol, LdMePE = lysodimethyl phosphatidyl ethanolamine
Glycerolipids: TG = triacylglycerol, MGDG = monogalactosyldiacylglycerol, MGMG = monogalactosylmonoacylglycerol, DG = diacylglycerol, SQMG = sulfoquinovosylmonoacylglycerol, SQDG = sulfoquinovosyldiacylglycerol,
Sphingolipids: CerG1 = simple Glc series, SM = sphingomyelin, Cer = ceramide, ST = sulfatide, So = sphingosine, phSM = sphingomyelin phytosphingosine, CerG2GNAc1 = Simple Glc series
Sterol lipids: ChE = cholesterol ester
Fatty acyls: FA = fatty acid

Supplementary Figure 8

Supplementary Figure 8. Individual lipid species significantly different between 6-month-old LacQ140 and WT mice.

Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid species expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent individual lipid species intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. One-way analysis of variance (ANOVA) was used to evaluate differences in lipid species intensity between groups and Tukey's multiple comparisons test was used for post-hoc pairwise comparisons (n=6 mice per group, *p < 0.05, Tukey's multiple comparisons test). To correct for multiple testing over all lipids analyzed, the Benjamini, Krieger and Yekutieli procedure was used with a false discovery rate of 5% (N=800 lipid species), FDR adjusted p-values are reported as q values. No individual species found to be different by one-way ANOVA were significant following correction of p-values. PS 22:6_22:6: $F_{(3, 20)}$ = 3.812,*P=0.0260, q=1, ns; PS 18:0_20:4: $F_{(3, 20)}$ = 3.349, *P=0.0396, q=1, ns; PI 18:0_20:4: $F_{(3, 20)}$ = 3.812, *P=0.0260, q=1, ns; PI 16:0_22:6: $F_{(3, 20)}$ = 3.466, *P=0.0356, q=1, ns. Abbreviations: PI = phosphatidylinositol, PS = phosphatidylserine

Supplementary Figure 9. Lipid subclasses detected in caudate putamen of 9-month-old mice.
Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid subclasses expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent summed lipid subclass intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. One-way analysis of variance (ANOVA) was used to evaluate differences in lipid subclass intensity between groups and Tukey's multiple comparisons test was used for post-hoc pairwise comparisons (n=6 mice per group, *p < 0.05, **p < 0.01, Tukey's multiple comparisons test). To correct for multiple testing the Benjamini, Krieger and Yekutieli procedure was used with a false discovery rate of 5% (N=24 subclasses) and FDR adjusted p-values are reported as q values. PS: $F_{(3, 20)}$ = 7.601, **P=0.0014, q=0.0125, PI: $F_{(3, 20)}$ = 5.707, **P=0.0054, q=0.0168, BisMePA: $F_{(3, 20)}$ = 6.086, **P=0.0041, q=0.0168, SM: $F_{(3, 20)}$ = 5.465, **P=0.0066, q=0.0168, Cer: $F_{(3, 20)}$ = 5.883, **P=0.0048, q=0.0168, MGDG: $F_{(3, 20)}$ = 9.350 , ***P=0.0005, q=0.0089. PE, TG and Hex1Cer were unchanged between LacQ140 and WT groups but had changes among other treatment groups (PE: $F_{(3, 20)}$ = 3.717, *P=0.0284, q=0.0563, ns, TG: $F_{(3, 20)}$ = 5.637, **P=0.0057, q=0.0168, Hex1Cer: $F_{(3, 20)}$ = 3.891, *P=0.0243, q=0.054, ns).

Abbreviations:
Glycerophospholipids: PC = phosphatidylcholine, PE = phosphatidylethanolamine, PS = phosphatidylserine, PI = phosphatidylinositol, PA = phosphatidic acid, BisMePA = bis-methyl phosphatidic acid, MePC = methylphosphocholine, PG = phosphatidylgylcerol, dMePE = dimethyl phosphatidylethanolamine, PEt = phosphatidylethanol, BiotinylPE = biotinyl-phosphoethanolamine, LPC = lysophosphatidylcholine, LPE = lysophosphatidylethanolamine, LPS = lysophosphatidylserine, LPI = lysophosphatidylinositol
Glycerolipids: TG = triacylglycerol, DG = diacylglycerol, MGDG = monogalactosyldiacylglycerol
Sphingolipids: Hex1Cer = monohexosylceramide, SM = sphingomyelin, ST = sulfatide, Cer = ceramide, CerPE = ceramide phosphorylethanolamine
Sterol lipids: ChE = cholesterol ester

Supplementary Figure 10

Supplementary Figure 10. Individual lipid species significantly changed between 9-month-old LacQ140 and WT mice.

Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid species expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent individual lipid species intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. One-way analysis of variance (ANOVA) was used to evaluate differences in lipid species intensity between groups and Tukey's multiple comparisons test was used for post-hoc pairwise comparisons (n=6 mice per group, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, Tukey's multiple comparisons test). To correct for multiple testing over all lipids analyzed, the Benjamini, Krieger and Yekutieli procedure was used with a false discovery rate of 5% (N=632 lipid species), FDR adjusted p-values are reported as q values. PC 36:5e: $F_{(3, 20)}$ = 10.68, ***P=0.0002, q=0.02, PE 16:1e_20:4: $F_{(3, 20)}$ = 18.14, ****P<0.0001, q=0.003, PE 16:0p_20:4: $F_{(3, 20)}$ = 16.97, ****P<0.0001, q=0.003, PI 18:1_20:4: $F_{(3, 20)}$ = 8.086, **P=0.0010, q=0.041, PI 16:0_20:4: $F_{(3, 20)}$ = 8.449, ***P=0.0008, q=0.039, PS 18:0_18:1: $F_{(3, 20)}$ = 8.226, ***P=0.0009, q=0.041, PS 18:1_22:6: $F_{(3, 20)}$ = 7.916, **P=0.0011, q=0.042, PS 18:0_20:4: $F_{(3, 20)}$ = 9.739, ***P=0.0004, q=0.023, Hex1Cer d18:1_18:0: $F_{(3, 20)}$ = 10.37, ***P=0.0002, q=0.02, Hex1Cer t18:0_18:0: $F_{(3, 20)}$ = 10.35, ***P=0.0003, q=0.02, Hex1Cer d18:1_25:1: $F_{(3, 20)}$ = 8.440, ***P=0.0008, q=0.039, SM d40:2: $F_{(3, 20)}$ = 8.052, **P=0.0010, q=0.041, TG 8:2COOH_18:1_18:1: $F_{(3, 20)}$ = 11.98, ***P=0.0001, q=0.017, MGDG 16:0_18:1: $F_{(3, 20)}$ = 11.28, ***P=0.0002, q=0.019.

Abbreviations:
Glycerophospholipids: PC = phosphatidylcholine, PE = phosphatidylethanolamine, PI = phosphatidylinositol, PS = phosphatidylserine
Sphingolipids: Hex1Cer = monohexosylceramide, SM = sphingomyelin
Glycerolipids: MGDG = monogalactosyldiacylglycerol, TG = triacylglycerol

Supplementary Figure 11

Supplementary Figure 11. Lipid subclasses detected in caudate putamen of 12-month-old mice.
Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid subclasses expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent summed lipid subclass intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. One-way analysis of variance (ANOVA) was used to evaluate differences in lipid subclass intensity between groups and Tukey's multiple comparisons test was used for post-hoc pairwise comparisons (n=6 mice per group, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, Tukey's multiple comparisons test). To correct for multiple testing the Benjamini, Krieger and Yekutieli procedure was used with a false discovery rate of 5% (N=29 subclasses), FDR adjusted p-values are reported as q values. PI: $F_{(4, 25)} = 4.316$, $**P=0.0086$, q=0.0338, PA: $F_{(4, 25)} = 5.537$, $**P=0.0025$, q=0.0184, BisMePS: $F_{(4, 25)} = 9.308$, $****P<0.0001$, q=0.0022, PIP2: $F_{(4, 25)} = 3.743$, $*P=0.0161$, q=0.0394, BiotinylPE: $F_{(4, 25)} = 3.446$, $*P=0.0225$, q=0.0451, LPE: $F_{(4, 25)} = 3.957$, $*P=0.0127$, q=0.035. TG: $F_{(4, 25)} = 8.593$, $***P=0.0002$, q=0.0022, MGMG: $F_{(4, 25)} = 2.765$, $*P=0.0497$, q=0.0887, ns, MGDG: $F_{(4, 25)} = 3.491$, $*P=0.0214$, q=0.0451. CerP: $F_{(4, 25)} = 5.185$, $**P=0.0035$, q=0.0193, Hex2Cer: $F_{(4, 25)} = 4.076$, $*P=0.0112$, q=0.035, ZyE: $F_{(4, 25)} = 4.256$, $**P=0.0092$, q= 0.0338.

Abbreviations:
Glycerophospholipids: PC = phosphatidylcholine, PE = phosphatidylethanolamine, PS = phosphatidylserine, PI = phosphatidylinositol, MePC = methylphosphocholine, PA = phosphatidic acid, BisMePA = bis-methyl phosphatidic acid, PG = phosphatidylgylcerol, BisMePS = bis-methylphosphatidylserine, PIP2 = phosphatidylinositol-bisphosphate, BiotinylPE = biotinyl-phosphoethanolamine, LPE = lysophosphatidylethanolamine, LPC = lysophosphatidylcholine, LPS = lysophosphatidylserine
Glycerolipids: TG = triacylglycerol, DG = diacylglycerol, MGMG = monogalactosylmonoacylglycerol, MGDG = monogalactosyldiacylglycerol
Sphingolipids: Hex1Cer = monohexosylceramide, SM = sphingomyelin, CerP = ceramide phosphate, Cer = ceramide, ST = sulfatide, SPH = sphingoid base, Hex2Cer = dihexosylceramide
Sterol lipids: ChE = cholesterol ester, ZyE = zymosterol
Fatty acyls: AcCa = acyl carnitine
Prenol lipids: Co = coenzyme

Supplementary Figure 12

Supplementary Figure 12. Individual lipid species significantly changed between 12-month-old LacQ140 and WT mice.

Lipids were extracted and analyzed by LC-MS/MS. Graphs show relative intensities for indicated lipid species expressed as a percent of total lipid intensity per sample for each genotype or treatment group. Plotted values represent individual lipid species intensity standardized to total amount of lipid detected in the same sample. Bar charts underneath individual points represent group means and error bars are ± standard deviation. One-way analysis of variance (ANOVA) was used to evaluate differences in lipid species intensity between groups and Tukey's multiple comparisons test was used for post-hoc pairwise comparisons (n=6 mice per group, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$, Tukey's multiple comparisons test). To correct for multiple testing over all lipids analyzed, the Benjamini, Krieger and Yekutieli procedure was used with a false discovery rate of 5% (N=735 lipid species), FDR adjusted p-values are reported as q values. PC 30:0: $F_{(4, 25)}$ = 7.601, $***P=0.0004$, q=0.0132, PC 20:3e_18:0: $F_{(4, 25)}$ = 5.567, $**P=0.0024$, q=0.0327, PE 16:0p_20:4: $F_{(4, 25)}$ = 7.879, $***P=0.0003$, q=0.0114, PE 20:1_18:1: $F_{(4, 25)}$ = 10.36, $****P<0.0001$, q=0.0038, PA 36:2e: $F_{(4, 25)}$ = 5.630, $**P=0.0023$, q=0.0327, PA 36:3e: $F_{(4, 25)}$ = 13.27, $****P<0.0001$, q=0.0007, TG 16:0_18:1_18:2: $F_{(4, 25)}$ = 8.357, $***P=0.0002$, q=0.0094, TG 18:1_18:1_18:2: $F_{(4, 25)}$ = 7.113, $***P=0.0006$, q=0.0182, TG 18:1_18:2_18:2: $F_{(4, 25)}$ = 6.249, $**P=0.0013$, q=0.0272, TG 16:0_16:1_18:1: $F_{(4, 25)}$ = 5.794, $**P=0.0019$, q=0.0306, TG 16:1_18:1_18:2: $F_{(4, 25)}$ = 4.983, $**P=0.0043$, q=0.0438, TG 16:0_16:0_18:1: $F_{(4, 25)}$ = 7.833, $***P=0.0003$, q=0.0114, TG 19:1_18:1_18:1: $F_{(4, 25)}$ = 6.060, $**P=0.0015$, q=0.0272, TG 18:0_16:0_18:1: $F_{(4, 25)}$ = 6.726, $***P=0.0008$, q=0.0217, TG 18:2_18:2_18:2: $F_{(4, 25)}$ = 5.352, $**P=0.0030$, q=0.0345, TG 56:3: $F_{(4, 25)}$ = 6.109, $**P=0.0014$, q=0.0272, Hex1Cer d18:1_18:0: $F_{(4, 25)}$ = 14.49, $****P<0.0001$, q=0.0005, Hex1Cer d18:2_22:0: $F_{(4, 25)}$ = 5.594, $**P=0.0023$, q=0.0327, Hex1Cer d18:1_22:1: $F_{(4, 25)}$ = 6.637, $***P=0.0009$, q=0.0219, Hex1Cer t18:0_22:1: $F_{(4, 25)}$ = 6.181, $**P=0.0013$, q=0.0272, Hex1Cer d40:2: $F_{(4, 25)}$ = 9.257, $****P<0.0001$, q=0.0058, Hex1Cer d41:2: $F_{(4, 25)}$ = 7.880, $***P=0.0003$, q=0.0114, SM d18:1_23:0: $F_{(4, 25)}$ = 5.007, $**P=0.0042$, q=0.0438, SM d41:1: $F_{(4, 25)}$ = 6.980, $***P=0.0006$, q=0.0197, ST d18:1_22:1: $F_{(4, 25)}$ = 6.263, $**P=0.0012$, q=0.0272, CerP t40:3: $F_{(4, 25)}$ = 14.26, $****P<0.0001$, q=0.0005.

Abbreviations:
Glycerophospholipids: PC = phosphatidylcholine, PE = phosphatidylethanolamine, PA = phosphatidic acid
Glycerolipids: TG = triacylglycerol
Sphingolipids: Hex1Cer = monohexosylceramide, SM = sphingomyelin, ST = sulfatide, CerP = ceramide phosphate

Supplementary Figure 13

a

b

c

Supplementary Figure 13. DEGs, reversed genes in 6- and 12-month-old LacQ140 mice
(a) Bar chart shows number of differentially expressed genes (DEGs) at 6-months (left) and 12-months (right). Total number of DEGs in LacQ140 mice (upregulated: red and downregulated: blue) are shown in the top bar; lower bars show number of DEGs in mice with mHtt repression (LacQ140_8M, LacQ140_2M, LacQ140_A) compared to LacQ140 mice (i.e., genes "reversed" with m*Htt* repression). **(b)** Venn diagram depicts genes "reversed" at 6-months for each respective m*Htt* repression group as shown in **Figure 6c**. **(c)** Venn diagram depicts genes "reversed" at 12-months for each respective m*Htt* repression group as shown in **Figure 6f**.