Electrophysiological signatures of veridical head direction in humans

Benjamin J. Griffiths[1,2], Thomas Schreiner[1], Julia Schaefer[1], Christian Vollmar[3], Elisabeth Kaufmann[3], Stefanie Quach[4], Jan Remi[3], Soheyl Noachtar[3] & Tobias Staudigl[1]

1. Department of Psychology, Ludwig-Maximilians-Universität München, Germany.
2. Centre for Human Brain Health, University of Birmingham, United Kingdom.
3. Epilepsy Center, Department of Neurology, Ludwig-Maximilians-Universität München, Germany.
4. Department of Neurosurgery, University Hospital Munich, Ludwig-Maximilians-Universität München, Germany.

Abstract

Information about heading direction is critical for navigation as it provides the means to orient ourselves in space. However, given that veridical head direction signals require physical rotation of the head and most human neuroimaging experiments depend upon fixing the head in position, little is known about how the human brain is tuned to such heading signals. To address this, we asked fifty-two healthy participants undergoing simultaneous EEG and motion tracking recordings (split into two experiments) and ten patients undergoing simultaneous intracranial EEG and motion tracking recordings to complete a series of orientation tasks in which they made physical head rotations to target positions. We then used a series of forward encoding models and linear mixed-effects models to isolate electrophysiological activity that was specifically tuned to heading direction. We identified a robust posterior central signature that predicts changes in veridical head orientation after regressing out confounds including cuing, visual input and muscular activity. Both source localisation and intracranial analysis implicated the medial temporal lobe as the origin of this effect. Subsequent analyses disentangled head direction signatures from signals relating to head rotation and those reflecting location-specific effects. Together, these results reveal a taxonomy of population-level head direction signals within the human brain that is reminiscent of those reported in the single units of rodents.

Supplementary materials: For convenience, the supplementary materials have been appended to this document, and can be found after the reference section.

Code availability: Openly available at https://github.com/benjaminGriffiths/human-hd.

Data availability: Data acquired from the healthy participants will be made available upon publication. Due to concerns about patient anonymity, data acquired from the patients can be provided upon reasonable request.

Acknowledgements: This work was supported by the European Research Council (https://erc.europa.eu/, Starting Grant 802681 awarded to TSt) and the Leverhulme Trust (https://www.leverhulme.ac.uk/, Early Career Fellowship ECF-2021-628 awarded to BJG). We thank all participants and in particular all patients who volunteered to participate in this study. We thank the staff and physicians at the Epilepsy Center, Department of Neurology, Ludwig-Maximilians-Universität, Munich, for assistance.

Competing interests: The authors declare no competing interest.

Introduction

Navigation is a complex cognitive phenomenon of which two basic forms can be distinguished: map-based and self-referential navigation (Buzsáki & Moser, 2013). While many have delved into how the human brain represents cognitive maps (Baumann & Mattingley, 2021; Bellmund et al., 2018; Epstein et al., 2017), comparatively few have explored how the human brain comprehends a sense of direction, core ingredient to both basic form of navigation. Consequently, our understanding of a cornerstone of human navigation is lacking. To redress this balance, we set out to explore how human electrophysiological activity is tuned to veridical (that is, true physical) head direction.

Our understanding of the neural code for veridical head direction is largely based on rodent studies. This work has demonstrated that single units across the rodent brain are tuned to current head direction (Muller et al., 1996; Ranck, 1984; Sharp et al., 2001; Taube, 2007; Taube & Bassett, 2003). These units selectively fire when the rodent faces a particular angle in the environment, regardless of the physical location of the rodent in the environment (Taube et al., 1990; Taube, 1995). Their firing is persistent over time (Taube & Bassett, 2003; Taube & Muller, 1998), continues in the absence of visual cues (Goodridge et al., 1998), and often precedes physical head rotation (Blair & Sharp, 1995; Zirkelbach et al., 2019). As head direction cells are found in abundance in subcortical and cortical brain regions (Hulse & Jayaraman, 2020), a coordinated population signal could be detectable in macroscopic electrophysiology (Kunz et al., 2019; Nau et al., 2020). Importantly, this representation of head direction is essential for navigation as the perturbation of the network destabilises representations of space (Butler et al., 2017; Calton et al., 2003; Winter et al., 2015). These findings have led to the conclusion that head direction acts as a 'neural compass' during spatial navigation (Winter et al., 2015).

Examining this neural compass in humans is not a trivial task. Imaging methods such as magnetoencephalography (MEG) and magnetic resonance imaging (MRI) require the body to be fixed in position to minimise artefacts. This prevents the generation of most self-referential cues, which is thought to be a profound problem for the study of navigation (Steel et al., 2021; Taube et al., 2013). Electroencephalography (EEG), however, has no such limitation. It has been used to successfully measure human brain activity during motor (e.g., Do et al., 2021; Gramann et al., 2021) and cognitive tasks (e.g., Griffiths et al., 2016; Piñeyro Salvidegoitia et al., 2019) while the body is in motion. As such, it is well suited to the exploration of how the human brain represents veridical head direction.

Here, we asked whether a signature of veridical head direction can be identified in the human scalp and intracranial EEG. Participants completed a series of orientation tasks in which they made physical head rotations between multiple computer monitors, with each task containing a unique manipulation that helped disentangle head direction from other confounds (see figure 1a-b). We used forward encoding models to predict intracranial and scalp EEG activity as a function of simultaneously recorded heading angle, and then used linear mixed-effect models to localise the electrophysiological signals that were specifically tuned to a given heading angle. We find converging evidence for a precisely tuned representation of head direction that emanates from posterior regions including the parietal cortex and parahippocampus, and shares numerous similar properties to the head direction cells of rodents (as described above).

Results

Scalp EEG activity is tuned to heading angle

In the first experiment, 32 participants completed four orientation tasks (see figure 1), each with distinct manipulations used to delineate head direction-related activity from activity

Figure 1. Experiment outline. (A) Participants completed a series of orientation tasks presented across five computer monitors. Participants began by fixating on a cross on the centre screen. This cross then moved to one of the other screens. In most tasks, participants were tasked with physically rotating their head towards the target screen. In most tasks, an auditory cue accompanied the movement of the fixation cross to help participants to track the cross. After a delay sufficient to allow participants to move into position and fixate, the cross returned to the centre screen. In experiment 2, an additional two screens were placed to the right of the five-monitor setup, with an angular orientation of 90° and 120° relative to the initial centre screen. **(B)** Three groups of participants completed four variations of these tasks. The first group of healthy participants and the group of patients completed Experiment 1. The second group of healthy participants completed Experiment 2. For full details of what each task entailed, see methods.

attributable to sensory cuing and visual input/saccades (see supplementary figure 1 for head
rotation time-series). We used cross-validated forward encoding models (FEM) to predict EEG
activity, with head motion recordings decomposed into basis sets of circular–Gaussian von-
Mises distributions (from here on termed "kernels") acting as model features. Six of these
FEMs were built, each using different kernel sizes (ranging from 10° to 60°) which allowed us
to approximate the precision of the tuning of the EEG signal to current heading angle. FEM
performance was quantified by correlating the FEM-predicted EEG signal with the observed
EEG signal. FEM performance was pooled across participants and linear mixed-effect models
were used to assess how much variance current heading angle explained of the observed
FEM performance after controlling for the confounds of auditory cuing (i.e., animal sounds that
pointed participants towards the target screens), visual input, saccadic activity, and muscular
activity (see methods for details).

For all tuning widths, EEG signal appeared to be tuned to changes to heading angle, and
to a degree significantly greater than chance (peak $z = 6.747$, $p_{FDR} < 0.001$; see figure 2a-b).
FEM performance was at its greatest over posterior central sensors and when using a kernel
width of 20° (see supplementary table 1 for statistics relating to all kernel widths and
regressors). When applying the FEM to source-space data, late visual and medial temporal
regions responded most strongly to changes in heading angle (see figure 2c). While EEG
activity was also tuned to both saccadic activity/visual input (peak $z = 2.731$, $p_{FDR} < 0.001$) and
muscular activity (peak $z = 2.958$, $p_{FDR} < 0.001$), the sizes of the effects were substantially
smaller than that of head direction (see figure 2a-b). Moreover, it is important to note that the
nature of the linear mixed-effect models means that these effects were statistically
independent of those for head direction. Altogether, this suggests that activity present over the
posterior central EEG is tuned to changes in heading angle – an effect that cannot be attributed
to cuing, muscular activity or visual input/saccadic activity.

Next, we examined the temporal dynamics of this tuned EEG activity, asking whether the
EEG activity precedes or follows physical changes in heading angle. We first built a "lagged"
encoding model, where the EEG time series was temporally shifted relative to the motion
tracking time series prior to building the FEM. By repeating this for multiple temporal lags, we
were able to build a time series which identified when the FEM best predicted EEG activity.
Here, it seemed that the lagged encoding model was best able to predict EEG activity when
considering EEG activity that preceded heading angle information by approximately 90ms
(peak $z = 9.415$, $p_{FDR} < 0.001$; see figure 2d). This aligns with animal and computational work
suggesting that signatures of head direction are anticipatory rather than responsive (Blair &
Sharp, 1995; Zirkelbach et al., 2019). We complemented our lagged FEM with an inverted
encoding model, which takes the weights from the training set and inverts them to produce an
estimate of heading angle for every sample. Representational similarity analysis (Kriegeskorte,

Figure 2. Electrophysiological activity tracks change in head angle. (A) Plot of the regressor coefficients (averaged over all scalp electrodes) derived from linear mixed effects models used to predict forward encoding model performance. Dots indicate the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. Colour indicates the tuning width of the kernel used in the forward encoding model, with darker colours representing the narrower tuning widths. For the head direction regressor, all tuning widths were significant ($p_{FDR} < 0.05$). No other regressor produced significant effects. **(B)** Topographic plots of the regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width (i.e., the best performing forward encoding model). Deeper red colours indicate larger coefficient values. Black dots indicate electrodes that reliably map onto forward encoding model predictions ($p_{FDR} < 0.05$). Each of the four regressors are plotted separately. **(C)** Source plots of the regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width. Deeper red colours indicate larger coefficient values. Each of the three regressors are plotted separately. Note that the EMG regressor was not included here as the process of beamforming will inherently project EMG components to outside the brain. **(D)** Time-series of regressor coefficients derived from linear mixed effects models used to predict lag-based forward encoding model performance. The deep red indicates the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. The dot indicates the lag at which predictive power of the regressor was greatest. **(E)** Time-series of regressor coefficients derived from linear mixed effects models used to predict inverted encoding model performance. The deep red indicates the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. The shaded grey areas indicate significance ($p_{FDR} < 0.05$).

2008) was then used to see when the predicted head angle differed most greatly between the
four outside monitor positions. In line with the lagged FEM, the inverted encoding model
showed that the predictive power of EEG activity for heading angle ramped up just before the
head began to rotate (peak z = 7.743, $p_{FDR} < 0.001$; see figure 2e). EEG activity continued to
predict heading angle throughout the head rotation and beyond as participants maintained the
new heading angle (albeit to a slightly lesser degree). This aligns with animal work showing
persistent but weaker head direction tuning after head rotation (Taube & Bassett, 2003).
Together, these results suggest that tuned EEG activity precedes the change in veridical
heading angle and is sustained after the head rotation is completed.

Figure 3. Intracranial electrophysiological activity tracks change in heading angle. (A) Plot of the head direction regressor coefficient derived from linear mixed effects models used to predict forward encoding model performance. Dots indicate the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. Colour indicates the tuning width of the kernel used in the forward encoding model, with darker colours representing the narrower tuning widths. For the frontal, parietal and temporal lobes, all tuning widths were significant ($p_{FDR} < 0.05$). **(B)** Time-series of regressor coefficients derived from linear mixed effects models used to predict lag-based forward encoding model performance. Plots are restricted to those regions which demonstrated significant predictive power in the previous analysis. The deep coloured line indicates the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. The dot indicates the lag at which predictive power of the regressor was greatest. **(C)** Details of plot match that of panel A, but here plotted for subregions of the medial temporal lobe. Only the parahippocampus demonstrated significant predictive power for tracking changes in head orientation ($p_{FDR} < 0.05$). **(D)** Details of plot match that of panel B, but here plotted for parahippocampal subregion of the medial temporal lobe.

Intracranial EEG activity is tuned to heading angle

Many key regions expressing head direction cells are found deep within the brain, which can be difficult to reliably measure in scalp EEG recordings. Therefore, we set out to repeat the first experiment in a group of ten patients with intracranial depth electrodes. The methodological and analytical approaches matched that of the scalp EEG experiment, with the single exception of the linear mixed-effect models. As patients have bespoke implantation schemes, models cannot be built for individual electrodes as done above. Therefore, we built models for several regions-of-interest (ROI) instead; one ROI for each cortical lobe of the brain, and one for each recorded subregion of the medial temporal lobe (a decision driven by the fact that subregions of the medial temporal lobe have unique functions in navigation; e.g., Baumann & Mattingley, 2021) [see methods for details]. As before, individual patients were modelled as random effects.

When probing individual lobes, all regions of interest demonstrated a tuned response to current heading angle except the occipital lobe (frontal lobe: peak tuning width = 10°, z = 6.697, $p_{FDR} < 0.001$; parietal lobe: peak tuning width = 15°, z = 3.388, $p_{FDR} < 0.001$; occipital lobe: peak tuning width = 10°, z = 2.156, $p_{FDR} = 0.079$; temporal lobe: peak tuning width = 10°, z = 5.836, $p_{FDR} < 0.001$; see figure 3a). Intriguingly, the recordings analysed here were more precisely tuned to current heading angle than the scalp EEG recordings – a possible effect of the bipolar referencing scheme used on the intracranial recordings isolating minute differences between pairs of contacts. As with the scalp recordings, lag-based models of the intracranial recordings suggested that changes in EEG signal preceded changes in head angle (frontal lobe: peak lag = 80ms, z = 7.475, $p_{FDR} < 0.001$; parietal lobe: peak lag = 100ms, z = 4.940, $p_{FDR} < 0.001$; peak lag = 130ms, z = 5.619, $p_{FDR} < 0.001$; see figure 3b).

When probing the subregions of the medial temporal lobe, only the parahippocampus demonstrated any substantial tuning to current head angle (parahippocampus: peak tuning width = 15°, z = 2.879, p_{FDR} = 0.012; hippocampus: peak tuning width = 15°, z = 2.108, p_{FDR} = 0.079; amygdala: peak tuning width = 15°, z = 1.843, p_{FDR} = 0.088; see figure 3c). As with the lobe-level analysis, the lag-based models of the parahippocampal signal suggested that changes in parahippocampus signal preceded changes in head angle (peak lag = 60ms, z = 3.634, p_{FDR} < 0.001; see figure 3d).

Taken together, these results corroborate what was observed in the scalp EEG recordings. Namely, several different regions are tuned to changes in head angle, including the parietal lobe and parahippocampus. Moreover, these regions all appeared to predict upcoming changes in head angle, suggesting that they anticipate rather than respond to a change in head angle. Within the medial temporal lobe, parahippocampus but not hippocampus showed substantial directional tuning, which aligns with animal work demonstrating HD cells outside hippocampus (Cullen & Taube, 2017).

A conceptual replication of heading angle-tuned scalp EEG activity

In the second experiment, 20 participants completed four orientation tasks that varied the position of the participants relative to the monitors. Two of these tasks involved sitting while the others involved standing (introducing a position change in the z-axis). One of the standing tasks also involved standing one metre away from the standard location (introducing a location change in the x-axis), while one of the sitting tasks involved sitting in the same location but rotated by 60° (a condition that will be discussed in more detail in the last results subsection). In the first instance, we used the same approach to forward encoding modelling as in Experiment 1 to both replicate and extend those findings. Once again, linear mixed-effect models were used to delineate the influence of head angle from potential confounds, in this case: position in the z-axis, location in the x-axis, and muscular activity.

Neatly mapping onto the findings from Experiment 1, we found that, after regressing out the influence of location and muscular activity, scalp EEG recordings demonstrated a tuning to current head angle significantly beyond what would be expected by than chance (peak z = 7.40, p_{FDR} < 0.001; see figure 4a-b; see supplementary table 2 for statistics relating to all kernel widths and regressors). These effects were most prominent over posterior central electrodes and were localised to occipital and medial temporal regions (albeit not reaching as far anteriorly as in Experiment 1; see figure 4c). Both the lag-based forward encoding model and the inverted encoding model also produced results matching Experiment 1, suggesting that changes in EEG activity precede changes in head angle (lag model: peak z = 5.65, p_{FDR} < 0.001; inverted model: peak z = 4.08, p_{FDR} < 0.001; see figure 4d-e).

The findings of Experiment 1 and their replication reported above suggest that human EEG activity is tuned to changes in current heading angle. However, it would be premature to suggest that, based on these results alone, this reflects a human analogue of the prototypical head direction signal observed in rodents. Specifically, we must probe whether the head direction effects uncovered here are (1) consistent over changes in location, and (2) separable from effects attributable to head rotation. The next two subsections address these issues in turn.

Tuned scalp EEG activity persists across locations within an environment

To delineate the effects of location-independent and location-specific head direction signals, we adapted the forward encoding modelling approach to consider how the models generalised across tasks. This generalisation was achieved by training the FEM on one task and applying it to data from another task. For EEG electrodes where the FEMs do generalise

Figure 4. Electrophysiological activity tracks change in head angle independent of location. (A) Plot of the regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance. Dots indicate the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. Colour indicates the tuning width of the kernel used in the forward encoding model, with darker colours representing the narrower tuning widths. For the head direction regressor, all tuning widths were significant ($p_{FDR} < 0.05$). No other regressor produced significant effects. **(B)** Topographic plots of the regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width (i.e., the best performing forward encoding model). Deeper green colours indicate larger coefficient values. Black dots indicate electrodes that reliably map onto forward encoding model predictions ($p_{FDR} < 0.05$). Each of the four regressors are plotted separately. **(C)** Source plots of the regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width. Deeper green colours indicate larger coefficient values. Each of the three regressors are plotted separately. Note that the EMG regressor was not included here as the process of beamforming will inherently project EMG components to outside the brain. **(D)** Time-series of regressor coefficients derived from linear mixed effects models used to predict lag-based forward encoding model performance. The deep green line indicates the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. The dot indicates the lag at which predictive power of the regressor was greatest. **(E)** Time-series of regressor coefficients derived from linear mixed effects models used to predict inverted encoding model performance. The deep green line indicates the estimated regressor coefficients. Error bars indicate the 95% confidence intervals. The shaded grey areas indicate significance ($p_{FDR} < 0.05$).

to tasks completed in differing locations (that is, the FEM performs better than what would be
expected by chance), one could conclude that the EEG signal predicted by the FEM is not
dependent on location. For electrodes where the models perform well within a given task but
do not generalise, one could conclude that the head angle effect is specific to a given location.
Linear mixed-effects models were used to explore the influence of location on head direction
signal, as well as regress out the influence of muscular activity (as done in the previous
analyses).

Figure 5. Electrophysiological representations of location-independent and location-dependent head direction signals. **(A)** Topographic plots of the two regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width (i.e., the best performing forward encoding model). Deeper green colours indicate larger coefficient values. Black dots indicate electrodes that reliably map onto forward encoding model predictions (p_{FDR} < 0.05). **(B)** Source plots of the two regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width. Deeper green colours indicate larger coefficient values.

This approach revealed that the generalised FEMs could reliably predict posterior central
EEG activity (peak z = 6.21, p_{FDR} < 0.001; see figure 5a), suggesting they make use of location-
independent heading angle information. Analysis of source-space data suggested these
effects originated from right retrosplenial and left inferior frontal areas (see figure 5b). Notably,
the linear mixed-effects models also found evidence to suggest some location-specific head
direction-related signals exist within the EEG (peak z = 5.73, p_{FDR} < 0.001). These effects were
positioned slightly further back on the EEG cap, with source analysis implicating the left
posterior parietal cortex (see figure 5a-b). These results suggest that distinct parts of the EEG
signal are tuned to location-independent and location-specific signatures of current heading
angle.

Tuned scalp EEG activity is distinct from head rotation-related EEG activity

A similar approach was used to delineate EEG activity tuned to head direction (i.e., that
anchored to the environment) from EEG activity tuned to head rotation (i.e., that anchored to
the body). In this instance, FEMs from one sitting condition were generalised to the other sitting
condition (where participants were rotated 60° to the right in the environment), and vice versa.
If the FEMs generalise, it would indicate that the effect is driven by head rotation (as the model
would be insensitive to the 60° rotation of the entire body within the environment). In contrast,
any part of the FEM that does not generalise would reflect head direction. Here, the linear
mixed-effects models suggested that the head direction effects in previous analyses were a
combination of "true" head direction signals and those tied to the rotation of the head. The
"true" head direction signals were the most prominent over temporal electrodes (peak z = 5.37,
p_{FDR} < 0.001; see figure 6a), with source-space analyses implicating the medial temporal lobe
and left frontal regions (see figure 6b). The head rotation signals were most prominent over
posterior central electrodes (peak z = 4.11, p_{FDR} < 0.001), and were localised to the posterior

Figure 6. Electrophysiological representations of environment-based head direction and body-based head rotation signals. (A) Topographic plots of the two regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width (i.e., the best performing forward encoding model). Deeper blue colours indicate larger coefficient values. Black dots indicate electrodes that reliably map onto forward encoding model predictions (p_{FDR} < 0.05). **(B)** Source plots of the two regressor coefficients derived from linear mixed effects models used to predict forward encoding model performance when using a 20° tuning width. Deeper blue colours indicate larger coefficient values.

196 parietal cortex (see figure 6a-b). These results suggest that both "true" head direction signals
197 and signals relating to head rotation are present in EEG activity.

Discussion

To date, it has remained unclear where the human brain is tuned to changes in veridical
head direction. To remedy this, we asked participants to complete a series of head rotation
tasks while undergoing scalp or intracranial EEG recordings with the aim of isolating the neural
signatures of human veridical head direction. Using a series of analytical models, we identified
a signature tuned to current heading angle that is independent from auditory cuing, visual input
and muscular activity. Anatomically speaking, both source-localised scalp EEG and
intracranial EEG implicated a wide network of regions in this tuning, including the medial
temporal lobe and parietal cortex. These findings were replicated in a second, independent
sample of healthy participants, who completed a series of additional tasks that regressed out
the influence of head rotation and location-specific effects to further isolate a neural signature
specifically related to veridical head direction. Altogether, these results provide a detailed
taxonomy of head direction-related signals that can be observed in free-moving human
participants.

On the similarities and dissimilarities to head direction cells found in rodents

For almost four decades, animal research has steadily advanced our knowledge of the
electrophysiological signatures of veridical head direction signals (for reviews, see Dudchenko
et al., 2019; Hulse & Jayaraman, 2020; Taube, 2007). Here, we aimed to expand this work to
electrophysiological population activity in humans and found numerous similarities (and the
occasional idiosyncrasy) to what has been reported in animal models. Below, we compare our
results from human population-level activity against what could be considered some of the
main features of rodent head direction cells:

1. *Anatomical overlap*: We observed directional tuning to current head angle in many
regions across the cortex. The strongest of these effects arose in the medial temporal
lobe, where head direction cells are observed in abundance (for reviews, see
Dudchenko et al., 2019; Hulse & Jayaraman, 2020; Taube, 2007). In line with the
majority of animal work, we observed directional tuning outside but not in the
hippocampus (see Cullen & Taube, 2017). That said, we observed no effect in the
anterior thalamus, the area that is arguably most densely packed with head direction
cells (Taube & Bassett, 2003). Perhaps direct intracranial recordings of the anterior
thalamus are better suited to reveal similar directional tuning as to what is reported in
other brain regions of humans, and the anterior thalamus of rodents. We did, however,
observe tuning effects in the frontal lobe, a region not featured in many maps of the
head direction network (e.g., Taube, 2007). During navigation, activity in the frontal lobe
has been attributed to several functions, such as goal tracking and route planning (Patai
& Spiers, 2021). Given that our experiment involved tracking the target direction and
intentionally orienting towards the target, the heightened tuning of frontal activity to
current heading angle (relative to tuning during free exploration in animal studies) is
perhaps not surprising.

2. *Independence from visual input*: Directional tuning of head direction cells is maintained
when visual input is removed (Goodridge et al., 1998). In our experiment, using virtual
reality goggles containing nothing but a fixation cross (eliminating changes in visual
input that could be used as visual cues as well as minimizing eye movements) and
linear mixed-effect modelling, we too could separate the influence of visual input and
saccade activity from that of head direction-related activity. While visual input-related
activity was centred on superior occipital and inferior parietal regions, activity tuned to

current head direction was centred in the medial temporal lobes. The fact that
population-level directional tuning was sustained in the absence of visual input brings
the current results further in line with the traditional definition of head direction-related
activity.

3. *Independence from location*: Cells tuned to head direction can be classified as
"traditional" head direction cells that fire independently of the animal's location within
an environment or "sensory" cells that are dependent on sensory input (Dudchenko et
al., 2019). We observed similar dissociable patterns of activity that relate to these two
signatures of head direction: "traditional" location-independent head direction-related
activity stemmed from the more anterior portion of the medial temporal lobe and left
inferior frontal gyrus, while location-specific head direction-related activity was more
closely related to activity in the posterior parietal cortex and retrosplenial cortex.
Notably, this spatial delineation overlaps with work in rodents, which identified location-
sensitive head direction-related activity within the retrosplenial cortex and its thalamic
connections (Jacob et al., 2017; Lomi et al., 2023) . Evidence that population-level
directional tuning was sustained across locations further supports the idea that a
portion of the tuned activity we observed matches that of "traditional" head direction
activity.

4. *Anchoring to the environment*: Head direction cells are anchored to the position of the
head within the environment, not to the position of the head with respect to the body
(Taube et al., 1990). We separated "body-based" tuning to head rotation from more
traditional head direction effect through experimental and statistical means and
identified distinct hubs of activity for the two processes. Once again, head direction-
related activity was most prevalent in the medial temporal lobes. In contrast, "body-
based" rotation-related activity was observed in the posterior parietal cortex (see figure
6b), aligning with reports that posterior parietal lesions do not impact tuning to head
direction (Calton et al., 2008) and supporting the idea that the posterior parietal cortex
is instead involved in translational updating (Medendorp et al., 2008).

5. *Anticipatory signatures*: In line with animal research, we observed the tendency for
tuned signatures of head direction to precede the physical head rotation (e.g., Blair et
al., 1998; Stackman & Taube, 1998). Intriguingly, the delay between neural signature
and physical head rotation observed here most closely matched the delay exhibited by
the lateral mammillary nucleus (~40-100ms; positioned at the early stages of the head
direction network) and not the delay exhibited by the postsubiculum (located later in
the head direction network hierarchy, showing almost instantaneous responses to
changes in head direction; Blair & Sharp, 1995), despite the regions identified in our
task being situated closer to the postsubiculum than the lateral mammillary nucleus. It
is unclear why this is the case, but perhaps (as with the frontal lobe effects discussed
above) the fact that participants were cued to rotate to a particular target meant that
anticipatory responses are earlier than those observed during free exploration.

6. *Tuning widths*: In rodents, individual head direction cells show a directional tuning of
around 60 to 120° (Taube, 1995; Taube et al., 1990). Population-level tuning observed
here, however, was substantially more precise (~10-20°; aligning with similar reports
from VR studies of human directional tuning; Nau et al., 2020). This discrepancy may
relate to the fact that tuning width tends to become more precise as one progresses
further down the head direction network hierarchy (Bassett & Taube, 2001; Stackman
& Taube, 1998). Given that scalp EEG will principally pick up on cortical activity at the
bottom end of this hierarchy, it is perhaps no surprise that the tuning we observed is
more precise. This idea is supported by the intracranial data suggesting that the medial
temporal lobe is more coarsely tuned (~15°) than neocortical sources (e.g., ~10° tuning

in frontal lobes), as well as 7T fMRI work suggesting a gradient of tuning widths along
the visual ventral stream which becomes coarser as one approaches the medial
temporal lobe (Nau et al., 2020).

7. *Persistence in the absence of movement*: In rodents, head direction cells fire persistently for at least ~600ms after head rotation has ceased (Taube & Muller, 1998). Our inverted encoding models came to similar conclusions, suggesting that tuned population activity is sustained for ~500ms after the offset of the head rotation (see figure 2e). Notably, however, tuned activity dissipates shortly after this. While not completely contradictory to the observed decline in head direction cell firing rates that accompany immobility (Shinder & Taube, 2014), the result is hard to reconcile with influential computational models which suggest that head direction-related activity should be persistent in the absence of movement (for review, see Taube & Bassett, 2003). We speculate this decline in tuning relates to participants anticipating the return to the centre screen, which would produce a signature of the imagined upcoming rotation (Shine et al., 2016), impairing the inverted encoding model's ability to predict current head angle.

8. *Population-level activity*: The most egregious dissimilarity between our results and those reported in rodents is the difference between population-level tuning and single-cell tuning. To what extent is tuned population-level activity reported here comparable to single-cell tuning? We speculate that there is a close relationship, with population-level activity arising as an emergent property of a network of head direction cells synchronising their action potentials. Such population-level activity may be essential for propagating head direction-related information across the head direction network hierarchy as synchronous firing would ensure neurons effectively impact downstream regions in the hierarchy (Hulse & Jayaraman, 2020). This synchronisation may be facilitated by theta (6-12Hz) and high-frequency (~150Hz) neural oscillations, which have been shown to co-ordinate head direction-related activity in rodents (Butler & Taube, 2017; Peyrache et al., 2015; Tsanov et al., 2011). Whether the level of synchronicity between head direction cells is relevant for behaviour, however, remains an intriguing open question.

Taken together, population-level activity tuned to current heading angle in free-moving humans shares several similar properties with individual head direction cells in animal models. However, open questions remain. Some pertain to the discrepancies between what was observed here and what has been reported in previous studies (see above). Others relate to facets of head direction cells not explored here. For example, head direction cells continue to fire when rodents are passively moved through an environment (e.g., Shinder & Taube, 2011), but it remains unknown if the same is true for population-level activity in humans. Moreover, the goal-directed nature of our task differs from free exploration used in rodent studies (and may explain differences related to anticipatory timing and persistence). It would be of interest to compare human population-level tuning to rodent single-unit tuning in tasks involving free exploration. Nonetheless, we feel that the similarities outweigh the differences, leading us to suggest that we can measure an analogue of head direction in population-level activity of free-moving humans.

On the link to existing research into human head direction-related signals

To date, our understanding of how the human brain represents head direction information comes from two distinct methodologies: one fixing the head and body in place, mostly for fMRI, and exploring directional tuning in virtual reality (VR), the other incorporating full-body and head movements with EEG recordings (also known as "mobile brain/body imaging"; MoBI) to assess how the brain tracks naturalistic movement.

The combination of fMRI and VR has revealed a whole host of regions (e.g., parahippocampus, retrosplenial cortex, parietal cortex) that are linked to the representation of head direction-related information (e.g., Baumann & Mattingley, 2010; Kim & Maguire, 2019; Vass & Epstein, 2013). Of particular relevance here, a recent study used forward encoding models to model "virtual head direction" information (that is, heading information within a virtual environment) and found evidence for the representation of current heading direction within ventral occipital, medial parietal and medial temporal areas (Nau et al., 2020). However, as participants were immobilised by necessity, the critical contributions of vestibular input to head direction information (Shinder & Taube, 2014; Stackman & Taube, 1997; Yoder et al., 2011) were excluded, questioning whether a full neural signature of head direction could be observed (Steel et al., 2021; Taube et al., 2013). Studies using the rare combination of human intracranial recordings and VR identified direction-related tuning of single units in the medial temporal lobe (Jacobs et al., 2010; Kunz et al., 2021), in line with the localization of the current population-level results.

The use of MoBI is less common, but key studies have explored how brain activity related to head rotation differs from other forms of body movement or when the head is stationary (e.g., Do et al., 2021; Gramann et al., 2021), No study, to our knowledge, however has used such an approach to explore how the brain is tuned to heading direction.

By combining the analytical approaches previously used in fMRI (Nau et al., 2020) with the methodological approaches used in MoBI (e.g., Do et al., 2021; Gramann et al., 2021), we have overcome many of the concerns about assessing head direction information in the human brain (Steel et al., 2021; Taube et al., 2013), and hope to have provided a deeper insight into how the human brain represents veridical head direction information.

Direction-related activity in non-human primates

Compared to the vast literature on head direction cells in rodents, few studies investigated head direction tuning of neural activity in non-human primates. We are aware of one study reporting HD cells in the pre-subiculum (Robertson et al., 1999), and another study demonstrating gravity-anchored HD-like cells in the anterior thalamus (Laurens et al., 2016). Intriguingly, an additional form of directionally-tuned cell has been reported in the entorhinal cortex of macaques; those tuned to saccade direction (Killian et al., 2015). While it was not possible to ultimately determine the reference-frame of these saccade-direction cells, the study does align with a growing body of work in non-human primates indicating tuning of neurons to visual space (Killian et al., 2012; Meister & Buffalo, 2018; Rolls, 1999; Rolls & O'Mara, 1995; Wirth et al., 2017). It would be interesting to investigate in future studies whether neural activity in humans is also tuned to saccade direction, in analogy to what has been shown for grid-like representations of visual space (Julian et al., 2018; Nau et al., 2018; Staudigl et al., 2018).

On the influence of electromyographic (EMG) artifacts

Given the sensitivity of EEG recordings to EMG artifacts, one is right to question whether the results we report here are driven by EMG activity. We would suggest that this is not the case for five reasons:

1. Using a combination of PCA and LMEs on the scalp EEG data, we modelled EMG activity as a statistically independent effect from our main regressors of interest, minimising EMG influence on the central effect.

2. EMG effects could not explain why the scalp EEG activity best predicted head orientation when considering EEG activity ~80ms before head rotation; if the central effect were attributable to EMG, the link between EEG and head orientation should be instantaneous. In the same vein, EMG effects would not be able to explain why scalp

EEG sensitivity to head orientation persisted up to 500ms after the head rotation was completed.

3. EMG effects could not explain why the strongest scalp-level effects were observed at Pz, an electrode in the EEG cap that is one of the furthest away from any source of EMG activity (which arises most commonly around the mastoids and other electrodes at the edge of the cap).

4. In the source-based analysis of the scalp EEG data, we used LCMV beamformers which will project any EMG activity to outside the brain, meaning any sources located within the brain are unlikely to reflect an EMG effect.

5. In the intracranial analysis, we used bipolar re-referencing which will subtract out any common signals between two neighbouring electrodes (including EMG), again meaning that any observed effect is unlikely to reflect an EMG effect.

In any EEG study, even in those with fixed head position, it is difficult, if not impossible, to claim with absolute certainty that there are, emphatically, no EMG effects in the final results and we don't set out to make such a claim here. We do however suggest that, because of the reasons outlined above, any muscular effect would be minimal and, as such, the effects observed here are more easily ascribed to a neural process rather than muscular one.

Summary

In conclusion, we find electrophysiological signatures of veridical head direction in population-level activity of humans that shows a strong correspondence with head direction cell activity reported in rodents. By combining physical head rotations with a series of forward encoding models, we are able to provide a unique insight into the role of vestibular movements in human spatial navigation, overcoming a key limitation that has plagued such study to date (Steel et al., 2021; Taube et al., 2013), and revealing a detailed taxonomy of human head direction-related signals.

Methods

Participants

In Experiment 1, 39 healthy participants were recruited (mean age = 24.8 years, 61.5% female). They received course credit or financial reimbursement in return for their participation. All participants gave written informed consent after being informed about the details of the study. The study was approved by the local ethics committee. Due to recording difficulties stemming from the EEG and motion tracking systems, data from some tasks of some participants was lost. Table 1 contains a breakdown of which participants were excluded from which tasks. For the intracranial recordings, 10 patients undergoing treatment for medication-resistant epilepsy were recruited. The patients, who volunteered to participate in the study, had depth electrodes implanted for diagnostic reasons. The study was approved by the ethics committee of the Medical Faculty of the Ludwig–Maximilian Universität. All patients gave written informed consent after being informed about the details of the study. Difficulties with EEG/motion tracking also resulted in the exclusion of participants (see Table 1). In experiment 2, 24 healthy participants were recruited (mean age = 24.7, 54.2% female). They too received course credit or financial reimbursement in return for their participation and gave written informed consent after being informed about the details of the study. The study was approved by the local ethics committee. See table 1 for details regarding excluded participants.

Experimental design: Experiment 1

Participants arrived at the lab at approximately 7pm in the evening, where they were first fitted with an EEG cap, and then tasked with a series of experiments. At around 11pm, they slept in the lab, and woke at approximately 7am for a second series of experiments. These experiments can be broadly split into

two categories: navigation and memory, both of which happened in the evening session and in the morning session. As this paper only relates to the navigation experiments, only the details of these tasks will be described.

Each navigation experiment followed the same typical pattern. Participants began by fixating on the centre screen. In all experiments, vertical screen positions match participants' line of sight. A stimulus was presented that they had to respond to by turning to one of four other screens (positioned at -60°, -30°, +30° and 60° relative to the centre screen, see figure 1a). Each experiment consisted of 160 trials, split across 4 blocks (i.e., 40 trials per block). Below, each variation of this experiment is outlined in the order they occurred:

1) *Uncued Head Rotation:* At stimulus onset, the fixation cross disappeared and appeared on one of the four flanking screens. Participants were instructed to turn their head to face the screen which the fixation cross appeared on and fixate upon the cross. No auditory cue was played.

2) *Cued Head Rotation:* At stimulus onset and as the fixation moved to one of the four flanking screens, one of four animal sounds were played. Each animal sound was associated with a single flanking screen. Participants were instructed to turn their head to face the screen which the fixation cross appeared on and fixate upon the cross.

3) *Cued Saccades:* Programmatically speaking, this experiment matched the *Cued Head Rotation* exactly. However, participants received different instructions. Specifically, they were instructed to saccade to whichever flanking screen the fixation cross appeared on and fixate upon the cross while keeping their head facing towards the centre screen.

4) *Cued Virtual Reality:* As with the *Cued Saccades* experiment, programmatically speaking, this experiment matched the *Cued Head Rotation* exactly. In this case however, participants wore cardboard virtual reality goggles where a fixation cross was presented directly in front of them. When the auditory cue was played, they were instructed to turn their head to face the screen which they believed was associated with the auditory cue, all the while fixating on the cross presented via the virtual reality goggles. Note that while the previous three tasks were conducted in the evening, prior to sleep, this task was conducted in the morning. This was done to avoid tiring the participant out before they took part in the evening memory task.

Experimental design: Experiment 2

This experiment focused solely on navigation with all tasks being completed in a single afternoon session. As in Experiment 1, each navigation task followed the same pattern. Participants fixated on the centre screen, before an auditory stimulus was presented that they had to respond to the associated screen (indicated by an animal image attached to the screen). In addition to the five screens present in

Table 1. A list of the number of participants for a given task and the reasons for those who were excluded

Experiment	Substage	N. Included	Reasons for exclusion
1 (Scalp EEG)	Uncued Head Rotation	33	Motion tracking error (n = 6)
	Cued Head Rotation	33	Motion tracking error (n = 6)
	Cued Saccades	32	Motion tracking error (n = 6) EEG trigger error (n = 1)
	Cued Virtual Reality	29	Motion tracking error (n = 7) Dropped out during sleep stage (n = 3)
1 (Intracranial EEG)	Uncued Head Rotation	10	Motion tracking error (n = 1)
	Cued Head Rotation	10	Motion tracking error (n = 1)
	Cued Saccades	6	Motion tracking error (n = 1) Eyetracker sync. issue (n = 3)
	Cued Virtual Reality	9	Motion tracking error (n = 1) Dropped out during sleep stage (n = 1)
2	Sitting Head Rotation	20	Motion tracking error (n = 4)
	Standing Head Rotation	20	Motion tracking error (n = 4)
	Sitting Offset Rotation	20	Motion tracking error (n = 4)
	Standing Relocated Rotation	20	Motion tracking error (n = 4)

Experiment 1 (see figure 1), a further two screens were added at 90° and 120° relative to the centre screen of Experiment 1. These two screens were associated with two additional animal sounds. These additional screens were only used in the third task. Each experiment consisted of 160 trials, split across 4 blocks (i.e., 40 trials per block). In all experiments, vertical screen positions match participants' line of sight. Below, each variation of this experiment is outlined:

1) *Cued Head Rotation:* This was a direct replication of the *Cued Head Rotation* task used in Experiment 1.

2) *Standing Head Rotation:* This task matched the *Cued Head Rotation* task with the single exception that participants stood instead of sat. Participants were instructed to rotate their head while keeping their body facing forwards.

3) *Cued Head Rotation at 60° offset:* This task matched the *Cued Head Rotation* task with the single exception that participants sat facing the screen positioned at 60° from the centre screen and were then tasked with rotating their head relative to this position (that is, they rotated to face the screens at 0°, 30°, 90° and 120°).

4) *Standing Head Rotation after relocation:* This task matched the *Standing Head Rotation* task. Here however, participants stood ~1 metre closer to the centre screen.

Note that, in this experiment, the order of the tasks was randomised such that tasks 2 and 4 could occur in either order. Task 1, however, always came first to help participants acclimatise to the task and match the position of this task in Experiment 1. Task 3 was always the third task as it gave the participants a chance to recover from standing.

Scalp EEG acquisition and pre-processing

The EEG was recorded using an EEGo EEG system (ANT Neuro Enschede, Netherlands) with 65 Ag/AgCl electrodes arranged in a 10/10 system layout (including left and right mastoids; using CPz as the reference and AFz as the ground). All electrode impedances were less than 20 kΩ prior to commencing the experiment. The sampling rate was set at 1,000Hz.

In preparation for the central analyses, the continuous EEG data was first high-pass (0.5Hz; Butterworth infinite impulse response [IIR]), low-pass (165Hz; Butterworth IIR) and band-stop filtered (49-51Hz; Butterworth IIR). Second, independent components analysis was conducted. In Experiment 1, we focused on only removing components that resembled eye blinks or saccades to preserve as much of the raw signal as possible. In Experiment 2, we took a more stringent approach and, in addition to removing eye blinks and saccades, removed probable electromuscular components. The strategy used did not impact the analysis outcome, as evidenced by the replication of the main effect across experiments (N.B., the more stringent approach used in Experiment 2 however may explain why the electromuscular regressor explained less variance in the linear mixed-effects model than it did in the first experiment). Third, the data was epoched around the onset of the auditory cue; these epochs began 1,500ms before the onset of the cue, and ended 4,500ms after the onset of the cue. In the case of the *Uncued Head Rotation* condition (in which no auditory cue was presented), the data was epoched around when the auditory cue would have, theoretically, occurred. Fourth, the data was visually inspected for artefactual trials and channels, which were marked and then removed from the data. Fifth, the artefactual channels that had been removed from the data in the previous step were interpolated using neighbouring channels. Sixth, the data was re-referenced using the average of all channels.

iEEG acquisition and pre-processing

Recordings were performed at the Epilepsy Center, Department of Neurology, University of Munich, Germany. Intracranial EEG was recorded from Spencer depth electrodes (Ad-Tech Medical Instrument, Racine, Wisconsin, United States). Electrodes carried up to 15 contacts with , distances of 5 or 10mm between contacts. Data were recorded using XLTEK Neuroworks software (Natus Medical, San Carlos, California, US) and an XLTEK EMU128FS amplifier, with voltages referenced to a parietal electrode site. The sampling rate was set at 1,000Hz.

For signal preprocessing, the continuous iEEG data was first high-pass (0.5Hz; Butterworth IIR), low-pass (165Hz; Butterworth IIR) and band-stop filtered (49-51Hz; Butterworth IIR), identical to the scalp EEG preprocessing. Second, the data was epoched in the same manner as the scalp EEG data. Third,

the data was then visually inspected for artefactual trials, in particular for interictal epileptic activity, and
channels which exhibited such activity were marked and then removed from the data. Fourth, the data
was re-referenced using a bipolar referencing scheme.

We estimated the locations of the intracranial contacts using the Lead-DBS software (Horn & Kühn,
2015). First, we co-registered the post-operative CT scan to pre-operative T1-weighted image using a
two-stage linear registration (rigid followed by affine) as implemented in Advanced Normalisation Tools
(Avants et al., 2008). Second, we manually recorded the native space co-ordinates of all electrodes for
the given patient with a specific focus on identifying and labelling those electrodes which sat in
subregions of medial temporal lobe. Third, we spatially normalised these scans to MNI space based on
the pre-operative T1-weighted image using the Unified Segmentation Approach as implemented in
SPM12 (Ashburner & Friston, 2005). Fourth, we manually recorded the MNI space co-ordinates of all
electrodes for the given patient. Fifth, we re-labelled the contacts based on the lobe they sat in (as
determined by the contact MNI co-ordinate and the lobe atlas available in *WFU_PickAtlas*;
https://www.nitrc.org/projects/wfu_pickatlas/). Any contacts deemed to be within the medial temporal
lobe were given the label of the visually identified subregion. See table 2 for a summary of electrodes
per region-of-interest and per patient.

Eye-tracking acquisition and pre-processing

Eye-tracking data was acquired using a Tobii Pro Spectrum system. This system was calibrated
immediately prior to the beginning of the experiment. The sampling rate was set as 600Hz.

Offline, the co-ordinate space data was smoothed using a sliding window average of 50ms. The data
was then converted to visual angles, epoched to auditory cue onset (using the same parameters as the
EEG data) and visually inspected for artifacts. Any trial which contained physiologically-implausible
changes in eye position were excluded. Moreover, any trial which was missing more than 25% of sample
points were excluded. Any missing data points that remained were linearly interpolated based on data
from the preceding and proceeding sample points. Note that only data from the three screens (center
screen +/- 30°) were used as the eye-tracker could not reliably track eye positions on the outside
screens. Lastly, the data was upsampled to match the EEG recording sample rate.

Head motion-tracking acquisition and pre-processing

Head rotations were recorded using a Polhemus Liberty system. A sensor was attached the EEG cap
and sat between the electrodes AFz and FPz. The reference sensor was placed ~75cm behind and to
the right of the participant. Similar positioning was used for the patients in Experiment 2, though some
variability was encountered due to the position of the patient's bandages. Polhemus's proprietary
software was used to record the change in position of the first sensor relative to the second. The
sampling rate was set as 240Hz.

Table 2. A summary of the number of contacts of in each region-of-interest, and for each participant. The temporal
lobe region-of-interest includes electrodes found in the medial temporal lobe. Note that the nature of the linear
mixed-effects models used in our main analyses means that N is defined by the total number of electrodes in a
given region-of-interest rather than the total number of participants with electrodes in a given region.

Patient	Lobular Region-of-Interest				MTL Region of Interest		
	Frontal	Parietal	Occipital	Temporal	Parahipp.	Hipp.	Amygdala
1	51	15	0	3	0	3	0
2	0	8	16	32	4	4	0
3	44	12	0	12	0	0	4
4	28	8	0	28	0	4	4
5	0	16	0	56	4	4	4
6	30	3	0	15	0	0	3
7	40	12	16	28	4	4	0
8	6	3	0	48	9	6	0
9	16	0	32	20	4	0	0
10	24	15	16	21	0	3	0
Total	239	92	64	263	25	28	15

Offline, the co-ordinate space data was epoched around the onset of the auditory cue (as done for the
EEG and eyetracking data). We then computed the heading angle of every sample by first estimating
the rotation point of the head and then calculating the angular distance between a vector from the
rotation point to the centre screen and a vector from the rotation point to the current heading direction.
The resulting head direction signal was then visually inspected for artefacts. Any trial which contained
physiologically-implausible changes in head angle was excluded. Lastly, the data was upsampled to
match the EEG recording sample rate.

Forward encoding model: Overview

Here, we provide a brief overview of the forward encoding model (FEM) approach and, below, we report
the specifics of each step. For a visualisation of the approach, see figure 7. The FEM we used took
inspiration from that used by Nau and colleagues (2020). First, we decomposed the real-time head angle
measurement into basis sets of circular–Gaussian von-Mises distributions (from here on termed "HD
kernels"). Second, taking these HD kernels and the recorded EEG signal, we estimated an electrode's
directional tuning (that is, the relative weighting of each HD kernel for a given EEG electrode) using
ridge regression. Finally, we applied these weights to held-out datasets from the same participant to test
the generalisability of the model.

Forward encoding model: Generating the model

We built a FEM for every participant, task, kernel width, and EEG electrode individually. Head angle
was modelled using a basis set of circular–Gaussian von-Mises distributions (as implemented in the
Circular Statistics Toolbox for Matlab; https://github.com/circstat/circstatmatlab), with each kernel
covering 180° (i.e., ±90° from the centre screen) with an angular resolution of 1°. To test the tuning width
of each electrode, we iterated our analysis across six models, which each used a bespoke kernel width
(6°, 10°, 15°, 20°, 30°, and 60°). The distance between each kernel was tied to the kernel width to control
for directional sensitivity across models (i.e., models with a large kernel width used fewer kernels). We
then estimated the magnitude of kernel activities for every time point based on the real-time head angle.
By doing this for all kernels and time points, we obtained a 2D matrix (termed 'X') that described head
angle as a function of time and directional tuning.

Forward encoding model: Training the model

The motion and EEG signal were split into four partitions; each partition contained the same number of
trials for each head rotation. The training dataset was then formed from three of these partitions, and
the testing dataset was the remaining partition. This was done in a cross-validated manner such that,
through iteration, each partition acted as the testing dataset for the other partitions.

To train the FEM, we used the matrix derived in the section above (X) in conjunction with the EEG
training data at a given electrode (y) to predict the HD kernel weightings (β) through ridge (i.e., L2-
regularised) regression. Ridge regression helps penalise excessively high coefficients that may come
about through the fact that the HD kernels for a given timepoint are not independent. However, the
regularisation parameter (λ) for this approach first needs to be estimated. To do this, we used cross-
validation; we took two of the three training partitions, estimated the HD kernel weightings (β) for ten
different values of λ (log-spaced between 1 and 10,000,000), and then assessed how well these values
generalised to the remaining training partition. To assess generalisability, we calculated the Pearson
correlation coefficient between the time course of the EEG signal and the time course predicted by the
β-weighted HD kernels for each value of λ. The value of λ that led to the highest Pearson correlation
coefficient (averaged across the cross-validations) was then used to estimate the final model weights.
The final model weights were derived using ridge regression (this time across all three partitions).

Forward encoding model: Testing the model

599 To test the validity of the FEM, we applied the model weights to head angle data held in the independent
600 testing dataset (one out of four partitions). This returned a prediction of the EEG time-series, which was
601 then correlated with the observed EEG time-series using Pearson's correlation co-efficient.

602 To aid comparison between the models using different kernel widths, we used a bootstrapping
603 procedure where we circularly-shifted the predicted EEG time-series by a random integer and then

Figure 7. Visual depiction of analysis pipeline for Experiment 1. The analysis began with the preparation of motion tracking and EEG signals. The continuous, univariate motion tracking signal were convolved with a series of kernels to provide a 2-dimensional design matrix where each column reflected a single kernel and each row reflected a single point in time (see top left). EEG was broken down into principal components and then reconstructed using one of five proportions of components that were least likely to reflect electromuscular (EMG) activity (see top right). The forward encoding model (FEM) was then trained, tested and validated (see middle row). The motion and EEG data were split into four folds. The model was trained on three of these folds (see middle left). The resulting model weights were then applied to the held-out fold and used to estimate EEG activity for this fold (see centre of middle row). The model was then validated by correlating the predicted EEG activity if the real EEG activity of the held-out fold (see middle right). The resulting correlation co-efficient was normalised against a chance distribution of correlation values derived from randomly time-shifting the predicted EEG time series relative to the real EEG activity. Lastly, linear mixed-effects (LME) models were used to investigate how distinct latent factors influenced FEM performance (see bottom row). The LME models were fit to the observed (or "true") data and to 200 instances of data were the sign of the FEM performance was flipped at random (see bottom left). The size of the resulting effect was then computed by comparing the LME weights derived from the "true" data to the chance data (see bottom right).

recomputed the correlation coefficient. By repeating this procedure 100 times, we generated a distribution of correlation coefficients between the observed and predicted time-series that we would expect by chance. We can then normalise the "true" correlation coefficient by subtracting the mean of the chance distribution and then dividing by the standard deviation of this distribution.

Forward encoding model: Modelling muscular activity

Muscular activity was modelled as a covariate in the linear mixed-effects (LME) model used for inferential analysis, and hence regressed out of the variables of interest. The full model is described below (see "*EEG encoding model: Group-level analysis for Experiment 1*"). Here, we just describe the component of the LME model used to regress out muscular activity.

To this end, we used principal components analysis (PCA) to break the EEG data into its principal components, and then ordered these components based on the relative likelihood that a given component contained muscular activity, informed by the components' topography. This likelihood

measure was computed by taking the mean of the size of the weights for electrodes on the outer rim of the cap and dividing this value by the mean of the size of the weights for all the remaining electrodes. The larger the resulting value, the more 'muscular' a component was judged to be, following the rationale that electrodes that sit at the edge of the cap are closest to the neck muscles involved in the head rotation. Based on this ordering procedure, we created five groups of components: the 20% least muscular components, the 40% least muscular components, and so on. For each of these groups, the EEG data was reconstructed in electrode-space using the selected components. The FEM was generated, trained, and tested for each of these datasets separately.

The performance of these FEMs can be thought of as being a linear sum of EEG signals that reflect muscular activity and EEG signals that do not reflect muscular activity (there can be no third option – the signal is either muscular or not). Based on the PCA approach used above, we can also conclude that the EEG signals that reflect muscular activity get progressively smaller as we remove more of the muscular components, while the EEG signals that do not reflect muscular activity remain constant. This can be expressed as:

$$y = a + bX + error$$

Where *y* represents FEM performance, *a* represents the constant (i.e., EEG signal that doesn't change as a function of the number of muscular components removed), *b* represents an unknown weighting, and *X* reflects the percentage of muscular components removed. This equation will form part of the central LME model and allow for muscular contributions to FEM performance to be regressed out of the central analyses.

Forward encoding model: Modelling saccade-related activity

In addition to creating a FEM which used head angle as a predictor of EEG activity, we also created a FEM which used eye position as a predictor of EEG activity. The eye position FEM was computed in the same manner as the head angle FEM with two exceptions: (1) visual angle was used as the predictor in place of head angle, and (2) only data from the screens as +/- 30° were used as the eye-tracker could not track eye positions on the outside screens. The eye position FEM was built on the data from the *Cued Saccades* task. As no head rotation occurred in this task, whenever we refer to FEM performance for this task, we refer to the eye position model.

Forward encoding model: Group-level analysis for Experiment 1

Through completion of the analysis steps described above, we obtained a z-score describing FEM performance relative to chance for every participant, task, electrode, kernel width and principal component subcategory. To assess whether these z-scores differed from chance across participants (i.e., z > 0), we used linear mixed-effect (LME) models. We computed an LME for every kernel-electrode pair. Each LME took data from all participants, all tasks and all principal components subcategories. The outcome variable of each LME was FEM performance. For predictors, random intercepts were created for each participant, as well as four fixed effects that represented latent factors. The fixed effects were as follows:

1. Head angle: A binary variable which denoted whether head rotation had occurred in the given task.
2. Auditory cuing: A binary variable which denoted whether an auditory cue had been played in the given task.
3. Visual input / saccadic activity: A binary variable which denoted whether visual input changed / a saccade occurred (the two are perfectly correlated) in the given task.
1. Muscular activity: A variable ranging from 0.2 to 1, which denoted the percentage of principal components included in the sample prior to the fitting of the encoding model.

See table 3 for a mapping of experiment conditions onto latent factors.

The LME returned beta weights for each regressor. To assess statistical significance, the resulting weights were compared to a surrogate distribution (200 permutations) of weights generated by randomly permuting the sign of the outcome variable for each sample and then refitting the model. The p-value was computed by calculating the proportion of surrogate weights which were larger in value than the 'true' model weights. This process was repeated for every kernel-electrode pair. False discovery rate

(FDR) correction was used to address the multiple comparison issue incurred by conducting analysis
on each resulting beta weight, kernel and electrode separately.

The LME used for the patients was near-identical to that used for the healthy participants. The key
difference was that, rather than using separate models for each electrode, separate models were used
for defined regions-of-interest. This approach was necessitated by the fact that implantation montages
are bespoke for each patient, leading to a variable number of contacts in each region across patients.
Regions-of-interest were defined as the frontal lobe, occipital lobe, parietal lobe, and temporal lobe (as
defined by the lobe options in WFU_PickAtlas; https://www.nitrc.org/projects/wfu_pickatlas/), and the
parahippocampus, hippocampus and amygdala (as defined by visual inspection of the patient scans.
As before, participants were included as random effects in the LME, avoiding issues in generalisability
that occur when using fixed-effect models.

Forward encoding model: Group-level analysis for Experiment 2

The concept behind the group-level analysis for Experiment 2 matched that of Experiment 1, but given
the different tasks, the latent factors inevitably changed. Here, the fixed effects were as follows:

2. Head angle: A binary variable which denoted whether head rotation had occurred in the given
task.
3. Position on the z-axis: A binary variable which denoted whether participants were sitting or
standing during the given task.
4. Relocation on the x-axis: A binary variable which denoted whether participants were
sitting/standing in the standard position during the given task, or whether they stepped one
metre closer to the screens.
5. Muscular activity: A variable ranging from 0.2 to 1, which denoted the percentage of principal
components included in the data prior to the fitting of the encoding model.

See table 4 for a mapping of experiment conditions onto latent factors.

Lag-based encoding model

To assess the temporal dynamics of the EEG signal tuned to head angle relative to the timing of physical
head rotation, we built a lag-based FEM. This FEM operates in the same manner as described above,
except that (1) the time series of the EEG is shifted by a given number of samples prior to fitting, and
(2) to reduce computational load, only kernels of 20° were used. The temporal shift of the EEG is referred
to as the lag. We fitted the lag-based model for 40 lags, ranging from when the EEG time series
preceded the motion tracking time series by 200ms to when the EEG time series followed the motion
tracking time series by 200ms. All other aspects of the analysis remained the same. False discovery
rate (FDR) correction was used to address the multiple comparison issue incurred by conducting
analysis on each electrode and lag separately.

Inverted encoding model

702 As a second means of assessing the temporal dynamics of the head angle-related EEG signal, we build
703 an inverted encoding model. This approach matched the main approach up until the section "*EEG
704 encoding model: Testing the model*". At this stage, the weights derived from 20° FEM were inverted (i.e.,
705 $inv(\beta * \beta') * \beta$). The weights were then applied to the EEG scalp data iteratively for each sample point.
706 This returned predicted head direction kernel values for every sample of every trial. Representational

Table 4. Mapping of experiment tasks onto latent variables for experiment 2.

Experimental task	Latent factor		
	Head angle	Position z-axis	Relocation x-axis
Cued Head Rotation	✓		
Standing Head Rotation	✓	✓	
Cued Head Rotation at 60° offset	✓		
Standing Head Rotation after relocation	✓	✓	✓

Table 5. Mapping of training/testing combinations onto latent variables for the dissociation of head rotation and head direction. In short, activity related to head rotation should generalise across sitting positions (i.e., when they faced different angles in the environment); head direction activity should not.

Training condition	Testing Condition	Latent factor	
		Head rotation	Head direction
Cued Head Rotation	Cued Head Rotation	✓	✓
Cued Head Rotation	Cued Head Rotation at 60° offset	✓	
Cued Head Rotation at 60° offset	Cued Head Rotation	✓	✓
Cued Head Rotation at 60° offset	Cued Head Rotation at 60° offset	✓	

similarity analysis (RSA) was then used to identify whether the kernel magnitudes at a given sample
could distinguish between the four monitor positions. To this end, a representational dissimilarity matrix
(RDM) was computed by computing Pearson correlations between the predicted kernel magnitudes of
every pair of trials. The RDM was then correlated with a model RDM which stated that the correlation
between pairs with the same target head direction would equal one and all other pairs would equal zero.
A surrogate distribution of chance correlation values was then computed by shuffling the labels of the
trials, rebuilding the model RDM and then correlating the shuffled model RDM with the observed RDM.
The correlation of the 'true' data was then normalised using the mean and standard deviation of the
surrogate data. By repeating these steps for every time point, we obtain a time series that describes the
extent to which the inverted model predicts head direction above and beyond what would be expected
by chance. False discovery rate (FDR) correction was used to address the multiple comparison issue
incurred by conducting analysis on each time point separately. The remainder of the analysis matches
that of the main approach.

Generalised encoding models

To disentangle the influences of head rotation and head direction signal, and location-specific and
location-independent head direction signal, we used generalised encoding models. We will use the head
rotation/head direction dissociation as the example here, but the same procedure was applied to the
location-specific/location-independent dissociation.

For the generalised encoding models, we trained the encoding model on one condition (e.g., Cued Head
Rotation) and applied the model weights to another condition (i.e., Cued Head Rotation at 60° offset).
In the case of head direction versus head rotation, if the weights lead to the test data outperforming

Table 6. Mapping of training/testing combinations onto latent variables for the dissociation of location-independent and location-specific head direction signals. In short, activity related to location-independent head direction signals should generalise across conditions, whereas location-specific signals should not.

Training condition	Testing Condition	Latent factor	
		Location independent	Location specific
Cued Head Rotation	Cued Head Rotation	✓	✓
Cued Head Rotation	Standing Head Rotation	✓	
Cued Head Rotation	Standing Rot. after relocation	✓	
Standing Head Rotation	Cued Head Rotation	✓	
Standing Head Rotation	Standing Head Rotation	✓	✓
Standing Head Rotation	Standing Rot. after relocation	✓	
Standing Rot. after relocation	Cued Head Rotation	✓	
Standing Rot. after relocation	Standing Head Rotation	✓	
Standing Rot. after relocation	Standing Rot. after relocation	✓	✓

what would be expected by chance, it can be said that the model tracks activity related to head rotation
as the FEM generalises to when participants are sitting at an offset. If the weights, however, lead to
poorer performance than what is observed when conducting cross-validated training and testing within
a condition, it can be said that the model tracks activity head direction. Notably, the two outcomes are
not mutually exclusive: generalised model performance can be significantly greater than chance, while
also being significantly poorer than within-condition performance. To test both ideas simultaneously,
linear mixed models were used with fixed effects denoting head rotation, head direction and muscular
activity (see table 5 for mapping of generalised encoding models onto factors). Random intercepts were
included for each participant. Statistical analysis matched what was reported above.

The same principles were applied to the dissociation of location-independent and location-specific head
direction signals. Table 6 describes how the generalised encoding models were mapped onto the fixed
effects of the linear mixed effect models.

References

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage*, *26*(3), 839–851.
    https://doi.org/10.1016/j.neuroimage.2005.02.018

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image
    registration with cross-correlation: Evaluating automated labeling of elderly and
    neurodegenerative brain. *Medical Image Analysis*, *12*(1), 26–41.
    https://doi.org/10.1016/j.media.2007.06.004

Bassett, J. P., & Taube, J. S. (2001). Neural Correlates for Angular Head Velocity in the Rat Dorsal
    Tegmental Nucleus. *The Journal of Neuroscience*, *21*(15), 5740–5751.
    https://doi.org/10.1523/JNEUROSCI.21-15-05740.2001

Baumann, O., & Mattingley, J. B. (2010). Medial Parietal Cortex Encodes Perceived Heading Direction
    in Humans. *Journal of Neuroscience*, *30*(39), 12897–12901.
    https://doi.org/10.1523/JNEUROSCI.3077-10.2010

Baumann, O., & Mattingley, J. B. (2021). Extrahippocampal contributions to spatial navigation in
    humans: A review of the neuroimaging evidence. *Hippocampus*, *January*, 1–18.
    https://doi.org/10.1002/hipo.23313

Bellmund, J. L. S., Gärdenfors, P., Moser, E. I., & Doeller, C. F. (2018). Navigating cognition: Spatial
    codes for human thinking. *Science*, *362*(6415). https://doi.org/10.1126/science.aat6766

Blair, H. T., Cho, J., & Sharp, P. E. (1998). Role of the Lateral Mammillary Nucleus in the Rat Head
    Direction Circuit: A Combined Single Unit Recording and Lesion Study. *Neuron*, *21*, 1387–
    1397.

Blair, H. T., & Sharp, P. (1995). Anticipatory head direction signals in anterior thalamus: Evidence for a
    thalamocortical circuit that integrates angular head motion to compute head direction. *The
    Journal of Neuroscience*, *15*(9), 6260–6270. https://doi.org/10.1523/JNEUROSCI.15-09-
    06260.1995

Butler, W. N., Smith, K. S., van der Meer, M. A. A., & Taube, J. S. (2017). The head-direction signal
    plays a functional role as a neural compass during navigation. *Current Biology*, 10.

Butler, W. N., & Taube, J. S. (2017). Oscillatory synchrony between head direction cells recorded
    bilaterally in the anterodorsal thalamic nuclei. *Journal of Neurophysiology*, *117*(5), 1847–1852.
    https://doi.org/10.1152/jn.00881.2016

Buzsáki, G., & Moser, E. I. (2013). Memory, navigation and theta rhythm in the hippocampal-
    entorhinal system. *Nature Neuroscience*, *16*(2), 130–138. https://doi.org/10.1038/nn.3304

Calton, J. L., Stackman, R. W., Goodridge, J. P., Archey, W. B., Dudchenko, P. A., & Taube, J. S.
    (2003). Hippocampal Place Cell Instability after Lesions of the Head Direction Cell Network.
    The Journal of Neuroscience, *23*(30), 9719–9731. https://doi.org/10.1523/JNEUROSCI.23-30-
    09719.2003

Calton, J. L., Turner, C. S., Cyrenne, D.-L. M., Lee, B. R., & Taube, J. S. (2008). Landmark control
    and updating of self-movement cues are largely maintained in head direction cells after
    lesions of the posterior parietal cortex. *Behavioral Neuroscience*, *122*(4), 827–840.
    https://doi.org/10.1037/0735-7044.122.4.827

Cullen, K. E., & Taube, J. S. (2017). Our sense of direction: Progress, controversies and challenges.
    Nature Neuroscience, *20*(11), 1465–1473. https://doi.org/10.1038/nn.4658

Do, T.-T. N., Lin, C.-T., & Gramann, K. (2021). Human brain dynamics in active spatial navigation.
    Scientific Reports, *11*(1), 13036. https://doi.org/10.1038/s41598-021-92246-4

Dudchenko, P. A., Wood, E. R., & Smith, A. (2019). A new perspective on the head direction cell
    system and spatial behavior. *Neuroscience & Biobehavioral Reviews*, *105*, 24–33.
    https://doi.org/10.1016/j.neubiorev.2019.06.036

Epstein, R. A., Patai, E. Z., Julian, J. B., & Spiers, H. J. (2017). The cognitive map in humans: Spatial
    navigation and beyond. *Nature Neuroscience*, *20*(11), 1504–1513.
    https://doi.org/10.1038/nn.4656

Goodridge, J. P., Dudchenko, P. A., Worboys, K. A., Golob, E. J., & Taube, J. S. (1998). Cue Control and Head Direction Cells. *Behavioral Neuroscience*, *112*(4), 749–761.

Gramann, K., Hohlefeld, F. U., Gehrke, L., & Klug, M. (2021). Human cortical dynamics during full-body heading changes. *Scientific Reports*, *11*(1), 18186. https://doi.org/10.1038/s41598-021-97749-8

Griffiths, B. J., Mazaheri, A., Debener, S., & Hanslmayr, S. (2016). Brain oscillations track the formation of episodic memories in the real world. *NeuroImage*, *143*, 256–266. https://doi.org/10.1101/042929

Horn, A., & Kühn, A. A. (2015). Lead-DBS: A toolbox for deep brain stimulation electrode localizations and visualizations. *NeuroImage*, *107*, 127–135. https://doi.org/10.1016/j.neuroimage.2014.12.002

Hulse, B. K., & Jayaraman, V. (2020). Mechanisms Underlying the Neural Computation of Head Direction. *Annual Review of Neuroscience*, *43*(1), 31–54. https://doi.org/10.1146/annurev-neuro-072116-031516

Jacob, P.-Y., Casali, G., Spieser, L., Page, H., Overington, D., & Jeffery, K. (2017). An independent, landmark-dominated head-direction signal in dysgranular retrosplenial cortex. *Nature Neuroscience*, *20*(2), 173–175. https://doi.org/10.1038/nn.4465

Jacobs, J., Kahana, M. J., Ekstrom, A. D., Mollison, M. V., & Fried, I. (2010). A sense of direction in human entorhinal cortex. *Proceedings of the National Academy of Sciences*, *107*(14), 6487–6492. https://doi.org/10.1073/pnas.0911213107

Julian, J. B., Keinath, A. T., Frazzetta, G., & Epstein, R. A. (2018). Human entorhinal cortex represents visual space using a boundary-anchored grid. *Nature Neuroscience*, *21*(2), 191–194. https://doi.org/10.1038/s41593-017-0049-1

Killian, N. J., Jutras, M. J., & Buffalo, E. A. (2012). A map of visual space in the primate entorhinal cortex. *Nature*, *491*(7426), 761–764. https://doi.org/10.1038/nature11587

Killian, N. J., Potter, S. M., & Buffalo, E. A. (2015). Saccade direction encoding in the primate entorhinal cortex during visual exploration. *Proceedings of the National Academy of Sciences*, *112*(51), 15743–15748. https://doi.org/10.1073/pnas.1417059112

Kim, M., & Maguire, E. A. (2019). Encoding of 3D head direction information in the human brain. *Hippocampus*, *29*(7), 619–629. https://doi.org/10.1002/hipo.23060

Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*, *2*(November), 1–28. https://doi.org/10.3389/neuro.06.004.2008

Kunz, L., Brandt, A., Reinacher, P. C., Staresina, B. P., Reifenstein, E. T., Weidemann, C. T., Herweg, N. A., Patel, A., Tsitsiklis, M., Kempter, R., Kahana, M. J., Schulze-Bonhage, A., & Jacobs, J. (2021). A neural code for egocentric spatial maps in the human medial temporal lobe. *Neuron*, *109*(17), 2781-2796.e10. https://doi.org/10.1016/j.neuron.2021.06.019

Kunz, L., Maidenbaum, S., Chen, D., Wang, L., Jacobs, J., & Axmacher, N. (2019). Mesoscopic Neural Representations in Spatial Navigation. *Trends in Cognitive Sciences*, *23*(7), 615–630. https://doi.org/10.1016/j.tics.2019.04.011

Laurens, J., Kim, B., Dickman, J. D., & Angelaki, D. E. (2016). Gravity orientation tuning in macaque anterior thalamus. *Nature Neuroscience*, *19*(12), 1566–1568. https://doi.org/10.1038/nn.4423

Lomi, E., Jeffery, K. J., & Mitchell, A. S. (2023). Convergence of direction, location and theta in the rat anteroventral thalamic nucleus. *BioRxiv*. https://doi.org/10.1101/2023.01.11.523585

Medendorp, W. P., Beurze, S. M., Van Pelt, S., & Van Der Werf, J. (2008). Behavioral and cortical mechanisms for spatial coding and action planning. *Cortex*, *44*(5), 587–597. https://doi.org/10.1016/j.cortex.2007.06.001

Meister, M. L. R., & Buffalo, E. A. (2018). Neurons in Primate Entorhinal Cortex Represent Gaze Position in Multiple Spatial Reference Frames. *The Journal of Neuroscience*, *38*(10), 2430–2441. https://doi.org/10.1523/JNEUROSCI.2432-17.2018

Muller, R. U., Ranck, J. B., & Taube, J. S. (1996). Head direction cells: Properties and functional
significance. *Current Opinion in Neurobiology*, *6*(2), 196–206. https://doi.org/10.1016/S0959-
4388(96)80073-0

Nau, M., Navarro Schröder, T., Bellmund, J. L. S., & Doeller, C. F. (2018). Hexadirectional coding of
visual space in human entorhinal cortex. *Nature Neuroscience*, *21*(2), 188–190.
https://doi.org/10.1038/s41593-017-0050-8

Nau, M., Navarro Schröder, T., Frey, M., & Doeller, C. F. (2020). Behavior-dependent directional
tuning in the human visual-navigation network. *Nature Communications*, *11*(1), 1–13.
https://doi.org/10.1038/s41467-020-17000-2

Patai, E. Z., & Spiers, H. J. (2021). The Versatile Wayfinder: Prefrontal Contributions to Spatial
Navigation. *Trends in Cognitive Sciences*, *25*(6), 520–533.
https://doi.org/10.1016/j.tics.2021.02.010

Peyrache, A., Lacroix, M. M., Petersen, P. C., & Buzsáki, G. (2015). Internally organized mechanisms
of the head direction sense. *Nature Neuroscience*, *18*(4), 569–575.
https://doi.org/10.1038/nn.3968

Piñeyro Salvidegoitia, M., Jacobsen, N., Bauer, A.-K. R., Griffiths, B. J., Hanslmayr, S., & Debener, S.
(2019). Out and about: Subsequent memory effect captured in a natural outdoor environment
with smartphone EEG. *Psychophysiology*, *October 2018*, e13331–e13331.
https://doi.org/10.1111/psyp.13331

Ranck, J. B. (1984). *Head direction cells in the deep layer of dorsal presubiculum in freely moving*
rats. 10, 599.

Robertson, R. G., Rolls, E. T., Georges-Fran�ois, P., & Panzeri, S. (1999). Head direction cells in the
primate pre-subiculum. *Hippocampus*, *9*(3), 206–219. https://doi.org/10.1002/(SICI)1098-
1063(1999)9:3<206::AID-HIPO2>3.0.CO;2-H

Rolls, E. T. (1999). Spatial view cells and the representation of place in the primate hippocampus.
Hippocampus, *9*(4), 467–480. https://doi.org/10.1002/(SICI)1098-1063(1999)9:4<467::AID-
HIPO13>3.0.CO;2-F

Rolls, E. T., & O'Mara, S. M. (1995). View-responsive neurons in the primate hippocampal complex.
Hippocampus, *5*(5), 409–424. https://doi.org/10.1002/hipo.450050504

Sharp, P. E., Blair, H. T., & Cho, J. (2001). The anatomical and computational basis of the rat head-
direction cell signal. *Trends in Neurosciences*, *24*(5), 289–294.

Shinder, M. E., & Taube, J. S. (2011). Active and passive movement are encoded equally by head
direction cells in the anterodorsal thalamus. *Journal of Neurophysiology*, *106*(2), 788–800.
https://doi.org/10.1152/jn.01098.2010

Shinder, M. E., & Taube, J. S. (2014). Self-motion improves head direction cell tuning. *Journal of*
Neurophysiology, *111*(12), 2479–2492. https://doi.org/10.1152/jn.00512.2013

Shine, J. P., Valdés-Herrera, J. P., Hegarty, M., & Wolbers, T. (2016). The Human Retrosplenial
Cortex and Thalamus Code Head Direction in a Global Reference Frame. *The Journal of*
Neuroscience, *36*(24), 6371–6381. https://doi.org/10.1523/JNEUROSCI.1268-15.2016

Stackman, R. W., & Taube, J. S. (1997). Firing Properties of Head Direction Cells in the Rat Anterior
Thalamic Nucleus: Dependence on Vestibular Input. *Journal of Neuroscience*, *17*(11), 4349–
4358.

Stackman, R. W., & Taube, J. S. (1998). Firing Properties of Rat Lateral Mammillary Single Units:
Head Direction, Head Pitch, and Angular Head Velocity. *Journal of Neuroscience*, *18*(21).

Staudigl, T., Leszczynski, M., Jacobs, J., Sheth, S. A., Schroeder, C. E., Jensen, O., & Doeller, C. F.
(2018). Hexadirectional Modulation of High-Frequency Electrophysiological Activity in the
Human Anterior Medial Temporal Lobe Maps Visual Space. *Current Biology*, *28*(20), 3325-
3329.e4. https://doi.org/10.1016/j.cub.2018.09.035

Steel, A., Robertson, C. E., & Taube, J. S. (2021). Current Promises and Limitations of Combined
Virtual Reality and Functional Magnetic Resonance Imaging Research in Humans: A
Commentary on Huffman and Ekstrom (). *Journal of Cognitive Neuroscience*, *33*(2), 159–166.
https://doi.org/10.1162/jocn_a_01635

Taube, J. S. (1995). Head Direction Cells Recorded in the Anterior Thalamic Nuclei of Freely Moving
Rats. *Journal of Neuroscience*, 17.

Taube, J. S. (2007). The Head Direction Signal: Origins and Sensory-Motor Integration. *Annual*
Review of Neuroscience, *30*(1), 181–207.
https://doi.org/10.1146/annurev.neuro.29.051605.112854

Taube, J. S., & Bassett, J. P. (2003). Persistent Neural Activity in Head Direction Cells. *Cerebral*
Cortex, *13*(11), 1162–1172. https://doi.org/10.1093/cercor/bhg102

Taube, J. S., Muller, R., & Ranck, J. (1990). Head-direction cells recorded from the postsubiculum in
freely moving rats. I. Description and quantitative analysis. *The Journal of Neuroscience*,
10(2), 420–435. https://doi.org/10.1523/JNEUROSCI.10-02-00420.1990

Taube, J. S., & Muller, R. U. (1998). Comparisons of head direction cell activity in the postsubiculum
and anterior thalamus of freely moving rats. *Hippocampus*, *8*(2), 87–108.
https://doi.org/10.1002/(SICI)1098-1063(1998)8:2<87::AID-HIPO1>3.0.CO;2-4

Taube, J. S., Valerio, S., & Yoder, R. M. (2013). Is Navigation in Virtual Reality with fMRI Really
Navigation? *Journal of Cognitive Neuroscience*, *25*(7), 1008–1019.
https://doi.org/10.1162/jocn_a_00386

Tsanov, M., Chah, E., Vann, S. D., Reilly, R. B., Erichsen, J. T., Aggleton, J. P., & O'Mara, S. M.
(2011). Theta-Modulated Head Direction Cells in the Rat Anterior Thalamus. *Journal of*
Neuroscience, *31*(26), 9489–9502. https://doi.org/10.1523/JNEUROSCI.0353-11.2011

Vass, L. K., & Epstein, R. A. (2013). Abstract Representations of Location and Facing Direction in the
Human Brain. *Journal of Neuroscience*, *33*(14), 6133–6142.
https://doi.org/10.1523/JNEUROSCI.3873-12.2013

Winter, S. S., Clark, B. J., & Taube, J. S. (2015). Disruption of the head direction cell network impairs
the parahippocampal grid cell signal. *Science*, *347*(6224), 870–874.
https://doi.org/10.1126/science.1259591

Wirth, S., Baraduc, P., Planté, A., Pinède, S., & Duhamel, J.-R. (2017). Gaze-informed, task-situated
representation of space in primate hippocampus during virtual navigation. *PLOS Biology*,
15(2), e2001045. https://doi.org/10.1371/journal.pbio.2001045

Yoder, R. M., Clark, B. J., Brown, J. E., Lamia, M. V., Valerio, S., Shinder, M. E., & Taube, J. S.
(2011). Both visual and idiothetic cues contribute to head direction cell stability during
navigation along complex routes. *Journal of Neurophysiology*, *105*(6), 2989–3001.
https://doi.org/10.1152/jn.01041.2010

Zirkelbach, J., Stemmler, M., & Herz, A. V. M. (2019). Anticipatory Neural Activity Improves the
Decoding Accuracy for Dynamic Head-Direction Signals. *The Journal of Neuroscience*,
39(15), 2847–2859. https://doi.org/10.1523/JNEUROSCI.2605-18.2019

Supplementary Figure 1. Saccades precede physical rotation during orienting tasks. Following the auditory cue (time = 0), participants made saccades/head rotations to the target screen. The coloured lines depicted mean angular position as a function of time, averaged over trials and participants. Blue lines reflect head position, turquoise lines reflect eye position. Filled lines reflect rotations to the first right-hand screen, dotted lines reflect rotations to the first left-hand screen. As the eye-tracker could not track eye position for the far left/far right screen, neither the head direction nor eye position data for these trials are plotted here.

Supplementary Table 1. Fixed effects for the basic linear mixed-effects model used in Experiment 1.

Regressor	Tuning Width	Peak Electrode	z	p_{FDR}
Head Direction	10	CP6	7.622	< 0.001
	15	FC2	7.621	< 0.001
	20	FC2	8.096	< 0.001
	30	C6	7.493	< 0.001
	45	Cz	7.332	< 0.001
	60	C6	7.140	< 0.001
Auditory Cue	10	C1	0.934	0.180
	15	T8	0.819	0.215
	20	C1	0.776	0.215
	30	Pz	1.092	0.130
	45	Cz	1.058	0.150
	60	T8	1.146	0.125
Visual Input / Saccadic Activity	10	Pz	2.472	0.005
	15	Pz	2.258	0.005
	20	Pz	2.501	0.010
	30	Pz	2.528	0.010
	45	Pz	2.190	0.010
	60	Pz	2.601	0.010
Electromuscular activity	10	AF8	2.847	< 0.001
	15	F8	2.761	< 0.001
	20	AF8	2.669	< 0.001
	30	AF8	3.017	< 0.001
	45	AF8	3.803	< 0.001
	60	AF8	2.745	< 0.001

Supplementary Table 2. Fixed effects for the basic linear mixed-effects model used in Experiment 2.

Regressor	Tuning Width	Peak Electrode	z	p_{FDR}
Head Direction	10	CP5	6.539	< 0.001
	15	CP5	6.653	< 0.001
	20	Cz	7.031	< 0.001
	30	CP1	7.404	< 0.001
	45	Cz	6.712	< 0.001
	60	Pz	6.524	< 0.001
Relocation x-axis	10	PO4	2.528	< 0.001
	15	P4	2.870	< 0.001
	20	P4	2.985	0.010
	30	P4	2.992	< 0.001
	45	P4	2.968	< 0.001
	60	P2	2.967	< 0.001
Relocation z-axis	10	FCz	0.351	0.375
	15	F2	1.077	0.145
	20	F2	1.069	0.135
	30	Cz	-0.617	0.715
	45	FCz	-0.525	0.685
	60	Cz	-0.513	0.715
Electromuscular activity	10	FC5	3.716	< 0.001
	15	FC5	3.659	< 0.001
	20	FC5	3.748	< 0.001
	30	F5	3.960	< 0.001
	45	FC3	3.891	< 0.001
	60	FC5	3.360	< 0.001