Can snare removal safeguard protected areas caught in the Southeast Asian snaring crisis? Promises and perspectives

Andrew Tilker*

Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

Re:wild, Austin, TX 78767 USA

tilker@izw-berlin.de

https://orcid.org/0000-0003-3630-8691

Jürgen Niedballa*

Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

camtrapr@gmail.com

https://orcid.org/0000-0002-9187-2116

Hung Luong Viet

WWF-Viet Nam, 150 Truong Gia Mo, Hue, Viet Nam

hung.luongviet@wwf.org.vn

Jesse F. Abrams

Global Systems Institute, University of Exeter, Exeter, EX4 4QE, UK

j.abrams@exeter.ac.uk

https://orcid.org/0000-0003-0411-8519

Lucile Marescot

CBGP, Univ Montpellier, CIRAD, INRAE, Institut Agro, IRD, Montpellier, France

lucile.marescot@cirad.fr

ORCID ID: 0000-0001-7625-5446

Nicholas Wilkinson

Independent consultant, Linton, Cambridgeshire, UK.

nicholasmwilkinson@gmail.com

Benjamin M. Rawson

World Wide Fund for Nature - Asia Pacific, Canberra, Australia

brawson@wwfint.org

Rahel Sollmann

Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

sollmann@izw-berlin.de

https://orcid.org/0000-0002-1607-2039

Andreas Wilting

Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

wilting@izw-berlin.de

https://orcid.org/0000-0001-5073-9186

* contributed equally

Abstract

Unsustainable snaring is causing biodiversity declines across tropical protected areas, resulting in

species extinctions and jeopardizing the health of forest ecosystems. Here, we used 11 years of ranger

collected data to assess the impact of intensive snare removal in two protected areas in Viet Nam.

Snare removal resulted in significant declines in snare occupancy (36.9, 95% BCI [4.6, 59.0] reduction

in percent area occupied), but snaring levels nonetheless remained high (31.4, [23.6, 40.8] percent

area occupied), and came with a substantial financial cost. Our results indicate that snare removal

remains an important component of efforts to protect tropical protected areas, but by itself, is likely

insufficient to address this threat. To achieve the target of zero snaring in protected areas, a

multifaceted approach will be necessary that combines short-term reactive snare removal with long-

term proactive programs that address the underlying drivers behind snaring.

Introduction

Unsustainable hunting has defaunated tropical forests around the world (Harrison et al, 2016; Benítez-López et al, 2017, 2019), and preventing further defaunation is a priority for addressing global biodiversity loss (Laurance et al 2012; Barlow et al 2018). Furthermore, because large vertebrates perform ecological functions that are integral to maintaining healthy tropical ecosystems (Lacher et al, 2019), the loss of forest fauna jeopardizes a number of ecological services linked to human well-being (Krause & Tilker et al, 2022). For example, defaunation has been linked to reduced carbon storage capacities in tropical forests (Bello et al, 2015; Osuri et al, 2016), increase in prevalence of zoonotic diseases (Young et al, 2014), and loss of ethnocultural identity for indigenous peoples (Bogoni et al, 2020).

Protected areas remain the most prominent conservation approach to safeguard pan-tropical wildlife, but the effectiveness of site-based conservation strategies varies (Watson et al, 2014; Coad et al, 2019) and has been insufficient to prevent the widespread decline of mammal and bird species in some regions (Benítez-López et al, 2017, 2019). Tropical defaunation has been particularly severe in Southeast Asia, where illegal unsustainable hunting realized through the setting of wire snares has depressed wildlife populations in many protected areas, and remains a significant and ongoing threat (Belecky & Gray, 2020; Gray et al, 2021). A recent study, for example, found that snaring is a more immediate and severe threat to Southeast Asian faunal communities than forest degradation in some areas (Tilker & Abrams et al, 2019). To prevent further defaunation in Southeast Asia, and thus maintain the ecosystem services and functions of its tropical forests, it is important to assess the effectiveness of conservation actions designed to counteract the ongoing Southeast Asian 'snaring crisis'.

Although several conservation actions have been proposed to counter snaring, snare removal remains the primary strategy employed in Southeast Asian protected areas (Belecky & Gray, 2020).

Snare removal is popular because it is straightforward and non-controversial (Belecky & Gray, 2020),

and it will likely continue to prevail as the main approach to address snaring across most protected

areas. Some studies have assessed the impact of snare removal on snaring levels, with both positive

(Linkie et al, 2015) and equivocal (Becker et al, 2013) findings. Most of these studies, however, have

been over relatively short time periods or in areas where snaring levels do not reach the high levels

reported for many parts of Southeast Asia (Jachmann, 2008; Becker et al, 2013; Watson et al, 2013;

Linkie et al, 2015). The extent to which snare removal reduces snaring pressure over longer time

horizons remains unexplored, hindering an objective evaluation of its effectiveness. For protected

areas operating under limited resources, this lack of knowledge is worrisome because employing

dedicated snare removal teams is expensive, potentially limiting investment in other strategies to

reduce snaring pressure.

Here, we used 11 years of patrol data from two contiguous protected areas in central Viet Nam to

assess the impact of snare removal on snaring levels over time. Like many protected areas in

Southeast Asia, the reserves are under high levels of snaring pressure, and snare removal has been

the main strategy to counteract this threat. Within this context, we seek to understand how snare

removal fits within a larger framework to counter snaring, including measures that prevent snares

from being set; we refer to these approaches as "reactive" and "proactive", respectively, even though

we acknowledge that these two categories are not always clear cut, and reactive patrolling may also

have proactive effects discouraging setting of snares (Moore et al, 2018; Dancer et al, 2022; results of

this study). To understand the financial investment needed for snare removal, we also investigated

the cost needed to implement these efforts, and possible sustainable opportunities to support this

undertaking. We discuss our findings within the wider context of a holistic framework for achieving

the long-term objective of zero snaring.

Methods

Collecting and processing patrol data

Ranger patrols were conducted in the contiguous Hue and Quang Nam Saola Nature Reserves by

Forest Guard teams, coordinated by World Wide Fund for Nature (WWF), together with local Forest

Protection Department staff. Threat data was input into Management Information System (MIST) and

Spatial Management and Reporting Tool (SMART) systems. MIST and SMART data were curated and

cleaned to create a database for analyses. We used patrol tracklog data to calculate monthly patrol

effort in a grid of 200 x 200 meter cells. Data for traps were converted into detection/non detection

matrices using monthly occasions using the same grid for analysis with occupancy models

(MacKenzie et al, 2017). See Supporting Information.

Description of the covariates

We considered the effect of five site covariates, which we assumed could influence snare site

selection by hunters, on snare occupancy: village density, remoteness, elevation, terrain ruggedness

(topographic ruggedness index), and topographic position index (TPI) (Figure S1). All covariates

were calculated in QGIS 2.18.9 (QGIS Development Team, 2016) and R version v4.1.3 (R Core Team,

2022). See Supporting Information.

Model description

We used a multi-season occupancy model (Mackenzie et al, 2017) to assess the distribution of snares

across both reserves. We implemented the model in a Bayesian framework in R using the *ubms*

package v1.1.9005 (Kellner et al, 2022). Monthly survey effort was calculated as the percentage of

each cell's area surveyed within a month. We divided the survey period into semesters (the first and

second half of each year) and used these as primary occasions. The second semester of 2012 and the

first semester of 2013 were removed due to missing data.

The final model included the effect of survey effort as a covariate on detection probability and the

five covariates on occupancy probability; a random effect of year on the coefficients of all occupancy

covariates and the occupancy intercept (allowing covariate coefficients and baseline occupancy to

vary between years); and a random effect of year-semester on detection probability (allowing

variation in detection probabilities, e.g. due to changes in staff). We found no evidence of spatial

autocorrelation. All models converged, and showed adequate fit (see Supporting Information).

We used parameter estimates to predict occupancy probabilities for each year and raster cell. From

these, we calculated the percentage of area occupied (PAO) as the percentage of cells predicted to be

occupied in each year for each posterior sample. The resulting distributions of PAO values were

summarized to obtain mean and 95% Bayesian credible intervals (BCIs) of PAO.

We also used occupancy models to assess the influence of past patrolling effort on snare occupancy.

Specifically, we fit separate single-season occupancy models to data from the second semester of

2016 and 2021, using the same covariate structure as described before, and adding as a covariate on

occupancy the total snare patrolling effort summed over the preceding six semesters, calculated for

each cell. We picked these two semesters as the approximate mid and endpoint of our time series.

Financial investment

We estimated the costs to remove a single snare by the Forest Guards, with data provided by WWF

(WWF internal and unpublished data), and the cost to set a snare by a local hunter, with cost

estimates provided by consultation with experts familiar with the study area (see Supporting

Information).

Results

Patrolling effort and snares removed

Between 2011 and 2021, the snare removal teams patrolled 3,040 days, covering 253,048 ha (Figure

1A). Patrol effort varied among and within years (Figure 1A; Figure S2). Overall, 99.2% of all 200 x

200 m cells received some level of effort within the study period (Figure S2). Patrols removed

118,151 snares from the two reserves (Figure 1A), with more snares removed from the Hue Saola

Nature Reserve compared to Quang Nam Saola Nature Reserve (Figure S3).

Snare occupancy over time

The multi-season occupancy model showed that the mean snare percent area occupied (PAO)

decreased by 36.9% (95% BCI [4.6, 59.0]) from 2011 to 2021 (Figure 1E & 1F; Figure S4). The largest

decline was 46.1% (95% BCI [21.8, 64.0]) between 2011 and 2016, after which PAO remained

approximately stable (Figure 1F). The decline in snare PAO was slightly stronger in the Quang Nam

Saola Nature Reserve (Figure 1D). The median snare occupancy across both sites during the last year

was 31.4% (95% BCI [23.6, 40.8], Figure 1C). Trap detection probability increased with the amount

of patrol effort (Figure 1B). Predicted snare occupancy increased with topographic position index

(TPI), elevation, remoteness, and ruggedness, and decreased with village density (Figure 1C; Table

S2).

Finally, snare occupancy probability decreased with increasing previous patrol effort (Figure 1G);

the effect was significant (95% BCIs did not overlap 0) in 2016 (-0.48, 95% BCI [-0.77, -0.24]) and

2021 (-0.67, 95% BCI [-0.99, -0.40]), but had a larger magnitude in 2021.

Financial investment

To achieve this level of snare reduction, it was necessary to invest approximately 220,000 USD

annually (WWF unpublished data), resulting in an average cost of 20.5 USD per snare removed; this

is more than an order of magnitude more expensive than the cost of setting a single snare (1.13 USD;

Figure 1H). During the initial years, patrol costs were covered by WWF, with financing provided

through international biodiversity conservation aid funds. In 2014 patrol costs began to be

181 supplemented by a Payment for Forest Environmental Service (PFES) initiative by the Vietnamese

182 government, and provided directly to the provincial management authorities. Some Forest Guard

183 costs began to be covered by PFES funding starting in 2015, increasing to 15% in 2021 (WWF

184 unpublished data).

186 **Discussion**

187 We show that, with intensive and sustained snare removal, snaring levels can be significantly reduced

188 within tropical protected areas. Snare removal is therefore an important component of strategies to

189 counter the widespread snaring in Southeast Asia, if employed at the appropriate spatial and

190 temporal scales. The constant effort needed to reduce snaring and maintain depressed snaring levels

191 may require conservation stakeholders to re-think how they approach site-based conservation,

192 especially since most initiatives are funded by short-term projects that last at best a few years.

193 Clearly, for protected areas facing high levels of snaring pressure, long-term engagement is needed

194 to counter this threat. Overall snare occupancy declined steadily for the first six years of patrolling,

195 and then appeared to plateau. This may indicate that, after an initial decline in snaring, subsequent

196 reductions are progressively more difficult to secure. Furthermore, declines in snaring were stronger

197 near the edges of the reserves and along access points. Such a finding highlights the importance of

198 increasing patrol effort in remote areas, especially since these are more likely to harbor

199 conservation-priority species (Tilker et al, 2020). In this way, robust monitoring of patrol data has

200 the potential to offer insights into patrolling strategies that can help inform adaptive management.

201 Our finding that prior patrolling decreased the probability of snare occupancy suggests that

202 patrolling acts as a deterrent to future snaring. Other studies have also indicated the deterrence value

203 of patrolling (Moore et al, 2028; Dancer et al, 2022). One possible explanation is that, with continued

204 snare removal, hunters experience increasingly diminishing rates of return, and thus the cost-benefit

ratio shifted so that it was less profitable to set snares in the study areas. In other parts of the world,

the threat of arrest or fines might also serve as a deterrent; in our sites in central Vietnam, however,

arrests for snaring are rare, due in part to legal loopholes that do not penalize people for having wire

materials inside protected areas (Belecky & Gray 2020; Gray et al, 2021), and in part because snaring

is seldom treated as a serious forest crime. We note that it is possible that local reductions in snaring

in our sites caused poaching to spill over to other protected areas which did not receive intensive

patrol effort; similar patterns of displacement have been documented for illegal logging in tropical

protected areas (Ford et al, 2020). Additional studies are needed to assess potential leakage and, if

present, ways to counteract it.

Despite the success of demonstrably reducing snaring levels in these protected areas, the snaring

levels that persist suggests that wildlife populations are still threatened by unsustainable offtake.

Snaring has already contributed to the loss or functional extinction of many larger vertebrates from

these areas (Tilker et al, 2019) – including the saola (*Pseudoryx nghetinhensis*), a Critically

Endangered bovid that these reserves were established to protect – and it seems likely that

remaining conservation-priority and snaring-sensitive species will continue to decline unless snaring

pressure is reduced further. However, we note that because species show different abilities to

recover from unsustainable harvest, largely due to life history characteristics (Cardillo et al, 2008), it

is possible that populations of some mammals and ground-dwelling birds will stabilize or increase

with this level of snare reduction. Long-term wildlife monitoring is needed to assess population

dynamics and potential recovery for individual species.

The financial investment necessary to attain such a level of snare reduction was substantial. With an

estimated 13 million snares in the protected areas of Viet Nam, Laos, and Cambodia alone (Belecky

& Gray, 2020), the cost to remove existing snares in Southeast Asian protected areas would likely

sum to hundreds of millions of dollars. Moreover, the high cost of snare removal prompts questions

about the long-term sustainability and scalability of this approach. It was only with a substantial and

230 continual investment from an external NGO that such intensive levels of snare removal were possible

231 in the two sites. Given the general lack of financial and personnel resources, capacity, and

232 government support that characterizes most protected areas in the region (Graham et al, 2021),

233 similar snare removal projects can likely only be implemented if NGO stakeholders work in

234 collaboration with government protected area staff. One option to increase the financial

235 sustainability of snare removal is to access sustainable financing schemes, such as Payment for

236 Ecosystem Services (PES, PFES in Viet Nam), Reducing Emissions from Deforestation and forest

237 Degradation (REDD+), or Carbon and Biodiversity offsets. The use of such incentive programs seems

238 to be increasing: In 2020, more than 80% of protected areas in Viet Nam obtained PFES funds at an

239 average of 323 USD per km^2 (Emerton et al, 2021), which based on Decree No. 156/2018/ND-CP can

240 be used for forest and biodiversity protection activities. However, these sustainable financing

241 schemes are unlikely to be a shortcut to complete financing of snare removal operations; in the two

242 reserves in our study, only 15% of the total costs could be covered by PFES in 2020 and 2021.

243 Furthermore, funding alone will not guarantee effective replication of intensive snare removal in

244 Southeast Asian protected areas. To guarantee transparency and accountability, protected area

245 managers should set clear targets for threat reduction, and these metrics should be monitored. Our

246 analytical workflow provides a streamlined approach to facilitate such robust analyses of patrol data

247 in the future.

248 Our results indicate that snare removal alone is likely insufficient to protect wildlife in Southeast

249 Asian protected areas. Although this is not unexpected, since snare removal does not address the

250 fundamental drivers behind snaring, our results provide strong empirical evidence that a

251 multifaceted approach is needed that combines short-term immediate responses within protected

252 areas, such as snare removal, with long-term approaches in rural and urban areas (Figure 2). Such

253 long-term responses could include demand reduction for illegal wildlife products (Shairp et al, 2016),

254 strengthened legislation that penalizes the possession of wire material inside protected areas

(Belecky & Gray, 2020), community engagement that promotes local guardianship (Viollaz et al,

2022), improved law enforcement coordination and capacity (Dudley et al, 2013), and engagement

with local judicial authorities to ensure appropriate prosecution and sentencing (Nurse & Nurse,

2015). Simultaneously, actions which support inclusive governance of protected areas and nature-

based and nature-positive development opportunities for buffer zone communities are required to

offset potential losses of income, ensure equitable benefit sharing, and harmonize both economic

advancement and biodiversity management and protection (Borrini et al, 2004; Andrade & Rhodes,

2012). Together, the activities needed for such a holistic approach will involve coordination among

multiple government and non-governmental stakeholders and will require considerable financial

investment.

Overall, our results are encouraging in that they show that sustained snare removal can be an

effective method for reducing snaring. Therefore, intensive snare removal schemes should be

expanded to other protected areas as a first and immediate response to safeguard highly threatened

wildlife populations. For these populations, snare removal could buy crucial time needed to

implement more proactive and long-term approaches, many of which could take time to become

effective. Ideally, these efforts should be funded through sustainable financing mechanisms to allow

for the long-term continuity necessary to suppress snaring. We acknowledge that, because the

drivers behind snaring are so complex, the impact of sustained and intensive snare removal in other

areas may differ from the results that we present here, nonetheless, the high demand for wild meat

in urban centers fueling snaring in central Vietnam is present across Southeast Asia (Lee et al, 2014;

Belecky & Gray, 2020), and we therefore believe that the situation in our sites is representative of the

larger snaring problem in the region. Perhaps most importantly, our findings illustrate the

importance of viewing snare removal as one component of a wider and multi-faceted conservation

response that addresses these underlying drivers; to rely solely on snare removal to safeguard

protected areas is unlikely to sufficiently address the threat. If we want to prevent the extinction of

many of the region's iconic megafaunal species, as well as to preserve the health of the tropical

ecosystems that Southeast Asian societies depend upon, snare removal operations will need to be

embedded within a more holistic and proactive framework that prevents snares from being set in the

first place.

284
Acknowledgements

We thank the staff of the WWF-CarBi project (phase I and phase II), the Thua Thien Hue Saola Nature

Reserve, and the Quang Nam Saola Nature Reserve for supporting data collection, including through

providing personnel for snare removal. The CarBi projects were and are still funded by the

International Climate Initiative of the Federal Ministry for the Environment, Nature Conservation,

Nuclear Safety and Consumer Protection (BMUV) through the German Development Bank (KfW). A.T.

would like to thank Barney Long for productive discussions over the years on snare removal within

the context of protected area management, Julie Louvrier for advice on study design and analysis,

and Lucy Emerton for input on protected area financing in Vietnam.

References

Andrade, G. S., & Rhodes, J. R. (2012). Protected areas and local communities: an inevitable

partnership toward successful conservation strategies? *Ecology and Society*, 17(4).

Barlow, J., França, F., Gardner, T.A., Hicks, C.C., Lennox, G.D., Berenguer, E., Castello, L., Economo, E.P.,

Ferreira, J., Guénard, B. & Graham, N. A. (2018). The future of hyperdiverse tropical ecosystems.

Nature, 559(7715), 517-526.

Becker, M., McRobb, R., Watson, F., Droge, E., Kanyembo, B., Murdoch, J., & Kakumbi, C. (2013).

Evaluating wire-snare poaching trends and the impacts of by-catch on elephants and large

carnivores. *Biological Conservation*, 158, 26-36.

Belecky, M. & Gray, T.N.E, (2020). Silence of the Snares: Southeast Asia's Snaring Crisis, WWF
International.

Bello, C., Galetti, M., Pizo, M.A., Magnago, L.F.S., Rocha, M.F., Lima, R.A., Peres, C.A., Ovaskainen, O. &
Jordano (2015). Defaunation affects carbon storage in tropical forests. *Science Advances*, 1(11),
e1501105.

Benítez-López, A., Alkemade, R., Schipper, A. M., Ingram, D. J., Verweij, P. A., Eikelboom, J. A. J., &
Huijbregts, M. A. J. (2017). The impact of hunting on tropical mammal and bird populations. *Science*,
356(6334), 180-183.

Benítez-López, A., Santini, L., Schipper, A. M., Busana, M., & Huijbregts, M. A. (2019). Intact but empty
forests? Patterns of hunting-induced mammal defaunation in the tropics. *PLoS Biology*, 17(5),
e3000247.

Bogoni, J. A., Peres, C. A., & Ferraz, K. M. (2020). Effects of mammal defaunation on natural ecosystem
services and human well being throughout the entire Neotropical realm. *Ecosystem Services*, 45,
101173.

Borrini, G., Kothari, A., & Oviedo, G. (2004). Indigenous and local communities and protected areas:
Towards equity and enhanced conservation: Guidance on policy and practice for co-managed
protected areas and community conserved areas (No. 11). IUCN, Gland, Switzerland.

Cardillo, M., Mace, G. M., Gittleman, J. L., Jones, K. E., Bielby, J., & Purvis, A. (2008). The predictability
of extinction: biological and external correlates of decline in mammals. *Proceedings of the Royal*
Society B: Biological Sciences, 275(1641), 1441-1448.

Coad, L., Watson, J.E., Geldmann, J., Burgess, N.D., Leverington, F., Hockings, M., Knights, K. & Di Marco,
M. (2019). Widespread shortfalls in protected area resourcing undermine efforts to conserve
biodiversity. *Frontiers in Ecology and the Environment*, 17(5), 259-264.

Dancer, A., Keane, A., Beale, C.M., Dobson, A.D., Amin, R., Freeman, R., Imong, I., Jones, K., Linkie, M.,

Long, B. & Collen, B. (2022). Evidence of deterrence from patrol data: Trialling application of a

differenced-CPUE metric. *Conservation Science and Practice*, 4(8), e12746.

Dudley, N., Stolton, S., & Elliott, W. (2013). Wildlife crime poses unique challenges to protected areas.

Parks, 19(1), 7-12.

Emerton, L., Nguyễn, V. D., Bui, T. M. N., & Roth, M. (2021). Review of PA financial status in Viet Nam:

'self-financing' needs, options & ways forward. Report to Conservation, Sustainable Use of Forest

Biodiversity & Ecosystem Services in Viet Nam. GIZ-Bio Project Phase II.

Ford, S. A., Jepsen, M. R., Kingston, N., Lewis, E., Brooks, T. M., MacSharry, B., & Mertz, O. (2020).

Deforestation leakage undermines conservation value of tropical and subtropical forest protected

areas. *Global Ecology and Biogeography*, 29(11), 2014-2024.

Graham, V., Geldmann, J., Adams, V. M., Grech, A., Deinet, S., & Chang, H. C. (2021). Management

resourcing and government transparency are key drivers of biodiversity outcomes in Southeast

Asian protected areas. *Biological Conservation*, 253, 108875.

Gray, T.N., Belecky, M., O'Kelly, H.J., Rao, M., Roberts, O., Tilker, A., Signs, M. & Yoganand, K. (2021).

Understanding and solving the South-East Asian snaring crisis. *The Ecological Citizen*, 4, 129-141.

Harrison, R.D., Sreekar, R., Brodie, J.F., Brook, S., Luskin, M., O'Kelly, H., Rao, M., Scheffers, B. & Velho,

N. (2016). Impacts of hunting on tropical forests in Southeast Asia. *Conservation Biology*, 30(5), 972-

981.

Jachmann, H. (2008). Illegal wildlife use and protected area management in Ghana. *Biological*

Conservation, 141(7), 1906-1918.

Kellner, K. F., Fowler, N. L., Petroelje, T. R., Kautz, T. M., Beyer Jr, D. E., & Belant, J. L. (2022). ubms: An
R package for fitting hierarchical occupancy and N-mixture abundance models in a Bayesian
framework. *Methods in Ecology and Evolution*, 13(3), 577-584.

Krause, T., & Tilker, A. (2022). How the loss of forest fauna undermines the achievement of the SDGs.
Ambio, 51(1), 103-113.

Lacher Jr, T.E., Davidson, A.D., Fleming, T.H., Gómez-Ruiz, E.P., McCracken, G.F., Owen-Smith, N., Peres,
C.A. and Vander Wall, S. B. (2019). The functional roles of mammals in ecosystems. *Journal of*
Mammalogy, 100(3), 942-964.

Laurance, W.F., Carolina Useche, D., Rendeiro, J., Kalka, M., Bradshaw, C.J., Sloan, S.P., Laurance, S.G.,
Campbell, M., Abernethy, K., Alvarez, P. & Arroyo-Rodriguez, V. (2012). Averting biodiversity collapse
in tropical forest protected areas. *Nature*, 489(7415), 290-294.

Lee, T. M., Sigouin, A., Pinedo-Vasquez, M., & Nasi, R. (2014). The harvest of wildlife for bushmeat and
traditional medicine in East, South and Southeast Asia: Current knowledge base, challenges,
opportunities and areas for future research. Occasional Paper 115. Bogor, Indonesia: CIFOR.

Linkie, M., Martyr, D. J., Harihar, A., Risdianto, D., Nugraha, R. T., Leader-Williams, N., & Wong, W. M.
(2015). Safeguarding Sumatran tigers: evaluating effectiveness of law enforcement patrols and local
informant networks. *Journal of Applied Ecology*, 52(4), 851-860.

MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., Bailey, L. L., & Hines, J. E. (2017). *Occupancy*
estimation and modeling: inferring patterns and dynamics of species occurrence. Elsevier.

Moore, J. F., Mulindahabi, F., Masozera, M. K., Nichols, J. D., Hines, J. E., Turikunkiko, E., & Oli, M. K.
(2018). Are ranger patrols effective in reducing poaching-related threats within protected areas?
Journal of Applied Ecology, 55(1), 99-107.

Nurse, A., & Nurse, A. (2015). "National wildlife legislation and law enforcement policies." In *Policing*

wildlife: Perspectives on the enforcement of wildlife legislation, pp. 63-82.

Osuri, A.M., Ratnam, J., Varma, V., Alvarez-Loayza, P., Hurtado Astaiza, J., Bradford, M., Fletcher, C.,

Ndoundou-Hockemba, M., Jansen, P.A., Kenfack, D., Sankaran, M. (2016). Contrasting effects of

defaunation on aboveground carbon storage across the global tropics. *Nature Communications*, 7(1),

1-7.

QGIS Development Team. (2016). QGIS Geographic Information System. Open Source Geospatial

Foundation. URL http://qgis.org

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Shairp, R., Veríssimo, D., Fraser, I., Challender, D., & MacMillan, D. (2016). Understanding urban

demand for wild meat in Vietnam: implications for conservation actions. *PloS One*, 11(1), e0134787.

Tilker, A., Abrams, J.F., Mohamed, A., Nguyen, A., Wong, S.T., Sollmann, R., Niedballa, J., Bhagwat, T.,

Gray, T.N., Rawson, B.M., Guegan, F., Kissing, J., Wegmann, M., & Wilting, A. (2019). Habitat

degradation and indiscriminate hunting differentially impact faunal communities in the Southeast

Asian tropical biodiversity hotspot. *Communications Biology*, 2(1), 1-11.

Tilker, A., Abrams, J.F., Nguyen, A., Hörig, L., Axtner, J., Louvrier, J., Rawson, B.M., Nguyen, H.A.Q.,

Guegan, F., Nguyen, T.V., Le, M., Sollmann, R., & Wilting, A. (2020). Identifying conservation priorities

in a defaunated tropical biodiversity hotspot. *Diversity and Distributions*, 26(4), 426-440.

Viollaz, J., Long, B., Trung, C.T., Kempinski, J., Rawson, B.M., Quang, H.X., Hiền, N.N., Liên, N.T.B., Dũng,

C.T., Huyền, H.T., McWhirter, R., Dung, N. T. T., & Gore, M. L. (2021). Using crime script analysis to

understand wildlife poaching in Vietnam. *Ambio*, 50(7), 1378-1393.

Watson, F., Becker, M. S., McRobb, R., & Kanyembo, B. (2013). Spatial patterns of wire-snare poaching:

Implications for community conservation in buffer zones around National Parks. *Biological*

Conservation, 168, 1-9.

Watson, J. E., Dudley, N., Segan, D. B., & Hockings, M. (2014). The performance and potential of

protected areas. *Nature*, 515(7525), 67-73.

Young, H.S., Dirzo, R., Helgen, K.M., McCauley, D.J., Billeter, S.A., Kosoy, M.Y., Osikowicz, L.M., Salkeld,

D.J., Young, T.P, & Dittmar, K. (2014). Declines in large wildlife increase landscape-level prevalence

of rodent-borne disease in Africa. *Proceedings of the National Academy of Sciences*, 111(19), 7036-

7041.

Figures

Figure 1: (A) Total number of snares and patrol effort (ha); (B) response of snare detection

probability to survey effort; (C) response of snare occupancy probability to covariates; thick black

line shows mean effect for all years, gray shaded area show 95% confidence intervals (B and C), and

light gray lines show mean effect for individual years (C only); (D) median snare occupancy

probability for survey period (2011-2021); (E) median change in snare occupancy probability across

the survey period; (F) Percentage of area occupied (PAO) values for individual years; black dot

indicates median and line indicates 50% posterior mass; based on predictions from multi-season

occupancy model (D - F); (G) response of snare occupancy to prior patrol effort for two time periods,

2016 and 2021; and (H) cost of setting and removing an individual snare in the study areas, with

removal cost divided by Payment for Forest Ecosystem Services and NGO aid funds.

Figure 2: Conceptual framework for achieving the target of zero snaring using a combination of
reactive and proactive approaches across different environments and time scales.

429