1 **PP1β opposes classic PP1 function, inhibiting spine maturation and promoting LTP**
2 Short title: PP1β is a memory enhancer
3

4 **Authors**

5 Karl Foley,[1,2] Cody McKee,[1,2] Archan Ganguly,[1] Daniel Barnett,[1] Nancy Ward,[1] Abigail
6 Mayer,[1,2] Yu Zhang,[1] Angus C. Nairn,[3] & Houhui Xia[1,2*]
7

8 **Affiliations**

9 [1]Department of Neuroscience, University of Rochester Medical Center, Rochester, NY 14642

10 [2]Department of Pharmacology and Physiology, University of Rochester Medical Center,
11 Rochester, NY 14642

12 [3]Department of Psychiatry, Yale University, New Haven, Connecticut, 06511
13

14 *Email: Houhui_xia@urmc.rochester.edu, Phone: 585-275-3853, Fax: 585-275-6654
15

Summary
Protein phosphatase 1 (PP1) regulates synaptic plasticity and has been described as a molecular
constraint on learning and memory. There are three neuronal isoforms, PP1α, PP1β, and PP1γ,
but little is known about their individual functions. PP1α and PP1γ are assumed to mediate the
effects of PP1 on learning and memory based on their enrichment at dendritic spines and their
preferential binding to neurabin and spinophilin, major PP1 synaptic scaffolding proteins.
However, it was recently discovered that human de novo PP1β mutations cause intellectual
disability, suggesting an important but ill-defined role for PP1β. In this study, we investigated
the functions of each PP1 isoform in hippocampal synaptic physiology using conditional CA1-
specific knockout mice. In stark contrast to classic PP1 function, we found that PP1β promotes
synaptic plasticity as well as spatial memory. These changes in synaptic plasticity and memory
are accompanied by changes in GluA1 phosphorylation, GluN2A levels, and dendritic spine
density and morphology, including silent synapse number. These functions of PP1β reveal a
previously unidentified signaling pathway regulating spine maturation and plasticity, broadening
our understanding of the complex role of PP1 in synaptic physiology.
31

32 **Keywords**
33 Protein phosphatase 1 isoforms (PP1); PP1β; PP1γ; synaptic transmission; spine; silent synapse;
34 Long term potentiation (LTP); memory
35

Introduction

Protein phosphatase 1 (PP1) is a highly conserved serine (S)-threonine (T) phosphatase that regulates a diverse array of cellular processes. The multifunctional roles of PP1 are mediated by distinct holoenzymes consisting of a PP1 catalytic subunit and one or more regulatory subunits (Bollen et al., 2010; Foley et al., 2021a; Heroes et al., 2013). The PP1 catalytic subunit is encoded by three genes, giving rise to PP1α, PP1β, and PP1γ isoforms which have high sequence similarity but differ in their binding affinity to regulatory proteins. The PP1 catalytic subunits possess several shared binding domains that accommodate substrates of varied physicochemical properties, while the regulatory subunits selectively restrict or promote PP1 activity towards specific substrates. Therefore, PP1 holoenzymes can dephosphorylate a wide range of substrates collectively, while individual holoenzymes maintain high substrate specificity. Further, PP1 isoforms participate in distinct but overlapping pools of PP1 holoenzymes.

In the brain, PP1 regulates synaptic transmission and plasticity (Aow et al., 2015; Foley et al., 2022; Gao et al., 2018; Hu et al., 2006; Hu et al., 2007; Jouvenceau et al., 2006; Morishita et al., 2001) and its partial inhibition improves spatial learning and memory in mice (Genoux et al., 2002). Given the potential therapeutic value of inhibiting PP1 in conditions with learning and memory impairments, identifying the associated PP1 holoenzyme and elucidating its molecular function is of great interest. Significant progress has been made through the identification of PP1 isoforms and regulatory proteins enriched at the synapse. Namely, neurabin (*PPP1R9A*) and spinophilin (*PPP1R9B*) are prominent synaptic scaffolding proteins associated with the actin cytoskeleton that target PP1 to the synapse (Terry-Lorenzo et al., 2002), preferentially binding PP1γ, followed by PP1α, and to a minimal extent PP1β (Carmody et al ., 2004). Consistent with this, PP1γ and PP1α, but not PP1β, are concentrated in spines (Bordelon et al., 2005). Selective disruption of the neurabin-PP1 holoenzyme via mutagenesis recapitulates the effect of PP1 partial inhibition on synaptic plasticity (Hu *et al.*, 2006; Hu *et al.*, 2007; Jouvenceau *et al.*, 2006). However, non-selective inhibition of PP1 has different effects on synaptic transmission than selective disruption of neurabin-PP1, suggesting there are other PP1 holoenzymes active at the synapse (Hu *et al.*, 2007). Consistent with this notion, all three PP1 isoforms and many additional PP1 regulatory proteins are present at the post-synaptic density (Bayes et al., 2012; Bayes et al., 2011; Foley *et al.*, 2021a; Li et al., 2016), although their distinct synaptic functions have not been investigated.

While the majority of studies on PP1 in synaptic plasticity have focused on neurabin and spinophilin, which preferentially bind PP1γ and PP1α, recent evidence suggests PP1β has a significant but ill-defined role in neuronal physiology. Several groups have discovered human de novo mutations in PP1β associated with severe intellectual disability (Gripp et al., 2016; Hamdan et al., 2014; Huckstadt et al., 2021; Lin et al., 2018; Ma et al., 2016; Umeki et al., 2019; Zambrano et al., 2017; Zhou et al., 2020), with none yet identified in PP1α and PP1γ. Further, mRNA expression data suggest PP1β could promote learning and memory: high-performing rodents in the Morris Water Maze have significantly lower levels of PP1γ (Haege et al., 2010) but higher levels of PP1β (Burger et al., 2007) in the hippocampus compared to poor-performing rodents.

In this study, we sought to define the functions of each PP1 isoform in hippocampal synaptic physiology. Using conditional CA1-specific knockout (KO) mice, we found that PP1β has

distinct and opposing functions to PP1α and PP1γ in regulating synaptic transmission and
plasticity as well as spatial learning and memory. Correspondingly, PP1β KO increases S831
GluA1 phosphorylation, decreases GluN2A levels, and increases dendritic spine density and
maturation, including silent synapse number.

Results
PP1 isoforms differentially regulate basal synaptic transmission
We have previously shown that genetic KO of PP1γ, but not PP1α, in neural precursor cells
(NPCs) leads to a reduction in basal synaptic transmission in the hippocampus (Foley *et al.*,
2022). Here, we report that KO of PP1β in NPCs is embryonic lethal, similar to previous
findings that global PP1β KO is lethal whereas global PP1α and PP1γ KO mice are viable
(Dickinson et al., 2016; International Mouse Phenotyping, 2022). We therefore generated CA1-
specific conditional KO mice (cKO) to compare the functions of each PP1 isoform by breeding
PP1 isoform floxed mice with a CaMKIIα-Cre transgenic line. We confirmed CA1-specific Cre
expression at 1-1.5 months of age through breeding with an Ai9 reporter line (Figure 1A).
Interestingly, cKO of each PP1 isoform had a distinct effect on hippocampal synaptic
transmission, as determined by recording field excitatory postsynaptic potentials (fEPSPs) from
CA3 Schaffer collateral-CA1 (Sch-CA1) synapses in acute hippocampal slices. PP1γ cKO
reduced synaptic transmission whereas PP1α cKO had no effect (Figure 1B-C), replicating the
findings in NPC-KO mice (Foley *et al.*, 2022) and the effect of disrupting neurabin-PP1
interaction (Hu *et al.*, 2007). To test if an effect of PP1α cKO is masked by PP1γ compensation,
we generated a PP1α/γ double cKO mouse. PP1α/γ cKO reduced synaptic transmission (Figure
1E) to a greater degree than PP1γ cKO alone (Figure 1F), suggesting that PP1α plays a
secondary, redundant role to PP1γ in regulating synaptic transmission, rather than no role at all.
Surprisingly, PP1β cKO increased synaptic transmission (Figure 1D), an unprecedented finding
in PP1 synaptic function. Overall, the findings suggest that PP1 isoforms play opposing roles in
regulating hippocampal synaptic transmission, with PP1γ and PP1α promoting transmission and
PP1β attenuating transmission.
106
PP1 isoforms differentially regulate LTP.
To investigate the roles of PP1 isoforms in synaptic plasticity, we induced LTP via a single bout
of high-frequency stimulation. Given the effects on basal synaptic transmission (Figure 1), we
chose to focus on PP1α/γ cKO and PP1β cKO mice. PP1α/γ cKO had no effect on classic LTP
induction (Figure 2C; 100 pulses, 100Hz) at Sch-CA1 synapses, but significantly increased the
potentiation following sub-threshold LTP stimulation (Figure 2B; 25 pulses, 100Hz). In contrast,
PP1β cKO significantly attenuated LTP (Figure 2A; 100 pulses, 100Hz). Similar to the effects
observed for basal synaptic transmission, the findings suggest PP1β opposes PP1α/PP1γ function
in regulating hippocampal synaptic plasticity.

Many different factors can impair postsynaptic depolarization and thus impair NMDAR activity
during LTP stimulation (Granger and Nicoll, 2014). To exclude impaired postsynaptic
depolarization as a cause of the LTP deficit in PP1β cKO mice, we induced LTP in CA1
pyramidal neurons using whole-cell patch clamp via the pairing protocol (Figure 2D). In this
approach, postsynaptic depolarization is controlled by holding the cell at 0 mV during LTP
stimulation. Further, we inhibited GABA$_A$ receptors by including 50μM picrotoxin in the bath to
exclude altered inhibitory signaling as a mechanism of the LTP deficit. Consistent with the LTP

deficit observed via field recordings, we found that PP1β cKO also significantly attenuated LTP
induced via the pairing protocol (Figure 2E).

PP1β cKO impairs spatial learning and memory
Changes in hippocampal LTP are often associated with changes in spatial learning and memory.
We therefore sought to test CA1-dependent learning and memory with the allocentric object-
place assay (Figure 3A), a variation of novel-object recognition (Langston and Wood, 2009).
PP1β cKO mice explored the novel object significantly less than their control littermates (Figure
3B), suggesting impaired short-term spatial memory. PP1α/γ cKO mice did not demonstrate any
change in novel object recognition, exploring the novel object to the same degree as their
littermate controls (Figure 3F). Neither PP1β cKO nor PP1α/γ cKO mice demonstrated any
alteration in the total time exploring both novel and familiar objects (Figure 3C, G).
Additionally, there was no change in locomotor activity or anxiety, as measured in open field
testing (Figure 3D-E, H-I).

PP1β regulates AMPAR phosphorylation and GluN2A levels
Our findings thus far demonstrate that PP1β has distinct, opposing functions to PP1α and PP1γ in
the CNS. We therefore sought to identify altered signaling pathways and measure glutamatergic
receptor levels. In immunoblotting experiments on CA1 microdissections, we observed no
change in several pathways involved in synaptic plasticity and previously shown to be regulated
by PP1, including pCREB, pCaMKII, pERK (Figure 4A). However, PP1β cKO mice had
significantly increased levels of phosphorylated GluA1 AMPAR at S831 (Figure 4B), a site
involved in LTP (Lee et al., 2000; Mammen et al., 1997) that increases AMPAR channel
conductance (Derkach et al., 1999) and contributes to GluA1 targeting to the post-synaptic
density (PSD) (Diering et al., 2016; Diering and Huganir, 2018) (but see (Hayashi et al., 2000)).
There was no significant change in phosphorylation at GluA1 S845 or GluA2 S880, which are
dephosphorylated during LTD (Hu *et al.*, 2007; Lee et al., 1998), and no change in total GluA1,
GluA2, or GluA3 levels (Figure 4B). We also observed a decrease in GluN2A levels in PP1β
cKO mice (Figure 4C). Meanwhile, there was no change in GluN2B pS1480, a known PP1 target
that regulates GluN2B lateral diffusion (Chiu et al., 2019), or total GluN2B or GluN1 levels
(Figure 4C).

PP1β regulates spine density, morphology, and silent synapses
Given the increase in synaptic transmission and GluA1 pS831 in PP1β cKO mice, we assessed
miniature excitatory postsynaptic currents (mEPSCs) in CA1 pyramidal cells to measure
synaptic AMPA receptor (AMPAR) content. Surprisingly, PP1β cKO mice showed a robust
increase in mEPSC frequency but not amplitude (Figure 4E-F). Interestingly, although the mean
amplitude was not significantly different, there was a significant shift in the cumulative
frequency, with a higher proportion of events in the 15-50 pA range in PP1β cKO cells (Figure
4E). A change in mEPSC frequency can reflect an increase in the docked vesicle pool or an
increase in synapse number. To test for a presynaptic cause, we measured paired-pulse
facilitation (PPF) at Sch-CA1 and observed no difference between PP1β cKO and control
littermates (Figure S1A). We also observed no difference between slice preparation protocols
(Figure S1B-E).

The lack of change in PPF combined with the postsynaptic KO approach strongly suggested an increase in synapse number. We therefore analyzed dendritic spine density in proximal dendrites of CA1 pyramidal neurons by confocal fluorescence microscopy of biocytin-filled neurons (Figure 4H). PP1β cKO mice had a significant increase in spine density (Figure 4I), as assessed by Imaris reconstruction of imaged dendrites. The increase in spine density was driven by an increase in mature mushroom spines (Figure 4J).

In addition to morphology, spines can be classified as functional or silent based on their expression or lack of expression of AMPARs, respectively. The specific increase in mature spines in PP1β cKO mice suggests that the increased mEPSC frequency could result from an increase in the number of functional, AMPAR-containing spines in addition to the increase in total number of spines. To test this possibility, we measured AMPAR- and NMDAR-mediated currents in response to minimal stimulation (Figure 4K). Stimulation intensity was lowered until approximately 50% of stimulations failed to elicit a response at -70 mV holding potential. We then measured the failure rate at +40 mV to determine the proportion of silent synapses, which exhibit a detectable response at +40 mV but not at -70 mV. PP1β cKO mice showed fewer silent synapses (Figure 4L), suggesting that PP1β cKO increases mEPSC frequency by increasing both the total number and AMPAR-expressing number of synapses.

Discussion

PP1 is classically viewed as a molecule of forgetfulness, with its inhibition improving spatial learning and memory and promoting LTP over LTD (Genoux *et al.*, 2002; Jouvenceau *et al.*, 2006). However, PP1 can have broad functions, given that PP1 refers to three isoforms and hundreds of PP1 holoenzymes exist with presumed distinct functions. Here, we demonstrate that PP1β overall opposes PP1γ and PP1α function in synaptic transmission, plasticity, and spatial learning and memory. Whereas PP1α/PP1γ cKO decreases synaptic transmission in Sch-CA1 synapses, consistent with neurabin-PP1 function, PP1β cKO increases synaptic transmission. Similarly, while PP1α/PP1γ cKO promotes LTP, PP1β cKO attenuates LTP. Finally, PP1β cKO impairs spatial learning and memory in a novel object recognition assay, whereas PP1α/PP1γ cKO has no effect. These effects are accompanied by changes in spine morphology, S831 GluA1 phosphorylation, and GluN2A levels. These novel functions of PP1β reveal a previously unrecognized complexity of PP1 function in synaptic physiology.

The opposing roles of PP1β help resolve discrepancies between the effects of non-selective PP1 inhibition and disruption of specific holoenzymes. We previously showed that non-selective PP1 inhibition has no effect on synaptic transmission whereas disruption of neurabin-PP1 decreases synaptic transmission (Hu *et al.*, 2007), suggesting the existence of an additional PP1 holoenzyme that opposes neurabin-PP1 function. Our findings suggest that a PP1β holoenzyme mediates this opposing role.

The impairment in spatial learning in PP1β cKO mice stands in stark contrast to the improvement seen with partial PP1 inhibition (Genoux *et al.*, 2002). One explanation is that the relevant PP1β holoenzyme remains sufficiently active in the partial inhibition paradigm. In the aforementioned studies, partial inhibition of PP1 was achieved via overexpression of a constitutively active inhibitor-1 fragment (I-1*) (Genoux *et al.*, 2002; Jouvenceau *et al.*, 2006). It is unclear if all PP1 holoenzymes are equally and proportionally inhibited by I-1*, or if the

partial inhibition primarily affects a subset of holoenzymes. I-1-PP1 interaction involves the RVXF binding domain and is predicted to involve the PP1 acidic groove (Liang et al., 2018; Terrak et al., 2004). I-1* may therefore have higher binding affinity to PP1 holoenzymes in which the acidic groove is fully exposed, such as neurabin-PP1 (Ragusa et al., 2010). Phosphoproteomic analysis of the I-1* transgenic mouse would likely resolve these questions and provide valuable insights into the mechanisms by which PP1 partial inhibition improves learning and memory.

The mechanism by which PP1β cKO alters synaptic transmission and plasticity appears to be explained by increased phosphorylation of S831 GluA1 and decreased GluN2A levels. Phosphorylation of GluA1 at S831 increases AMPAR channel conductance (Derkach *et al.*, 1999) and contributes to GluA1 targeting to the PSD under basal conditions (Diering *et al.*, 2016; Diering and Huganir, 2018), consistent with the observed increase in synaptic transmission. Increased targeting of GluA1 to the PSD via S831 phosphorylation may help account for the decrease in silent synapses in PP1β cKO mice, though the contribution of GluA1 phosphorylation to unsilencing has not been directly investigated. LTP is accompanied by an increase in GluA1 S831 phosphorylation (Lee *et al.*, 2000; Mammen *et al.*, 1997), suggesting the increase in pS831 in PP1β cKO may also contribute to the LTP deficit by saturating the contribution of pS831 to potentiation. However, dual-phosphomimetic (S831D/S845D) GluA1 mice show a decrease in the LTP threshold (Makino et al., 2011). On the other hand, the effect of S831D and S845D alone on LTP has not been reported, therefore it is unknown if one or both phosphorylation sites mediates the decreased LTP threshold.

The decrease in total GluN2A levels in PP1β cKO is consistent with the impairment of LTP and spatial learning and memory (Kiyama et al., 1998; Sakimura et al., 1995). While the relative contributions of GluN2A and GluN2B to LTP are still debated, classic pharmacologic and genetic ablation studies have demonstrated that GluN2A contributes significantly to LTP at Sch-CA1 (Shipton and Paulsen, 2014). Interestingly, GluN2B-containing NMDARs are enriched in small spines, and GluN2A-containing NMDARs in larger spines (Sobczyk et al., 2005), opening the possibility that the observed potentiation deficit could vary between spine subtypes. Single-synapse plasticity studies are needed to test this possibility.

How does PP1β regulate GluA1 S831 phosphorylation and GluN2A levels? Each function of PP1 is mediated by a specific holoenzyme, consisting of the PP1 catalytic subunit and regulatory protein. It is currently unclear which PP1β holoenzyme, or holoenzymes, mediates the synaptic functions defined in this study, as the neuronal regulatory proteins that bind PP1β specifically have not been characterized. PP1β may directly dephosphorylate GluA1, or act on other relevant regulatory proteins. For example, PP1β cKO may increase GluA1 S831 phosphorylation by promoting CaMKII and PKC activity. We did not observe a change in CaMKII T286 phosphorylation in PP1β cKO mice, but we cannot exclude the possibility that other phosphorylation sites are affected. Similarly, PP1β may directly dephosphorylate GluN2A, a GluN2A scaffolding protein, or another regulatory protein. Several kinases can affect GluN2A surface expression and recycling, including GSK3β, DYRK1A, PKC, CaMKII, and CDK5, either through direct action on GluN2A or its scaffolding proteins (Gardoni et al., 2001; Gardoni et al., 2003; Grau et al., 2014; Li et al., 2001; Mauceri et al., 2007; Monaco et al., 2018; Mota Vieira et al., 2020; Yong et al., 2021). If the PP1β cKO impairments are mediated by unopposed

activity of a kinase, it may be possible to rescue the deficits via inhibition of the corresponding kinase.

Although the identities and functions of PP1β holoenzymes in the central nervous system are largely unstudied, MYPT1-PP1β is an attractive candidate holoenzyme. MYPT1-PP1β, also known as myosin phosphatase, regulates the activity of myosin II proteins, including non-muscle myosin II, via inhibitory dephosphorylation of the myosin light chain (MLC) regulatory subunit (Ito et al., 2004). MLC has been implicated in spine maturation, LTP, as well as NMDAR trafficking (Amparan et al., 2005; Bajaj et al., 2009; Hodges et al., 2011; Rex et al., 2010). Given that PP1β opposes classic PP1 function, identifying neuronal PP1β holoenzymes and elucidating their functions presents an exciting avenue for further research.

Human de novo mutations in PP1β cause intellectual disability (Gripp *et al.*, 2016; Hamdan *et al.*, 2014; Huckstadt *et al.*, 2021; Lin *et al.*, 2018; Ma *et al.*, 2016; Umeki *et al.*, 2019; Zambrano *et al.*, 2017; Zhou *et al.*, 2020). The clinical profile of patients with these mutations resembles a Rasopathy, specifically Noonan syndrome with loose anagen hair. Consistent with this classification, PP1β mutations have been reported to promote Ras signaling (Young et al., 2018). We did not observe a change in pERK, which is downstream of Ras signaling, suggesting PP1β activity may not be important for basal Ras-MEK-ERK signaling and supporting the notion that PP1β mutations represent a gain-of-function. On the other hand, PP1β may be important for activity-induced Ras-MEK-ERK signaling, presenting another means by which PP1β cKO could impair LTP. Further investigation of the neuronal functions of PP1β will improve our understanding of the pathophysiology of these mutations and help generate potential therapeutic targets.

Acknowledgments

This work was supported by the National Institutes of Health (NIH) NIH F30 MH122046 to K.F., NIH DA10044 to A.C.N., R01 MH109719 and National Science Foundation (NSF) IOS-1457336 to H.X.

Author contributions:
Conceptualization: KF, HX
Methodology: KF, CM, DB, NW, HX
Investigation: KF, NW, YZ
Validation: AG, AM
Visualization: KF
Resources: ACN, HX
Supervision: HX
Writing—original draft: KF, HX
Writing—review & editing: CAN

Declaration of interests
The authors declare no competing interests.

Figure Legends

Fig. 1. PP1 isoforms play opposing roles in regulating hippocampal synaptic transmission.
(**B-E**) (**A**) CA1-specific Cre expression shown by breeding with Ai9 mice, which express tdTomato following Cre-recombination. Field recordings of Sch-CA1 synapses were conducted to assess input-output (IO) curves. (**B**) PP1α cKO does not significantly alter synaptic transmission compared to control littermates (two-way ANOVA, genotype: p=0.6504; Regression: Extra sum-of-squares F-test (F-test), slope: p=0.7769; N=4 (control) and 3 (cKO) mice, n=9 and 10 slices), whereas (**C**) PP1γ cKO significantly decreases transmission (two-way ANOVA, genotype: p<0.01; F-test, slope: p<0.001; N=5 and 3, n=9 and 8) and (**D**) PP1β cKO significantly increases transmission (two-way ANOVA, genotype: p<0.01; F-test, slope: p<0.0001; N=4 and 4, n=11 and 14). (**E**) PP1α/γ cKO decreases transmission (two-way ANOVA, genotype: p=<0.0001; F-test, slope: p<0.0001; N=9 and 10, n=20 and 15). (**F**) Comparison of changes in IO curves for PP1 isoform cKO mice normalized to control littermates. PP1α/γ cKO decreases transmission to a greater degree than PP1γ cKO (two-way ANOVA, genotype: p<0.001; F-test, slope: p<0.0001).

Fig. 2. PP1 isoforms play opposing roles in regulating hippocampal synaptic plasticity. (**A**) PP1β cKO attenuates LTP induced by 100 pulses at 100 Hz stimulation (two-tailed t-test: p<0.001, N=3 (control) and 4 (cKO) mice, n=5 and 8 slices). (**B**) PP1α/γ cKO promotes sub-threshold LTP induced by 25 pulses at 100 Hz stimulation (two-tailed t-test: p<0.05, N=3 and 3, n=5 and 6). (**C**) PP1α/γ cKO does not alter LTP induced by 100 pulses at 100 Hz stimulation (two-tailed t-test: p=0.30, N=3 and 4, n=4 and 5). (**D**) Examples of whole-cell recording LTP experiments from control (left) and PP1β cKO (right). LTP was delivered by pairing 100 pulses at 100 Hz at 0 mV holding potential (x-axis dotted line) before 10 min of recording to avoid washout. Y-axis dotted line denotes average baseline response. (**E**) PP1β cKO attenuates LTP induced via the pairing protocol (two-tailed t-test: p<0.001, N=3 (control) and 5 (cKO) mice, n=5 and 6 cells). LTP magnitude was measured as the average fEPSP or EPSC 25-30 min after LTP induction normalized to the baseline.

Fig. 3. PP1β cKO impairs spatial learning and memory. (**A**) Overview of the allocentric place-object task, a novel object recognition variant sensitive to hippocampal deficits. In the test phase, the novelty of the object is based on its location in the arena. (**B**) PP1β cKO are impaired in novel object recognition compared to their control littermates (two-tailed t-test: p<0.05, N=5 (control) and 6 (cKO) mice, 3 litters). There was no difference in (**C**) total time exploring objects between groups (two-tailed t-test: p=0.98, N=5 and 6). (**D**) Locomotor activity (two-tailed t-test: p=0.82, N=8 and 7), or (**E**) anxiety (two-tailed t-test: p=0.66, N=8 and 7), as measured by time spent in the center during an open field test. (**F**) There was no difference in novel object recognition between PP1α/γ cKO and control littermates (two-tailed t-test: p=0.79, N=6 and 8, 4 litters). PP1α/γ cKO mice showed no difference in (**G**) total time exploring objects between groups (two-tailed t-test: p=0.81, N=6 and 8), (**H**) locomotor activity (two-tailed t-test: p=0.97, N=8 and 10), or (**I**) anxiety (two-tailed t-test: p=0.98, N=8 and 10). Locomotor activity was measured as cumulative distance traveled at 1 hour. Anxiety was measured by time spent in the center of the open field arena over 10 minutes. The standard error of the mean is displayed as error bars or shaded lines.

Fig. 4. PP1β cKO increases GluA1 phosphorylation, spine density, and spine maturation.
(**A-D**) Western blot results from CA1 microdissections of PP1β cKO mice and littermate
controls. 3 mice per group. Phosphorylation was normalized to total protein, and total protein
was normalized to β-tubulin. (**A**) Signaling pathways relevant to synaptic plasticity. No
significant differences were observed. (**B**) AMPAR phosphorylation and total protein levels.
S831 GluA1 phosphorylation was significantly increased in PP1β cKO (two-tailed t-test:
$p<0.05$). (**C**) NMDAR phosphorylation and total protein levels. Total GluN2A levels were
significantly decreased (two-tailed t-test: $p<0.01$). (**D**) Quantification of comparisons. Following
normalization to total protein or β-tubulin, intensities were normalized to control samples. (**E-G**)
Results from mEPSC recordings from CA1 pyramidal neurons. (**E**) PP1β cKO has no effect on
mean mEPSC amplitude (two-tailed t-test: $p=0.31$, N=7 (control) and 6 (cKO) mice, n=26 and
31) but alters the amplitude distribution (Kolmogorov-Smirnov: $p<0.0001$, 1000 randomly
sampled events per group). (**F**) PP1β cKO significantly increases mEPSC frequency (two-tailed
t-test: $p<0.01$, N=7 and 6, n=26 and 31) and correspondingly shifts the frequency distribution
(Kolmogorov-Smirnov: $p<0.0001$, 1000 randomly sampled events per group). (**G**) Example
mEPSC traces from control (top) and a PP1β cKO (bottom). (**H-J**) Spine number and
morphology of CA1 pyramidal neurons. (**H**) Examples of dendritic spines in biocytin-filled CA1
pyramidal cells. (**I**) PP1β cKO significantly increases spine density (two-tailed t-test: $p<0.05$,
N=4 (control) and 3 (cKO) mice, n=23 and 19 dendrites). (**J**) PP1β cKO alters spine morphology
(two-way ANOVA, genotype: $p<0.05$), significantly increasing mushroom spines (Šidák,
$p<0.001$), as categorized by Imaris reconstruction. (**K**) Examples of minimal stimulation
experiments from control (top) and PP1β cKO (bottom) mice. Failures are shown in grey,
successful responses in black. Insets show waveforms from averaged responses at -70 mV (top
left) and +40 mV (bottom right) holding potential, scale bars 5 pA, 10 ms. Y-axis dotted lines
denote 5 pA thresholds used for initial classification. X-axis dotted line denotes the switch to
+40 mV holding potential. All events were manually reviewed. (**L**) PP1β cKO decreases the
proportion of silent synapses in CA1 pyramidal cells (two-tailed t-test: $p<0.05$, N=4 and 7, n=8
and 10 cells).

**Fig. S1. PP1β cKO mEPSC effects are independent of neurotransmitter release and acute
hippocampal slice preparation methodology.**
(**A**) PP1β cKO has no effect on paired-pulse facilitation (two-way ANOVA: $p=0.39$, N=3
(control) and 3 (cKO) mice, n=10 and 10 slices). (**B-C**) mEPSC results from slices prepared in
NMDG-based cutting solution. PP1β cKO has no effect on mEPSC amplitude (two-tailed t-test:
$p=0.25$, N=3 and 3, n=15 and 15 cells) but increases mEPSC frequency (two-tailed t-test:
$p<0.05$, N=3 and 3, n=15 and 15). (**D-E**) mEPSC results from slices prepared in sucrose-based
cutting solution. PP1β cKO has no effect on mEPSC amplitude (two-tailed t-test: $p=0.82$, N=3
and 3, n=11 and 16) but increases mEPSC frequency (two-tailed t-test: $p<0.01$, N=3 and 3, n=11
and 16).

377 **STAR Methods**
378 *Key resources table*

Reagent or Resource	Source	Identifier
Antibodies		
β-tubulin	Cell Signaling Technology	5666
CaMKII pT286	Cell Signaling Technology	12716
CaMKII	EMD Millipore	04-1079
CREB pS133	Cell Signaling Technology	9191
CREB	Cell Signaling Technology	9104
ERK	Cell Signaling Technology	9102
GluA1 pS831	Sigma	A4352
GluA1 pS845	EMD Millipore	AB5849
GluA1	EMD Millipore	AB1504
GluA2 pS880	EMD Millipore	07-294-I
GluA2	EMD Millipore	MAB397
GluA2/3	Upstate	07-598
GluN1	Invitrogen	32-0500
GluN2B pS1480	ThermoFisher	PA1-4733
GluN2B	Santa Cruz Biotechnology	1469
GluN2A	Santa Cruz Biotechnology	1468
Chemicals, Peptides, and Recombinant Proteins		
Picrotoxin	Tocris	1128
Tetrodotoxin	Tocris	1069
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Jackson Laboratory	JAX: 000664
Mouse: CaMKIIα-Cre (T29-1)	Jackson Laboratory	JAX: 005359
Mouse: Ai9 reporter	Jackson Laboratory	JAX:007909
Mouse: PP1α cKO	Liu et al. (2015)	NA
Mouse: PP1β cKO	Liu *et al.* (2015)	NA
Mouse: PP1γ cKO	Liu *et al.* (2015)	NA
Software and Algorithms		
AzureSpot Pro (2.0.062)	Azure Biosystems	https://azurebiosystems.com/software/azurespotpro/
Prism (9.3.1)	GraphPad	https://www.graphpad.com/
R (4.0.2)	R Foundation for Statistical Computing	https://www.r-project.org/
Behavioral Observation Research Interactive Software (BORIS) (7.9.21)	Friard et al. (2016)	https://www.boris.unito.it/
Recording Artist Package	Rick Gerkin	http://github.com/rgerkin/recording-artist
Igor Pro 8	Wavemetrics	https://www.wavemetrics.com/Products/igorpro/igorpro.htm
Imaris (9.5.0)	Oxford Instruments	https://imaris.oxinst.com/

379
380
Mice
CaMKIIα-Cre (T29-1; JAX 005359) were purchased from Jackson laboratory. Floxed PP1 isoform mice
were generated as described previously (Liu *et al.*, 2015). Ai9 mice (JAX 007909) were a gift from A.

Majewska at the University of Rochester. Mice were maintained by breeding with wild-type C57BL/6J
(JAX 000664). Animals were group-housed (2-5 mice per cage) on a 12-hour light/dark cycle with food
and water ad libitum. For all experiments, control littermates (PP1(fl/fl)) were compared to KO
littermates (PP1(fl/fl); CaMKIIα-Cre). All animal procedures were conducted in accordance with the
National Institutes of Health Guide for the Care and Use of Laboratory Animals and with protocols
approved by the University of Rochester Institutional Animal Care and Use Committee.
390
Acute hippocampal slice preparation
Acute hippocampal slices were prepared from 1-1.5 month-old mice of both sexes as described previously
(Foley *et al.*, 2022; Foley et al., 2021b). After brief isoflurane anesthetization and decapitation, the brain
was rapidly dissected and sectioned in an ice-cold high sucrose solution containing, in mM: 215 sucrose,
2.5 KCl, 26 $NaHCO_3$, 1.6 NaH_2PO_4, 20 D-glucose, 1 $CaCl_2$, 4 $MgCl_2$, and 4 $MgSO_4$. 400 μm thick
coronal (field recordings) or 300 μm thick transverse (whole-cell recordings) slices were sectioned on a
Leica VT1200S vibratome with carbon steel blades (Electron Microscopy Sciences). After sectioning,
slices were recovered in artificial cerebrospinal fluid (ACSF) at 34°C for 30 min, followed by an
additional 1-hour incubation at room temperature (RT). ACSF contained, in mM: 124 NaCl, 2.5 KCl, 1.0
NaH_2PO_4, 26 $NaHCO_3$, 10 D-glucose, 1.3 $MgSO_4$, and 2.5 $CaCl_2$ for field recordings; and 124 NaCl, 2.5
KCl, 1.2 NaH_2PO_4, 24 $NaHCO_3$, 5 HEPES, 12.5 D-glucose, 2 $MgSO_4$, and 2 $CaCl_2$ for whole-cell
recordings. For a subset of mEPSC recordings (Figure S1B,C), slices were prepared in NMDG-based
ACSF containing, in mM: 92 NMDG, 2.5 KCl, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 D-glucose, 5
Na ascorbate, 2 thiourea, 3 Na pyruvate, 10 $MgSO_4$, and 0.5 $CaCl_2$. Solutions were continuously aerated
with carbogen (95% O_2, 5% CO_2).
406
Electrophysiology
Field recordings in acute hippocampal slices were performed at Sch-CA1 synapses in RT ACSF at a flow
rate of 2-3 ml/min. Recording (~1 MΩ) and stimulating (<1MΩ) electrodes were pulled from borosilicate
glass and filled with ACSF. The recording electrode was placed in CA1 stratum radiatum, 200 μm medial
to the stimulating electrode, with both electrodes approximately 100 μm below the slice surface.
Responses were elicited every 15 sec. Input-output (IO) curves were measured by comparing the fiber
volley amplitude to the fEPSP slope. Paired-pulse facilitation (PPF) ratios were measured by comparing
the second fEPSP amplitude to the first. Long-term potentiation (LTP) was induced via 100 pulses, or 25
pulses (sub-threshold LTP), at 100 Hz after at least 10 min stable baseline recording.
416
Whole-cell recordings were performed on CA1 pyramidal neurons in ACSF held at 28°C (LTP; minimal
stimulation) or 32°C (mEPSCs). Picrotoxin (50μM) was included in the ACSF for LTP and minimal
stimulation, and picrotoxin (100μM) and tetrodotoxin (1μM) for mESPC recordings. 3-5 MΩ recording
electrodes were pulled from borosilicate glass and filled with, in mM: 140 $CsMeSO_3$, 8 CsCl, 7 Na_2
phosphocreatine, 0.25 EGTA, 10 HEPES, 0.1 spermine, 2 Mg ATP, 0.3 Na GTP, at 7.32-7.36 pH and
294-298 mOsm for LTP and minimal stimulation, and 115 $CsMeSO_3$, 20 CsCl, 10 Na_2 phosphocreatine,
0.6 EGTA, 10 HEPES, 2.5 $MgCl_2$, 5 QX-314-Br, 4 Mg ATP, 0.3 Na GTP, at 7.32-7.36 pH and 294-298
mOsm for mEPSC recordings. Stimulation electrodes were filled with ACSF. Responses were elicited
every 10 sec and access and input resistance were continuously monitored. Cells were excluded from
analysis if access resistance varied by more than 25%. CA1 pyramidal cells were identified via location
(stratum pyramidale) and morphology under differential interference contrast microscopy with a 60x
water-immersion objective on an Olympus BX51WI. For whole-cell LTP experiments, cells were
depolarized to 0 mV during stimulation with 100 pulses at 100 Hz. For minimal stimulation experiments,
stimulation intensity was reduced until ~50% of stimulations failed to elicit an AMPAR-mediated
response at a holding potential of -70 mV. We then measured the composite AMPAR- and NMDAR-
mediated response by holding the cell at +40 mV. The percentage of silent synapses was calculated using
the equation $1-ln(F_{-70})/ln(F_{+40})$, where F_{-70} is the failure rate at -70 mV holding potential and F_{+40} is the
failure rate at +40 mV holding potential (Isaac et al., 1995; Liao et al., 1995). The stochastic nature of the

release probability creates variability in the failure rate which can lead to mathematically negative values in the calculation of silent synapses (Favaro et al., 2018). Nevertheless, we excluded four PP1β cKO neurons which had calculated silent synapses proportions of <-0.15. No PP1β control neurons were excluded from the minimal stimulation analysis. For biocytin-fill experiments, neurons were held in whole-cell configuration for at least 10 min.

For all electrophysiology experiments, stimulation intensity was controlled by an ISO-Flex stimulus isolator (AMPI) and timing controlled by a Master-8 pulse stimulator (AMPI). Data were collected with a MultiClamp 700 series amplifier (Axon Instruments), PCI-6221 data acquisition device (National Instruments), and Igor Pro 7 (Wavemetrics) with a customized software package (Recording Artist, http://github.com/rgerkin/recording-artist).

Spine density and morphology
Biocytin-filled neurons were fixed in 4% paraformaldehyde (PFA) for 24-48 hours and stained with conjugated streptavidin (Vector Laboratories) for 48 hours. Slices were then mounted with Prolong Diamond Antifade (Invitrogen) mounting media and allowed to cure for at least 24 hours. Confocal images of secondary dendrites within 200 μm of the cell body were acquired on an Olympus FV1000MP microscope system controlled by Fluoview software using a 60x oil immersion objective (Olympus UPlanSApo, 1.35 NA). Images were acquired at a resolution of 1024 x 1024 pixels at 3x zoom. For quantification of spine density and morphology, z-stacks of biocytin-filled neurons were deconvolved and reconstructed in 3D in Imaris (v9.3, Bitplane). Dendrites were reconstructed using semi-automated filament tracing in autopath mode. Spines were then identified using automatic seed points and manually verified. Spine identification parameters were set at 3.5 μm maximum spine length, 0.138 μm minimum spine head diameter, and branched spines were allowed. Spine morphology was classified in Imaris using the classify spines wizard algorithm. The general criteria for classification were: (i) Stubby spines have a large head without an appreciable neck; (ii) Mushroom spines have a large head greater than the width of the neck; (iii) Thin spines have a large head greater than the width of the neck, but smaller than mushroom spines; and (iv) Filopodia are long and lack an appreciable spine head. These criteria were achieved algorithmically with the following parameters in order, in μm: (i) Stubby: $\min(\text{neck}_{width}) > 0.3$ & $\text{length}(\text{spine}) < 0.5$ & $\max(\text{head}_{width}) < \max(\text{neck}_{width})$ | neck not detected; (ii) Mushroom: $\max(\text{head}_{width}) > \min(\text{neck}_{width})*2$ & neck detected | $\max(\text{head}_{width}) > 0.38$; (iii) Thin: $\text{length}(\text{neck}) > \text{mean}(\text{neck}_{width})$ & $\max(\text{head}_{width}) < \text{length}(\text{neck})$ | $\max(\text{head}_{width}) > 0.14$; and (iv) Filopodia: remaining spines. Note that spine morphology classification in Imaris is obligatorily ordered, such that spines that are classified as Stubby cannot be classified as Mushroom, and so on. Spine morphology was manually verified.

SDS-PAGE and immunoblotting
CA1 microdissection was performed in ice-cold PBS after rapid extraction of the brain from 1-1.5 month-old mice anesthetized with isoflurane. Dissected tissue was frozen in liquid nitrogen and stored at -80°C until lysate preparation. Total protein lysates were prepared by homogenizing samples in RIPA buffer, containing 50 mM Tris-HCl pH 8.0, 1% NP-40, 0.5% Na deoxycholate, 0.1% SDS, 150 mM NaCl, 1 mM EDTA, 10 mM NaF, protease inhibitors (Pierce, A32965), and phosphatase inhibitor cocktails (Santa Cruz, 45055). Samples were agitated for 40 min at 4°C and centrifuged at 6,000 g for 20 min. The soluble fraction was used for protein concentration quantification (Pierce BCA) and then boiled in Laemmli buffer. Equal amounts of protein per sample were loaded and resolved by SDS-PAGE on 8-15% polyacrylamide gels and transferred to polyvinylidene fluoride membrane. Membranes were blocked with 5% bovine serum albumin or 5% non-fat dried milk in Tris-buffered saline with 0.1% Tween 20 (TBS-T) and washed with TBS-T. After primary antibody incubation (4°C), washing, secondary antibody incubation (RT), and final washing, immunoreactive signals were visualized by chemiluminescence on an Azure Biosystems Imaging System. Densitometry analysis was performed on AzureSpot Pro software (Azure Biosystems). The following antibodies were used: β-tubulin (Cell Signaling Technology (CST), 5666), CaMKII pT286 (CST, 12716), CaMKII (Millipore, 04-1079), CREB pS133 (CST, 9191), CREB

486   (CST, 9104), ERK (CST, 9102), GluA1 pS831 (Sigma, A4352), GluA1 pS845 (Millipore, AB5849),
487   GluA1 (Millipore, AB1504), GluA2 pS880 (Millipore, 07-294-I), GluA2 (Millipore, MAB397), GluA2/3
488   (Upstate, 07-598), GluN1 (Invitrogen, 32-0500), GluN2B pS1480 (PA1-4733), GluN2B (Santa Cruz,
489   1469), GluN2A (Santa Cruz, 1468).
490
Behavioral procedures
Behavioral experiments were performed on 1-1.5 month-old male mice, using approximately equal
numbers of control and KO littermates from each litter. Three to four litters were used per mouse line. All
experiments were carried out following 2-3 days of handling (Gouveia and Hurst, 2013) in the light phase
(8 am-12 pm). Testing chambers and objects were thoroughly cleaned with 70% ethanol after each
experiment. Open-field testing was performed for 1 hour in a chamber equipped with infrared photocells
(Opto-Varimex-5, Columbus Instruments). Mouse position was automatically detected via beam
interruption.
499
The allocentric object-place assay was performed following standard novel-object recognition procedures
(Langston and Wood, 2009; Leger et al., 2013). The outer walls of the testing chamber were covered with
four different colors or patterns for spatial cues. Sessions were split into three phases of 10 min each:
habituation, sample, and test, with 5-10 min between phases to allow for thorough cleaning of the
chambers. Three copies of the objects ($A_{1,2,3}$, $B_{1,2,3}$) were used to avoid any lingering cues between
phases. In the habituation phase, mice were placed into the chamber facing the southern wall and allowed
to explore the chamber in the absence of any objects. In the sample phase, two different objects (A_1-B_1)
were placed in the chamber and mice were again placed facing the southern wall. In the test phase, two
copies of one of the two sampling phase objects (A_2-A_3; B_2-B_3) were placed in the chamber and mice
were placed into the chamber facing the eastern or western wall. Therefore, in the test phase, one of the
objects was presented in the same location as in the sample phase (familiar; e.g. A_2 at A_1's position),
whereas the other object occupied a different location than in the sample phase (novel; e.g. A_3 at B_1's
position). Testing phase object pair (A or B) and site of entry (eastern/western wall) were
counterbalanced. The objects used were 50 mL conical tubes (inverted) and 70 mL culture flasks filled
with white or black sand. Exploration was defined as the mouse being within 2.5 cm of the object and
directing its nose at the object while sniffing or whisking. Sessions were monitored via overhead video
recording and analyzed in BORIS (Friard *et al.*, 2016).
517
Statistical analyses
GraphPad Prism (9.3.1) was used for statistical analyses and numerical data visualization. Statistical
significance between means was calculated using unpaired, two-tailed t-tests or ANOVAs. Two-way
ANOVAs were used for IO and PPF comparisons. IO curves were also fit with least squares regression
curves without weighting and the slopes of the regressions were compared via extra sum-of-squares F-
tests. LTP magnitude was quantified as the change in fEPSP slope at 25-30 min post-induction compared
to the 5 (whole-cell) or 10 (field) minute baseline. Tukey and Šidák post-hoc comparisons were
performed for one- and two-way ANOVAs, respectively. The arithmetic mean and standard error of the
mean are displayed in all figures unless otherwise noted.

References

Amparan, D., Avram, D., Thomas, C.G., Lindahl, M.G., Yang, J., Bajaj, G., and Ishmael, J.E. (2005). Direct interaction of myosin regulatory light chain with the NMDA receptor. J Neurochem *92*, 349-361. 10.1111/j.1471-4159.2004.02869.x.

Aow, J., Dore, K., and Malinow, R. (2015). Conformational signaling required for synaptic plasticity by the NMDA receptor complex. Proc Natl Acad Sci U S A *112*, 14711-14716. 10.1073/pnas.1520029112.

Bajaj, G., Zhang, Y., Schimerlik, M.I., Hau, A.M., Yang, J., Filtz, T.M., Kioussi, C., and Ishmael, J.E. (2009). N-methyl-D-aspartate receptor subunits are non-myosin targets of myosin regulatory light chain. J Biol Chem *284*, 1252-1266. 10.1074/jbc.M801861200.

Bayes, A., Collins, M.O., Croning, M.D., van de Lagemaat, L.N., Choudhary, J.S., and Grant, S.G. (2012). Comparative study of human and mouse postsynaptic proteomes finds high compositional conservation and abundance differences for key synaptic proteins. PLoS One *7*, e46683. 10.1371/journal.pone.0046683.

Bayes, A., van de Lagemaat, L.N., Collins, M.O., Croning, M.D., Whittle, I.R., Choudhary, J.S., and Grant, S.G. (2011). Characterization of the proteome, diseases and evolution of the human postsynaptic density. Nat Neurosci *14*, 19-21. 10.1038/nn.2719.

Bollen, M., Peti, W., Ragusa, M.J., and Beullens, M. (2010). The extended PP1 toolkit: designed to create specificity. Trends Biochem Sci *35*, 450-458. 10.1016/j.tibs.2010.03.002.

Bordelon, J.R., Smith, Y., Nairn, A.C., Colbran, R.J., Greengard, P., and Muly, E.C. (2005). Differential localization of protein phosphatase-1alpha, beta and gamma1 isoforms in primate prefrontal cortex. Cereb Cortex *15*, 1928-1937. 10.1093/cercor/bhi070.

Burger, C., Lopez, M.C., Feller, J.A., Baker, H.V., Muzyczka, N., and Mandel, R.J. (2007). Changes in transcription within the CA1 field of the hippocampus are associated with age-related spatial learning impairments. Neurobiol Learn Mem *87*, 21-41. 10.1016/j.nlm.2006.05.003.

Carmody, L.C., Bauman, P.A., Bass, M.A., Mavila, N., Depaoli-Roach, A.A., and Colbran, R.J. (2004). A protein phosphatase-1gamma1 isoform selectivity determinant in dendritic spine-associated neurabin. J Biol Chem *279*, 21714-23. 10.1074/jbc.M402261200.

Chiu, A.M., Wang, J., Fiske, M.P., Hubalkova, P., Barse, L., Gray, J.A., and Sanz-Clemente, A. (2019). NMDAR-Activated PP1 Dephosphorylates GluN2B to Modulate NMDAR Synaptic Content. Cell Rep *28*, 332-341 e335. 10.1016/j.celrep.2019.06.030.

Derkach, V., Barria, A., and Soderling, T.R. (1999). Ca2+/calmodulin-kinase II enhances channel conductance of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate type glutamate receptors. Proc Natl Acad Sci U S A *96*, 3269-3274. 10.1073/pnas.96.6.3269.

Dickinson, M.E., Flenniken, A.M., Ji, X., Teboul, L., Wong, M.D., White, J.K., Meehan, T.F., Weninger, W.J., Westerberg, H., Adissu, H., et al. (2016). High-throughput discovery of novel developmental phenotypes. Nature *537*, 508-514. 10.1038/nature19356.

Diering, G.H., Heo, S., Hussain, N.K., Liu, B., and Huganir, R.L. (2016). Extensive phosphorylation of AMPA receptors in neurons. Proc Natl Acad Sci U S A *113*, E4920-4927. 10.1073/pnas.1610631113.

Diering, G.H., and Huganir, R.L. (2018). The AMPA Receptor Code of Synaptic Plasticity. Neuron *100*, 314-329. 10.1016/j.neuron.2018.10.018.

Favaro, P.D., Huang, X., Hosang, L., Stodieck, S., Cui, L., Liu, Y.Z., Engelhardt, K.A., Schmitz, F., Dong, Y., Lowel, S., and Schluter, O.M. (2018). An opposing function of paralogs in balancing developmental synapse maturation. Plos Biol *16*, e2006838. 10.1371/journal.pbio.2006838.

Foley, K., Altimimi, H., Hou, H., Zhang, Y., McKee, C., Papasergi-Scott, M.M., Yang, H.,
Mayer, A., Ward, N., MacLean, D.M., et al. (2022). PP1 inhibitor-2 promotes PP1γ positive
regulation of synaptic transmission. Biorxiv, 2022.2002.2010.480004.
10.1101/2022.02.10.480004.
Foley, K., McKee, C., Nairn, A.C., and Xia, H. (2021a). Regulation of Synaptic Transmission
and Plasticity by Protein Phosphatase 1. J Neurosci *41*, 3040-3050. 10.1523/JNEUROSCI.2026-
20.2021.
Foley, K.F.W., Barnett, D., Cory-Slechta, D.A., and Xia, H. (2021b). Early Low-Level Arsenic
Exposure Impacts Post-Synaptic Hippocampal Function in Juvenile Mice. Toxics *9*.
10.3390/toxics9090206.
Friard, O., Gamba, M., and Fitzjohn, R. (2016). BORIS: a free, versatile open source
event logging software for video/audio coding and live observations. Methods Ecol Evol *7*,
1325-1330. 10.1111/2041-210x.12584.
Gao, J., Hu, X.D., Yang, H., and Xia, H. (2018). Distinct Roles of Protein Phosphatase 1 Bound
on Neurabin and Spinophilin and Its Regulation in AMPA Receptor Trafficking and LTD
Induction. Mol Neurobiol *55*, 7179-7186. 10.1007/s12035-018-0886-2.
Gardoni, F., Bellone, C., Cattabeni, F., and Di Luca, M. (2001). Protein kinase C activation
modulates alpha-calmodulin kinase II binding to NR2A subunit of N-methyl-D-aspartate
receptor complex. J Biol Chem *276*, 7609-7613. 10.1074/jbc.M009922200.
Gardoni, F., Mauceri, D., Fiorentini, C., Bellone, C., Missale, C., Cattabeni, F., and Di Luca, M.
(2003). CaMKII-dependent phosphorylation regulates SAP97/NR2A interaction. J Biol Chem
278, 44745-44752. 10.1074/jbc.M303576200.
Genoux, D., Haditsch, U., Knobloch, M., Michalon, A., Storm, D., and Mansuy, I.M. (2002).
Protein phosphatase 1 is a molecular constraint on learning and memory. Nature *418*, 970-975.
10.1038/nature00928 PMID - 12198546.
Gouveia, K., and Hurst, J.L. (2013). Reducing mouse anxiety during handling: effect of
experience with handling tunnels. PLoS One *8*, e66401. 10.1371/journal.pone.0066401.
Granger, A.J., and Nicoll, R.A. (2014). Expression mechanisms underlying long-term
potentiation: a postsynaptic view, 10 years on. Philos Trans R Soc Lond B Biol Sci *369*,
20130136. 10.1098/rstb.2013.0136.
Grau, C., Arato, K., Fernandez-Fernandez, J.M., Valderrama, A., Sindreu, C., Fillat, C., Ferrer,
I., de la Luna, S., and Altafaj, X. (2014). DYRK1A-mediated phosphorylation of GluN2A at
Ser(1048) regulates the surface expression and channel activity of GluN1/GluN2A receptors.
Front Cell Neurosci *8*, 331. 10.3389/fncel.2014.00331.
Gripp, K.W., Aldinger, K.A., Bennett, J.T., Baker, L., Tusi, J., Powell-Hamilton, N., Stabley, D.,
Sol-Church, K., Timms, A.E., and Dobyns, W.B. (2016). A novel rasopathy caused by recurrent
de novo missense mutations in PPP1CB closely resembles Noonan syndrome with loose anagen
hair. Am J Med Genet A *170*, 2237-2247. 10.1002/ajmg.a.37781.
Haege, S., Galetzka, D., Zechner, U., Haaf, T., Gamerdinger, M., Behl, C., Hiemke, C., and
Schmitt, U. (2010). Spatial learning and expression patterns of PP1 mRNA in mouse
hippocampus. Neuropsychobiology *61*, 188-196. 10.1159/000297736.
Hamdan, F.F., Srour, M., Capo-Chichi, J.M., Daoud, H., Nassif, C., Patry, L., Massicotte, C.,
Ambalavanan, A., Spiegelman, D., Diallo, O., et al. (2014). De novo mutations in moderate or
severe intellectual disability. Plos Genet *10*, e1004772. 10.1371/journal.pgen.1004772.
Hayashi, Y., Shi, S.-H., Esteban, J.A., Piccini, A., Poncer, J.-C., and Malinow, R. (2000).
Driving AMPA Receptors into Synapses by LTP and CaMKII: Requirement for GluR1 and PDZ

Domain Interaction. Science *287*, 2262-2267. 10.1126/science.287.5461.2262 PMID -
10731148.
Heroes, E., Lesage, B., Görnemann, J., Beullens, M., Meervelt, L., and Bollen, M. (2013). The
PP1 binding code: a molecular☐lego strategy that governs specificity. FEBS Journal *280*, 584-
595. 10.1111/j.1742-4658.2012.08547.x PMID - 22360570.
Hodges, J.L., Newell-Litwa, K., Asmussen, H., Vicente-Manzanares, M., and Horwitz, A.R.
(2011). Myosin IIb activity and phosphorylation status determines dendritic spine and post-
synaptic density morphology. PLoS One *6*, e24149. 10.1371/journal.pone.0024149.
Hu, X.D., Huang, Q., Roadcap, D.W., Shenolikar, S.S., and Xia, H. (2006). Actin-associated
neurabin-protein phosphatase-1 complex regulates hippocampal plasticity. J Neurochem *98*,
1841-1851. 10.1111/j.1471-4159.2006.04070.x.
Hu, X.D., Huang, Q., Yang, X., and Xia, H. (2007). Differential regulation of AMPA receptor
trafficking by neurabin-targeted synaptic protein phosphatase-1 in synaptic transmission and
long-term depression in hippocampus. J Neurosci *27*, 4674-4686. 10.1523/JNEUROSCI.5365-
06.2007.
Huckstadt, V., Chinton, J., Gomez, A., Obregon, M.G., and Gravina, L.P. (2021). Noonan
syndrome with loose anagen hair with variants in the PPP1CB gene: First familial case reported.
Am J Med Genet A *185*, 1256-1260. 10.1002/ajmg.a.62089.
International Mouse Phenotyping, C. (2022). www.mousephenotype.org.
Isaac, J.T., Nicoll, R.A., and Malenka, R.C. (1995). Evidence for silent synapses: implications
for the expression of LTP. Neuron *15*, 427-434. 10.1016/0896-6273(95)90046-2.
Ito, M., Nakano, T., Erdodi, F., and Hartshorne, D.J. (2004). Myosin phosphatase: structure,
regulation and function. Mol Cell Biochem *259*, 197-209.
10.1023/b:mcbi.0000021373.14288.00.
Jouvenceau, A., Hedou, G., Potier, B., Kollen, M., Dutar, P., and Mansuy, I.M. (2006). Partial
inhibition of PP1 alters bidirectional synaptic plasticity in the hippocampus. Eur J Neurosci *24*,
564-572. 10.1111/j.1460-9568.2006.04938.x.
Kiyama, Y., Manabe, T., Sakimura, K., Kawakami, F., Mori, H., and Mishina, M. (1998).
Increased thresholds for long-term potentiation and contextual learning in mice lacking the
NMDA-type glutamate receptor epsilon1 subunit. J Neurosci *18*, 6704-6712.
Langston, R.F., and Wood, E.R. (2009). Associative recognition and the hippocampus:
Differential effects of hippocampal lesions on object-place, object-context and object-place-
context memory. Hippocampus *20*, 1139-1153. 10.1002/hipo.20714 PMID - 19847786.
Lee, H.K., Barbarosie, M., Kameyama, K., Bear, M.F., and Huganir, R.L. (2000). Regulation of
distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. Nature
405, 955-959. 10.1038/35016089.
Lee, H.K., Kameyama, K., Huganir, R.L., and Bear, M.F. (1998). NMDA induces long-term
synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in
hippocampus. Neuron *21*, 1151-1162. 10.1016/s0896-6273(00)80632-7.
Leger, M., Quiedeville, A., Bouet, V., Haelewyn, B., Boulouard, M., Schumann-Bard, P., and
Freret, T. (2013). Object recognition test in mice. Nat Protoc *8*, 2531-2537.
10.1038/nprot.2013.155.
Li, B.S., Sun, M.K., Zhang, L., Takahashi, S., Ma, W., Vinade, L., Kulkarni, A.B., Brady, R.O.,
and Pant, H.C. (2001). Regulation of NMDA receptors by cyclin-dependent kinase-5. Proc Natl
Acad Sci U S A *98*, 12742-12747. 10.1073/pnas.211428098.

Li, J., Wilkinson, B., Clementel, V.A., Hou, J., O'Dell, T.J., and Coba, M.P. (2016). Long-term
potentiation modulates synaptic phosphorylation networks and reshapes the structure of the
postsynaptic interactome. Sci Signal *9*, rs8. 10.1126/scisignal.aaf6716.
Liang, C.T., Lin, Y.S., Huang, Y.C., Huang, H.L., Yang, J.Q., Wu, T.H., Chang, C.F., Huang,
S.J., Huang, H.B., and Lin, T.H. (2018). Characterization of the interactions between inhibitor-1
and recombinant PP1 by NMR spectroscopy. Sci Rep *8*, 50. 10.1038/s41598-017-18383-x.
Liao, D., Hessler, N.A., and Malinow, R. (1995). Activation of postsynaptically silent synapses
during pairing-induced LTP in CA1 region of hippocampal slice. Nature *375*, 400-404.
10.1038/375400a0.
Lin, C.H., Lin, W.D., Chou, I.C., Lee, I.C., Fan, H.C., and Hong, S.Y. (2018). Epileptic spasms
in PPP1CB-associated Noonan-like syndrome: a case report with clinical and therapeutic
implications. BMC Neurol *18*, 150. 10.1186/s12883-018-1157-6.
Liu, R., Correll, R.N., Davis, J., Vagnozzi, R.J., York, A.J., Sargent, M.A., Nairn, A.C., and
Molkentin, J.D. (2015). Cardiac-specific deletion of protein phosphatase 1beta promotes
increased myofilament protein phosphorylation and contractile alterations. J Mol Cell Cardiol
87, 204-213. 10.1016/j.yjmcc.2015.08.018.
Ma, L., Bayram, Y., McLaughlin, H.M., Cho, M.T., Krokosky, A., Turner, C.E., Lindstrom, K.,
Bupp, C.P., Mayberry, K., Mu, W., et al. (2016). De novo missense variants in PPP1CB are
associated with intellectual disability and congenital heart disease. Hum Genet *135*, 1399-1409.
10.1007/s00439-016-1731-1.
Makino, Y., Johnson, R.C., Yu, Y., Takamiya, K., and Huganir, R.L. (2011). Enhanced synaptic
plasticity in mice with phosphomimetic mutation of the GluA1 AMPA receptor. Proc Natl Acad
Sci U S A *108*, 8450-8455. 10.1073/pnas.1105261108.
Mammen, A.L., Kameyama, K., Roche, K.W., and Huganir, R.L. (1997). Phosphorylation of the
alpha-amino-3-hydroxy-5-methylisoxazole4-propionic acid receptor GluR1 subunit by
calcium/calmodulin-dependent kinase II. J Biol Chem *272*, 32528-32533.
10.1074/jbc.272.51.32528.
Mauceri, D., Gardoni, F., Marcello, E., and Di Luca, M. (2007). Dual role of CaMKII-dependent
SAP97 phosphorylation in mediating trafficking and insertion of NMDA receptor subunit NR2A.
J Neurochem *100*, 1032-1046. 10.1111/j.1471-4159.2006.04267.x.
Monaco, S.A., Ferguson, B.R., and Gao, W.J. (2018). Lithium Inhibits GSK3beta and Augments
GluN2A Receptor Expression in the Prefrontal Cortex. Front Cell Neurosci *12*, 16.
10.3389/fncel.2018.00016.
Morishita, W., Connor, J.H., Xia, H., Quinlan, E.M., Shenolikar, S., and Malenka, R.C. (2001).
Regulation of synaptic strength by protein phosphatase 1. Neuron *32*, 1133-1148.
10.1016/s0896-6273(01)00554-2.
Mota Vieira, M., Nguyen, T.A., Wu, K., Badger, J.D., 2nd, Collins, B.M., Anggono, V., Lu, W.,
and Roche, K.W. (2020). An Epilepsy-Associated GRIN2A Rare Variant Disrupts CaMKIIalpha
Phosphorylation of GluN2A and NMDA Receptor Trafficking. Cell Rep *32*, 108104.
10.1016/j.celrep.2020.108104.
Ragusa, M.J., Dancheck, B., Critton, D.A., Nairn, A.C., Page, R., and Peti, W. (2010).
Spinophilin directs protein phosphatase 1 specificity by blocking substrate binding sites. Nat
Struct Mol Biol *17*, 459-464. 10.1038/nsmb.1786.
Rex, C.S., Gavin, C.F., Rubio, M.D., Kramar, E.A., Chen, L.Y., Jia, Y., Huganir, R.L.,
Muzyczka, N., Gall, C.M., Miller, C.A., et al. (2010). Myosin IIb regulates actin dynamics

during synaptic plasticity and memory formation. Neuron *67*, 603-617.

10.1016/j.neuron.2010.07.016.

Sakimura, K., Kutsuwada, T., Ito, I., Manabe, T., Takayama, C., Kushiya, E., Yagi, T., Aizawa,

S., Inoue, Y., Sugiyama, H., and et al. (1995). Reduced hippocampal LTP and spatial learning in

mice lacking NMDA receptor epsilon 1 subunit. Nature *373*, 151-155. 10.1038/373151a0.

Shipton, O.A., and Paulsen, O. (2014). GluN2A and GluN2B subunit-containing NMDA

receptors in hippocampal plasticity. Philos Trans R Soc Lond B Biol Sci *369*, 20130163.

10.1098/rstb.2013.0163.

Sobczyk, A., Scheuss, V., and Svoboda, K. (2005). NMDA receptor subunit-dependent [Ca2+]

signaling in individual hippocampal dendritic spines. J Neurosci *25*, 6037-6046.

10.1523/JNEUROSCI.1221-05.2005.

Terrak, M., Kerff, F., Langsetmo, K., Tao, T., and Dominguez, R. (2004). Structural basis of

protein phosphatase 1 regulation. Nature *429*, 780-784. 10.1038/nature02582.

Terry-Lorenzo, R.T., Carmody, L.C., Voltz, J.W., Connor, J.H., Li, S., Smith, F.D., Milgram,

S.L., Colbran, R.J., and Shenolikar, S. (2002). The neuronal actin-binding proteins, neurabin I

and neurabin II, recruit specific isoforms of protein phosphatase-1 catalytic subunits. J Biol

Chem *277*, 27716-27724. 10.1074/jbc.M203365200.

Umeki, I., Niihori, T., Abe, T., Kanno, S.I., Okamoto, N., Mizuno, S., Kurosawa, K., Nagasaki,

K., Yoshida, M., Ohashi, H., et al. (2019). Delineation of LZTR1 mutation-positive patients with

Noonan syndrome and identification of LZTR1 binding to RAF1-PPP1CB complexes. Hum

Genet *138*, 21-35. 10.1007/s00439-018-1951-7.

Yong, X.L.H., Zhang, L., Yang, L., Chen, X., Tan, J.Z.A., Yu, X., Chandra, M., Livingstone, E.,

Widagdo, J., Vieira, M.M., et al. (2021). Regulation of NMDA receptor trafficking and gating by

activity-dependent CaMKIIalpha phosphorylation of the GluN2A subunit. Cell Rep *36*, 109338.

10.1016/j.celrep.2021.109338.

Young, L.C., Hartig, N., Boned Del Rio, I., Sari, S., Ringham-Terry, B., Wainwright, J.R., Jones,

G.G., McCormick, F., and Rodriguez-Viciana, P. (2018). SHOC2-MRAS-PP1 complex

positively regulates RAF activity and contributes to Noonan syndrome pathogenesis. Proc Natl

Acad Sci U S A *115*, E10576-E10585. 10.1073/pnas.1720352115.

Zambrano, R.M., Marble, M., Chalew, S.A., Lilje, C., Vargas, A., and Lacassie, Y. (2017).

Further evidence that variants in PPP1CB cause a rasopathy similar to Noonan syndrome with

loose anagen hair. American Journal of Medical Genetics Part A *173*, 565-567.

10.1002/ajmg.a.38056 PMID - 27868344.

Zhou, P., Zhu, L., Fan, Q., Liu, Y., Zhang, T., Yang, T., Chen, J., Cheng, Q., Li, T., and Chen, L.

(2020). A case report of Noonan syndrome-like disorder with loose anagen hair 2 treated with

recombinant human growth hormone. Am J Med Genet A *182*, 1967-1971.

10.1002/ajmg.a.61638.

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1