1 **Multivariate models of animal sex: breaking binaries leads to a better understanding of**
2 **ecology and evolution**
3

4 J. F. McLaughlin*, Kinsey M. Brock*[†#], Isabella Gates[‡], Anisha Pethkar[‡], Marcus Piattoni[‡],
5 Alexis Rossi[‡], Sara E. Lipshutz[‡]

6
7 The first two authors contributed equally to this work.

8
9 *Department of Environmental Science, Policy, and Management, College of Natural
10 Resources, University of California, Berkeley, CA, USA

11
12 [†]Museum of Vertebrate Zoology, University of California, Berkeley, CA, USA

13
14 [#]Department of Biology, San Diego State University, San Diego, CA, USA

15
16 [‡]Department of Biology, Loyola University Chicago, Chicago, IL, USA
17
18
19
20 Keywords: sexual selection, polymorphism, biological sex, terminology, reproduction, sexual
21 phenotype

22
23
24 ***Abstract***
25

'Sex' is often used to describe a suite of phenotypic and genotypic traits of an organism related
to reproduction. However, these traits – gamete type, chromosomal inheritance, physiology,
morphology, behavior, etc. – are not necessarily coupled, and the rhetorical collapse of variation
into a single term elides much of the complexity inherent in sexual phenotypes. We argue that
consideration of 'sex' as a constructed category operating at multiple biological levels opens up
new avenues for inquiry in our study of biological variation. We apply this framework to three
case studies that illustrate the diversity of sex variation, from decoupling sexual phenotypes to
the evolutionary and ecological consequences of intrasexual polymorphisms. We argue that
instead of assuming binary sex in these systems, some may be better categorized as
multivariate and nonbinary. Finally, we conduct a meta-analysis of terms used to describe
diversity in sexual phenotypes in the scientific literature to highlight how a multivariate model of
sex can clarify, rather than cloud, studies of sexual diversity within and across species. We
argue that such an expanded framework of 'sex' better equips us to understand evolutionary
processes, and that as biologists it is incumbent upon us to push back against
misunderstandings of the biology of sexual phenotypes that enact harm on marginalized
communities.

42
43

Introduction: Sex is not a single trait

In sexually reproducing species, the common assumption is that there are two sexes, strictly classified as female or male (Fisher 1930; Hurst 1996; Jaffe 1996). This assertion is supported by the cellular mechanisms of sexual reproduction, in which multiple (usually two) parental gametes combine genotypes to create an offspring with a novel genetic makeup. In animals, these gametes are of different sizes (e.g., large ova and small sperm), a condition called anisogamy. However, the binary classification of gametic sex breaks down when we consider the broader diversity of gametic phenotypes. For instance, hermaphroditic species possess both gamete types required for reproduction, and do not have separate sexes (Jarne and Auld 2006). Of the estimated 1.2 million animal species, roughly 5-6% are hermaphroditic (Jarne and Auld 2006). Outside of animals, systems of sex determination rely on genetic markers that determine compatibility between equally-sized gametes, a condition known as isogamy (Togashi and Cox 2011) found in fungi (Kothe 1996; Lee et al. 2010; Billiard et al. 2011, 2012), algae (Perrin 2012; Tillmann and Hoppenrath 2013), and amoebozoa (Douglas et al. 2016).

Operationally, the term 'sex' has two meanings - one, as a reproductive process that refers to the transmission of genetic information to the next generation, and another, as a categorical term that encompasses a broad collection of gametic, genetic, hormonal, anatomic, and behavioral traits (Gross 1996; Whitfield 2004; Engqvist and Taborsky 2016; Mank 2022). Whereas some biologists argue that gametes are the only meaningful sex categories (Goymann et al. 2023), we find several limitations to this gametic sex definition, particularly for ecologists and evolutionary biologists. First, gametes are rarely measured directly, with researchers instead relying on genetic and phenotypic proxies. Second, though biologists have drawn a direct connection between the evolution of gamete size and other sexual phenotypes (e.g., Kalmus 1932), these traits (gametes, genotype, hormones, anatomy, behavior) are not universally coupled. Joan Roughgarden summarized this problem: "the biggest error in biology today is uncritically assuming that the gamete size binary implies a corresponding binary in body type, behavior, and life history" (2013). Finally, selection typically acts on phenotypic traits, and variation in these traits is the raw material of evolution Thus, if we want to understand the evolution of diversity, we need to expand, rather than collapse, our definition of sex beyond binary categorization.

As a categorical term, 'sex' is often semantically flattened into a binary, univariate model, for which individuals are classified as either 'female' or 'male' (Figure 1A) in gonochoristic species (i.e., an individual produces one type of gamete throughout its lifespan). A more expansive definition of sex is bimodal - with most individuals falling within one of two peaks of a trait distribution (Figure 1B). However, even a bimodal, univariate model is an oversimplification, since 'sex' comprises multiple genetic and phenotypic traits, with variable distributions (see Figure 1C and case studies). Individuals may possess different combinations of chromosome type, gamete size, hormone level, morphology, and social roles, which do not always align or persist across an organism's lifespan (Karkazis 2019; Griffiths 2021). Reliance on strict binary categories of sex fails to accurately capture the diverse and nuanced nature of sex.

Here we propose a model of animal sex as multivariate (Figure 1C), encompassing
multiple traits that may or may not follow a binary distribution. Together, the distributions of
these traits– gametic, hormonal, morphological, behavioral, etc.– comprise the sexual
phenotype of an individual within a species. We lay out the shortcomings of assuming and
analyzing sex as binary across these biological levels of organization. We then provide three
case studies that illustrate the diversity of sex variation: decoupling sexual phenotypes in 'sex-
role reversed' birds, the evolutionary consequences of a chromosomal inversion resulting in
more than two sexes, and the ecological and evolutionary consequences of intrasexual color
polymorphism in lizards. We argue that rethinking the default assumption of binary sex can help
us better understand the evolutionary and ecological contexts in which these animals operate.
We conduct a meta-analysis of terminology used to describe sexual phenotypes in scientific
literature. Finally, we offer recommendations for future research and education on sex diversity.
100
We are certainly not the first, nor the last biologists to interrogate the definition of sex
(Hoekstra 1990; Fausto-Sterling, 2012; Roughgarden 2013; Karkazis 2019), or the nature of
biological categories more broadly (Cadena and Zapata 2021). We rely on literature mainly
pertaining to organismal biology, spanning the fields of molecular genetics, neuroendocrinology,
behavior, ecology, and evolution. We find it useful to clarify that our focus is primarily on
animals, and vertebrates in particular. We also differentiate between the concepts of sex and
gender (Goymann and Brumm 2018). Though the term 'gender' has occasionally been used to
describe sexual phenotypes (Roughgarden 2013; Kutschera 2016), this term describes
multifaceted and abstract self-interpretations of identity, an internal personal experience that
expresses itself in the context of an individual's specific culture, place, and time (Thorne et al.
2019). We validate the important role of gender in the human experience, especially in regard to
transgender, nonbinary, and gender nonconforming (TNGC) identities. However, in studies of
non-human animals it is key to differentiate between the specifically cultural role of gender and
the observed traits of other organisms that we broadly call 'sex' (Goymann and Brumm 2018).
Thus, throughout this paper we focus solely on sex, as anything similar to the human
experience of gender in other organisms is fundamentally inaccessible to us.
117

118
Figure 1: Three models of sexual phenotypes, demonstrated by their distribution across a
hypothetical population: A) A strict binary, for which all individuals are unambiguously grouped
into one of two categories. Whereas some traits, such as gamete size, operate this way, a
binary does not accurately model the distribution of most phenotypic traits. B) A bimodal model,
with most individuals falling around two peaks, or means, of a single sexual trait with continuous
variation. C) A multivariate model of sex as a collection of traits, represented as individual lines,

that contribute to the overall sex phenotype. Each trait has its own distribution, which may or
may not be bimodal, or coincide with other traits.
127
128 ***Sex at the genetic level***
129
The genetics of sexual determination vary widely (Bachtrog et al. 2014; Figure 2). In the case of
highly specialized sex chromosomes with reduced recombination, biologists classify individuals
as homogametic (i.e., have two of the same sex chromosome type) or heterogametic (having
two distinct chromosome types), with sexual phenotypes following from those genotypes.
However, even these 'simple' systems include intermediate (intersex) phenotypes (Vaiman and
Pailhoux 2000), as additional genes outside the sex chromosomes modulate developmental
pathways. The most widely familiar systems are XX/XY, where the presence of certain sex-
determining genes, such as *sry* on the Y chromosome in most mammals, activates diverging
pathways early in the developmental process (Bachtrog et al. 2014; Stévant et al. 2018) (Figure
2A). ZZ/ZW systems operate similarly, but with the ova-producing individuals being
heterogametic (Bachtrog et al. 2014) (Figure 2B). In some taxa, rather than a small number of
genes on the smaller chromosome, it is the dose-dependent effects of homogametic
chromosomes that lead to higher expression of genes that start a particular development
pathway (Smith et al. 2009). Dose-dependent effects also underlie the haplodiploid systems of
many insects, where unfertilized and fertilized ova develop into different social and reproductive
classes (Beye et al. 2003; Verhulst et al. 2010; Figure 2C). Some combination of sex-
determining genes and dosage dependence may be at play in species for which the smaller sex
chromosome and most of its associated genes have been lost entirely (Sutou et al. 2001; Just
et al. 2002; Kuroiwa et al. 2010).
149
In polygenic systems, multiple loci on several chromosomes, both sex-linked and
autosomal, interact to determine sexual phenotypes. Polymorphisms in the sex chromosomes
can override sex-determining genes (Figure 2D; Veyrunes et al. 2010; Saunders et al. 2014;
Zhao et al. 2017) or lead to multiple reproductive morphs (Sandkam et al. 2021). In several
species of cichlid fish, sex is determined by the interplay of both ZZ/ZW and XX/XY
chromosomes, resulting in four genotypes but two phenotypes (Ser et al. 2010; Moore et al.
2022; Figure 2E). In species with these polygenic systems, the number of reproductive
phenotypes is not necessarily the same as the number of genotypes in the population,
demonstrating how genotype alone is not sufficient in defining sexual categories.
159
Not all sexual differentiation is controlled by genetic mechanisms, in part or whole
(Bachtrog et al. 2014). Temperature-dependent sex determination is common among reptiles
(Figure 2F) (Sarre et al. 2004) and fish (Baroiller et al. 2009). In its simplest form, the
temperature of incubation determines which hormonal pathways are triggered in a developing
egg (Sarre et al. 2004). In other systems, temperature can override genetic pathways, resulting
in animals with the genetic markers usually associated with one sex, but the phenotypic traits of
another (Quinn et al. 2009; Holleley et al. 2015; Whiteley et al. 2017; Wiggins et al. 2020). Other
environmental variables may likewise interact with genetic sex determination, possibly through
epigenetic factors (Cabej 2018; Piferrer 2021), but these last mechanisms are only just

beginning to be explored. Altogether, evidence from these diverse species reveals that genetic
sex is not the universal binary often assumed, and that genetic sex is one variable among the
many traits that comprise multivariate sex.

Figure 2: Mechanisms of sex determination vary widely across animals. The most familiar (top row) are the relatively simple (**A**) XX/XY (in purple, here shown with northern elephant seal, *Mirounga angustirostris*, as an example taxon) and (**B**) ZZ/ZW (in green, shown with royal flycatcher, *Onychorhynchus coronatus*). (**C**) Haplodiploid sex determination involves sex being determined by having one or two copies of the genome, and is found predominantly in insects such as the pictured *Dasymutilla occidentalis*. (**D**) A mutation in the X chromosome of some African pygmy mice (*Mus minutoides*) overrides the presence of a Y chromosome, and (**E**) in some cichlids (shown here with *Metriaclima mbenjii*), both the absence of the W chromosome and presence of the Y are required for an individual to develop 'male' characteristics. (**F**) In central bearded dragons (*Pogona vitticeps*), the ZZ genotype is overridden by high temperatures, leading to phenotypically 'female' individuals capable of laying eggs. (**G**) The white-throated sparrow (*Zonotrichia albicollis*) is found in two color morphs linked to a large inversion on chromosome two (in turquoise), which is paired with the Z and W genotype to determine which individuals typically reproduce with each other.

Sex at the endocrine level

Hormones influence a number of sexual phenotypes, including 1) morphological traits such as gonads, genitals, ornamentation, and body size, 2) reproductive states such as estrous cycling and gametogenesis, and 3) behaviors such as courtship displays, receptivity to mating, aggression, and parental care (Adkins-Regan 2005). So-called 'sex' hormones typically refer to steroid hormones produced in the testes and ovaries, and include estrogens, progestogens, as well as several androgens (testosterone, androstenedione, dihydrotestosterone, and 11-ketotestosterone). However, many of these hormones are also produced outside of the gonads, including in the adrenal cortex, fat, and brain (Schmidt et al. 2008, Saldanha et al. 2011). Furthermore, variation in other elements of endocrine signaling pathways (e.g., receptors, enzymes, steroid binding proteins) beyond circulating hormone levels may contribute to the expression of sexual phenotypes (Adkins-Regan 2005, Hau 2005). For instance, in male zebra finches (*Taeniopygia guttata*), testosterone is converted to 17β-estradiol via the enzyme aromatase in the brain, which then regulates directed singing (Walters et al. 1991). In another songbird, dark-eyed juncos (*Junco hyemalis*), variation in the expression of genes encoding the androgen receptor, estrogen receptor, and aromatase enzyme correlates with aggression in both sexes (Rosvall et al. 2012). Thus, androgens and estrogens are not strictly associated with regulating the expression of 'male typical' and 'female typical' morphologies and behaviors. However, a common misconception is that androgens regulate 'masculine' phenotypes and estrogens regulate 'feminine' phenotypes, which has led to the widely used terms 'masculinization' and 'feminization'. We argue that this terminology is circular and confusing. Steroid hormones are not sex-specific - both females and males possess the capacity to produce, metabolize, and respond to these hormones (Staub and De Beer 1997). Thus, testosterone is not a 'male hormone,' and estrogen is not a 'female hormone'.

    Whereas it is often the case that average peak hormone levels may be higher or lower in one sex, this depends on various contexts, including when animals are developing (organizational effects) and when they are breeding (activational effects; Phoenix et al. 1959), as well as in response to social stimuli (Wingfield et al. 1990; Goymann et al. 2019). Reflecting back on our multivariate framework, testosterone levels in circulation are typically higher in adult

breeding males than in females for many species (Goymann and Wingfield 2014; Van Leeuwen
and Bladh 2016), reflecting the bimodal distribution depicted in Figure 1B. However, there are
many hormonal patterns that follow the distributions depicted in Figure 1C. For example,
testosterone in many temperate bird species peaks seasonally during territorial establishment
and mating, but decreases to basal, non-detectable levels during the non-breeding season,
when gonads regress (Wingfield et al. 2019). Within an individual, testosterone levels vary
substantially throughout the day (Panico et al. 1990; Brambilla et al. 2009; Greives et al. 2021).
Given their daily, seasonal, and cross-species variation, hormone levels are not a binary
indicator of sex.
228
229 ***Sex at the morphological level***
230
Organismal biologists often assign binary sex to species based on their primary sexual
characteristics, despite the diversity of genital morphology within and among taxa. For example,
spotted hyenas (*Crocuta crocuta*) mate, give birth, and urinate through a urogenital canal at the
tip of a hypertrophied clitorophallus (Glickman et al. 2006), which they may use to signal social
status (Hofer and East 1995) or to facilitate post-copulatory sperm choice (Holekamp 2006). In
Neotrogla barkflies, females have a highly elaborate penis (gynosome) that retrieves sperm
from the male's vagina (Yoshizawa et al. 2014). In color polymorphic lizards, male color morphs
have distinct hemipene morphology associated with their alternative mating strategies (Gilman
et al. 2019). Phallus polymorphism is common in hermaphroditic gastropods, in which some
individuals have a functional penis (euphally), some have a non-functional penis (hemiphally),
and some have no penis (aphally) (Leonard et al. 2007). In birds, only 3% of species even have
phalluses (external genitalia) (Brennan 2022). As with phallic diversity, animal morphologies
involved in birthing and brooding exhibit extraordinary intra- and intersexual variation. Male
pregnancy has evolved several times in syngnathid fishes (seahorses and pipefish) where
gestation in males requires morphological and physiological changes in paternal tissues to
incubate developing embryos in a specialized brooding pouch, analogous to structures in
viviparous females (Carcupino et al. 2002). In many birds, for which biparental care is common
(Clutton-Brock 2019), any adult can develop brood patches to incubate eggs (Tucker 1943). In
mammals (humans included), genital structures are not formed de novo in males and females,
but develop from a shared anatomical basis that can differentiate into forms other than the two
most commonly observed pathways, leading to intermediate phenotypes that are unfortunately
pathologized (Quigley et al. 1995; Dreger 2006; Grimstad et al. 2021). Thus, as a result of
variation in developmental pathways, external genitalia are not necessarily concordant with sex
chromosome makeup, and the presence, shape, and size of genitalia do not always fall into
binary categories of morphological sex.
256
Secondary sexual characteristics include ornamentation, color, weaponry, and many
other sexual morphologies that are not directly involved in reproduction. Although these traits
are expected to be sexually dimorphic, there are many exceptions. For example,
polymorphisms, both sex-limited and not, present a classic example of more than two sexual
phenotypes (Mank 2022). The three male morphs of the ruff (*Philomachus pugnax*), a bird in the
sandpiper family, exhibit extremely divergent size and plumage ornamentation, which

correspond to alternative mating strategies and chromosomal genotypes (Küpper et al. 2016). In
many odonates there are multiple female morphs, consisting of one or more 'gynomorphs' that
vary in color, and one 'andromorph' that closely resembles males in color, sexual
ornamentation, and behavior, presumably to avoid mating harassment (Gallesi et al. 2015).
These so-called 'andromorphs' exhibit morphological and behavioral variation distinct from
males and 'gymnomorph' females (Paulson 1998; Fincke et al. 2005; Gallesi et al. 2015),
resulting in an overall phenotype that does not conform to a binary distribution and likely arises
due to a variety of factors beyond male mimicry (Fincke et al. 2005). For both primary and
secondary sexual morphologies, there are many cases in which the distribution of these traits is
continuous, rather than binary.
273
274 ***Sex at the behavioral level***
275
Scientific notions of how animals 'should' adaptively behave originate with Darwin in the
Victorian era (Darwin 1871). The Darwin-Bateman paradigm posits that 'traditional' sex roles
arise from anisogamy, such that mate competition is stronger in the sex with the smaller
gametes (Bateman 1948; Dewsbury 2005). Trivers (1972) extended this paradigm with the
hypothesis that gametic investment drives postzygotic parental investment, which should be
higher in females because of larger gamete size. Theories on the evolution of sex roles have
dominated sexual selection research, but not without criticism (Gowaty et al. 2012; Ah-King
2013; Tang-Martínez 2016), as others have argued for the importance of ecology, sex ratio, and
life history traits on the evolution of sex-specific behaviors (Kokko and Jennions 2008; Mokos et
al. 2021; Kappeler et al. 2022). Indeed, there are many examples of competitive females and
choosy males across animal groups (reviewed in Hare and Simmons 2018; Edward and
Chapman 2011). Outside of mammals, many examples in fish, frogs, and birds contradict the
notion that the sex with the larger gamete suffers higher mating costs and invests more
resources into the next generation, such as the repeated evolution of male parental care in frogs
(Townsend et al. 1984; Gross 2005; Furness and Capellini 2019).
291
Within species, alternative reproductive strategies are a classic example of
discontinuous variation in behavior (Gross 1996; Sinervo and Lively 1996; Oliveira et al. 2008;
Stiver et al. 2015), and can arise via different mechanisms of genotype, condition dependence,
and social environment. For example, in the bluegill sunfish (*Lepomis macrochirus*), there are
three different reproductive strategies found in males, one of which overlaps with courtship
soliciting behaviors in females (Dominey 1980). The diversity of sexual behaviors also includes
the common occurrence of same-sex sexual behavior (SSB) in animals (Monk et al. 2019).
SSBs have been observed across a large number of animals (Bailey and Zuk 2009), but were
frequently unreported by early naturalists due to societal biases (Russell et al. 2012).
301
Compared to the other traits we review, sex studied at the social level seems to more
readily acknowledge variation outside of binary categories, perhaps because plasticity is a
fundamental aspect of behavioral research. Furthermore, several recent studies in animal
behavior and evolutionary biology have acknowledged that the positionality of the scientists
themselves influences their research (Ahnesjö et al. 2020; Tang-Martínez 2020; Ah-King 2022;

Pollo and Kasumovic 2022), making more space for scientists to reflect on how their identities
shape their scientific practices.

We next provide three case studies, which highlight how a de-constructed approach to
sex can lead to asking novel empirical questions. These examples come from our own study
systems and research expertise. Rather than considering these systems as rare exceptions, we
can use them as models to understand the diversity of sexual phenotypes across multiple
biological levels.

Case study 1: 'Sex-role reversal' and the decoupling of sexual phenotypes
Binary expectations of sexual phenotypes are often met with exceptions, and one classic
example are birds with polyandrous mating systems, including black coucals (*Centropus grillii*),
spotted sandpipers (*Actitis macularius*), buttonquails (Turnicidae), and northern jacanas (*Jacana*
spinosa). In these systems, individuals with larger gametes typically face stronger sexual
selection for mating opportunities (Fritzsche et al. 2021), decoupling sexual phenotypes from
anisogamous expectations. These systems are often called 'sex-role reversed' because they
defy 'traditional' expectations of sex roles historically found in mammals - male competition and
female parental care. However, sex role terminology imposes binary categorization, limiting our
understanding of continuous variation in phenotypes (Ah-King and Ahnesjö 2013). As we
describe below, polyandrous mating systems are not simply the 'reverse' of polygynous mating
systems, but rather, they are novel evolutionary outcomes to be studied in their own right, not
solely in relation to a polygynous foil. We propose that studying so-called 'exceptional' systems
can help reveal unexamined routes of phenotypic evolution, expanding the conceptual
framework for how we study sexual diversity.

Here, we review examples from northern jacanas (*Jacana spinosa*), a socially
polyandrous shorebird and a focal species of one of our co-authors. Jacanas allow us to
evaluate: To what extent do the sexes differ in the first place? For which traits, and at what
developmental and seasonal timepoints? At the genetic level, jacanas have ZW/ZZ sex
determination and females are heterogametic - this categorization, as for most birds, is binary.
At the endocrine level, circulating testosterone concentrations are context-dependent. Females
have significantly lower testosterone than males in the courtship stage, but sex differences in
testosterone disappear when males are conducting parental care (Lipshutz and Rosvall 2020a).
Thus, the degree of sexual dimorphism in testosterone varies with male breeding stage, a
pattern that applies broadly across independent origins of social polyandry (Eens and Pinxten
2000; Lipshutz and Rosvall 2020b). The gonadal size of territorial males also fluctuates between
reproductive stages of courtship and parental care, whereas territorial females remain in a
continuous state of fertility (Lipshutz and Rosvall 2020a). For this species, the cyclical variation
in hormonal and gonadal traits reveals that these sexual phenotypes, and their degree of sexual
dimorphism, are dynamic.

At the morphological level, sexual size dimorphism is biased towards females, who have
more exaggerated ornamentation, weaponry, and body size (Emlen and Wrege 2004) There is
some evidence that wing spur length positively correlates with testosterone in females, but not

males (Lipshutz and Rosvall 2020a), suggesting that hormonal sensitivity may vary between the
sexes. At the social level, female jacanas defend territories that encompass multiple male mates
(i.e. social polyandry), and males conduct the majority of parental care (Emlen and Oring 1977).
Despite expectations that 'role-reversed' male jacanas should be choosy, virtually nothing is
known about mate choice in this system. Thus far, competition for territories seems to be the
predominant determinant of mating success in jacanas (Emlen and Wrege 2004). Larger
females and males gain access to breeding territories, whereas floater females and males do
not reproduce. Thus, reproductive access varies within and among both sexes, and there are
non-reproductive phenotypes for both females and males. Altogether, the genetic, endocrine,
morphological, and social traits of 'sex-role reversed' species such as northern jacanas
demonstrate that sexual phenotypes are dynamic, complex, and support a multivariate
framework of sexual diversity (Figure 1C, 3).
363

364
365 **Figure 3:** Multivariate sex in the northern jacana. 'Sex' includes phenotypic variation at multiple
366 levels, from genetic makeup, anatomy and morphology, hormone levels, and social roles and
367 behaviors.
368
369 ***Case study 2: The evolutionary consequences of more than two sexes***
370
How does the number of operative sexes influence the evolution of a species? White-throated
sparrows (*Zonotrichia albicollis*), have been described as 'the bird with four sexes', with both tan
and white stripe morphs occurring in ZZ and ZW individuals. This species mates disassortatively
by color morph, with tan stripe morphs predominantly mating with white stripe morphs (Hedrick
et al. 2018). The more aggressive white-stripe morph has a large inversion on chromosome 2
that functions as a 'supergene', which tan-stripe individuals lack (Tuttle et al. 2016; Falls and
Kopachena 2020; Maney et al. 2020;Figure 2G). These morphs are further associated with

differences in testosterone and estrogen (Maney and Küpper 2022). Population persistence as
a whole requires that all four morphs be present (Campagna 2016; Maney et al. 2020). A similar
system occurs in hybrid *Pogonomyrmex* ants, where each colony requires three sexes to
operate, and four to persist beyond a single generation (Parker 2004).
382
There are significant evolutionary consequences to having more than two operative
sexes. A key parameter in population genetics is effective population size (N_e; Charlesworth
2009), which is fundamentally impacted by the number of sexes in a given population (Caballero
1994). Assuming an equal sex ratio, the absolute maximum of N_e will be equal to $\frac{N}{S_O}$, where N is
the census population size and S_o is the operative number of sexes. As N_e influences the
strength of selection, genetic drift, and genetic diversity of a population (Charlesworth 2009), the
operative number of sexes substantially impacts the evolutionary trajectory of a lineage. The
number of sexes in a system also shapes genome evolution (Charlesworth and Mank 2010),
mating kinetics (Power 1976; Iwasa and Sasaki 1987), and sexual selection (Emlen and Oring
1977), and thus a framework of sex that accounts for diversity in sexual phenotypes allows us to
more accurately model evolutionary processes (Mokos et al. 2021).
394
We can gain novel insights by applying this framework to other taxa with genetic
polymorphisms. The previously mentioned ruff also has a genetic polymorphism linked to
multivariate phenotypic differences in reproductive strategies (Lank et al. 1995; Jukema and
Piersma 2006). It is typical to frame such systems as having *subtypes* of two sexes– in this
case, three males and one female. But if we are willing to consider the white-throated sparrows
and *Pogonomyrmex* systems as having more than two sexes, why not apply a similar
framework to the ruff and other polymorphic systems (Galeotti et al. 2003; Roulin 2004)? Our
binary framing of sex in these systems may interfere with understanding their evolutionary
trajectories.
404
405   ***Case study 3: Ecological and evolutionary consequences of multimodal sex in color***
406   ***polymorphic lizards***
407
Color polymorphism, the evolution of two or more genetically-determined color morphs within a
single population (Ford 1945; Sinervo et al. 2001; Andrade et al. 2019), is a classic example of
multivariate and multimodal sexual diversity (Figure 1C; Mank 2022). Intrasexual color morphs
have evolved in every major animal group across the tree of life, and are particularly common in
lizards (Cote et al. 2008; Stuart-Fox et al. 2021; Brock, McTavish, et al. 2022a). In lizards,
intrasexual color morphs occur in both females and males, and are often associated with
alternative sexual phenotypes comprising discontinuous variation in genital morphology (Gilman
et al. 2019), clutch size (Sinervo et al. 2000), mate preference (Pérez i de Lanuza et al. 2013),
mating strategies (Gross 1982; Sinervo and Lively 1996), and endocrine profiles (Huyghe et al.
2009; Brock et al. 2020). These distinct sexual phenotypes give rise to ecological specialization
among morphs, including thermal preference (Thompson et al. 2023), microhabitat selection
(BeVier et al. 2022), diet (Lattanzio and Miles 2016), and non-sexual behaviors (Brock and
Madden 2022; Brock, Chelini, et al. 2022b). Although each morph utilizes a specialized set of
resources, in aggregate, greater morph diversity can contribute to a greater niche breadth in

polymorphic populations relative to monomorphic populations (Pérez i de Lanuza et al. 2018;
Forsman et al. 2008); Figure 4). For example, in *Podarcis* wall lizards with three genetically-
determined throat color morphs (Figure 4), morphs have different preferred body temperatures
and microhabitat use that results in morphs that use a narrow space within the overall species'
environmental niche space (Pérez i de Lanuza and Carretero 2018; Thompson et al. 2023). As
an ecological consequence, the overall greater niche breadth of polymorphic populations is
hypothesized to confer greater capacity for range expansion, less susceptibility to environmental
change, and a larger buffer to extinction risk (Forsman et al. 2008). Evolutionary consequences
of multimodal morphs include punctuated patterns of trait evolution, greater mobility on the
adaptive landscape, and increased diversification rates (West-Eberhard 1986; Corl et al. 2010;
Hugall and Stuart-Fox 2012; Brock, McTavish, et al. 2022a). New species could form from
different morphs if they become reproductively isolated or fixed within a population (West-
Eberhard 1986; Gray and McKinnon 2007); (Seehausen et al. 2008; Corl et al. 2010; McLean
and Stuart-Fox 2014; Brock, Madden, et al. 2022c). In addition to facilitating speciation,
character release associated with morph fixation could produce punctuated accelerations of
morphological change (Eldredge 1976; West-Eberhard 1986; Corl et al. 2010). Thus, the
ecological differences that emerge from intrasexual polymorphism are a crucial component of
biodiversity within a species, which has far-reaching consequences for population persistence in
a rapidly changing world. Collapsing intrasexual polymorphisms into a female-male binary
erases extensive multivariate phenotypic variation (Figure 4), including its ecological and
evolutionary consequences.
443

445   **Figure 4:** Illustration of alternative patterns of population-level environmental niche breadth
446   (dotted line) distributions for A) monomorphic, B) dimorphic, and C) polymorphic populations of
447   wall lizards (*Podarcis*). Different color morphs (solid color lines) have non-overlapping
448   environmental resource use that, in the aggregate, result in an overall greater niche breadth in
449   an A) monomorphic population compared to B) dimorphic and C) polymorphic populations.
450
These case studies demonstrate that breaking the sex binary - and expanding to a multivariate
model of sex - is a useful framework for understanding animal ecology and evolution. Each of
these animal systems possesses a different set of genetic, endocrine, morphological, and
behavioral sexual traits, which do not fit neatly into binary distributions. Once we detach our
assumptions of binary sex classifications, we can ask more expansive questions such as 'how
does variation within and among these sexual traits shape ecological and evolutionary
processes?'
458
459   ***Literature survey: How does terminology for sexual diversity change over time?***
460
The language used in published research forms the foundation of how we as scientists think
about, discuss, and investigate the natural world. To track how terminology usage has changed
over time in ecology and evolutionary biology, we conducted a systematic review on Web of
Science of search terms associated with genetic, endocrine, morphological, and behavioral
levels of sex. Our search terms included the following: "andromorph/gynomorph" (60 papers),
"female/male hormone" (40) "female-like/male-like" (124), "feminize/masculinize" (539) as well
as outdated and transphobic language, including "transsexual", "transvestite" and "she-male"
(7). We also surveyed the conflation of sex and gender (1047). Additional details are available in
the Supplementary Materials.

Figure 5. Literature survey for terms (a) by taxonomy and (b) over time. The search was
conducted on 18 October 2022.

We initially identified 2,222 papers, 1,817 of which were exclusive to non-human
animals, could be assigned a taxonomic class, and were available to Loyola University Chicago
online (Table S1, Supplementary Materials). We used a chi-square test to evaluate taxonomic
bias associated with each term, and a line of best fit to examine change in search term usage
over time.

We found that most terms were significantly associated with a taxonomic class, except
for masculinize/feminize and gender/sex conflation (Table S2). Andromorph/gynomorph are
almost exclusively used in insects, particularly damselflies. "Female-like" and "male-like" are
most commonly used in birds. Application of the outdated and offensive term "she-male" was
confined to reptiles, garter snakes (*Thamnophis* sp.) in particular. Most terms in our literature
survey increased in frequency over time, except for "she-male" (Figure S2).The conflation of sex
and gender increased in the past three decades (Figure 5b), consistent with findings from a
previous study (Goymann and Brumm 2018).

Terminology describing sex can have different meanings for different groups of people.
Language is influenced by culture, and this can shift rapidly over time. What is popular today
may not necessarily be seen as appropriate in a few years. Thus, it is important for biologists to
be explicit in how we use these terms in our work. Sex is complex, so talking about it requires
context and specificity - are we inferring sexes genetically? From their gonads? By color?
Taking this path forward is important to not only reduce harm and improve science, but also to
encourage and facilitate more research in systems and species that do not fit our own societal
perceptions of what is "normal" when it comes to sexual phenotypes.

The interplay of science and society

Multivariate models of sex reveal overlapping but not necessarily coincident phenotypes at
every biological level within an individual, from the molecular to the behavioral (Maney, 2016). In
zoology, we impose a binary categorization of sex as an emergent property of many traits.
Whereas some of these traits do typically have a bimodal distribution (some chromosomes,
gametes), others demonstrate largely continuous or multimodal variation (hormone levels
[(Wingfield et al. 1990), morphology [Mank 2022], behavior [Dominey 1980)]), suggesting that
most animals can best be studied from the framework of multiple phenotypic axes– some
categorical, but most continuous. Even the basic inclusion of sex as a variable is missing from
many studies, particularly in fields related to human health (Woitowich et al. 2020; Garcia-
Sifuentes and Maney 2021). However, uncritically applying a simple binary without considering
the mechanisms shaping sex-specific effects can confound inferences (Casto et al. 2022) and
when applied to humans, completely erases the biological realities of trans, nonbinary, and
gender-nonconforming (TGNC) and intersex people (Cheung et al. 2021; Phiri-Ramongane and
Khine 2022).

Though we focus on animals, embracing complex models of sex has also led to greater
understanding of fungi and plants (Lee et al. 2010; Billiard et al. 2011, 2012). For example,
exploration of the diversity of fungal sex has illuminated the evolution of the process of sexual
reproduction (Lee et al. 2010), the advantages and disadvantages of anisogamy (Billiard et al.
2011), and the evolutionary success of fungi in general (Naranjo-Ortiz and Gabaldón 2020).
Here, we have argued that applying such a lens to animals may lead to similarly novel insights.
Some readers may consider our examples of sexual diversity as rare anomalies – exceptions to
the evolutionary rule, which do not require a novel framework to explain. We acknowledge some
variations in sexual phenotypes may be relatively less common in animals, but that does not
minimize their importance in our understanding of evolution. When we regard diversity as a
deviation from the norm, we ignore its successful role in generating novel solutions to
evolutionary challenges. Our frameworks of sex need to accommodate and embrace this
diversity, lest we fall prey to a deterministic model of evolution, where the ideal endpoint is
always binary.
529
As biologists, we seek to understand the vast diversity of life in all its wonderful forms -
yet the lens we use fundamentally shapes that which we can observe. The historical legacies of
sexism, racism, queerphobia, and ableism have deeply influenced the frameworks we use to
study nature (Branch et al. 2022; Graves et al. 2022; Kamath et al. 2022), and the exclusion of
certain groups of people from the scientific enterprise has delayed important discoveries such
as the prevalence of female bird song (Haines et al. 2020) and the existence of hemiclitores in
snakes (Folwell et al 2022a; Folwell et al 2022b). Challenging these foundations is difficult but
vital to both increasing inclusion in biology (Hales 2020; Casper et al. 2022) and dismantling
assumptions that interfere with our ability to observe and interpret the natural world (Monk et al.
2019; Ahnesjö et al. 2020; Kamath et al. 2022; Packer and Lambert 2022).
540
We recognize that TGNC and intersex people are valid regardless of the sexual diversity
we find in nature. Human rights cannot be defined solely by biology - this is a prime example of
the appeal to nature fallacy. However, opposition to the inclusion of TGNC and intersex people
in society is often based in the language of biology. Especially in the United States, legislation
targeting TGNC people is increasingly undergirded with the simplistic binary model of 'biological'
sex (such as OH HB454 §3129.02 2021, WV HB3293 §18-2-25d.b1 2021, MO SB22 §191.1720
2022, and TX HB672 88R §71.004.1A 2022). There is pressure for scientists to avoid making
the politics of our work explicit, especially those of us who do not directly study social issues.
However, our science is being weaponized to discriminate against marginalized groups (Ha et
al. 2014). It is imperative that we biologists challenge the misuse and abuse of this language
(Miyagi et al. 2021) and confront how our scientific models impact society (Bazzul and Sykes
2011). As biologists, it is our responsibility to dispel misconceptions and recognize the rich
tapestry of diversity in nature.
554
555
556 ***Acknowledgements***
557

Thanks to Kelsey Lewis and Sam Sharpe for the opportunity to contribute to this symposium. Aqdas Aftab, Wolfgang Goymann, and Kim Rosvall provided fruitful conversations discussing sex and gender diversity. Claire Curry and Elliott Smith provided vital assistance in designing the systematic review and mapping analysis. Thank you to Savvy Cornett in the Hofmann Lab at UT Austin and Kwasi Wrensford in the UC Berkeley Animal Behavior Reading Group for invitations to discuss the preprint version of this paper. Anusha Bishop, Anne Chambers, Jay Falk, Ambika Kamath, Daniel Olivera, Erin Westeen, Alejandro Rico-Guevara and four anonymous reviewers provided valuable feedback on the manuscript. Thank you to the Trans Formations Project, whose database of legislation was very helpful in understanding the use of the term 'biological sex' in recent legislation – we encourage interested readers to follow their tracking efforts at https://www.transformationsproject.org/.

Conflict of Interest

This research was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest.

References

Adkins-Regan E. 2005. Hormones and animal social behavior. Monographs in Behavior and Ecology. Princeton (NJ): Princeton University Press.

Ah-King M. 2013. On anisogamy and the evolution of "sex roles." Trends Ecol Evol 28:1-2.

Ah-King M. 2022. The female turn. Singapore: Springer Nature.

Ah-King M, Ahnesjö I. 2013. The "sex role" concept: an overview and evaluation. Evol Biol 40:461–70.

Ahnesjö I, Brealey JC, Günter KP, Martinossi-Allibert I, Morinay J, Siljestam M, Stångberg J, Vasconcelos P. 2020. Considering gender-biased assumptions in evolutionary biology. Evol Biol 47:1–5.

Andrade, P, Pinho C, Pérez i de Lanuza G, Afonso S, Brejcha J, Rubin C-J, Wallerman O, Pereira P, Sabatino SJ, Bellati A, et al. 2019. Regulatory changes in pterin and carotenoid genes underlie balanced color polymorphisms in the wall lizard. PNAS 116:5633–42.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman T-L, Hahn MW, Kitano J, Mayrose I, Ming R, et al. 2014. Sex determination: why so many ways of doing it? PLoS Biol 12:e1001899.

Bailey NW, Zuk M. 2009. Same-sex sexual behavior and evolution. Trends Ecol Evol 24:439–46.

Baroiller JF, D'Cotta H, Saillant E. 2009. Environmental effects on fish sex determination and differentiation. Sex Dev 3:118–35.

Bateman AJ. 1948. Intra-sexual selection in *Drosophila*. Heredity (Edinb) 2:349–68.

Bazzul J, Sykes H. 2011. The secret identity of a biology textbook: straight and naturally sexed. Cult Stud Sci Educ 6:265–86.

BeVier GT, Ayton C, Brock KM. 2022. It ain't easy being orange: lizard colour morphs occupying highly vegetated microhabitats suffer greater ectoparasitism. Amphib-reptil 43:287-97.

Beye M, Hasselmann M, Fondrk MK, Page RE, Omholt SW. 2003. The gene *csd* is the primary signal for sexual development in the honeybee and encodes an SR-type protein. Cell

114:419–29.

Billiard S, López-Villavicencio M, Devier B, Hood ME, Fairhead C, Giraud T. 2011. Having sex,
yes, but with whom? Inferences from fungi on the evolution of anisogamy and mating
types. Biol Rev Camb Philos Soc 86:421–42.

Billiard S, López-Villavicencio M, Hood ME, Giraud T. 2012. Sex, outcrossing and mating types:
unsolved questions in fungi and beyond. J Evol Biol 25:1020–38.

Brambilla DJ, Matsumoto AM, Araujo AB, McKinlay JB. 2009. The effect of diurnal variation on
clinical measurement of serum testosterone and other sex hormone levels in men. J Clin
Endocrinol Metab 94:907–13.

Branch HA, Klingler AN, Byers KJRP, Panofsky A, Peers D. 2022. Discussions of the "not so
fit": how ableism limits diverse thought and investigative potential in evolutionary biology.
Am Nat 200:101–13.

Brennan PLR. 2022. Bird genitalia. Curr Biol 32:R1061–62.

Brock KM, Baeckens S, Donihue CM, Martín J, Pafilis P, Edwards DL. 2020. Trait differences
among discrete morphs of a color polymorphic lizard, Podarcis erhardii. PeerJ 8:e10284.

Brock KM, Madden IE. 2022. Morph-specific differences in escape behavior in a color
polymorphic lizard. Behav Ecol Sociobiol 76:104.

Brock KM, Chelini MC, Ayton C, Madden IE, Ramos C. 2022a. Colour morph predicts social
behaviour and contest outcomes in a polymorphic lizard *(Podarcis erhardii)*. Anim Behav
191:91-103.

Brock KM, McTavish EJ, Edwards DL. 2022b. Color polymorphism is a driver of diversification in
the lizard family Lacertidae. Syst Biol 71:24–39.

Brock KM, Indiana Madden, Rosso A, Ramos C, Degen R, Stadler SR, Ayton C, Fernandez
MEL, Servin JR. 2022c. Patterns of colour morph diversity across populations of Aegean
Wall Lizard, *Podarcis erhardii* (Bedriaga, 1882). HNO 15:361–64.

Caballero A. 1994. Developments in the prediction of effective population size. Heredity 73 (Pt
6):657–79.

Cabej NR. 2018. Epigenetic principles of evolution. 2nd edition. Amsterdam: Elsevier Science.

Cadena CD, Zapata F. 2021. The genomic revolution and species delimitation in birds (and
other organisms): why phenotypes should not be overlooked. Ornithology 138:ukaa069.

Campagna L. 2016. Supergenes: the genomic architecture of a bird with four sexes. Curr Biol
26:R102-24.

Capel B. 2017. Vertebrate sex determination: evolutionary plasticity of a fundamental switch.
Nat Rev Genet 18:675–89.

Carcupino M, Baldacci A, Mazzini M, Franzoi P. 2002. Functional significance of the male brood
pouch in the reproductive strategies of pipefishes and seahorses: a morphological and
ultrastructural comparative study on three anatomically different pouches. J Fish Biol
61:1465–80.

Casper AMA, Rebolledo N, Lane AK, Jude L, Eddy SL. 2022. "It's completely erasure": A
qualitative exploration of experiences of transgender, nonbinary, gender nonconforming,
and questioning students in biology courses. CBE Life Sci Educ 21:ar69.

Casto KV, Leininger E, Tan T. 2022. Teaching about sex and gender in neuroscience: more
than meets the "XY." J Undergrad Neurosci Educ 17:A1-3.

Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and patterns
of molecular evolution and variation. Nat Rev Genet 10:195–205.

Charlesworth D, Mank JE. 2010. The birds and the bees and the flowers and the trees: lessons
from genetic mapping of sex determination in plants and animals. Genetics 186:9–31.

Cheung AS, Lim HY, Cook T, Zwickl S, Ginger A, Chiang C, Zajac JD. 2021. Approach to
interpreting common laboratory pathology tests in transgender individuals. J Clin
Endocrinol Metab 106:893–901.

Clutton-Brock TH. 2019. The Evolution of Parental Care. Princeton (NJ): Princeton University

Press.

Corl A, Davis AR, Kuchta SR, Comendant T, Sinervo B. 2010. Alternative mating strategies and
the evolution of sexual size dimorphism in the side-blotched lizard, *Uta stansburiana*: a
population-level comparative analysis. Evolution 64:79–96.

Cote J, Le Galliard JF, Rossi JM, Fitze PS. 2008. Environmentally induced changes in
carotenoid-based coloration of female lizards: a comment on Vercken et al. Journal of
Evolutionary Biology 21:1165–72.

Darwin C. 1871. The descent of man: and selection in relation to sex. London: J. Murray.

Dewsbury DA. 2005. The darwin-bateman paradigm in historical context. Integr Comp Biol
45:831–37.

Dominey WJ. 1980. Female mimicry in male bluegill sunfish—a genetic polymorphism? Nature
284:546–48.

Douglas TE, Strassmann JE, Queller DC. 2016. Sex ratio and gamete size across eastern North
America in Dictyostelium discoideum, a social amoeba with three sexes. J Evol Biol
29:1298–1306.

Dreger AD. 2006. Intersex and human rights: the long view. In: International Library of Ethics,
Law and the New Medicine. Dordrecht: Kluwer Academic Publishers. p. 73–86.

Edward DA, Chapman T. 2011. The evolution and significance of male mate choice. Trends
Ecol Evol 26:647–54.

Eens M, Pinxten R. 2000. Sex-role reversal in vertebrates: behavioural and endocrinological
accounts. Behav Processes 51:135–47.

Eldredge N. 1976. Differential evolutionary rates. Paleobiology 2:174–77.

Emlen ST, Oring LW. 1977. Ecology, sexual selection, and the evolution of mating systems.
Science 197:215–23.

Emlen ST, Wrege PH. 2004. Size dimorphism, intrasexual competition, and sexual selection in
Wattled Jacana (*Jacana jacana*), a sex-role-reversed shorebird in Panama. Auk
121:391–403.

Engqvist L, Taborsky M. 2016. The evolution of genetic and conditional alternative reproductive
tactics. Proc Biol Sci 283:20152945.

Falls JB, Kopachena JG. 2020. White-throated Sparrow (*Zonotrichia albicollis*). Birds of the
World Online.

Fausto-Sterling A. 2012. Sex/gender: biology in a social world. New York City (NY): Routledge.

Fincke OM, Jödicke R, Paulson DR, Schultz TD. 2005. The evolution and frequency of female
color morphs in Holarctic Odonata: why are male-like females typically the minority? Int J
Odonatol 8:183–212.

Fisher RA. 1930. The genetical theory of natural selection. Oxford: Oxford University Press.

Fritzsche K, Henshaw JM, Johnson BD, Jones AG. 2021. The 150th anniversary of The
Descent of Man: Darwin and the impact of sex-role reversal on sexual selection research.
Bio J Linn Soc 134:525–40.

Folwell M, Sanders K, Crowe-Riddell J. 2022a. The squamate clitoris: a review and directions
for future research. Integr Comp Bio 62:559–68.

Folwell MJ, Sanders KL, Brennan PLR, Crowe-Riddell JM. 2022b. First evidence of hemiclitores
in snakes. Proc Roy Soc B 289:20221702.

Ford EB. 1945. Polymorphism. Biol Rev Camb Philos Soc 20:73–88.

Forsman A, Ahnesjö J, Caesar S, Karlsson M. 2008. A model of ecological and evolutionary
consequences of color polymorphism. Ecology 89:34–40.

Furness AI, Capellini I. 2019. The evolution of parental care diversity in amphibians. Nat
Commun 10:4709.

Galeotti P, Rubolini D, Dunn PO, Fasola M. 2003. Colour polymorphism in birds: causes and
functions. J Evol Biol 16:635–46.

Gallesi MM, Sacchi R, Hardersen S. 2015. Does wing shape of andromorph females of

Calopteryx splendens (Harris, 1780) resemble that of males? Int J Odonatol 18:305–15.

Garcia-Sifuentes Y, Maney DL. 2021. Reporting and misreporting of sex differences in the biological sciences. Elife 10.

Gilman CA, Corl A, Sinervo B, Irschick DJ. 2019. Genital morphology associated with mating strategy in the polymorphic lizard, *Uta stansburiana*. J Morphol 280:184–92.

Glickman SE, Cunha GR, Drea CM, Conley AJ, Place NJ. 2006. Mammalian sexual differentiation: lessons from the spotted hyena. Trends Endocrinol Metab 17:349–56.

Gowaty PA, Kim Y-K, Anderson WW. 2012. No evidence of sexual selection in a repetition of Bateman's classic study of *Drosophila melanogaster*. Proc Nat Acad Sci USA 109:11740–45.

Goymann W, Brumm H. 2018. Let's talk about sex - not gender. Bioessays 40:e1800030.

Goymann W, Moore IT, Oliveira RF. 2019. Challenge Hypothesis 2.0: a fresh look at an established idea. Bioscience 69:432–42.

Goymann W, Wingfield JC. 2014. Male-to-female testosterone ratios, dimorphism, and life history—what does it really tell us? Behav Ecol 25:685–99.

Goymann W, Brumm H, Kappeler. 2023. Biological sex is binary, even though there is a rainbow of sex roles. BioEssays 45:2200173.

Graves JL, Kearney M, Barabino G, Malcom S. 2022. Inequality in science and the case for a new agenda. Proc Nat Acad Sci USA 119:e2117831119.

Gray SM, McKinnon JS. 2007. Linking color polymorphism maintenance and speciation. Trends Ecol Evol 22:71–79.

Greives T, Eshleman M, Galante H, Elderbrock E, Deimel C, Hau M. 2021. Early nighttime testosterone peaks are correlated with GnRH-induced testosterone in a diurnal songbird. Gen Comp Endocrinol 312:113861.

Griffiths PE. 2021. What are biological sexes? PhilSci Archive.

Grimstad F, Boskey ER, Taghinia A, Estrada CR, Ganor O. 2021. The role of androgens in clitorophallus development and possible applications to transgender patients. Andrology 9:1719–28.

Gross MR. 1982. Sneakers, satellites and parentals: Polymorphic mating strategies in North American sunfishes. Z Tierpsychol 60:1–26.

Gross MR. 1996. Alternative reproductive strategies and tactics: diversity within sexes. Trends Ecol Evol 11:92–98.

Gross MR. 2005. The evolution of parental care. Q Rev Biol 80:37–45.

Ha NQ, Dworkin SL, Martínez-Patiño MJ, Rogol AD, Rosario AD, Sánchez FJ, Wrynn A, Vilain. 2014. Hurdling over sex? Sport, science, and equity. Arch Sex Behav 43:1035–42.

Hales KG. 2020. Signaling inclusivity in undergraduate biology courses through deliberate framing of genetics topics relevant to gender identity, disability, and race. CBE Life Sci Educ 19:es2.

Hare RM, Simmons LW. 2019. Sexual selection and its evolutionary consequences in female animals. Biol Rev Proc Camb Philos Soc 94:929–56.

Hau M. 2007. Regulation of male traits by testosterone: implications for the evolution of vertebrate life histories. BioEssays 29:133–44.

Hedrick PW, Tuttle EM, Gonser RA. 2018. Negative-assortative mating in the white-throated sparrow. J Hered 109:223–31.

Hoekstra RF. 1990. The evolution of male-female dimorphism: Older than sex? J Genet 69:11–15.

Hofer H, East ML. 1995. Virilized sexual genitalia as adaptations of female spotted hyaenas. Rev Suisse Zool 102.

Holekamp KE. 2006. Spotted hyenas. Current Biology 16:R944–5.

Holleley CE, O'Meally D, Sarre SD, Marshall Graves JA, Ezaz T, Matsubara K, Azad B, Zhang X, Georges A. 2015. Sex reversal triggers the rapid transition from genetic to

temperature-dependent sex. Nature 523:79–82.

Hugall AF, Stuart-Fox D. 2012. Accelerated speciation in colour-polymorphic birds. Nature
485:631–34.

Hurst LD. 1996. Why are there only two sexes? Proceedings of the Royal Society of London
Series B: Biological Sciences 263:415–22.

Huyghe K, Husak JF, Herrel A, Tadić Z, Moore IT, Van Damme R, Vanhooydonck B. 2009.
Relationships between hormones, physiological performance and immunocompetence in
a color-polymorphic lizard species, *Podarcis melisellensis*. Horm Behav 55:488–94.

Iwasa Y, Sasaki A. 1987. Evolution of the number of sexes. Evolution 41:49–65.

Jaffe K. 1996. The dynamics of the evolution of sex: Why the sexes are, in fact, always two?
Interciencia 21:259-67.

Jarne P, Auld JR. 2006. Animals mix it up too: The distribution of self-fertilization among
hermaphroditic animals. Evolution 60:1816–24.

Jukema J, Piersma T. 2006. Permanent female mimics in a lekking shorebird. Biol Lett 2:161–
64.

Just W, Baumstark A, Hameister H, Schreiner B, Reisert I, Hakhverdyan M, Vogel W. 2002. The
sex determination in *Ellobius lutescens* remains bizarre. Cytogenet Genome Res 96:146–
53.

Kalmus H. 1932. Über den Erhaltungswert der phänotypischen (morphologischen) Anisogamie
und die Entstehung der ersten Geschlechtsunterschiede. Biologisches Zentralblatt 52:
716–726.

Kamath A, Velocci B, Wesner A, Chen N, Formica V, Subramaniam B, Rebolleda-Gómez M.
2022. Nature, data, and power: how hegemonies shaped this special section. Am Nat
200:81–88.

Kappeler PM, Benhaiem S, Fichtel C, Fromhage L, Höner OP, Jennions MD, Kaiser S, Krüger
O, Schneider JM, Tuni C, van Schaik J, Goymann W. 2022. Sex roles and sex ratios in
animals. Biol Rev Camb Philos Soc, early view.

Karkazis K. 2019. The misuses of "biological sex." Lancet 394:1898–99.

Kokko H, Jennions MD. 2008. Parental investment, sexual selection and sex ratios. J Evol Biol
21:919–48.

Kothe E. 1996. Tetrapolar fungal mating types: sexes by the thousands. FEMS Microbiol Rev
18:65–87.

Küpper C, Stocks M, Risse JE, Dos Remedios N, Farrell LL, McRae SB, Morgan TC, Karlionova
N, Pinchuk P, Verkuil YI, et al. 2016. A supergene determines highly divergent male
reproductive morphs in the ruff. Nat Genet 48:79–83.

Kuroiwa A, Ishiguchi Y, Yamada F, Shintaro A, Matsuda Y. 2010. The process of a Y-loss event
in an XO/XO mammal, the Ryukyu spiny rat. Chromosoma 119:519–26.

Kutschera U. 2016. Sex versus gender in sea urchins and leeches two centuries after Lamarck
1816. J Marine Sci Res Dev 6:5.

Lank DB, Smith CM, Hanotte O, Burke T, Cooke F. 1995. Genetic polymorphism for alternative
mating behaviour in lekking male ruff *Philomachus pugnax*. Nature 378:59–62.

Lattanzio MS, Miles DB. 2016. Trophic niche divergence among colour morphs that exhibit
alternative mating tactics. R Soc Open Sci 3:150531.

Lee SC, Ni M, Li W, Shertz C, Heitman J. 2010. The evolution of sex: A perspective from the
fungal kingdom. Microbiol Mol Biol Rev 74:298–340.

Leonard JL, Westfall JA, Pearse JS. 2007. Phally polymorphism and reproductive biology in
Ariolimax (Ariolimax) buttoni (Pilsbry and Vanatta, 1896) (Stylommatophora: Arionidae)*.
Malb 23:121–35.

Lipshutz SE, Rosvall KA. 2020a. Testosterone secretion varies in a sex- and stage-specific
manner: Insights on the regulation of competitive traits from a sex-role reversed species.
Gen Comp Endocrinol 292:113444

Lipshutz SE, Rosvall KA. 2020b. Neuroendocrinology of Sex-Role Reversal. Integr Comp Biol
60:692–702.
Lorenzi V, Earley RL, Rodgers EW, Pepper DR, Grober MS. 2008. Diurnal patterns and sex
differences in cortisol, 11-ketotestosterone, testosterone, and 17beta-estradiol in the
bluebanded goby (*Lythrypnus dalli*). Gen Comp Endocrinol 155:438–46.
Maney DL. 2016. Perils and pitfalls of reporting sex differences. Philos Trans R Soc B
371:20150119.
Maney DL, Küpper C. 2022. Supergenes on steroids. Phil Trans R Soc B 377:20200507.
Maney DL, Merritt JR, Prichard MR, Horton BM, Yi SV. 2020. Inside the supergene of the bird
with four sexes. Horm Behav 126:104850.
Mank JE. 2022. Sex-specific morphs: the genetics and evolution of intra-sexual variation. Nat
Rev Genet 1–9.
McLean CA, Stuart-Fox D. 2014. Geographic variation in animal colour polymorphisms and its
role in speciation. Biol Rev Camb Philos Soc 89:860–73.
Miyagi M, Guthman EM, Sun SED-K. 2021. Transgender rights rely on inclusive language.
Science 374:1568–69.
Mokos J, Scheuring I, Liker A, Freckleton RP, Székely T. 2021. Degree of anisogamy is
unrelated to the intensity of sexual selection. Sci Rep 11:19424.
Monk JD, Giglio E, Kamath A, Lambert MR, McDonough CE. 2019. An alternative hypothesis
for the evolution of same-sex sexual behaviour in animals. Nat Ecol Evol 3:1622–31.
Moore EC, Cicotto PJ, Peterson EN, Lamm MS, Albertson RC, Roberts RB. 2022. Polygenic
sex determination produces modular sex polymorphism in an African cichlid fish. Proc
Nat Acad Sci USA 119:e2118574119.
Naranjo-Ortiz MA, Gabaldón T. 2020. Fungal evolution: cellular, genomic, and metabolic
complexity. Biol Rev Cam Phil Soc 95:1198–232.
Oliveira, R. F., Taborsky, M., & Brockmann, H. J. (Eds.). (2008). Alternative reproductive tactics:
an integrative approach. Cambridge: Cambridge University Press.
Packer M, Lambert MR. 2022. What's gender got to do with it? Dismantling the human
hierarchies in evolutionary biology and environmental toxicology for scientific and social
progress. Am Nat 200:114–28.
Panico S, Pisani P, Muti P, Recchione C, Cavalleri A, Totis A, Berrino F. 1990. Diurnal variation
of testosterone and estradiol: a source of bias in comparative studies on breast cancer. J
Endocrinol Invest 13:423–26.
Parker JD. 2004. A major evolutionary transition to more than two sexes? Trends Ecol Evol
19:83–86.
Paulson DR. 1998. Possible morphological and behavioral male mimicry in a libellulid dragonfly,
Erythrodiplax umbrata (L.)(Anisoptera: Libellulidae). Odonatologica 27:249–52.
Pérez i de Lanuza G, Font E, Carazo P. 2013. Color-assortative mating in a color-polymorphic
lacertid lizard. Behav Ecol 24:273–79.
Pérez i de Lanuza G, Carretero MA. 2018. Partial divergence in microhabitat use suggests
environmental-dependent selection on a colour polymorphic lizard. Behav Ecol Sociobiol
72:138.
Pérez i de Lanuza G, Sillero N, Carretero MA. 2018. Climate suggests environment-dependent
selection on lizard colour morphs. J Biogeo 45:2791–802.
Perrin N. 2012. What uses are mating types? The "developmental switch" model. Evolution
66:947–56.
Phiri-Ramongane B, Khine AA. 2022. Role of clinical laboratories in reporting results of
transgender individuals on hormonal therapy. Journal of Endocrinology, Metabolism and
Diabetes of South Africa 27:8–13.
Phoenix CH, Goy RW, Gerall AA, Young WC. 1959. Organizing action of prenatally
administered testosterone propionate on the tissues mediating mating behavior in the

female guinea pig. Endocrinology 65:369–82.

Piferrer F. 2021. Epigenetic mechanisms in sex determination and in the evolutionary transitions
862   between sexual systems. Philos Trans R Soc Lond B Biol Sci 376:20200110.

Pollo P, Kasumovic MM. 2022. Let's talk about sex roles: what affects perceptions of sex
864   differences in animal behaviour? Anim Behav 183:1–12.

Power HW. 1976. On forces of selection in the evolution of mating types. Am Nat 110:937–44.

Quigley CA, De Bellis A, Marschke KB, el-Awady MK, Wilson EM, French FS. 1995. Androgen
868   receptor defects: historical, clinical, and molecular perspectives. Endocr Rev 16:271–321.

Quinn AE, Radder RS, Sarre SD, Georges A, Ezaz T, Shine R. 2009. Isolation and
870   development of a molecular sex marker for *Bassiana duperreyi*, a lizard with XX/XY sex
871   chromosomes and temperature-induced sex reversal. Mol Genet Genomics 281:665–72.

Rosvall KA, Bergeon Burns CM, Barske J, Goodson JL, Schlinger BA, Sengelaub DR,
873   Ketterson ED. 2012. Neural sensitivity to sex steroids predicts individual differences in
874   aggression: implications for behavioural evolution. Proc Biol Sci 279:3547–55.

Roughgarden J. 2013. Evolution's rainbow: diversity, gender, and sexuality in nature and
876   people. 2nd edition. Berkeley (CA): University of California Press.

Roulin A. 2004. The evolution, maintenance and adaptive function of genetic colour
878   polymorphism in birds. Biol Rev Camb Philos Soc 79:815–48.

Russell DGD, Sladen WJL, Ainley DG. 2012. Dr. George Murray Levick (1876–1956):
880   unpublished notes on the sexual habits of the Adélie penguin. Polar Rec 48:387–93.

Saldanha CJ, Remage-Healey L, Schlinger BA. 2011. Synaptocrine signaling: steroid synthesis
882   and action at the synapse. Endocr Rev 32:532–49.

Sandkam BA, Almeida P, Darolti I, Furman BLS, van der Bijl W, Morris J, Bourne GR, Breden F,
884   Mank JE. 2021. Extreme Y chromosome polymorphism corresponds to five male
885   reproductive morphs of a freshwater fish. Nat Ecol Evol 5:939–48.

Sarre SD, Georges A, Quinn A. 2004. The ends of a continuum: genetic and temperature-
887   dependent sex determination in reptiles. Bioessays 26:639–45.

Saunders PA, Perez J, Rahmoun M, Ronce O, Crochet P-A, Veyrunes F. 2014. XY females do
889   better than the XX in the African pygmy mouse, *Mus minutoides*. Evolution 68:2119–27.

Schmidt KL, Pradhan DS, Shah AH, Charlier TD, Chin EH, Soma KK. 2008. Neurosteroids,
891   immunosteroids, and the Balkanization of endocrinology. Gen Comp Endocrinol 157:266–
892   74.

Seehausen O, Terai Y, Magalhaes IS, Carleton KL, Mrosso HDJ, Miyagi R, van der Sluijs I,
894   Schneider MV, Maan ME, Tachida H, et al. 2008. Speciation through sensory drive in
895   cichlid fish. Nature 455:620–26.

Ser JR, Roberts RB, Kocher TD. 2010. Multiple interacting loci control sex determination in lake
897   Malawi cichlid fish. Evolution 64:486–501.

Sinervo B, Lively CM. 1996. The rock–paper–scissors game and the evolution of alternative
899   male strategies. Nature 380:240–43.

Sinervo B, Svensson E, Comendant T. 2000. Density cycles and an offspring quantity and
901   quality game driven by natural selection. Nature 406:985–88.

Sinervo B, Bleay C, Adamopulou C. 2001. Social causes of correlational selection and the
903   resolution of a heritable throat color polymorphism in a lizard. Evolution 55:2040–52.

Smith CA, Roeszler KN, Ohnesorg T, Cummins DM, Farlie PG, Doran TJ, Sinclair AH. 2009.
905   The avian Z-linked gene *DMRT1* is required for male sex determination in the chicken.
906   Nature 461:267–71.

Staub NL, De Beer M. 1997. The role of androgens in female vertebrates. Gen Comp
908   Endocrinol 108:1–24.

Stévant I, Papaioannou MD, Nef S. 2018. A brief history of sex determination. Mol Cell
910   Endocrinol 468:3–10.

Stiver KA, Harris RM, Townsend JP, Hofmann HA, Alonzo SH. 2015. Neural gene expression

profiles and androgen levels underlie alternative reproductive tactics in the ocellated
wrasse, *Symphodus ocellatus*. Ethology 121:152–67.
Stuart-Fox D, Aulsebrook A, Rankin KJ, Dong CM, McLean CA. 2021. Convergence and
divergence in lizard colour polymorphisms. Biol Rev Camb Philos Soc 96:289–309.
Sutou S, Mitsui Y, Tsuchiya K. 2001. Sex determination without the Y chromosome in two
Japanese rodents *Tokudaia osimensis osimensis* and *Tokudaia osimensis spp*. Mamm
Genome 12:17–21.
Tang-Martínez Z. 2016. Rethinking Bateman's Principles: challenging persistent myths of
sexually reluctant females and promiscuous males. J Sex Res 53:532–59.
Tang-Martínez Z. 2020. The history and impact of women in animal behaviour and the ABS: a
North American perspective. Anim Behav 164:251–60.
Thompson A, Kapsanaki V, Liwanag HEM, Pafilis P, Wang IJ, Brock KM. 2023. Some like it
hotter: differential thermal preferences among lizard color morphs. J Therm Bio
113:103532.
Thorne N, Yip AK-T, Bouman WP, Marshall E, Arcelus J. 2019. The terminology of identities
between, outside and beyond the gender binary - a systematic review. Int J Transgend
20:138–54.
Tillmann U, Hoppenrath M. 2013. Life cycle of the pseudocolonial dinoflagellate *Polykrikos*
kofoidii (Gymnodiniales, Dinoflagellata). J Phycol 49:298–317.
Togashi T, Cox PA. 2011. The Evolution of Anisogamy: A Fundamental Phenomenon
Underlying Sexual Selection. Cambridge University Press.
Townsend DS, Stewart MM, Pough FH. 1984. Male parental care and its adaptive significance
in a neotropical frog. Anim Behav 32:421–31.
Trivers RL. 1972. Parental investment and sexual selection In: Campbell B, editor. Sexual
selection and the descent of man, 1871–1971. Chicago (IL): Aldine.
Tucker BW. 1943. Brood-patches and the physiology of incubation. Br Birds 37:22–8.
Tuttle EM, Bergland AO, Korody ML, Brewer MS, Newhouse DJ, Minx P, Stager M, Betuel A,
Cheviron ZA, Warren WC, et al. 2016. Divergence and functional degradation of a sex
chromosome-like supergene. Curr Biol 26:344–50.
Vaiman D, Pailhoux E. 2000. Mammalian sex reversal and intersexuality: deciphering the sex-
determination cascade. Trends Genet 16:488–94.
Van Leeuwen AM, Bladh ML. 2016. Textbook of laboratory and diagnostic testing: practical
application of nursing process at the bedside. Philadelphia (PA): F.A. Davis.
Verhulst EC, Beukeboom LW, van de Zande L. 2010. Maternal control of haplodiploid sex
determination in the wasp *Nasonia*. Science 328:620–23.
Veyrunes F, Chevret P, Catalan J, Castiglia R, Watson J, Dobigny G, Robinson TJ, Britton-
Davidian J. 2010. A novel sex determination system in a close relative of the house
mouse. Proc Biol Sci 277:1049–56.
Walters MJ, Collado D, Harding CF. 1991. Oestrogenic modulation of singing in male zebra
finches: differential effects on directed and undirected songs. Anim Behav 42:445–52.
West-Eberhard MJ. 1986. Alternative adaptations, speciation, and phylogeny (a review). Proc
Natl Acad Sci USA 83:1388–92.
Whiteley SL, Holleley CE, Ruscoe WA, Castelli M, Whitehead DL, Lei J, Georges A,
Weisbecker V. 2017. Sex determination mode does not affect body or genital
development of the central bearded dragon (*Pogona vitticeps*). Evodevo 8:25.
Whitfield J. 2004. Everything you always wanted to know about sexes. PLoS Biol 2:e183.
Wiggins JM, Santoyo-Brito E, Scales JB, Fox SF. 2020. Gene dose indicates presence of sex
chromosomes in collared lizards (*Crotaphytus collaris*), a species with temperature-
influenced sex determination. Herp 76:27–30.
Wingfield JC, Goymann W, Jalabert C, Soma KK. 2019. Concepts derived from the Challenge
Hypothesis. Horm Behav 115:104550.

Wingfield JC, Hegner RE, Dufty AM, Ball GF. 1990. The "Challenge Hypothesis": theoretical
implications for patterns of testosterone secretion, mating systems, and breeding
strategies. Am Nat 136:829–46.
Woitowich NC, Beery A, Woodruff T. 2020. A 10-year follow-up study of sex inclusion in the
biological sciences. Elife 9.
Yoshizawa K, Ferreira RL, Kamimura Y, Lienhard C. 2014. Female penis, male vagina, and
their correlated evolution in a cave insect. Curr Biol 24:1006–10.
Zhao L, Quinn A, Ng ET, Veyrunes F, Koopman P. 2017. Reduced activity of *SRY* and its target
enhancer *Sox9-TESCO* in a mouse species with X*Y sex reversal. Sci Rep 7:41378.
972