Title: *Pseudomonas aeruginosa* Siderophores Damage Lung Epithelial Cells and Promote

Inflammation

Authors: Donghoon Kang, Natalia V. Kirienko

Author Affiliations: Department of BioSciences, Rice University, Houston, TX, USA

Corresponding Author: Natalia V. Kirienko; kirienko@rice.edu

Abstract:

Multidrug-resistant *Pseudomonas aeruginosa* is a common nosocomial respiratory pathogen

that continues to threaten the lives of mechanically-ventilated patients in intensive care units and

those with underlying comorbidities such as cystic fibrosis or chronic obstructive pulmonary

disease. For over 20 years, studies have repeatedly demonstrated that the major siderophore

pyoverdine is an important virulence factor for *P. aeruginosa* in invertebrate and mammalian hosts

in vivo. Despite its physiological significance, an *in vitro,* mammalian cell culture model to

characterize the impact and the molecular mechanism of pyoverdine during infection has only

recently been developed. In this study, we adapt a previously-established murine macrophage-

based model for human bronchial epithelial cells. We demonstrate that pyoverdine-rich

conditioned medium from *P. aeruginosa* disrupts epithelial integrity in a manner that depends on

protease activity and the type II secretion system. Disrupting pyoverdine production, whether

genetically or chemically, mitigates this damage. Interestingly, this damage did not require

exotoxin A or PrpL (protease IV), two previously-characterized toxins regulated by pyoverdine.

We also examined the effects of exposure to purified pyoverdine on lung epithelial cells. While

pyoverdine accumulates within cells, the siderophore is largely sequestered inside early

endosomes, showing little cytotoxicity. This is in contrast to other, more membrane-permeable

iron chelators and siderophores such as pyochelin. However, pyoverdine may indirectly contribute

to lung inflammation by potentiating these iron chelators in promoting the production of

proinflammatory cytokines.

Introduction:

Multidrug-resistant *Pseudomonas aeruginosa* is one of the most common respiratory

pathogens infecting patients in intensive care units, especially those who are mechanically-

ventilated or have chronic airway disorders like cystic fibrosis (CF) or chronic obstructive

pulmonary disease (COPD) (Bhagirath et al., 2016; Hassett et al., 2014; Kollef et al., 2014; Lyczak

et al., 2002; Murphy, 2009). This pathogen's intrinsic resistance to several classes of antibiotics

and exceptional ability to form biofilms on medical devices and airway tissue pose a serious

challenge for medical intervention (Anderson and O'Toole, 2008; Moreau-Marquis et al., 2008).

In addition to colonizing the respiratory tract, *P. aeruginosa* actively deploys numerous virulence

factors and toxins to damage host tissue, compromising pulmonary function (Curran et al., 2018).

One of the major virulence factors produced by this pathogen is the siderophore pyoverdine.

Several groups have proposed possible mechanisms of pyoverdine-dependent virulence during

lung infection. First and foremost, as a siderophore, pyoverdine scavenges ferric iron, providing a

crucial nutritional need. Generally, iron acquisition serves an important function during *P.*

aeruginosa infection by promoting bacterial growth and biofilm formation (Banin et al., 2005;

Kang and Kirienko, 2018), and *P. aeruginosa* mutants lacking various iron uptake systems exhibit

attenuation of virulence during murine lung infection (Minandri et al., 2016). Pyoverdine-mediated

iron uptake further promotes *P. aeruginosa* virulence by derepressing the alternative sigma factor

PvdS, which activates the transcription of several virulence genes such as those encoding the

translational inhibitor exotoxin A, exoprotease PrpL (protease IV), and pyoverdine biosynthetic

enzymes (Lamont et al., 2002; Visca et al., 2007).

Of the two major siderophores secreted by *P. aeruginosa*, pyoverdine and pyochelin,

pyoverdine is distinct in its ability to directly chelate iron from host ferroproteins such as

transferrin and lactoferrin (Dumas et al., 2013; Xiao and Kisaalita, 1997). We have recently used

a *Caenorhabditis elegans* nematode host model to demonstrate that pyoverdine may also directly

chelate intracellular iron, disrupting mitochondrial homeostasis (Kang et al., 2018; Kirienko et al.,

2015; Kirienko et al., 2013). Some combination of these various pathogenic functions makes

pyoverdine production necessary for full *P. aeruginosa* virulence against murine hosts during

acute lung infection (Imperi et al., 2013; Kang et al., 2019; Meyer et al., 1996; Minandri et al.,

2016; Takase et al., 2000).

Recently, we established the first-reported *in vitro* cell culture model for pyoverdine-dependent

virulence, where murine macrophages are treated with conditioned medium (also referred to as

bacterial filtrate) from *P. aeruginosa* grown in serum-free cell culture medium (Kang and Kirienko,

2020). Under these conditions, *P. aeruginosa* exhibits robust pyoverdine production, yet the

siderophore is not required for bacterial growth (**Figure 1A, B**), allowing pyoverdine's distinct

role in virulence outside of its role in iron acquisition to be studied. This pyoverdine-rich

conditioned medium from wild-type *P. aeruginosa* PAO1 is cytotoxic towards murine

macrophages and murine alveolar macrophages (**Figure 1C**), and in clinical isolates, pyoverdine

content in conditioned medium positively correlates with cytotoxicity (Kang and Kirienko, 2020).

In this report, we adapt this *in vitro* pyoverdine virulence model for human bronchial epithelial

cells to examine the consequences of pyoverdine production (i.e., pyoverdine and known

pyoverdine-regulated virulence factors) during *P. aeruginosa* lung infection. We demonstrate that

pyoverdine-rich conditioned medium from *P. aeruginosa* disrupts epithelial integrity in a protease-

dependent manner. Genetic or chemical disruption of pyoverdine production mitigates this damage.

Interestingly, this phenomenon depends on type II secretion but is independent of exotoxin A or

PrpL production. We also examine the effects of purified pyoverdine exposure in lung epithelial

cells. While pyoverdine accumulates within cells, the siderophore is largely sequestered within

early endosomes and exhibits low cytotoxicity, in contrast to other more membrane-permeable

iron chelators and siderophores such as pyochelin. However, we also demonstrate that pyoverdine

may indirectly affect cell viability and promote neutrophilic inflammation in the presence of these

chelators.

Results:

Pyoverdine-rich conditioned medium disrupts epithelial integrity

To investigate the role of pyoverdine production during *P. aeruginosa* lung infection, we

treated human bronchial epithelial cells (16HBE) with pyoverdine-rich conditioned medium from

P. aeruginosa PAO1 grown in serum-free cell growth medium (Eagle's Minimum Essential

Medium). To visualize the integrity of the epithelial monolayer, we prelabeled cells with a plasma

membrane stain. Within 1 h, the conditioned medium severely damaged the monolayer, causing

more than half the cells to detach **(Figure 1D)**. This disruption was significantly attenuated in

16HBE cells treated with identically-prepared material from an isogenic pyoverdine biosynthetic

mutant (PAO1*ΔpvdF*) but not a pyochelin mutant (PAO1*ΔpchBA*) **(Figure 1D, E)**. In the absence

of pyoverdine production, preventing pyochelin biosynthesis ($\Delta pvdF\Delta pchBA$) did not confer

further protection to 16HBE cells. One difference between these two siderophores is their affinity

for ferric iron. Due to an exceptionally high affinity for the metal, pyoverdine is uniquely able to

remove iron from host ferroproteins (Dumas et al., 2013; Xiao and Kisaalita, 1997). We examined

whether host iron chelation was important for the damage we observed in the epithelial monolayer.

To hinder pyoverdine's ability to bind iron, we pretreated the conditioned medium with a ferric

iron mimetic, gallium (Ga^{3+}). However, even with the addition of excess gallium, there was no

significant rescue **(Figure S1)**. Interestingly, we observed that removing large macromolecules

greater than 10 kDa via centrifugal filtration or supplementing the conditioned medium with fetal

bovine serum (FBS) abrogated the damage to the epithelial monolayer **(Figure S1)**. A common

target for these two methods of intervention would be secreted *P. aeruginosa* proteases. Consistent

with this hypothesis, supplementing the conditioned medium with a commercially-sourced

protease inhibitor cocktail (capable of inhibiting serine and cysteine proteases) significantly

lessened cell detachment **(Figure 1F, G, Figure S1)**.

type II secretion is important for P. aeruginosa virulence against 16HBE cells

Based on these results, we posited that the pyoverdine-regulated serine protease PrpL (protease

IV) was primarily responsible for the damage we observed in 16HBE cells. To test this hypothesis,

we prepared conditioned medium from an array of *P. aeruginosa* MPAO1 transposon mutants

carrying insertions in *toxA* (exotoxin A), *prpL*, *xcpP*, or *xcpQ*, the latter two encoding proteins in

the Xcp type II secretion system that is responsible for the secretion of exotoxin A and PrpL

(Filloux, 2011). Due to a lack of a suitable wild-type strains, these transposon mutants were

compared to those carrying mutations in antibiotic resistance genes (chloramphenicol

acetyltransferase *cat* and cephalosporinase *ampC*) that were irrelevant to our phenomenon of

interest. Conditioned medium from neither *toxA* nor *prpL* mutant resulted in significant attenuation

of cell detachment **(Figure 2A, B)**. However, we observed significant rescue in 16HBE cells

treated with conditioned medium from the two type II secretion mutants **(Figure 2A, B)**,

suggesting that this secretion system may transport additional pyoverdine-regulated proteases and

other toxins to damage epithelial cells.

To further demonstrate the importance of pyoverdine production in *P. aeruginosa* virulence

against lung epithelial cells, we took advantage of two clinical strains of *P. aeruginosa* isolated

from pediatric CF patients, PA2-9 and PA3-29. These two strains were selected from a collection

of CF isolates on the basis of their high *in vitro* pyoverdine production and virulence against the

nematode host *C. elegans* **(Figure S2)** (Kang et al., 2019). These two isolates were also amenable

to genetic manipulation, allowing us to generate mutants with transposon insertions in pyoverdine

biosynthetic enzymes **(Figure 2C, D, Table 1)**. As expected, 16HBE cells treated with conditioned

medium from wild-type PA2-9 exhibited severe disruption to the epithelial monolayer **(Figure**

2E). This disruption was attenuated in the pyoverdine biosynthetic mutant (PA2-9 Tn) **(Figure 2E,**

F). Importantly, we were also able to accomplish significant rescue by growing wild-type PA2-9

in the presence of the pyoverdine biosynthetic inhibitor 5-fluorocytosine (5-FC) at concentrations

where 5-FC does not affect bacterial growth **(Figure 2C, E, F)**. Notably, 5-FC has been

consistently shown to rescue various hosts from pyoverdine-dependent virulence (Costabile et al.,

2016; Imperi et al., 2013; Kang et al., 2019; Kang et al., 2021a; Kirienko et al., 2016).

In contrast, we did not observe any discernible damage to the epithelial monolayer by

conditioned medium from PA3-29, even though the strain exhibited comparable levels of

pyoverdine production and bacterial growth to that of PA2-9 **(Figure 2C-F)**. Whole genome

sequence analysis of PA3-29 revealed a 9 base-pair deletion in the Xcp type II secretion system

gene XcpW. This is consistent with our observation that the type II secretion system is important

for the disruption of epithelial integrity by *P. aeruginosa*.

Pyoverdine translocates into 16HBE cells but is sequestered within early endosomes

Next, we wanted to investigate the consequences of exposing 16HBE cells to pyoverdine in

the absence of other virulence factors. In brief, pyoverdine-rich bacterial filtrate was subjected to

two purification steps to separate small molecules by polarity: absorption and elution from a

nonpolar polymeric resin (Amberlite XAD-4) and high-performance liquid chromatography

(HPLC) via a C-18 reverse-phase column **(Figure S3)**. We tested whether this purified material

was cytotoxic to 16HBE cells using a resazurin-based cell viability assay and compared its toxicity

to other known iron-chelating molecules, including the ferric iron chelator ciclopirox olamine, the

ferrous iron chelator 1, 10-phenanthroline, and deferoxamine, a siderophore produced by

Streptomyces spp. While all other iron-chelators exhibited time- and dose-dependent cytotoxicity

towards 16HBE cells, pyoverdine remained largely nontoxic even after 72 h treatment at 200 µM

(Figure 3A, Figure S4A, B).

Because pyoverdine is considerably larger than these iron-chelators, with a molecular weight

of ~1365 g/mol, it may be unable to cross cellular membranes. We examined whether pyoverdine

can translocate into 16HBE cells by taking advantage of its intrinsic spectral properties. After 24

h, 16HBE cells treated with purified pyoverdine accumulated pyoverdine **(Figure 3B, C)**.

Consistent with previous studies, this intracellular fluorescence was enhanced when pyoverdine

was pre-saturated with gallium and was quenched when pyoverdine was pre-saturated with iron

(Figure 3B, C) (Kang et al., 2018; Kang and Kirienko, 2020). Importantly, pyoverdine

fluorescence did not colocalize with a plasma membrane stain, indicating that pyoverdine was

primarily localized to the cytosol **(Figure S4C)**. However, we also observed that pyoverdine

fluorescence formed distinct punctae within the cell. Based on previous observations in murine

macrophages (Kang and Kirienko, 2020), we hypothesized that pyoverdine was sequestered within

early endosomes. Supporting this hypothesis, pyoverdine colocalized with fluorophore-conjugated

dextran, a well-established endosomal marker **(Figure 3D, Figure S4D)** (Oliver et al., 1984).

To investigate whether there was an endocytosis-independent route for pyoverdine

internalization, we utilized giant plasma membrane vesicles (GPMVs) derived from 16HBE cells.

We anticipated that these vesicles lacked the cellular machinery for complex molecular transport.

Remarkably, these vesicles exhibited substantial accumulation of pyoverdine fluorescence

following treatment. In contrast to the punctae observed in living cells, this fluorescence was

evenly distributed throughout the vesicle **(Figure S5A-C)**. Pyoverdine within these vesicles

remained fluorescent in the presence of excess ferric iron, unlike pyoverdine in solution **(Figure**

S5A). However, FITC-conjugated dextran showed a similar distribution as pyoverdine in GPMVs

(Figure S5D), suggesting that GPMVs may be more permeable to macromolecules than intact

cells, as previously suggested by others (Skinkle et al., 2020; Zemljic Jokhadar et al., 2018).

Altogether, while we were able to detect accumulation of intracellular pyoverdine in 16HBE cells,

it appears to be sequestered within early endosomes, and it remains unclear whether pyoverdine

can translocate across nonporous cellular membranes. These results are largely consistent with our

observations that pyoverdine, unlike other iron-chelating molecules, exhibits low cytotoxicity

towards 16HBE cells.

Intracellular iron chelation activates an inflammatory response in 16HBE cells

While pyochelin exhibits lower affinity towards ferric iron than pyoverdine, it is also

substantially smaller, with a molecular weight of ~325 g/mol. We hypothesized that pyochelin

may be able to enter 16HBE cells and chelate intracellular iron. While we were not able to visualize

pyochelin internalization into cells (due to its lack of distinct spectral properties), one consequence

of iron deprivation in epithelial cells would be a proinflammatory transcriptional response. Several

studies have demonstrated that iron chelation by various siderophores such as deferoxamine and

enterobactin promotes the production of proinflammatory cytokines, most notably interleukin

(IL)-8 in lung epithelial cells, intestinal epithelial cells, and oral keratinocytes (Choi et al., 2004;

Holden et al., 2014; Lee et al., 2007). To reaffirm these findings, we treated 16HBE cells with

ciclopirox olamine (CPX), 1,10-phenanthroline (PHE), pyochelin (PCH), pyoverdine (PVD), or

deferoxamine (DFO) and measured the mRNA levels of genes involved in neutrophilic

inflammation. We first observed that total RNA yield (from phenol-chloroform extraction) in these

cells corresponded with the resazurin-based cell viability assay **(Figure 3A, Figure S4A, B)**. Cells

treated with cytotoxic iron chelators such as ciclopirox olamine, phenanthroline, and deferoxamine

yielded substantially lower quantities of RNA, while cells treated with pyoverdine had quantities

comparable to that of media control **(Figure 4A)**. Notably, RNA yield in cells treated with

commercially-sourced pyochelin resembled that of ciclopirox olamine and phenanthroline. By

qRT-PCR, we measured the expression of genes encoding components of the inflammasome,

namely *NLRP3* and *NLRP1*, and those encoding the major proinflammatory cytokines produced

by lung epithelial cells IL-1β (*IL1B*), IL-8 (*IL8*), and tumor necrosis factor alpha (*TNF*). Notably,

all these genes have been associated with inflammation during lung infection (Moldoveanu et al.,

2009). With the exception of pyoverdine, all iron chelators induced the expression of these

proinflammatory genes **(Figure 4B)**. To ensure that this phenomenon was due to intracellular iron

chelation, we repeated the experiment with pyochelin and deferoxamine pre-saturated with excess

gallium (1:2 stoichiometric ratio). Cells treated with gallium-bound pyochelin or deferoxamine

exhibited RNA yields comparable to that of media control, suggesting that gallium inhibited the

cytotoxic effects of the siderophores **(Figure 4C)**. Furthermore, pretreating the siderophores with

gallium resulted in a significant decrease in proinflammatory gene expression **(Figure 4D)**,

demonstrating that the chelator-induced inflammatory response was due to iron chelation rather

than other nonspecific reactions or contaminants in the commercially-sourced material.

Finally, we investigated if pyoverdine can indirectly promote lung inflammation by

potentiating other iron chelating molecules. Due to its exceptionally high affinity for iron,

pyoverdine is likely to remove iron from other more cell-permeable siderophores or outcompete

them for trace iron in the extracellular milieu, increasing the pool of apo-siderophores to disrupt

host cell homeostasis and promote inflammation. To test this hypothesis, we treated 16HBE cells

with deferoxamine, pyoverdine, or both. Cells treated with both siderophores exhibited greater cell

death and increased expression of proinflammatory genes compared to those treated with

deferoxamine alone **(Figure 4E, F)**. Considering that pyoverdine alone did not affect cell viability

or transcription of proinflammatory genes, these results suggest that pyoverdine was able to

potentiate deferoxamine-mediated damage by increasing the labile pool of apo-deferoxamine.

Discussion:

Arguably one of the greatest challenges to combating *P. aeruginosa* infections is the sheer

multitude of virulence factors the bacterium can utilize to either directly or indirectly cause host

damage. These include small molecule virulence factors (e.g., siderophores, quorum-sensing

molecules), small molecule toxins (e.g., pyocyanin), factors involved in biofilm formation and

motility (e.g, exopolysaccharides, type IV pili, flagella), and more than twenty proteinaceous

toxins that either directly kill host cells (e.g., exotoxin A, exoenzyme S, exotoxin T, exotoxin U)

or damage host tissue (e.g., elastase LasA, elastase LasB, PrpL/protease IV, alkaline protease)

(Hall et al., 2016; Hauser, 2009; Lee and Zhang, 2015; Michalska and Wolf, 2015; Thi et al., 2020).

This complexity casts a dark shadow over the prospects of epidemiological or therapeutic

intervention. In an ideal situation, we would be able to reliably predict a pathogen's ability to cause

disease through our evolving molecular surveillance tools such as whole genome sequencing and

mass spectrometry and therapeutically impair pathogenesis through anti-virulence drugs that

inhibit the production or function of key virulence factors and toxins. In *P. aeruginosa*, the only

feasible way to approach these strategies would be to unravel the regulation of virulence factors in

the bacterium and to target virulence networks rather than individual factors.

The results we report in this study support the idea of targeting the alternative sigma factor

PvdS, as has been investigated by others (Imperi reference). PvdS regulates the production of

several secreted toxins such as the translational inhibitor exotoxin A and secreted protease PrpL.

Exotoxin A, arguably one of the most extensively studied toxins in *P. aeruginosa*, inhibits protein

synthesis (Michalska and Wolf, 2015), inducing airway epithelial cell death (Plotkowski et al.,

2002) and potentially inhibiting cell junction repair in the presence of *P. aeruginosa* elastase

(Azghani, 1996). Exotoxin A also contributes to *P. aeruginosa* virulence in various murine

infection models (Hirakata et al., 1993; Miyazaki et al., 1995; Pillar and Hobden, 2002).

Importantly, Ochsner and colleagues have demonstrated that *P. aeruginosa* exhibits a > 95%

decrease in exotoxin A production in the absence of PvdS (Ochsner et al., 1996). PrpL contributes

to the bacterium's overall extracellular proteolytic activity that is responsible for the responsible

for the degradation of extracellular matrix and cell junction proteins and host defense proteins such

as surfactant proteins that contribute to innate immunity during *P. aeruginosa* lung infections

(Malloy et al., 2005). Furthermore, PrpL has been shown to distinctly contribute to *P. aeruginosa*

virulence during ocular infections (Engel et al., 1998; Engel et al., 1997) and degrade IL-22 to

subvert lung epithelial immune response (Bradshaw et al., 2018; Guillon et al., 2017).

Of course, PvdS is best known for its role in pyoverdine biosynthesis and is indispensable for

the production of pyoverdine biosynthetic enzymes. In addition to scavenging trace iron in the

environment or directly from host ferroproteins, pyoverdine is involved in a positive feedback loop

where the uptake of iron-bound pyoverdine by its outer membrane receptor derepresses PvdS by

the inner membrane protein FpvR, increasing the production of pyoverdine, exotoxin A, and PrpL

(Lamont et al., 2002; Visca et al., 2007). We have demonstrated here that pyoverdine production

promotes the production of an additional secreted toxin, possibly a protease secreted by the type

II secretion system, to damage the lung epithelium. Pyoverdine may also indirectly contribute to

neutrophilic inflammation by removing iron from other more cell permeable siderophores such as

pyochelin and enterobactin, the latter in the context of polymicrobial infections with

Enterobacteriaceae such as the respiratory pathogen *K. pneumoniae* (Holden et al., 2016).

Importantly, while *P. aeruginosa* may lose the ability to produce pyoverdine during lung infection

with the emergence of social cheaters or due to a transition in iron acquisition strategy (Andersen

et al., 2015; De Vos et al., 2001; Marvig et al., 2014), several surveys of patient sputum samples

and clinical isolates have revealed that a large fraction of strains still exhibit substantial pyoverdine

production (Haas et al., 1991; Kang et al., 2019; Martin et al., 2011; Mayer-Hamblett et al., 2014),

demonstrating that pyoverdine may still represent an important target for therapeutic intervention.

Fortunately, the FDA-approved antimycotic drug, 5-fluorocytosine (5-FC) inhibits *pvdS*

expression in *P. aeruginosa* and attenuates virulence during murine lung infection. Imperi and

colleagues first identified 5-FC in a screen for small molecules that inhibit pyoverdine production

(Imperi et al., 2013). We have independently identified a chemical analogue of 5-FC, 5-

fluorouracil – another inhibitor of *pvdS* inhibitor (Ueda et al., 2009), in a small molecule screen

for compounds that rescue *C. elegans* from *P. aeruginosa* in a pyoverdine-dependent pathogenesis

model (Kang et al., 2021b; Kirienko et al., 2016). We have also previously reported that 5-FC

synergizes with gallium nitrate, another FDA-approved drug, to inhibit *P. aerugin*osa growth and

virulence against *C. elegans* (Kang et al., 2021a). Our findings in this study suggest that in addition

to its bactericidal and biofilm-inhibitory activities (Goss et al., 2018; Kaneko et al., 2007), gallium

may also function as an anti-inflammatory agent during lung infection by inhibiting intracellular

iron chelation by pyochelin.

The benefits of suppressing neutrophilic inflammation during lung infection, particularly

chronic lung infection, has been well documented. While the mechanisms employed by neutrophils

to kill and remove pathogens, such as the production of neutrophil elastases, are important for host

defense, they also cause host tissue damage by degrading extracellular matrix proteins (Kruger et

al., 2015; Twigg et al., 2015). During chronic infections (such as those in CF patients), these host

defense factors continue to cause airway damage while the pathogen persists, exacerbating

pulmonary function decline (Cantin et al., 2015). While lung inflammation is mediated by many

factors in CF patients, a strategy to specifically inhibit the NLRP3 inflammasome by therapeutics

such as MCC950 is currently being investigated. This approach has shown promising results in

murine infection studies (Hosseinian et al., 2015; McElvaney et al., 2019). NLRP3 inflammasomes

have been shown to be activated by intracellular iron chelation in lung epithelial cells. We observed

this transcriptional response not only in wild-type 16HBE cells but also those carrying mutations

in the cystic fibrosis transmembrane conductance regulator (CFTR G551D, CFTR ΔF508), two of

the most frequently identified mutations in CF patients **(Figure S6)** (Estivill et al., 1997). Although

gallium has been broadly associated with anti-inflammatory properties (Apseloff, 1999; de

Albuquerque Wanderley Sales et al., 2021; Zhang et al., 2022), it has not been investigated for

inhibiting pathogen-associated inflammation. Based on recent revelations that bacterial

siderophores promote inflammation (Choi et al., 2004; Holden et al., 2016; Holden et al., 2014;

Lee et al., 2007), this therapeutic avenue may merit consideration.

Materials and Methods:

Bacterial Strains and Growth Conditions

P. aeruginosa PAO1 and siderophore biosynthetic mutants were provided by Dr. Dieter Haas.

The CF isolates PA2-9 and PA3-29 were provided by Dr. Carolyn Cannon (Kang et al., 2019).

MPAO1 transposon mutants were from a commercially available library distributed by the

University of Washington (Jacobs et al., 2003).

To produce pyoverdine-rich conditioned medium, a LB overnight culture of *P. aeruginosa* was

diluted 20-fold into 2 mL of serum-free Eagle's Minimum Essential Medium in a 6-well plate. The

plate was sealed with a Breath-Easy sealing membrane (Diversified Biotech, Dedham, MA) and

grown statically at 37 °C for 16 h. Pyoverdine production (Ex. 405 nm; Em. 460 nm) and bacterial

growth (OD 600nm) were measured spectrophotometrically. Bacteria was then removed by

centrifugation and the supernatant was treated with an antibiotic cocktail to kill residual bacteria

(100 µg/mL amikacin, 100 µg/mL gentamicin, 100 µg/mL tobramycin).

Cell Culture

16HBE human bronchial epithelial cells were provided by Dr. Carolyn Cannon (Shah et al.,

2020). 16HBE wild-type and CFTR mutant cell lines were provided by Dr. Gang Bao. All 16HBE

cells were passaged in Eagle's Minimum Essential Medium (EMEM) (MilliporeSigma, Burlington,

MA) supplemented with 10% fetal bovine serum, penicillin/streptomycin, and MEM non-essential

amino acids.

For experiments with *P. aeruginosa* conditioned medium, 4×10^6 cells were seeded into each

well of a collagen-coated 12-well plate and grown at 37 °C for 24 h in a CO_2-jacketed incubator

by which they reached 100% confluence. To visualize the epithelial monolayer, cells were stained

with 2.5 µg/mL CellMask Orange plasma membrane stain (Invitrogen, Carlsbad, CA) for 1 h prior

to conditioned medium exposure. Following treatment, the medium was aspirated and the

monolayer was imaged on a Cytation5 Multimode Reader (Biotek, Winnoski, VT) using a RFP

filter cube. Percentage image area covered by fluorescent cells was quantified using ImageJ.

For cell viability measurements, 440µM resazurin (ThermoFisher Scientific, Waltham, MA)

in phosphate buffered saline was diluted 10-fold into the treatment medium, and cells were

incubated for 1.5 h. The medium was collected and briefly centrifuged to remove cells. 150 µL of

the supernatant was transferred to a 96-well plate, and resorufin (reduced resazurin) fluorescence

(Ex. 560 nm; Em. 590 nm) was measured on a Cytation5 Multimode Reader (Biotek, Winnoski,

VT).

Pyoverdine Purification

A LB overnight culture of *P. aeruginosa* PAO1 was diluted 100-fold into 300 mL of M9

medium (1% w/v 5X M9 Salts (BD Difco, Franklin Lakes, NJ), 1.5% w/v Bacto Casamino Acids

with low iron and salt content (BD Difco, Franklin Lakes, NJ), 1 mM $MgSO_4$, 1 mM $CaCl_2$) in a

2 L flask and grown aerobically for 24 h at 37 °C. Bacteria was then removed by centrifugation

and filtration through a 0.22 µm membrane. The filtrate was incubated with 10% w/v amberlite

XAD-4 resin at RT for 4 h. After rinsing the resin with copious amounts of water, pyoverdine was

eluted in 50% methanol. This eluent was diluted in water to 15% MeOH and loaded onto a Luna

Omega 5 µm Polar C18 LC prep column (Phenomenex, Torrance, CA) for high-performance

liquid chromatography on a 1220 Infinity LC system (Agilent Technologies, Santa Clara, CA).

Pyoverdine was eluted from the column by a 0-100% methanol gradient across 4 h at a flowrate

of 5 mL/min. Fractions were collected every other minute for pyoverdine content analysis **(Figure**

S3B). The fractions with the highest pyoverdine content were pooled. Methanol was removed from

this material using a SpeedVac vacuum concentrator. The final purified product was analyzed by

HPLC on an analytical column to verify sample purity **(Figure S3C)**.

Confocal Laser Scanning Microscopy

8×10^6 16HBE cells were seeded into each well of a collagen-coated 6-well plate and

grown at 37 °C for 24 h in a CO_2-jacketed incubator by which they reached 100% confluence.

After treatment, cells were washed in serum-free EMEM and detached from the microtiter plate

by trypsin-EDTA solution (MilliporeSigma, Burlington, MA). After inactivating the trypsin with

media containing 10% fetal bovine serum, the cells were concentrated via centrfiguation and

transferred onto a glass side with a 3% noble agar pad. These slides were visualized under a

LSM800 AiryScan confocal laser scanning microscope (Zeiss, Oberkochen, Germany).

Pyoverdine fluorescence was visualized via a 405 nm laser line using the channel conditions for

Pacific Blue. FITC-dextran fluorescence was visualized via a 488 nm laser line using channel

conditions for FITC. Dextran-Texas Red fluorescence was visualized via a 561 nm laser line using

channel conditions for Texas Red. CellMask Deep Red plasma membrane stain fluorescence was

visualized via a 640 nm laser line using channel conditions for Alex Fluor 660.

Giant plasma membrane vesicles from 16HBE cells were generated as previously described

(Gerstle et al., 2018). In brief, GPMV production was induced in a 100% confluent t-75 flask of

16HBE cells by a vesiculation buffer containing 1.9 mM DTT and 27.6 mM formaldehyde.

Pyoverdine or dextran treatment was directly applied to this GPMV/buffer. After treatment,

GPMVs were labeled with CellMask Deep Red plasma membrane stain then washed in

vesiculation buffer with centrifugation (16,000 g for 15 min) in between. GPMVs were transferred

onto a glass side with a 3% noble agar pad for confocal microscopy.

qRT-PCR

8×10^6 16HBE cells were seeded into each well of a collagen-coated 6-well plate and

grown at 37 °C for 24 h in a CO_2-jacketed incubator by which they reached 100% confluence.

After treatment, the medium was aspirated and cells were directly treated with TRI reagent

(Molecular Research Center, Cincinnati, OH) for phenol/chloroform/guanidinium thiocyanate

RNA extraction according to the manufacturer's protocols. Total RNA yield was quantified on a

Nanodrop spectrophotometer (ThermoFisher Scientific, Waltham, MA). cDNA synthesis was

performed on a Bio-RAD T100 Thermo Cycler (Bio-RAD, Hercules, CA) using a reverse

transcription kit (Applied Biosystems, Waltham, MA). qRT-PCR was performed on a Bio-RAD

CFX Connect Real-Time System (Bio-RAD, Hercules, CA) using a universal qPCR master mix

(New England Biolabs, Ipswich, MA). qPCR primer sequences are available upon request.

Acknowledgements:

This study was supported by funding from the Cystic Fibrosis Foundation (KIREIN20I0 to NVK;

KANG19H0, KANG22H0 to DK), National Institutes of Health (R35GM129294 to NVK), and

American Heart Association (903591 to DK). The authors declare no conflict of interest.

Figure Legends:

Figure 1. Pyoverdine-rich conditioned medium causes lung epithelial cells to detach from the

monolayer. (A, B) Bacterial growth **(A)** and pyoverdine production **(B)** by *P. aeruginosa* PAO1

siderophore biosynthetic mutants (PAO1*ΔpvdF* – pyoverdine; PAO1*ΔpchBA* – pyochelin;

PAO1*ΔpvdFΔpchBA* – pyoverdine and pyochelin) after 16 h growth in serum-free EMEM. **(C)**

MH-S murine alveolar macrophage viability after 2 h exposure to conditioned media from PAO1

siderophore mutants grown in EMEM. **(D)** Fluorescent micrographs of 16HBE human bronchial

epithelial cells after ~30 min exposure to conditioned media from PAO1 siderophore mutants

grown in EMEM. Cells were prelabeled with CellMask Orange plasma membrane stain. **(E)**

Quantification of percentage micrograph area covered by fluorescent cells. **(F, G)** Fluorescent

micrographs of 16HBE cells after 30 min exposure to wild-type PAO1 conditioned medium

pretreated with EDTA-free cOmplete Protease Inhibitor Cocktail or solvent control **(F)** and

quantification of percent cell coverage **(G)**. Error bars in **A, B, C, E, G** represent SEM between

four biological replicates. * corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on

Student's *t* test.

Figure 2. Damage to the epithelial monolayer is attenuated in conditioned medium from type

II secretion-deficient strains. (A) Fluorescent micrographs of 16HBE cells after 30 min exposure

to conditioned media from MPAO1 transposon mutants grown in EMEM. Cells were prelabeled

with CellMask Orange plasma membrane stain. **(B)** Quantification of percentage micrograph area

covered by fluorescent cells. **(C, D)** Bacterial growth **(C)** and pyoverdine production **(D)** by *P.*

aeruginosa cystic fibrosis isolates PA2-9 and PA3-29, their pyoverdine biosynthetic mutant

counterparts (Tn), and wild-type strain supplemented with 100 µM after 16 h growth in EMEM.

(E, F) Fluorescent micrographs of 16HBE cells after 30 min exposure to conditioned media from

P. aeruginosa cystic fibrosis isolates PA2-9 or PA3-29 **(E)** and quantification of percent cell

coverage **(F)**. Error bars in **B, C, D, F** represent SEM between four biological replicates. *

corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on Student's *t* test.

Figure 3. Pyoverdine translocates into lung epithelial cells and localizes to early endosomes.

(A) 16HBE cell viability after 72 h treatment of ciclopirox olamine, 1, 10-phenanthroline,

pyoverdine, or deferoxamine at 100 µM in serum-free EMEM. **(B)** Confocal micrographs of

16HBE cells exposed to 100 µM purified pyoverdine, pyoverdine with excess $Ga(NO_3)_3$,

pyoverdine with excess $FeCl_3$, or media control for 24 h. Cells were trypsinated prior to imaging.

(C) Quantification of pyoverdine fluorescence within 30 individual cells. **(D)** Confocal

micrographs of 16HBE cells treated with 100 µM pyoverdine-gallium and dextran-Texas Red

(10,000 MW). Error bars in **C** represent standard deviation. * corresponds to $p < 0.01$ based on

Student's *t* test.

Figure 4. Iron chelators promote the expression of proinflammatory genes in lung epithelial

cells. (A) Total RNA yield in 16HBE cells treated with ciclopirox olamine (CPX), 1,10-

phenanthroline (PHE), or pyochelin (PCH) for 30 h and cells treated with pyoverdine (PVD) or

deferoxamine (DFO) for 60 h normalized to that of media control. All treatments were at 100 µM

in serum-free EMEM. **(B)** Proinflammatory gene expression (*NLRP3, IL1B, NLRP1, IL8, TNF*)

normalized to *ACTB* expression in 16HBE cells treated with iron chelators. mRNA levels were

measured via qRT-PCR. **(C, D)** Total RNA yield **(C)** and proinflammatory gene expression **(D)** in

16HBE cells treated with iron chelators with or without excess $Ga(NO_3)_3$ supplementation. **(E, F)**

16HBE cell viability **(E)** and proinflammatory gene expression **(F)** after 60 h treatment with

pyoverdine, deferoxamine, or both molecules. Error bars in **A-F** represent SEM between three

biological replicates. * corresponds to $p < 0.01$, # corresponds to $p < 0.05$, and NS corresponds to

$p > 0.05$ based on Student's *t* test.

Supplementary Figure Legends:

Figure S1. Protease activity in pyoverdine-rich conditioned medium is important for the

disruption of epithelial integrity. (A) Fluorescent micrographs of 16HBE cells after ~30 min

exposure to conditioned media from *P. aeruginosa* PAO1 grown in EMEM. Conditioned media

were supplemented with 250 µM $Ga(NO_3)_3$, 5% fetal bovine serum (FBS), or EDTA-free

cOmplete Protease Inhibitor Cocktail, or had macromolecules (> 10 kDa) removed by centrifugal

filtration. Cells were prelabeled with CellMask Orange plasma membrane stain. **(B)** Quantification

of percentage micrograph area covered by fluorescent cells. Error bars represent SEM between

four biological replicates. * corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on

Student's *t* test.

Figure S2. PA2-9 and PA3-29 are highly virulent CF isolates. (A) Pyoverdine production in

M9 media by 69 multidrug-resistant *P. aeruginosa* strains isolated from pediatric cystic fibrosis

patients. **(B)** Percent *C. elegans* death following Liquid Killing by *P. aeruginosa* cystic fibrosis

isolates. Black bar represents the median pyoverdine production/*C. elegans death*. Red bar

represents PA2-9, blue bar represents PA3-29. Survey data was adapted from (Kang et al., 2019).

Figure S3. Purification of pyoverdine from *P. aeruginosa*. (A) Summary of the pyoverdine

purification pipeline. **(B)** Representative chromatogram from the HPLC purification step of the

pipeline. Red box depicts the predominant pyoverdine-containing fractions that were collected. **(C)**

Analysis of the final purified product via HPLC.

Figure S4. Pyoverdine accumulates in early endosomes of lung epithelial cells. (A, B) 16HBE

human bronchial epithelial cell viability after 48 **(A)** or 72 h **(B)** treatment of ciclopirox olamine,

1, 10-phenanthroline, pyoverdine, or deferoxamine in serum-free EMEM. **(C)** Confocal

micrographs of 16HBE cells exposed to 100 µM purified pyoverdine, pyoverdine with excess

$Ga(NO_3)_3$, pyoverdine with excess $FeCl_3$, or media control for 24 h. Cells were labeled with

CellMask Deep Red plasma membrane stain and trypsinated prior to imaging. **(D)** Confocal

micrographs of 16HBE cells treated with 100 µM pyoverdine-gallium and dextran-Texas Red

(10,000 MW). Bottom row shows an enlarged micrograph of one representative cell.

Figure S5. Pyoverdine translocates into giant plasma membrane vesicles. (A) Pyoverdine

fluorescence quenching by $FeCl_3$ supplementation in GPMVs treated with 100 µM pyoverdine

(GPMV - PVD) or pyoverdine diluted in buffer control (Free PVD). **(B, C)** Confocal micrographs

of GPMVs **(B)** and 16HBE cells **(C)** treated with 100 µM pyoverdine or solvent control for 24 h.

GPMVs were labeled with CellMask Deep Red plasma membrane stain prior to imaging. 16HBE

cells were trypsinated prior to imaging. **(D, E)** Confocal micrographs of GPMVs **(B)** and 16HBE

cells **(C)** treated with FITC-dextran (3000 MW) or solvent control for 24 h. GPMVs were labeled

with CellMask Deep Red plasma membrane stain prior to imaging. 16HBE cells were trypsinated

prior to imaging.

Figure S6. Deferoxamine promotes the expression of proinflammatory genes in 16HBE

CFTR mutants. (A) Total RNA yield in wild-type 16HBE cells and 16HBE cells carrying

mutations (G551D, ΔF508) in the cystic fibrosis transmembrane conductance regualator (CFTR)

after 48 h treatment with 100 µM deferoxamine in serum-free EMEM. **(B)** Proinflammatory gene

expression (*NLRP3, IL1B, NLRP1, IL8, TNF*) normalized to *ACTB* expression in cells treated with

deferoxamine. mRNA levels were measured via qRT-PCR. Error bars represent SEM between

three biological replicates.

References:

Andersen, S.B., Marvig, R.L., Molin, S., Krogh Johansen, H., and Griffin, A.S. (2015). Long-

term social dynamics drive loss of function in pathogenic bacteria. Proc Natl Acad Sci U S A

112, 10756-10761.

Anderson, G.G., and O'Toole, G.A. (2008). Innate and induced resistance mechanisms of

bacterial biofilms. Curr Top Microbiol Immunol *322*, 85-105.

Apseloff, G. (1999). Therapeutic uses of gallium nitrate: past, present, and future. Am J Ther *6*,

327-339.

Azghani, A.O. (1996). Pseudomonas aeruginosa and epithelial permeability: role of virulence

factors elastase and exotoxin A. Am J Respir Cell Mol Biol *15*, 132-140.

Banin, E., Vasil, M.L., and Greenberg, E.P. (2005). Iron and Pseudomonas aeruginosa biofilm

formation. Proc Natl Acad Sci U S A *102*, 11076-11081.

Bhagirath, A.Y., Li, Y., Somayajula, D., Dadashi, M., Badr, S., and Duan, K. (2016). Cystic

fibrosis lung environment and Pseudomonas aeruginosa infection. BMC Pulm Med *16*, 174.

Bradshaw, J.L., Caballero, A.R., Bierdeman, M.A., Adams, K.V., Pipkins, H.R., Tang, A.,

O'Callaghan, R.J., and McDaniel, L.S. (2018). Pseudomonas aeruginosa Protease IV Exacerbates

Pneumococcal Pneumonia and Systemic Disease. mSphere *3*.

Cantin, A.M., Hartl, D., Konstan, M.W., and Chmiel, J.F. (2015). Inflammation in cystic fibrosis

lung disease: Pathogenesis and therapy. J Cyst Fibros *14*, 419-430.

Choi, E.Y., Kim, E.C., Oh, H.M., Kim, S., Lee, H.J., Cho, E.Y., Yoon, K.H., Kim, E.A., Han,

W.C., Choi, S.C.*, et al.* (2004). Iron chelator triggers inflammatory signals in human intestinal

epithelial cells: involvement of p38 and extracellular signal-regulated kinase signaling pathways.

J Immunol *172*, 7069-7077.

Costabile, G., d'Angelo, I., d'Emmanuele di Villa Bianca, R., Mitidieri, E., Pompili, B., Del

Porto, P., Leoni, L., Visca, P., Miro, A., Quaglia, F.*, et al.* (2016). Development of inhalable

hyaluronan/mannitol composite dry powders for flucytosine repositioning in local therapy of

lung infections. J Control Release *238*, 80-91.

Curran, C.S., Bolig, T., and Torabi-Parizi, P. (2018). Mechanisms and Targeted Therapies for

Pseudomonas aeruginosa Lung Infection. Am J Respir Crit Care Med *197*, 708-727.

de Albuquerque Wanderley Sales, V., Timoteo, T.R.R., da Silva, N.M., de Melo, C.G., Ferreira,

A.S., de Oliveira, M.V.G., de Oliveira Silva, E., Dos Santos Mendes, L.M., Rolim, L.A., and

Neto, P.J.R. (2021). A Systematic Review of the Anti-inflammatory Effects of Gallium

Compounds. Curr Med Chem *28*, 2062-2076.

De Vos, D., De Chial, M., Cochez, C., Jansen, S., Tummler, B., Meyer, J.M., and Cornelis, P.

(2001). Study of pyoverdine type and production by Pseudomonas aeruginosa isolated from

cystic fibrosis patients: prevalence of type II pyoverdine isolates and accumulation of

pyoverdine-negative mutations. Arch Microbiol *175*, 384-388.

Dumas, Z., Ross-Gillespie, A., and Kummerli, R. (2013). Switching between apparently

redundant iron-uptake mechanisms benefits bacteria in changeable environments. Proceedings

Biological sciences *280*, 20131055.

Engel, L.S., Hill, J.M., Moreau, J.M., Green, L.C., Hobden, J.A., and O'Callaghan, R.J. (1998).

Pseudomonas aeruginosa protease IV produces corneal damage and contributes to bacterial

virulence. Invest Ophthalmol Vis Sci *39*, 662-665.

Engel, L.S., Hobden, J.A., Moreau, J.M., Callegan, M.C., Hill, J.M., and O'Callaghan, R.J.

(1997). Pseudomonas deficient in protease IV has significantly reduced corneal virulence. Invest

Ophthalmol Vis Sci *38*, 1535-1542.

Estivill, X., Bancells, C., and Ramos, C. (1997). Geographic distribution and regional origin of

272 cystic fibrosis mutations in European populations. The Biomed CF Mutation Analysis

Consortium. Hum Mutat *10*, 135-154.

Filloux, A. (2011). Protein Secretion Systems in Pseudomonas aeruginosa: An Essay on

Diversity, Evolution, and Function. Front Microbiol *2*, 155.

Gerstle, Z., Desai, R., and Veatch, S.L. (2018). Giant Plasma Membrane Vesicles: An

Experimental Tool for Probing the Effects of Drugs and Other Conditions on Membrane Domain

Stability. Methods Enzymol *603*, 129-150.

Goss, C.H., Kaneko, Y., Khuu, L., Anderson, G.D., Ravishankar, S., Aitken, M.L., Lechtzin, N.,

Zhou, G., Czyz, D.M., McLean, K.*, et al.* (2018). Gallium disrupts bacterial iron metabolism and

has therapeutic effects in mice and humans with lung infections. Sci Transl Med *10*.

Guillon, A., Brea, D., Morello, E., Tang, A., Jouan, Y., Ramphal, R., Korkmaz, B., Perez-Cruz,

M., Trottein, F., O'Callaghan, R.J.*, et al.* (2017). Pseudomonas aeruginosa proteolytically alters

the interleukin 22-dependent lung mucosal defense. Virulence *8*, 810-820.

Haas, B., Murphy, E., and Castignetti, D. (1991). Siderophore synthesis by mucoid Pseudomonas

aeruginosa strains isolated from cystic fibrosis patients. Can J Microbiol *37*, 654-657.

Hall, S., McDermott, C., Anoopkumar-Dukie, S., McFarland, A.J., Forbes, A., Perkins, A.V.,

Davey, A.K., Chess-Williams, R., Kiefel, M.J., Arora, D.*, et al.* (2016). Cellular Effects of

Pyocyanin, a Secreted Virulence Factor of Pseudomonas aeruginosa. Toxins (Basel) *8*.

Hassett, D.J., Borchers, M.T., and Panos, R.J. (2014). Chronic obstructive pulmonary disease

(COPD): evaluation from clinical, immunological and bacterial pathogenesis perspectives. J

Microbiol *52*, 211-226.

Hauser, A.R. (2009). The type III secretion system of Pseudomonas aeruginosa: infection by

injection. Nat Rev Microbiol *7*, 654-665.

Hirakata, Y., Furuya, N., Tateda, K., Kaku, M., and Yamaguchi, K. (1993). In vivo production of

exotoxin A and its role in endogenous Pseudomonas aeruginosa septicemia in mice. Infect

Immun *61*, 2468-2473.

Holden, V.I., Breen, P., Houle, S., Dozois, C.M., and Bachman, M.A. (2016). Klebsiella

pneumoniae Siderophores Induce Inflammation, Bacterial Dissemination, and HIF-1α

Stabilization during Pneumonia. MBio *7*.

Holden, V.I., Lenio, S., Kuick, R., Ramakrishnan, S.K., Shah, Y.M., and Bachman, M.A. (2014).

Bacterial siderophores that evade or overwhelm lipocalin 2 induce hypoxia inducible factor 1α

and proinflammatory cytokine secretion in cultured respiratory epithelial cells. Infect Immun *82*,

3826-3836.

Hosseinian, N., Cho, Y., Lockey, R.F., and Kolliputi, N. (2015). The role of the NLRP3

inflammasome in pulmonary diseases. Ther Adv Respir Dis *9*, 188-197.

Imperi, F., Massai, F., Facchini, M., Frangipani, E., Visaggio, D., Leoni, L., Bragonzi, A., and

Visca, P. (2013). Repurposing the antimycotic drug flucytosine for suppression of Pseudomonas

aeruginosa pathogenicity. Proc Natl Acad Sci U S A *110*, 7458-7463.

Jacobs, M.A., Alwood, A., Thaipisuttikul, I., Spencer, D., Haugen, E., Ernst, S., Will, O., Kaul,

R., Raymond, C., Levy, R.*, et al.* (2003). Comprehensive transposon mutant library of

Pseudomonas aeruginosa. Proc Natl Acad Sci U S A *100*, 14339-14344.

Kaneko, Y., Thoendel, M., Olakanmi, O., Britigan, B.E., and Singh, P.K. (2007). The transition

metal gallium disrupts Pseudomonas aeruginosa iron metabolism and has antimicrobial and

antibiofilm activity. The Journal of Clinical Investigation *117*, 877-888.

Kang, D., Kirienko, D.R., Webster, P., Fisher, A.L., and Kirienko, N.V. (2018). Pyoverdine, a

siderophore from Pseudomonas aeruginosa, translocates into C. elegans, removes iron, and

activates a distinct host response. Virulence *9*, 804-817.

Kang, D., and Kirienko, N.V. (2018). Interdependence between iron acquisition and biofilm

formation in Pseudomonas aeruginosa. J Microbiol *56*, 449-457.

Kang, D., and Kirienko, N.V. (2020). An In Vitro Cell Culture Model for Pyoverdine-Mediated

Virulence. Pathogens *10*.

Kang, D., Revtovich, A.V., Chen, Q., Shah, K.N., Cannon, C.L., and Kirienko, N.V. (2019).

Pyoverdine-Dependent Virulence of Pseudomonas aeruginosa Isolates From Cystic Fibrosis

Patients. Front Microbiol *10*, 2048.

Kang, D., Revtovich, A.V., Deyanov, A.E., and Kirienko, N.V. (2021a). Pyoverdine Inhibitors

and Gallium Nitrate Synergistically Affect Pseudomonas aeruginosa. mSphere, e0040121.

Kang, D., Zhang, L., and Kirienko, N.V. (2021b). High-Throughput Approaches for the

Identification of Pseudomonas aeruginosa Antivirulents. mBio *12*.

Kirienko, D.R., Revtovich, A.V., and Kirienko, N.V. (2016). A High-Content, Phenotypic

Screen Identifies Fluorouridine as an Inhibitor of Pyoverdine Biosynthesis and Pseudomonas

aeruginosa Virulence. mSphere *1*.

Kirienko, N.V., Ausubel, F.M., and Ruvkun, G. (2015). Mitophagy confers resistance to

siderophore-mediated killing by Pseudomonas aeruginosa. Proc Natl Acad Sci U S A *112*, 1821-

1826.

Kirienko, N.V., Kirienko, D.R., Larkins-Ford, J., Wählby, C., Ruvkun, G., and Ausubel, F.M.

(2013). Pseudomonas aeruginosa disrupts Caenorhabditis elegans iron homeostasis, causing a

hypoxic response and death. Cell Host Microbe *13*, 406-416.

Kollef, M.H., Chastre, J., Fagon, J.Y., Francois, B., Niederman, M.S., Rello, J., Torres, A.,

Vincent, J.L., Wunderink, R.G., Go, K.W.*, et al.* (2014). Global prospective epidemiologic and

surveillance study of ventilator-associated pneumonia due to Pseudomonas aeruginosa. Crit Care

Med *42*, 2178-2187.

Kruger, P., Saffarzadeh, M., Weber, A.N., Rieber, N., Radsak, M., von Bernuth, H., Benarafa,

C., Roos, D., Skokowa, J., and Hartl, D. (2015). Neutrophils: Between host defence, immune

modulation, and tissue injury. PLoS Pathog *11*, e1004651.

Lamont, I.L., Beare, P.A., Ochsner, U., Vasil, A.I., and Vasil, M.L. (2002). Siderophore-

mediated signaling regulates virulence factor production in Pseudomonasaeruginosa. Proc Natl

Acad Sci U S A *99*, 7072-7077.

Lee, H.J., Lee, J., Lee, S.K., Lee, S.K., and Kim, E.C. (2007). Differential regulation of iron

chelator-induced IL-8 synthesis via MAP kinase and NF-kappaB in immortalized and malignant

oral keratinocytes. BMC Cancer *7*, 176.

Lee, J., and Zhang, L. (2015). The hierarchy quorum sensing network in Pseudomonas

aeruginosa. Protein Cell *6*, 26-41.

Lyczak, J.B., Cannon, C.L., and Pier, G.B. (2002). Lung infections associated with cystic

fibrosis. Clin Microbiol Rev *15*, 194-222.

Malloy, J.L., Veldhuizen, R.A., Thibodeaux, B.A., O'Callaghan, R.J., and Wright, J.R. (2005).

Pseudomonas aeruginosa protease IV degrades surfactant proteins and inhibits surfactant host

defense and biophysical functions. Am J Physiol Lung Cell Mol Physiol *288*, L409-418.

Martin, L.W., Reid, D.W., Sharples, K.J., and Lamont, I.L. (2011). Pseudomonas siderophores in

the sputum of patients with cystic fibrosis. Biometals *24*, 1059-1067.

Marvig, R.L., Damkiær, S., Khademi, S.M., Markussen, T.M., Molin, S., and Jelsbak, L. (2014).

Within-host evolution of Pseudomonas aeruginosa reveals adaptation toward iron acquisition

from hemoglobin. MBio *5*, e00966-00914.

Mayer-Hamblett, N., Rosenfeld, M., Gibson, R.L., Ramsey, B.W., Kulasekara, H.D., Retsch-

Bogart, G.Z., Morgan, W., Wolter, D.J., Pope, C.E., Houston, L.S.*, et al.* (2014). Pseudomonas

aeruginosa in vitro phenotypes distinguish cystic fibrosis infection stages and outcomes. Am J

Respir Crit Care Med *190*, 289-297.

McElvaney, O.J., Zaslona, Z., Becker-Flegler, K., Palsson-McDermott, E.M., Boland, F.,

Gunaratnam, C., Gulbins, E., O'Neill, L.A., Reeves, E.P., and McElvaney, N.G. (2019). Specific

Inhibition of the NLRP3 Inflammasome as an Antiinflammatory Strategy in Cystic Fibrosis. Am

J Respir Crit Care Med *200*, 1381-1391.

Meyer, J.M., Neely, A., Stintzi, A., Georges, C., and Holder, I.A. (1996). Pyoverdin is essential

for virulence of Pseudomonas aeruginosa. Infect Immun *64*, 518-523.

Michalska, M., and Wolf, P. (2015). Pseudomonas Exotoxin A: optimized by evolution for

effective killing. Front Microbiol *6*, 963.

Minandri, F., Imperi, F., Frangipani, E., Bonchi, C., Visaggio, D., Facchini, M., Pasquali, P.,

Bragonzi, A., and Visca, P. (2016). Role of Iron Uptake Systems in Pseudomonas aeruginosa

Virulence and Airway Infection. Infect Immun *84*, 2324-2335.

Miyazaki, S., Matsumoto, T., Tateda, K., Ohno, A., and Yamaguchi, K. (1995). Role of exotoxin

A in inducing severe Pseudomonas aeruginosa infections in mice. J Med Microbiol *43*, 169-175.

Moldoveanu, B., Otmishi, P., Jani, P., Walker, J., Sarmiento, X., Guardiola, J., Saad, M., and Yu,

J. (2009). Inflammatory mechanisms in the lung. J Inflamm Res *2*, 1-11.

Moreau-Marquis, S., Stanton, B.A., and O'Toole, G.A. (2008). Pseudomonas aeruginosa biofilm

formation in the cystic fibrosis airway. Pulm Pharmacol Ther *21*, 595-599.

Murphy, T.F. (2009). Pseudomonas aeruginosa in adults with chronic obstructive pulmonary

disease. Curr Opin Pulm Med *15*, 138-142.

Ochsner, U.A., Johnson, Z., Lamont, I.L., Cunliffe, H.E., and Vasil, M.L. (1996). Exotoxin A

production in Pseudomonas aeruginosa requires the iron-regulated pvdS gene encoding an

alternative sigma factor. Mol Microbiol *21*, 1019-1028.

Oliver, J.M., Berlin, R.D., and Davis, B.H. (1984). Use of horseradish peroxidase and

fluorescent dextrans to study fluid pinocytosis in leukocytes. Methods Enzymol *108*, 336-347.

Pillar, C.M., and Hobden, J.A. (2002). Pseudomonas aeruginosa exotoxin A and keratitis in

mice. Invest Ophthalmol Vis Sci *43*, 1437-1444.

Plotkowski, M.C., Povoa, H.C., Zahm, J.M., Lizard, G., Pereira, G.M., Tournier, J.M., and

Puchelle, E. (2002). Early mitochondrial dysfunction, superoxide anion production, and DNA

degradation are associated with non-apoptotic death of human airway epithelial cells induced by

Pseudomonas aeruginosa exotoxin A. Am J Respir Cell Mol Biol *26*, 617-626.

Shah, K.N., Shah, P.N., Mullen, A.R., Chen, Q., Southerland, M.R., Chirra, B., DeBerardinis,

R.J., and Cannon, C.L. (2020). N-Acetyl cysteine abrogates silver-induced reactive oxygen

species in human cells without altering silver-based antimicrobial activity. Toxicol Lett *332*,

118-129.

Skinkle, A.D., Levental, K.R., and Levental, I. (2020). Cell-Derived Plasma Membrane Vesicles

Are Permeable to Hydrophilic Macromolecules. Biophys J *118*, 1292-1300.

Takase, H., Nitanai, H., Hoshino, K., and Otani, T. (2000). Impact of siderophore production on

Pseudomonas aeruginosa infections in immunosuppressed mice. Infect Immun *68*, 1834-1839.

Thi, M.T.T., Wibowo, D., and Rehm, B.H.A. (2020). Pseudomonas aeruginosa Biofilms. Int J

Mol Sci *21*.

Twigg, M.S., Brockbank, S., Lowry, P., FitzGerald, S.P., Taggart, C., and Weldon, S. (2015).

The Role of Serine Proteases and Antiproteases in the Cystic Fibrosis Lung. Mediators Inflamm

2015, 293053.

Ueda, A., Attila, C., Whiteley, M., and Wood, T.K. (2009). Uracil influences quorum sensing

and biofilm formation in Pseudomonas aeruginosa and fluorouracil is an antagonist. Microb

Biotechnol *2*, 62-74.

Visca, P., Imperi, F., and Lamont, I.L. (2007). Pyoverdine siderophores: from biogenesis to

biosignificance. Trends Microbiol *15*, 22-30.

Xiao, R., and Kisaalita, W.S. (1997). Iron acquisition from transferrin and lactoferrin by

Pseudomonas aeruginosa pyoverdin. Microbiology *143 (Pt 7)*, 2509-2515.

Zemljic Jokhadar, S., Klancnik, U., Grundner, M., Svelc Kebe, T., Vrhovec Hartman, S., Liovic,

M., and Derganc, J. (2018). GPMVs in variable physiological conditions: could they be used for

therapy delivery? BMC Biophys *11*, 1.

Zhang, C., Yang, B., Biazik, J.M., Webster, R.F., Xie, W., Tang, J., Allioux, F.M., Abbasi, R.,

Mousavi, M., Goldys, E.M.*, et al.* (2022). Gallium Nanodroplets are Anti-Inflammatory without

Interfering with Iron Homeostasis. ACS Nano *16*, 8891-8903.

Figure 1

Figure 2

Figure 3

Figure 4