Title: *In vitro* Lung Epithelial Cell Model Reveals Novel Roles for *Pseudomonas aeruginosa*

Siderophores

Authors: Donghoon Kang[1], Qi Xu[1,2], Natalia V. Kirienko[1]

Author Affiliations:

[1]Department of BioSciences, Rice University, Houston, TX, USA

[2]Department of Bioengineering, Rice University, Houston, TX, USA

Corresponding Author: Natalia V. Kirienko; kirienko@rice.edu

Abstract:

Multidrug-resistant *Pseudomonas aeruginosa* is a common nosocomial respiratory pathogen

that continues to threaten the lives of patients with mechanical ventilation in intensive care units

and those with underlying comorbidities such as cystic fibrosis or chronic obstructive pulmonary

disease. For over 20 years, studies have repeatedly demonstrated that the major siderophore

pyoverdine is an important virulence factor for *P. aeruginosa* in invertebrate and mammalian hosts

in vivo. Despite its physiological significance, an *in vitro,* mammalian cell culture model to

characterize the impact and molecular mechanism of pyoverdine-mediated virulence has only been

developed very recently. In this study, we adapt a previously-established, murine macrophage-

based model for human bronchial epithelial cells (16HBE). We demonstrate that conditioned

medium from *P. aeruginosa* induced rapid 16HBE cell death through the pyoverdine-dependent

secretion of cytotoxic rhamnolipids. Genetic or chemical disruption of pyoverdine biosynthesis

decreased rhamnolipid production and mitigated cell death. Consistent with these observations,

chemical depletion of lipid factors or genetic disruption of rhamnolipid biosynthesis was sufficient

to abrogate conditioned medium toxicity. Furthermore, we also examine the effects of purified

pyoverdine exposure on 16HBE cells. While pyoverdine accumulated within cells, the siderophore

was largely sequestered within early endosomes, showing minimal cytotoxicity. More membrane-

permeable iron chelators, such as the siderophore pyochelin, decreased epithelial cell viability and

upregulated several proinflammatory genes. However, pyoverdine potentiated these iron chelators

in activating proinflammatory pathways. Altogether, these findings suggest that the siderophores

pyoverdine and pyochelin play distinct roles in virulence during acute *P. aeruginosa* lung infection.

Key Words: *Pseudomonas aeruginosa*, Virulence, Siderophores, Lung Epithelial Cells,

Inflammation, Pyoverdine, Pyochelin, Rhamnolipids

Introduction:

Multidrug-resistant *Pseudomonas aeruginosa* is one of the most common Gram-negative,

respiratory pathogens, and infects mechanically-ventilated patients in intensive care units or those

with cystic fibrosis (CF) or chronic obstructive pulmonary disease (COPD) (1-5). This pathogen's

intrinsic resistance to several classes of antibiotics and exceptional ability to form biofilms on

medical devices and airway tissue pose a serious challenge for medical intervention (6, 7). In

addition to colonizing the respiratory tract, *P. aeruginosa* actively deploys numerous virulence

factors and toxins that damage host tissue, affecting pulmonary function (8). Two of the major

virulence factors produced by this pathogen are the siderophores pyoverdine and pyochelin.

Several have proposed possible mechanisms of siderophore-dependent virulence during *P.*

aeruginosa lung infection (9-12). As siderophores, both pyoverdine and pyochelin scavenge ferric

iron and provide the pathogen with this essential micronutrient during infection. However,

pyoverdine exhibits orders of magnitude higher affinity for ferric iron and is distinctly able to

chelate the metal from host ferroproteins such as transferrin and lactoferrin (13, 14). Generally,

iron acquisition serves an important function during infection by promoting bacterial growth and

biofilm formation (15, 16), and *P. aeruginosa* mutants lacking various iron uptake systems exhibit

attenuation of virulence during murine lung infection (17). It is important to note however that

these iron uptake systems do not contribute equally, and of the two siderophores, pyoverdine plays

a greater role in *P. aeruginosa* virulence (17).

Pyoverdine-mediated iron uptake further promotes *P. aeruginosa* virulence by derepressing

the alternative sigma factor PvdS, which activates the transcription of several virulence genes such

as those encoding the translational inhibitor exotoxin A, the exoprotease PrpL (protease IV), and

pyoverdine biosynthetic enzymes (12, 18). Furthermore, we have recently used a *Caenorhabditis*

elegans nematode model to demonstrate that pyoverdine likely directly chelates host iron,

disrupting mitochondrial homeostasis (19-21). Pyoverdine's well documented role in acute lung

infection is likely mediated by a combination of these various pathogenic functions (17, 22-25).

Recently, we established the first-reported *in vitro* cell culture model for pyoverdine-dependent

virulence, where murine macrophages were treated with conditioned medium from *P. aeruginosa*

grown in serum-free cell culture medium (26). Under these conditions, *P. aeruginosa* exhibited

robust pyoverdine production, yet the siderophore was not required for bacterial growth (**Fig. 1A,**

B; Fig. S1A, B), allowing for the study of pyoverdine's role in virulence. This pyoverdine-rich

conditioned medium from wild-type *P. aeruginosa* PAO1 was cytotoxic towards murine

macrophages, including murine alveolar macrophages (**Fig. S1C**); in clinical isolates; pyoverdine

content in the conditioned medium positively correlated with cytotoxicity (26).

In this report, we adapted this *in vitro* pyoverdine virulence model for human bronchial

epithelial cells to examine the consequences of pyoverdine production (i.e. exposure to pyoverdine

and/or pyoverdine-regulated virulence factors) during *P. aeruginosa* lung infection. Conditioned

medium from *P. aeruginosa* caused acute cell death and severe damage to the epithelial monolayer

in a pyoverdine-, but not pyochelin-, dependent manner. Interestingly, this damage did not require

host iron chelation nor production of the two known pyoverdine-regulated toxins, exotoxin A or

PrpL. Instead, pyoverdine production promoted the secretion of cytotoxic rhamnolipids that have

previously been shown to permeabilize host membranes (27). Consistent with this observation,

chemical depletion of lipids or genetic disruption of rhamnolipid production was sufficient to

abrogate toxicity from conditioned medium on 16HBE cells. Importantly, the pyoverdine

biosynthetic inhibitor 5-fluorocytosine effectively inhibited rhamnolipid production and mitigated

P. aeruginosa virulence in two highly virulent clinical isolates. We also examined the effects of

exposing 16HBE cells to purified pyoverdine. While pyoverdine accumulated within cells, the

siderophore was largely sequestered within early endosomes, showing minimal cytotoxicity. More

membrane-permeable iron chelators such as the siderophore pyochelin, decreased epithelial cell

viability and upregulated several proinflammatory pathways. However, pyoverdine potentiated

these iron chelators in activating proinflammatory pathways. Altogether, these findings suggest

that the siderophores pyoverdine and pyochelin play distinct roles in virulence during acute *P.*

aeruginosa lung infections.

Results:

Pyoverdine-rich, conditioned medium induces rapid cell death and damages the epithelial

monolayer

To investigate the role of pyoverdine production during *P. aeruginosa* lung infection, we

treated human bronchial epithelial cells (16HBE) with bacteria-free, pyoverdine-rich conditioned

medium from *P. aeruginosa* PAO1 grown in serum-free cell growth medium (Eagle's Minimum

Essential Medium or EMEM). To visualize the integrity of the epithelial monolayer, we prelabeled

cells with a plasma membrane stain. Within 30 min, the conditioned medium severely damaged

the monolayer, causing detachment of more than half of the cells **(Fig. 1C, D)**. This disruption was

significantly attenuated in 16HBE cells treated with identically prepared material from an isogenic

pyoverdine biosynthetic mutant (PAO1*ΔpvdF*) but not a pyochelin mutant (PAO1*ΔpchBA*) **(Fig.**

S1D, E). Preventing the biosynthesis of both pyoverdine and pyochelin (PAO1*ΔpvdFΔpchBA*)

conferred no further protection to 16HBE cells than the disruption of pyoverdine alone **(Fig. S1D,**

E). To determine whether cell detachment was caused by cell death (rather than from degradation

of the extracellular matrix via bacterial proteases and other factors), we labeled cells with cell-

permeant (Hoechst 33342; labels all cells) and cell-impermeant (Sytox Orange; labels only dead

cells) nucleic acid stains. Exposure to pyoverdine-rich, conditioned medium from wild-type cells

caused rapid (within 15 min) membrane permeabilization and internalization of the cell-

impermeant nucleic acid stain, suggesting cell death **(Fig. 1E, F)**. In contrast, conditioned medium

from the pyoverdine mutant exhibited substantially less cytotoxicity **(Fig. 1E, F)**.

One key difference between pyoverdine and pyochelin is their affinity for ferric iron. Due to

an exceptionally high affinity for the metal, pyoverdine is uniquely able to remove iron from host

ferroproteins (13, 14) and induce a lethal hypoxic response in a *C. elegans* nematode model (19).

We thus examined whether host iron chelation was important for the cell death we observed in the

epithelial monolayer. To hinder pyoverdine's ability to bind iron, we pretreated the conditioned

media with gallium (Ga^{3+}) or ferric iron (Fe^{3+}), either of which would prevent pyoverdine from

scavenging iron from the epithelial cells. Surprisingly, even with the addition of excess metal,

there was no significant rescue **(Fig. 1G, H; Fig. S2)**. Interestingly, we observed that removing

material with a molecular mass greater than 10 kDa (via centrifugal filtration) prevented damage

to the epithelial monolayer **(Fig. 1G, H, Fig. S2)**, suggesting that macromolecules or molecular

complexes were responsible for conditioned medium cytotoxicity. To investigate the role of

proteinaceous toxins, we pre-treated the conditioned medium with proteinase K. Degradation of

secreted proteins did not significantly attenuate conditioned medium cytotoxicity **(Fig. 1G, H)**.

We could not determine whether proteinase K treatment mitigated cell detachment; treatment

alone caused considerable damage to the extracellular matrix, making it very difficult to

unambiguously assess contributions **(Fig. S2)**. Consistent with this observation, genetic disruption

of the two pyoverdine-regulated toxins, exotoxin A and PrpL, or the type II secretion system

through which these toxins are secreted, did not affect conditioned medium toxicity **(Fig. S3)**. We

did however observe that removing lipids from the conditioned medium by chloroform extraction

abrogated cytotoxicity and prevented damage to the epithelial monolayer **(Fig. 1G, H; Fig. S2)**.

Based on these results, we directly measured lipid content in the conditioned medium using the

lipophilic dye FM 1-43. This probe is nonfluorescent in aqueous solution but is highly-fluorescent

upon binding to lipid membranes (28). Chloroform extraction and removal of macromolecules

(>10 kDa) significantly depleted lipid content **(Fig. 1I)**. Importantly, conditioned medium from

the pyoverdine biosynthetic mutant also contained significantly less lipid material than that from

wild-type PAO1 **(Fig. 1J)**.

Pyoverdine regulates the production of cytotoxic rhamnolipids

Based on previous studies (27, 29), we posited that the relevant secreted lipid factors were

rhamnolipids. We recently demonstrated that *P. aeruginosa* secretes rhamnolipids that rapidly

induce membrane rupture and permeabilization in a wide range of host cells, including murine

macrophages, human bronchial epithelial cells, and erythrocytes (27). Chloroform extraction and

centrifugal filtration, two treatments that would likely remove these rhamnolipids from

conditioned medium, decreased cytotoxicity **(Fig. 1G, H)**. To test this hypothesis, we used the

rhamnolipid biosynthetic mutant MPAO1*rhlA* to measure the lipid content of conditioned medium.

This mutant lacked lipid content in its conditioned medium but showed no apparent defects in

bacterial growth or pyoverdine production compared to the control strain MPAO1*cat* (transposon

inserted in an extraneous gene encoding a chloramphenicol acetyltransferase, **Fig. 2A-C)**,

suggesting that much of the lipid material being detected by FM 1-43 was due to rhamnolipid

production. Importantly, the conditioned medium from MPAO1*rhlA* neither damaged the 16HBE

epithelial monolayer nor induced cell death **(Fig. 2D-G)**. We observed similar results (i.e., reduced

lipid content and lower toxicity) for mutants with transposons inserted into genes encoding the

RhlRI quorum sensing system (MPAO1*rhlR*, MPAO1*rhlI*) **(Fig. 2A-G)**, indicating that

rhamnolipid production is primarily regulated via this signaling pathway in EMEM (30).

To test whether quorum-sensing is impaired in pyoverdine biosynthetic mutants, we took

advantage of an *Escherichia coli*-based bioluminescent reporter that responds to extracellular

quorum-sensing molecules (31), specifically N-butanoyl-L-homoserine lactone (C4-HSL) that is

produced by the RhlRI system (32). Using this reporter, we quantified C4-HSL concentrations in

the conditioned medium of pyoverdine mutants but did not observe significant difference in C4-

HSL production between the mutants and their pyoverdine-producing counterparts **(Fig. 2H; Fig.**

S4). For the *rhlI* C4-HSL biosynthetic mutant, the bioluminescent output in the conditioned

medium was comparable to that of media control **(Fig. S4)**, indicating that the reporter was

selectively responding to extracellular C4-HSL. These results imply that pyoverdine regulates

rhamnolipid production through an alternative pathway.

Furthermore, we examined the effects of exposing 16HBE cells to purified rhamnolipids (27).

At concentrations comparable to those seen in conditioned medium from wild-type PAO1 or

MPAO1*cat* **(Fig. 2I; Fig. S5A)**, purified rhamnolipids were sufficient to kill 16HBE cells **(Fig.**

2J, K; Fig. S5C, E). However, cells treated with purified rhamnolipids displayed less detachment

(Fig. S5B, D), suggesting that other secreted factors (e.g., proteases) in the conditioned medium

were required for full damage to the epithelial monolayer. Consistent with these findings, heat

denaturation of the conditioned medium did not affect rhamnolipid content **(Fig. S5F)** or

cytotoxicity of the material **(Fig. S5H, J),** but it did attenuate cell detachment from the monolayer

(Fig. S5G, I). Since the type II secretion system is responsible for the secretion of several *P.*

aeruginosa protein toxins and the majority of secreted proteases **(Fig. S3E)** (33), we hypothesized

that this system would contribute to cell detachment. We treated 16HBE cells with the conditioned

medium from a mutant (MPAO1*xcpQ*) lacking the outer membrane transporter for these secreted

proteins but saw no significant attenuation in cell detachment **(Fig. S3A, C)**. It is important to note

that while protease production in the type II secretion mutant was substantially impaired, it was

not abolished **(Fig. 3E)**. If extracellular proteases were involved in epithelial damage, they could

be secreted through other means.

Pyoverdine mutants do not exhibit changes in mexEF expression

While investigating possible secreted toxins that may facilitate rhamnolipid-mediated

epithelial damage, we serendipitously identified two presumably isogenic MPAO1 mutants with

transposon insertions in *plcH*, a gene that encodes a hemolytic phospholipase C: PW2536 and

PW2537. Sanger sequencing verified the presence of the transposon in the *plcH* gene in each of

the strains. (PW2536 – 1711/2193nt; PW2537 – 1049/2193nt;). Despite this, incongruous results

were obtained when conditioned medium was tested for cytotoxicity against 16HBE cells;

conditioned medium from PW2536 showed reduced toxicity, while identically-prepared medium

from PW2537 did not **(Fig. 3A-D).** To address this discrepancy, each strain was subjected to

whole-genome sequencing, which revealed two monogenic single nucleotide polymorphisms

(SNPs) between the two strains. This caused two amino acid substitutions in the transcriptional

regulator MexT in PW2536. MexT is known to regulate the *mexEF-oprN* multidrug efflux pump

operon **(Fig. 3E)** (34). The mutations in PW2536 led to the overexpression of *mexEF* **(Fig. 3F)**, a

phenotype previously associated with decreased rhamnolipid production **(Fig. 3G)** (35, 36).

Based on these observations, we tested whether pyoverdine biosynthetic mutants with poor

rhamnolipid production **(Fig. 3G)** had background mutations in MexT or increased *mexEF*

transcription. Neither the MPAO1 transposon mutant (MPAO1*pvdF*) nor the PAO1 in-frame

deletion mutant (PAO1*ΔpvdF*) had altered MexT protein sequence or changes in *mexEF*

expression **(Fig. 3E, F)**, suggesting that pyoverdine regulates rhamnolipid production through an

unknown *mexEF*- and C4-HSL- **(Fig. S4)** independent pathway.

5-Fluorocytosine inhibits pyoverdine and rhamnolipid production in highly virulent P. aeruginosa

clinical isolates

Since genetic disruption of pyoverdine biosynthesis decreased rhamnolipid production, we

hypothesized the same could be accomplished using a chemical inhibitor. To that end, we tested

whether the pyoverdine biosynthetic inhibitor and FDA-approved antimycotic drug 5-

fluorocytosine (5-FC) (23, 37, 38) could inhibit rhamnolipid production in several *P. aeruginosa*

strains, including PAO1 and two clinical strains isolated from pediatric CF patients, PA2-72 and

PA2-61. These two strains were selected from a large collection of CF isolates for their high *in*

vitro pyoverdine production and virulence against the nematode host *C. elegans* **(Fig. 4A, B)** (22).

These isolates also exhibited substantial *in vivo* pyoverdine production during acute murine lung

infection where they caused host mortality (22). When these strains were grown in EMEM, 5-FC

significantly impaired pyoverdine and rhamnolipid production without overtly affecting bacterial

growth **(Fig. 4C-F)** with the exception of PA2-61, where the drug induced planktonic cell

aggregation **(Fig. 4D)**, confounding bacterial growth measurement by optical density.

5-FC significantly attenuated 16HBE cell detachment and death after exposure to conditioned

medium from each of the three strains **(Fig. 4G-J)**. These findings are consistent with previous

work where the inhibition of pyoverdine biosynthesis by 5-FC was sufficient to rescue invertebrate

and mammalian hosts from *P. aeruginosa* virulence (22, 23, 39).

Pyoverdine translocates into 16HBE cells but is sequestered within early endosomes

Next, we wanted to investigate the consequences of exposing 16HBE cells to pyoverdine in

the absence of other virulence factors. In brief, pyoverdine-rich bacterial filtrate was subjected to

two purification steps to separate small molecules by polarity: absorption and elution from a

nonpolar polymeric resin (Amberlite XAD-4) and high-performance liquid chromatography

(HPLC) via a C-18 reverse-phase column **(Fig. 5A-C)**. We tested whether this purified material

was toxic to 16HBE cells using a resazurin-based cell viability assay and compared its toxicity to

other known iron-chelating molecules, namely the ferric iron chelator ciclopirox olamine, the

ferrous iron chelator 1,10-phenanthroline, or the siderophores pyochelin (from *P. aeruginosa*) or

deferoxamine (from *Streptomyces spp.*). While all other iron chelators exhibited time- and dose-

dependent cytotoxicity towards 16HBE cells, pyoverdine remained largely nontoxic **(Fig. 5D)**

even after 72 h treatment at 200 µM **(Fig. S6A, B)**.

Because pyoverdine is considerably larger than these iron chelators, with a molecular weight

of ~1,365 g/mol, it may be unable to translocate across cellular membranes. We took advantage of

pyoverdine's intrinsic spectral properties to examine whether pyoverdine could translocate into

16HBE cells. After 24 h, 16HBE cells treated with purified pyoverdine showed considerable

internalization of pyoverdine **(Fig. 5E, F)**. Consistent with previous studies, this intracellular

fluorescence was enhanced when pyoverdine was presaturated with gallium and was quenched

when pyoverdine was presaturated with iron **(Fig. 5E, F)** (21, 26). Importantly, pyoverdine

fluorescence did not colocalize with that of the plasma membrane stain, indicating that pyoverdine

was within the cell **(Fig. S6C)**. However, we also observed that pyoverdine fluorescence formed

distinct punctae within the cell. Based on previous observations in murine macrophages (26), we

hypothesized that pyoverdine was sequestered within early endosomes. Supporting this hypothesis,

pyoverdine colocalized with fluorophore-conjugated 10 kDa dextran, a well-established

endosomal marker **(Fig. 5G, Fig. S6D)** (40). These results are consistent with our observations

that pyoverdine, unlike other iron chelating molecules, exhibited low cytotoxicity towards 16HBE

cells.

Iron chelation activates a proinflammatory response in 16HBE cells

While pyochelin exhibits lower affinity towards ferric iron than pyoverdine, it is also

substantially smaller, with a molecular weight of ~325 g/mol. We hypothesized that pyochelin

may be able to enter 16HBE cells and chelate intracellular iron. While we were not able to visualize

pyochelin within cells (due to its lack of distinct spectral properties), one likely consequence of

iron deprivation in epithelial cells would be a proinflammatory transcriptional response. Several

studies have demonstrated that iron chelation by various siderophores such as deferoxamine or

enterobactin promotes the production of proinflammatory cytokines, most notably interleukin

(IL)-8 in lung epithelial cells, intestinal epithelial cells, or oral keratinocytes (41-43). To reaffirm

these findings, we treated 16HBE cells with various iron chelators and measured the mRNA levels

of genes involved in neutrophilic inflammation. We first observed that total RNA yield (from

phenol-chloroform extraction) in these cells corresponded with the resazurin-based cell viability

assay **(Fig. 5D)**. Cells treated with small molecule (< 1,000 MW), cytotoxic iron chelators

(ciclopirox olamine, phenanthroline, pyochelin, or deferoxamine) yielded lower quantities of RNA,

while cells treated with pyoverdine had RNA quantities comparable to that of media control **(Fig.**

6A). Using qRT-PCR, we measured the expression of genes encoding components of the

inflammasome, namely *NLRP3* and *NLRP1*, and those encoding the major proinflammatory

cytokines produced by lung epithelial cells, specifically IL-1β (*IL1B*), IL-8 (*IL8*), and tumor

necrosis factor alpha (*TNF*). All of these genes have been associated with inflammation during

lung infection (44). With the exception of pyoverdine, all iron chelators induced the expression of

these proinflammatory genes **(Fig. 6B)**. For *IL8*, we validated the qRT-PCR results by ELISA to

determine whether transcriptional activation led to increased cytokine production. Cells treated

with cytotoxic iron chelators exhibited time-dependent increase in IL-8 secretion that correlated

to *IL8* mRNA levels **(Fig. 6C, Fig. S7)**. Cells treated with pyoverdine exhibited IL-8 secretion

comparable to that of media control **(Fig. 6C)**. To ensure that the observed proinflammatory

response was due to iron chelation, we presaturated pyochelin and deferoxamine with excess

gallium (1:2 stoichiometric ratio) prior to exposure. Cells treated with gallium-bound pyochelin or

deferoxamine yielded RNA quantities comparable to that of media control, suggesting that gallium

inhibited the cytotoxic effects of the siderophores **(Fig. 6D)**. Furthermore, pretreating the

siderophores with gallium resulted in a significant decrease in proinflammatory gene expression

(Fig. 6E), demonstrating that the siderophore-induced inflammatory response was due to iron

chelation rather than other nonspecific reactions or contaminants in the commercially-sourced

material.

Finally, we investigated whether pyoverdine can indirectly promote lung inflammation by

potentiating other iron chelating molecules. Due to its exceptionally high affinity for iron,

pyoverdine is likely to remove iron from other, more cell-permeable siderophores or outcompete

them for trace iron in the extracellular milieu, increasing the pool of apo-siderophores that can

promote inflammation. To test this hypothesis, we treated 16HBE cells with deferoxamine,

pyoverdine, or both. Cells treated with both siderophores exhibited higher expression of

proinflammatory genes compared to those treated with deferoxamine alone **(Fig. 6F)**. Considering

that pyoverdine alone did not affect the transcription of proinflammatory genes, these results

suggest that pyoverdine enhanced deferoxamine-mediated damage by removing iron from

deferoxamine and effectively increasing the pool of apo-deferoxamine.

Discussion:

Arguably one of the greatest challenges to combating *P. aeruginosa* infections is the sheer

multitude of virulence factors produced by the bacterium that contribute to pathogenesis. These

include small molecule virulence factors (e.g., siderophores, quorum-sensing molecules), factors

involved in biofilm formation and motility (e.g, exopolysaccharides, type IV pili, flagella), and

more than twenty toxins that either directly kill host cells (e.g., exotoxin A, exoenzyme S, exotoxin

T, exotoxin U) or damage host tissue (e.g., elastase LasA, elastase LasB, PrpL, alkaline protease)

(32, 45-48). This complexity casts a dark shadow over the prospects of epidemiological or

therapeutic intervention. Ideally, we would be able to reliably predict a pathogen's ability to cause

disease through our evolving molecular surveillance tools such as whole-genome sequencing and

mass spectrometry and therapeutically impair pathogenesis through antivirulence drugs that inhibit

the production or function of key virulence factors and toxins. In *P. aeruginosa*, the only feasible

way to approach these strategies would be to unravel the regulation of virulence factors in the

bacterium and to target virulence networks rather than individual factors.

The results we report in this study suggest that the alternative sigma factor PvdS may be a

promising target for therapeutic intervention during *P. aeruginosa* lung infections. PvdS has

already been shown to regulate the production of several secreted toxins, such as the translational

inhibitor exotoxin A and the secreted protease PrpL. Exotoxin A, arguably one of the most

extensively studied toxins in *P. aeruginosa*, inhibits protein synthesis (46, 49), inducing airway

epithelial cell death (50) and inhibiting cell junction repair in the presence of *P. aeruginosa* elastase

(51). Exotoxin A also contributes to *P. aeruginosa* virulence in various murine infection models

(52-54). PrpL degrades host defense factors, like surfactant proteins and IL-22, that contribute to

lung innate immunity (55-57). PrpL has also been shown to directly contribute to *P. aeruginosa*

virulence during ocular infections (58, 59).

PvdS is best known for its role in pyoverdine biosynthesis and is indispensable for the

production of pyoverdine biosynthetic enzymes. In addition to scavenging trace iron in the

environment or directly from host ferroproteins, pyoverdine is involved in a positive feedback loop

where the uptake of iron-bound pyoverdine by its outer membrane receptor, FpvR, derepresses

PvdS, increasing the production of pyoverdine, exotoxin A, and PrpL (12, 18). Here, we have

demonstrated that pyoverdine promotes the production of an additional secreted material,

rhamnolipids, that rapidly induce cell death. Secreted rhamnolipids have been shown to assemble

into micellar structures (60) that directly interact with host membranes, causing rapid plasma

membrane rupture and cell death (29, 61, 62) and damaging the lung epithelium (63, 64). It remains

unclear however how pyoverdine regulates rhamnolipid production and whether this mechanism

is linked to quorum-sensing, the primary mode of rhamnolipid regulation in the bacterium. While

certain studies suggest that *P. aeruginosa* quorum-sensing affects pyoverdine production (65-69),

an inverse relationship has yet to be explored.

In addition to regulating secreted toxins, pyoverdine may also indirectly contribute to

inflammation by removing iron from other more cell permeable siderophores such as pyochelin,

deferoxamine, or enterobactin, the latter in the context of polymicrobial infections with

Enterobacteriaceae such as the respiratory pathogen *K. pneumoniae* (70). Importantly, while *P.*

aeruginosa may lose the ability to produce pyoverdine during lung infection with the emergence

of social cheaters or due to a transition in iron acquisition strategy (71-73), several surveys of

patient sputum samples and clinical isolates have revealed that a large fraction of strains still

exhibit substantial pyoverdine production (22, 74-76), demonstrating that pyoverdine may be an

important target for therapeutic intervention.

We took advantage of the FDA-approved antimycotic drug, 5-fluorocytosine (5-FC), that

inhibits *pvdS* expression in *P. aeruginosa* and attenuates virulence during murine lung infection.

Imperi and colleagues first identified 5-FC in a screen for small molecules that inhibit pyoverdine

production (23). We have independently identified a chemical analogue of 5-FC, 5-fluorouracil –

another *pvdS* inhibitor, in a small molecule screen for compounds that rescue *C. elegans* from *P.*

aeruginosa in a pyoverdine-dependent pathogenesis model (37, 77). We also recently reported that

5-FC synergizes with another FDA-approved drug, gallium nitrate, to inhibit *P. aerugin*osa growth

and virulence against *C. elegans* (38). Our findings in this study suggest that in addition to its

bactericidal and biofilm-inhibitory activities (78, 79), gallium could also function as an anti-

inflammatory agent during lung infection by inhibiting intracellular iron chelation by pyochelin

and mitigating not only epithelial cell death, but also activation of proinflammatory pathways such

as the NLRP3 inflammasome and IL-8 production. These newly discovered roles for pyochelin

further demonstrate that pyoverdine and pyochelin play distinct roles in *P. aeruginosa* virulence.

Previous studies have shown that pyochelin triggers reactive oxygen species (ROS) production in

an iron-dependent manner in host cells during infection (80) or in other microbes during

interbacterial competition (81-83). While pyochelin may not regulate additional virulence

pathways, its ability to permeate membranes makes this siderophore a distinctly effective tool for

direct host cell damage.

The benefits of suppressing pyochelin-mediated neutrophilic inflammation during lung

infection, particularly chronic lung infection, has been well documented. While the mechanisms

neutrophils employ to kill and remove pathogens such as the production of neutrophil elastases are

important for host defense, they can also cause tissue damage by degrading extracellular matrix

proteins (84, 85). During chronic infections (such as those in CF patients), these host defense

factors continue to cause airway damage while the pathogen persists, exacerbating the decline in

pulmonary function (86). While lung inflammation is mediated by many factors (cytokines and

chemokines), for CF patients, a strategy to specifically inhibit the NLRP3 inflammasome by

therapeutics such as MCC950 is currently being investigated with promising results in murine

infection studies (87, 88). It is important to note that NLRP3 inflammasome priming (i.e.,

transcriptional upregulation of *NLRP3*, encoding the major component of the inflammasome, and

IL1B, encoding pro-IL-1β) (89) was a pathway that was activated by intracellular iron chelation in

lung epithelial cells **(Fig. 6B)**. We observed this transcriptional response in not only wild-type

16HBE cells but also those carrying mutations in the cystic fibrosis transmembrane conductance

regulator (CFTR G551D, CFTR ΔF508), two of the most frequently identified mutations in CF

patients **(Fig. S8)** (90). While gallium has been broadly associated with anti-inflammatory

properties (91-93), studies have yet to specifically explore gallium's role in inhibiting pathogen-

associated inflammation. Considering recent findings that bacterial siderophores promote

inflammation (41-43, 70), this therapeutic avenue may merit consideration.

Materials and Methods:

Bacterial Strains and Growth Conditions

See list of bacterial strains in **Table 1**. All MPAO1 transposon insertion sites were verified by

Sanger sequencing (94). Insertions were determined by adapting a previously established method

for the MAR2xT7 transposon library in PA14 (95) and primers in **Table S2**. To produce

pyoverdine-rich conditioned medium, an LB overnight culture of *P. aeruginosa* was diluted 20-

fold into 2 mL of serum-free Eagle's Minimum Essential Medium (EMEM) (Millipore Sigma, St.

Louis, MO) in a 6-well plate. The plate was sealed with a Breathe-Easy sealing membrane

(Diversified Biotech, Dedham, MA) and grown statically at 37 °C for 18 h. Pyoverdine production

(Ex. 405 nm; Em. 460 nm) and bacterial growth (Abs. 600 nm) were measured

spectrophotometrically on a Cytation5 Multimode Reader (Biotek, Winnoski, VT). Bacteria was

then removed by centrifugation and the supernatant was treated with an antibiotic cocktail to kill

residual bacteria (100 µg/mL amikacin, 100 µg/mL gentamicin, 100 µg/mL tobramycin).

Cell Culture

Wild-type and mutant 16HBE cells **(Table 1)** were passaged in Eagle's Minimum Essential

Medium (EMEM) supplemented with 10% fetal bovine serum (Corning, Corning, NY),

penicillin/streptomycin (Millipore Sigma), and MEM non-essential amino acids (Millipore Sigma).

For experiments with *P. aeruginosa* conditioned medium, 4×10^6 cells were seeded into each

well of a collagen (type I from calf skin - Millipore Sigma)-coated 12-well plate and grown at

37 °C for 24 h in a CO_2-jacketed incubator by which they reached 100% confluence. To visualize

the epithelial monolayer, cells were stained with 2.5 µg/mL CellMask Orange plasma membrane

stain (Invitrogen, Carlsbad, CA) for 1 h prior to conditioned medium exposure. Following

treatment, the medium was aspirated and the monolayer was imaged on a Cytation5 Multimode

Reader using a RFP filter cube. Percentage image area covered by fluorescent cells was quantified

using ImageJ. To visualize cell death, cells were prelabeled with 20 µM Hoechst 33342

(ThermoFisher Scientific, Waltham, MA) for 30 min then exposed to conditioned medium in the

presence of 2.5 µM Sytox Orange (Invitrogen). Following treatment, the medium was aspirated,

and the monolayer was imaged on a Cytation5 Multimode Reader using DAPI (for Hoechst 33342)

and RFP (for Sytox Orange) filter cubes. Images were exported and quantified for mean blue or

red fluorescence intensity on ZEN Blue image analysis software (Zeiss, Oberkochen, Germany).

For 16HBE cell viability measurements following iron chelator treatment, 440 µM resazurin

(ThermoFisher Scientific) in phosphate buffered saline was diluted 10-fold into the treatment

medium, and cells were incubated for 1.5 h. The medium was collected and briefly centrifuged to

remove cells. 150 µL of the supernatant was transferred to a 96-well plate, and resorufin (reduced

resazurin) fluorescence (Ex. 560 nm; Em. 590 nm) was measured on a Cytation5 Multimode

Reader.

N-butanoyl-L-homoserine lactone (C4-HSL) Measurement

To measure C4-HSL concentration, one part spent culture medium from *P. aeruginosa* grown

in EMEM was mixed with three parts *E. coli* JM109 pSB536 in LB medium. This LB medium

was inoculated with the *E. coli* reporter strain by diluting an overnight culture 5-fold. 150 µL of

the *P. aeruginosa* conditioned medium-reporter strain mixture was transferred to each well in a

96-well plate and incubated at 37 °C for 2 h in a Cytation5 Multimode Reader. Bioluminescence

measurements were taken every 15 min.

P. aeruginosa whole genome sequence analysis

Bacterial genomic DNA was purified from overnight culture using DNeasy UltraClean

Microbial Kit (Qiagen, Hilden, Germany). Paired end Illumina sequencing was performed by the

Microbial Genome Sequencing Center (MiGS, Pittsburgh, PA) for at least 40X genome coverage.

To compare the two *plcH* transposon mutants, raw sequencing reads from PW2537 were first

assembled via SPAdes (96) and annotated via Prokka using PAO1 (NC_002516.2) as the reference

genome (97). Genomic polymorphisms in PW2536 were identified using *breseq* (98).

Pyoverdine Purification

A LB overnight culture of *P. aeruginosa* PAO1 was diluted 100-fold into 300 mL of M9

medium (1% w/v 5X M9 Salts (BD Difco, Franklin Lakes, NJ), 1.5% w/v Bacto Casamino Acids

with low iron and salt content (BD Difco), 1 mM $MgSO_4$, 1 mM $CaCl_2$) in a 2 L flask and grown

aerobically for 24 h at 37 °C. Bacteria were then removed by centrifugation and filtration through

a 0.22 µm membrane. The filtrate was incubated with 10% w/v amberlite XAD-4 resin

(MilliporeSigma) at room temperature for 4 h with constant agitation. After rinsing the resin with

copious amounts of water, pyoverdine was eluted in 50% methanol. This eluent was diluted in

water to 15% methanol and loaded onto a Luna Omega 5 µm Polar C18 LC prep column

(Phenomenex, Torrance, CA) for high-performance liquid chromatography on a 1220 Infinity LC

system (Agilent Technologies, Santa Clara, CA). Pyoverdine was eluted from the column by a 0-

100% methanol gradient across 4 h at a flowrate of 5 mL/min. Fractions were collected every other

minute for pyoverdine content analysis **(Fig. 5B)**. The fractions with the highest pyoverdine

content were pooled. Methanol was evaporated using a SpeedVac vacuum concentrator. The final

purified product was analyzed by HPLC on an analytical column to verify sample purity **(Fig. 5C)**.

Confocal Laser Scanning Microscopy

8×10^6 16HBE cells were seeded into each well of a collagen-coated 6-well plate and

grown at 37 °C for 24 h in a CO_2-jacketed incubator by which they reached 100% confluence.

After treatment, cells were washed in serum-free EMEM and detached from the microtiter plate

by trypsin-EDTA solution (MilliporeSigma). After inactivating the trypsin with media containing

10% fetal bovine serum, the cells were concentrated via centrifugation and transferred onto a glass

side with a 3% noble agar pad. These slides were visualized under a LSM800 AiryScan confocal

laser scanning microscope (Zeiss). Pyoverdine fluorescence was visualized via a 405 nm laser line

using the channel conditions for Pacific Blue. Dextran-Texas Red (Invitrogen) fluorescence was

visualized via a 561 nm laser line using channel conditions for Texas Red. CellMask Deep Red

plasma membrane stain (Invitrogen) fluorescence was visualized via a 640 nm laser line using

channel conditions for Alex Fluor 660.

qRT-PCR

For *P. aeruginosa*, bacterial cells were collected from 12 mL of EMEM culture by

centrifugation. The pellet was resuspended in 2 mL of TRI reagent (Molecular Research Center,

Cincinnati, OH) for phenol/chloroform/guanidinium thiocyanate RNA extraction according to

manufacturer's protocols (bromochloropropane phase separation followed by isopropanol RNA

precipitation). Prior to phase separation, bacterial cells were homogenized with 0.1 mm zirconia

beads by vigorous vortexing. To remove DNA contaminants in bacterial RNA extracts, samples

were treated with DNase I (ThermoFisher Scientific) at 37 °C for 30 min, followed by 75 °C

enzyme heat denaturation for 10 min. For 16HBE cells grown and treated in 6-well plates, the

treatment medium was aspirated, and cells were incubated in TRI reagent at room temperature

for 15 min to lyse cells prior to RNA extraction.

For both bacterial and human cell RNA, cDNA synthesis was performed on a Bio-Rad T100

Thermo Cycler (Bio-Rad, Hercules, CA) using a reverse transcription kit (Applied Biosystems,

Waltham, MA). qRT-PCR was performed on a Bio-Rad CFX Connect Real-Time System (Bio-

Rad) using a universal qPCR master mix (New England Biolabs, Ipswich, MA). All qPCR primer

sequences are in **Table S2**. For *P. aeruginosa* genes, cDNA amplification (Ct value) was

normalized to that of housekeeping gene *gyrB*. For 16HBE genes, cDNA amplification was

normalized to that of *ACTB*.

Strains	Relevant Genotype	Source or Reference
Pseudomonas aeruginosa Strains		
PAO1	WT	D. Haas
PAO1*ΔpvdF*	*ΔpvdF*	D. Haas
PAO1*ΔpchBA*	*ΔpchBA*	D. Haas
PAO1*ΔpvdFpchBA*	*ΔpvdFpchBA*	D. Haas
MPAO1 PW2280	*cat* (ISlacZ/hah)	Jacobs et al., 2003
MPAO1 PW5033	*pvdF* (ISlacZ/hah)	Jacobs et al., 2003
MPAO1 PW3078	*toxA* (ISphoA/hah)	Jacobs et al., 2003
MPAO1 PW8077	*prpL* (ISphoA/hah)	Jacobs et al., 2003
MPAO1 PW6887	*rhlA* (ISphoA/hah)	Jacobs et al., 2003
MPAO1 PW6882	*rhlR* (ISlacZ/hah)	Jacobs et al., 2003
MPAO1 PW6880	*rhlI* (ISphoA/hah)	Jacobs et al., 2003
MPAO1 PW5085	*pvdS* (ISlacZ/hah)	Jacobs et al., 2003
MPAO1 PW6223	*xcpQ* (ISphoA/hah)	Jacobs et al., 2003
MPAO1 PW2537	*plcH* (ISlacZ/hah)	Jacobs et al., 2003
MPAO1 PW2536	*plcH* (ISlacZ/hah)	Jacobs et al., 2003
PA2-72	WT (CF Isolate)	Kang et al., 2019
PA2-61	WT (CF Isolate)	Kang et al., 2019
Other Bacterial Strains		
Escherichia coli JM109 pSB536	*PahyI*::*luxABCDE* regulated by constitutive AhyR	Winson et al., 1998
Human Bronchial Epithelial Cells		
16HBE	WT CFTR	G. Bao
16HBE G551D	CFTR G551D	G. Bao
16HBE ΔF508	CFTR ΔF508	G. Bao

Table 1. List of bacterial strains and cell lines used in this study.

Acknowledgements:

This study was supported by funding from the Cystic Fibrosis Foundation (KIRIEN20I0 to NVK;

XU23H0 to QX; KANG19H0, KANG22H0 to DK), National Institutes of Health (R35GM129294

to NVK), and American Heart Association (903591 to DK). The authors declare no conflict of

interest.

Figure Legends:

Fig. 1. Pyoverdine-rich conditioned medium kills 16HBE cells and damages the epithelial

monolayer. (A, B) Bacterial growth **(A)** or pyoverdine production **(B)** by WT *P. aeruginosa* PAO1

or pyoverdine biosynthetic mutant (PAO1*ΔpvdF*) in serum-free EMEM. **(C)** Fluorescent

micrographs of 16HBE human bronchial epithelial cells after 30 min exposure to conditioned

medium from WT PAO1 or PAO1*ΔpvdF* grown in EMEM. Cells were prelabeled with CellMask

Orange plasma membrane stain. **(D)** Quantification of percentage micrograph area covered by

fluorescent cells. **(E)** Fluorescent micrographs of 16HBE cells after 15 min exposure to

conditioned medium from WT PAO1 or PAO1*ΔpvdF* in the presence of Sytox Orange nucleic acid

stain (red). Cells were prelabeled with Hoechst 33342 nucleic acid stain (blue). **(F)** Quantification

of Sytox Orange mean fluorescence intensity normalized to that of Hoechst 33342. **(G)** Fluorescent

micrographs of 16HBE cells after 15 min exposure to conditioned medium from WT PAO1 in the

presence of Sytox Orange nucleic acid stain (red). Conditioned medium was pretreated with 200

µM Ga(NO$_3$)$_3$, 200 µM FeCl$_3$, or 100 µg/mL proteinase K for 24 h, had macromolecules depleted

via a 10 kDa centrifugal filter, or had lipids depleted by chloroform (CHCl$_3$) extraction. Cells were

prelabeled with Hoechst 33342 nucleic acid stain (blue). **(H)** Quantification of Sytox Orange mean

fluorescence intensity normalized to that of Hoechst 33342. **(I)** Quantification of lipids by FM 1-

43 fluorescent labeling in conditioned medium from WT PAO1 (control) or macromolecule- (<10

kDa) or lipid- (CHCl$_3$) depleted material. **(J)** Quantification of lipids in conditioned medium from

WT PAO1 or PAO1*ΔpvdF* by FM 1-43 labeling. All error bars represent SEM from at least three

biological replicates. * corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on Student's

t-test **(D, F, J)** or one-way ANOVA with Dunnett's multiple comparisons test **(H, I)**.

Fig. 2. Pyoverdine regulates the production of rhamnolipids. (A, B) Bacterial growth **(A)** or

pyoverdine production **(B)** by MPAO1 transposon mutants in serum-free EMEM. **(C)**

Quantification of lipids in conditioned medium from MPAO1 transposon mutants by FM 1-43

fluorescent labeling. **(D)** Fluorescent micrographs of 16HBE cells after 30 min exposure to

conditioned medium from MPAO1 transposon mutants. Cells were prelabeled with CellMask

Orange plasma membrane stain. **(E)** Fluorescent micrographs of 16HBE cells after 15 min

exposure to conditioned medium from MPAO1 transposon mutants in the presence of Sytox

Orange nucleic acid stain (red). Cells were prelabeled with Hoechst 33342 nucleic acid stain (blue).

(F) Quantification of percentage micrograph area covered by fluorescent cells in **(D)**. **(G)**

Quantification of Sytox Orange mean fluorescence intensity normalized to that of Hoechst 33342

in **(E)**. **(H)** Bioluminescence produced by *E. coli* JM109 pSB536 (N-butanoyl-L-homoserine

lactone reporter strain) grown in media supplemented with conditioned medium from MPAO1

transposon mutants. **(I)** Quantification of lipids in EMEM with purified rhamnolipids (RHL) by

FM 1-43 labeling. **(J)** Fluorescent micrographs of 16HBE cells after 15 min exposure to purified

rhamnolipids from *P. aeruginosa* in the presence of Sytox Orange nucleic acid stain (red). Cells

were prelabeled with Hoechst 33342 nucleic acid stain (blue). **(K)** Quantification of Sytox Orange

mean fluorescence intensity normalized to that of Hoechst 33342 in **(J)**. All error bars represent

SEM from three biological replicates. * corresponds to $p < 0.01$, # corresponds to $p < 0.05$, and

NS corresponds to $p > 0.05$ based on one-way ANOVA with Dunnett's multiple comparisons test

(C, F, G) or Tukey's multiple comparisons test **(H)** – see **Fig. S4** or Student's *t*-test **(K)**.

Fig. 3. *mexEF* overexpression mutant exhibits decreased rhamnolipid production. (A)
Fluorescent micrographs of 16HBE cells after 30 min exposure to conditioned medium from *plcH*
transposon mutants. Cells were prelabeled with CellMask Orange plasma membrane stain. **(B)**
Quantification of percentage micrograph area covered by fluorescent cells. **(C)** Fluorescent
micrographs of 16HBE cells after 15 min exposure to conditioned medium from *plcH* transposon
mutants in the presence of Sytox Orange nucleic acid stain (red). Cells were prelabeled with
Hoechst 33342 nucleic acid stain (blue). **(D)** Quantification of Sytox Orange mean fluorescence
intensity normalized to that of Hoechst 33342 in. **(E)** Alignment of MexT protein sequences from
MPAO1*plcH* (PW2536), MPAO1*plcH* (PW2537), MPAO1*pvdF*, PAO1, or PAO1*ΔpvdF*. MexT
sequences were acquired from paired-end Illumina sequencing (PW2536, PW2537, PAO1,
PAO1*ΔpvdF*) or PCR-Sanger sequencing (MPAO1*pvdF*). Amino acid substitution sites are
highlighted yellow. Amino acid substitutions in PW2536 are labeled red. **(F)** *mexE* and *mexF*
mRNA levels compared to that of WT PAO1. mRNA levels were measured by qRT-PCR. **(G)**
Rhamnolipid content in conditioned medium from 14 h EMEM cultures measured by FM 1-43
labeling. All error bars represent SEM from three biological replicates. * corresponds to $p < 0.01$
based on Student's *t*-test **(B, D)** or one-way ANOVA with Tukey's multiple comparisons test **(G)**.

Fig. 4. 5-Fluorocytosine (5-FC) inhibits pyoverdine and rhamnolipid production in highly
virulent cystic fibrosis isolates. (A, B) Pyoverdine production **(A)** or *C. elegans* death **(B)** by
multidrug-resistant *P. aeruginosa* strains isolated from pediatric cystic fibrosis patients. Black bar
represents the median pyoverdine production or *C. elegans death*. Red bar represents PA2-72, blue
bar represents PA2-61. Survey data was adapted from (22). **(C)** Bacterial growth of PAO1, PA2-

72, or PA2-61 in EMEM with or without 100 µM 5-FC. **(D)** Photograph of EMEM culture after

18 h incubation. **(E, F)** Pyoverdine **(E)** or rhamnolipid **(F)** production of PAO1, PA2-72, or PA2-

61 in EMEM with or without 100 µM 5-FC. **(G)** Fluorescent micrographs of 16HBE cells after 40

min (PAO1, PA2-72) or 20 min (PA2-61) exposure to EMEM conditioned medium. Cells were

prelabeled with CellMask Orange plasma membrane stain. **(H)** Fluorescent micrographs of 16HBE

cells after 20 min (PAO1, PA2-72) or 10 min (PA2-61) exposure to EMEM conditioned medium

in the presence of Sytox Orange nucleic acid stain (red). Cells were prelabeled with Hoechst 33342

nucleic acid stain (blue). **(I)** Quantification of percentage micrograph area covered by fluorescent

cells in **(G)**. **(J)** Quantification of Sytox Orange mean fluorescence intensity normalized to that of

Hoechst 33342 in **(H)**. All error bars represent SEM from four biological replicates. * corresponds

to $p < 0.01$ and NS corresponds to $p > 0.05$ based on one-way ANOVA with Sidak's multiple

comparisons test.

Fig. 5. Pyoverdine translocates into 16HBE cells and localizes to early endosomes. (A)

Summary of the pyoverdine purification pipeline. **(B)** Representative chromatogram from the

HPLC purification step of the pipeline. Red box depicts the predominant pyoverdine-containing

fractions that were collected. **(C)** Analysis of the final purified product via HPLC. **(D)** 16HBE cell

viability after 48 h treatment with 100 µM ciclopirox olamine, 1, 10-phenanthroline, or pyochelin,

or 72 h treatment of 100 µM pyoverdine or deferoxamine in serum-free EMEM. **(E)** Confocal

micrographs of 16HBE cells exposed to 100 µM purified pyoverdine, pyoverdine with excess

Ga(NO$_3$)$_3$, pyoverdine with excess FeCl$_3$, or media control for 24 h. Cells were trypsinated prior

to imaging. **(F)** Quantification of pyoverdine fluorescence within 30 individual cells. **(G)** Confocal

micrographs of 16HBE cells treated with 100 µM pyoverdine-gallium and dextran-Texas Red

(10,000 MW). Error bars in **(D)** represent SEM from three biological replicates. Error bars in **(C)**

represent standard deviation. * corresponds to $p < 0.01$ based on one-way ANOVA with Dunnett's

multiple comparisons test.

Fig. 6. Small molecule iron chelators promote the expression of proinflammatory genes in

16HBE cells. (A) Total RNA yield in 16HBE cells treated with ciclopirox olamine (CPX), 1,10-

phenanthroline (PHE), or pyochelin (PCH) for 30 h or cells treated with pyoverdine (PVD) or

deferoxamine (DFO) for 60 h normalized to that of media control. All treatments were at 100 µM

in serum-free EMEM. **(B)** Proinflammatory gene expression (*NLRP3, IL1B, NLRP1, IL8, TNF*) in

cells treated with iron chelators compared to that of media control. mRNA levels were measured

by qRT-PCR. **(C)** IL-8 protein concentration in the supernatants of 16HBE cells treated with iron

chelators. IL-8 was quantified by ELISA. **(D, E)** Total RNA yield **(D)** or proinflammatory gene

expression **(E)** in 16HBE cells treated with iron chelators with or without excess $Ga(NO_3)_3$

supplementation. **(F)** Proinflammatory gene expression after 60 h treatment with pyoverdine,

deferoxamine, or both molecules. All error bars represent SEM from three biological replicates. *

corresponds to $p < 0.01$, # corresponds to $p < 0.05$, and NS corresponds to $p > 0.05$ based on one-

way ANOVA with Dunnett's **(A, C)**, Sidak's **(D, E)**, or Tukey's **(F)** multiple comparisons test.

Supplemental Figure Legends:

Fig. S1. Disruption of pyochelin biosynthesis does not mitigate epithelial damage. (A, B)

Bacterial growth **(A)** or pyoverdine production **(B)** of *P. aeruginosa* PAO1 siderophore

biosynthetic mutants (PAO1*ΔpvdF* – pyoverdine; PAO1*ΔpchBA* – pyochelin;

PAO1*ΔpvdFΔpchBA* – pyoverdine and pyochelin) after 18 h growth in serum-free EMEM. **(C)**

MH-S murine alveolar macrophage viability after 2 h exposure to conditioned medium from PAO1

siderophore mutants grown in EMEM. Cell viability was measured using a resazurin-based assay.

(D) Fluorescent micrographs of 16HBE cells after 30 min exposure to conditioned medium from

PAO1 siderophore mutants. Cells were prelabeled with CellMask Orange plasma membrane stain.

(E) Quantification of percentage micrograph area covered by fluorescent cells. All error bars

represent SEM from four biological replicates. * corresponds to $p < 0.01$ and NS corresponds to p

> 0.05 based on one-way ANOVA with Tukey's multiple comparisons test.

Fig. S2. Lipid extraction abrogates epithelial damage by pyoverdine-rich conditioned medium. **(A)** Fluorescent micrographs of 16HBE cells after 30 min exposure to conditioned medium from WT PAO1. Conditioned medium was pretreated with 200 µM $Ga(NO_3)_3$, 200 µM $FeCl_3$, or 100 µg/mL proteinase K for 24 h, had macromolecules depleted via a 10 kDa centrifugal filter, or had lipids depleted by chloroform ($CHCl_3$) extraction. Cells were prelabeled with CellMask Orange plasma membrane stain. **(B)** Quantification of percentage micrograph area covered by fluorescent cells. Error bars represent SEM from three biological replicates. * corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on one-way ANOVA with Dunnett's multiple comparisons test.

Fig. S3. Type II secretion system toxins do not contribute to lung epithelial damage. (A) Fluorescent micrographs of 16HBE cells after 30 min exposure to conditioned medium from MPAO1 transposon mutants. Cells were prelabeled with CellMask Orange plasma membrane stain. **(B)** Fluorescent micrographs of 16HBE cells after 15 min exposure to conditioned medium from MPAO1 transposon mutants in the presence of Sytox Orange nucleic acid stain (red). Cells were

prelabeled with Hoechst 33342 nucleic acid stain (blue). **(C)** Quantification of percentage

micrograph area covered by fluorescent cells in **(A)**. **(D)** Quantification of Sytox Orange mean

fluorescence intensity normalized to that of Hoechst 33342 in **(B)**. **(E)** Protease activity in

conditioned medium from MPAO1 transposon mutants. Proteolytic activity was measured by

fluorescence release from cleavage of FITC-conjugated casein. All error bars represent SEM from

three biological replicates. * corresponds to $p < 0.01$ and NS corresponds to $p > 0.05$ based on

one-way ANOVA with Dunnett's multiple comparisons test.

Fig. S4. Pyoverdine biosynthetic mutants do not exhibit impaired C4-HSL quorum-sensing.

(A) Bioluminescence produced by *E. coli* JM109 pSB536 (N-butanoyl-L-homoserine lactone

reporter strain) grown in media supplemented with conditioned medium from wild-type PAO1 or

PAO1*ΔpvdF*. **(B)** Bioluminescence at the kinetics **(A)** endpoint. **(C)** Bioluminescence produced

by the *E. coli* reporter strain grown in media supplemented with conditioned medium from

MPAO1 transposon mutants. **(D)** Bioluminescence at the kinetics **(C)** endpoint. All error bars

represent SEM from at least three biological replicates. * corresponds to $p < 0.01$ and NS

corresponds to $p > 0.05$ based on one-way ANOVA with Tukey's multiple comparisons test.

Fig. S5. Purified rhamnolipids are sufficient for 16HBE cell death but not for epithelial

monolayer damage. (A) Quantification of rhamnolipids in purified samples (RHL) or *P.*

aeruginosa conditioned medium from WT PAO1, PAO1*ΔpvdF*, MPAO1*cat*, or MPAO1*pvdF*. **(B)**

Fluorescent micrographs of 16HBE cells after 30 min exposure to purified rhamnolipids in EMEM.

Cells were prelabeled with CellMask Orange plasma membrane stain. **(C)** Fluorescent

micrographs of 16HBE cells after 15 min exposure to purified rhamnolipids in EMEM in the

presence of Sytox Orange nucleic acid stain (red). Cells were prelabeled with Hoechst 33342

nucleic acid stain (blue). **(D)** Quantification of percentage micrograph area covered by fluorescent

cells in **(B)**. **(E)** Quantification of Sytox Orange mean fluorescence intensity normalized to that of

Hoechst 33342 in **(C)**. **(F)** Quantification of rhamnolipids in conditioned medium from WT PAO1

or conditioned medium after heat denaturation (85 °C for 1 h). **(G)** Fluorescent micrographs of

16HBE cells after 30 min exposure to conditioned medium from WT PAO1 or conditioned

medium after heat denaturation. Cells were prelabeled with CellMask Orange plasma membrane

stain. **(H)** Fluorescent micrographs of 16HBE cells after 15 min exposure to conditioned medium

from WT PAO1 or conditioned medium after heat denaturation in the presence of Sytox Orange

nucleic acid stain (red). Cells were prelabeled with Hoechst 33342 nucleic acid stain (blue). **(I)**

Quantification of percentage micrograph area covered by fluorescent cells in **(G)**. **(J)**

Quantification of Sytox Orange mean fluorescence intensity normalized to that of Hoechst 33342

in **(H)**. All error bars represent SEM from three biological replicates. # corresponds to $p < 0.05$

and NS corresponds to $p > 0.05$ based on Student's *t*-test.

Fig. S6. Pyoverdine accumulates in early endosomes of lung epithelial cells. (A, B) 16HBE

cell viability after 48 **(A)** or 72 h **(B)** treatment with ciclopirox olamine, 1, 10-phenanthroline,

pyoverdine, or deferoxamine in serum-free EMEM. **(C)** Confocal micrographs of 16HBE cells

exposed to 100 µM purified pyoverdine, pyoverdine with excess $Ga(NO_3)_3$, pyoverdine with

excess $FeCl_3$, or media control for 24 h. Cells were labeled with CellMask Deep Red plasma

membrane stain and trypsinated prior to imaging. **(D)** Confocal micrographs of 16HBE cells

treated with 100 µM pyoverdine-gallium and dextran-Texas Red (10,000 MW). Bottom row shows

an enlarged micrograph of one representative cell. All error bars represent SEM from three

biological replicates.

Fig. S7. Small molecule iron chelators promote IL-8 production in 16HBE cells. (A) *IL8*

mRNA levels in 16HBE cells treated with iron chelators (ciclopirox olamine – CPX, 1,10-

phenanthroline – PHE, pyochelin – PCH, pyoverdine – PVD) for 30 h compared to that of media

control. mRNA levels were measured by qRT-PCR. **(B)** IL-8 protein concentration in the

supernatants of 16HBE cells treated with iron chelators for 48 h. IL-8 was quantified by ELISA.

(C) Correlation between *IL8* gene expression and IL-8 protein production. All error bars represent

SEM from three biological replicates.

Fig. S8. Deferoxamine promotes the expression of proinflammatory genes in 16HBE CFTR

mutants. (A) Total RNA yield in wild-type 16HBE cells and 16HBE cells carrying mutations

(G551D, ΔF508) in the cystic fibrosis transmembrane conductance regulator (CFTR) after 48 h

treatment with 100 µM deferoxamine in serum-free EMEM. **(B)** Expression of proinflammatory

genes (*NLRP3, IL1B, NLRP1, IL8, TNF*) or iron-regulated genes (*NDRG1*, *TFRC*) in 16HBE cells

treated with deferoxamine compared to that of media control. mRNA levels were measured by

qRT-PCR. All error bars represent SEM from three biological replicates.

References:

1.   Kollef MH, Chastre J, Fagon JY, Francois B, Niederman MS, Rello J, Torres A, Vincent

JL, Wunderink RG, Go KW, Rehm C. 2014. Global prospective epidemiologic and

surveillance study of ventilator-associated pneumonia due to Pseudomonas aeruginosa.

Crit Care Med 42:2178-87.

2.   Lyczak JB, Cannon CL, Pier GB. 2002. Lung infections associated with cystic fibrosis.

Clin Microbiol Rev 15:194-222.

3.   Bhagirath AY, Li Y, Somayajula D, Dadashi M, Badr S, Duan K. 2016. Cystic fibrosis

lung environment and Pseudomonas aeruginosa infection. BMC Pulm Med 16:174.

4.   Hassett DJ, Borchers MT, Panos RJ. 2014. Chronic obstructive pulmonary disease

(COPD): evaluation from clinical, immunological and bacterial pathogenesis

perspectives. J Microbiol 52:211-26.

5.   Murphy TF. 2009. Pseudomonas aeruginosa in adults with chronic obstructive pulmonary

disease. Curr Opin Pulm Med 15:138-42.

6.   Anderson GG, O'Toole GA. 2008. Innate and induced resistance mechanisms of bacterial

biofilms. Curr Top Microbiol Immunol 322:85-105.

7.   Moreau-Marquis S, Stanton BA, O'Toole GA. 2008. Pseudomonas aeruginosa biofilm

formation in the cystic fibrosis airway. Pulm Pharmacol Ther 21:595-9.

8.   Curran CS, Bolig T, Torabi-Parizi P. 2018. Mechanisms and Targeted Therapies for

Pseudomonas aeruginosa Lung Infection. Am J Respir Crit Care Med 197:708-727.

9.   Cézard C, Farvacques N, Sonnet P. 2015. Chemistry and biology of pyoverdines,

Pseudomonas primary siderophores. Curr Med Chem 22:165-86.

10.   Vasil ML, Ochsner UA. 1999. The response of Pseudomonas aeruginosa to iron:

genetics, biochemistry and virulence. Mol Microbiol 34:399-413.

11.   Cornelis P, Dingemans J. 2013. Pseudomonas aeruginosa adapts its iron uptake strategies

in function of the type of infections. Front Cell Infect Microbiol 3:75.

12.   Visca P, Imperi F, Lamont IL. 2007. Pyoverdine siderophores: from biogenesis to

biosignificance. Trends Microbiol 15:22-30.

13.   Xiao R, Kisaalita WS. 1997. Iron acquisition from transferrin and lactoferrin by

Pseudomonas aeruginosa pyoverdin. Microbiology 143 (Pt 7):2509-15.

14.   Dumas Z, Ross-Gillespie A, Kummerli R. 2013. Switching between apparently redundant

iron-uptake mechanisms benefits bacteria in changeable environments. Proc Biol Sci

280:20131055.

15.   Banin E, Vasil ML, Greenberg EP. 2005. Iron and Pseudomonas aeruginosa biofilm

formation. Proc Natl Acad Sci U S A 102:11076-81.

16.   Kang D, Kirienko NV. 2018. Interdependence between iron acquisition and biofilm

formation in Pseudomonas aeruginosa. J Microbiol 56:449-457.

17.   Minandri F, Imperi F, Frangipani E, Bonchi C, Visaggio D, Facchini M, Pasquali P,

Bragonzi A, Visca P. 2016. Role of Iron Uptake Systems in Pseudomonas aeruginosa

Virulence and Airway Infection. Infect Immun 84:2324-35.

18.   Lamont IL, Beare PA, Ochsner U, Vasil AI, Vasil ML. 2002. Siderophore-mediated

signaling regulates virulence factor production in Pseudomonasaeruginosa. Proc Natl

Acad Sci U S A 99:7072-7.

19.   Kirienko NV, Kirienko DR, Larkins-Ford J, Wählby C, Ruvkun G, Ausubel FM. 2013.

Pseudomonas aeruginosa disrupts Caenorhabditis elegans iron homeostasis, causing a

hypoxic response and death. Cell Host Microbe 13:406-16.

20.   Kirienko NV, Ausubel FM, Ruvkun G. 2015. Mitophagy confers resistance to

siderophore-mediated killing by Pseudomonas aeruginosa. Proc Natl Acad Sci U S A

112:1821-6.

21.   Kang D, Kirienko DR, Webster P, Fisher AL, Kirienko NV. 2018. Pyoverdine, a

739   siderophore from Pseudomonas aeruginosa, translocates into C. elegans, removes iron,

740   and activates a distinct host response. Virulence 9:804-817.

741   22.   Kang D, Revtovich AV, Chen Q, Shah KN, Cannon CL, Kirienko NV. 2019. Pyoverdine-

742   Dependent Virulence of Pseudomonas aeruginosa Isolates From Cystic Fibrosis Patients.

743   Front Microbiol 10:2048.

744   23.   Imperi F, Massai F, Facchini M, Frangipani E, Visaggio D, Leoni L, Bragonzi A, Visca

745   P. 2013. Repurposing the antimycotic drug flucytosine for suppression of Pseudomonas

746   aeruginosa pathogenicity. Proc Natl Acad Sci U S A 110:7458-63.

747   24.   Takase H, Nitanai H, Hoshino K, Otani T. 2000. Impact of siderophore production on

748   Pseudomonas aeruginosa infections in immunosuppressed mice. Infect Immun 68:1834-

749   9.

750   25.   Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. 1996. Pyoverdin is essential for

751   virulence of Pseudomonas aeruginosa. Infect Immun 64:518-23.

752   26.   Kang D, Kirienko NV. 2020. An In Vitro Cell Culture Model for Pyoverdine-Mediated

753   Virulence. Pathogens 10.

754   27.   Xu Q, Kang D, Meyer MD, Pennington CL, Gopal C, Schertzer JW, Kirienko NV. 2023.

755   Cytotoxic rhamnolipid micelles drive acute virulence in Pseudomonas aeruginosa.

756   bioRxiv doi:10.1101/2023.10.13.562257:2023.10.13.562257.

757   28.   Wu Y, Yeh FL, Mao F, Chapman ER. 2009. Biophysical characterization of styryl dye-

758   membrane interactions. Biophys J 97:101-9.

759   29.   Jensen PO, Bjarnsholt T, Phipps R, Rasmussen TB, Calum H, Christoffersen L, Moser C,

760   Williams P, Pressler T, Givskov M, Hoiby N. 2007. Rapid necrotic killing of

polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of

rhamnolipid by Pseudomonas aeruginosa. Microbiology (Reading) 153:1329-1338.

30.   Pearson JP, Pesci EC, Iglewski BH. 1997. Roles of Pseudomonas aeruginosa las and rhl

quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes. J

Bacteriol 179:5756-67.

31.   Winson MK, Swift S, Fish L, Throup JP, Jorgensen F, Chhabra SR, Bycroft BW,

Williams P, Stewart GS. 1998. Construction and analysis of luxCDABE-based plasmid

sensors for investigating N-acyl homoserine lactone-mediated quorum sensing. FEMS

Microbiol Lett 163:185-92.

32.   Lee J, Zhang L. 2015. The hierarchy quorum sensing network in Pseudomonas

aeruginosa. Protein Cell 6:26-41.

33.   Filloux A. 2011. Protein Secretion Systems in Pseudomonas aeruginosa: An Essay on

Diversity, Evolution, and Function. Front Microbiol 2:155.

34.   Kohler T, Epp SF, Curty LK, Pechere JC. 1999. Characterization of MexT, the regulator

of the MexE-MexF-OprN multidrug efflux system of Pseudomonas aeruginosa. J

Bacteriol 181:6300-5.

35.   Kohler T, van Delden C, Curty LK, Hamzehpour MM, Pechere JC. 2001. Overexpression

of the MexEF-OprN multidrug efflux system affects cell-to-cell signaling in

Pseudomonas aeruginosa. J Bacteriol 183:5213-22.

36.   Vaillancourt M, Limsuwannarot SP, Bresee C, Poopalarajah R, Jorth P. 2021.

Pseudomonas aeruginosa mexR and mexEF Antibiotic Efflux Pump Variants Exhibit

Increased Virulence. Antibiotics (Basel) 10.

37.   Kirienko DR, Revtovich AV, Kirienko NV. 2016. A High-Content, Phenotypic Screen

784   Identifies Fluorouridine as an Inhibitor of Pyoverdine Biosynthesis and Pseudomonas

785   aeruginosa Virulence. mSphere 1.

786   38.   Kang D, Revtovich AV, Deyanov AE, Kirienko NV. 2021. Pyoverdine Inhibitors and

787   Gallium Nitrate Synergistically Affect Pseudomonas aeruginosa. mSphere

788   doi:10.1128/mSphere.00401-21:e0040121.

789   39.   Costabile G, d'Angelo I, d'Emmanuele di Villa Bianca R, Mitidieri E, Pompili B, Del

790   Porto P, Leoni L, Visca P, Miro A, Quaglia F, Imperi F, Sorrentino R, Ungaro F. 2016.

791   Development of inhalable hyaluronan/mannitol composite dry powders for flucytosine

792   repositioning in local therapy of lung infections. J Control Release 238:80-91.

793   40.   Oliver JM, Berlin RD, Davis BH. 1984. Use of horseradish peroxidase and fluorescent

794   dextrans to study fluid pinocytosis in leukocytes. Methods Enzymol 108:336-47.

795   41.   Holden VI, Lenio S, Kuick R, Ramakrishnan SK, Shah YM, Bachman MA. 2014.

796   Bacterial siderophores that evade or overwhelm lipocalin 2 induce hypoxia inducible

797   factor 1α and proinflammatory cytokine secretion in cultured respiratory epithelial cells.

798   Infect Immun 82:3826-36.

799   42.   Lee HJ, Lee J, Lee SK, Lee SK, Kim EC. 2007. Differential regulation of iron chelator-

800   induced IL-8 synthesis via MAP kinase and NF-kappaB in immortalized and malignant

801   oral keratinocytes. BMC Cancer 7:176.

802   43.   Choi EY, Kim EC, Oh HM, Kim S, Lee HJ, Cho EY, Yoon KH, Kim EA, Han WC, Choi

803   SC, Hwang JY, Park C, Oh BS, Kim Y, Kimm KC, Park KI, Chung HT, Jun CD. 2004.

804   Iron chelator triggers inflammatory signals in human intestinal epithelial cells:

involvement of p38 and extracellular signal-regulated kinase signaling pathways. J

Immunol 172:7069-77.

44.   Moldoveanu B, Otmishi P, Jani P, Walker J, Sarmiento X, Guardiola J, Saad M, Yu J.

2009. Inflammatory mechanisms in the lung. J Inflamm Res 2:1-11.

45.   Hauser AR. 2009. The type III secretion system of Pseudomonas aeruginosa: infection by

injection. Nat Rev Microbiol 7:654-65.

46.   Michalska M, Wolf P. 2015. Pseudomonas Exotoxin A: optimized by evolution for

effective killing. Front Microbiol 6:963.

47.   Hall S, McDermott C, Anoopkumar-Dukie S, McFarland AJ, Forbes A, Perkins AV,

Davey AK, Chess-Williams R, Kiefel MJ, Arora D, Grant GD. 2016. Cellular Effects of

Pyocyanin, a Secreted Virulence Factor of Pseudomonas aeruginosa. Toxins (Basel) 8.

48.   Thi MTT, Wibowo D, Rehm BHA. 2020. Pseudomonas aeruginosa Biofilms. Int J Mol

Sci 21.

49.   Ochsner UA, Johnson Z, Lamont IL, Cunliffe HE, Vasil ML. 1996. Exotoxin A

production in Pseudomonas aeruginosa requires the iron-regulated pvdS gene encoding

an alternative sigma factor. Mol Microbiol 21:1019-28.

50.   Plotkowski MC, Povoa HC, Zahm JM, Lizard G, Pereira GM, Tournier JM, Puchelle E.

2002. Early mitochondrial dysfunction, superoxide anion production, and DNA

degradation are associated with non-apoptotic death of human airway epithelial cells

induced by Pseudomonas aeruginosa exotoxin A. Am J Respir Cell Mol Biol 26:617-26.

51.   Azghani AO. 1996. Pseudomonas aeruginosa and epithelial permeability: role of

virulence factors elastase and exotoxin A. Am J Respir Cell Mol Biol 15:132-40.

52.   Hirakata Y, Furuya N, Tateda K, Kaku M, Yamaguchi K. 1993. In vivo production of

exotoxin A and its role in endogenous Pseudomonas aeruginosa septicemia in mice.

Infect Immun 61:2468-73.

53.   Pillar CM, Hobden JA. 2002. Pseudomonas aeruginosa exotoxin A and keratitis in mice.

Invest Ophthalmol Vis Sci 43:1437-44.

54.   Miyazaki S, Matsumoto T, Tateda K, Ohno A, Yamaguchi K. 1995. Role of exotoxin A

in inducing severe Pseudomonas aeruginosa infections in mice. J Med Microbiol 43:169-

75.

55.   Malloy JL, Veldhuizen RA, Thibodeaux BA, O'Callaghan RJ, Wright JR. 2005.

Pseudomonas aeruginosa protease IV degrades surfactant proteins and inhibits surfactant

host defense and biophysical functions. Am J Physiol Lung Cell Mol Physiol 288:L409-

18.

56.   Bradshaw JL, Caballero AR, Bierdeman MA, Adams KV, Pipkins HR, Tang A,

O'Callaghan RJ, McDaniel LS. 2018. Pseudomonas aeruginosa Protease IV Exacerbates

Pneumococcal Pneumonia and Systemic Disease. mSphere 3.

57.   Guillon A, Brea D, Morello E, Tang A, Jouan Y, Ramphal R, Korkmaz B, Perez-Cruz M,

Trottein F, O'Callaghan RJ, Gosset P, Si-Tahar M. 2017. Pseudomonas aeruginosa

proteolytically alters the interleukin 22-dependent lung mucosal defense. Virulence

8:810-820.

58.   Engel LS, Hobden JA, Moreau JM, Callegan MC, Hill JM, O'Callaghan RJ. 1997.

Pseudomonas deficient in protease IV has significantly reduced corneal virulence. Invest

Ophthalmol Vis Sci 38:1535-42.

59.   Engel LS, Hill JM, Moreau JM, Green LC, Hobden JA, O'Callaghan RJ. 1998.

850   Pseudomonas aeruginosa protease IV produces corneal damage and contributes to

851   bacterial virulence. Invest Ophthalmol Vis Sci 39:662-5.

852   60.   Gdaniec BG, Bonini F, Prodon F, Braschler T, Kohler T, van Delden C. 2022.

853   Pseudomonas aeruginosa rhamnolipid micelles deliver toxic metabolites and antibiotics

854   into Staphylococcus aureus. iScience 25:103669.

855   61.   McClure CD, Schiller NL. 1992. Effects of Pseudomonas aeruginosa rhamnolipids on

856   human monocyte-derived macrophages. J Leukoc Biol 51:97-102.

857   62.   Van Gennip M, Christensen LD, Alhede M, Phipps R, Jensen PO, Christophersen L,

858   Pamp SJ, Moser C, Mikkelsen PJ, Koh AY, Tolker-Nielsen T, Pier GB, Hoiby N,

859   Givskov M, Bjarnsholt T. 2009. Inactivation of the rhlA gene in Pseudomonas aeruginosa

860   prevents rhamnolipid production, disabling the protection against polymorphonuclear

861   leukocytes. APMIS 117:537-46.

862   63.   Zulianello L, Canard C, Kohler T, Caille D, Lacroix JS, Meda P. 2006. Rhamnolipids are

863   virulence factors that promote early infiltration of primary human airway epithelia by

864   Pseudomonas aeruginosa. Infect Immun 74:3134-47.

865   64.   Read RC, Roberts P, Munro N, Rutman A, Hastie A, Shryock T, Hall R, McDonald-

866   Gibson W, Lund V, Taylor G, et al. 1992. Effect of Pseudomonas aeruginosa

867   rhamnolipids on mucociliary transport and ciliary beating. J Appl Physiol (1985)

868   72:2271-7.

869   65.   Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, Kumar N,

870   Schembri MA, Song Z, Kristoffersen P, Manefield M, Costerton JW, Molin S, Eberl L,

Steinberg P, Kjelleberg S, Høiby N, Givskov M. 2003. Attenuation of Pseudomonas

aeruginosa virulence by quorum sensing inhibitors. EMBO J 22:3803-15.

66.   Kang D, Turner KE, Kirienko NV. 2017. PqsA Promotes Pyoverdine Production via

Biofilm Formation. Pathogens 7.

67.   Stintzi A, Evans K, Meyer JM, Poole K. 1998. Quorum-sensing and siderophore

biosynthesis in Pseudomonas aeruginosa: lasR/lasI mutants exhibit reduced pyoverdine

biosynthesis. FEMS Microbiol Lett 166:341-5.

68.   Thomann A, de Mello Martins AG, Brengel C, Empting M, Hartmann RW. 2016.

Application of Dual Inhibition Concept within Looped Autoregulatory Systems toward

Antivirulence Agents against Pseudomonas aeruginosa Infections. ACS Chem Biol

11:1279-86.

69.   R YR-R, Salvador MJ. 2020. Phenotypic detection of quorum sensing inhibition in

Pseudomonas aeruginosa pyoverdine and swarming by volatile organic products. Future

Microbiol 15:1147-1156.

70.   Holden VI, Breen P, Houle S, Dozois CM, Bachman MA. 2016. Klebsiella pneumoniae

Siderophores Induce Inflammation, Bacterial Dissemination, and HIF-1α Stabilization

during Pneumonia. MBio 7.

71.   Marvig RL, Damkiær S, Khademi SM, Markussen TM, Molin S, Jelsbak L. 2014.

Within-host evolution of Pseudomonas aeruginosa reveals adaptation toward iron

acquisition from hemoglobin. MBio 5:e00966-14.

72.   De Vos D, De Chial M, Cochez C, Jansen S, Tummler B, Meyer JM, Cornelis P. 2001.

Study of pyoverdine type and production by Pseudomonas aeruginosa isolated from

cystic fibrosis patients: prevalence of type II pyoverdine isolates and accumulation of

pyoverdine-negative mutations. Arch Microbiol 175:384-8.

73. Andersen SB, Marvig RL, Molin S, Krogh Johansen H, Griffin AS. 2015. Long-term

social dynamics drive loss of function in pathogenic bacteria. Proc Natl Acad Sci U S A

112:10756-61.

74. Martin LW, Reid DW, Sharples KJ, Lamont IL. 2011. Pseudomonas siderophores in the

sputum of patients with cystic fibrosis. Biometals 24:1059-67.

75. Haas B, Murphy E, Castignetti D. 1991. Siderophore synthesis by mucoid Pseudomonas

aeruginosa strains isolated from cystic fibrosis patients. Can J Microbiol 37:654-7.

76. Mayer-Hamblett N, Rosenfeld M, Gibson RL, Ramsey BW, Kulasekara HD, Retsch-

Bogart GZ, Morgan W, Wolter DJ, Pope CE, Houston LS, Kulasekara BR, Khan U,

Burns JL, Miller SI, Hoffman LR. 2014. Pseudomonas aeruginosa in vitro phenotypes

distinguish cystic fibrosis infection stages and outcomes. Am J Respir Crit Care Med

190:289-97.

77. Kang D, Zhang L, Kirienko NV. 2021. High-Throughput Approaches for the

Identification of Pseudomonas aeruginosa Antivirulents. mBio 12.

78. Kaneko Y, Thoendel M, Olakanmi O, Britigan BE, Singh PK. 2007. The transition metal

gallium disrupts Pseudomonas aeruginosa iron metabolism and has antimicrobial and

antibiofilm activity. The Journal of Clinical Investigation 117:877-888.

79. Goss CH, Kaneko Y, Khuu L, Anderson GD, Ravishankar S, Aitken ML, Lechtzin N,

Zhou G, Czyz DM, McLean K, Olakanmi O, Shuman HA, Teresi M, Wilhelm E,

Caldwell E, Salipante SJ, Hornick DB, Siehnel RJ, Becker L, Britigan BE, Singh PK.

2018. Gallium disrupts bacterial iron metabolism and has therapeutic effects in mice and

humans with lung infections. Sci Transl Med 10.

80.   Britigan BE, Rasmussen GT, Cox CD. 1994. Pseudomonas siderophore pyochelin

enhances neutrophil-mediated endothelial cell injury. Am J Physiol 266:L192-8.

81.   Ong KS, Cheow YL, Lee SM. 2017. The role of reactive oxygen species in the

antimicrobial activity of pyochelin. J Adv Res 8:393-398.

82.   Adler C, Corbalan NS, Seyedsayamdost MR, Pomares MF, de Cristobal RE, Clardy J,

Kolter R, Vincent PA. 2012. Catecholate siderophores protect bacteria from pyochelin

toxicity. PLoS One 7:e46754.

83.   Jenul C, Keim KC, Jens JN, Zeiler MJ, Schilcher K, Schurr MJ, Melander C, Phelan VV,

Horswill AR. 2023. Pyochelin biotransformation by Staphylococcusaureus shapes

bacterial competition with Pseudomonas aeruginosa in polymicrobial infections. Cell Rep

42:112540.

84.   Kruger P, Saffarzadeh M, Weber AN, Rieber N, Radsak M, von Bernuth H, Benarafa C,

Roos D, Skokowa J, Hartl D. 2015. Neutrophils: Between host defence, immune

modulation, and tissue injury. PLoS Pathog 11:e1004651.

85.   Twigg MS, Brockbank S, Lowry P, FitzGerald SP, Taggart C, Weldon S. 2015. The Role

of Serine Proteases and Antiproteases in the Cystic Fibrosis Lung. Mediators Inflamm

2015:293053.

86.   Cantin AM, Hartl D, Konstan MW, Chmiel JF. 2015. Inflammation in cystic fibrosis lung

disease: Pathogenesis and therapy. J Cyst Fibros 14:419-30.

87.   Hosseinian N, Cho Y, Lockey RF, Kolliputi N. 2015. The role of the NLRP3

inflammasome in pulmonary diseases. Ther Adv Respir Dis 9:188-97.

88.   McElvaney OJ, Zaslona Z, Becker-Flegler K, Palsson-McDermott EM, Boland F,

Gunaratnam C, Gulbins E, O'Neill LA, Reeves EP, McElvaney NG. 2019. Specific

Inhibition of the NLRP3 Inflammasome as an Antiinflammatory Strategy in Cystic

Fibrosis. Am J Respir Crit Care Med 200:1381-1391.

89.   McKee CM, Coll RC. 2020. NLRP3 inflammasome priming: A riddle wrapped in a

mystery inside an enigma. J Leukoc Biol 108:937-952.

90.   Estivill X, Bancells C, Ramos C. 1997. Geographic distribution and regional origin of

272 cystic fibrosis mutations in European populations. The Biomed CF Mutation

Analysis Consortium. Hum Mutat 10:135-54.

91.   de Albuquerque Wanderley Sales V, Timoteo TRR, da Silva NM, de Melo CG, Ferreira

AS, de Oliveira MVG, de Oliveira Silva E, Dos Santos Mendes LM, Rolim LA, Neto

PJR. 2021. A Systematic Review of the Anti-inflammatory Effects of Gallium

Compounds. Curr Med Chem 28:2062-2076.

92.   Zhang C, Yang B, Biazik JM, Webster RF, Xie W, Tang J, Allioux FM, Abbasi R,

Mousavi M, Goldys EM, Kilian KA, Chandrawati R, Esrafilzadeh D, Kalantar-Zadeh K.

2022. Gallium Nanodroplets are Anti-Inflammatory without Interfering with Iron

Homeostasis. ACS Nano 16:8891-8903.

93.   Apseloff G. 1999. Therapeutic uses of gallium nitrate: past, present, and future. Am J

Ther 6:327-39.

94.   Jacobs MA, Alwood A, Thaipisuttikul I, Spencer D, Haugen E, Ernst S, Will O, Kaul R,

Raymond C, Levy R, Chun-Rong L, Guenthner D, Bovee D, Olson MV, Manoil C. 2003.

Comprehensive transposon mutant library of Pseudomonas aeruginosa. Proc Natl Acad

Sci U S A 100:14339-44.

95.   Liberati NT, Urbach JM, Miyata S, Lee DG, Drenkard E, Wu G, Villanueva J, Wei T,

Ausubel FM. 2006. An ordered, nonredundant library of Pseudomonas aeruginosa strain

PA14 transposon insertion mutants. Proc Natl Acad Sci U S A 103:2833-8.

96.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. J Comput Biol 19:455-77.

97.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

30:2068-9.

98.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved

microbes from next-generation sequencing data using breseq. Methods Mol Biol

1151:165-88.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6