Redefining hematopoietic progenitors cells and reaffirming lineage commitment by transcriptional atlas

Yongjian Yue[1]

[1]Shenzhen Institute of Respiratory Disease, the First Affiliated Hospital of Southern University of Science and Technology, Shenzhen, Guangdong, China

* Correspondence: Yongjian Yue
Address: Shenzhen People's Hospital, The First Affiliated Hospital of Southern University of Science and Technology, No. 1017 Dongmen North Road, Luohu District, Shenzhen 518020, China.
Email: yueyongj@163.com
Tel: +86 15919909356

Abstract

Tracing the differentiation of the hematopoietic progenitor cells(HPCs) in human blood, particularly with the scarcity in adults have been challenging owing to unable capture the whole differentiation stages and subgroups of HPCs. Deciphering the hematopoiesis mechanism that underlies progenitor cells definition and fate decision are critical to understanding fundamental questions of hematopoietic lineage commitment. Often, the use of classic and traditional surface markers lacks the sensitivity and specificity to clearly define the hematopoietic progenitors. The lineage definition and commitment of hematopoietic progenitors in adult peripheral blood remains to be elucidated by a reliable transcriptional atlas. Here, single cell RNA sequencing (scRNA seq) was used to define the lineage of HPCs and reaffirm the hierarchy of hematopoiesis in adults peripheral blood. A total of 10450 adults peripheral blood CD34+ cells were enriched and sequenced, including most differentiation stages and subgroups of HPCs. HPCs lineage were redefined by unique makers and key factor genes. We are the first to clearly depict the transcriptional atlas and defined each subgroups of adult peripheral blood progenitor. We recovered the key transcript factors in lineages fate decision, and restructured the lineage commitment model of HPCs that can reconcile previous models and explain the conflict among functional studies. Several fate decision key factors of myeloid and lymphoid progenitors were identified as major switches that controlled hematopoietic lineage formation. Moreover, we are also the first to identify the cell fate decision transcriptional factors in each branches among hematopoietic hierarchy formation. The reaffirmed lineage commitment also revealed hematopoiesis was a continuum procedure controlled by the activation and suppression of transcription factors, with identified boundary. With these information, an elaborate and comprehensive transcriptional atlas of all types of hematopoietic progenitors was then constructed. Our results will enable the whole map of the transcriptional signal and pathway among lineage commitment. Importantly, this newly created atlas can enable accurate definition, hierarchy, and biological insights of hematopoietic progenitors for facilitating new research development in cell engineering, regenerative medicine, and

disease treatment, which we called Artificial Operation of HPCs. Moreover, this open a wide variety of possibilities for artificial precise control progenitor cells differentiation and targeted treatment of immune disease, providing hope for untreatable diseases.

Graphical abstract

The reaffirmed lineage commitment model of HPCs

Highlights

- Classic and traditional surface markers are often speciosity, inefficient and inaccurate for defining hematopoietic progenitor cells.
- Multiple hematopoietic lineages progenitor cells of adult peripheral blood were clearly depicted and redefined by unique makers and key transcript factors.
- Key fate decision transcript factors and genes in each branches among hematopoietic hierarchy formation were recovered by transcriptional atlas.
- Several key factors, including NFE2, LYL1, MYB, CDK4, ETV6, TESPA1, and GATA2, and RUNX1 of myeloid progenitors and RUNX2, HOXA9, JUN/FOS, LST1, TCF4, DDIT4, HOPX, KLF10, HOXA3, TSC22D1, and BCL11A of lymphoid progenitors, were identified as major regulators that controlled hematopoietic lineage formation.
- Hematopoiesis is a continuum procedure controlled by the activation and suppression of transcription factors, with relative identified boundary.
- The lineage commitment model of HPCs was reaffirmed that can reconcile previous models and explain the conflict among functional studies.
- Transcriptional atlas of HPCs help to provide a well explanation of the immune reconstitution in adult and on its cytokines secretion and regulation.
- Application of this atlas can enable artificial precise control progenitor cell

77   differentiation, in vitro accurate culture of each lineages cells, and targeted

78   treatment of immune cell disease to be made possible for future development.

Introduction

Deciphering the hematopoiesis mechanism that underlies progenitor cells definition and fate decision are critical to understanding fundamental questions of hematopoietic lineage commitment. Toward the diversities of peripheral blood cells in the human body, it is like the multifarious stars existing in our galaxy. Although we know their existence, the exact characterization of these cell subsets remains a challenge.

Among the various cells types in peripheral blood (PB), hematopoietic stem cells (HSCs) are one of such rare cell subsets. These cells possess self-renewal capacity and can differentiate to all blood lineage cells. Such that, HSCs reside at the top of a hierarchy and differentiate to all blood lineage cells with tree-like pattern in classical hematopoietic hierarchical models [1]. Single-cell sequencing technique has been successfully implemented to explore molecular heterogeneity and hierarchy of HSCs[2-4]. The lineage commitment is affected by transcription factors (TFs) with a continuum fate decision procedure[5, 6]. Though *in vitro* and *in vivo* studies have reaffirmed their critical role of transcription factors in lineage commitment, the function and regulatory mechanism of HSCs is still unclear [7]. In addition, the transcriptional atlas and the differentiation among subtypes of progenitors in adult peripheral blood remains to be elucidated.

The reason for the molecular mechanisms that controlled lineage commitment and progenitor behaviors remaining an enigma is due to various reasons. Firstly, the definition of hematopoietic progenitor cells (HPCs) is not standardized. Based on the classical and traditional definition, the characterization of HPCs is based on the presence of surface markers, such as CD34, CD38, FLT3 and KIT[5]. Similarly, the combination of surface markers has been proposed for the identification toward common lymphoid progenitors (CLP) and common myeloid progenitors (CMP). These markers include CD10 (MME) and CD38. Yet, owing to the rare and transient nature, the subtypes of CLP and CMP from HSCs have never been stringently identified and accurately captured. Secondly, the functional studies were also used surface markers for the characterization of HPCs. These include the selection of CD34

positive cell from bone marrow or cord blood. However, this approach can be affected

by the type of culture methods and surface markers being used for the identification.

Simply, the use of classical and traditional markers may be not insufficient, such as

KIT, to differentiate mature subset cells. Thus, the use of combined surface markers

for HPCs may cause biased and arbitrary.

In recent years, many studies have identified the lineage by using single cell

sequencing. However, their is still several limitations of the transcriptional atlas being

defined. The scRNA expression data of CD34 positive cells showed two well-defined

subtypes of progenitor CLP and CMP being scattered and mixed distribution in

T-SNE plots. This result implies that the transcriptional atlas for these two types of

cells are not distinguishable by unsupervised dimensionality reduction and clustering.

When compared with others mature cells, HPCs should exist as a relatively

homogeneous progenitor population. The idea classification and defined subset of

lineage cells should be separately distinguishable by unsupervised dimensionality

reduction which carried the PCA key markers of genes as factors.

In addition, the hematopoietic hierarchical models are also inconsistent among

the literature studies[1, 8, 9]. Most of the established lineage fate decision models

were based on *in-vitro* / *in-vivo* functional studies or cell fate induces experiments[10,

11]. The use of cellular conditions can often lead to inaccurate interpretations of HPCs

cell fate decision because cell lineages were induced by diverse experimental

conditions and cannot be permissive to all possible branches of lineage commitment

[11]. Similarly, single cell studies would have modified some aspects of classical

hematopoiesis hierarchical model, as the hematopoietic progenitors are changing

continuously among the progenitors differentiation process[5, 12]. In addition,

progenitors cells were defined by classical markers and selected by surface markers[5,

6]. As each type of progenitors is very rare in whole blood, the captured cell numbers

are often very low with less than one hundred cells[5]. This implies that some key

progenitor subgroups may be missed. Most of the bone marrow HPCs was

quiescent status progenitors. sc-RNA Seq of bone marrow progenitors can't clearly

depict the whole lineage commitment of hematopoietic progenitors. Thus, this would

affect the reliability of results. As the classic definition of hematopoietic progenitors

remains inefficient and inaccurate, this can cause the disorganization among the

subgroups and without a reliable identification of the boundary among them[2].

Moreover, unsupervised dimensionality reduction and clustering of lineage cells

demonstrated farraginous model[4]. The measurement of lineage definition, fate

decision, and establishing the reliable lineage commitment between them are still a

challenge[8, 13].

Herein, we redefined the adult hematopoietic progenitors, and recovered the key

transcript factors in lineages fate decision, and reaffirmed the lineage commitment

model of HPCs by single-cell sequencing. Our HPCs atlas defined the cellular biology

of differentia stage progenitors among hematopoietic lineages generation. Our study

can help to open new perspectives for the characterization, definition, and

classification of HPCs.

Results

Single-cell identified two subtypes of progenitor cells in adults peripheral blood.

In order to enrich most of the subset of HPCs and later progenitors, such as CLP

and CMP, lymphocyte and granulocyte were removed by surface marker panel

(Lineage−) and progenitor cells were enriched by negative deletion (Fig. 1A). 10X

genomics scRNA was conducted and totally 79,655 cells were captured after quality

filters of mitochondrial RNA, low gene expression cells and erythrocyte cell.

Unsupervised dimensionality reduction and clustering analysis showed 17 clusters by

Seurat and UMAP (Fig. 1B). Total 3180 CD34 positive cells were identified in the

dataset. Besides, samples from all donors were highly reproducible for cell

composition. The HPCs marker of CD34 mainly expressed in two nearby clusters 6

and 14 with the co-expression of FLT3 and KIT (Fig. 1C).

To confirm whether the two clusters are subset of progenitor cells, the different

expression of lymphocyte and granulocyte surface markers and known HSC related

genes were compared (Fig. 1C). Both of the two clusters were found to express KIT

and FLT3, but do not contained the expression of lymphocyte and granulocyte surface

markers. Based on this, the two clusters can then be well distinguishable by three

markers, defined with CD34+CD38+MME- and CD34+CD38-MME+ (Fig. 1C).

However, a portion of the cells can positively expressed CD38 or MME in the two

subgroups, respectively. KIT (CD117) and FLT3 were mainly expressed in one of the

CD34+ clusters, with low positive percentages but also high expressed in the other

clusters. This indicates that the progenitors of CD34+KIT and CD34+FLT3 belonging

to two different clusters which may control the two differentiation directions.

We further analyzed the significantly identified genes among the two clusters

with the other clusters. As found, we presented 16 progenitors related genes which

expressed in more than 80% cells in the two clusters (Fig. 1C). Several HSCs key

regulate genes were identified, such as high expression of *CDK6, SERPINB1, SPINT2,*

IMPDH2, TXN, SOX4, MYADM, HMGA1, CDK2AP1, CD82, TSC22D1, LYL1, and

middle expression of SPI1[6], IFITM3, HMGB1, and PPIA. Based on the surface

markers and the above hematopoietic stem cells functional key gens, we can confirm

that the two identified clusters are hematopoietic progenitor cells. We also identified

three new progenitor key genes of *PPIB, HMGA1,*and *EGFL7*. It is very obviously

that the gene of EGFL7 are very high and specific expressed in the two progenitors

clusters, so that EGFL7 is a key functional gene in hematopoietic progenitor cells

lineage differentiation and mature (Fig. 1D). The single cell sequencing identified two

subgroups which are progenitor cells with category and functional heterogeneity.

The differentiation of transcript markers among progenitors

Our single cell expression atlas level showed the two clusters of myeloid

progenitors (CD38+) and lymphoid progenitors (MME+) presented heterogeneity in

these progenitors, suggesting that functional of the two-cluster subset are diversity. To

determine the diversity, re-clustering of analysis was conducted and 11 clusters was

generated (Fig 2A). Based on our finding, the subgroup of cells aggregated in two

main directions. One is the cluster 0 called lymphoid progenitors, which showed the

up-regulated expression of MME. The other is myeloid progenitors comprised with

cluster 1, cluster 2, and cluster 5, with the gradual up-regulation of CD38 and KIT (Fig 2B). The three genes expression demonstrated contrary among the direction of myeloid progenitors and lymphoid progenitors. But the expression is not whole positive of the two direction cell types. This implies that CD38 and MME can be used as the markers distinguished myeloid progenitors and lymphoid progenitor differentiation directions, but not suitable to use in subtypes classification.

The efficient and accurately of classical hematopoietic progenitors markers which be used to distinguished CMP, GMP (Granulocyte-monocyte progenitors) and CLPs, were also re-analyzed. Our data demonstrated the expression of THY1 (CD90) is very rare (only 17 cells) in the two progenitor clusters which may cause as the progenitors were isolated from adult PBMC. To confirm whether the traditional classification markers THY1 and CD49f (ITGA6) expressed in progenitor, we analyzed four bone marrow single cell projects data of GSE120221, GSE117498, GSE145531 and GSE181989 (Fig. S1). Total 8296 CD34 positive cells were extracted and merged of the four projects data. The results showed positive expression cell numbers of THY1 is only 84 and CD49f is 824, and the distribution is inconsistent. This is similar with the other identified markers of FLT3, CD38 and KIT (Fig. S1). The different progenitor types of cells (CMP or CLP) were not distinguishable by unsupervised dimensionality reduction and clustering in bone marrow CD34+ cells. In addition, our data showed that the use of CD49f, THY1, FLT3 and KIT to classify or define the progenitor cells groups were very artificial and subjective. With the progenitors of CD34+FLT3+ cells defined a mixed population of progenitors, this finding may help to provide a well explanation that previously functional study of selected progenitors (such as CD34+FLT3+) presented multiple differential potential.

The essential regulate genes of HPCs lineages differentiation

To confirm the essential genes in HPCs status maintain and differentiation. We compared the expressions among three single cells data sheets, including bone marrow, PBMC and our CD34- cells subsets. The genes of SERPINB1, CDK6, HMGA1, IMPDH2, SPINT2, TXN, and EGFL7 were highly expressed in bone

marrow and in our identified progenitor cells when compared with mature PBMC

cells (Fig S2A). Those genes also high expressed in most of the subgroups of our

CD34+ cells. This indicates those genes are essential to regulating genes in HPCs

status for maintenance and differentiation.

Classifications of the lineages differentiation

The transcriptomes among membrane marker-defined and transcriptomes-defined

cells were thoroughly compared. Traditionally, membrane markers of progenitor CMP

were defined with CD38+MME-FLT3+, while megakaryocytic–erythroid progenitors

(MEP) were defined with CD38+MME-FLT3-. However, single cell clusters based on

these markers were not distinguishable among the subgroups. As the expressions of

CD38 and FLT3 were contradictory, this demonstrated FLT3 positive cells were

negative of CD38 expression in most clusters (Fig 2B). With the CD38-FLT3+

subgroups progenitors were more original than the FLT3 negative cells, they are more

likely to be CMPs. The expression is changed gradually and presented a continuum

procedure. Although the expression tendency of FLT3+ was consistent with

previously definition, it remained inefficient and inaccurate to classify the

differentiation stages among the CMPs lineages. On the other hand, our result

demonstrated clearly that CMPs lineages clusters can be distinguishable with the

expression of KLF1, GATA1 and CSF3R (Fig 2B).

Our results showed KLF1 and GATA1 were uniquely expressed in the three

clusters of 3, 8 and 9. This indicated that three clusters are likely to be either

megakaryocyte progenitors (MkPs) or erythrocyte progenitors (ErP) of MEPs

progenitors. The expression of KLF1 and GATA1 only presented in the late stage of

CMPs directions, which indicated those two genes are the keys regulated genes in the

differentiation from CMPs to MkPs or ErP. The other gene of GATA2 was widely

expressed in most of the others clusters but mainly highly expressed in CMPs clusters.

Interestingly, the expression of CSF3R disappeared in the MEPs or MkPs clusters

(Fig 2B). During the lineage differentiation of CMPs, GATA2 play a very important

role in all types of myeloid cell lines. The procedure of MEPs or MkPs differentiation

was managed by up-regulated KLF1 and GATA1, and down-regulated with CSF3R.

Hence, the cluster 5 was MEPs would act as the key progenitor to control the

differentiation to MkPs or ErP progenitors.

As AVP was known to be uniquely expressed in HSC, our data showed both of

AVP and CSF3R were mainly expressed in cluster 1 and 2 (Fig 2C and 2D). The high

expression of CSF3R was consistent with G-CSF mobilization in peripheral blood

HSC transplantation. Those clusters were the original multipotent clusters named as

HPCs, which can prime to lineages of CLPs and CMPs. MPO was uniquely expressed

in neutrophils in the border of cluster 1 (Fig S2B and S2C). This indicated that the

granulocyte-monocyte progenitor branches were also present in the CMPs clusters.

Our results showed the expression of IGHM, MME, and HOXA9[14] was presented

in cluster 0, which confirmed the cluster was lymphoid progenitor cells (Fig 2C). The

expression level is increasing gradually of the transcription factors HOXA9 from

cluster 1 to cluster 0 (Fig 2C).

Based on traditional surface markers, our data showed the classification of the

progenitor subgroups or stages among lineage differentiation were inefficient and

inaccurate. Importantly, this demonstrated that the expression and activation of

transcription factors combined with unique marker genes would be more efficient for

lineage definition of HPCs.

Differentiation and enrichment analysis

Differentiation and enrichment analysis was conducted among the clusters. As the

expression changed gradually and presented in a continuum procedure, simple

analysis on the different genes among clusters, especially when compared with nearby

clusters, may not be efficient. To overcome this, we conducted the differentiation

analysis among cluster 0 (CLPs) and cluster 3 (MEPs). Both of the two clusters are

original clusters of CLPs or MEPs. The biology procedure analysis of Gene Ontology

showed the high genes expression of CLPs were enriched in the regulation of

leukocyte and T cell activation (Fig. 2E). While for CMPs, the genes expressions were

highly enriched in mononuclear cell differentiation, erythrocyte differentiation and

homeostasis (Fig. 2F). This was consistent with our previous results on CLPs and

CMPs progenitors differentiation to two directions with strictly lineage affiliated by

unsupervised dimensionality reduction and clustering.

Dynamics of progenitors cells subpopulations

It is clearly visible that CD38 and MME subgroups predominantly occupied the

two directions of the progenitor differentiation, respectively. To confirm the trajectory

change of the cells, trajectory analysis was conducted and three directions of

progenitor cells differentiation were presented. Along the trajectory of right direction

(State 1) which was specific expressed MME, only the cluster 0 and 4 were

distributed (Fig. 3A and 3B). In contrast, the left direction (State 3) was composed by

the CD38 positive expression clusters, such as cluster 1, 2, 3 and 5 (Fig. 3A and 3B)

(Fig. S3). The expression tendency of MME and FLT3 was consistent with CLP

differentiation, with contrary of KIT and CD38 which expressed with CMP lineages.

Heatmap analysis showed the expression tendency of selected marker genes in

CMPs and CLPs. The CLPs branch highly expressed lymphoid marker genes of MME,

IGHM and FLT3 (Fig. 3C). The CMPs branch was highly expressed with myeloid

marker genes of GATA1, KLF1 and CD38. The expression of MME and CD38 were

clearly inverse during the CLP and CMP lineages priming procedures (Fig. 3D). The

last branch was highly expressed with NKG7, CD68 and FCGR3A (CD16), which are

the markers of NK cells, monocyte and macrophages progenitor cells. This indicated

lineage priming of these three types of cells may be distinct with CLPs and CMPs. In

addition, the expression of transcription factors was distinguished among CLPs and

CMPs directions. The expression of KLF1 and GATA1 only appeared in the later

stage of myeloid progenitors named MEPs. This indicated that the HPCs lineages

priming is likely controlled by key transcription factors. This may also indicated that

the unperturbed adult hematopoiesis depends on replenishment from multiple

potential progenitor.

Transcription factors promote the progenitor lineage priming with key receptors

To identify and characterize the committed cellular status, the critical co-expression of transcription factors for lineage fates were re-analyzed. Our study showed the CMP multipotent progenitors, which co-express with transcription factors GATA2, can differentiate into the lineage of myeloid and megakaryocyte cells (Fig. 2B). The expression of KLF1 was only present in the later stage of MEPs status. Furthermore, the expression of CSF3R was mainly showed in the original progenitors and almost disappeared in the later stage of MEPs progenitor. The cytokine receptors will co-express or co-silence with the relevant lineage key regulators, which indicated the progenitor lineages priming was multiple genes regulate procedure.

Likewise, the transcription factors regulation by SCENIC analysis showed the expression of transcription factors was completely distributed differently among CLP and CMP progenitors priming directions (Fig. 4A). The activation and expression of key regulators was much different among the priming stage of progenitors, such as GATA1 (Fig. 4B). As shown in our analysis, the transcription factors of BCL11A, YY1 and REL were the key regulators in CLP priming. And the KLF1, GATA1, and GATA2 were found to play a critical role during the lineage procedure of CMPs (Fig. 4C, Fig. S4).

To confirm which cluster was a megakaryocyte progenitor, the analysis on the expression of marker genes was performed. When compared with the key different genes and transcription factors among cluster 8 and cluster 9, the expression of KLF1 and GATA1 were gradually up-regulated in cluster 9 (Fig. 4A). The expression of KLF1 and GATA1 were up-regulated when the progenitor cells differentiate to megakaryocyte and erythrocyte lineages (Fig. S5). Our results also showed the expression of KIT, MS4A2 and TPSB2 were up-regulated or unique expression in cluster 8, indicating that the cluster was consisted of mast cell and basophil progenitors (Fig. S5).

G-CSF mobilization various type of progenitors in peripheral blood

Various types of progenitors were shown to exist in the PBMC of adults in the

quiescent status. To examine whether any different among the mobilization and quiescent (unperturbed) progenitors, single cell sequencing was conducted in the G-CSF mobilization peripheral blood progenitors. A total of 10450 CD34+ cells were enriched and sequenced, among which 16 clusters subgroups were showed after reclustering (Fig. 5A). Our results indicated that the total number of all types of progenitors were remarkable increased. All kinds of subtyped progenitors were shown to be existed in the quiescent status when compared before and after mobilization (Fig. 5B). This signifies that all types of progenitors were existed in the quiescent status of peripheral blood.

Since the progenitor was shown to gradually differentiate to mature stages in the peripheral blood, it would be expected fold changes of the more primitive progenitors to be higher than mature progenitors. Based on our analysis, the highest cell numbers and folder changed among the clusters were found to be in cluster 0 (Fig. 5C). The distribution and characterization of the featured markers were consistent with previously CD34 subgroups (Fig. 5D). Additionally, the AVP was expressed in cluster 0 which was consistent with the analyzed results in the quiescent status of the progenitor cells. Hence, cluster 0 was the most original progenitor in the peripheral blood that can be identified as HPCs. In contrast, the expression of THY1 (CD49f) were only identified in very few cells. Gene expression analysis also demonstrated four genes, including *AVP*, *SOX4*, *C1QTNF4*, and *JUN*, were high expressed in cluster 0 (Fig. 5F).

The progenitors were clustering into several directions which also consistent with the cluster distributions of quiescent status (Fig. 5A and Fig. 5D). One new direction of cluster 3 was presented in the all cells merged populations (Fig. 5E). The markers genes of that cluster were MPO and CSF3R. As myeloperoxidase (MPO) is the most abundant protein in human neutrophils and plays an important role in neutrophil antimicrobial responses[15, 16]. The other marker gene PRSS57 of neutrophils was also presented and up-regulated expression in the cluster[17]. So the cluster 3 was the progenitor of neutrophils, this was shown to be in a completely different direction with megakaryocyte or erythrocyte progenitors. Based on this analysis, it further

showed NK cells progenitors and monocyte progenitor cells were not clustering

nearby the other lineages (Fig. 5A).

In summary, there are three clusters which nearby the original progenitor of HPCs.

It is clearly shown that during the HPCs priming and differentiation procedures, the

HPCs priming into three progenitor cell types coinstantaneous and later maturing to

diversity hematopoietic lineages. The lymphoid progenitors, neutrophils progenitors,

and megakaryocytic–erythroid progenitors were presented at the same time point and

stage during the HPCs first step of priming process.

Re-definition of hematopoietic progenitor cells (HPCs)

As previously shown, cluster 0 was the most original progenitor in the peripheral

blood that can be identified and defined as HPCs. All of the later mature progenitors

are found to be developed from those original cells. Based on this rationale, we

re-defined the three directions with common lymphoid progenitors (CLPs), GAPs

(GATA genes controlled progenitors), and neutrophilic and monocyte progenitors

(NMPs), which cannot be characterized by CD38, FLT3, and KIT (Fig. 6A). The

differentiation of GAPs lineages was independently with the other two branches. The

shape of each branch was controlled by the expression of key regulated genes which

was shown to function like a switch. Through the transcription factors expression

switches on-or-off and further gradually accumulated, generally synergistic with key

receptor genes expression and signals activation, the original progenitor cells will be

primed and differentiated to various progenitors.

During the myeloid progenitor differentiation procedure, the expression of

GATA2 and CSF3R genes is the first branch being formed and presented (Fig. 6).

This was shown to be followed by the down-regulation of CSF3R genes and

up-regulation of GATA2, which implies the progenitors were priming to oligo-potent

progenitor GAPs. The expression of GATA1 and KLF1 were presented and gradually

up-regulated at later stage. This subgroup was defined as GAPs mature stage named

with megakaryocytic–erythroid progenitors (MEPs). With the differentiation of MEPs

enhancement, the second branch was presented in the mature stage of MEPs. Toward

the final stages of the MEPs, the expression of KIT, GATA1, and KLF1 controlled the differentiation of progenitor to megakaryocyte progenitors, erythrocyte progenitors (Pro-ME), and mast cell or basophil progenitors (MBPs). In contrast to the other branch of myeloid progenitor differentiation process, the expression of GATA2 and KLF1 were restrained during differentiation trajectories when the progenitors were priming to neutrophilic and monocyte progenitors (NMPs, Fig 6A). In the lymphoid progenitor branch priming, critical transcription factor of HOXA9 and HOPX were identified and was shown to gradually up-regulated during the process of CLPs lineage differentiation (Fig. 6B). Similarly, the marker genes IGHM and MME of lymphocyte were also found to be up-regulated. This result provided strong evidence that the whole peripheral blood progenitors include a gradient of heterogeneity hematopoietic progenitors. The progenitor cells fate decisions and lineage commitment were mainly controlled by a combination of critical TFs and key receptor genes (Fig. 6). Also, the hierarchical organization of hematopoiesis was a continuum procedure, with key gene expressions presented and gradually being regulated. After the switch genes were activated and up-regulated, the progenitor cells would then differentiated into the next stages of priming. During this process, the decision of cell fate will be controlled and divided into different lineage branches. The fate deciding boundaries were then identified among the differentiation stages. The expression of GATA2 and CSF3R genes was shown to be the boundary for GAPs and NMPs. This finding is the first to validate that hematopoiesis was controlled by a switch-like model with identified relatively boundary, opening a new perspective on hematopoiesis (Fig. 6C).

Cell Atlas and characterization of all re-defined subtype progenitors cells

To clearly define the subgroups of progenitors hematopoietic, the characterization of all re-defined subtype progenitors cells were re-analyzed. Each of subgroups presents specific highly expression genes or negative express genes. These activated genes may enrich in some signal pathways which play a critical role in the lineages differentiation.

Pan markers of HPCs and unique makers of each subgroups

Our study showed the classical markers of CD49f and THY1 were not found in the multipotent HPCs progenitors. In contrast, ITGA4 (CD49d) was highly expressed in most of the progenitors, which is key molecular regulation of HSC homing[18]. Some reported HSC functional related marker genes was broadly expressed in most of the progenitor cells, such as PTPRC, HMGB1, CD74, PFN1, TXN, ZFP36L2, SERPINB1, and HSP90AA1 (Table 1 and Fig. 1B)[19-25]. These genes were shown to mediate the self-renewal, differentiation, mobilization, and maintenance of hematopoietic progenitor cells. In addition, two new genes of PPIA and EIF1 were identified. These genes were similarly to previous reported genes that pan-expressed in progenitors and highly expressed in matured subgroups of peripheral blood cells. A few progenitors' specific genes of CDK6, SOX4, CD82, EGFL7, IMPDH2, SPINK2, TSC22D1, FOXP1 were also identified in most progenitors[26-32]. Among these specific genes, EGFL7 was identified as being pan-expressed in HPCs, CLPs and MEPs. As most of these genes were confirmed by functional validation, our data demonstrated that those high expression genes CD34 progenitor subsets were essential genes in progenitor status and differentiation.

Several key factors were identified, including NFE2, LYL1, MYB, RUNX1, CDK4, ETV6, TESPA1, and GATA2 of myeloid progenitors and RUNX2, HOXA9, JUN/FOS, LST1, TCF4, DDIT4, HOPX, KLF10, HOXA3, TSC22D1, and BCL11A of lymphoid progenitors (Fig. 6B). Interestingly, the two factors of RUNX2 and RUNX1 showed completely reverse expression tendency. Our data revealed RUNX2 monitored the differentiation of NMPs and GAPs, while RUNX1 controlled the differentiation of CLPs. Importantly, these key factors were shown to play reversely functional role during the differentiation of the HPCs. Through the up-regulated combination of NFE2, LYL1, MYB, RUNX1 , CDK4, ETV6, TESPA1, GATA2 and in restraining the expression of RUNX2, HOXA9, JUN/FOS, LST1, TCF4, DDIT4, HOPX, KLF10, HOXA3, TSC22D1, and BCL11A, the HPCs will diverse to differentiate to myeloid progenitors. Together with the activation of other receptors

and unique transcript factors in each lineages progenitors, such as CSF3R for

neutrophilic and GATA2 for GAPs, the whole hematopoietic lineage commitment was

precisely controlled and gradually formed (Table 1, Fig. 6C). However, some

unidentified regulators may still play a role during the process of hematopoietic

lineage commitment. Thus, larger progenitor cell numbers and effective enrichment

methods would be helpful to uncover those factors.

Our result also showed the selected specific marker genes and key transcript

factors of each lineage progenitors (Table 1). Additionally, the negatively expressed

genes were listed to distinguish each subgroup and clearly define the lineages

progenitors (Table 1). It was identified that SLC40A1 was a unique new marker and

key regulator of the whole lineages during GAPs differentiation. Our studies showed

TAL1 only expressed in MEPs progenitors, which is the later stages of GAPs. LYL1

was up-regulated in the lineages of GAPs but decreased in the CLPs lineages. Our

studies further confirmed that those two genes were critical regulators during myeloid

lineages differentiation.

Pathways of each subgroups

Diversity pathways affect and control the differentiation of HPCs. Our current

analyses results showed the JUN and JUND were enriched and highly expressed in

most of the progenitors (Fig. 5 and 5F). This indicated that the JUN pathway may

play a critical role in differentiation, mobilization, and maintenance of hematopoietic

progenitor cells. The DEPTOR gene was found to be highly expressed in the clusters

of MEPs. As DEPTOR was a negative regulator of the mTORC1 and mTORC2

signaling pathways[33]. That meaning the mTOR pathway should be suppressed in

the MEPs mature and differentiation. The expression of NFKBIZ was very high

expressed in the progenitors of monocyte and macrophages[34], that meaning the

NFKB pathway was suppressed in the monocyte and macrophages mature and

differentiation. The other key regulators and pathways of each subgroup will be

shown in future.

Functional validation and primary culture

Previously studies already conducted many functional validations of HSCs key regulated genes. We hypothesis that our HPCs definitions and hematopoiesis model were right, that meanings the functional studies phenotype results would be consistent or can be explanation by our reaffirmed model. And furthermore, those conflicting and confusing results can be clearly interpretation by the reaffirmed hematopoiesis hierarchical organization model.

Our analysis results were found to be consistent with other functional studies results. KLF1 and GATA1 were well-known play a key role in the erythroid and megakaryocytic differentiation[35, 36]. The expression of GATA1 progenitors generated mast cells, eosinophils, megakaryocytes and erythroid cells, and lacking expression can drive generated monocytes, neutrophils and lymphocytes[35]. Our results revealed that GATA1 was unique expressed in the later stages of MEPs and negatively expressed in the monocytes, neutrophils and lymphocytes progenitors (Fig. 5D). The result implies that the key transcript factor of GATA1 was very critical for the MEPs progenitor lineages differentiation. Other evidence is transcript factor NFE2. It was reported that NFE2 acts as a critical regulator in hematopoietic cells. The knockout of NFE2 in mouse model displayed mild erythroid abnormalities and major defect in megakaryocyte biogenesis[37, 38]. Likewise, our results revealed NFE2 was up-regulated in the NMPs and GAPs lineage directions (Fig. 6B).

Based on findings from primary culture, M-CSF was shown to be critical for the survival of macrophage, which is consistent with our results that the receptor CSF1R of M-CSF only expression in monocyte progenitors and matured macrophages (data not shown). In addition, SCGF, also known as CLEC11A, is a secreted glycoprotein that exhibits erythroid burst (BFU-E) promoting activity in the absence of SCF/c-kit Ligand[39, 40]. Our results showed stem cell growth factor (SCGF) was highly expression in NMP and GAP progenitors, especially in the whole lineages of MEPs that can explain SCGF burst promote erythroid activity. Furthermore, SCGF cooperates with GM-CSF, or G-CSF to enhance the growth of granulocyte-macrophage colonies (CFU-GM)[39, 40]. Because the CSF3R receptor of

G-CSF was highly expression in progenitors of myeloid, SCGF cooperates with G-CSF would activate and induced the progenitor differentiation to neutrophils and macrophages but reduce MEPs lineages generation.

Discussion

Taken together, we demonstrated the transcription atlas data of original status HPCs in adult peripheral blood with large number of HPCs. Our study further reaffirmed and established a comprehensive model of hematopoiesis that can reconcile previous models and explain the conflict among functional studies. This helps to provide a detailed insight into lineage commitment from HPCs into almost branches of human peripheral blood progenitors.

Currently, there is without a standard consensus for the definition of all HPCs lineages. This is owing to the inconsistent classification with conflict functional outcomes among the current literatures[9, 13]. Our study succeed enriched the large numbers and almost whole subgroups of HPCs. This will be particularly attractive for large single cells studies to establish the comprehensive model of hematopoiesis with clearly HPCs definition and reliable lineage commitment. All of the hematopoietic progenitors were reaffirmed by unique makers and key factor genes, which can be well distinguishable. We are the first clearly definition all lineages of HPCs by unsupervised and inartificial clustering, which provide the unambiguous transcription atlas of each type of HPCs lineages. The unique markers genes of progenitor subgroups were highly consistent with previously characterization descriptions, such as KLF1 of MEPs, HPGDS of mast cell, and MPO of neutrophilic[15, 41, 42].

It is widely known that the HPCs would differentiate into common myeloid progenitors (CMPs) and common lymphoid progenitors (CLPs). Our data showed none of the MME positive cells co-expressed CD38, suggesting that classification of CLPs using solely CD38+ and MME+ was not sufficient. CMPs are defined with

CD34+CD38+MME- which is consistent with classic definition, but our single cell

expression atlas showed the CD34+CD38-MME+ progenitor cells is inconsistent with

the traditional define of CLPs[4, 5]. Based on our study, we identified two subgroups

of progenitors by single cell sequencing which are not consistent with classical

definition, and showed specific classify surface or characterization markers genes,

such as GATA1, MPO and KLF1. Likewise, we provided a clearly evidence that the

classical surface markers of CD38, KIT, THY1, CD49f and FLT3 were non-specific

and inefficient to identify HSCs. This remind us the traditional lineage definition of

HSC by above markers was imperfect and need to be improved. Evidently, the

combination of transcriptional factors with unique marker genes would be more

efficient for lineage definition of HPCs.

Our classification and definition of HPCs also help to explain the conflict and

mechanism of *in vitro* and *in vivo* progenitor cell studies. It is well known that KLF1

and GATA1 are the keys regulate genes of erythrocyte or megakaryocyte

differentiation[43, 44]. Functional study have been shown that the upregulated

transcriptional factor of GATA-1 can affect mast cells, eosinophils, megakaryocytes

and erythroid cells generation, and downregulation to affect monocytes, neutrophils

and lymphocytes generation[35]. Our finding reaffirmed the definition of GAPs

lineages to show GATA1 to be solely expressed in the MEPs and up-regulated during

maturation. Previously studies reported TAL1 and LYL1 are essential for maintenance

of adult HSC function [45-47]. Since the induced pluripotent stem cells (iPSCs) was

established through activation of four transcript factors, reprogram of pluripotent stem

cells was widely investigated in various somatic cells[48]. GATA1 and TAL1 have

been validated in the reprogram of megakaryocytes and erythroid cells[43, 49].

Although they finally induced mature erythroid cells based on those key transcript

factors, but there are still many functional capacity limited during differentiation.

The CSF3R was mainly expressed in the original clusters (HPCs), which was the

receptor of G-CSF [50]. The underlying molecular mechanisms of G-CSF induces

enhancement of hematopoietic remain unclear. Our results provide a new model of

hematopoietic through scRNA of unperturbed and G-CSF mobilized progenitors

lineages in peripheral blood. All types of the progenitors were remarkable increased,

especially the HPCs. G-CSF directly activates the downstream activation signals of

HPCs through bind with CSF3R, which converted the quiescent status HPCs to

activation progenitors. After the mobilization procedures, HPCs was priming and

differentiation to all kinds of progenitors which can differentiate to mature myeloid

and lymphocyte cells. As key transcriptional factors and regulators controlled the

whole hematopoietic lineage commitment formation, the use of primary cultures also

showed factors of G-CSF and SCGF to induce the lineages formation. Both of these

mechanisms would cooperate to play key roles during the lineage commitment

formation. That reminds us two questions: First, how did the key transcript factors

was activated during the lineage commitment procedure? Secondly, the receptors

activated the signaling cascade and transcript factors, or the transcript factors were

activated independently and factors enhance the commitment formation?

In brief, the unique markers were highly consistent with characterization of most

progenitor subgroups. Secondly, the definition can very well explanation the conflict

of functional studies. Thirdly, the expression of distinguished receptor of cytokines

can perfectly explain the primary culture phenomenon of progenitor subgroups, such

as CSF3R for neutrophilic and CSF1R for monocyte culture[51]. These three aspects

strongly support the definition and classification of HPCs to be distinctly verifiable,

reliable and reasonable.

Moreover, our transcriptional atlas revealed the major role of each progenitor

subgroups in hematopoiesis and the fate decision of downstream various

hematopoietic cells. Through the expression changes of the three key transcription

factors of KIT, KLF1 and GATA1, the later stage of MEPs can differentiate into two

directions. The expression of transcriptional factors, both in original and mature stage

of progenitors, revealed a basic split between myeloid-lymphocyte development and

megakaryocyte-erythroid development. This provides a strong evidence that the transcription factors were the key regulators in each stages of the progenitor's differentiation. Generally, progenitor cells can be priming by key transcription factor activation, which derived progenitor differentiated to downstream specific mature lineages[52]. But previously study showed that transcriptional programs were not strictly lineage affiliated, and expression changes between HSCs and early progenitors occur independently of lineage choice[53]. That may cause by not reliable classification and definition of the subgroups and different stage of progenitor among lineage commitment, which based on the traditional cell surface markers in most of those investigations. We demonstrated the unique markers or transcriptional factors which control the progenitors differentiate to all kind of blood lineages in a continuum differentiation procedure. Our results strongly supported that in an ideally cell fate decision model, the activation of transcription factors and expression of progenitor lineage unique markers demonstrated a consistent lineage affiliated with relatively obvious subgroups boundary.

Our analysis on the transcriptional atlas revealed that the transcriptional factors activation and suppression play a critical role in the differentiation among lineage commitment. The bias during HPCs differentiation was gradually accumulated by activation of multiple lineage-specifying TFs, such as NFE2, LYL1, MYB, and RUNX1 of myeloid and RUNX2, HOXA9, and BCL11A of lymphoid progenitors. These TFs play critical role during the progenitor priming with dynamic entail quantitative expression and abundances change. By the multiple TFs activation and suppression, the downstream signal pathway would be activated. The stepwise controlled the differentiation of each branch during HPCs lineage commitment. During the process, individual progenitors would gradually acquire lineage biases along multiple directions during homeostasis. Our atlas results showed that key transcriptional factors priming HPCs for distinct downstream lineages, which finally generated and formatted the whole lineage commitment. The expression of key TFs could be considered as the relative reliable boundary among stages during progenitor

mature. Through the control of TFs genes, the artificial precise committed progenitor

cell differentiation can be made possible.

It was reported the HSC lineage commitment is a continuous process in recently

single cell sequencing studies[5, 12]. But HSC heterogeneity and unclearly definition

of HSC subgroups caused the mechanism of hierarchy formation still an enigma. The

established hematopoietic lineage models were inconsistent and incompatible. This

was due to the limitation of captured cell numbers and inefficient definition by

identification using surface markers. Based on our clear definition and atlas of HPCs,

it revealed that a gradient of hematopoietic progenitors existed during the lineage

commitment and can be distinguished by the expression of unique TFs. Those

progenitors gradually acquired lineage biases, divergence and prime to each branch.

Our elaborate and comprehensive transcriptional atlas of HPCs can provide accurate

definition, hierarchy, and biological insights of hematopoietic progenitors. As our

study further explained how the diversified progenitor cells were primed, induced,

and differentiated to the whole lineages of immune system cells, this enable the

transcriptional atlas of HPCs to provide new perspective and well explanation about

how the immune was reconstituted in adult. Recently, cytokines storm was reported in

severe infection and coronavirus[54, 55]. Our atlas can clearly demonstrated the

expression cytokines receptor in specific progenitor. By controlling the original stage

and generation of cytokines, especially suppress the initial cytokines activation of the

cascade activation, this could inhibit or reduce the effect of the follow-up cascade

activation. Technically, this would be helpful in controlling the excessive activation

by cytokines and preventing the occurrence of cytokines storm as a new therapeutic

strategy. Our reaffirmed taxonomy will enhance more accurate prevention and

treatment as well as immune monitoring in health and disease.

There were some limitations in our current study. Firstly, our study did not define

the original progenitors with HSCs. Even though the characterization of these

progenitors was most likely originated with stem cells, the markers of CD49f and

THY1 were found to be negatively expressed. Our study showed two type of
progenitor cells that are not clustering nearby the other lineages. One is the NK cells
progenitors and the other is monocyte progenitors. Monocyte progenitors have more
than two types of status progenitors, which located far away from the main cell
populations. We speculated two possibilities to cause this phenomenon. One may be
from the limitation of our enrichment method. This can cause the loss of major status
progenitors from these two types of cells. This will have an effect to cause only the
capture from the later stages. Secondly, the original source of NK cells progenitors
and the monocyte progenitors were found to be independent and distinct with all of
other progenitor cell types. As the whole atlas of our definition was established by
unsupervised dimensionality reduction and clustering from statistical approach, the
accuracy and efficiency should be considered. Artificial intelligence and deep learning
would provide a more accurate, reliable, and distinguishable definition for each status
and all progenitor subgroups during the whole procedure of hematopoiesis.

Importantly, our study helped to re-defined hematopoietic progenitor cells and
reaffirming lineage commitment by transcriptional atlas. We are the first to clearly
depict the transcriptional atlas and defined each subgroups of adult peripheral blood
progenitor. Our results will enable the whole map of the transcriptional signal and
pathway among lineage commitment. Moreover, we are also the first to identify the
cell fate decision transcriptional factors in each branches among hematopoietic
hierarchy formation. The precise control on the hematopoietic progenitor cells
priming and differentiation can then be interfere by the key transcriptional factors and
receptors signals, which we called Artificial Operation of HPCs.

In summary, our transcriptional atlas of HPCs established a comprehensive
continuum and key regulators controlled model of the human adult peripheral blood
hematopoiesis. Toward this transcriptional atlas, it provides accurate definition,
hierarchy, and biological insights of hematopoiesis and lineage commitment of HPCs.
Importantly, this can enable new research development in cell engineering,

regenerative medicine, and disease treatment. Also, this can have critical insight and

implications for the etiology and treatment of hematologic disorders. Moreover, this

open a wide variety of possibilities for artificial precise control progenitor cells

differentiation and targeted treatment of immune disease, providing hope for

untreatable diseases.

Acknowledgments

We acknowledge Professor Chen Qiu, Rongchang Chen, and Yingyun Fu for supporting this research. We also thank Lipeng Chen for bioinformatics and statistical analysis of sequencing data, Qijun Huang and Guoqiang Li for sample collection, and Sujin Chen for draw the schematic diagram.We also thank Chiwai Man of CUHK and Qing Sun for manuscript revsie. Moreover, we are grateful to the subjects for their participation in this study.

Author Contributions

Y.Y conceived the original idea, designed the study, and performed experiments on HPC's enrichment.Y.Y conducted the bioinformatics and statistical analysis of sequencing data.Y.Y wrote the manuscript and revised the manuscript. All authors approved the final draft of the manuscript.

Declaration of Interests

The authors have declared no competing financial interest.

Funding

The correspondence author Dr. Yongjian Yue did not even get one coin supported from any funding or organizations in the study.

Data Availability Statement

The datasets used or analyzed during the current study are available from the corresponding author on reasonable request.

Methods

Sample collection and HPCs enrichment

Totally, 8mL peripheral blood was draw from each donor. Prior to blood collection, mobilizated donors will received subcutaneous recombinant human granulocyte colony-stimulating factor with 300ug in each 12 hours for five days. After five days of

continuous treatment, a total of 10mL of peripheral blood was collected in heparinized

tubes from the stem cell donor, and kept at room temperature. PBMC and total white

blood cells were obtained according to standard protocols. The protocol of peripheral

blood mononuclear cell (PBMC) isolation was based on separation by density

gradient centrifugation with Ficoll® (Cytiva, GE), a highly branched polysaccharide.

Lineages (Lin−, CD3-CD19-CD56-) negative cells were removed by STEMCELL

reagents (15272). Self-design method was applied to enrich the HPCs. Ethical

approval was obtained from the ethics committee of the Shenzhen People's Hospital,

as per the ethical standards recommended in the 1964 Declaration of Helsinki.

Informed consent was obtained from the subjects in the study.

Single cell RNA preparation and sequencing

RNA-sequencing was also performed on human enriched HPCs (Lin−) purified from

10 donors, including one mobilization PBMC progenitor cells by G-CSF. The single

cells suspension was loaded and captured by Chromium microfluidic chips and

barcoded with a 10× Chromium Controller (10X Genomics). Barcoded single cells

RNA was subsequently reverse-transcribed and sequencing libraries constructed with

reagents from a Chromium Next GEM Single Cell 5' reagent kits v2 according to the

manufacturer's instructions (10X Genomics). Sequencing was conducted by Illumina

Nova 6000 (PE150).

Single-cell RNA-seq data processing

The raw sequencing reads of all samples were demultiplexed and aligned to the

reference genome hg38 by Cellranger pipeline. All downstream single-cell

unsupervised dimensionality reduction and clustering was conducted by Seurat[56].

Gene Ontology (GO) enrichment analysis of marker genes was implemented by the

cluster Profiler R package. Trajectory analysis was conducted by Monocle2 [57].

Transcription factors regulation inference and gene regulatory networks analyses

were performed using SCENIC [58].

Integrated analysis with online single cell dataset

The online scRNA data of GSE120221[59], GSE117498[4], GSE145531[60], and GSE181989[61] was downloaded from GSE database. Integrated analysis was performed by Seurat.

References

1. Cheng H, Zheng Z, Cheng T: **New paradigms on hematopoietic stem cell differentiation**. *Protein & cell* 2020, **11**(1):34-44.
2. Xie X, Liu M, Zhang Y, Wang B, Zhu C, Wang C, Li Q, Huo Y, Guo J, Xu C *et al*: **Single-cell transcriptomic landscape of human blood cells**. *Natl Sci Rev* 2021, **8**(3):nwaa180.
3. Ranzoni AM, Tangherloni A, Berest I, Riva SG, Myers B, Strzelecka PM, Xu J, Panada E, Mohorianu I, Zaugg JB *et al*: **Integrative Single-Cell RNA-Seq and ATAC-Seq Analysis of Human Developmental Hematopoiesis**. *Cell stem cell* 2021, **28**(3):472-487 e477.
4. Pellin D, Loperfido M, Baricordi C, Wolock SL, Montepeloso A, Weinberg OK, Biffi A, Klein AM, Biasco L: **A comprehensive single cell transcriptional landscape of human hematopoietic progenitors**. *Nature communications* 2019, **10**(1):2395.
5. Velten L, Haas SF, Raffel S, Blaszkiewicz S, Islam S, Hennig BP, Hirche C, Lutz C, Buss EC, Nowak D *et al*: **Human haematopoietic stem cell lineage commitment is a continuous process**. *Nat Cell Biol* 2017, **19**(4):271-281.
6. Sugimura R, Jha DK, Han A, Soria-Valles C, da Rocha EL, Lu YF, Goettel JA, Serrao E, Rowe RG, Malleshaiah M *et al*: **Haematopoietic stem and progenitor cells from human pluripotent stem cells**. *Nature* 2017, **545**(7655):432-438.
7. Zhu J, Emerson SG: **Hematopoietic cytokines, transcription factors and lineage commitment**. *Oncogene* 2002, **21**(21):3295-3313.
8. Jacobsen SEW, Nerlov C: **Haematopoiesis in the era of advanced single-cell technologies**. *Nat Cell Biol* 2019, **21**(1):2-8.
9. Haas S, Trumpp A, Milsom MD: **Causes and Consequences of Hematopoietic Stem Cell Heterogeneity**. *Cell stem cell* 2018, **22**(5):627-638.
10. Manz MG, Miyamoto T, Akashi K, Weissman IL: **Prospective isolation of human clonogenic common myeloid progenitors**. *Proceedings of the National Academy of Sciences of the United States of America* 2002, **99**(18):11872-11877.
11. Akashi K, Traver D, Miyamoto T, Weissman IL: **A clonogenic common myeloid progenitor that gives rise to all myeloid lineages**. *Nature* 2000, **404**(6774):193-197.
12. Buenrostro JD, Corces MR, Lareau CA, Wu B, Schep AN, Aryee MJ, Majeti R, Chang HY, Greenleaf WJ: **Integrated Single-Cell Analysis Maps the Continuous Regulatory Landscape of Human Hematopoietic Differentiation**. *Cell* 2018, **173**(6):1535-1548 e1516.
13. Laurenti E, Gottgens B: **From haematopoietic stem cells to complex differentiation landscapes**. *Nature* 2018, **553**(7689):418-426.
14. Huang Y, Sitwala K, Bronstein J, Sanders D, Dandekar M, Collins C, Robertson G, MacDonald J, Cezard T, Bilenky M *et al*: **Identification and characterization of Hoxa9 binding sites in**

hematopoietic cells. *Blood* 2012, **119**(2):388-398.

15.   Metzler KD, Fuchs TA, Nauseef WM, Reumaux D, Roesler J, Schulze I, Wahn V,
Papayannopoulos V, Zychlinsky A: **Myeloperoxidase is required for neutrophil extracellular**
trap formation: implications for innate immunity. *Blood* 2011, **117**(3):953-959.

16.   Schultz J, Kaminker K: **Myeloperoxidase of the leucocyte of normal human blood. I. Content**
and localization. *Archives of Biochemistry and Biophysics* 1962, **96**(3):465-467.

17.   Perera NC, Wiesmuller KH, Larsen MT, Schacher B, Eickholz P, Borregaard N, Jenne DE: **NSP4**
is stored in azurophil granules and released by activated neutrophils as active
endoprotease with restricted specificity. *Journal of immunology* 2013, **191**(5):2700-2707.

18.   Li D, Xue W, Li M, Dong M, Wang J, Wang X, Li X, Chen K, Zhang W, Wu S *et al*: **VCAM-1(+)**
macrophages guide the homing of HSPCs to a vascular niche. *Nature* 2018,
564(7734):119-124.

19.   Becker-Herman S, Rozenberg M, Hillel-Karniel C, Gil-Yarom N, Kramer MP, Barak A, Sever L,
David K, Radomir L, Lewinsky H *et al*: **CD74 is a regulator of hematopoietic stem cell**
maintenance. *PLoS biology* 2021, **19**(3):e3001121.

20.   Sundaramoorthy P, Wang Q, Zheng Z, Jiao Y, Chen BJ, Doan PL, Chao NJ, Kang Y: **Thioredoxin**
mitigates radiation-induced hematopoietic stem cell injury in mice. *Stem cell research &*
therapy 2017, **8**(1):263.

21.   Zheng J, Lu Z, Kocabas F, Bottcher RT, Costell M, Kang X, Liu X, Deberardinis RJ, Wang Q, Chen
GQ *et al*: **Profilin 1 is essential for retention and metabolism of mouse hematopoietic stem**
cells in bone marrow. *Blood* 2014, **123**(7):992-1001.

22.   Zhang L, Prak L, Rayon-Estrada V, Thiru P, Flygare J, Lim B, Lodish HF: **ZFP36L2 is required for**
self-renewal of early burst-forming unit erythroid progenitors. *Nature* 2013,
499(7456):92-96.

23.   Benarafa C, LeCuyer TE, Baumann M, Stolley JM, Cremona TP, Remold-O'Donnell E: **SerpinB1**
protects the mature neutrophil reserve in the bone marrow. *Journal of leukocyte biology*
2011, **90**(1):21-29.

24.   Altmann S, Lange S, Pommerencke J, Murua Escobar H, Bullerdiek J, Nolte I, Freund M,
Junghanss C: **High Mobility Group Box 1-Protein expression in canine haematopoietic cells**
and influence on canine peripheral blood mononuclear cell proliferative activity. *Veterinary*
immunology and immunopathology 2008, **126**(3-4):367-372.

25.   Kruta M, Sunshine MJ, Chua BA, Fu Y, Chawla A, Dillingham CH, Hidalgo San Jose L, De Jong B,
Zhou FJ, Signer RAJ: **Hsf1 promotes hematopoietic stem cell fitness and proteostasis in**
response to ex vivo culture stress and aging. *Cell stem cell* 2021, **28**(11):1950-1965 e1956.

26.   Saito-Reis CA, Balise VD, Pascetti EM, Jiminez M, Gillette JM: **Tetraspanin CD82 regulates**
S1PR1-mediated hematopoietic stem and progenitor cell mobilization. *Stem cell reports*
2021, **16**(10):2422-2431.

27.   Karigane D, Kobayashi H, Morikawa T, Ootomo Y, Sakai M, Nagamatsu G, Kubota Y, Goda N,
Matsumoto M, Nishimura EK *et al*: **p38alpha Activates Purine Metabolism to Initiate**
Hematopoietic Stem/Progenitor Cell Cycling in Response to Stress. *Cell stem cell* 2016,
19(2):192-204.

28.   Scheicher R, Hoelbl-Kovacic A, Bellutti F, Tigan AS, Prchal-Murphy M, Heller G,
Schneckenleithner C, Salazar-Roa M, Zochbauer-Muller S, Zuber J *et al*: **CDK6 as a key**
regulator of hematopoietic and leukemic stem cell activation. *Blood* 2015, **125**(1):90-101.

29. Doulatov S, Vo LT, Chou SS, Kim PG, Arora N, Li H, Hadland BK, Bernstein ID, Collins JJ, Zon LI
et al: **Induction of multipotential hematopoietic progenitors from human pluripotent stem**
cells via respecification of lineage-restricted precursors. *Cell stem cell* 2013, **13**(4):459-470.
30. Calvanese V, Capellera-Garcia S, Ma F, Fares I, Liebscher S, Ng ES, Ekstrand S,
Aguade-Gorgorio J, Vavilina A, Lefaudeux D *et al*: **Mapping human haematopoietic stem cells**
from haemogenic endothelium to birth. *Nature* 2022, **604**(7906):534-540.
31. Naudin C, Hattabi A, Michelet F, Miri-Nezhad A, Benyoucef A, Pflumio F, Guillonneau F,
Fichelson S, Vigon I, Dusanter-Fourt I *et al*: **PUMILIO/FOXP1 signaling drives expansion of**
hematopoietic stem/progenitor and leukemia cells. *Blood* 2017, **129**(18):2493-2506.
32. Yu J, Ershler M, Yu L, Wei M, Hackanson B, Yokohama A, Mitsui T, Liu C, Mao H, Liu S *et al*:
TSC-22 contributes to hematopoietic precursor cell proliferation and repopulation and is
epigenetically silenced in large granular lymphocyte leukemia. *Blood* 2009,
113(22):5558-5567.
33. Peterson TR, Laplante M, Thoreen CC, Sancak Y, Kang SA, Kuehl WM, Gray NS, Sabatini DM:
DEPTOR is an mTOR inhibitor frequently overexpressed in multiple myeloma cells and
required for their survival. *Cell* 2009, **137**(5):873-886.
34. Totzke G, Essmann F, Pohlmann S, Lindenblatt C, Janicke RU, Schulze-Osthoff K: **A novel**
member of the IkappaB family, human IkappaB-zeta, inhibits transactivation of p65 and its
DNA binding. *The Journal of biological chemistry* 2006, **281**(18):12645-12654.
35. Drissen R, Buza-Vidas N, Woll P, Thongjuea S, Gambardella A, Giustacchini A, Mancini E, Zriwil
A, Lutteropp M, Grover A *et al*: **Distinct myeloid progenitor-differentiation pathways**
identified through single-cell RNA sequencing. *Nature immunology* 2016, **17**(6):666-676.
36. Siatecka M, Bieker JJ: **The multifunctional role of EKLF/KLF1 during erythropoiesis**. *Blood*
2011, **118**(8):2044-2054.
37. erythroid abnormalitiesGasiorek JJ, Blank V: **Regulation and function of the NFE2**
transcription factor in hematopoietic and non-hematopoietic cells. *Cellular and molecular*
life sciences : CMLS 2015, **72**(12):2323-2335.
38. Lecine P, Villeval JL, Vyas P, Swencki B, Xu Y, Shivdasani RA: **Mice lacking transcription factor**
NF-E2 provide in vivo validation of the proplatelet model of thrombocytopoiesis and show
a platelet production defect that is intrinsic to megakaryocytes. *Blood* 1998,
92(5):1608-1616.
39. Hiraoka A, Yano Ki K, Kagami N, Takeshige K, Mio H, Anazawa H, Sugimoto S: **Stem cell growth**
factor: in situ hybridization analysis on the gene expression, molecular characterization and
in vitro proliferative activity of a recombinant preparation on primitive hematopoietic
progenitor cells. *The hematology journal : the official journal of the European Haematology*
Association 2001, **2**(5):307-315.
40. Hiraoka A, Ohkubo T, Fukuda M: **Production of human hematopoietic survival and growth**
factor by a myeloid leukemia cell line (KPB-M15) and placenta as detected by a monoclonal
antibody. *Cancer research* 1987, **47**(19):5025-5030.
41. Jiang J, Faiz A, Berg M, Carpaij OA, Vermeulen CJ, Brouwer S, Hesse L, Teichmann SA, Ten
Hacken NHT, Timens W *et al*: **Gene signatures from scRNA-seq accurately quantify mast cells**
in biopsies in asthma. *Clinical and experimental allergy : journal of the British Society for*
Allergy and Clinical Immunology 2020, **50**(12):1428-1431.
42. Klimchenko O, Mori M, Distefano A, Langlois T, Larbret F, Lecluse Y, Feraud O, Vainchenker W,

Norol F, Debili N: **A common bipotent progenitor generates the erythroid and**
megakaryocyte lineages in embryonic stem cell-derived primitive hematopoiesis. *Blood*
2009, **114**(8):1506-1517.

43.   Capellera-Garcia S, Pulecio J, Dhulipala K, Siva K, Rayon-Estrada V, Singbrant S, Sommarin
Mikael NE, Walkley Carl R, Soneji S, Karlsson G *et al*: **Defining the Minimal Factors Required**
for Erythropoiesis through Direct Lineage Conversion. *Cell reports* 2016, **15**(11):2550-2562.

44.   Wontakal SN, Guo X, Smith C, MacCarthy T, Bresnick EH, Bergman A, Snyder MP, Weissman
SM, Zheng D, Skoultchi AI: **A core erythroid transcriptional network is repressed by a master**
regulator of myelo-lymphoid differentiation. *Proceedings of the National Academy of*
Sciences 2012, **109**(10):3832-3837.

45.   Bergiers I, Andrews T, Vargel Bolukbasi O, Buness A, Janosz E, Lopez-Anguita N, Ganter K,
Kosim K, Celen C, Itir Percin G *et al*: **Single-cell transcriptomics reveals a new dynamical**
function of transcription factors during embryonic hematopoiesis. *eLife* 2018, **7**.

46.   Souroullas GP, Salmon JM, Sablitzky F, Curtis DJ, Goodell MA: **Adult hematopoietic stem and**
progenitor cells require either Lyl1 or Scl for survival. *Cell stem cell* 2009, **4**(2):180-186.

47.   Chan WY, Follows GA, Lacaud G, Pimanda JE, Landry JR, Kinston S, Knezevic K, Piltz S,
Donaldson IJ, Gambardella L *et al*: **The paralogous hematopoietic regulators Lyl1 and Scl are**
coregulated by Ets and GATA factors, but Lyl1 cannot rescue the early Scl-/- phenotype.
Blood 2007, **109**(5):1908-1916.

48.   Takahashi K, Yamanaka S: **Induction of Pluripotent Stem Cells from Mouse Embryonic and**
Adult Fibroblast Cultures by Defined Factors. *Cell* 2006, **126**(4):663-676.

49.   Moreau T, Evans AL, Vasquez L, Tijssen MR, Yan Y, Trotter MW, Howard D, Colzani M,
Arumugam M, Wu WH *et al*: **Large-scale production of megakaryocytes from human**
pluripotent stem cells by chemically defined forward programming. *Nature communications*
2016, **7**(1).

50.   Beekman R, Touw IP: **G-CSF and its receptor in myeloid malignancy.** *Blood* 2010,
115(25):5131-5136.

51.   Hume DA, MacDonald KP: **Therapeutic applications of macrophage colony-stimulating**
factor-1 (CSF-1) and antagonists of CSF-1 receptor (CSF-1R) signaling. *Blood* 2012,
119(8):1810-1820.

52.   Hoppe PS, Schwarzfischer M, Loeffler D, Kokkaliaris KD, Hilsenbeck O, Moritz N, Endele M,
Filipczyk A, Gambardella A, Ahmed N *et al*: **Early myeloid lineage choice is not initiated by**
random PU.1 to GATA1 protein ratios. *Nature* 2016, **535**(7611):299-302.

53.   Laurenti E, Doulatov S, Zandi S, Plumb I, Chen J, April C, Fan JB, Dick JE: **The transcriptional**
architecture of early human hematopoiesis identifies multilevel control of lymphoid
commitment. *Nature immunology* 2013, **14**(7):756-763.

54.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X *et al*: **Clinical features of**
patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020,
395(10223):497-506.

55.   Fajgenbaum DC, Longo DL, June CH: **Cytokine Storm.** *New England Journal of Medicine* 2020,
383(23):2255-2273.

56.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, Hao Y, Stoeckius M,
Smibert P, Satija R: **Comprehensive Integration of Single-Cell Data.** *Cell* 2019,
177(7):1888-1902 e1821.

57. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen
TS, Rinn JL: **The dynamics and regulators of cell fate decisions are revealed by**
pseudotemporal ordering of single cells. *Nature biotechnology* 2014, **32**(4):381-386.
58. Aibar S, González-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H, Hulselmans G, Rambow F,
Marine J-C, Geurts P, Aerts J *et al*: **SCENIC: single-cell regulatory network inference and**
clustering. *Nature Methods* 2017, **14**(11):1083-1086.
59. Oetjen KA, Lindblad KE, Goswami M, Gui G, Dagur PK, Lai C, Dillon LW, McCoy JP, Hourigan CS:
Human bone marrow assessment by single-cell RNA sequencing, mass cytometry, and flow
cytometry. *JCI insight* 2018, **3**(23).
60. Zhu C, Lian Y, Wang C, Wu P, Li X, Gao Y, Fan S, Ai L, Fang L, Pan H *et al*: **Single-cell**
transcriptomics dissects hematopoietic cell destruction and T-cell engagement in aplastic
anemia. *Blood* 2021, **138**(1):23-33.
61. Tonglin H, Yanna Z, Xiaoling Y, Ruilan G, Liming Y: **Single-Cell RNA-Seq of Bone Marrow Cells**
in Aplastic Anemia. *Front Genet* 2021, **12**:745483.
965
966

967   **List of Table and Legends**

969   **Table 1. The selected specific marker genes of all re-defined subgroups**

Progenitor Cell Types	Markers	Negative
Pan-Progenitors	CD49d, EIF1,PPIA, HMGB1, CD74, PFN1, TXN, ZFP36L2, SERPINB1,HSP90AA1	
HPCs-Progenitors(C0)	CD34, CDK6, SOX4, **CD82**, EGFL7, IMPDH2, SPINK2, TSC22D1,SPI1(PU.1)	
CLP(C4,C13)	BCL11A, DNTT, RUNX2, MME, HOXA9, NKG7,CCR7	GATA2, SELL
NMPs(C3, neutrophilic,)	NFE2, MYB, LYL1, RUNX1, MPO, CSF3R, CLEC11A, MGST1	RUNX2, **SLC40A1**
GAPs (C1)	GATA2, NFE2, LYL1, MYB, RUNX1, **SLC40A1**	RUNX2, GATA1, KLF1
MEP(C2,C10,C14)	GATA2, NFE2, LYL1, MYB, GATA1,SLC40A1, CLEC11A, TAL1, KLF1, FCER1A, CPA3	SPINK2, CSF3R
C10 (Mast cell or Basophil, MBPs)	MS4A2, TPSB2, HPGDS, **HDC,KIT,** LMO4, T P S A B 1	
C14(Pro-ME)	HBD, TYMS, MCM7	
C5 (Pro-NK)	FCGR3A, GNLY, NKG7	
Monocytes and Macrophages(C6, C7)	FCGR3A, CD68, CST3, LYZ, SPI1, CEBPB, EGR1, CD86, PECAM1, CSF1R	

Figure 1. Two subtypes of progenitors were identified by unsupervised

dimensionality reduction and clustering analysis of single cell data. A, Schematic

showing the progenitor cell enrichment and sequencing work flow. B,17 clusters were

presented by UMAP anlaysis. The progenitors marker of CD34 mainly expressed in

two nearby clusters 6 and 14. C, The heatmap results showed the different expression

surface markers and known progenitors cell related markers in all clusters. Progenitor

cells key regulate genes were identified, such as high expression of CDK6,

SERPINB1, and SOX4. D, Violin plot presented the expression of HSCs traditional

classification marker genes and known progenitor cell marker genes.

Figure 2. Transcriptome differentiation and characterization among re-clustered

CD34 positive subgroups progenitor cells. A, UMAP analysis showed the CD34

positive subgroup cells aggregation in two mainly distinguished directions. B, Violin

plots presented the expression unique marker genes, which demonstrated GATA2

expression of common myeloid progenitors (CMPs) and MME expression of common

lymphoid progenitors (CLPs) characterization. C, The heatmap results showed the

different expression marker genes confirmed the two distinguished directions belong

different lineage cells of CLPs and CMPs. D, Feature plot of AVP and CSF3R

expression showed cluster 1 was the most original progenitors in all of CD34 positive

cells. E, The biology procedure analysis of Gene Ontology showed the high

expression genes of CLPs were enriched in regulation of leukocyte and T cell

activation. F, CMPs mainly high expression genes were enriched in mononuclear cell

differentiation, erythrocyte differentiation and homeostasis.

Figure 3. Dynamics of myeloid progenitor and lymphoid progenitor cells

transcriptomes. A, trajectory analysis showed all clusters distributed in three

branches. B, The expression tendency of unique markers genes was completely

difference among the two lineages of myeloid and lymphoid progenitor. MME and

FLT3 were up-regulated with CLP differentiation branches, with contrary of KIT and

GATA2 up-regulated in CMP lineages. C, Heatmap analysis showed the expression

tendency of selected marker genes in CMP and CLP. D, The expression of MME and

CD38 were clearly inverse during the CLP and CMP lineages priming procedures.

1004

Figure 4. Difference transcription factors promote the progenitor lineage

priming. A, Heatmap results showed the different expression transcription factors

among all progenitors subgroups. B, The activation and expression of key regulators

GATA1 was obviously different among the priming stage of progenitors. C, Feature

plots of the key transcription factors. BCL11A, YY1 and REL were the key regulators

in CLP priming. KLF1, GATA1, and GATA2 were played a critical role in the lineage

priming of CMPs.

1012

Figure 5. The characterization of progenitors subtype in quiescent status and

activated status. A, 16 clusters were presented of integrated CD34 positive

progenitor cells by UMAP analysis. B, The cells distribution among the quiescent

(PBMC) and activate (G-CSF) status, which showed all kinds of progenitors were

persisting existence in quiescent status of adult peripheral blood. C, The percentage

changes column plot of CLP and CMP clusters. The cell numbers and folder were

most obvious changed in the primitive cluster 0. D, Violin plot showed the feature

markers distribution and characterization was consistent with quiescent status. E,

Feature plots of unique marker genes. F, Dot plots presented the expression of unique

marker genes and lineages specific genes.

1023

Figure 6. New hematopoietic hierarchy of hematopoietic progenitor cells

differentiation. A, T-SNE plots showed the re-defined progenitor clusters. B, Dot

plots presented the expression of fate decision transcript factor genes of myeloid

progenitors and lymphoid progenitors. C, Reaffirmed lineage commitment model by

scRNA transcriptional atlas. In the new hematopoietic hierarchy, the HPCs priming

into three progenitor cell types coinstantaneous and later maturing to diversity

hematopoietic lineages. The first progenitor cell type is lymphoid progenitors (CLPs)

branches differentiated to T cell, B cell and NK cells.   The second progenitor cell

types GAPs   branches differentiated to a middle stage of megakaryocytic–erythroid

progenitors (MEPs) first, which controlled by GATA2 and GATA1. In the later stages,

MEPs differentiated to two directions controlled by KIT and KLF1. The KIT high

expression branches will be derived to mast cell or basophil, eosinophil progenitors

(MBPs). The other branch will differentiated to megakaryocytic and erythroid

pro-progenitors (Pro-ME). The third progenitor cell type is neutrophilic and monocyte

progenitors (NMPs), which differentiated to neutrophilic and monocyte. Dashed lines

meanings haven't strong directly evidence of the differentiation relationship which

indicated partial lineage connections. The shaded area indicates the GATA2 and

GATA1 expression domain.

1042

Classical hierarchy of hematopoiesis

HPCs

CLP   CMP

Pro-TBNK   GMP   MEP

T-cells B-cells NK-cells Basophil Pro-Monocyte Megakaryocyte Erythrocyte
                  Neutrophilic   Eosinophil

Reaffirming hierarchy of hematopoiesis

HPCs

HOXA9   CSF3R   NFE2
GAPs
GATA2

CLP   NMPs   MEPs
GATA1

Pro-TBNK   Pro-Monocyte Pro-Neu   Pro-ME   MBPs

T-cells B-cells NK-cells Macrophages Megakaryocyte Basophil Eosinophil
                  Dendritic cell Neutrophilic Erythrocyte Mastcell