Subcortical correlates of consciousness with human single neuron recordings

Michael Pereira[1,2,3*], Nathan Faivre[1,4*], Fosco Bernasconi[1*], Nicholas Brandmeir[5], Jacob Suffridge[2,6], Kaylee Tran[2], Shuo Wang[2,6,7], Victor Finomore[2], Peter Konrad[5], Ali Rezai[2,5*], Olaf Blanke[1,8*]

Affiliations
Laboratory of Cognitive Neuroscience, Neuro-X Institute & Brain Mind Institute, Faculty of Life Sciences, Swiss Federal Institute of Technology (EPFL), Geneva 1202, Switzerland
Department of Neurosciences, WVU Rockefeller Neuroscience Institute, Morgantown, West Virginia
University Grenoble Alpes, Inserm U1216, CHU Grenoble Alpes, Grenoble Institut Neurosciences, 38000 Grenoble, France
University Grenoble Alpes, University Savoie Mont Blanc, CNRS, LPNC, 38000 Grenoble, France
Departments of Neurosurgery, WVU Rockefeller Neuroscience Institute, Morgantown, West Virginia
Department of Computer Science and Electrical Engineering, West Virginia University, Morgantown, WV, 26506, USA
Department of Radiology, Washington University in St. Louis, St. Louis, MO, 63110, USA
Department of Clinical Neurosciences, University Hospital Geneva, Geneva, Switzerland
* equal contribution

Correspondence:
Michael Pereira
michael.pereira@univ-grenoble-alpes.fr
Grenoble Institut Neuroscience
INSERM U1216
Bâtiment Edmond J. Safra
31 Chem. Fortuné Ferrini
38700 La Tronche
France

Keywords: Neural correlates of consciousness, detection, deep brain stimulation, single neuron activity

Author Contributions: MP, NF, FB, AR and OB developed the study concept and contributed to the study design. Neurosurgical procedures were carried out by NB, PK, and AR. Data collection was performed by MP, NF, FB, JS, and KT. Data analysis was carried out by MP, NF, FB. SW provided methodological support and VF supervised data collection. OB supervised the project. MP and NF drafted the paper; all authors provided critical revisions and approved the final version of the paper for submission.

Declaration of interest: The authors declare no competing interests.

Abstract

Subcortical brain structures such as the subthalamic nucleus or the thalamus are involved in regulating motor and cognitive behavior. However, their contribution to perceptual consciousness is still unclear, due to the inherent difficulties of recording subcortical neuronal activity in humans. Here, we asked neurological patients undergoing surgery for deep brain stimulation to detect weak vibrotactile stimuli applied on their hand while recording single neuron activity from the tip of a microelectrode. We isolated putative single neurons in the subthalamic nucleus and thalamus. A significant proportion of neurons modulated their activity while participants were expecting a stimulus. We found that the firing rate of 23% of these neurons differed between detected and undetected stimuli. Our results provide direct neurophysiological evidence of the involvement of the subthalamic nucleus and the thalamus for the detection of vibrotactile stimuli, thereby calling for a less cortico-centric view of the neural correlates of consciousness.

Introduction

Current methods to investigate the *neural correlates of consciousness* aim at contrasting the neural activity associated with different percepts under constant sensory stimulation to identify the minimal set of neuronal events sufficient for a specific conscious percept to occur (Koch et al., 2016; Seth & Bayne, 2022). Typically, this involves asking participants to report whether a stimulus with an intensity around detection threshold is present or not. Taking advantage of the wealth of invasive electrophysiology recordings available, researchers have documented such correlates with detection tasks in rodents (Schmack et al., 2021), birds (Nieder et al., 2020) and non-human primates Shepherd (de Lafuente & Romo, 2006). However, the use of animal models to study consciousness raises specific ethical concerns (Mazor et al., 2023), and requires interpreting behavioral responses with caution (Birch et al., 2022). Research into the neural correlates of consciousness in human volunteers is enriched by the analysis of fine-grained subjective reports to rule out various confounds (e.g attention, memory, report), but suffers from less spatially and temporally resolved physiological measurements. Indeed, only very few studies have found such correlates at the single neuron level (Fried et al., 1997; Gelbard-Sagiv et al., 2018; Pereira et al., 2021; Quiroga et al., 2008; Reber et al., 2017) and only in cortical regions. The role of subcortical structures for perceptual consciousness is theoretically relevant (Aru et al., 2020; Dehaene & Changeux, 2011; Schiff, 2008; Seth & Bayne, 2022; Ward, 2013) with some empirical support from detection studies in non-human primates (Haegens et al., 2014; Tauste Campo et al., 2019; Vazquez et al., 2012, 2013), as well as functional imaging or local field potentials in humans (Kronemer et al., 2022; Levinson et al., 2021). Nonetheless, it remains unknown how the firing rate of subcortical neurons changes when a stimulus is consciously perceived. Single-unit recordings provide a much higher temporal resolution than functional imaging, which helps assess how the neural correlates of consciousness unfold over time. Contrary to local field potentials, single-unit recordings can expose the variety of functional roles of neurons within subcortical regions, thereby offering a potential for a better mechanistic understanding of perceptual consciousness. Here, we recorded individual neurons from the subthalamic nucleus (STN) and thalamus of human participants during 36 deep brain stimulation surgeries. Participants detected vibrotactile stimuli provided at the perceptual threshold and we tested how neurons in both subcortical structures were modulated by the task, the onset of the stimulus or the detection or not of the stimulus.

Results

Task and behavior

Deep brain stimulation surgeries provide a unique opportunity to record the activity of single neurons in subcortical structures of the human brain. Microelectrode recordings are performed routinely after patients are awakened from anesthesia, to allow electrophysiologists and neurosurgeons to identify the target brain structure along the planned trajectory (Figures 1B, S1). During this procedure, we attached a vibrotactile stimulator to the palm of the hand contralateral to the microelectrode recordings and estimated the stimulus intensity corresponding to participants' individual tactile detection threshold. Once stable neuronal activity could be recorded in the target brain region (STN or thalamus), we proceeded to the main experiment, which comprised one or two sessions of 71 trials (total: 48 sessions). Each trial started with an audio "go" cue, followed by a vibrotactile stimulus applied at any time between 0.5 s and 2.5 s after the end of the cue (i.e. stimulation window), except for 20% of catch trials in which no stimulus was applied (Figure 1A). After a random delay ranging from 0.5 to 1 s, a "respond" cue was played, prompting participants to verbally report whether they felt a vibration or not. Therefore, none of the reported analyses are confounded by motor responses. Using a staircase procedure, the stimulus intensity was kept around the detection threshold over the whole experiment. When possible, participants were trained to perform the task prior to the surgery.

Figure 1. Task and behavior. A. Task timeline. Each trial started with an auditory start cue, followed by a 0.5 s delay. Next, the stimulus could occur anytime during a 2 s stimulation window. After a variable 0.5 to 1 s delay, a response cue prompted patients to answer whether or not they detected the stimulus. B. Two example sets of 1 s long microelectrode recordings along the surgical tract showing specific firing for the subthalamic nucleus (left) and the motor thalamus (right). The depth at which the research data was collected is represented as a red dot (see Figure S1 for anatomical correspondence). C. Number of hits, misses, correct rejections (C.R.), and false alarms (F.A.) collected during the main experiment. D. Averages of the absolute vibrotactile intensity in hits and misses in arbitrary units (values cannot be compared between participants). In panels C and D, each small dot represents a participant with Parkinson's Disease (PD, in green) or essential tremor (ET, in orange). Big dots represent averages; error bars represent 95% confidence intervals.

When analyzing tactile perception, we ensured that our results were not contaminated by fluctuating attention and arousal due to the surgical procedure. Based on objective criteria, we excluded specific series of trials from analyses and focused on trials where hits and misses occurred in commensurate proportions (see methods). This procedure led us to keep 36 sessions out of 48 with a mean of 24.0 [95% confidence interval = 22.0, 25.9] hit trials and 22.7 [20.8, 24.5] miss trials. Permutation tests at the single-participant level indicated that detected and missed stimuli were of similar intensity except in 5 sessions for which the intensity of detected stimuli was higher on average. Likewise, detected and missed stimuli had similar onsets, except in 1 session for whom stimuli with late onsets were predominantly missed, and in 2 sessions for whom stimuli with early onsets were predominantly missed. The hit rate was comparable between participants with Parkinson's disease (0.51 [0.49, 0.53]) and essential tremor (0.52 [0.51, 0.53], Wilcoxon rank sum test: W = 114.5, p = 0.45; Figure 1C). Catch trials were separated into 9.1 [8.1 10.1] correct rejections and 2.1 [1.7, 2.6] false alarms, with an equivalent false alarm rate between participants with Parkinson's disease (0.24 [0.19, 0.28]) and essential tremor (0.24 [0.18, 0.30], Wilcoxon rank sum test: W = 145, p = 0.76). Intraoperative behavior was similar to the behavior observed during the training session (Figure S2) and similar to what we found recently in a cohort of healthy participants using the same task (Pereira et al., 2021).

Figure 2. Representative cue- and task-responsive neurons in distinct patients. A-C. Upper panels: firing rates time-locked to the onset of a trial (300 ms long auditory cue; vertical purple shade), compared to a 500 ms pre-cue baseline ("B"). Two significance windows were tested: the post-cue window (500 ms after cue offset; grey horizontal bar; cue-selective neurons) or the stimulation window (800 ms to 2800 ms post-cue; black horizontal bar; task-selective neurons). Asterisks represent statistical significance (p < 0.05).

*Shaded areas indicate bootstrapped standard errors. Inset: corresponding action potentials (shaded area indicates standard deviation; vertical bar corresponds to 100 µV, duration: 2.5 ms). Lower panels: raster plot with trials sorted by stimulus onset (dashed lines) and type: hits (blue), misses (red), correct rejections (C.R.; green), and false alarms (F.A.; black). **A**. Cue-selective neuron in the thalamus. **B**. Cue- and task-selective neurons in the STN. **C**. Task-selective neuron in the thalamus. **D**. Sagittal view of recording locations for thalamic (squares) and subthalamic (circles) targets (see Figure S4A for a coronal view) for patients for which we could obtain anatomical images. Filled circles or squares are cue/task-selective neurons. Legend: VL: ventral lateral thalamus, VPlm: ventral posterior lateral and medial thalamus, VPi: ventral posterior inferior thalamus, STN: subthalamic nucleus, SN: substantia nigra, GPi/e: globus pallidus internalis / externalis,*

Neuronal firing was modulated by the task

We performed a total of 48 (STN: 25, Thal: 23) successful microelectrode recording sessions during 36 surgeries for deep brain stimulation electrode implantation. We isolated 50 putative single neurons (STN: 26, Thal: 24) according to spike sorting metrics (Figure S3A-G). We ensured that all neurons showed stable spike amplitudes during the recording (Figure S3H-J). We also ensured that for every analysis, a minimum of 20 trials per condition were kept after removing artifacts. First, we looked for cue-selective neurons that modulate their firing rate during the 500 ms delay following the end of the "go" cue, compared to a 500 ms pre-cue baseline period. There were 8 / 44 (18 %) cue-selective neurons (Figure 2A; 6 neurons were removed from the analysis due to an insufficient number of trials). We confirmed that these 8 cue-selective neurons could not have been obtained by chance by comparing this number to a null distribution obtained by permuting the sign of the trial-wise differences 1000 times (sign permutation test: $p < 0.001$). The proportion of cue-selective neurons was not significantly different in the STN (21%) and thalamus (15%; difference: $p = 0.31$, sign permutation test) and 6 out of 8 neurons showed a decrease in firing rate compared to the pre-cue baseline.

***Figure 3. Representative stimulus-responsive neurons in distinct patients. A-C**. Upper panels: firing rate time-locked to the onset of the stimulus (100 ms vibrotactile stimulation; blue sinusoid) for all trials. Green trace represents corresponding activity for catch trials. Thick horizontal black segments show significant time windows. Shaded areas indicate bootstrapped standard errors. Inset: corresponding action potentials (shaded area indicates standard deviation; vertical bar corresponds to 100 µV, duration: 2.5 ms).*

*Lower panels: raster plot. The 300 ms pre-stimulus baseline was used only for statistics. **D**. Sagittal view of recording locations for thalamic (squares) and subthalamic (circles) targets (see Figure S4B for a coronal view) for patients for which we could obtain anatomical images. . Filled circles or squares are sensory-selective neurons. Legend: VL: ventral lateral thalamus, VPlm: ventral posterior lateral and medial thalamus, VPi: ventral posterior inferior thalamus, STN: subthalamic nucleus, SN: substantia nigra, GPi/e: globus pallidus internalis / externalis,*

Next, we investigated how many neurons showed task-selective modulations by comparing firing rates during the 2 s stimulation window to the 500 ms pre-cue baseline, indicating a modulation of their firing rate when a stimulus is expected. There were 9 / 44 (20 %) task-selective neurons (sign permutation test: p = 0.003 with a similar proportion in the STN (21 %) and thalamus (20 %; Figure 2B-D). Interestingly, 8 out of 9 neurons decreased their firing rate relative to the pre-cue baseline. In both regions, a significant proportion (44 %; sign permutation test: p < 0.001) of the task-selective neurons were also cue-selective, modulating their firing rate before any sensory stimulation necessary for a decision occurred. Therefore, these cue- and task-selective neurons are unlikely to be involved in decision-related action selection or cancellation (Bastin et al., 2014; Mosher et al., 2021) but should be involved in the detection task *per se*.

Neuronal firing was modulated by the stimulus

We then searched for neurons that modulate their firing rate after the stimulus onset compared to a 300 ms pre-stimulus baseline while correcting for possible drifts in the firing rate during the trial (see methods). We found 8 / 37 such stimulus-selective neurons (22%, sign permutation test: p = 0.011; Figure 3A-D; 13 neurons were removed due to an insufficient number of trials), with 29% in the STN and 15% in the thalamus. These differences occurred 211 ms ± 34 after the stimulus onset, lasted for an average of 126 ms ± 28, and 6 out of 8 neurons showed a decrease in firing rate after the stimulus onset. These results show that subthalamic and thalamic neurons are modulated by stimulus onset, irrespective of whether it was reported or not, even though no immediate motor response was required.

***Figure 4. Representative perception-selective neurons in distinct patients. A-C** Upper panels: firing rate time-locked to the onset of the stimulus (100 ms vibrotactile stimulation; blue sinusoid) for hits (light blue) and misses (red). Thick horizontal black segments show significant time windows. Shaded areas indicate bootstrapped standard errors. Inset: corresponding action potentials (shaded area indicates*

standard deviation; vertical bar corresponds to 100 µV, duration: 2.5 ms). Lower panels: raster plot for hits (light blue) and misses (red). See Figure S5 for all neurons that are perception- or sensory-selective. **D.** *Sagittal view of recording locations for thalamic (squares) and subthalamic (circles) targets (see Figure S4C for a coronal view) for patients for which we could obtain anatomical images. Filled circles or squares are perception-selective neurons. Legend: VL: ventral lateral thalamus, VPlm: ventral posterior lateral and medial thalamus, VPi: ventral posterior inferior thalamus, STN: subthalamic nucleus, SN: substantia GPi/e: globus pallidus internalis / externalis.*

Neuronal firing was modulated by tactile perception

Having identified subcortical neurons that were cue-, task- or stimulus-selective, we next sought to assess the role of these structures in conscious detection by comparing firing rates time-locked to detected vs missed stimuli. Of the 50 neurons recorded, 35 were associated with periods of high-quality behavior, allowing us to assume tactile stimulation at the perceptual threshold. We found 8 neurons (23 %) showing a significant difference between hits and misses after stimulus onset (trial permutation test: $p = 0.0060$; Figure 4A-D, S5). Each neuron was found in a different participant. The proportion of these perception-selective neurons was similar in the STN (27 %) and the thalamus (20 %; difference: $p = 0.529$; trial permutation test). These differences in firing rates occurred on average 160 ms ± 36 after the stimulus onset and lasted for an average of 100 ms ± 10. We note that the timings of sensory and perception effects in Figures 3 and 4 showed a bimodal distribution with an early cluster (149 ms for sensory neurons; 121 ms for perception neurons; c.f. methods) and a later cluster (330 ms for sensory neurons; 315 ms for perception neurons; Figure 5). We note that, 6 out of 8 neurons had higher firing rates for missed trials than hit trials. None of the aforementioned neurons showed sustained differences between the highest and lowest stimulus amplitudes nor between early and late stimulus onset within the 2 s stimulus window (Figure S6). Our control analyses confirm that our results do not stem from slight differences in stimulus amplitudes due to the staircase procedure or spurious differences induced by the start or response cues. Qualitatively, we found little overlap between task-, stimulus- and perception-selective neurons (one neuron selective for all three categories, one task- and perception-selective neuron, two stimulus- and perception-selective neurons and one task- and perception-selective neuron; Figure S7). This result suggests that these two subcortical structures display neurons with several different functional roles. We also found no clear indication that neurons with a beta-band oscillatory component were more or less selective.

Figure 5. *Number of neurons with a significant effect over time for sensory-selective neurons (green) and perception-selective neurons (red). Arrows show the corresponding early and late cluster centroids (green for sensory; red for perception).*

Discussion

The importance of cortico-subcortical loops for physiological and cognitive functions is well-established (Shepherd & Yamawaki, 2021) Yet, while the role of subcortical structures in perceptual consciousness is largely acknowledged (Aru et al., 2020; Dehaene & Changeux, 2011; Koch et al., 2016; Shepherd & Yamawaki, 2021; Ward, 2013), it remains poorly described in humans. This limit is likely due to the difficulty of recording subcortical activity in awake humans capable of providing conscious reports under controlled experimental conditions. We report the first intraoperative recordings of subcortical neurons in awake individuals during a detection task. By imposing a delay between the end of the tactile stimulation window and the subjective report, we ensured that neuronal responses reflected stimulus detection and not mere motor responses. In addition, because stimuli were applied on the palm, we asked participants to provide detection responses orally to avoid confounding neural activity related to sensory and motor processes of the upper limb. Our main result is that the activity of subcortical neurons co-varies with subjective reports following the presentation of detected vs missed tactile stimuli. This result confirms that the neuronal underpinnings of tactile detection can be observed at the scale of single neurons in humans (Fried et al., 1997; Gelbard-Sagiv et al., 2018; Pereira et al., 2021; Quiroga et al., 2008; Reber et al., 2017) but also shows for the first time that they are not limited to the cortex.

Our findings that neurons in the thalamus modulate their activity according to tactile detection adds to the existing evidence in favor of the role of the thalamus for perceptual consciousness. Indeed, thalamic activity and more precisely thalamocortical loops are often considered key to gate sensory stimuli to conscious access (Ward, 2013). In non-human primates, for example, oscillatory thalamic activity predicts tactile detection (Haegens et al., 2014), and functional interactions between the somatosensory thalamus and the cortex increase when a tactile stimulus is detected (Tauste Campo et al., 2019). In humans, thalamic local field potentials and fMRI activity were higher for seen vs unseen stimuli (Kronemer et al., 2022; Levinson et al., 2021) and causal effects of thalamic stimulation on the levels of consciousness have been found (Schiff et al., 2007). Future studies with higher neuronal yields will be helpful in assessing the contribution of distinct thalamic territories to tactile consciousness, focusing notably on the ventral caudal part, which contains

neurons with tactile receptive fields. Moreover, given the prominent role of the thalamus in multisensory processing (Yang et al., 2025), it will be interesting to assess if it is specifically involved in tactile consciousness or if it has a supramodal contribution, akin to what is found in the cortex (Filimonov et al., 2022; Noel et al., 2018; Sanchez et al., 2020).

Concerning the subthalamic nucleus, a possibility is that perception-selective neurons determine stimulus detection through the regulation of decisional processes. Indeed, previous studies reported a modulatory role of subthalamic activity on decisional processes, notably by elevating the decisional threshold on accumulated sensory evidence (Bogacz & Gurney, 2007; Cavanagh et al., 2011; Green et al., 2013; Herz et al., 2016). In a recent study in which we measured the activity of cortical neurons in a similar task, we showed that evidence accumulation is also at play during conscious detection (Pereira et al., 2021). Based on this finding, we proposed that percepts fade in and out of consciousness when evidence accumulated by cortical neurons passes a given threshold (Pereira et al., 2022). The present results, therefore, indicate that the contribution of subthalamic neurons to decisional processes is not limited to discrimination tasks or motor planning, but may also regulate the threshold at which accumulated evidence gives rise to a conscious percept. Considering the inhibitory role of the subthalamic nucleus on the cortex (Mink, 1996), the fact that many of the perception-selective neurons we found had higher firing rate for misses than for hits suggests a role in elevating that threshold, similar to what is found in decision tasks manipulating conflict or cautiousness and requiring immediate responses (Bastin et al., 2014; Cavanagh et al., 2011; Frank et al., 2007; Herz et al., 2016; Mosher et al., 2021). Thus, our results suggest that the STN plays an important role in a subcortical network gating conscious access, although it might not encode conscious content *per se* (Aru et al., 2012).

Apart from perception-selective neurons, we also found a distinct population of neurons in both the STN and thalamus that modulated their firing rate both after the cue and during the task, and therefore much before the stimulus onset. These neurons cannot be involved in detection-related processes but could instead be involved in task switching (Hikosaka & Isoda, 2010; Laquitaine et al., 2024). Micro-electrode recordings could therefore reconcile the multiple roles ascribed to subcortical structures by exposing the variety of functional roles of the neurons inside those regions. We also found neurons that modulated their firing rates after the stimulus onset, irrespective of detection, similar to animal works in the STN (Al Tannir et al., 2023; Coizet et al., 2009) and thalamus (Tauste Campo et al., 2019; Vazquez et al., 2012). Our results should be taken with caution as they are based on a small number of neurons due to the high complexity of intraoperative recordings, and because the number of trials we could collect was not sufficient to test the computational mechanisms underlying the neuronal activity we recorded. Interestingly, we found that both sensory and perception effects were could be grouped into an early and late cluster. The early cluster's average timing around 150 ms post-stimulus corresponds to the onset of a putative cortical correlate of tactile consciousness, the somatosensory awareness negativity (Dembski et al., 2021). Similar electroencephalographic markers are found in the visual and auditory modality. It is unclear, however, whether these markers are related to perceptual consciousness or selective attention (Dembski et al., 2021). The later cluster is centered around 300 ms and could correspond to a well-known electroencephalographic marker, the P3b (Polich, 2007) whose association with perceptual consciousness has been questioned (Dembski et al., 2021; Pitts et al., 2014) although brain activity related to consciousness has been observed at similar timing even in the absence of report demands (Sergent et al., 2021; Stockart et al., 2024). It is also

important to note that these clusters contain neurons with both increased and decreased firing rates following stimulus onset, similar to what was observed previously in the posterior parietal cortex (Pereira et al., 2021). Future studies ruling out the presence of motor preparation triggered by perceived stimuli (Bennur & Gold, 2011; Fang et al., 2024; Twomey et al., 2016) and verifying that similar neuronal activity occurs in the absence of task-demands (no-reports; (Tsuchiya et al., 2015)) or attention (Wyart & Tallon-Baudry, 2008) will be useful to support that subcortical neurons contribute specifically to perceptual consciousness.

In sum, our study provides neurophysiological evidence from single neurons in humans that subcortical structures play a significant role in tactile detection either by themselves (Ward, 2013) or through their numerous connections with the cortex (Dehaene & Changeux, 2011). A comprehensive account of the neural correlates of consciousness should, therefore, not be cortico-centric but also consider subcortical contributions.

Methods

Participants

We recorded high impedance electrophysiological signals from microelectrodes inserted intraoperatively in the subthalamic nucleus of 32 participants with Parkinson disease or essential tremor undergoing deep brain stimulation electrode implantation surgeries (N = 36; 4 participants had two surgeries, one for each side). For individuals with Parkinson's disease, the age at the time of the recording was 60.4 ± 2.7 years and the average UPDRS III score was 40.6 ± 3.0 prior to surgery and was reduced to 20.8 ± 2.8 after the surgery (p = 0.0015, z = 3.18). We also recorded intraoperatively in the thalamus of individuals with essential tremor undergoing deep brain stimulation surgeries. The age at the time of the recording was 68.9 ± 3.2 years and the average TETRA motor score was 20.1 ± 2.9 prior to surgery. The study was approved by the institutional review board of the West Virginia University Hospital (WVU02HSC17; #1709745061) and all participants provided written informed consent prior to any data collection.

Experimental procedure

Participants performed a tactile detection task programmed in Matlab using the Psychophysics toolbox (Brainard, 1997; Kleiner, s. d.; Pelli, 1997). When possible, participants were trained a few days before the surgery (N = 18 / 36 surgeries). Participants sat in a reclining chair in a quiet room (training session) or were lying in the operating room (main session). Every trial started with a 300 ms long auditory "go" cue delivered through an external loudspeaker placed near the participants. Following the end of the go cue and a delay of 500 ms, a 100 ms vibrotactile stimulus could be delivered at any time during a two second stimulation window (i.e., uniform distribution between 0.8 and 2.8 s after the onset of the go cue; Figure 1A) on the lateral palm contralateral to the deep brain implant. Stimuli were applied using a MMC3 Haptuator vibrotactile device from TactileLabs Inc. (Montréal, Canada) driven by a 230 Hz sinusoid audio signal. Participants reported orally whether they felt the stimulus or not and whether they were confident in their answer or not after an auditory "respond" cue played one second after the end of the stimulation window. The participants responses could thus consist in "yes, sure", "yes, unsure", "no, sure" and "no, unsure". The task was stopped after two sessions of 71 trials, or before in case of discomfort or other clinical constraints. As –upon waking from anesthesia– most participants did not use both confidence levels, confidence data was therefore not analyzed.

To keep the vibrotactile stimulus intensity around the detection threshold, we first conducted a rough threshold search by presenting a series of stimuli whose intensity decreased by steps of 5% until participants reported not feeling them anymore. Then we presented series of low intensity stimuli whose intensities increased by step of 5% until participants reported feeling them again. These procedures were repeated until the experimenter deemed the results satisfying. We took the average between the thresholds obtained during these procedures as a seed for the main task. During the main task, a 1up/1down adaptive staircase procedure (Levitt, 1971) ensured that the intensity was kept around the perceptual threshold by increasing the intensity by 5% after miss trial and decreasing the intensity by 5% after a hit trial. Of note, the absolute stimulus intensity is not informative and cannot be compared across patients and sessions, as it varied according to different factors (e.g. the length of the cable or the manner with which the tactile stimulator was strapped onto the palm).

Surgical procedure

STN or thalamus targets and trajectories were defined preoperatively using CranialSuite (Neurotargeting Inc., Nashville, US) based on MRI scans. Both targets were then defined with respect to the AC-PC (commissural) line using standard atlas-based methods and refined based on individual anatomy. The entry point was chosen approximately 2 to 3 cm from the midline and 1 cm anterior from the coronal suture and adjusted to individual anatomy in order to avoid traversing brain sulci, lateral ventricles or the medial bridging veins. Scalp incisions and burr-hole drilling were performed under local (lidocaine) and general (propofol) anesthesia and a microelectrode (FHC, Maine, US) was inserted through a guide cannula using a microdrive placed either on a Leksell frame (N = 13 surgeries) or a 3D printed mould (N = 23 surgeries).

Electrophysiology

Once the microelectrode reached the target brain structure (STN or thalamus), the speed of the microdrive was reduced and neuronal activity was streamed to a loudspeaker, allowing the electrophysiologist to verify the depth of the preplanned trajectory. The main research task was initiated when a neuron showed stable activity for a few tens of seconds and the anatomical localization was confirmed by the electrophysiologist. Recording depths were saved and used offline to define the anatomical localization (see Anatomical localization section). Electrophysiological data were recorded from the 5 mm tip of the microelectrode, referenced to the guide cannula and an adaptive line noise canceller was applied. Data were digitized either using a Guideline 4000 LP+ amplifier (FHC, Maine, US) at 30 kHz (N = 21 surgeries), or using a Guideline 5 amplifier (FHC, Maine, US) at 32 kHz and resampled offline to 30 kHz (N = 14 surgeries).

Anatomical localization

For 34 / 50 neurons, preoperative MRI and postoperative CT scans (co-registered in patient native space using CranialSuite) were available to precisely reconstruct surgical trajectories and recording locations (for the remaining 16 neurons, localizations were based on neurosurgical planning and confirmed by electrophysiological recordings at various depths). Recording depths were inspected along the trajectories in patient native space, projected to an MNI-coordinate space and compared against the Ilinsky atlas (Ilinsky et al., 2018) which delineates distinct thalamic sub-territories based on a marker of γ-aminobutyric acid on sections post-mortem human brains.

Behavioral analyses

We used R 4.1.2 (Team R, 2020) and the tidyverse (Wickham et al., 2019) package to analyze behavioral data. Permutation tests were performed by permuting hit and miss trials over 1000 iterations for each participant. Non-parametric p-values were estimated by counting the permutations for which the difference between hits and misses was higher in the observed compared to the shuffled data.

As titrating and keeping the vibrotactile stimulation intensity to the perceptual level after anesthesia was a challenging task, we took great care in keeping only the highest quality recordings. We estimated the trial-by-trial hit-rate using a sliding window of 11 trials (for the first and last 5 trials, we mirrored trials to avoid border effects). Any trial with a hit-rate out of the]25, 75[% range were removed from further analysis comparing hit to miss trials. If less than 10 hit and 10 miss trials were kept by this procedure, the session (and its corresponding neurons) was removed from subsequent analyses (13 / 48 sessions; 27 %).

Spike sorting and firing rate estimation

Each microelectrode recording was filtered between 300 and 3000 Hz and visually inspected. Artifacts such as cross-talk from the participants' vocal responses were marked and replaced by noise with a standard deviation matching the second pre- and post-artifact. We performed this procedure to avoid spuriously lowering the thresholds for neuronal spike detection. The timing of these artifactual epochs were saved in order to reject affected trials in later analyses. Neuronal spikes were detected and clustered using an online semi-automatic spike sorting algorithm (Osort) (Rutishauser et al., 2006). Each resulting cluster of neurons was inspected based on common metrics such as spike waveform, percentage of inter-spike interval below 3 ms, signal-to-noise ratio and power spectral densities and possibly merged with other clusters. Finally, the resulting curated neurons were labeled as *putative single neuron* or *multiunit*, depending on the spike waveforms, peak amplitude distribution and the percentage of inter-spike interval below 3 ms. Electrophysiological signals were realigned either to the onset of the "go" cue (Figures 2) or to the onset of the stimulus (Figures 3-4), which was precisely obtained by applying a matched filter to a copy of the audio signal used to drive the vibrotactile stimulator we simultaneously recorded with the electrophysiological data. We estimated instantaneous firing rates using a sliding Gaussian kernel with a standard deviation of 40 ms and 1 ms steps. When displaying the resulting average firing rates over time, we estimated the standard error of the mean using a bootstrap procedure with 1000 resamplings.

Identification of selective neurons

To deal with possibly non-parametric distributions, we used Wilcoxon rank sum test or sign test instead of t-tests to test differences between distributions. We used permutation tests instead of Binomial tests to test whether a reported number of neurons could have been obtained by chance. For each analysis, we verified that the reported number of neurons could not have been obtained by chance by comparing this number to a null distribution using permutation tests (Maris & Oostenveld, 2007). For paired tests with respect to a baseline, we randomly flipped the sign of the difference between the firing rate during the trial and during the baseline (*sign permutation test* in the main text) and for unpaired tests, we randomly shuffled the conditions (i.e. a hit trial could be randomly assigned to a hit or a miss trial; *trial permutation test* in the main text). To obtain a p-value, we compared the number of selective neurons to a null distribution obtained by randomly permuting the data 1000 times. This procedure allowed us to show that the number of selective neurons could not have been obtained by chance while controlling for multiple comparisons over time. Similarly, to test whether the proportion of neurons was different in the STN compared to the thalamus, we compared the absolute difference in the proportion of neurons in each anatomical location to a null distribution obtained by random permutations.

To identify cue-selective neurons we compared the number of spikes in a 500 ms baseline preceding the "go" cue to the number of spikes in a 500 ms period following the offset of the "go cue" using a two-tailed non-parametric sign test. Similarly, we identified task-responsive neurons by comparing the mean number of spikes in a 500 ms baseline preceding the "go" cue to the mean number of spikes during the 2 s stimulation window and performing a permutation test. We compared the differences in the proportion of selective neurons in the STN and thalamus, to the same differences observed in the shuffled data to assess its significance. Finally, we also compared the number of cue- and task-selective neurons to the same number observed in the shuffled data to assess whether the overlap was significant.

To identify sensory-selective neurons, we assumed that subcortical signatures of stimulus detection ought to be found early following its onset and looked for differences in the firing rates during the first 400 ms post-stimulus onset compared to a 300 ms pre-stimulus baseline. To correct for possible drifts occurring during the trial, we subtracted the average cue-locked activity from catch trials to the cue-locked activity of each stimulus-present trials before realigning to stimulus onset. We defined a cluster as a set of adjacent time points for which the firing rates were significantly different between hits and misses, as assessed by a non-parametric sign rank test. A putative neuron was considered sensory-selective when the length of a cluster was above 80 ms, corresponding to twice the standard deviation of the smoothing kernel used to compute the firing rate. Whether for the shuffled data or the observed data, if more than one cluster was obtained, we discarded all but the longest cluster. This permutation test allowed us to control for multiple comparisons across time and participants.

For perception-selective neurons, we looked for differences in the firing rates between hit and miss trials during the first 400 ms post-stimulus onset. We defined a cluster as a set of adjacent time points for which the firing rates were significantly different between hits and misses as assessed by a non-parametric Wilcoxon rank sum test. As for sensory-selective neurons, a putative neuron was considered perception-selective when the length of a cluster was above 80 ms, corresponding to twice the standard deviation of the smoothing kernel used to compute the firing rate and we discarded all but the longest cluster.

To measure bimodal timings of effect latencies, we fitted a two-component Gaussian mixture distribution to the data in Figure 5 by minimizing the mean square error with an interior-point method. We took the best of 20 runs with random initialization points and verified that the resulting mean square error was markedly (> 4 times) better than using a single component.

Data and code availability

Data and code necessary to replicate our results are available online (https://gitlab.com/michael.pereira/subcortical-ncc).

Further information and requests should be directed to and will be fulfilled by the lead contact, Michael Pereira (michael.pereira@univ-grenoble-alpes.fr).

Acknowledgments: MP was supported by two Postdoc.Mobility fellowships from the Swiss National Science Foundation (P2ELP3_187974; P400PM_199251). NF has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement no. 803122). OB is supported by the Bertarelli Foundation, the Swiss National Science Foundation, and the European Science Foundation.

References

Al Tannir, R., Pautrat, A., Baufreton, J., Overton, P. G., & Coizet, V. (2023). The Subthalamic Nucleus : A Hub for Sensory Control via Short Three-Lateral Loop Connections with the Brainstem? *Current Neuropharmacology*, *21*(1), 22-30. https://doi.org/10.2174/1570159X20666220718113548

Aru, J., Bachmann, T., Singer, W., & Melloni, L. (2012). Distilling the neural correlates of consciousness. *Neuroscience & Biobehavioral Reviews*, *36*(2), 737-746. https://doi.org/10.1016/j.neubiorev.2011.12.003

Aru, J., Suzuki, M., & Larkum, M. E. (2020). Cellular Mechanisms of Conscious Processing. *Trends in Cognitive Sciences*, *24*(10), 814-825. https://doi.org/10.1016/j.tics.2020.07.006

Bastin, J., Polosan, M., Benis, D., Goetz, L., Bhattacharjee, M., Piallat, B., Krainik, A., Bougerol, T., Chabardès, S., & David, O. (2014). Inhibitory control and error monitoring by human subthalamic neurons. *Translational Psychiatry*, *4*(9), e439-e439. https://doi.org/10.1038/tp.2014.73

Bennur, S., & Gold, J. I. (2011). Distinct Representations of a Perceptual Decision and the Associated Oculomotor Plan in the Monkey Lateral Intraparietal Area. *Journal of Neuroscience*, *31*(3), 913-921. https://doi.org/10.1523/JNEUROSCI.4417-10.2011

Birch, J., Broom, D. M., Browning, H., Crump, A., Ginsburg, S., Halina, M., Harrison, D., Jablonka, E., Lee, A. Y., Kammerer, F., Klein, C., Lamme, V., Michel, M., Wemelsfelder, F., & Zacks, O. (2022). How Should We Study Animal Consciousness Scientifically? *Journal of Consciousness Studies*, *29*(3), 8-28. https://doi.org/10.53765/20512201.29.3.008

Bogacz, R., & Gurney, K. (2007). The Basal Ganglia and Cortex Implement Optimal Decision Making Between Alternative Actions. *Neural Computation*, *19*(2), 442-477. https://doi.org/10.1162/neco.2007.19.2.442

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433-436. https://doi.org/10.1163/156856897X00357

Cavanagh, J. F., Wiecki, T. V., Cohen, M. X., Figueroa, C. M., Samanta, J., Sherman, S. J., & Frank, M. J. (2011). Subthalamic nucleus stimulation reverses mediofrontal influence over decision threshold. *Nature Neuroscience*, *14*(11), 1462-1467. https://doi.org/10.1038/nn.2925

Coizet, V., Graham, J. H., Moss, J., Bolam, J. P., Savasta, M., McHaffie, J. G., Redgrave, P., & Overton, P. G. (2009). Short-Latency Visual Input to the Subthalamic Nucleus Is Provided by the Midbrain Superior Colliculus. *The Journal of Neuroscience*, *29*(17), 5701-5709. https://doi.org/10.1523/JNEUROSCI.0247-09.2009

Dehaene, S., & Changeux, J.-P. (2011). Experimental and Theoretical Approaches to Conscious Processing. *Neuron*, *70*(2), 200-227. https://doi.org/10.1016/j.neuron.2011.03.018

de Lafuente, V., & Romo, R. (2006). Neural correlate of subjective sensory experience gradually builds up across cortical areas. *Proceedings of the National Academy of Sciences*, *103*(39), 14266-14271. https://doi.org/10.1073/pnas.0605826103

Dembski, C., Koch, C., & Pitts, M. (2021). Perceptual awareness negativity : A physiological correlate of sensory consciousness. *Trends in Cognitive Sciences*, *25*(8), 660-670. https://doi.org/10.1016/j.tics.2021.05.009

Fang, Z., Dang, Y., Ling, Z., Han, Y., Zhao, H., Xu, X., & Zhang, M. (2024). The involvement of the human prefrontal cortex in the emergence of visual awareness. *eLife*, *12*, RP89076. https://doi.org/10.7554/eLife.89076

Filimonov, D., Railo, H., Revonsuo, A., & Koivisto, M. (2022). Modality-specific and modality-general electrophysiological correlates of visual and auditory awareness : Evidence from a bimodal ERP experiment. *Neuropsychologia*, *166*, 108154. https://doi.org/10.1016/j.neuropsychologia.2022.108154

Frank, M. J., Samanta, J., Moustafa, A. A., & Sherman, S. J. (2007). Hold Your Horses : Impulsivity, Deep Brain Stimulation, and Medication in Parkinsonism. *Science*, *318*(5854), 1309-1312. https://doi.org/10.1126/science.1146157

Fried, I., MacDonald, K. A., & Wilson, C. L. (1997). Single Neuron Activity in Human Hippocampus and Amygdala during Recognition of Faces and Objects. *Neuron*, *18*(5), 753-765. https://doi.org/10.1016/S0896-6273(00)80315-3

Gelbard-Sagiv, H., Mudrik, L., Hill, M. R., Koch, C., & Fried, I. (2018). Human single neuron activity precedes emergence of conscious perception. *Nature Communications*, *9*(1), 2057. https://doi.org/10.1038/s41467-018-03749-0

Green, N., Bogacz, R., Huebl, J., Beyer, A.-K., Kühn, A. A., & Heekeren, H. R. (2013). Reduction of Influence of Task Difficulty on Perceptual Decision Making by STN Deep Brain Stimulation. *Current Biology*, *23*(17), 1681-1684. https://doi.org/10.1016/j.cub.2013.07.001

Haegens, S., Vazquez, Y., Zainos, A., Alvarez, M., Jensen, O., & Romo, R. (2014). Thalamocortical rhythms during a vibrotactile detection task. *Proceedings of the National Academy of Sciences*, *111*(17), E1797-E1805. https://doi.org/10.1073/pnas.1405516111

Herz, D. M., Zavala, B. A., Bogacz, R., & Brown, P. (2016). Neural Correlates of Decision Thresholds in the Human Subthalamic Nucleus. *Current Biology*, *26*(7), 916-920. https://doi.org/10.1016/j.cub.2016.01.051

Hikosaka, O., & Isoda, M. (2010). Switching from automatic to controlled behavior : Cortico-basal ganglia mechanisms. *Trends in Cognitive Sciences*, *14*(4), 154-161. https://doi.org/10.1016/j.tics.2010.01.006

Ilinsky, I., Horn, A., Paul-Gilloteaux, P., Gressens, P., Verney, C., & Kultas-Ilinsky, K. (2018). Human Motor Thalamus Reconstructed in 3D from Continuous Sagittal Sections with Identified Subcortical Afferent Territories. *Eneuro*, *5*(3), ENEURO.0060-18.2018. https://doi.org/10.1523/ENEURO.0060-18.2018

Kleiner, M. (s. d.). *What's new in Psychtoolbox-3?* 89.

Koch, C., Massimini, M., Boly, M., & Tononi, G. (2016). Neural correlates of consciousness : Progress and problems. *Nature Reviews Neuroscience*, *17*(5), 307-321. https://doi.org/10.1038/nrn.2016.22

Kronemer, S. I., Aksen, M., Ding, J. Z., Ryu, J. H., Xin, Q., Ding, Z., Prince, J. S., Kwon, H., Khalaf, A., Forman, S., Jin, D. S., Wang, K., Chen, K., Hu, C., Agarwal, A., Saberski, E., Wafa, S. M. A., Morgan, O. P., Wu, J., … Blumenfeld, H. (2022). Human visual consciousness involves large scale cortical and subcortical networks independent of task report and eye movement activity. *Nature Communications*, *13*(1), 7342. https://doi.org/10.1038/s41467-022-35117-4

Laquitaine, M., Polosan, M., Kahane, P., Chabardes, S., Yelnik, J., Fernandez-Vidal, S., Domenech, P., & Bastin, J. (2024). Optimal level of human intracranial theta activity for behavioral switching in the subthalamo-medio-prefrontal circuit. *Nature Communications*, *15*(1), 7827. https://doi.org/10.1038/s41467-024-52290-w

Levinson, M., Podvalny, E., Baete, S. H., & He, B. J. (2021). Cortical and subcortical signatures of conscious object recognition. *Nature Communications*, *12*(1), 2930. https://doi.org/10.1038/s41467-021-23266-x

Levitt, H. (1971). Transformed Up Down Methods in Psychoacoustics. *The Journal of the Acoustical Society of America*, *49*(2B), 467-477. https://doi.org/10.1121/1.1912375

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177-190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Mazor, M., Brown, S., Ciaunica, A., Demertzi, A., Fahrenfort, J., Faivre, N., Francken, J. C., Lamy, D., Lenggenhager, B., Moutoussis, M., Nizzi, M.-C., Salomon, R., Soto, D., Stein, T., & Lubianiker, N. (2023). The Scientific Study of Consciousness Cannot and Should Not Be Morally Neutral. *Perspectives on Psychological Science*, *18*(3), 535-543. https://doi.org/10.1177/17456916221110222

Mink, J. W. (1996). The Basal Ganglia : Focused Selection and Inhibition of Competing Motor Programs. *Progress in Neurobiology*, *50*(4), 381-425. https://doi.org/10.1016/S0301-0082(96)00042-1

Mosher, C. P., Mamelak, A. N., Malekmohammadi, M., Pouratian, N., & Rutishauser, U. (2021). Distinct roles of dorsal and ventral subthalamic neurons in action selection and cancellation. *Neuron*, *109*(5), 869-881.e6. https://doi.org/10.1016/j.neuron.2020.12.025

Nieder, A., Wagener, L., & Rinnert, P. (2020). A neural correlate of sensory consciousness in a corvid bird. *Science*, *369*(6511), 1626-1629. https://doi.org/10.1126/science.abb1447

Noel, J.-P., Simon, D., Thelen, A., Maier, A., Blake, R., & Wallace, M. T. (2018). Probing Electrophysiological Indices of Perceptual Awareness across Unisensory and Multisensory Modalities. *Journal of Cognitive Neuroscience*, *30*(6), 814-828. https://doi.org/10.1162/jocn_a_01247

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics : Transforming numbers into movies. *Spatial Vision*, *10*(4), 437-442.

Pereira, M., Megevand, P., Tan, M. X., Chang, W., Wang, S., Rezai, A., Seeck, M., Corniola, M., Momjian, S., Bernasconi, F., Blanke, O., & Faivre, N. (2021). Evidence accumulation relates to perceptual consciousness and monitoring. *Nature Communications*, *12*(1), 3261. https://doi.org/10.1038/s41467-021-23540-y

Pereira, M., Perrin, D., & Faivre, N. (2022). A leaky evidence accumulation process for perceptual experience. *Trends in Cognitive Sciences*, S1364661322000614. https://doi.org/10.1016/j.tics.2022.03.003

Pitts, M. A., Padwal, J., Fennelly, D., Martínez, A., & Hillyard, S. A. (2014). Gamma band activity and the P3 reflect post-perceptual processes, not visual awareness. *NeuroImage*, *101*, 337-350. https://doi.org/10.1016/j.neuroimage.2014.07.024

Polich, J. (2007). Updating P300 : An integrative theory of P3a and P3b. *Clinical Neurophysiology*, *118*(10), 2128-2148. https://doi.org/10.1016/j.clinph.2007.04.019

Quiroga, R. Q., Mukamel, R., Isham, E. A., Malach, R., & Fried, I. (2008). Human single-neuron responses at the threshold of conscious recognition. *Proceedings of the National Academy of Sciences*, *105*(9), 3599-3604. https://doi.org/10.1073/pnas.0707043105

Reber, T. P., Faber, J., Niediek, J., Boström, J., Elger, C. E., & Mormann, F. (2017). Single-Neuron Correlates of Conscious Perception in the Human Medial Temporal Lobe. *Current Biology*, *27*(19), 2991-2998.e2. https://doi.org/10.1016/j.cub.2017.08.025

Rutishauser, U., Schuman, E. M., & Mamelak, A. N. (2006). Online detection and sorting of extracellularly recorded action potentials in human medial temporal lobe recordings, in vivo. *Journal of Neuroscience Methods*, *154*(1-2), 204-224. https://doi.org/10.1016/j.jneumeth.2005.12.033

Sanchez, G., Hartmann, T., Fuscà, M., Demarchi, G., & Weisz, N. (2020). Decoding across sensory modalities reveals common supramodal signatures of conscious perception. *Proceedings of the National Academy of Sciences*, *117*(13), 7437-7446. https://doi.org/10.1073/pnas.1912584117

Schiff, N. D. (2008). Central Thalamic Contributions to Arousal Regulation and Neurological Disorders of Consciousness. *Annals of the New York Academy of Sciences*, *1129*(1), 105-118. https://doi.org/10.1196/annals.1417.029

Schiff, N. D., Giacino, J. T., Kalmar, K., Victor, J. D., Baker, K., Gerber, M., Fritz, B., Eisenberg, B., O'Connor, J., Kobylarz, E. J., Farris, S., Machado, A., McCagg, C., Plum, F., Fins, J. J., & Rezai, A. R. (2007). Behavioural improvements with thalamic stimulation after severe traumatic brain injury. *Nature*, *448*(7153), 600-603. https://doi.org/10.1038/nature06041

Schmack, K., Bosc, M., Ott, T., Sturgill, J. F., & Kepecs, A. (2021). Striatal dopamine mediates hallucination-like perception in mice. *Science*, *372*(6537), eabf4740. https://doi.org/10.1126/science.abf4740

Sergent, C., Corazzol, M., Labouret, G., Stockart, F., Wexler, M., King, J.-R., Meyniel, F., & Pressnitzer, D. (2021). Bifurcation in brain dynamics reveals a signature of conscious processing independent of report. *Nature Communications*, *12*(1), 1149. https://doi.org/10.1038/s41467-021-21393-z

Seth, A. K., & Bayne, T. (2022). Theories of consciousness. *Nature Reviews Neuroscience*, *23*(7), 439-452. https://doi.org/10.1038/s41583-022-00587-4

Shepherd, G. M. G., & Yamawaki, N. (2021). Untangling the cortico-thalamo-cortical loop : Cellular pieces of a knotty circuit puzzle. *Nature Reviews Neuroscience*, *22*(7), 389-406. https://doi.org/10.1038/s41583-021-00459-3

Stockart, F., Msheik, R., Robin, A., Jurkovičová, L., Goueytes, D., Rouy, M., Mareček, R., Hoffmann, D., Mudrik, L., Roman, R., Brázdil, M., Minotti, L., Kahane, P., Pereira, M., & Faivre, N. (2024). *Cortical evidence accumulation for perceptual experience occurs irrespective of reports*. https://doi.org/10.1101/2024.03.20.585198

Tauste Campo, A., Vázquez, Y., Álvarez, M., Zainos, A., Rossi-Pool, R., Deco, G., & Romo, R. (2019). Feed-forward information and zero-lag synchronization in the sensory thalamocortical circuit are

modulated during stimulus perception. *Proceedings of the National Academy of Sciences*, *116*(15), 7513-7522. https://doi.org/10.1073/pnas.1819095116

Tsuchiya, N., Wilke, M., Frässle, S., & Lamme, V. A. F. (2015). No-Report Paradigms : Extracting the True Neural Correlates of Consciousness. *Trends in Cognitive Sciences*, *19*(12), 757-770. https://doi.org/10.1016/j.tics.2015.10.002

Twomey, D. M., Kelly, S. P., & O'Connell, R. G. (2016). Abstract and Effector-Selective Decision Signals Exhibit Qualitatively Distinct Dynamics before Delayed Perceptual Reports. *Journal of Neuroscience*, *36*(28), 7346-7352. https://doi.org/10.1523/JNEUROSCI.4162-15.2016

Vazquez, Y., Salinas, E., & Romo, R. (2013). Transformation of the neural code for tactile detection from thalamus to cortex. *Proceedings of the National Academy of Sciences*, *110*(28), E2635-E2644. https://doi.org/10.1073/pnas.1309728110

Vazquez, Y., Zainos, A., Alvarez, M., Salinas, E., & Romo, R. (2012). Neural coding and perceptual detection in the primate somatosensory thalamus. *Proceedings of the National Academy of Sciences*, *109*(37), 15006-15011. https://doi.org/10.1073/pnas.1212535109

Ward, L. M. (2013). The thalamus : Gateway to the mind: Gateway to the mind. *Wiley Interdisciplinary Reviews: Cognitive Science*, *4*(6), 609-622. https://doi.org/10.1002/wcs.1256

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wyart, V., & Tallon-Baudry, C. (2008). Neural Dissociation between Visual Awareness and Spatial Attention. *The Journal of Neuroscience*, *28*(10), 2667-2679. https://doi.org/10.1523/JNEUROSCI.4748-07.2008

Yang, M., Keller, D., Dobolyi, A., & Valtcheva, S. (2025). The lateral thalamus : A bridge between multisensory processing and naturalistic behaviors. *Trends in Neurosciences*, *48*(1), 33-46. https://doi.org/10.1016/j.tins.2024.11.005