Title: Mutation of *gabra1* is associated with hypermotility and abnormal expression of proteins critical for ion homeostasis and synaptic vesicle transport.

Authors: Nayeli G. Reyes-Nava[1], Isaiah Perez[1], Brian Grajeda[1,¶], Igor L. Estevao[1,¶], Cameron C. Ellis[1,¶], Sourav Roy[1], and Anita M. Quintana[1,*].

Affiliations:

[1] Department of Biological Sciences, Border Biomedical Research Center, The University of Texas at El Paso, El Paso, Texas, United States of America

¶These authors contributed equally

*Corresponding author

Email: amquintana8@utep.edu (AQ)

Abstract

Mutation of the *GABRA1* gene is associated with neurodevelopmental defects and epilepsy phenotypes. *GABRA1* encodes for the α1 subunit of the gamma-aminobutyric acid type A receptor (GABA$_A$R), which regulates the fast inhibitory impulses of the nervous system. Multiple model systems have previously been developed to understand the mechanism by which mutations in *GABRA1* cause disease, but these models have produced complex and incongruent data. Thus, additional model systems are required to validate and substantiate previously published results. We investigated the behavioral patterns associated with a non-sense mutation of the zebrafish *gabra1* (*sa43718* allele) gene. The *sa43718* allele has a 90% decrease in total *gabra1* mRNA expression, which is associated with light induced seizure-like behavior. Mutation of *gabra1* was accompanied by increased mRNA expression of *gabra4,* which encodes for the alpha-4 subunit of the GABA$_A$R. Despite increased expression at the RNA level, Gabra4 protein was not increased according to proteomics analysis. Thus, implying that RNA expression patterns of alpha sub-units may not accurately reflect the mechanism underlying seizure. Interestingly, proteomics analysis identified significant enrichment of genes that regulate proton transport, ion homeostasis, vesicle transport, and mitochondrial protein complexes. Collectively, our analysis validates that mutation of *gabra1* results in seizure like phenotypes and provides a blueprint of putative proteins which may mediate these phenotypes *in vivo*.

Introduction

The gamma-aminobutyric acid type A receptor (GABA$_A$R) is a multi-subunit ion channel that mediates inhibitory synapses of the nervous system. The GABA$_A$R can be composed of unique combinations of any of the following: 6 α subunits, 3 β subunits, 3 γ subunits, and one ε, δ, θ, or π subunits. The most common GABA$_A$R in mammals consists of 2 α subunits, 2 β subunits, and γ subunit. Each of the subunits is encoded by an independent gene, which are located on different chromosomes. Mutations in many of these subunits have been associated with seizure phenotypes in humans [1] and most recently we used whole exome sequencing to identify a putative heterozygous mutation in the *GABRA1* gene (c.875C>T) associated with seizure phenotypes [2]. *GABRA1* encodes for the α1 subunit of the GABA$_A$R and to date, over 30 different variants in the *GABRA1* gene have been reported and associated with neurological disorders and neurodevelopmental defects [2]–[11].

Because of its indicated role in neural development, several murine models have been developed to investigate the physiological and behavioral functions of *GABRA1* in mammals, however, the phenotypic outcomes are heterogeneous. For instance, deletion of *Gabra1* or the knock-in of the p.Ala332Asp allele cause behavioral phenotypes that range from tremors/absence-like seizures to the development of myoclonic seizures at postnatal stages [12]–[15]. Noteworthy, these phenotypes are generally strain, age, and sex specific, which makes interpretation of results difficult. Additional zebrafish models have been characterized including a germline mutant and a morpholino mediated transient knockdown. These two studies yielded contrasting results whereby hypermotility

was observed in a nonsense mutant of *gabra1* and hypomotility was observed after

transient knockdown [2], [16]. Thus, additional complementary models are warranted.

In 2016, the zebrafish mutation project was completed by the Wellcome Sanger

Institute, where over 40,000 mutant alleles, covering 60% of zebrafish protein-coding

genes, were generated [17]. However, only a small number of alleles have been

characterized or associated with a specific phenotype. An allele carrying a nonsense

mutation in the *gabra1* gene (sa43718) was generated and remains uncharacterized. We

hypothesized that a more comprehensive analysis of this allele coud discern the disparate

phenotypes observed across other zebrafish approaches. We therefore performed

behavioral and molecular analysis of the sa43718 allele.

Our results indicate that homozygous *gabra1$^{sa43718/sa43718}$* larvae show hyperactive

locomotion (seizure-like behavior) upon a light stimulus. These behavioral changes are

associated with a moderate increase in the expression of *gabra4* transcript, but no change

at the level of Gabra4 protein expression. However, differential expression of proteins

identified via proteomics approaches revealed abnormal expression of proteins that

regulate proton transport, sodium ion homeostasis across the plasma membrane, and the

mitochondrial inner membrane protein complex and respiratory chain.

Methods

Experimental model and animal husbandry

The *gabra1$^{sa43718/+}$* allele (N=10 female fish) was obtained from Sanger Institute through

the Zebrafish International Resource Center (ZIRC). Female carriers were outcrossed

with wildtype (AB) males to generate independent families of heterozygous male and female carriers. For all experiments, zebrafish larvae were obtained by mating adult heterozygous *gabra1*$^{sa43718/+}$ fish. Collected zebrafish embryos were maintained in E3 media (5mM NaCl, 0.17mM KCl, 0.33mM CaCl, 0.33mM MgSO4, 0.05% methylene blue, pH 7.4) at 28°C under 14:10 light: dark cycle. All experimental procedures were performed at 5 days post fertilization (DPF). All animals were maintained and used in accordance with the guidelines from the University of Texas at El Paso Institutional Animal Care and Use Committee (Animal Protocol Number 811869-5). Euthanasia and anesthesia were performed according with the American Veterinary Medical Association (AVMA) guidelines for the euthanasia, 2020 edition.

Genotyping

Genotyping of the *gabra1*$^{sa4371/sa4371}$ allele was performed by PCR amplification and restriction enzyme digest. Adult fin clips and larval tails were lysed in lysis buffer (30ul) (50mM NaOH) for 5 minutes at 95°C followed by 10 minutes at 4°C. DNA was then neutralized by addition of 500mM Tris-HCL solution (6ul) and used directly for PCR. The fragment of interest (237bp) was amplified by standard PCR at an annealing temperature of 60°C (forward: TTGTGACTCAAAGCCACGAG and reverse: TGAGACGAGAACCATCGTCA). PCR amplicon containing the *sa43718* allele region, was digested with XhoI (NebBiolabs) at 37°C according to manufacturer's guidelines. The mutation present in the *sa43718* inhibits cleavage by XhoI allowing differentiation of wildtype, heterozygous, and homozygous offspring.

Behavioral analysis and pentylenetetrazole (PTZ) treatment

Behavioral analysis was performed using the ZebraBox (ViewPoint Behavioral Technology, Montreal, Canada). Briefly, embryos were obtained from natural spawning and raised to 5 DPF. Larvae (5 DPF) were individually tracked for swim speed and total distance swam in a 96-well plate. The behavioral analysis consisted of 15 minutes divided into 5-minute intervals of dark/light/dark conditions. All larvae were acclimated to the chamber for 1 hour prior to data acquisition [2]. Data was collected each minute, for a total of 15 minutes and total distance traveled (mm) and speed (mm/s) was calculated as previously described [2]. Experiments were performed in biological duplicates using a minimum of N=30 larvae per trial. Treatment with 10mM pentylenetetrazole (PTZ) (Millipore-Sigma) was performed as previously described [2]. Genotyping of each larvae was performed after behavioral analysis.

Quantitative real time PCR (QPCR)

Total RNA was isolated from brain homogenates at 5 DPF from wildtype, heterozygous, and homozygous larvae with TRIzol reagent (Invitrogen) according to manufacturer's instructions. RNA was reverse transcribed using the Verso cDNA Synthesis Kit (ThermoFisher Scientific) and total RNA (500ng) was normalized across samples. Gene expression was measured in biological quadruplicates using a pool of at least N=5 larvae per biological replicate. For each individual biological replicate, technical replicates were performed as internal controls. The Applied Biosystems StepOne Plus machine with Applied Biosystems associated software was used for quantitative PCR (qPCR) analysis. Sybr green (ThermoFisher Scientific) based primer pairs were designed for each gene analyzed: *gabra1* (FWD: TCGGGAGTCCAGATTTTGCT, REV: AGAGCGTGTAACCGAAGTCA), g*abra2a* (FWD: GATGGCTACGACAACAGGCT, REV:

TGTCCATCGCTGTCGGAAAA), *gabra3* (FWD: GCTGAAGTTCGGGAGCTATG, REV:

GGAGCTGATGGTCTCTTTGC), *gabra4* (FWD: GACTGCGATGTACCCCACTT, REV:

ATCCAGGTCGGAGTCTGTTG), *gabra5* FWD: CATGACAACACCCAACAAGC, REV:

CAGGGCCTTTTGTCCATTTA), *gabra6a* (FWD: TCGCGTACCCATCTTTCTTC, REV:

CCCTGAGCTTTTCCAGAGTG), *gabra6b* (FWD: CGGAGGAGTGCTGAAGAAAC, REV:

GGGAAAAGGATGCGTGAGTA), *gabrb2* (FWD: CCCGACACCTATTTCCTCAA, REV:

TCTCGATCTCCAGTGTGCAG), *gabrg2* (FWD: ACACCCAATAGGATGCTTCG, REV:

AGCTGCGCTTCCACTTGTAT). The *rpl13a* gene was used as a reference gene for $2^{\Delta\Delta ct}$

quantification. Statistical analysis performed using a *t-test*.

Proteomic analysis

Protein isolation

For protein analysis, total protein was obtained from a pool of whole brain homogenates

(n=9) per genotype. Two biological replicates per genotype were collected. Brains were

excised and stored for protein isolation in 1X Cell Lysis Buffer (ThermoFisher Scientific)

with protease inhibitors cocktail (ThermoFisher Scientific). DNA was isolated from the tail

tissue for genotyping and protein was isolated from brain homogenates using manual

homogenization. Protein quantification was performed using Precision Red Protein Assay

Reagent (Cytoskeleton) according to manufacturer's instructions. Samples were then

sent to the Biomolecule Analysis and Omics Unit (BAOU) at The University of Texas El

Paso for sample processing and proteomic analysis.

Sample preparation for proteomics

Proteins were removed from 1X Cell Lysis Buffer by trichloroacetic acid (TCA) protein

precipitation. Fifty microliters of 100% TCA (Sigma/Millipore - cat# T6399-5G) was added

to 200 µL of sample in 1x Cell Lysis Buffer (Safe Seal microcentrifuge tubes, Sorenson

BioScience, cat. No. 12030) and incubated at 4°C for 10 minutes. Total protein ranged

from 2 – 5 µg per sample. The precipitated proteins were pelleted by centrifugation for

five minutes at 14,000 rcf and the supernatant was discarded. The protein pellet was

subsequently washed a total of three times with 200 µL LCMS grade acetone by

resuspension and pelleting, the acetone supernatant was discarded each time. Residual

acetone was evaporated with a heating block at 95°C for 2 minutes. Samples were stored

as a pellet at -80°C until processing for proteomic analysis. Stored pellets were

resuspended in 8M Urea prior and subjected to enzymatic tryptic digestion using the

PreOmics iST sample preparation kit (catalog no. iST 96x P.O.00027). Briefly,

denaturation, alkylation, and reduction were performed with 10 µL LYSE buffer per 1 µg

of protein and heated at 80 °C for 20 min while being mixed every 5-minutes. Remaining

droplets on the cap were given a brief centrifugation (RT; 300 rcf; 10 sec) and samples

were sonicated ten times at 30-second on/off intervals for a total of 5 minutes. Fifty

microliters of resuspended DIGEST solution were added to the samples for protein

digestion. Sample-filled microtubes were gently vortexed, centrifuged and kept for 90

minutes at 37°C in a heating block. Throughout the 90-minute incubation period, samples

were gently mixed every 10 minutes. One hundred microliters of STOP solution were

added and mixed by pipetting ten times. Samples were then transferred to the cartridge

and centrifuged for 1 minute at 3,800 rcf, followed by subsequent 200 µL washes with

WASH 1 and WASH 2 solutions according to the manufacturer's instructions. Peptides

were eluted in two cycles of 100 μL of ELUTE solution and dried completely in a vacuum

evaporator (Savant; Thermo Fisher Scientific) for 90 minutes at 45 °C and 100-mTorr and

stored at -80°C until LC-MS/MS acquisition.

Liquid chromatography–tandem mass spectrometry (LC-MS/MS)

Peptides were resuspended in 4% acetonitrile (ACN) with 0.1% formic acid (FA) at a

concentration of 1μg/μL. Peptides were separated by a Dionex Ultimate 3000 UHPLC

system (Thermo Scientific) tandem with a Q-Exactive Plus Hybrid Quadrupole-Orbitrap

Mass Spectrometer (Thermo Scientific) with Xcalibur software (v. 3.0.63) for data

acquisition in positive mode. Peptides were separated on a C18 Acclaim PepMap nanoLC

column (75 μm x 50 cm nanoViper, PN 164942, Thermo Fisher Scientific) equilibrated

with 4% solvent B (99.9% acetonitrile, 0.1% formic acid) and 96% solvent A (99.9% H2O,

0.1% formic acid) kept at 55°C throughout the entire acquisition. One microliter of

peptides was loaded onto the column for 15-minutes at a flow rate of 0.5 μL/min and

eluted with a multi-step gradient. The flow rate was reduced to 0.3μL/min over the course

of 15 min, and solvent B was set to 20% over 100-minutes, then increased to 32% and

maintained for 20-minutes before increasing to 95% over 1-minute. The column was

washed with 95% solvent B at a flow rate of 0.4 μL/min for 4-minutes to remove/clean any

remaining peptides. The column was then re-equilibrated with 4% solvent B at 0.5μL/min

until a 180-minutes total runtime. Blank injections were added after each biological

replicate, using a 60-minute two sawtooth gradient from 4-95% solvent B, and column re-

equilibration at 4% solvent B. The mass spectrometer was set to top10 data-dependent

acquisition (DDA), with a scan range of 375 to 1500 m/z and a full MS resolution of

70,000; AGC target 3e6. Ions were fragmented with NCE at 27 and collected with an AGC

target of 1e5 at 17,500 resolution, 2 m/z isolation window, maximum IT of 60 ms. Charged

exclusion ions were unassigned, 1, 6–8, and >8 charges.

Bioinformatics data analysis

After proteomics analysis, Proteome Discover (PD) 2.5.0.400 (Fisher Scientific) was

utilized to identify the proteins from each peptide mixture. The database for Danio rerio

was downloaded from UniProtKB; http://www.uniprot.org/ on 21 October 2021 with a

database 61,623 sequences. A contaminant dataset was run in parallel composed of

trypsin autolysis fragments, keratins, standards found in CRAPome repository and in-

house contaminants. PD analysis parameters are as follows: false-discovery rate (FDR)

of 1%, HCD MS/MS, fully tryptic peptides only, up to 2 missed cleavages, parent-ion mass

of 10 ppm (monoisotopic); fragment mass tolerance of 0.6 Da (in Sequest) and 0.02 Da

(in PD 2.1.1.21) (monoisotopic). Two-high confidence peptides per protein were applied

for identifications. PD dataset was processed through Scaffold Q+S 5.0.1. Scaffold

(Proteome Software, Inc., Portland, OR 97219, USA) was used to probabilistically

validate protein identifications derived from MS/MS sequencing results using the

X!Tandem and Protein Prophet. Data was transferred to Scaffold LFQ (Proteome

Software, Portland, Oregon, USA) which was used to validate and statistically compare

protein identifications derived from MS/MS search results. A protein threshold of 95%,

peptide threshold of 95%, and a minimum number of 2 peptides were used for protein

validation. Normalized weighted spectral counts were used when comparing the samples.

To ascertain p-values, Fisher's Exact was run with a control FDR level q * .05 with

standard Benjamini-Hochberg correction.

Results

Characterization of the sa43718 allele

The sa43718 allele results in a single base pair substitution in exon 6, which is predicted to result in a premature stop codon (Fig 1A). Exon 6 encodes part of the extracellular domain. We used Sanger sequencing to validate the single base pair substitution shown in Fig 1B (Sibling) &B'(sa43718). Based on the sequence change present in the sa43718 allele, we developed a restriction enzyme based genotyping strategy. The sa43718 allele abolishes an XhoI cleavage site and therefore cannot be cleaved by XhoI in a restriction digest. We performed PCR amplification using primers that flank the substituted nucleotide and used restriction digest to identify wildtype, heterozygous, and homozygous larvae. DNA from wildtype siblings was completely digested as predicted (Fig 1C, lane 2), while DNA from homozygous larvae was completely undigested, and DNA from heterozygous carriers was partially digested (Fig 1C, lane 4 and 3, respectively).

Fig 1. The sa43718 allele carries a single non-sense mutation in the *gabra1* gene. (A) Schematic representation of the *gabra*^{sa43718} allele. Homozygous individuals of the sa43718 allele (*gabra1*$^{sa43718/sa43718}$) carry a single base pair substitution (C>T) in the *gabra1* gene (ENSDART0000010000), predicted to result in a premature stop. Schematic of exon structure of the *gabra1* gene, not to scale. The sa43718 site is indicated with a red asterisk within exon 6, which encodes the extracellular domain of Gabra1. (B-B') Chromatograph showing confirmation of the sa43718 allele through Sanger sequencing of *gabra1*$^{sa43718/+}$ offspring. (B) Wildtype sibling sequence. (B'): *gabra1*$^{sa43718/sa43718}$ sequence. (C) Restriction enzyme digest gel showing digestion of PCR fragments of genomic *gabra1.* The sa43718 allele abolishes the XhoI recognition site. Therefore, digestion of homozygous (sa43718/sa43718) sample results in an uncut fragment, whereas digestion of wildtype (sibling) sample results in a fully digested fragment. LDR: 1kb plus ladder. (D) The offspring of *gabra1*$^{sa43718/+}$ fish was genotyped by tail-clipping at 3 days post fertilization (DPF). Larvae were raised separately according to their genotypes (sibling n=11,

sa43718/+ n=13, sa43718/sa43718 n=14), and their survival was monitored until day 22.
Premature death of homozygous carriers of the sa43718 allele (sa43718/sa43718*)* was observed
starting at day 7 post fertilization, when compared to the wildtype (sibling) and heterozygous
(sa43718*/+)* siblings*.*

Homozygous mutation in zebrafish *gabra1* has been associated with early lethality.

Consequently, we performed a survival assay and monitored the survival for 22 DPF. We

observed a 50% survival rate in wildtype and heterozygous siblings, but all homozygous

carriers of the sa43718 allele died by 22 DPF (Fig 1D). The remaining heterozygous

individuals survived into adulthood.

258   Expression of GABA$_A$ receptor subunits in the sa43718 allele.

Based on our previous studies, we hypothesized that the expression of other GABA$_A$R

receptor subunits were disrupted in the sa43718 allele. We first measured the expression

of β2 and γ2 subunits, as these are present in the most common form of the GABA$_A$R

receptor, alongside α1. We validated that the sa43718 allele reduced the expression of

gabra1 (Fig 2A), but that decrease was not associated with decreased expression of β2

and γ2 transcripts (Fig 2A). We therefore hypothesized that these subunits were in

complex with other α subunits to compensate for the loss of α1. We measured the

expression of α2-6 (including 6a and 6b due to gene duplication). We observed a mild

increase in *gabra4* (α4) gene expression, which was statistically significant (Fig 2B).

However, we did not detect a significant difference in any other α subunit (Fig 2B).

Fig 2. Non-sense mutation of *gabra1* results in decreased expression of *gabra1*. (A)
Quantitative real time PCR (qPCR) analyzing the relative mRNA expression of major GABA$_A$R
subunits, α1 (*gabra1),* β2 *(gabrb2)* and, γ2 *(gabrg2)* at 5 days post fertilization (DPF). n≥5 larvae

per group with five biological replicates. Error bars represent the standard error of the mean of

biological replicates. *p=1.00511E-07. (B) Relative expression of genes encoding for alpha type

subunits, *gabra2a* (α2), *gabra3* (α3), *gabra4* (α4), *gabra5* (α5), *gabra6a* (α6a), and *gabra6b* (α6b),

was analyzed in wildtype siblings (sibling) and homozygous (sa43718/sa43718) mutants. n≥5

larvae per group in four independent biological replicates. Error bars represent the standard error

of the mean. *p=0.002884.

279 Mutation of *gabra1* results in seizure-like behavior

We have previously reported hypoactivity at 5 DPF upon knockdown of *gabra1* [2]. In

contrast, seizure-like behavior at juvenile stages has been reported in a zebrafish model

of *gabra1* loss of function [16]. To begin to rectify these differences, we monitored

swimming behavior after a light stimulus, which has been shown to induce seizure like

behavior in a previous allele of *gabra1* mutation. We measured baseline activity for a

period of 5 minutes in the dark after a 1-hour acclimation period. During this time, we did

not observe any differences in baseline swimming patterns according to total distance

swam (mm) or swim speed (mm/s) in the sa43718 allele (Fig 3A&A'). However, after light

stimulus, we detected a marked increase in swim speed in the sa43718 allele (Fig 3A&A').

Swim patterns were increased in mutant larvae during the entire light stimulus (Fig

3C&C'). Importantly, increased speed (velocity) was associated with rapid circling

behavior, hyperactivity burst, and whirlpool behavior (Fig 3B) indicative of a seizure-like

behavior as defined by the zebrafish behavioral glossary [18]. Because we did not

observe behavioral changes in heterozygous carriers in comparison to wildtype siblings,

all data shown here are from sibling wildtype or homozygous carriers of the sa43718

allele.

Fig 3. gabra1$^{sa43718/sa43718}$ larvae undergo seizure-like behavior upon light stimuli. (A & A') Behavioral analysis of sa43718 allele and wildtype siblings at 5 days post fertilization. (A) Swim speed (mm/s) over 15 minutes (900 seconds) in dark-light-dark transitions [5 minutes (0-300 seconds) in the dark, 5 minutes (300-600 seconds) in the light, and 5 minutes (600-900 seconds) in the dark) was analyzed using the Zebrabox technology. Data was collected every minute (60 seconds). *gabra1$^{sa43718/sa43718}$* larvae undergo hyperlocomotion upon light stimuli when compared to their wildtype siblings (Sibling). *p<0.05. (A') Total distance swam over 15 minutes in dark-light-dark transitions. Homozygous carriers (sa43718/sa43718) larvae showed increased distance swam across the light phase, showing a statistically significant increase at seconds 480-540 (minute 9), when compared to their wildtype counterparts (sibling). *gabra1$^{sa43718/sa43718}$* larvae then show decreased distance swam under dark conditions (600-660s) when compared to their wildtype siblings (sibling) *p<0.05. (B) Representative images of swimming track of a single larvae, representing each genotype, generated from the Viewpoint Zebralab Tracking software in dark (180-240s), light (480-540s), and dark (600-660s) conditions. Green lines indicate movements between 4-8mm/s and red lines indicate burst movements (>8mm/s). (C) Average total speed representing overall behavioral responses in dark-light-dark conditions of *gabra1* wildtype (sibling) and homozygous (sa43718/sa43718) larvae. Homozygous carriers (sa43718/sa43718) showed increased average swim speed in light conditions when compared to their wildtype (sibling) counterparts. (C') Average total distance swam representing overall behavioral responses in dark-light-dark conditions of *gabra1* wildtype (sibling) and homozygous (sa43718/sa43718) larvae. No statistically significant changes were observed in homozygous larvae (sa43718/sa43718) when compared to their wildtype siblings for average distance measurements. All behavioral analysis was performed using a minimum of 30 larvae per biological replicate, over three independent biological replicates. Representative analysis of a single biological replicate is shown here. Sibling (n=13) and *gabra1$^{sa43718/sa43718}$* (n=26).

Gabra1 sa43718 response to PTZ

We have previously demonstrated that reduced *gabra1* expression does not prohibit receptor responses to pentylenetetrazole (PTZ), a GABA$_A$R antagonist [2]. We also observed increased expression of *gabra4* and therefore, we sought to understand if the

sa43718 allele would respond to PTZ, which may indicate the presence of a functional

receptor. As shown in Fig 4A, treatment with PTZ induces increased swimming distance

and speed in wildtype larvae. Within the first five minutes post treatment, in dark

conditions, wildtype larvae exhibit an average 2-fold change in total distance swam and

a 1.54-fold change in swim speed. However, the sa43718 allele had a slightly reduced

response to PTZ in the dark accounting for a 1.7-fold change in total distance swam and

a 1.2-fold change in swim speed in dark conditions (minutes 1-5) (Fig 4C-D). We

continued to monitor the speed and distance swam after light stimulus in PTZ treated

larvae. After light stimulus, wildtype sibling larvae showed a 1.5-fold change in average

swim speed and a 2.5-fold change in average total distance swam (Fig 4C-D). Whereby,

the sa43718 allele exhibited a 1.4-fold change in average swim speed and 1.7-fold

change in average distance swam in response to PTZ treatment (Fig 4C-D). The reduced

response to PTZ treatment by the sa43718 allele was statistically significant in dark

conditions for speed ($p=0.03$) and distance ($p=0.002$).

Fig 4. Locomotion response to pentylenetetrazole (PTZ) in the sa43718 allele. (A-B).
Behavioral responses of the sa43718 allele at 5 days post fertilization (DPF) were analyzed using
the Zebrabox. (A) Comparison of wildtype sibling (Sibling) larvae response to treatment with 10uM
PTZ (10uM PTZ) or no treatment (NO PTZ) at 5 days post fertilization. NO PTZ n=18, 10uM PTZ
n=16. *p<0.05 (B) Comparison of homozygous carriers (sa43718/sa43718) response to treatment
with 10uM PTZ (10uM PTZ) or no treatment (NO PTZ) at 5 days post fertilization. NO PTZ n=13,
10uM PTZ n=14. *p<0.05. Swim speed (mm/s) was analyzed using the distance swam and
swimming duration data generated by ViewPoint software. Larvae were monitored for 5 minutes
in the dark (0-300 s), 5 minutes in the light (300-600 s), and 5 minutes in the dark (600-900 s).
Data was collected every minute (60 s). Error bars represent the standard error of the mean. (C)
Average fold change (total distance) response to treatment with PTZ in dark (0-300s) and light
(300-600s) conditions by wildtype (sibling) and sa43718/sa43718 mutants. *p=0.002. (D) Average
fold change (swim speed) response to treatment with PTZ in dark (0-300s) and light (300-600s)

conditions by wildtype (sibling) and sa43718/sa43718 mutants. *p=0.03. Error bars represent
standard deviation in C-D.

Proteomic analysis identifies differentially expressed proteins (DEPs) in the sa43718 allele.

To begin to understand the underlying molecular mechanisms by which seizure

phenotypes occurred, we performed proteomics analysis. We isolated protein from whole

brain homogenates isolated from wildtype and homozygous larvae. A total of 3500

proteins were identified in the wildtype sibling brain homogenate, 3511 were identified in

heterozygous animals, and 3552 were identified in the homozygous carriers of the

sa43718 allele. A complete list of identified proteins is in S1 file. From these proteins, we

identified a total of 173 DEPs in homozygous mutants relative to wildtype clutch mates.

Of these 81 were up-regulated and 93 were down-regulated. These DEPs are

summarized in the volcano plot shown inf Fig 5A. A complete list of DEPs can be found

in file S2. DEPs of interest include Synaptotagmin, Complexin 4a, Dynactin subunit 1,

Dynamin GTPase, Secretogranin II, and a Vat1 homolog. Each of these proteins has a

function in synaptic fusion, transport, exocytosis, endocytosis, or direct interactions with

small synaptic vesicles. Interestingly, we also noted a significant number of DEPs

involved in microtubule growth, microtubule binding, vesicle recycling, retrograde

transport along microtubules, or microtubule folding. These include microtubule

associated protein, RP/EB family member 3b, Dynactin subunit 1, Dynamin GTPase, and

Tubulin-specific chaperone A.

We further annotated DEPs by biological process (Fig 5B), molecular function (Fig

5C), and cellular process (Fig 5D). According to biological process, DEPs were clustered

in Gene Ontology (GO) groups associated with proton transport including sodium and

potassium export across the plasma membrane. Some examples of DEPs in this category

include Slc82b, Atp1a1b, Atp1b3a, and Atp1a1a.3. All these DEPs are important for

sodium and potassium homeostasis consistent with GO analysis. GO analysis by

molecular function also indicated molecular carrier activity and active transport

mechanisms, consistent with the GO analysis according to biological process. GO

analysis by cellular process indicated DEPs in the respiratory chain, mitochondrial protein

containing complexes, the cytoskeleton, proton transport, and exocytosis. These data

collectively indicate mitochondrial transport, synaptic vesicle regulation, and proton

homeostasis are dysregulated after mutation of *Gabra1*.

Fig 5. Proteomic analysis of sa43718 allele carriers. (A) Volcano plot of differentially

expressed proteins between wildtype siblings and homozygous (sa43718/sa43718)

larvae. Volcano plot show –log10 P-values from the normalized proteomics data exported

from Scaffold LFQ versus log2foldchange (FC) across each contrast. Thresholds set were

p-value ≤ 0.05 and Log2FC ≤ -1.5 or ≥ 1.5. (B-D) Gene Ontology terms enriched by

proteins with significant expression (p-value <0.05) between wildtype sibling and sa43718

homozygous mutant larvae at 5 days post fertilization. Terms are separated by (B)

Biological Process, (C) Molecular Function, (D) Cellular Process.

Discussion

The GABA$_A$R has previously been associated with seizure and epileptic

phenotypes [1], [9], [15]. In support of this notion, we reported a heterozygous missense

mutation in humans that is associated with a seizure phenotyes [2]. Multiple model

systems have been developed to study early *GABRA1* function. These include mouse

and zebrafish models. Survival of homozygous knockouts in mice and zebrafish is

compromised [12], [16]. Due to external fertilization and early juvenile survival of

homozygous mutants, zebrafish provide a window of opportunity to analyze

developmental and juvenile onset of seizures [19]–[24]. However, in previous studies

germline nonsense mutation of zebrafish *gabra1* and morpholino mediated knockdown of

gabra1 demonstrated different behavioral phenotypes after a light stimulus [2], [16]. Here

we characterized the phenotypes of a novel allele, known as the sa43718 allele. The

sa43718 allele is a single base pair change in exon 6, which results in a premature stop

codon and leads to a greater than 90% decrease in *gabra1* expression, indicating the

possibility of nonsense mediated decay of the mutated transcript. The sa43718 allele

reduces viability and homozygous mutants die by 22 DPF. This is consistent with the

viability of other zebrafish [16] and mouse models [13].

We observed seizure like (hypermotility, whirlpool behavior) behavior upon light

stimuli in the sa43718 allele. These data are supported by previously published works,

which established seizure like behavior at the juvenile stage in an additional zebrafish

germline mutant [16]. Similar results have also been documented in mouse models of

Gabra1 mutation, although these data are sex and strain specific [12]–[14]. In our

experiments, the zebrafish larvae have not yet determined sex and therefore, sex is not

a variable considered in our experiments. Hyperactivity and seizure like behavior in the

sa43718 allele contrasts with our previous study using morpholino mediated knockdown

[2]. There are several reasons for these differences. For example, we validated an

approximate 50% knockdown of *gabra1* in morphants, which would most likely reflect a

haploinsufficency or dosage effect. In addition, *gabra1* morphants demonstrated with

decreased expression of other GABA$_A$ receptor subunits. We did not observe statistically

significant differences in other subunits, except for the *gabra4* subunit. However, this was

only at the mRNA level, as proteomics did not reveal differences in the Gabra4 protein

level in the sa43718 allele.

Given the normal expression of other GABA$_A$R subunits, it is not surprising that

sa43718 larvae responded to PTZ treatment. Mutant larvae were capable of a response

to 10uM PTZ, which has been shown to induce seizures in zebrafish [25], [26]. These

data are consistent with our previous work, which demonstrated that *gabra1* knockdown

did not affect response to PTZ [2]. Additionally, Samarut and colleagues demonstrated

that the overall brain morphology of *gabra1* mutant brains was normal. Despite an

overwhelming normal morphology, Samarut and colleagues did show 460 differentially

expressed genes in *gabra1* mutant brains. Of these genes, 3 GABA$_A$R subunits, in

addition to *gabra1*, were down regulated [16]. We observed increased expression of

gabra4 here and in a previous study we found *gabra6* to be up regulated at the

transcriptional level after morpholino mediated knockdown [2]. These data are partially

supported by murine studies that show abnormal expression of receptor subunits.

Although data from murine studies are mutant, strain, and gender specific [27], [28]. We

performed proteomics analysis to determine if these changes translate to differential

expression of proteins. We did not identify a single GABA$_A$R subunit differentially

expressed in the sa43718 allele at the protein level. These data are consistent with the

ability of *gabra1* mutants to respond to PTZ, indicating the presence of a receptor.

Our proteomics analysis identified proteins with functions in synaptic vesicle

transport, microtubule organization/growth, mitochondrial function, and regulation of

potassium and sodium homeostasis as differentially expressed. A subset of these data

are consistent with data from Samarut and colleagues, who noted decreased expression

of kinesin motor proteins [16]. However, the functional significance of differential

expression of transcripts is unclear because we did not validate up-regulation of Gabra4

protein despite increased transcript. Thus, there are some differences from our protein

data relative to previous results. For example, we did not identify statistically significant

changes in neuroligins 2 and 4, nor other GABA$_A$ subunits. We cannot rule out differential

expression of these proteins, however, in our analysis we did not find significant changes

in protein expression. Of note, we identified abnormal expression of synaptotagmin,

secretogranin II, and synapsin IIb; all of which are important for synaptic vesicle release

at the presynaptic membrane. Synapsin I was also downregulated and has been

associated with epileptic phenotypes [29]–[32]. Furthermore, we identified

downregulation of proteins implicated in mitochondrial complex I deficiency, including

NADH:ubiquinone oxidoreductase core subunit S2 and NADH:ubiquinone

oxidoreductase core subunit A10. Remarkably, mutations in the genes encoding these

proteins are associated with Leigh syndrome, one of the most common mitochondrial

diseases, in which epilepsy is a primary phenotype [33], [34]. Interestingly, mutation of

GABRA1 in humans have been prematurely diagnosed as a primary mitochondrial

disorder prior to exome sequencing [23]. Interestingly, none of the DEPs discovered

inhibited the response to PTZ, a GABA$_A$R agonist, suggesting that a residual receptor is

present. Nonetheless, our proteomics data implicates neurotransmitter release, vesicle

transport, microtubule structure, proton homeostasis, and regulation of mitochondrial
proteins as putative underlying mechanisms resulting in seizure-like phenotypes after
mutation of *gabra1*.

Here we have characterized an additional germline mutant of the zebrafish *gabra1*
gene. We observed early lethality, seizure-like hypermotility after light stimuli, and
abnormal expression of proteins involved in vesicle transport, synaptic vesicle release,
exocytosis, and endocytosis. Our data support the hypermotility and expression patterns
previously described in a secondary zebrafish germline mutant [16]. Collectively, these
data suggest that models derived from germline mutation of *gabra1* rather than transient
knockdown are more physiologically relevant to human disease.

Acknowledgements

The authors would like to thank the members of the Quintana Lab, past and present, for
discussions, animal care, genotyping, and general laboratory maintenance. A special
thank you to Pamela Rodriguez and Carla Canales for their efforts in the laboratory and
their explicit help with genotyping embryos related to this manuscript. The work provided
herein could not be completed without the support of The University of Texas El Paso
core facilities and Border Biomedical Research Center (BBRC). We graciously thank Dr.
Robert Kirken, Dr. Igor C. Almeida, and Dr. Renato Aguilera for their directorships of the
BBRC and core facilities.

References

1. Macdonald RL, Gallagher MJ, Feng HJ, Kang J. GABAA receptor epilepsy mutations. Biochem Pharmacol. 2004 Oct;68(8):1497–506.

2. Reyes-Nava NG, Yu HC, Coughlin CR, Shaikh TH, Quintana AM. Abnormal expression of GABAA receptor subunits and hypomotility upon loss of gabra1 in zebrafish. Biol Open [Internet]. 2020 Apr 15 [cited 2020 Apr 14];9(4). Available from: https://bio.biologists.org/content/9/4/bio051367

3. Steudle F, Rehman S, Bampali K, Simeone X, Rona Z, Hauser E, et al. A novel de novo variant of GABRA1 causes increased sensitivity for GABA in vitro. Sci Rep. 2020 Feb 11;10(1):2379.

4. Choi YJ, Lee SY, Yang KS, Park JY, Yoon SZ, Yoon SM. Polymorphism rs4263535 in GABRA1 intron 4 was related to deeper sedation by intravenous midazolam. J Int Med Res. 2015 Oct;43(5):686–98.

5. Cossette P, Liu L, Brisebois K, Dong H, Lortie A, Vanasse M, et al. Mutation of *GABRA1* in an autosomal dominant form of juvenile myoclonic epilepsy. Nat Genet. 2002 Jun;31(2):184–9.

6. Farnaes L, Nahas SA, Chowdhury S, Nelson J, Batalov S, Dimmock DM, et al. Rapid whole-genome sequencing identifies a novelGABRA1variant associated with West syndrome. Cold Spring Harb Mol Case Stud. 2017 Sep;3(5).

7. Kodera H, Ohba C, Kato M, Maeda T, Araki K, Tajima D, et al. De novo GABRA1 mutations in Ohtahara and West syndromes. Epilepsia. 2016 Apr;57(4):566–73.

8. Epi4K Consortium, Epilepsy Phenome/Genome Project, Allen AS, Berkovic SF,

Cossette P, Delanty N, et al. De novo mutations in epileptic encephalopathies.

Nature. 2013 Sep 12;501(7466):217–21.

9. Hirose S. Chapter 3 - Mutant GABAA receptor subunits in genetic (idiopathic)

epilepsy. In: Steinlein OK, editor. Progress in Brain Research [Internet]. Elsevier;

2014 [cited 2018 Jul 30]. p. 55–85. (Genetics of Epilepsy; vol. 213). Available from:

http://www.sciencedirect.com/science/article/pii/B978044463326200003X

10. Johannesen K, Marini C, Pfeffer S, Møller RS, Dorn T, Niturad CE, et al. Phenotypic

spectrum of GABRA1: From generalized epilepsies to severe epileptic

encephalopathies. Neurology. 2016 13;87(11):1140–51.

11. Maillard PY, Baer S, Schaefer É, Desnous B, Villeneuve N, Lépine A, et al. Molecular

and clinical descriptions of patients with GABAA receptor gene variants (GABRA1,

GABRB2, GABRB3, GABRG2): A cohort study, review of literature, and genotype–

phenotype correlation. Epilepsia. 2022;63(10):2519–33.

12. Ye GL, Baker KB, Mason SM, Zhang W, Kirkpatrick L, Lanthorn TH, et al. GABAA

Receptor α1 Subunit (Gabra1) Knockout Mice: Review and New Results. In: Kalueff

AV, Bergner CL, editors. Transgenic and Mutant Tools to Model Brain Disorders

[Internet]. Totowa, NJ: Humana Press; 2010 [cited 2020 Apr 9]. p. 65–90.

(Neuromethods). Available from: https://doi.org/10.1007/978-1-60761-474-6_4

13. Arain FM, Boyd KL, Gallagher MJ. Decreased viability and absence-like epilepsy in

mice lacking or deficient in the GABAA receptor α1 subunit. Epilepsia. 2012

Aug;53(8):e161-165.

14. Arain F, Zhou C, Ding L, Zaidi S, Gallagher MJ. The Developmental Evolution of the
Seizure Phenotype and Cortical Inhibition in Mouse Models of Juvenile Myoclonic
Epilepsy. Neurobiol Dis. 2015 Oct;82:164–75.

15. Maljevic S, Møller RS, Reid CA, Pérez-Palma E, Lal D, May P, et al. Spectrum of
GABAA receptor variants in epilepsy: Curr Opin Neurol. 2019 Apr;32(2):183–90.

16. Samarut É, Swaminathan A, Riché R, Liao M, Hassan-Abdi R, Renault S, et al. γ-
Aminobutyric acid receptor alpha 1 subunit loss of function causes genetic
generalized epilepsy by impairing inhibitory network neurodevelopment. Epilepsia.
2018 Nov 1;59(11):2061–74.

17. ZMP | Busch Lab [Internet]. [cited 2022 Oct 24]. Available from:
https://zmp.buschlab.org/

18. Kalueff AV, Gebhardt M, Stewart AM, Cachat JM, Brimmer M, Chawla JS, et al.
Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. Zebrafish.
2013 Mar;10(1):70–86.

19. Canzian J, Müller TE, Franscescon F, Michelotti P, Fontana BD, Costa FV, et al.
Modeling psychiatric comorbid symptoms of epileptic seizures in zebrafish. J
Psychiatr Res. 2019 Dec 1;119:14–22.

20. Grone BP, Baraban SC. Animal models in epilepsy research: legacies and new
directions. Nat Neurosci. 2015 Mar;18(3):339–43.

21. Yaksi E, Jamali A, Diaz Verdugo C, Jurisch-Yaksi N. Past, present and future of
zebrafish in epilepsy research. FEBS J. 2021;288(24):7243–55.

22. Stewart AM, Desmond D, Kyzar E, Gaikwad S, Roth A, Riehl R, et al. Perspectives of zebrafish models of epilepsy: what, how and where next? Brain Res Bull. 2012 Feb 10;87(2–3):135–43.

23. Reyes-Nava NG. Zebrafish: A Comprehensive Model to Understand the Mechanisms Underlying Neurodevelopmental Disorders. In: Costa, Andres., Villalba, Eugenio., editors. Horizons in Neuroscience Research. Nova Biomedical; 2018.

24. Gawel K, Langlois M, Martins T, van der Ent W, Tiraboschi E, Jacmin M, et al. Seizing the moment: Zebrafish epilepsy models. Neurosci Biobehav Rev. 2020 Sep 1;116:1–20.

25. Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. Neuroscience. 2005 Jan 1;131(3):759–68.

26. Afrikanova T, Serruys ASK, Buenafe OEM, Clinckers R, Smolders I, Witte PAM de, et al. Validation of the Zebrafish Pentylenetetrazol Seizure Model: Locomotor versus Electrographic Responses to Antiepileptic Drugs. PLOS ONE. 2013 Jan 14;8(1):e54166.

27. Zhou C, Huang Z, Ding L, Deel ME, Arain FM, Murray CR, et al. Altered cortical GABAA receptor composition, physiology, and endocytosis in a mouse model of a human genetic absence epilepsy syndrome. J Biol Chem. 2013 Jul 19;288(29):21458–72.

28. Zhou C, Ding L, Deel ME, Ferrick EA, Emeson RB, Gallagher MJ. Altered Intrathalamic GABAA Neurotransmission in a Mouse Model of a Human Genetic Absence Epilepsy Syndrome. Neurobiol Dis. 2015 Jan;73:407–17.

29. Fassio A, Raimondi A, Lignani G, Benfenati F, Baldelli P. Synapsins: From synapse
to network hyperexcitability and epilepsy. Semin Cell Dev Biol. 2011 Jun
1;22(4):408–15.

30. Garcia CC. Identification of a mutation in synapsin I, a synaptic vesicle protein, in a
family with epilepsy. J Med Genet. 2004 Mar 1;41(3):183–6.

31. Giannandrea M, Guarnieri FC, Gehring NH, Monzani E, Benfenati F, Kulozik AE, et
al. Nonsense-Mediated mRNA Decay and Loss-of-Function of the Protein Underlie
the X-Linked Epilepsy Associated with the W356× Mutation in Synapsin I. PLOS
ONE. 2013 Jun 20;8(6):e67724.

32. Barbieri R, Contestabile A, Ciardo MG, Forte N, Marte A, Baldelli P, et al. Synapsin
I and Synapsin II regulate neurogenesis in the dentate gyrus of adult mice.
Oncotarget. 2018 Apr 10;9(27):18760–74.

33. Lopriore P, Gomes F, Montano V, Siciliano G, Mancuso M. Mitochondrial Epilepsy,
a Challenge for Neurologists. Int J Mol Sci. 2022 Oct 30;23(21):13216.

34. Schubert Baldo M, Vilarinho L. Molecular basis of Leigh syndrome: a current look.
Orphanet J Rare Dis. 2020 Jan 29;15(1):31.

Supporting information

594 **S1 File. Complete list of identified proteins.**

595 **S2 File. Complete list of differentially expressed proteins.**

Funding

Partial funding for this project was provided by K01NS099153 to AMQ, R03DE029517 to AMQ, NIMHD grant No 5U54MD007592 to University of Texas El Paso, RISE program grant No R25GM069621-11 from NIGMS, and NIGMS linked awards RL5GM118969, TL4GM118971, and UL1GM118970 to the University of Texas El Paso. NGR was partially supported by Keelung-Hong Fellowship. We thank the Biomolecule Analysis and Omics Unit (BAOU) at BBRC/UTEP, for the full access to the nanoUHPLC-ESI-Q Exactive Plus orbitrap MS system used in this study. AMQ was provided pilot grant funds that partially funded this award as a component of the 5U54MD007592 award. The content is solely the responsibility of the authors and does not represent the official views of the funding agencies.

Author contributions

NGR performed experiments, provided experimental conceptualization, wrote aspects of the manuscript, performed zebrafish care and maintenance, supervised trainees that provided input/data analysis or acquisition. AMQ performed data analysis, wrote portions of the manuscript, obtained funding, provided mentorship to NGR, and performed data analysis. IP performed survival analysis and produced graphs related to survival. BG, ILE, and CCE performed mass spectrometry, protein isolation, and data analysis for proteomics. SR supervised and provided additional correction for BG data analysis.

Data availability

All data and reagents are available upon request from the corresponding author.

Competing interests

Authors declare no competing or financial interests.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5