Conformational Stability of Peroxidase from latex of *Artocarpus lakoocha*: Influence of pH,

Chaotropes, and Temperature

Kirti Shila Sonkar[a,c]*, Manendra Pachauri[b,d], Amit Kumar[b,e], Medicherla V.

Jagannadham[a]

[a]Molecular Biology Unit, Institute of Medical Sciences, Banaras Hindu University, Varanasi-

221005, India.

[b]Department of Biochemistry, University of Delhi South Campus, New Delhi- 110021, India

[c]Present address: Molecular Pathology Group, International Centre for Genetic Engineering and

Biotechnology (ICGEB), 34149 Trieste, Italy.

[d]Present address: Molecular Medicine Group, International Centre for Genetic Engineering and

Biotechnology (ICGEB), 34149 Trieste, Italy.

[e]Present address: Department of Biochemistry, Jacobs School of Medicine and Medical Sciences

The State University of New York- Buffalo, New York 14203, USA.

*Correspondence should be addressed to k.s.sonkar@gmail.com

Correspondence:

Kirti Shila Sonkar

Molecular Biology Unit,

Institute of Medical Sciences, Banaras Hindu University,

Varanasi- 221005, India.

Mobile: +91-9899032960

E-mail: k.s.sonkar@gmail.com

Abstract

A novel heme-peroxidase has been extracted from the latex of the medicinal plant *Artocarpus lakoocha (A. lakoocha)*, known for its potential anti-inflammatory and wound healing properties. To study its stability, structure, and dynamics, this protein was analyzed using far-UV circular dichroism, fluorescence spectroscopy, and activity measurements. The results demonstrated the presence of three folding states: the **native state** (N) at neutral pH, **intermediate states** including molten globule (MG) at pH 2 and acid-unfolded (UA) at pH 1.5 or lower, and acid-refolded (A) at pH 0.5, along with alkaline denatured (UB) at pH 8-12 and the third **denatured state** (D) at GuHCl concentrations exceeding 5 M. Absorbance studies indicated the presence of free heme in the pH range of 1-2. The protein showed stability and structural integrity across a wide pH range (3-10), temperature (70 °C), and high concentrations of GuHCl (5 M) and urea (8 M). This study is the first to report multiple 'partially folded intermediate states' of *A. lakoocha* peroxidase, with varying amounts of secondary structure, stability, and compactness. These results demonstrate the high stability of *A. lakoocha* peroxidase and its potential for biotechnological and industrial applications, making it a valuable model system for further studies on its structure-function relationship.

Keywords: *Artocarpus lakoocha*, Circular dichroism, Molten globule, Peroxidase.

1. Introduction

Artocarpus lakoocha (*A. lakoocha*) is a medicinal plant of the Moraceae family. It is a widely used medicinal plant of India for the treatment of many diseases like intestinal fluke *Haplorchis taichui* (Wongsawad, Wongsawad, Luangphai, & Kumchoo, 2005), taeniasis (Charoenlarp, Radomyos, & Bunnag, 1989), and cosmetics (Theorell, 1941). In an earlier report, we purified and characterized a novel peroxidase from *A. lakoocha.* The preliminary biophysical studies on *A. lakoocha* peroxidase revealed that the isolated novel enzyme from *A. lakoocha* is a stable heme-peroxidase with anti-inflammatory and wound healing properties (Sonkar et al., 2015). These broad applications provide a stimulus for detailed characterization of this peroxidase regarding folding behavior. Proteins undergo extensive conformational changes as part of their functionality; tracing these changes is important for understanding the mechanism by which proteins function.

Folding and conformational stability of proteins are primarily based on their amino acid sequence as well as cellular external variables such as temperature, pH, and chemical denaturants (Anfinsen, 1973; Masson & Lushchekina, 2022; Mohamed, El-Badry, Drees, & Fahmy, 2008). To predict the behavior of protein folding, it is important to understand the effect of these external variables on the thermodynamics of the folding process. Several other forces such as hydrogen bonding, disulfide bonding, entropic effects, hydrophobic, van der Waals, and electrostatic interactions, stabilize the folded state of the protein.

From unfolded chain to functional three dimensional state, the protein undergoes various non-native intermediate forms that are involved in various cell phenomena such as chaperone action (Díaz-Villanueva, Díaz-Molina, & García-González, 2015; Uversky, 2010), amyloid fibril formation in pathological cells (Almeida & Brito, 2020). A better understanding of protein folding mechanisms can help in treating diseases associated with protein misfolding, such as Alzheimer's, Parkinson's, and prion diseases (Rochet & Lansbury, 2000; Stefani & Dobson, 2003). The protein folding pathway is characterized by the accumulation of many intermediates between the native (N) and fully unfolded (U) states. Understanding these intermediate states is important to know how and when various forces come into play in directing the protein to attain three-dimensional functional structures. Studying these intermediates will help to explain the mechanism of folding of protein into the specific biologically active conformations (Benaim & Villalobo, 2002). Protein folding is a sequentially ordered process with the existence of stable

'molten globule' (MG) conformations between fully folded and unfolded states [3]. Molten

globules serve as a common equilibrium intermediate state in the protein folding pathway, which

is acquired under mild denaturing conditions. These molten globule states are believed to give

information on early events in the folding process, at a stage when specific side-chain

interactions are not yet formed. Molten globule was considered less compact than its native state

and more compact than its unfolded state with prominent secondary structure but slight or no

tertiary packing (Baldwin & Rose, 2013; Bhattacharyya & Varadarajan, 2013). The

characteristics of the molten globule state are an exposed hydrophobic surface that binds a

hydrophobic dye. This dye helps to yield experimental evidence concerning the structure of

intermediates throughout the protein folding state (Judy & Kishore, 2019). The exact role of

molten globule state is controversial among the proteins. The same protein may have varied

molten globule states in the folding pathway (Bychkova, Dolgikh, Balobanov, & Finkelstein,

2022). The experimental techniques typically used to detect molten globule states are far-UV and

near-UV CD spectroscopy, which detect the secondary and tertiary structure of protein. The

hydrophobic dye, such as ANS (8-anilino-naphtalene-1-sulfonate) binding experiments, detects

the formation of a loose hydrophobic core and estimates the extent of hydrophobic area exposed

to the solvent.

With this background, the conformational stability of *A. lakoocha* peroxidase from the latex of *A.*

lakoocha has been examined in a wide range of pH and temperature and in the presence of

various chemical denaturants (GuHCl). The present analysis sheds light on the biophysical

properties of *A. lakoocha* peroxidase to find the correlation between its structure and function

and the root and logic behind its distinct physiochemical properties. The conformational stability

of the native protein as well as intermediate states have been evaluated, compared, and

discussed. Several intermediates with distinct spectroscopic properties are populated under

various conditions.

Our initial biochemical analysis has revealed that the isolated peroxidase has a molecular mass of

53 kDa, and its molecular structure consists of four disulfide bridges, fourteen tyrosine residues,

and seventeen tryptophan residues, with an extinction coefficient ($\varepsilon_{280}^{1\%}$) of the enzyme at 16.3

(Sonkar et al., 2015). This peroxidase showed remarkable stability under various denaturing

conditions. In this section, conformational studies, and other factors responsible for such high

stability of peroxidase against various denaturing conditions, such as high concentrations of urea

and guanidine hydrochloride and a wide range of pH and temperature, have been reported.
Several equilibrium intermediate states with different stabilities have been obtained and
characterized to increase the understanding of the folding aspects of the peroxidase molecule.
The structural statistics about transition-state intermediates have been examined to identify the
whole protein folding method of peroxidase from the latex of *A. lakoocha*. These studies
establish a stable folded state of protein, which reflects its functional stability, and are aimed at
providing an insight into the folding of proteins in general, which may find application in
biotechnology and other fields of science.

2. Materials and methods

2.1 Materials

Peroxidase was purified from fresh latex of *Artocarpus lakoocha* plant using the method publish
ealier (Sonkar et al., 2015). ANS (8-anilino-1-naphthalene sulphonate) and GuHCl were
purchased from Sigma Chemicals, USA. Other reagents used were of analytical grade. The
samples were centrifuged before spectroscopic measurements, and the exact concentration of the
protein and pH of the buffer were determined before the measurements were taken.

2.2 Methods

2.2.1 Absorbance spectroscopy

Spectroscopic analysis was performed using a Beckman DU-640B spectrophotometer with a
constant-temperature cell holder. A protein concentration of 1 mM was used for all absorbance
measurements. The concentration of the enzyme was determined spectrophotometrically using
an extinction coefficient ($\varepsilon 280 1\%$) of 16.3 $M^{-1}cm^{-1}$. Far-UV circular dichroism (CD)
measurements were done using a JASCO J-815 spectropolarimeter (JASCO Corporation, Tokyo,
Japan) equipped with a constant temperature cell holder. The instrument was calibrated using
ammonium (+)-10-camphor sulfonate, and conformational changes in the secondary structure of
the peroxidase were monitored in the 200-260 nm range. After subtracting appropriate blanks,
mean residue ellipticity was calculated using the equation provided by Balasubramanian et al
(Balasubramanian & Kumar, 1976) as

$$[\theta] \text{ MRW} = \theta \text{obs x MRW}/10cl$$

Where θobs is the observed ellipticity in degrees, MRW represents mean residue weight, c
represents the concentration of protein (mg/ml), and l represents the path length in centimeters.

The mean weight of amino acid residues (MRW) was taken as 112. Sensitivities of 1 and 2

millidegrees/cm were used for far-UV CD measurements.

2.2.2 Fluorescence spectroscopy

Fluorescence measurements were carried out using an Eclipse Cary Varian UV-Vis

spectrofluorometer (Varian Inc., Palo Alto, California, USA) attached to a Peltier temperature

controller. For intrinsic tryptophan fluorescence, the protein was excited at 292 nm, and the

fluorescence emission was monitored in the 300-400 nm range. For ANS fluorescence, the

excitation was done at 380 nm, and the emission was monitored between 400-600 nm. For all

experiments, a protein concentration of 1 mM and excitation and emission slit widths of 10 and 5

nm were used, respectively. All the fluorescence spectra were recorded at 25 ℃ unless stated

otherwise. The activity of *A. lakoocha* peroxidase under various conditions of pH and chemical

denaturant was monitored using hydrogen peroxide as a hydrogen acceptor, and other

compounds such as guaiacol as a hydrogen donor. Twenty micrograms of peroxidase were added

to 0.2 mM of hydrogen peroxide (substrate I) with guaiacol in a 50 mM acetate buffer (substrate

II). The rate of change of absorbance of guaiacol was determined at 470 nm.

2.2.3 pH denaturation of A. lakoocha peroxidase

Acid denaturation of peroxidase was observed at a broad pH range using various buffers, such as

KCl–HCl buffer (pH 0.5–1.5), Gly–HCl buffer (pH 2–3.5), sodium acetate buffer (pH 4.0–5.5),

sodium phosphate buffer (pH 6.0–8.0), Tris–HCl buffer (pH 8.5–10.5), and Gly–NaOH buffer

(pH 11–12.5). The total concentration of all the buffers used in this study was 50 mM. A stock

solution of the protein was added to the appropriate buffer, and the mixture was incubated for 24

hr at 25 °C. The pH of the final suspension and the concentration of the protein in each sample

were confirmed by measurement at the final steps.

2.2.4 8-anilino-1-naphthalene sulfonate (ANS) binding assay

ANS is highly soluble in methanol (Hawe, Sutter, & Jiskoot, 2008), and a minimal amount of

methanol does not affect the structure of the protein, as *A. lakoocha* peroxidase shows activity

and stability up to 50 % methanol (Sonkar et al., 2015). Based on this, ANS solution was

prepared in methanol. An extinction coefficient value of 5000M−1 cm−1 at 350 nm was used for

ANS concentration measurement. A molar ratio of 1:100 for protein to ANS was used, and the

suspension was incubated for 30 minutes in the dark. Subsequently, the extrinsic fluorescence of

the protein-ANS complex was monitored by recording the emission spectrum between 400-600

nm using an excitation wavelength of 380 nm. Protein concentration of 1 mM and slit widths of

excitation and emission of 10 and 5 nm were used, respectively.

2.2.5   Guanidine hydrochloride and urea induced unfolding

Concentrations of GuHCl solutions were determined from the refractive index of the solution. As

the concentration of the chemical denaturant increased, denaturation of the enzyme was observed

at different pH conditions. To reach equilibrium, the peroxidase sample was incubated at the

corresponding denaturant concentration for 24 hr at 25 °C. To obtain transition midpoints, as

well as to judge the cooperativity or non-cooperativity of the transitions, the method used by

Shirley (1995) was followed for the analysis of unfolding transitions. To compare the results

from different measurements (molar ellipticity, fluorescence emission maximum, and relative

activity), all the results were normalized and expressed as a fraction unfolded. The formula used

for fraction unfolded calculation (Fu) is:

196            $\mathbf{F_u = (F_{obs} - F_n) / (F_u - F_n)}$

Where F_{obs} represent the observed signal at a given denaturant concentration, F_n represents the

signal for the native protein, and Fu represents the respective signal of the denatured protein. The

values for Fn and Fu are obtained by extrapolating the linear dependence of the signal on the

concentration of the denaturant before and after the transition, respectively. Denaturant blanks

are used to ensure that the observed signal is specific to the peroxidase and not due to the

presence of the denaturant. This equation is used in a technique called spectroscopic

denaturation, which is used to study the stability and unfolding of proteins.

2.2.6   Thermal unfolding of peroxidase

Temperature-induced denaturation of the enzyme was performed under specified conditions as a

function of increasing temperature. The enzyme samples are heated to increasing temperatures

and kept at each temperature for 15 minutes before measurements are taken. The temperature of

the samples is monitored using a thermocouple and a digital multimeter to ensure that the desired

temperature is reached and maintained. Additionally, the samples are checked for protein

aggregation using light scattering measurements, which can indicate if the proteins have

undergone irreversible denaturation due to heat. This method allows for the study of how

temperature affects the stability and folding of the enzyme, which can provide insight into its
function and potential for thermal inactivation.

2.2.7 Data analysis

The denaturation curves were plotted with the ratio of fluorescence intensity at the emission
wavelength maxima of the native protein and the denatured protein against the denaturant
molarity or temperature. A two-state $N \rightleftharpoons U$ unfolding mechanism is assumed, meaning that at
any point on the curve, only the folded and unfolded conformations are present in significant
concentrations. The method used by Shirley (Shirley, 1995) is followed to obtain the transition
midpoints and judge the cooperativity or non-cooperativity of the transitions. For comparison of
results obtained from different measurements such as molar ellipticity, fluorescence emission
maximum, and relative activity, all the results are normalized and expressed as the fraction of
unfolded protein. The fraction of unfolded protein, at any denaturant concentration, is calculated
using the equation: fraction unfolded = (Fobs – Fn)/(Fu – Fn), as previously mentioned.

Denaturation curves of urea and guanidine hydrochloride (GuHCl) were monitored using
CD spectroscopy. Denaturation experiments were conducted using Far-UV CD spectroscopy
with an acquisition range of 215–260 nm. Data at 222 nm were used for fitting purposes,
assuming a two-state model. This allowed for the determination of experimental equilibrium m
(m_{urea} and m_{GuHCl}) and C_m (concentration at which half-denaturation occurs) using following
Equation:

$$S = [S_N + m_N \cdot [\text{denaturants}] + (S_U + m_U) \cdot \left(\exp \frac{m_{denaturant} \cdot ([denaturant] - Cm)}{R.T} \right)] / \left[1 + (\exp \frac{m_{denaturant} \cdot ([denaturant] - Cm)}{R.T}) \right]$$

In this model, S represents the experimental CD signal as a function of denaturants
concentration. S_N and S_U are the fitted CD signals for the Native and Unfolded states at 0 M
denaturants, respectively. m_N and m_U represent the slopes of the native and unfolded state
baselines, while $m_{denaturants}$ describe the unfolding cooperativity. *R* is the ideal gas constant, and *T*
is the experimental temperature (298.15 K) (Pacheco-García et al., 2022). This model provides a
simplified approach for comparing the cooperativity of reversible chemical unfolding in *A.*
lakoocha peroxidase.

3. Results

3.1 Absorbance spectroscopy

To elucidate the structure-function relationship of *A. lakoocha* peroxidase and its impact on
enzymatic activity, various biophysical techniques were used. These approaches facilitate the
observation of enzyme intermediates under varying pH conditions and during the denaturation
process. The absorbance spectra of peroxidase showed four bands (Fig. 1A) in the visible and
ultraviolet (UV) regions: the α-band at ~630 nm, the β-band at ~502 nm, the γ-band or Soret
band at 404 nm, and the absorbance spectra of peroxidase at 280 nm. These bands can be used as
a tool to investigate the structural and functional properties of *A. lakoocha* peroxidase. Changes
in the wavelength or intensity of these bands upon ligand binding or unfolding can provide
insight into the conformational changes of the enzyme. For example, substantial changes at 404
nm were observed upon ligand binding or unfolding, indicating changes in the tertiary and/or
quaternary structure of the enzyme. However, changes at 280 nm were not observed under native
or unfolded conditions, indicating that the secondary structure of the enzyme remains unchanged.
The shifts at the 404 nm wavelength are evaluated to study the conformational change of the
peroxidase. The absorbance spectra of *A. lakoocha* peroxidase in its native state, at pH 2 and in
its unfolded state (6 M GuHCl) (Fig. 1B) are compared to observe the changes in the spectra.
Under native conditions, the absorbance spectra show a soret peak at 404 nm, which is absent in
the unfolded state. This indicates that the conformational change of the enzyme occurs upon
unfolding. Additionally, fig. 1C shows the peroxidase absorbance maxima measured at 404 nm
under varying pH. The bell-shaped pattern suggests that the peroxidase is stable in the range of
pH 3.5 to 8.5, which corresponds to the enzyme's physiological pH range. This information is
useful for understanding the stability and activity of the enzyme under different conditions.
263

3.2 . Fluorescence spectroscopy

The fluorescence spectra of peroxidase under native conditions at pH 2, 11.0, and with 6 M
GuHCl have been shown in fig. 1D. The intrinsic fluorescence spectrum of peroxidase in its
native state has an emission maximum at 333.07 nm, indicating a hydrophilic environment of its
tryptophan residues. Under acidic and alkaline conditions, the fluorescence emission maximum
remains the same, but compared to the native protein, the intensity of fluorescence increases by
57.4% at pH 2 (178.43 to 280.88) and decreases by 16.42% at pH 11 (178.43 to 149.13). This
indicates that changes in pH can affect the microenvironment of the tryptophan residues and

influence their fluorescence properties. On complete unfolding of the protein in the presence of 6

M GuHCl, the emission maximum of peroxidase suffers a red shift of 8 nm (333.07 nm to 341.07

nm) along with an approximately 175% increase in fluorescence intensity (from 178.43 to

491.04) compared to the native condition. This suggests that upon unfolding, the tryptophan

residues are exposed to a more hydrophobic environment, resulting in an increase in fluorescence

intensity and a red shift in the emission maximum. The effect of different pH on *A. lakoocha*

peroxidase was also observed by the intrinsic fluorescence emission scale. The wavelength

maximum of peroxidase as a function of pH has been shown in (Fig. 1E). The transition curve

did not show much change in the emission maximum under varying pH and appeared to occur

through three state transitions. The first transition shows a red shift of 1 nm on decreasing pH

from 4.0 to 2.0 under acid unfolded state. The second transition occurred between 6.0 and 8.5 pH

along with a red shift of 2nm. On increasing the pH to the basic region, a third transition occurs

in the range of 9.0 to 10.5 pH, with a blue shift of 4 nm in the fluorescence emission maxima,

resulting in the alkaline denatured state of peroxidase. The pH-induced transition is non-

cooperative with a slight change in the transition maxima because tryptophan residues, even in

their native state, seem to be exposed to the solvent. Changes in the fluorescence intensity as a

function of pH are shown in fig. 1F. The fluorescence intensity spectra displayed two transition

phases. The first transition occurs from the native state to acid unfolded state (between pH 4.0 to

pH 2), with a transition mid-point at pH 3.0, where the fluorescence intensity is increased

marginally. The second transition towards basic pH is incomplete and takes place from pH 8.0 to

12.0. This study of the intrinsic fluorescence spectra of peroxidase under different pH conditions

provides insight into the conformational changes occurring in the enzyme and its stability at

different pH.

3.3. *ANS binding assay*

ANS (8-anilino-1-naphthalene sulphonate) is also known as a polarity-sensitive extrinsic

fluorescent probe. The effect of pH on the ANS fluorescence of peroxidase at different pH is

shown in fig. 2A and B. ANS emission spectra were recorded with excitation at 380 nm. At pH

2.5 proteins show partial ANS binding. On further decreasing the pH from 2.5 to 1.0, an increase

in fluorescence intensity was observed. At pH 2 enzymes showed maximum ANS binding.

Compared to native state, at pH 2 ANS fluorescence intensity showed up to 5-fold increase and
fluorescence maxima exhibited the red shift from 481 nm (pH 2) to 508.05 nm (pH 7). ANS
fluorescence decreases on either side of 2 pH and at the alkaline denatured state (UB state, pH 8-
12), ANS binding is almost absent and stable up to pH 12.

3.4. Circular dichroism

The effect of varying pH on the secondary structure of peroxidase was also studied. The CD
spectra of *A. lakoocha* peroxidase at acidic, neutral, and denatured conditions (6 M GuHCl) are
presented in fig. 4A. At pH 7 in the far-UV region, the CD spectrum of peroxidase comprises
strong negative ellipticity at 208 and 222 nm, suggesting that the molecule may consist of α-
helix and β-sheet rich regions and belong to the α + β class of proteins (Kumar, Tripathi, de
Moraes, Caruso, & Jagannadham, 2014; Manavalan, 1983). Fig. 2C displays that the mean
residue ellipticity at 222 nm is -5300.2 deg.cm^2 dmol^{-1} in the native condition (Fig. 4A) while
under an acidic state at 1.0 pH, the ellipticity at 222 nm shifts to -3779.9 deg.cm^2 dmol^{-1}.
However, under similar conditions, all the spectral features of peroxidase are lost in the presence
of 6 M GuHCl, indicating that the enzyme drops the secondary structural organization under
denaturing conditions.
The structural changes of peroxidase according to pH are indicated by shifts in ellipticity at 208
and 222 nm, as shown in fig. 2D. These structural changes display that the secondary structural
contents of proteins remain the same in the range of 4.0-9.5 pH and decrease on either side of
this range, without changing the shape of the spectra. The changes are similar in magnitude at
lower pH and in the basic range.

3.5. Guanidine hydrochloride induced unfolding of peroxidase

Guanidine hydrochloride (GuHCl) induced the unfolding of peroxidase under neutral conditions,
which, when examined by alterations in secondary structure, absorbance, and fluorescence,
showed structural stability till 3.0 M GuHCl (Fig. 3A). Under natural conditions, GuHCl-induced
unfolding results in cooperative sigmoidal curves (Fig. 5A) with a transition midpoint at 4.5 ±
0.1 M GuHCl. However, at pH 2, enzymes displayed structural stability up to 1.5 M GuHCl (Fig.
3B). The transitions as measured by different spectroscopic techniques are cooperative and non-
coincidental (Fig. 3B). This observation illustrated that fluorescence intensity is lost first,

followed by losses in secondary structure. Loss in secondary structure took place between 2.0

and 5.0 M of GuHCl, whereas the changes in intrinsic fluorescence occurred between 1.5 and 5.0

M of GuHCl at pH 2. The transition mid-points (Cm) of denaturation followed by far UV CD are

3.5 ± 0.1 M, and by absorbance and fluorescence intensity is approximately 3.25 ± 0.1 M. The

effect of GuHCl on the alkaline denatured state of a protein at pH 11 is similarly studied. The

transitions as measured by different spectroscopic techniques are cooperative and coincidental

(Fig. 3C). Loss of secondary structure and intrinsic fluorescence took place between 2.5 and 5.0

M of GuHCl, and the enzyme showed structural stability until 2.0 M of GuHCl.

When ANS fluorescence was used to monitor the denaturation process in the native and alkaline

denatured states at pH 11, peroxidase did not specify any ANS binding, even after a prolonged

exposure. That may be due to an absence of contact with the hydrophobic surface. These

characteristics indicate that *A. lakoocha* peroxidase is more susceptible to denaturation in acidic

conditions in comparison to alkaline conditions. So, it is meaningful to observe the structural

integrity of the molecule at an acidic pH. Fig. 3D shows the ANS binding pattern of peroxidase

with increasing concentrations of GuHCl, at pH 2. Under the acid denatured state, a decrease in

ANS fluorescence intensity by increasing the molarity of GuHCl was observed with a red shift in

its wavelength maximum with GuHCl at pH 2 (Fig. 3E).

3.6.   *Urea induced unfolding of peroxidase*

Under neutral conditions, urea does not cause any structural agitation in the protein, as the

enzyme retains its maximum structural and functional parameters even at 5.5 M urea

concentration (Fig. 4A). The urea induced unfolding studies are therefore carried out at low and

high pH, where enzymes are more susceptible to denaturants. However, urea induced unfolding

at a lower pH yielded some interesting facts. The denaturation is followed by absorbance, far UV

CD and fluorescence. At pH 2, the urea induced unfolding is cooperative, but non-coincidental

(Fig. 4B). The transition mid-points for intrinsic fluorescence and absorbance are 4.25 ± 0.1 M

and for secondary structure are 4.5 ± 0.3 M. The loss in secondary structure occurred between

2.0 and 7.0 M urea. But, at pH 11, where the enzyme is in an alkaline denatured state, the

unfolding transitions by different probes are cooperative, with a transition mid-point at $4.75 \pm$

0.1M (Fig. 4C).

The cooperative but non-coincidental transitions at pH 2, along with an increase in ANS binding

at low concentrations of urea (Fig. 4D and E) indicate that a discrete intermediate state exists in

the urea induced unfolding of peroxidase at pH 2. An increase in ANS fluorescence intensity

coupled with a red shift in the wavelength maximum of peroxidase at pH 2 is shown in fig. 4D

and E.

3.7.   Thermal denaturation of peroxidase

Temperature, the oldest mode of protein denaturation, can induce structural changes and hence

provide ample information about a protein molecule. The temperature induced unfolding of

peroxidase at pH 7 was measured (Fig. 4F). The transition midpoints of temperature induced

unfolding of peroxidase are very high and come in the range of 62.5 ± 0.5 °C. Further, the

structure of the protein melts at a much higher temperature, at 70 °C, indicating marked

structural agitations already induced by the increased concentration of protons.

4.   Discussion

In the era of proteomics, it is becoming very important to understand the structure-function

relationship of proteins using various biophysical tools (Neet & Lee, 2002). Despite the

importance of peroxidase, relatively few biophysical studies have been carried out on this class

of proteins. Because of the dependency of catalytic activity on pH, the stability of an enzyme

determines its functionality and may limit its wide application. Therefore, pH denaturation of

peroxidase was monitored by various spectroscopic techniques to probe the different aspects of

conformational changes in its structure and the identification of populated intermediates at

extremes of pH, during denaturation.

4.1.   Effect of pH on the absorbance spectra of A. lakoocha peroxidase

The absorbance technique is used to measure the absorption of radiation by a sample as a

function of wavelength or frequency. The absorbance spectra of the protein at different pH levels

are shown in Fig. 1A, which shows spectra of the protein under neutral conditions that are

similar to the characteristic heme peroxidase spectra. The absorption bands in the visual

spectrum region (α- and β-peaks) and at the boundary of the visual and UV parts (Soret γ) are

due to the intensively colored prosthetic groups of peroxidases. Other absorption bands of

peroxidase solution are located in the UV region: φ-band with a maximum of 280 nm, indicating

a significant contribution from tryptophan and tyrosine residues also due to the light absorption

by aromatic amino acids in proteins. As revealed from Fig. 1A, peroxidase spectra do not show

much change in the range of pH 2.5 to 8.5, which indicates good stability of peroxidase in this

state. But at lower pH, heme peroxidase spectra express a left shift with a declined absorbance at

404 nm due to denaturation, which is a clear indication of free heme in the range of pH 1 to 2.

This result specifies that at lower pH peroxidase releases its prosthetic group (heme). Whereas

beyond pH 8, the heme group of peroxidase shows OH^- or H_2O binding, with an increase in pH,

which results in the right shifting of soret band. At higher pH, the Q-band of peroxidase gets split

and the soret peak shifts to the right. Though the shift in absorbance intensity at the 404 nm

(Soret peak) gives information regarding the stability and activity of peroxidase, this shift at 404

nm wavelength is investigated further to follow the conformational change of the protein in the

present investigation.

The spectra in Fig. 1B show a decrease in the soret band of peroxidase at pH 2 and 6 M GuHCl,

indicating the loss of the heme group and denaturation of the protein. Fig. 1C shows that the

enzyme retains proteolytic activity in the pH range of 3.5-8.5 and drops on either side of it,

suggesting that the native structure and functional stability of the protein are maintained in this

pH range. The decline in proteolytic activity outside of this range is likely due to the repulsion of

columbic forces from the positive charges of the polypeptide chains.

410   *4.2.   Fluorescence spectroscopy*

The fluorescence emission maximum of chromophores, such as tyrosine and tryptophan residues,

can change when exposed to different solvent environments. The quantum yield of fluorescence

can also decrease when the chromophores interact with quenching agents in the solvent or within

the protein. In the case of *A. lakoocha* peroxidase, the fluorescence spectra under different pH

and denaturing conditions are shown in fig. 1D. The tryptophan excitation spectra reflect the

average environment of the tryptophan and shows a red shift in the emission maximum when the

chromophores are exposed to solvent. The results of the experiment, as observed in fig. 1E,

indicate that the tryptophan residues become exposed to solvents due to the unfolding and

disclosure of chromophores to the solvent, particularly under denaturing conditions with 6 M

GuHCl. Additionally, the intensity of fluorescence increases at pH 2 but decreases at pH 11,

likely due to the presence of a heme group in the peroxidase that acts as a main quencher and

quenches the energy in the native state. However, upon denaturation with 6 M GuHCl, the

fluorescence intensity is amplified compared to the acidic condition, indicating the complete

exposure of tryptophan residues to an aqueous environment.

The observations suggest that the fluorescence intensity of the peroxidase protein

increases as the protein loses iron upon denaturation. The pH-induced transition of peroxidase in

the alkaline range, as monitored by fluorescence intensity, seems to be cooperative, with a

transition mid-point at pH 3. The fluorescence wavelength maxima remained largely unaffected

throughout different pH values (Fig. 1E), but the fluorescence intensity revealed changes at

different pH values (Fig. 1F). Lower pH values resulted in higher fluorescence intensity, while

higher pH values resulted in a decrease in fluorescence intensity, likely due to the quenching of

tryptophan fluorescence by specific quenching groups such as protonated carboxyl and imidazole

groups.

434 *4.3. Acid-induced changes in peroxidase*

ANS binding is a tool used to measure the exposure of hydrophobic surfaces in enzymes. The

magnitude of ANS binding can indicate the structural integrity of a protein. In the case of *A.*

lakoocha peroxidase, at pH levels above 3, no binding of ANS to peroxidase was observed.

However, at lower pH levels, some binding was detected, albeit with a lower magnitude. As the

pH decreased from 3 to 2.5, an increase in ANS binding was observed, as indicated by an

increase in fluorescence intensity from 21.76 to 149.78 and a blue shift of the emission

maximum from 483 nm to 482 nm. On further decreasing pH to lower values, the protein gets

refolded by pH 0.5, driving the molecule into acid refolded state caused by the shielding of

columbic repulsions by counterions., as shown in Fig. 2A. However, at pH 2 proteins show

maximum ANS binding along with a significant amount of secondary structure with no tertiary

contacts. This sudden increase in ANS fluorescence intensity is due to maximum exposure of

hydrophobic surfaces. That is one of the properties of molten globule state. This result indicates

the presence of molten globule state at pH 2. The results of the experiment as observed in fig. 2A

show that the peroxidase retains its proteolytic activity and native structure within a pH range of

3.0-10.0, as indicated by the absence of ANS binding in this range, suggesting that the protein

has a relatively stable and rigid native conformation in a wide range of pH values but becomes
more flexible and susceptible to changes in acidity.
The pH-induced conversion of this protein shows the first transition phase in between and pH 2.5
and 4.0 (midpoint 3.25 plus second transition in the middle of pH 1 and 2.5 (midpoint 1.5). As
observed by absorbance and fluorescence spectra, at pH 7.5 and above, protein turns into an
alkaline denatured state (UB), besides at pH 1.5 or lower, the protein unfolds to form acid-
unfolded state (UA). While on further decreasing the pH to 0.5 induces refolding of the UA-state
and the resulting state has been termed as acid-refolded 'A-state' (Fig.2).

Circular dichroism

The use of circular dichroism (CD) spectroscopy and the hydrophobic dye ANS allowed for the
study of the structure and stability of *A. lakoocha* peroxidase. The Far-UV CD spectra of the
enzyme (Fig. 2C) showed it to be consistent with the $\alpha + \beta$ class of proteins and possibly contain
distinct α and β-rich domains. The peroxidase was found to be quite stable in terms of secondary
structure, with only a 50% decrease in ellipticity at lower pH. However, under denaturing
conditions, the peroxidase lost all prominent peaks in aromatic regions, indicating the loss of
secondary structural features and the enzyme being in an unfolded state. The shifts in ellipticity
at varying pH were found to not follow a simple two-state transition but rather a three-step
transition (Fig. 2D). The first transition from the native state to acid unfolded state occurred
between pH 4 - 1, with a half bell shape in the range of pH 1 - 4, revealing the loss of some
prominent peaks. This loss of secondary structure, as well as reduced activity, represents the
enzyme acid unfolded state. The second transition occurred between pH 4 to 8 under native
conditions, and the third transition occurred between pH 8 – 12 with a mid-point at 10.5 pH. The
use of hydrophobic dye ANS also revealed that despite the melting of its secondary structure, a
large amount of hydrophobic clusters were left over in the acid-unfolded state (Fig. 2A).
474

4.4. GuHCl induced unfolding of peroxidase

The analysis of denaturation curves induced by GuHCl can provide information about the
conformational stability of the enzyme. The denaturation of peroxidase by GuHCl at neutral pH
is a cooperative, two-state process with a high transition mid-point of about 4.5 ± 0.1M (fig. 3A).
However, at pH 2 and 11.0, (fig. 3B and C) the transition mid-points are lower as compared to

the same parameters at pH 7. When ANS fluorescence was used to monitor the denaturation
process under different conditions, enzyme only reveals ANS binding with acidic condition.
Where the decrease in ANS fluorescence intensity coupled with red shift in its wavelength
maximum was found, as shown in fig.3D. At pH 2 maximum ANS binding as well as other
spectroscopic properties exist due to increase of distance from the heme group, which serves as a
quencher in heme containing peroxidase. This kind of quenching effect of heme has also been
reported in other heme containing peroxidases, like lactose peroxidase and horse radish
peroxidase (Carvalho et al., 2003; Zelent B, 2010). ANS binding to peroxidase at pH 2 in the
presence of ≥2.5M GuHCl suggests the absence of surface hydrophobic patches, since no ANS
binding was observed. While decreasing the concentration of GuHCl, one intermediate was
found at 1.0 M GuHCl, here the protein shows maximum (Fig.3E) ANS fluorescence intensity,
which indicates the exposure of hydrophobic residues in this state. That's in line with the idea of
molten globule state (MG) at the particular GuHCl concentration. While there is absence of ANS
binding to peroxidase at 3 M GuHCl and above, which suggests the absence of surface
hydrophobic patches and refolding and stabilization due to specific stabilizing interactions
between the intermediate state and GuHCl.

4.5. Urea induced unfolding of peroxidase

The stability of the enzyme peroxidase against urea was evaluated under different pH conditions.
Results revealed that at neutral pH, the enzyme retained its native structure and function even
when exposed to 5.5M urea (Fig .4A), indicating stability against urea. However, at an acidic pH
(2), urea caused the enzyme to unfold cooperatively, with transition midpoints for secondary and
tertiary structures at 4.5 ± 0.3M and 4.25 ± 0.1M urea, respectively (Fig. 4B). The unfolding
process was observed using far-UV CD and fluorescence, which revealed cooperative and non-
overlapping transitions (Fig. 4B) along with an increase in ANS binding at low concentrations of
urea (Fig. 4D & E). Additionally, ANS fluorescence intensity, which is highest at pH 2 in the
absence of denaturant, decreased parallel with an increase in urea concentration. The sigmoidal
change in fluorescence intensity with urea at pH 2 is attributed to configurational changes in the
amino acids. The unfolding process at pH 2 showed the same changes in far-UV CD and
wavelength maxima as at pH 7 (Fig. 4B). However, complete loss of activity above 5 M urea, a
sigmoidal increase in fluorescence intensity, ANS binding, and conversion to beta-sheet-like

spectra seen by far-UV CD above 3.5 M urea were due to perturbation of the enzyme's functional

structure (Fig. 4B). These results suggest that a discrete intermediate state exists in the urea-

induced unfolding of peroxidase.

Overall, this study found that the enzyme peroxidase unfolds differently when exposed to

urea and guanidine hydrochloride (GuHCl) under similar conditions. The results showed that

urea is less effective than GuHCl in reducing the mid-points of unfolding at a given

concentration. This difference in behavior is likely due to the ionic character of the two

denaturants. Urea and GuHCl interact with peptide bonds but have different electrostatic and

hydrophobic properties. GuHCl, which is positively charged, may stabilize the protein through

its anionic effect, while urea does not have this effect. Additionally, the counter-anion of GuHCl,

chloride, may shield electrostatic repulsion and allow forces that favor the folding and stability

of the protein. However, the stabilizing effect of GuHCl may not always be predominantly ionic

in nature (Zarrine-Afsar, Mittermaier, Kay, & Davidson, 2006). The study also discussed the

effects of temperature on the structure and activity of peroxidase from *A. lakoocha*. It was found

that the enzyme retained approximately 45% of its structure under neutral conditions and had a

high degree of stability, with a transition midpoint (Tm) of 62.5±0.5°C (Fig. 4F). The complete

unfolding parameters of *A. lakoocha* peroxidase are summarized in Table 1. The accumulation of

intermediates in the GuHCl and urea induced unfolding pathways is presented here, which

indicates that peroxidase may possess two domains that are behaving differently in various

denaturing conditions, leading to the formation of a number of intermediate states.

5. Conclusion

In this study, we observed three different folding states of a peroxidase from the latex of *A.*

lakoocha using various spectroscopic techniques under different pH, denaturing conditions, and

temperatures. At pH 7, the enzyme is in its **native state,** with proteolytic activity, intact

secondary structure, and without any hydrophobic patches. At lower pH values, the enzyme

transitions through intermediate states such as the **molten globule** state, the acid-unfolded state

(**UA state**) and acid refolded state **(A state)**. In the alkaline denatured state (**UB state**), ANS

binding is almost absent, indicating that *A. lakoocha* peroxidase is more susceptible to

denaturation under acidic conditions, than alkaline conditions. Fluorescence intensity decreases

at higher pH in the UB state due to a reduced distance between tryptophan and specific

quenching groups, leading to quenching of tryptophan fluorescence. Under denaturing conditions, the enzyme can reach a completely denatured state when exposed to GuHCl concentrations above 5 M. At low concentrations (1.0 M), GuHCl acts as an anion, while at concentrations of 2.5 M or higher, it behaves as a chaotrope. Further experiments revealed that incubation of the peroxidase in 8 M urea overnight resulted in only a 60% loss of secondary structure, whereas incubation with 6 M GuHCl led to a complete loss of secondary structure. Our study demonstrates that the purified peroxidase from the latex of *A. lakoocha* is a robust model system for investigating protein folding pathways. The identification of these intermediate states represents a significant advancement in the field of protein folding and lays the foundation for future research into the enzyme's physiological substrates, structural characteristics, and potential applications in the biotechnology and pharmaceutical industries.

Author Contribution

K.S. purified the protein and performed biophysical experiments, with assistance from M.P. and A.K. for data analysis. The experimental design was planned by K.S. and J.M. The manuscript was written by K.S. with input from all authors. We thank Prof. Suman Kundu from the Department of Biochemistry at UDSC Delhi for providing the necessary infrastructure.

Conflict of interest

The authors declare no competing financial interests.

Acknowledgements

Financial assistance from the CSIR, Government of India, in the form of fellowships to KS is acknowledged. Grants and funds from UGC and DBT for infrastructure are also acknowledged.

References

Almeida, Z. L., & Brito, R. M. (2020). Structure and aggregation mechanisms in amyloids. *Molecules, 25*(5), 1195.

Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. *Science, 181*(4096), 223-230.

Balasubramanian, D., & Kumar, C. (1976). Recent studies of the circular dichroism and optical rotatory dispersion of biopolymers. *Applied Spectroscopy Reviews, 11*(2), 223-286.

Baldwin, R. L., & Rose, G. D. (2013). Molten globules, entropy-driven conformational change and protein folding. *Curr Opin Struct Biol, 23*(1), 4-10. doi:10.1016/j.sbi.2012.11.004 S0959-440X(12)00184-4 [pii]

Benaim, G., & Villalobo, A. (2002). Phosphorylation of calmodulin. Functional implications. *Eur J Biochem, 269*(15), 3619-3631. doi:3038 [pii]

Bhattacharyya, S., & Varadarajan, R. (2013). Packing in molten globules and native states. *Curr Opin Struct Biol, 23*(1), 11-21. doi:10.1016/j.sbi.2012.10.010

S0959-440X(12)00180-7 [pii]

Bychkova, V. E., Dolgikh, D. A., Balobanov, V. A., & Finkelstein, A. V. (2022). The Molten Globule State of a Globular Protein in a Cell Is More or Less Frequent Case Rather than an Exception. *Molecules, 27*(14), 4361.

Carvalho, A. S., Melo, E. P., Ferreira, B. S., Neves-Petersen, M. T., Petersen, S. B., & Aires-Barros, M. R. (2003). Heme and pH-dependent stability of an anionic horseradish peroxidase. *Arch Biochem Biophys, 415*(2), 257-267. doi:S0003986103002753 [pii]

Charoenlarp, P., Radomyos, P., & Bunnag, D. (1989). The optimum dose of Puag-Haad in the treatment of taeniasis. *J Med Assoc Thailand, 72*(2), 71-73.

Díaz-Villanueva, J. F., Díaz-Molina, R., & García-González, V. (2015). Protein folding and mechanisms of proteostasis. *International journal of molecular sciences, 16*(8), 17193-17230.

Hawe, A., Sutter, M., & Jiskoot, W. (2008). Extrinsic fluorescent dyes as tools for protein characterization. *Pharmaceutical research, 25*, 1487-1499.

Judy, E., & Kishore, N. (2019). A look back at the molten globule state of proteins: Thermodynamic aspects. *Biophysical Reviews, 11*, 365-375.

Kumar, R., Tripathi, P., de Moraes, F. R., Caruso, I. P., & Jagannadham, M. V. (2014). Identification of folding intermediates of streblin, the most stable serine protease: biophysical analysis. *Appl Biochem Biotechnol, 172*(2), 658-671. doi:10.1007/s12010-013-0565-8

Manavalan, P., and Jonson jr, W. C. . (1983). Sensitivity of circular dichroism to protein tertiary structure class. *Nature, 305*, 831-832.

Masson, P., & Lushchekina, S. (2022). Conformational Stability and Denaturation Processes of Proteins Investigated by Electrophoresis under Extreme Conditions. *Molecules, 27*(20), 6861.

Mohamed, S. A., El-Badry, M. O., Drees, E. A., & Fahmy, A. S. (2008). Properties of a cationic peroxidase from Citrus jambhiri cv. Adalia. *Applied biochemistry and biotechnology, 150*, 127-137.

Neet, K. E., & Lee, J. C. (2002). Biophysical characterization of proteins in the post-genomic era of proteomics. *Mol Cell Proteomics, 1*(6), 415-420.

Pacheco-García, J. L., Loginov, D. S., Naganathan, A. N., Vankova, P., Cano-Muñoz, M., Man, P., & Pey, A. L. (2022). Loss of stability and unfolding cooperativity in hPGK1 upon gradual structural perturbation of its N-terminal domain hydrophobic core. *Scientific reports, 12*(1), 17200.

Rochet, J. C., & Lansbury, P. T., Jr. (2000). Amyloid fibrillogenesis: themes and variations. *Curr Opin Struct Biol, 10*(1), 60-68. doi:S0959-440X(99)00049-4 [pii]

Shirley, B. A. (1995). Urea and guanidine hydrochloride denaturation curves. *Methods Mol Biol, 40*, 177-190. doi:10.1385/0-89603-301-5:177

Sonkar, K. S., Manendra Pachauri, Amit Kumar, Ankita Shukla, Monika Patel, & Jagannadham., M. V. (2015). Heme-peroxidase from medicinal plant Artocarpus lakoocha: Purification, characterization and wound healing studies. *Biocatal Agricu Biotechnol, 4*, 180-190.

Stefani, M., & Dobson, C. M. (2003). Protein aggregation and aggregate toxicity: new insights
into protein folding, misfolding diseases and biological evolution. *J Mol Med (Berl),*
81(11), 678-699. doi:10.1007/s00109-003-0464-5
Theorell, H. (1941). Crystalline peroxidase. *Enzymologia, 10*, 250.
Uversky, V. N. (2010). Targeting intrinsically disordered proteins in neurodegenerative and
protein dysfunction diseases: another illustration of the D(2) concept. *Expert Rev*
Proteomics, 7(4), 543-564. doi:10.1586/epr.10.36
Wongsawad, C., Wongsawad, P., Luangphai, P., & Kumchoo, K. (2005). In vitro effects of
aqueous extract from Artocarpus takoocha Roxb on tegumental surface of Haplorchis
taichui. *Southeast Asian J Trop Med Public Health, 36 Suppl 4*, 112-116.
Zarrine-Afsar, A., Mittermaier, A., Kay, L. E., & Davidson, A. R. (2006). Protein stabilization
by specific binding of guanidinium to a functional arginine-binding surface on an SH3
domain. *Protein Science, 15*(1), 162-170.
Zelent B, S. K., Vanderkooi JM. . (2010). Differential scanning calorimetry and fluorescence
study of lactoperoxidase as a function of guanidinium-HCl, urea, and pH. . *Biochim*
Biophys Acta, (7)(1804), 1508-1515.

655   Figure captions:

658   Fig.1.   pH-induced conformational changes of *A. lakoocha* peroxidase: (A) UV
659   absorbance spectra at different pH, (B) Spectra at pH 7.0 and at 6 M GuHCl have
660   been compared with spectra at pH 2.0 (free of heme), (C) The absorbance
661   spectrum at 404 nm under different pH conditions., (D) fluorescence spectra at
662   different pH and in 6.0M GuHCl, (E) fluorescence wavelength maxima, and (F)
663   fluorescence intensity at different pH.

665   Fig.2.   Effect of pH on the ANS fluorescence of *A. lakoocha* peroxidase: (A) ANS
666   binding to peroxidase was studied at pH 1.0, pH 1.5, pH 2.0, pH 2.5, pH 7.0, and
667   in the presence of 6M GuHCl, and (B) For intrinsic fluorescence measurement,
668   samples were incubated for 24 h at 25oC. ANS emission spectra were recorded
669   with excitation at 380nm., pH induced conformational changes of *A. lakoocha*
670   peroxidase: (C) far-UV CD spectra at different pH and in 6.0 M GuHCl, and (D)
671   Ellipticity at 208 nm and 222 nm as a function of pH.

673   Fig.3.   GuHCl induced unfolding of A. lakoocha peroxidase at different pH: (A) pH 7.0,
674   (B) pH 2.0, and (C) pH 11.0, by following far-UV CD, fluorescence, and UV
675   absorbance, Effect of pH on the ANS fluorescence of A. lakoocha peroxidase: (D)
676   ANS binding spectra, as a function of increasing concentration of GuHCl at pH
677   2.0 (E) ANS fluorescence intensity and wavelength maxima at pH 2.0, as a
678   function of increasing concentration of GuHCl.

680   Fig.4.   Urea induced unfolding of A. lakoocha peroxidase at different pH: (A) pH 7.0,
681   (B) pH 2.0, and (C) pH 11.0, by following far-UV CD, Fluorescence and UV
682   absorbance,   Effect of pH on the ANS fluorescence of A. lakoocha peroxidase:
683   (D) ANS binding spectra, as a function of increasing concentration of urea at pH
684   2.0, (E) ANS fluorescence intensity and wavelength maxima at pH 2.0, as a
685   function of increasing concentration of urea, and (F)Temperature induced
686   unfolding of A. lakoocha peroxidase at pH 7.0: The structural changes are
687   followed by UV absorbance, far- UV CD, and fluorescence.

692   .

696   Fig.1

697
698

699   Fig.2

701   Fig.3

703   Fig.4

711 **Table.1:** Unfolding parameters of *A. lakoocha* peroxidase

Condition	Denaturant	Method	Transition mid-point
pH 7.0 (Native state)	GuHCl	CD[Θ]$_{222}$	4.5 ± 0.1 M
		Fluorescence	4.5 ± 0.1 M
		absorbance	4.5 ± 0.1 M
	Urea	CD[Θ]$_{222}$	6.5 ± 0.1 M
		Fluorescence	6.5 ± 0.1 M
		absorbance	6.5 ± 0.1 M
	Temperature	CD[Θ]222	65.0 ± 0.5°C
		Fluorescence	62.5 ± 0.5°C
		absorbance	62.5 ± 0.5°C
pH 2.0 (Molten globule state)	GuHCl	CD[Θ]$_{222}$	3.5 ± 0.1 M
		Fluorescence	3.25 ± 0.1 M
		absorbance	3.25 ± 0.1 M
	Urea	CD[Θ]$_{222}$	4.5 ± 0.3 M
		Fluorescence	4.25 ± 0.1 M
		absorbance	4.25 ± 0.1 M
pH 11.0 (Alkaline denatured state)	GuHCl	CD[Θ]$_{222}$	3.5 ± 0.1 M
		Fluorescence	3.5 ± 0.1 M
		absorbance	3.5 ± 0.1 M
	Urea	CD[Θ]$_{222}$	4.75 ± 0.1 M
		Fluorescence	4.75 ± 0.1 M
		absorbance	4.75 ± 0.1 M

715