LIMBARE: an Advanced Linear Mixed-effects Breakpoint Analysis with Robust Estimation Method with Applications to Longitudinal Ophthalmic Studies

Authors:

TingFang Lee[1,2], Joel S Schuman[1,3,4,5], Maria de los Angeles Ramos Cadena[1], Yan Zhang[2], Gadi Wollstein[1,3,5*], Jiyuan Hu[2].

Affiliations:

1. Department of Ophthalmology, NYU Langone Health, New York, NY, United States.
2. Departments of Population Health, NYU Langone Health, New York, NY, United States.
3. Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn, NY, United States.
4. Department of Electrical and Computer Engineering, NYU Tandon School of Engineering, Brooklyn, NY, United States.
5. Center for Neural Science, NYU College of Arts and Sciences, New York, NY, United States.

Supporting Grant: NIH R01-EY013178 (Bethesda, MD), R01-EY030770, P30-EY013079, U54-MD000538, R33-AG057382, unrestricted grant from Research to Prevent Blindness.

Conflict of Interest: Dr. Schuman receives royalties for intellectual property licensed by Massachusetts Institute of Technology and Massachusetts Eye and Ear Infirmary to Zeiss.

22   *** Corresponding Author:** Gadi Wollstein, Department of Ophthalmology, NYU Langone

23   Health, 222 East 41st St., New York, NY, 10017 United States; Gadi.Wollstein@nyulangone.org

Abstract

Purpose: Broken stick analysis is a widely used approach for detecting unknown breakpoints where association between measurements is non-linear. We propose LIMBARE, an advanced **li**near **m**ixed-effects **b**reakpoint **a**nalysis with **r**obust **e**stimation, especially designed for longitudinal ophthalmic studies. LIMBARE accommodates repeated measurements from both eyes and overtime, and effectively address the presence of outliers.

Methods: The model setup of LIMBARE and computing algorithm for point and confidence interval estimates of the breakpoint was introduced.   The performance of LIMBARE and other competing methods was assessed via comprehensive simulation studies and application to a longitudinal ophthalmic study with 216 eyes (145 subjects) followed for an average of 3.7±1.3 years to examine the longitudinal association between structural and functional measurements.

Results: In simulation studies, LIMBARE showed the smallest bias and mean squared error (MSE) for estimating the breakpoint, with empirical coverage probability of corresponding CI estimate closest to the nominal level for scenarios with and without outlier data points. In the application to the longitudinal ophthalmic study, LIMBARE detected two breakpoints between visual field mean deviation (MD) and retinal nerve fiber layer thickness (RNFL) and one breakpoint between MD and cup to disc ratio (CDR), while the cross-sectional analysis approach only detected one and none, respectively.

Conclusions: LIMBARE enhances breakpoint estimation accuracy in longitudinal ophthalmic studies, while cross-sectional analysis approach is not recommended for future studies.

Translational Relevance: Our proposed method and companion software R package provides a valuable computational tool for advancing longitudinal ophthalmology research and exploring the association relationships between ophthalmic variables.

1 Introduction

Broken stick analysis is a powerful analytical tool for investigating piecewise-linear association between continuous variables, particularly in cases where non-linear associations are present. The goal of the analysis is to accurately detect breakpoints at which the effect size of the association changes, allowing for piecewise-linear association inference to be made before and after these breakpoints. In ophthalmology, broken stick analysis has been used to study the association between structural and functional measurements in glaucoma[1-3], where varying levels of association between structural measurements, such as average RNFL thickness, and functional measurements obtained with visual field tests[4-9] have been observed at different stages of disease severity. These differences may be due to the "floor effect" of structural measurements beyond which no further structural damage can be measured, leading to changes in structural-function association[10].

Current statistical approaches for broken stick analysis in ophthalmology mainly employ cross-sectional approaches where observations from one eye per subject from a single visit are included in the analysis[9]. The approach negates the possibility of maximizing available data by using information obtained from both eyes of the same subject and evaluating longitudinal processes such as disease trajectory. Alternatively, aggregate of measurements across all visits from both eyes are used without considering the inter-eye correlation and within-subject correlation between repeated measurements[8]. Without considering the correlations of repeated measurements from two eyes within a subject, the breakpoint estimation may be misleading.

To address these shortfalls, we propose a new and advanced linear mixed-effects breakpoint analysis with robust estimation (LIMBARE) that can provide accurate broken stick analysis for ophthalmic studies that assess association relationship between measurements longitudinally

sampled from both eyes. Specifically, LIMBARE is built upon the segmented mixed model,

which accounts for repeated measurements overtime and between eyes, and additional

confounders commonly confronted in human studies[11, 12]. LIMBARE further incorporates the

least trimmed squares (LTS) strategy[13, 14] to mitigate the impact of outliers and provide a robust

and accurate point estimate of breakpoints. We also propose a new standard error estimator for

LIMBARE to improve the coverage probability of the confidence interval estimate. With

LIMBARE (method workflow illustrated in Figure 1), the point and confidence interval (CI)

estimates of breakpoints, together with the association effect sizes quantified as slopes of the

piecewise linear mixed effects model are reported.

A simulation study was conducted to evaluate the performance of LIMBARE against

competing methods. We also applied these methods to a longitudinal study examining the

association between visual field mean deviation (MD) and OCT RNFL thickness, and between

MD and cup to disc ratio, to examine the piecewise association relationship. Our results

demonstrate that LIMBARE improves the accuracy of breakpoint estimation, and we recommend

the use of this method in future ophthalmic studies.

2 Methods

87
2.1 Model specification of LIMBARE

We used a segmented mixed-effects model (SMM) to model the longitudinal, non-linear

association between two variables of interest. Following the notation of Muggeo (2003)[11], two

variables, denoted by x_{ijk} and y_{ijk}, are observed from subject i $(= 1, \dots, n)$ at visit time t_{ijk} $(j =$

$1, \dots, n_i)$ for eye k $(=$ "OD", "OS"). The model also includes time-dependent covariates, denoted

by z_{ijk}, such as a constant vector 1, visit time t_{ijk} and other confounders (See Figure 1 for an

illustration of the data input). Our primary goal is to detect unknown yet important breakpoints in

the association between x_{ijk} and y_{ijk}. The SMM regression model is given by Equation (1):

$$y_{ijk} = \xi_0 \cdot x_{ijk} + \sum_{p=1}^{P} \xi_p \cdot \left(x_{ijk} - \varphi_p\right) \cdot I_{(x_{ijk} > \varphi_p)} + \beta^t \cdot z_{ijk} + \gamma_i^t \cdot s_{ij} + \gamma'_{ij} \cdot s'_{ijk} + \varepsilon_{ijk} \quad (1)$$

In this equation, $\Xi = \left(\xi_1, \cdots, \xi_p\right)^t$ is the vector of slope changes between P breakpoints

$\varphi_1, \cdots, \varphi_P$ $(\varphi_1 < \cdots < \varphi_P)$, $\Psi = (\xi_0, \xi_0 + \xi_1, \cdots, \xi_0 + \xi_1 + \xi_2 + \cdots + \xi_P)^t$ is then the vector of

$P + 1$ slopes (association coefficients) for each of the segments, β is the regression coefficients

of the covariates in z_{ijk} (including intercept), and $\gamma_i \sim N\left(0, \Sigma_\gamma\right)$ and $\gamma'_{ij} \sim N\left(0, \Sigma_{\gamma'}\right)$ are the

random effects for covariate s_{ij} and s'_{ijk} respectively. The residual errors ε_{ijk} are assumed to be

independent and normally distributed with mean zero and constant variance σ^2. The model

reduces to a standard linear mixed-effects model when no breakpoint is detected (i.e., $P = 0$).

We estimated the unknown breakpoints $\Phi = (\varphi_1, \cdots, \varphi_P)^t$, and the other unknown parameters

$\Theta = \left(\beta^t, \xi_0, \xi_1, \cdots, \xi_P, \text{Vech}(\Sigma_\gamma), \text{Vech}(\Sigma_{\gamma'}), \sigma^2\right)^t$, along with their corresponding 95%

confidence intervals (CIs), using the algorithms detailed in **Algorithms 1** and **2** below.

2.2 Accurate estimation of breakpoints and other parameters

Point estimation: While the specification of the regression model is straightforward, estimating

the unknown parameters are challenging due to parameter identifiability problem between Φ and

Θ. In the case of the standard linear mixed-effects model, maximum likelihood estimation (MLE)

approach is commonly used to estimate unknown parameters using the derived log-likelihood

function[15]. However, the strategy is not applicable for segmented mixed effects model in

Equation (1) because the corresponding log-likelihood function is not differentiable at $\varphi_p's$.

Moreover, MLE is sensitive to outliers, which are common in ophthalmic studies due to

measurement error, reading error, instrumental error, etc. To overcome these challenges, we

propose a robust estimation approach that allows for the detection of multiple breakpoints,

accounting for potential outliers.

Given a value $\varphi_p^{(s)}$, the linear piece $\left(x_{ijk} - \varphi_p\right) \cdot I\left(x_{ijk} > \varphi_p\right)$ can be approximated via the

first order Taylor expansion around $\varphi_p^{(s)}$, i.e., $\left(x_{ijk} - \varphi_p\right) \cdot I\left(x_{ijk} > \varphi_p\right) \approx \left(x_{ijk} - \varphi_p^{(s)}\right) \cdot$

$I\left(x_{ijk} > \varphi_p^{(s)}\right) + (\varphi_p - \varphi_p^{(s)})(-1)I\left(x_{ijk} > \varphi_p^{(s)}\right)$, where $(-1)I(x_{ijk} > \varphi_p^{(s)})$ can be considered

as the first derivative of $\left(x_{ijk} - \varphi_p\right) \cdot I\left(x_{ijk} > \varphi_p\right)$. This motivates us to solve the estimation

problem by iteratively estimating breakpoints Φ and other parameters Θ. We further introduce

the Least Trimmed Squares (LTS) technique to the iteration algorithm to reduce the impact of

outliers in the data on parameter estimation[16, 17].

Specifically, given a predetermined proportion of outlier data points $\alpha \in (0,1)$, the iterative

procedure for parameter estimation of LIMBARE is given as Algorithm 1 (A1):

Algorithm 1 (A1) Point estimate of parameters for LIMBARE framework

1. Randomly sample $100 \times (1 - \alpha)\%$ observations from the full data set.
2. For m'th iteration:
   a. Set initial values of $\Phi^{(0)} = \left(\varphi_1^{(0)}, \cdots, \varphi_P^{(0)}\right)$;
   b. Given the current values of $\varphi_p^{(s)}, p = 1, \cdots P$, calculate $U_{ijk,p}^{(s)} = (x_{ijk} - \varphi_p^{(s)}) \cdot I_{\left(x_{ijk} > \varphi_p^{(s)}\right)}$ and
      $V_{ijk,p}^{(s)} = -I_{\left(x_{ijk} > \varphi_p^{(s)}\right)}$.
   c. The model can be re-expressed as $y_{ijk} = \xi_0 \cdot x_{ijk} + \sum_{p=1}^{P} \xi_p \cdot U_{ijk,p}^{(s)} + \sum_{p=1}^{P} \eta_p \cdot V_{ijk,p}^{(s)} + \beta \cdot z_{ijk} +$
      $\gamma_i \cdot s_{ij} + \gamma'_{ij} \cdot s'_{ijk} + \varepsilon_{ijk}$. Obtain the MLE estimate of parameter Θ from the model and denote it as
      $\Theta^{(s)}$;
   d. Update φ_p as $\varphi_p^{(s+1)} = \frac{\hat{\eta}_p^{(s)}}{\hat{\xi}_p^{(s)}} + \varphi_p^{(s)}$;
   e. Repeat steps b-d until $\hat{\eta}_p^{(s)} \to 0$. Denote the resulting parameters as $\Phi^{[m]}$ and $\Theta^{[m]}$.
3. Use the m'th estimates and corresponding model to predict the full dataset and calculate the squared
   residuals $(y_{ijk} - \hat{y}_{ijk})^2$ for each observation.
4. Select the $100 \times (1 - \alpha)\%$ observations that have the smallest squared residuals and repeat steps 2-3
   until the sampled observations achieve the smallest squared residuals.

The proposed iterative procedure consists of two layers. The outer layer (Steps 1,3, and 4)

employs the LTS technique to account for the potential impact of outliers on the model fitting.

The inter layer (a-e of Step 2) uses the iterative optimization algorithm of the segmented mixed

model, which we will discuss later in the competing methods section. To determine the trimming

proportion of outliers, we assessed and compared three values of α: 0(no outlier), 0.05, and 0.10

for model fitting and parameter estimation.

Estimation of standard error (SE) and confidence interval (CI): While LTS is effective in

reducing or eliminating the influence of outliers on point estimation, it can lead to

underestimation of the standard errors of the unknown parameters[18, 19]. To address this issue, we

propose Algorithm 2 (A2) to improve the estimates of SEs and CIs of unknown parameters,

including breakpoints and their associated slopes.

Algorithm 2 (A2) SE and 95% CI estimate of LIMBARE framework
1. Denote the observed maximal and minimal residuals from **A1** as ϵ_M and ϵ_m respectively. Estimate standard deviation of the residual error as $\hat{\sigma} = (\frac{\epsilon_M}{\text{invNorm}\left(1-\frac{\alpha}{2}\right)} + \frac{\epsilon_m}{\text{invNorm}\left(\frac{\alpha}{2}\right)})/2$ where invNorm() is the inverse cumulative distribution function of the standard normal distribution.
2. Generate calibrated residual errors from the recovered residual distribution function $\hat{\varepsilon}_{ijk} \sim N(0, \ \hat{\sigma}^2)$.
3. Update the full dataset by adding the calibrated errors to the predicted values: $y_{new,ijk} = \hat{y}_{ijk} + \hat{\varepsilon}_{ijk}$ and re-run step 2 of A1 accordingly.
4. Employ the standard errors (SE) of Φ, Θ from the resulted model as the calibrated SEs.
5. Derive the 95% CI of Φ, Θ with point estimates from A1 $\pm 1.96 \cdot$ SE.

The algorithm approximates the $\alpha/2 \times 100\%$ and $(1 - \alpha/2) \times 100\%$ quantiles of the

distribution of model residuals with the observed maximal and minimum residuals, since the

observed dataset is trimmed with $\alpha \times 100\%$. By recovering of distribution of the residuals, each

observation's residual is calibrated and used as input for the estimation of SE and CI.

Hypothesis tests: To test the null hypothesis that the slope of each regression piece is equal to

zero, we use Wald test statistics calculated from the point and SE estimates of the slope

parameters $\Psi = (\xi_0, \xi_0 + \xi_1, \cdots, \xi_0 + \xi_1 + \xi_2 + \cdots + \xi_P)^t$. Specifically, we divide the point

estimate of the slope by the corresponding SE to obtain the Wald test statistics. This statistics is

then used to examine the null hypothesis for each regression piece.

Figure 1 illustrates the input, algorithm and the output of the proposed method.

2.3 Competing methods

To evaluate the performance of our proposed method, we compared it with three alternative

methods commonly used for estimating breakpoints in longitudinal studies. The first method is

the segmented mixed model (**SMM**) proposed by Muggeo et al[11, 20]. SMM uses an iterative

optimization algorithm to estimate the model parameters, but does not account for outlier data

points. Therefore, we also evaluated the effect of outliers on the estimates of breakpoints

obtained with this method. The second method is the robust mixed-effects segmented model

(**RMESR**) proposed by Zhou et. al.(2020)[19]. RMESR combines the estimates from both

segmented linear models and linear mixed-effects models, with the breakpoint estimated based

on the segmented linear model even though the data comes from a longitudinal design. To

accommodate outliers in the dataset, RMESR uses the least trimmed squares technique. The third

method is the conventional method, named **CROSS**, in which longitudinal measurements are

aggregated cross-sectionally and analyzed with a segmented linear regression model[11]. The

detailed description of each method can be found in **Appendix A** of the **Supplementary**

material.

2.4 Simulation setup

Our simulation was designed to imitate the structural and functional data from real longitudinal

ophthalmic studies. For both Scenario 1 (without outliers) and 2 (with outliers), we simulated

two breakpoints, φ_1 and φ_2, for the variable x (referred to as the "breakpoint variable" hereafter).

We also included a time variable, t, to record the time from baseline visit in years, and

considered random effects between visits of the same subject and between eyes within subjects.

To generate the outcome variable y, we used the following model:

$$y_{ijk} = \xi_0 \cdot x_{ijk} + \xi_1\big(x_{ijk} - \varphi_1\big)I\big(x_{ijk} > \varphi_1\big) + \xi_2\big(x_{ijk} - \varphi_2\big)I\big(x_{ijk} > \varphi_2\big) + \beta_0 + \beta_1 \cdot t_{ijk}$$

$$+ r_i \, \text{ID}_i + r'_{ij}\text{Eye:ID}_{ij} + \varepsilon_{ijk}$$

Here, the start slope and slope changes (ξ_0, ξ_1, ξ_2) were set to $(1, 2, 3)$, and the breakpoints

(φ_1, φ_2) were set to $(-10, 2)$, while the intercept the time slope (β_0, β_1) were set to $(50, -0.5)$.

We set x_{ijk} to following a normal distribution $N(-4, 6^2)$, and t_{ij} to take values $(0, 1, 2, 3)^t$. The

random effects r_i, r'_{ij}, and ε_{ijk} were generated from normal distributions with means 0 and

variances 1, 4, and 0.25, respectively. We simulated data from 3-4 visits and from two eyes for

60% of the total of 300 subjects. We conducted N = 1000 replications to evaluate the

performance of our proposed method, LIMBARE, and compared it with the SMM, RMESR, and

CROSS methods.

To simulate outliers under Scenario 2, we randomly selected 5% of observations from the

simulated dataset and added $(\max(y) - \min(y))/4$ to the outcome variable y and $(\max(x) -$

$\min(x))/4$ to the breakpoint variable x.

We used the finite sample bias and mean squared error (MSE), and the empirical coverage

probability (ECP), as comparison metrics for assessing the point and confidence interval

estimates of the model parameters, respectively.

2.5 The longitudinal ophthalmic study

This longitudinal clinical study examined 216 eyes (163 open angle glaucoma, 44 glaucoma

suspects, and 9 healthy eyes) from a cohort of 145 subjects aiming to investigate changes in

ocular structure and function over time. The study was approved by the IRB of the University of

Pittsburgh and NYU Langone Health, and all participants provided informed consent[8]. Each

subject underwent comprehensive ophthalmic examinations including qualified visual fields,

OCT optic nerve head (ONH) and macular scans, in at least 5 visits over an average period of 3.7

years (**Table 1**). We evaluated the piecewise association between visual field mean deviation

(MD) and two structural measures: retinal nerve fiber layer (RNFL) thickness and cup to disc

ratio (CDR). The proportion of data trimming α was set to 0.05 for the MD-RNFL association,

and 0.1 for the MD-CDR association. To control for potential confounding factors, such as time

from baseline visit (in year), age at baseline, signal strength of scans, and eye, all break-point

analysis models were adjusted accordingly. We used Bayesian information criterion (BIC) to

determine the optimal number of breakpoints for each association and each method. The model

with the smallest BIC was selected, and corresponding estimated breakpoints, slopes were

reported.

3. Results

3.1 Simulation results

We present the estimation performance of all assessed methods under two scenarios, including

mean value, empirical bias, and MSE for point estimates, mean standard errors (SEs) and

empirical coverage probability (ECP) for 95% confident interval estimates of the detected

breakpoints in Error! Reference source not found.**2** and Error! Reference source not found.**3**.

Additionally, **Table S1** and **Table S2** in **Appendix B** of the **Supplementary material**

demonstrate the estimation performance for the estimates of start slope and slope changes for all

assessed methods.

Under the first scenario, which involves synthetic datasets without outliers (**Error! Reference**

source not found.2 and **S1**), the proposed method LIMBARE and SMM are statistically

efficient, as indicated by their negligible bias and small MSE for point estimates. In comparison,

the point estimate of RMESR and CROSS has a larger bias and MSE than other methods. The CI

estimates of LIMBARE and SMM have satisfactory ECPs close to the nominal level of 95%.

However, the CI estimates of RMESR and CROSS tend to have much lower ECPs than nominal.

The under-coverage of RMESR is mainly due to the underestimation of SEs, while the poor

performance of the CI estimates of CROSS is largely driven by the bias of its point estimate.

Under the second scenario, which involves synthetic datasets with outliers (**Error! Reference**

source not found.3 and **S2**), the proposed method LIMBARE has the markedly smallest bias

and MSE for point estimates, indicating its statistical efficiency and superior robustness to

outliers. The point estimates of all competing methods have a large bias and MSE, indicating the

failure to accurately detect breakpoints and estimate slopes under this scenario. The ECPs of all

competing methods are substantially lower than nominal, while LIMBARE maintains ECP

closest to the nominal. However, we detected a slight over-coverage of the CI estimate of

LIMBARE.

3.2 Broken stick analysis of the longitudinal ophthalmic study

We conducted a piecewise association analysis between visual field mean deviation (MD) and

RNFL thickness, as well as MD and CDR, using four different methods: LIMBARE, SMM,

RMESR, and CROSS. **Figures 2** and **3**, along with **Tables 4** and **5**, present the results of these

analyses.

For the MD-RNFL association (**Figure 2** and **Table 4**), LIMBARE, SMM, and RMESR

consistently identified two breakpoints, while CROSS detected only one. Specifically,

LIMBARE indicated that RNFL thickness decreases significantly and rapidly at a rate of 1.9

[1.52,2.29] μm/dB before reaching the breakpoint of -4.31 dB (from top right to bottom left in

Figure 2, which represents the disease spectrum from mild to severe glaucoma). After that,

RNFL thickness still significantly decreases but at a slower rate of 1.05 [0.81,1.29] μm/dB until

the next breakpoint of -15.70 dB. The association between MD and RNFL approaches a plateau

with insignificant rate of change when MD is smaller than -15.70 dB. In contrast, CROSS

captured only one breakpoint at MD= -4.08 dB, which is inconsistent with the other three

methods. The estimated slope from CROSS indicated that RNFL thickness decreases

significantly with MD when MD is larger than -4.08 dB, which is not typically observed

clinically. Noticeably, RMESR reported the narrowest confidence interval estimates for both

breakpoints and slopes, which suggests the possibility of under-coverage of the ground truth of

these parameters (**Table 4**).

For the MD-CRD association (**Figure 3** and **Table 5**)**,** LIMBARE, SMM, and RMESR identified

one breakpoint while Cross detected none. LIMBARE and RMESR, which were designed to

accommodate outliers, reported breakpoints close to each other at -8.61 dB and -9.83 dB,

respectively. The corresponding slope estimates from LIMBARE and RMESER indicated that

CDR increases at the rate of 0.0131 [0.0107,0.0155] per dB (from right to left of Figure 3) and

0.0099 [0.0089,0.0109] per dB before MD decreases to the breakpoint. After that, the increasing

trend slows down with a rate of 0.0023[0.0008,0.0038] per dB and 0.0022[0.0014,0.0030] per

dB. Yet, RMESR reported much narrower confidence interval estimates for both breakpoints and

slopes. In comparison, SMM showed that CDR increases at the rate 0.0159 [0.0140,0.0178] per

dB when MD decreases before reaching the breakpoint of -12.46 dB. After the breakpoint,

CDR decreases at the rate 0.0019 [-0.0011,0.0048] per dB as MD declines. The patterns

identified by LIMBARE and RMESR complied with clinical observation, while SMM showed a

trend that is less frequently observed. RMESR resulted the narrowest confidence interval

estimates compared with LIMBARE.

Overall, the results suggest that LIMBARE is a robust method for identifying breakpoints in the

association between MD and RNFL thickness or MD and CDR by providing both point and

confidence interval estimates. Conventional methods like CROSS may miss these breakpoints,

leading to inaccurate or incomplete understanding of the relationship between these variables.

The identified breakpoints and slopes may provide valuable clinical insights into the study of

glaucoma.

4. Conclusion

In this manuscript, we proposed a novel advanced linear mixed-effects breakpoint analysis with

robust estimation (LIMBARE) for longitudinal data analysis. The proposed method provides

efficient point estimation of breakpoints and other association parameters, largely mitigating the

impact of outliers, and improving the estimation of confidence intervals for both breakpoints and

slopes. Simulation studies show that LIMBARE outperforms three competing methods in terms

of negligible bias, smallest mean squared error, and empirical coverage probabilities closest to

the nominal level of 95% under the scenarios with and without outliers.

In the application to broken stick analysis to a longitudinal clinical study, the piecewise

associations between MD and RNFL thickness estimated by LIMBARE, SMM, and RMESR

were comparable and consistent with clinical observations. The piecewise associations between

MD and CDR estimated by LIMBARE and RMESR also complied with clinical observation,

while SMM exhibited a less likely clinical pattern. The results of CROSS, however, were

inconsistent with three other methods.

In conclusion, LIMBARE is recommended for future longitudinal ophthalmic studies as it

exhibited the most favorable results in both simulation scenarios and illustrated patterns that

comply with clinical observation. Furthermore, a user-friendly R package "LIMBARE" is

publicly available at https://github.com/JiyuanHu/. The conventional cross-sectional method is

not capable of capturing breakpoints in longitudinal studies and its use is not recommended.

Tables and Figures

Table 1. Characteristics of the longitudinal clinical study. This table presents the demographic

and clinical characteristics of the study population.

Parameters	Ns or Mean ± standard deviation (SD)
Eyes/subjects	216/145
N visits during f/u	5.8±0.9
N observations	1221
Age (years)	69.7±10.6
follow up (years)	3.7±1.3
ONH signal strength	8.3±0.8
baseline RNFL (μm)	73.8±14.4
baseline MD (dB)	-4.23±6.29
baseline CDR	0.67±0.15

293 **Table 2:** Simulation results for Scenario 1: synthetic dataset without outliers (breakpoints at -10

294 and 2). The table shows comparisons of point estimates, standard error estimates, and confidence

295 interval estimates of breakpoints among different statistical methods.

	$\varphi_1 = -10$					$\varphi_2 = 2$				
	Est	Bias	MSE	SE	ECP	Est	Bias	MSE	SE	ECP
LIMBARE	-10.001	-5E-04	0.0010	0.0291	0.923	2.0000	<0.0001	0.0005	0.0193	0.928
SMM	-10.001	-7E-04	0.0008	0.0284	0.943	1.9999	-0.0001	0.0004	0.0189	0.942
RMESR	-10.038	-0.038	0.0027	0.0189	0.485	2.0340	0.0340	0.0017	0.0126	0.349
CROSS	-5.9814	4.0186	19.7530	0.8238	0.141	-1.6171	-3.6171	16.5240	0.8236	0.141

Est, bias and MSE represents the mean value, bias and mean squared error of the point estimate,
respectively. SE represents mean value of the standard error estimate of the breakpoint; ECP
represents the empirical coverage probability of the 95% confidence interval estimate.

Table 3: Simulation results for Scenario 2: synthetic dataset with outliers (breakpoints at -10 and

2). The table shows comparisons of point estimates, standard error estimates, and confidence

interval estimates of breakpoints among different statistical methods.

| | $\varphi_1 = -10$ | | | | | $\varphi_2 = 2$ | | | | |
	Est	Bias	MSE	SE	ECP	Est	Bias	MSE	SE	ECP
LIMBARE	-9.9963	0.0037	0.0817	0.0705	0.996	2.0080	0.0080	0.1169	0.0662	0.998
SMM	-10.336	-0.336	7.2465	0.8715	0.747	1.7874	-0.2126	4.8063	0.7424	0.778
RMESR	-10.073	-0.073	1.2084	0.0197	0.597	1.9793	-0.0207	0.6337	0.0139	0.567
CROSS	-5.5309	4.4691	28.4270	1.2464	0.218	-1.9612	-3.9612	22.8790	1.7822	0.231

304 Est, bias and MSE represents the mean value, bias and mean squared error of the point estimate,
305 respectively. SE represents mean value of the standard error estimate of the breakpoint; ECP
306 represents the empirical coverage probability of the 95% confidence interval estimate.
307

309 **Table 4.** Parameter estimates [95% CIs] for the broken stick association analysis between MD

310 and RNFL. Breakpoints are reported as dB and slopes as µm/dB.

LIMBARE	Breakpoints	-15.70 [-17.86, -13.53]		-4.31 [-6.19, -2.42]
	Slopes	-0.21 [-0.63, 0.22]	1.05* [0.81, 1.29]	1.90* [1.52, 2.29]
SMM	Breakpoints	-15.73 [-17.87, -13.58]		-4.93 [-6.25, -3.62]
	Slopes	-0.16 [-0.51, 0.18]	0.95* [0.70, 1.2]	2.19* [1.90, 2.48]
RMESR	Breakpoints	-15.64 [-16.95, -14.34]		-5.47 [-7.16, -3.78]
	Slopes	-0.19 [-0.42, 0.05]	1.04* [0.87, 1.21]	1.64* [1.40, 1.87]
CROSS	Breakpoint	-4.08 [-6.26, -1.89]		
	Slopes	0.48 [-0.03, 0.98]		3.91* [1.93, 5.90]

311 * p-value <0.05 for the statistical test against the null hypothesis of slope equal to zero.

312 **Table 5**. Parameter estimates [95% CIs] for the broken stick association analysis between MD
313 and CDR. Breakpoints are reported as dB and slopes as cup-to-disc-ratio/dB.

LIMBARE	Breakpoints	-8.61 [-10.09, -7.14]	
	Slopes	-0.0023* [-0.0038, -0.0008]	-0.0131* [-0.0155, -0.0107]
SMM	Breakpoint	-12.46 [-13.85, -11.07]	
	Slopes	0.0019 [-0.0011, 0.0048]	-0.0159* [-0.0178, -0.014]
RMESR	Breakpoint	-9.83 [-10.69, -8.98]	
	Slopes	-0.0022* [-0.0030, -0.0014]	-0.0099* [-0.0109, -0.0089]
CROSS	Slope (breakpoint not detected)	-0.0109* [-0.0141, -0.0077]	

314 * p-value <0.05 for the statistical test against the null hypothesis of slope equal to zero.

Figure 1. Schematic framework of the linear mixed-effects breakpoint analysis with robust

estimation (LIMBARE) method. Data input: the input data is in long format and includes values

of the variables of interest (x, y), time (t) and other confounders such as age. LIMBARE: The

LIMBARE comprises two parts: Algorithms A1 and A2 are used in each part to detect outliers

and obtain robust point (Part 1) and 95% confidence interval (CI) (Part 2) estimates of

breakpoints, association slopes, and other unknown parameters. Applications: the output is the

piecewise association inference between longitudinal variables, which can be applied to identify

changes in association patterns for the study of interested variables.

Figure 2. Estimated piecewise associations between visual field mean deviation and RNFL

thickness and from all assessed methods.

336 **Figure 3.** Estimated piecewise associations between visual field mean deviation and cup to disc

337 ratio from all assessed methods.

Reference

1.	Shah NN, Bowd C, Medeiros FA, et al. Combining Structural and Functional Testing for Detection of Glaucoma. *Ophthalmology* 2006;113:1593-1602.
2.	Harwerth RS, Wheat JL, Fredette MJ, Anderson DR. Linking structure and function in glaucoma. *Progress in Retinal and Eye Research* 2010;29:249-271.
3.	Johnson CA, Cioffi GA, Liebmann JR, Sample PA, Zangwill LM, Weinreb RN. The Relationship Between Structural and Functional Alterations in Glaucoma: A Review. *Seminars in Ophthalmology* 2000;15:221-233.
4.	Alasil T, Wang K, Yu F, et al. Correlation of retinal nerve fiber layer thickness and visual fields in glaucoma: a broken stick model. *Am J Ophthalmol* 2014;157:953-959.
5.	Liu WW, McClurkin M, Tsikata E, et al. Three-dimensional Neuroretinal Rim Thickness and Visual Fields in Glaucoma: A Broken-stick Model. *J Glaucoma* 2020;29:952-963.
6.	Murphy MC, Conner IP, Teng CY, et al. Retinal Structures and Visual Cortex Activity are Impaired Prior to Clinical Vision Loss in Glaucoma. *Sci Rep* 2016;6:31464.
7.	Park K, Kim J, Lee J. The Relationship Between Bruch's Membrane Opening-Minimum Rim Width and Retinal Nerve Fiber Layer Thickness and a New Index Using a Neural Network. *Transl Vis Sci Technol* 2018;7:14.
8.	Ramos Cadena MdlA, Wollstein G, Lucy K, et al. Disease Severity Threshold (Tipping Point) at which RNFL and GCIPL Glaucoma Progression Rates Fall. *Investigative Ophthalmology & Visual Science* 2018;59:3501-3501.
9.	Wollstein G, Kagemann L, Bilonick RA, et al. Retinal nerve fibre layer and visual function loss in glaucoma: the tipping point. *Br J Ophthalmol* 2012;96:47-52.
10.	Sung MS, Heo H, Park SW. Structure-function relationship in advanced glaucoma after reaching the RNFL floor. *Journal of Glaucoma* 2019;28:1006-1011.
11.	Muggeo VM. Estimating regression models with unknown break-points. *Stat Med* 2003;22:3055-3071.
12.	Muggeo VM, Atkins D, Gallop R, Dimidjian S. Segmented mixed models with random changepoints: A maximum likelihood approach with application to treatment for depression study. *Statistical Modeling* 2014;14:293-313.
13.	Rousseeuw P. Multivariate Estimation With High Breakdown Point. *Mathematical Statistics and Applications* 1985;B:283-297.
14.	Rousseeuw PJ, Leroy AM. *Robust regression and outlier detection*: John wiley & sons; 2005.
15.	Jiang J, Nguyen T, SpringerLink (Online service). *Linear and generalized linear mixed models and their applications*. Second edition. ed; 2007:1 online resource (xiv, 343 pages).
16.	Dhakal CP. DEALING WITH OUTLIERS AND INFLUENTIAL POINTS WHILE FITTING REGRESSION. *Journal of Institute of Science and Technology* 2017;22:61-65.
17.	Ozsahin DU, Mustapha MT, Mubarak AS, Ameen ZS, Uzun B. Impact of Outliers and Dimensionality Reduction on the Performance of Predictive Models for Medical Disease Diagnosis. *2022 International Conference on Artificial Intelligence in Everything (AIE)*; 2022:79-86.
18.	Peddada SD, Haseman JK. Analysis of nonlinear regression models: a cautionary note. *Dose-response* 2005;3:dose-response. 003.003. 005.

383   19.   Zhou X, Gao Y, Yao W, Yu N. A Robust Segmented Mixed Effect Regression Model for
384   Baseline Electricity Consumption Forecasting. *Journal of Modern Power Systems and Clean*
385   *Energy* 2020;10:71-80.
386   20.   Muggeo VM. Regression Models with Break-Points / Change-Points Estimation. 2022.
387
388
389

Data Input

Long format of longitudinal dataset for repeated measurements

y: outcome of interest;
x: breakpoint variable;
time: time from baseline;
age: an example of other covariates.

Subject_ID	Eye	y	x	time	age
3	OD	74.15	4.97	0	67
3	OD	73.63	4.75	0.8	67
3	OD	40.28	-10.03	1.6	67
3	OS	61.47	0.99	0	67
3	OS	47.73	-3.69	0.8	67
3	OS	57.29	0.07	1.6	67
4	OD	38.49	-13.06	0	76
4	OD	59.14	0.54	0.88	76
⋮	⋮	⋮	⋮	⋮	⋮

LIMBARE

Part 1: Breakpoints and slopes estimations, and outliers detection

Part 2: 95% CI estimates for breakpoints and slopes

Trim large residuals

Distribution of residuals

Distribution of trimmed residuals

The standard error of residuals were underestimated due to trimming the large residuals during RSMM

Distribution of calibrated residuals

Recover the standard error using Algorithm 2

Applications