1 **Early life exposure to broccoli sprouts confers stronger protection against enterocolitis**
2 **development in an immunological mouse model of inflammatory bowel disease.**
3

4 Lola Holcomb[1$], Johanna M. Holman[2$], Molly Hurd[3], Brigitte Lavoie[3], Louisa Colucci[4], Benjamin
5 Hunt[5], Timothy Hunt[5], Marissa Kinney[2], Jahnavi Pathak[1], Gary M. Mawe[3], Peter L. Moses[3,6],
6 Emma Perry[7], Allesandra Stratigakis[8], Tao Zhang[8], Grace Chen[9], Suzanne L. Ishaq[1*], Yanyan
7 Li[1*]
8
[1] Graduate School of Biomedical Sciences and Engineering, University of Maine, Orono, Maine,
USA 04469
[2] School of Food and Agriculture, University of Maine, Orono, Maine, USA 04469
[3] Larner College of Medicine, University of Vermont, Burlington, Vermont, USA 05401
[4] Department of Biology, Husson University, Bangor, Maine, USA 04401
[5] Department of Biology, University of Maine, Orono, Maine, USA 04469
[6] Finch Therapeutics, Somerville, Massachusetts, USA 02143
[7] Electron Microscopy Laboratory, University of Maine, Orono, Maine, USA 04469
[8] School of Pharmacy and Pharmaceutical Sciences, SUNY Binghamton University, Johnson
City, New York, USA 13790
[9] Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, Michigan,
USA 48109
21
22
23 [$] these authors contributed equally.
24
25 **\*Corresponding authors:**
26 Suzanne L. Ishaq, University of Maine, School of Food and Agriculture, Orono, ME 04469,
27 sue.ishaq@maine.edu, 1-207-581-2770
28 Yanyan Li, University of Maine, School of Food and Agriculture, Orono, ME 04469,
29 yanyan.li@maine.edu, 1-207-581-3134

30
31

32 **Keywords**: Crohn's Disease, cruciferous vegetables, sulforaphane, glucoraphanin, gut
33 microbiota, dietary bioactives, 16S rDNA, interleukin-10 knockout

34
35

**Abstract**

Crohn's Disease (CD) is a presentation of Inflammatory Bowel Disease (IBD) that

manifests in childhood and adolescence, and involves chronic and severe enterocolitis, immune

and gut microbiome dysregulation, and other complications. Diet and gut-microbiota-produced

metabolites are sources of anti-inflammatories which could ameliorate symptoms. However,

questions remain on how IBD influences biogeographic patterns of microbial location and

function in the gut, how early life transitional gut communities are affected by IBD and diet

interventions, and how disruption to biogeography alters disease mediation by diet components

or microbial metabolites. Many studies on diet and IBD use a chemically induced ulcerative

colitis model, despite the availability of an immune-modulated CD model. Interleukin-10-

knockout (IL-10-KO) mice on a C57BL/6 background, beginning at age 4 or 7 weeks, were fed a

control diet or one containing 10% (w/w) raw broccoli sprouts, which was high in the sprout-

sourced anti-inflammatory sulforaphane. Diets began 7 days prior to, and for 2 weeks after

inoculation with *Helicobacter hepaticus,* which triggers Crohn's-like symptoms in these immune-

impaired mice. The broccoli sprout diet increased sulforaphane in plasma; decreased weight

stagnation, fecal blood, and diarrhea associated; and increased microbiota richness in the gut,

especially in younger mice. Sprout diets resulted in some anatomically specific bacteria in

younger mice, and reduced the prevalence and abundance of pathobiont bacteria which trigger

inflammation in the IL-10-KO mouse, for example; *Escherichia coli* and *Helicobacter*. Overall,

the IL-10-KO mouse model is responsive to a raw broccoli sprout diet and represents an

opportunity for more diet-host-microbiome research.

**Importance**

To our knowledge, IL-10-KO mice have not previously been used to investigate the

interactions of host, microbiota, and broccoli, broccoli sprouts, or broccoli bioactives in resolving

symptoms of CD. We showed that a diet containing 10% raw broccoli sprouts increased the

plasma concentration of the anti-inflammatory compound sulforaphane, and protected mice to

varying degrees against negative disease symptoms, including weight loss or stagnation, fecal

blood, and diarrhea. Younger mice responded more strongly to the diet, further reducing

symptoms, as well as increased gut bacterial community richness, increased bacterial

community similarity to each other, and more location-specific communities than older mice on

the diet intervention. Crohn's Disease disrupts the lives of patients, and requires people to alter

dietary and lifestyle habits to manage symptoms. The current medical treatment is extremely

expensive, and a dietary intervention represents an affordable, accessible, and simple strategy

to reduce the burden of symptoms.

**Introduction**

Inflammatory Bowel Diseases (IBD) affect over 6 million people globally, with more than

25% of cases reported in the United States (1). The varied combination of symptoms, the

intensity of their presentation, and their multifactorial origin make symptoms difficult (2) and

extremely expensive for patients to manage (3), and detracts from quality of life (4, 5). Crohn's

Disease (CD) is one of the primary immune-disordered presentations of IBD, which typically

presents during late adolescence and early adulthood (6), has strongly associated genetic (7)

and environmental (8, 9) risk factors, and involves a loss of function of the innate immune

system which disrupts host-microbial interactions in the gastrointestinal (GI) tract (10). CD

treatments attempt to suppress the immune response to alleviate inflammation as a method of

reducing various symptoms, and to return patients to as close to homeostasis as possible.

However, some patients (11) and mouse models (12) respond poorly to single-strategy

treatments. Diet can play an important, economical, and accessible role in the prevention and/or

management of IBD as a source of anti-inflammatory metabolites (13, 14), and broadly for

influencing the robustness of the gut microbiome (15–17). However, there are gaps in

knowledge about how IBD affects gut microbial ecology, including taxonomic and metabolic

biogeography (18–20), how gut microbiota mediate response to disease (21) or diet (22, 23),

and how dietary components and microbial metabolites improve symptoms (24, 25).

<u>Inflammation in CD is severe and disrupts host-microbial interactions</u>

Inflammation in CD is chronic, relapsing, occurs throughout the GI tract, and is

complicated to treat because it involves several breakdowns of the innate immune system (10,

26): decreased expression of the MUC1 gene reduces coverage of mucin in the ileum (27) and

reduces efficacy of tight junction proteins (28), which can allow microbial translocation from the

gut to other tissues (10, 26). Additionally, poor absorption of bile salts in the ileum then causes

damage to the colon (29). CD patients mount weak acute inflammation and low neutrophil

counts in response to infection (10), and may be unable to clear infections. This results in

infiltration of fecal material through the mucosal lining of the gut, dysregulation of the adaptive

immune response, and results in chronic inflammation (10, 30). The adaptive immune response

in CD primarily involves excessive recruitment of effector T-cells (Th1 and Th17) by

inflammatory cytokines (interleukins -12, -18, and -23) which are upregulated in CD lesions (6,

31, 32).

The duration of inflammation, damage to intestinal walls, and other negative outcomes

associated with CD increase the risk of developing gastrointestinal cancers, such as colorectal

cancer (33). While CD symptoms can develop and present at any age, they are most commonly

observed in individuals between the ages of 15 and 30, and approximately 20-30% of cases

occur in children and adolescents under the age of 18 (6, 8). Children and adolescents

diagnosed with CD encounter distinct challenges, as their gut microbiome is disrupted during

critical periods of anatomical, immunological, and microbiological development (34–36).

Children experience additional complications such as malnutrition, anemia, impaired growth and

development, psychological disorders, bone demineralization, and delayed puberty (37). Dietary

treatments for CD could ameliorate age-related aspects of disease (37).

Diet can be a source of anti-inflammatories to complement medical strategies

Diets which are high in raw, cooked, or fermented cruciferous vegetables can reduce

inflammation and cancer risk in the gut of many people (14, 38), in part because these

vegetables contain sulfur-containing glucosinolates that can be metabolized to other beneficial

compounds (39, 40). Isothiocyanates are a class of bio-active compounds derived from

glucosinolates, which can reduce inflammation including in IBD patients (14, 40, 41).

Sulforaphane (SFN), the most well-studied isothiocyanate, is produced from the glucosinolate

glucoraphanin (GLR), and inhibits the immune factor NF-kB and downregulates multiple

inflammatory signaling pathways *ex vivo* and *in vivo* (14, 41–44).

Purified GLR or SFN have been demonstrated to protect against chemical-induced

ulcerative colitis in mice (38, 45–47) and some pathogenic bacteria in the gut (48, 49). However,

purified SFN is unstable (50, 51), and pure GLR induces extremely variable rates of conversion

in humans (40, 51). The conversion of GLR to SFN is achieved by plant-sourced enzymes when

raw vegetables are chewed or chopped, and by microbially sourced enzymes in the gut

especially if vegetables are cooked to heat-inactivate plant enzymes (38, 46, 49). GLR is

concentrated in broccoli and especially immature sprouts, and a whole-food strategy using

broccoli sprouts increases gut microbial richness, increases sulforaphane in colon tissues and

plasma, and reduces inflammation in chemically induced models of colitis (38, 45, 46). Previous

diet research used mouse models lacking the complex immune-dysfunction component of CD

(52, 53). Research must elucidate how the immune system of CD patients will react to diet-

induced microbiota changes (24, 54), as CD patients have historically been advised to avoid

cruciferous vegetables (54) and sulfur-rich diets (55).

CD disrupts microbial community development from adolescence to early adulthood

The composition of the gut microbiota changes from transitional microbial communities

in childhood to a more stable community in adulthood (16), with distinct variations in the

composition of gut microbiota between people who are healthy or have CD. For example,

pediatric patients exhibit decreased bacterial α-diversity and β-diversity compared to healthy

children (34–36)), and while no specific gut microbiota alterations were consistently reported, a

gain in *Enterococcus* and a significant decrease in *Anaerostipes, Blautia, Coprococcus,*

*Faecalibacterium, Roseburia, Ruminococcus*, and *Lachnospira* have been noted (34–36). There

are also variations across different age groups with CD, from infants through elderly patients

(56–58). Older patients tend to exhibit a higher prevalence of colitis, whereas younger patients

are more likely to present with ileal disease (59), which may result in age-specific and location-

specific changes to gut microbiota. A more thorough understanding of microbial location in the

gut (biogeography) may provide more insight into the interactions between host and microbiota

in the development of CD.

Diet studies using immunological models of IBD are lacking

The interleukin 10 (-/-) knockout (IL-10-KO) mouse is commonly used as a non-

chemical, genetic model well-suited to studying the immune factors, inflammation, and

microbiota of CD (60, 61). Interleukins (IL) are signaling proteins that regulate inflammatory and

immune responses and mediate host-microbial tolerance (62). IL-10 also stimulates the growth

and differentiation of numerous cell types, suppresses macrophage activation, inhibits

inflammatory cytokine production, displays multiple mechanisms of control of Th1 cells (63), and

influences the innate immune response to microorganisms without which there is breakdown of

host-microbial relations in IBD (64). IL-10-KO are raised in pathogen-free conditions and

develop chronic enterocolitis upon exposure to microorganisms which act commensally in

immune-competent mice, and their response resembles the transmural inflammation of CD,

complete with the formation of granulomas, crypt abscess, mucosal hyperplasia, as well as

aberrant immune cell response (65, 66).

To our knowledge, IL-10-KO mice have not previously been used to investigate the

interactions of host, microbiota, and broccoli, broccoli sprouts, or broccoli bioactives in reducing

inflammation, modifying the immune response, and supporting GI tract microbial systems in CD

patients. The objective of this study was to evaluate IL-10-KO mice as a model for studying the

broccoli sprout bioactives. We hypothesized that a medium-high concentration (10% w/w) of raw

broccoli sprouts in the diet, which contains both the glucoraphanin (GLR) precursor and the anti-

inflammatory byproduct SFN, would protect mice from the effects of inflammation triggered by

microbial conventionalization even in a host with disrupted responses to commensal bacteria.

We hypothesized that sprouts would alter the gut microbiota and in location-specific patterns,

and increase the abundance of potentially beneficial taxa while reducing the abundance of

putative pathogens.

It is standard to conventionalize IL-10-KO at approximately 8 weeks of age (67), to

mimic adolescence and typical timing for the onset of symptoms. In conducting two replications

of a diet trial, logical constraints during the pandemic resulted in the use of mice beginning the

diet at 4-weeks (~1 week after weaning (68)) and 7-weeks of age, and conventionalized at 5

and 8 weeks, respectively. We observed a previously unreported effect of young age on the

effectiveness of the response to broccoli sprouts in the diet in resolving the symptoms of

enterocolitis, and added an *a posteriori* hypothesis that younger mice with transitional gut

microbial communities might be more amenable to changes from this diet (68, 69).

**Results**

Raw broccoli sprouts alleviated disease characteristics of immune-modulated enterocolitis

The experimental design was structured as a prevention paradigm (Figure 1) in which

specific-pathogen-free, homozygous IL-10-KO mice were given *ad libitum* either a standard

chow diet (5LOD irradiated) or the treatment diet consisting of 10% (w/w) raw broccoli sprouts,

which were balanced for micronutrients and fiber. Mice were acclimated to diets for 7 days,

continuing for a further 16 days during the induction of colitis via microbial colonization and

symptom onset. Raw broccoli sprouts contain concentrated amounts of GLR, some of which is

metabolized to SFN by mastication, or in this case, diet preparation, when the broccoli enzyme

myrosinase is released from tissues (70). The control diet contained no GLR or SFN, and the

broccoli sprout diet contained on average 4 µg of GLR and 85 µg of SFN per gram of diet

sampled.

Trial 1 included 9 mice (n = 5 in the treatment group and 4 in the control group) starting

the diet at 4 weeks of age, and being exposed to diverse microbiota at 5 weeks of age. Trial 2

included 11 mice (n = 5 in the treatment group and 6 in the control group) starting the diet at 7

weeks of age, and being exposed at 8 weeks which is the standard age for inducing symptoms

in IL-10-KO microbiota trials. The younger mice consuming the broccoli sprout diet continued to

gain weight once enterocolitis was induced compared to their baseline weight, and the younger

mice fed the control diet plateaued around 120% of body weight (Figure 2A, ANOVA $p < 0.05$).

The older mice had a similar weight stagnation during enterocolitis regardless of diet (Figure

2B).

Consumption of the broccoli diet significantly attenuated the development of enterocolitis

in IL-10-KO mice regardless of age group, as indicated by lower Disease Activity Index (DAI)

scores (Figure 2, ANOVA $p < 0.05$). DAI scores were calculated based on three characteristics:

fecal blood (presence/absence), fecal blood severity, and fecal consistency

(firm/loose/diarrhea). The higher the DAI score, the more severe the disease was.

Lipocalin (LCN2), a neutrophil protein that binds bacterial siderophores, serves as a

biomarker for intestinal inflammation (71). Analysis of fecal samples from both age groups on

the day of microbial exposure for conventionalization (D0), appearance of symptoms (D10), and

end of the trial (D15/16) revealed a markedly lower LCN2 concentration in the group of mice fed

the 10% raw broccoli sprouts diet compared to the control diet at both D10 and D15/16 (Figure

3A, ANOVA $p < 0.05$). This was confirmed by serum lipocalin on D15/16, which showed a

marked decrease in LCN2 concentration in the group of mice fed the 10% raw broccoli sprouts

diet compared to the control diet (Figure 3B, ANOVA $p < 0.05$). Similarly on D15/16, pro-

inflammatory cytokines IL-1β, IL-6, and TNF-α were significantly reduced in the plasma of mice

consuming broccoli sprouts (Figure 3C, ANOVA $p < 0.05$). CCL4 (known as macrophage

inflammatory protein 1β (MIP1β)) which regulates the migration and activation of immune cells

during inflammatory processes was not different in younger mice and not evaluated in older

mice.

Tissue from the ileum, proximal colon, and distal colon were collected for histological

scoring (0 = no signs, to 2 = significant signs) of epithelial damage, architectural changes,

infiltration of mononuclear cells into the lamina propria, infiltration of polymorphonuclear cells in

the lamina propria or into the epithelium, as well as abscesses, ulcers, erosion, and branched

crypts which were scored together (Figure S1). Diet was not a significant factor for most scoring

criteria, even when data were subset by trial/age, or subset by organ. The exception to this was

infiltration of mononuclear cells into the lamina propria, which was significantly lower (linear

regression model (lm), $p = 0.02$) in the younger broccoli sprout mice compared to control mice

in trial 1.

SFN was absorbed from the gut by the mice consuming broccoli sprouts in both age

groups, and was found in high concentrations in their plasma, while control mice exhibited no

circulating SFN (Figure 3D, ANOVA $p < 0.05$).

<u>Age significantly influenced bacterial community responsiveness to raw broccoli sprouts</u>
Of the three experimental factors analyzed in this study, we report that age is the most

significant factor (relative to diet and anatomical location) in driving the richness of gut bacterial

communities, as well as their similarity to each other. The younger trial 1 mice that started

consuming the raw broccoli sprouts at 4 weeks of age had greater observed bacterial richness

by 6 weeks of age than their counterparts fed a control diet (Figure 4A, permANOVA $p < 0.05$).

However, mice in trial 2, starting at 7 weeks, showed no significant difference in bacterial

richness compared to their control group by age 9 weeks (Figure 4B, permANOVA $p > 0.05$).

When comparing bacterial communities within each age group and within each of the four

anatomical locations studied, the broccoli sprout diet increased bacterial richness in the cecum

and the proximal colon of the younger mice, compared to the younger controls (Figure 4B,

Wilcox tests $p < 0.05$). There was no difference between bacterial richness in any gut location in

the older mice consuming broccoli sprouts versus the control diet.

When comparing bacterial community similarity between all samples (beta diversity),

age was again the strongest explanatory factor when comparing bacterial taxa

presence/absence (unweighted, Figures S2, S3) and bacterial taxa presence and abundance

(weighted, Figures 5, 6) metrics. Within the younger trial 1 group, bacterial communities

clustered separately by diet (Figure 5; Table 1, permANOVA $p < 0.001$). This was, in part, due

to significant taxonomic dissimilarity between the two diet groups in the cecum and the distal

colon (Figure 6, Table 1), while differences between diet groups within the proximal colon were

approaching significance. Within the older trial 2 mice, diet was also the most significant driver

of bacterial community similarity, but to a lesser extent than in the younger mice (Figure 5;

Table 1, permANOVA: lower F values, $p < 0.03$). Overall, anatomical location was a significant

factor in bacterial community similarity in older mice, however; within each of the four

anatomical locations studied, there was no significant effect between the diets ($p > 0.05$),

indicating smaller-scale changes across the GI tract communities.

The bacterial taxa which drove the differences between age, diet, and anatomical

treatment were primarily different sequence variants (SVs) identified to the Muribaculaceae and

Lachnospiraceae families, common mouse commensals, abundant in younger broccoli sprout

fed mice (Figure S4). The lactic acid-fermenting genus *Limosilactobacillus* was found in older mice,

and was more abundant in older mice consuming broccoli sprouts (Figure S4). *Helicobacter typhlonius*,

which is pathogenic in immunocompromised mice, was abundant in control mice of both ages,

and in older mice consuming sprouts, while *Escherichia* was found in many locations in the

control mice but only a few older mice consuming sprouts (Figure S4).

Bacterial taxa which were abundant in more than 70% of samples in the groups subset

by trial/age and diet were considered part of the "core" community. SVs in the Muribaculaceae

family made up much of the core microbiota specific to each group, as well as several SVs

identified to Lachnospiraceae, *Bacteroidetes*, *Helicobacter*, and others (Figure S5). *Helicobacter*

was more prevalent and abundant in control mice of both ages.

The Source Tracker algorithm was used to determine if the cecum could be the 'source'

for bacterial population 'sinks' in the colon, as a proxy for the mouse model's applicability to the

human gut anatomical features and microbial communities. A total of 38 SVs were identified as

possibly sourced from the cecum (Figure S6). Common mouse commensals, but not the

putative GSL-converting bacteria in the broccoli sprout fed gut samples, were among those taxa

identified as sourced in the cecum and sinking in the proximal or distal colon.

<u>Raw broccoli sprout diet reduced the abundance of putative pathogens</u>

SFN is bacteriostatic against common gut pathogens, including *Escherichia coli*,

*Klebsiella pneumonia*, *Staphylococcus aureus*, *S. epidermidis*, *Enterococcus faecalis*, *Bacillus*

*cereus* as well as non-pathogenic *Bacillus,* and *Helicobacter* spp., which can be commensal in

immune-competent mice (48, 72). Using that previous literature, we quantified differences in

abundance in putative pathogens, which were higher in control groups, across more locations in

the gut, and in higher abundance than the broccoli sprout groups (Figure 7). This trend was

more pronounced in older control mice which exhibited an abundance of *Helicobacter* SVs

through the cecum, proximal colon, and distal colon, as compared to the younger controls

(Figure 7, ANOVA p=0.147). Within the broccoli sprout groups, the older mice also appeared to

have a greater abundance of *Helicobacter* SVs compared to their younger counterparts in the

distal colon and though it was not statistically significant (Figure 7, ANOVA p=0.657).

296
<u>Raw broccoli sprout diet increased certain putative GLR metabolizing taxa</u>

Several bacterial taxa perform myrosinase-like enzymatic activity and metabolize GLR

into SFN, and after consulting the current literature, we identified 19 taxa to analyze in detail

(38, 85, 86). From our samples, we identified 562 SVs belonging to the following genera:

*Bacillus, Bacteroides, Enterococcus, Lactobacillus, Lactococcus, Pseudomonas,* and

*Staphylococcus* (Figure 8). Though, we did not find *Bifidobacterium, Listeria, Pediococcus,*

*Streptomyces, Aerobacter, Citrobacter, Enterobacter, Escherichia, Salmonella,*

*Paracolobactrum, Proteus,* or *Faecalibacterium.* There were few previously identified putative

GSL-converting taxa found in the younger mice fed the broccoli sprout diet, however, there were

higher read counts for *Bacteroides* and *Lactobacillus* in the younger controls as compared with

older controls or broccoli-fed groups (Figure 8, ANOVA $p < 0.01$). Both diet groups of older mice

exhibited high read counts of *Bacillus* and *Enterococcus* SVs (Figure 8).

The reads from putative GSL-converting genera were identified to species-level using

NCBI BLASTN (Fig. 9). The younger mice consuming sprouts contained *Bacteroides sartorii*

and *B. acidifaciens* in the cecum and distal colon, and *B. caecimuris*, *B. sartorii*, and *B.*

*acidifaciens* in the proximal colon. Most of the SVs in the ileum could not be identified to species

level. In the older trial, similar GSL-converting taxa were present, with five of the identified

species having high abundance (Fig. 9). The older sprout-fed mice contained many

*Lactobacillus intestinalis* reads in the ileum, with fewer in the cecum, proximal and distal colon;

many *B. caecimuris* and *B. acidifaciens* reads in the cecum, and proximal and distal colon; and

*B. sartorii* in the proximal and distal colon.

By contrast, the younger control group had four of the identified GSL-converting species

(Fig. 9): *Lactobacillus gasseri* in the ileum, proximal and distal colon; *B. sartorii*, *B. caecimuris*,

and *B. acidifaciens* in the cecum, proximal and distal colon). The older control group also had

abundant *L. gasseri* in the ileum, but much less in the proximal and distal colon; *B. acidifaciens*

was dominant in the cecum, proximal and distal colon; *L. intestinalis* was present in the ileum,

cecum, and distal colon; and *B. sartorii* was abundant in cecum, proximal and distal colon (Fig.

9). The older control mice had a minor abundance of *E. casseliflavus*, *E. faecalis*, *B.*

*thetaiotaomicron, E. cecorum*, and *S. aureus* (data not shown).

The presence of *Bacteroides thetaiotaomicron* (*B. theta*) was further investigated
through qPCR amplification of its SusR4 regulon, which contains the BT2159-BT2156 operon
and the regulatory gene BT2160, from samples along the digestive tract (ileum, cecum,
proximal & distal colon) of mice consuming broccoli sprout and control diets. For *B. theta*, this
operon has been demonstrated as necessary and sufficient for GSL metabolism and ICT
production both *in vivo* and *in vitro (73)*. These genes code for enzymes (a sugar phosphate
isomerase, glycosyl hydrolase, and two oxidoreductases) that support the transformation of
GLR to the bioactive SFN. We found copies of *B. theta* genes BT2156 in the ileum, cecum, and
distal colon, with a small number of copies in the proximal colon (Figure S7). BT2159 had some
copies in the cecum in both diet cohorts of mice, with other gene copies present in low
quantities across diet and anatomical location The highest copy number of any gene, and only
the significant comparison by diet, was BT2156 in the distal colon of broccoli sprout fed mice
(ANOVA, difference = 92362 gene copies more than control mice, p = 0.00499, controlling for
Trial).

Liou et al. found that *B. theta* sulforaphane production required BT2158 together with
either BT2156 or BT2157 but was most effective when all four operon genes acted in concert
(73). Thus, at each anatomical location, we characterized the distribution of the presence of the
four operon genes (Figure 10A). For the young mice (Trial 1) across all anatomical sites, we
found that the broccoli diet on average shifted the proportion of mice with a high operon
indication (max operon mean copy count > 10,000) from 24% in controls to 64% in the
treatment group (Figure 10B). For the older mice (Trial 2), the impact of the broccoli diet was not
as prevalent across anatomical sites, but was indicated in the cecum and distal colon shifting
the combined middle and high operon indication (max operon mean copy count > 100) in the
ileum from 83% in control mice to 100%, from 66% in control mice to 80% in the cecum, and
from 83% in control mice to 100% in the distal colon (Figure 10A).

**Discussion**

IL-10-KO is an effective model for evaluating host, diet, immune response, and microbiota

This pilot study evaluated IL-10 deficient mice as a model for using dietary broccoli sprout bioactives to reduce inflammation, modify the immune response, and support GI tract microbiota. Inflammatory diseases such as CD are associated with changes in the presence, abundance, and functionality of someone''s microbiome (56, 57, 74), and therapies which improve the functionality of the gut microbiome have the potential to mediate other symptoms (37, 45). Specific pathogen free IL-10-KO mice acquire gut microbiota as they age (75), but they never fully develop diverse, commensal gut microbiota and this appears to be attributable to the spontaneous generation of enterocolitis at the 4 - 6 weeks of age (76). Diet is an important driver of microbiota development in early life (16), and consumption of fiber-rich foods has been associated with microbiota stabilization (77), because the wealth of microbial byproducts is demonstrated to recruit and maintain a diverse microbial community (78), reduce inflammation, improve nutrition, and stimulate gut motility – aspects which are limited in IBD (56, 79, 80). In our study, raw broccoli sprouts in the diet provided during the critical period of microbiome stabilization and enterocolitis development in these IL-10-KO mice resulted in an increase in microbial diversity, increased abundance of potentially beneficial bacteria, and a reduction of potentially pathogenic bacteria. We did not see an improvement in histological damage, which may have been precluded by the use of raw sprouts: feeding pure SFN results in high absorption in the stomach which decreases over the GI tract (81).

Dextran sodium sulfate (DSS) is an established and widely used model for studying SFN and prevention of IBD in animals (45, 52, 53), and results in a disease profile similar in progression and morphology to human Ulcerative Colitis (52, 82). DSS modifies the expression of tight junction proteins in intestinal epithelial cells, leading to a leaky epithelial barrier (83), goblet cell depletion, erosion, ulceration, and infiltration of neutrophils into the lamina propria and submucosa (84), triggering the innate immune response (85, 86). However, Bhattacharyya

et al. reported DSS results in an excess of ROS species which quickly dephosphorylate Hsp27

resulting in direct IkBa dissociation, circumventing the canonical inflammatory pathway (87). As

DSS instigates colitis through chemical and physical damage, without a direct interaction on the

immune system, it is inadequate for investigating the specific immunohistopathology present in

Crohn's Disease. While a few studies have used IL-10-KO mice to study the effects of certain

mineral additives (88, 89), isomaltodextrin supplementation (90), or high-fat diets on colitis (91,

92), the use of this model for diet studies is still nascent. We demonstrated that a medium-high

concentration of raw broccoli sprouts in the diet of IL-10-KO mice induced increased bacterial

richness, SFN in the plasma, and collectively reduced the symptoms of enterocolitis.

<u>Younger IL-10-KO mice were more responsive to diet compared to adolescent counterparts</u>

We observed a previously unreported effect of age in early life – a three week difference

– on the heightened response of younger mice to broccoli sprouts. The gut microbiota and

immune systems in younger mice were more responsive to raw broccoli sprouts, and we

hypothesized that this was driven by the instability of the developing gut microbiota and its

amenability to selective pressures (68, 93, 94).

Mice wean at approximately 20-22 days of life (3 weeks) which is a critical period for the

co-development of the immune system and gut microbiota, and is coordinated by immune

factors in milk, growth of the GI tract, and improvement of epithelial barrier function (95). By 28 -

35 days of life (4 - 5 weeks), juvenile mice are beginning sexual development and still have a

changing gut microbiome (96). By days 35 - 42 (5 - 6 weeks), they begin adolescence (97)

during which their gut microbiome stabilizes unless it is perturbed (96). Thus, our younger mice

were still in a transitional state of life and older mice were just beginning a period of life marked

by stability of the immune system and gut microbiome. Our findings are consistent with previous

human studies that suggest that the gut microbiota of infants and children are more plastic than

that of adults (16): starting around age 3, gut microbiota fall into long-term patterns driven by

consistency in diet and lifestyle (16, 77, 98, 99). Collectively, this may indicate that dietary

interventions to restructure the microbiome, would be more effective in children and adolescents

than in adults, which is supported by other literature (37).

Also, compared to the controls, the younger mice fed a diet containing broccoli sprouts

had more bacterial genera meet the ad hoc core cutoff for prevalence and abundance. The gut

microbiomes of pediatric CD patients show divergence from healthy siblings early in the disease

and a more definable "disease state microbiome" than adults with CD. We found that the older

broccoli-fed mice had fewer genera that met the core standard, which could suggest that the

core gut microbial community diminishes with age and interventions would need to be

individualized. In adults, undergoing dietary change is one of the few external factors that can

"destabilize," or subject the adult gut microbiota to change (16, 99), which may be useful in

removing gut microbial communities which are not providing functional benefits and recruiting a

different community instead.

Biogeographical patterns emergent in younger mice consuming broccoli sprouts

Anatomical location within the gut, including different organs and solid-associated, liquid-

associated, and mucosal-associated locations, select for different bacterial communities due to

environmental conditions and anatomical features specific to each location, i.e. biogeography

(100–102), even within the first few days of life (103). In IL-10-KO mice, which are raised in

pathogen-free conditions and acquire some gut microbiota, there is no biogeographical signal in

the gut, and the damage to the epithelium caused by inflammation or the lack of IL-10 may

preclude the ability of a typical biogeographical signal to form. The addition of a conventionally

raised mouse to the IL-10-KO mouse cages, and placement in a conventional mouse room, is

enough to stimulate microbial transmission (104) and for IL-10-KO mice to acquire new gut

microbiota even as they react negatively to them.

The third experimental factor of this study, and the one that showed to be the least

significant driver of microbial community dissimilarity, was the anatomical location of sample

derivation. We found that the number of bacterial taxa (SVs) in the cecum and proximal colon of

the younger trial 1 mice fed broccoli sprouts was significantly elevated, as compared to the

control trial 1 mice, and significant community dissimilarity between the broccoli sprout diet

group and the control group in both the cecum and the distal colon. There were no significant

dietary treatment effects on richness or bacterial community similarity in any anatomical location

studied in the older trial 2 mice (Figure 4B). The cecum, a muscular and oblong pouch emerging

from the junction of the ileum and proximal colon, houses a diverse community of fibrolytic

bacteria, and specialized peristaltic movements allow for fiber in the ileocecal junction to be

pulled into the cecum, as well as separate large and small fibers to allow for retention of

particles to induce more microbial fermentation (105, 106). Mice have a large cecum, even in IL-

10-KO mice this organ acquires a diverse microbiota (75, 107). Studies on IBD models report

significant inflammation and changes to cecal microbiota in mice (45, 108, 109). It is likely that

given the short duration of this experimental period, the cecum and parts of the colon that also

house diverse microbiota were the only organs which were suited to recruit differential bacterial

communities fast enough to be statistically significant. While humans have a small cecum

relative to our size, and do suffer inflammation there in IBD patients, the effect on IBD of the

microbiota there is unknown.

Raw broccoli sprouts encouraged some putatively beneficial taxa

CD has specifically been associated with a reduction in *Faecalibacterium prausnitzii*, a

putative glucosinolate metabolizer, and reintroduction of this bacteria resulted in decreased

inflammation (110). However, many gut microorganisms are only useful to the host in specific

metabolic conditions (111, 112), which can obfuscate our understanding of which

microorganisms may be useful to try and maintain in order to preserve beneficial interactions

with the host. Further, commensalism cannot be assumed for any taxa, as IL-10-KO mice

infected with *Helicobacter* also exhibited higher abundance of non-commensal *Bacteroides* and

*Lactobacillus* species (113).

SFN is bacteriostatic against bacterial taxa which may provide some benefit to mice,

including *Bacteroides (114)*, a genera containing known glucosinolate metabolizers (73), and

non-pathogenic *Bacillus* or *Enterococcus*. Across many of our experimental groups, we found

several bacterial genera, including *Bacteroides* (but not *Bacteroides thetaiotaomicron*),

*Enterococcus casseliflavus*, *Lactobacillus gasseri*, and others in very low abundance, that may

perform myrosinase-like activity in the digestive tract (115). *Enterococcus casseliflavus* are

gram-positive, motile facultative anaerobes sourced from vegetables, with high glucosinolate

degradation efficiency (116). One SV was matched to *E. casseliflavus* based on BLASTN

results in the older mice, and favored the broccoli mice (1164 count) compared to the control

(113 count). Although GLR was low in the broccoli diet (4 µg per gram of diet sampled), *E.*

*casseliflavus* may have been responsible for some of the conversion to SFN.

*Lactobacillus gasseri* was the most abundant species with putative myrosinase-like

enzymatic activity. A probiotic strain with implications for fighting pathogens, *L. gasseri* has

hydrolyzed GLR to SFN in the cecum of rats (39, 117). Four SVs were mapped to *L. gasseri*

using BLASTN. These sequences were most abundant in the control for both trials, with the

highest counts occurring in the younger control mice.

Contrary to expectations, except for *Bacteroides caecimuris*, *Enterococcus casseliflavus*,

and *Lactobacillus intestinalis*, the species identified from GSL converting genera tended to be

found in control treatments for both age groups of mice, although it is important to note that the

use of 16S rDNA precluded any assessment of microbial activity. There was a greater

abundance of putative GSL converting taxa associated with the older mice (7-9 weeks-old),

except in the cecum where the abundances were higher for the younger mice. The high

concentration of SFN in the diet, and low concentration of the GLR precursor, may have

precluded the need for metabolizing-competent bacteria, and it may be that the taxa present in

younger mice were sensitive to the presence of SFN while the community present in older mice

was not. In contrast, mice on the broccoli sprout diet overall demonstrated higher indications of

the presence of the operon BT2159-BT2156, suggesting *B. theta* may be the primary GLR

convertor in these mice, and suggesting that 16S rDNA taxonomic identification on its own may

not be an accurate metric for presumptive GLR conversion in the gut. Using the presence of the

operon as a metric, the use of a broccoli diet to recruit functional *B. theta* was most strongly

indicated for young mice (4-6 weeks-old), supporting our *a posteriori* hypothesis.

<u>Raw broccoli sprouts reduced putative proinflammatory bacteria</u>

The causal relationship between potentially pathogenic bacteria and the onset of IBD

symptoms remains murky (56, 118), with postulations about the involvement of various

microorganisms, including *Mycobacterium avium* subspecies *paratuberculosis*, adherent-

invasive *Escherichia coli, Mycoplasma* spp., *Helicobacter pylori*, and *Chlamydia* spp. However,

the role of proinflammatory bacteria in exacerbating flare-ups is clear, and proinflammatory

bacteria have been identified in and found to be more abundant in IBD patients as compared to

healthy children (34) or adults (56), although often newly-diagnosed children exhibit similar fecal

microbial communities compared to healthy siblings (119).

SFN induces colonocytes to produce several antimicrobial compounds (120), something

which is lacking in CD patients with abnormally functioning Paneth cells (121). SFN itself is

bacteriostatic against several bacteria, including *Escherichia coli*, *Klebsiella pneumonia*,

*Staphylococcus aureus*, *S. epidermidis*, *Enterococcus faecalis*, *Bacillus cereus,* and

*Helicobacter* spp. (48, 49, 72). Interestingly, our results suggest that consumption of broccoli

sprouts may prevent the accumulation of putatively harmful, pro-inflammatory bacteria. When

compared to the controls, the broccoli-fed mice exhibited less prevalence and abundance of

*Helicobacter*, which is a non-harmful commensal in conventionally raised and immune

competent mice (122), but acts as a pro-inflammatory bacterial genus in IL-10-KO mice with

little ability to moderate their own response to the presence of gut microorganisms (64). The

specific species of *Helicobacter* which colonies IL-10-KO mice can also affect the host's

immune response and severity of symptoms (123).

<u>Whole food strategies provide multiple mechanisms of benefit and harm reduction</u>

In many but not all people or circumstances, high-fiber diets increase microbial diversity,

recruit beneficial symbiotic species in the gut, and reduce inflammation by providing

antioxidants against reactive oxygen species or antimicrobials against microbial competitors

(120, 124). Dietary fiber has the potential to reduce the risk of Crohn's flare-ups by up to 40%

(125), and high-fiber foods can stimulate beneficial microbial byproducts in the gut (124).

However, during a flare-up, high-fiber foods can worsen symptoms, and certain fiber types can

exacerbate symptoms in IBD patients (126). Opting for foods rich in soluble fiber, which can

help slow down digestion and ease diarrhea, is recommended for individuals with Crohn's

disease. On the other hand, foods containing insoluble fiber can increase water content in the

gut, leading to rapid digestion, watery diarrhea, stomach cramps, or gas, and in severe cases,

may cause blockages. Using purified GLR or SFN have been evaluated as a means of inducing

health benefits while avoiding fiber in IBD patients, with varying effectiveness, e.g., (40, 51).

Feeding SFN directly to mice, thus eliminating fiber risk, does not appear to change

cecal butyrate (127), and while butyrate treatments to the gut are often successful in patients

with ulcerative colitis it is unclear whether butyrate is helpful for patients with CD as cellular

damage may prevent colonocytes from utilizing it (128). We balanced the 10% raw broccoli

sprout/90% control diet to the 100% control diet by total fiber, and the control diet contains oats

which are a prime source for microbially-produced butyrate (129). While mature broccoli is a

reliable source of fiber and stimulates butyrate production by gut bacteria, there is no literature

to suggest that broccoli sprouts would stimulate more microbial fermentation and byproducts

than the equivalent grams of fiber from oats. Further, we found no putative butyrate-producing

genera in our sprout-fed mice: no *Bifidobacterium*, *Clostridium*, or *Butyrococcus*, and

*Bacteroides* were only high in the younger control mice. Similarly, *Bifidobacterium*, *Clostridium*,

and *Bacteroides* can dissociate bilt salts in the colon, which otherwise exacerbate symptoms in

IBD patients, and we found *Lactobacillus intestinalis,* also capable of this (130), in our sprout-

fed mice. Finally, mucin degrading capacity of gut bacteria is reduced in CD and UC patients

(131), and alterations to the gut microbiota could have action here.

Although some studies have indicated positive outcomes with anti-inflammatory dietary

interventions in children and adolescents with Crohn's disease, their effectiveness may vary

among individuals. Factors such as adherence to the prescribed diet, individual tolerance to

specific foods, and the involvement of healthcare professionals in closely monitoring progress

all play critical roles. Diet can also trigger symptoms of IBD, however; It is important to note that

there is no singular food or food group that can be universally linked to every case of Crohn's

disease, as different individuals may experience gut irritation and inflammation from different

foods. Children and adolescents with Crohn's disease may exhibit distinct manifestations

compared to adults, often presenting with more extensive disease involvement, including

perianal disease and structural complications (34–36). Consequently, personalized

recommendations are often needed, and the unique nutritional requirements of pediatric and

adolescent populations must be addressed to ensure adequate nutrient intake for growth,

development, and overall health (37).

Limitations and Future Directions

Due to the short duration of the experiment, we were unable to discern if the increase in

diversity persists into adulthood, and if this would preclude the development of more intense

symptoms and damage to the intestinal epithelium. We lacked sufficient power to determine sex

specific differences in response to diet (90, 132), immune function (133), and microbial

acquisition (134). We also lacked data to determine microbial activity, beneficial or detrimental.

Future research will be required to examine the effect of cooking preparation on the

concentrations of precursors and bioactives which are contained in the diet and available in the

gut, as well as microbially sourced bioactives, and will require metabolomics to fully explore this.

The myrosinase enzyme present in broccoli and sprouts can metabolize the GLR precursor to

bioactive SFN when the plant tissues are cut or chewed, but most of the precursor is converted

to biologically inactive SFN nitrile by the epithiospecifier protein, also present in broccoli (14).

SFN was high in our raw sprout diet, which was an effect of the preparation process stimulating

release of myrosinase. However, SFN is an unstable molecule and will not persist in processed

raw sprouts unless kept frozen. These enzymes can be inactivated by cooking, preserving the

stable precursor in the diet and allowing gut microbiota to perform the conversion in the

intestines (38).

Future research will be needed into the feasibility and adoptability of this intervention.

The difficulty in managing CD symptoms is costly to patients (135), and healthcare systems

(136, 137). Further, many IBD patients may be told to avoid fiber for fear of exacerbating

symptoms or aggravating tender intestines (24, 54). Given the importance of diet in supporting

host health and microbiota, we underscore the need for understanding dietary preferences, and

for incorporating nutritional intervention as part of a holistic treatment for gastrointestinal

inflammation.

**Materials and Methods**

<u>Diet</u>

Multiple lots of Jonathan's Sprouts™ (Rochester, MA, U.S.) broccoli sprouts were

purchased from a nearby grocery store (Bangor, ME, U.S.) and stored in a -80▢ freezer until

freeze-drying at the University of Maine Pilot Plant (Orono, ME, U.S.) to lower moisture content

and reduce enzymatic activity. The control diet was 100% irradiated 5LOD diet irradiated 5LOD

rodent diet powder (ScottPharma Inc., Marlborough, MA, #50795) mixed with ultra-pure water.

The freeze-dried sprouts were crushed by mortar and pestle into a fine powder and mixed with

the 5LOD powder and water to a concentration of 10% by weight to form the treatment diet, as

that concentration reliably produces consistent anti-inflammatory results (46). Diet pellets were

formed using a silicone mold to ensure consistent sizing, dried at room temperature for up to 48

hours in a chemical safety hood to facilitate moisture evaporation, and after drying were stored

in Ziploc bags in a -10□ freezer until future use. The sprouts are washed prior to packaging, and

freezing, freeze-drying, and drying reduces microbial biomass, growth, and activity, and may kill

some bacteria through desiccation. However, we did not verify that diets were sterile after

formulating them, as the goal of the experiment was to introduce microbiota to mice.

The fiber content of the control diet and broccoli sprout-supplemented diet was ~5%. No

GLR or SFN was found in the control and the sprout diet contained an average 4 µg of GLR and

85 µg of SFN (induced when raw diets are crushed and myrosinase is released from broccoli

tissue vesicles) per gram of dried diet sampled (Figure S8), using LC/MS (38).

IL-10 Mouse Model

All experimental protocols were approved by the Institutional Animal Care and Use

Committee of the University of Vermont (PROTO202000040), in Burlington, Vermont, U.S. and

the Institutional Biosafety Committee of the University of Maine (protocol SI-092220). Two

replicate trials were conducted using male and female IL-10-KO mice (*Mus musculus*) on a

C57BL/6 background (B6.129P2-Il10tm1Cgn/J, strain 2251; Jackson Laboratories, Bar Harbor,

Maine, U.S.). IL-10-KO mice are a well-demonstrated model for Crohn's research and the

C57BL/6 genetic line shows some genetic-based resistance to developing symptoms (61),

which allowed us to focus on environmental triggers of disease.

Trial 1 included 9 mice (n = 4, control; n = 5 sprout) starting at 4 weeks of age and post-

weaning. Trial 2 included 11 mice (n = 6, control; n =5 sprout) starting at 7 weeks of age. The

experimental design was structured as a prevention paradigm (Figure 1). A colony of

homozygous IL-10-KO mice was maintained in a barrier facility which is specific pathogen free,

and especially *Helicobacter (H.) hepaticus* free. For the duration of the trial, they were fed either

the 5LOD control diet or the treatment diet consisting of 10% (w/w) raw broccoli sprouts and

90% control diet, which was balanced for micronutrients and fiber (5% of diet). Mice had access

to tap water *ad libitum*.

Mice were acclimated to diets for 7 days prior to the induction of colitis, which was

achieved by moving mice to a conventional mouse room and adding an *H. hepaticus* positive

mouse in the cage for the remainder of the experiment. The *H. hepaticus* positive mouse

transfers commensal bacteria, as well as species which may cause infection in

immunocompromised mice, via feces and grooming, and bacteria induce colitis over the course

of 10 days (61, 67). While *H. hepaticus* is widespread in research mouse colonies and causes

disease in immunocompromised mice, it does not cause symptoms in healthy, conventional

mice (122). The conventional mice were fed an irradiated RMH 3000 diet (Prolab® IsoPro®)

prior to being co-housed with IL-10-KO mice, at which point they consumed the control or 10%

broccoli sprout diets based on the cage treatment. Day 0 of each trial was set as the day the IL-

10 mice were exposed to the *H. hepaticus* positive mouse, such that experimental Day 10 is

when symptoms appear (Figure 1). Animals were euthanized by carbon dioxide and cervical

dislocation on Day 16.

Assessment of Disease Activity Index

Disease severity was assessed every other day beginning on Day 0 when mice were co-

housed with the *H. hepaticus* positive mouse, using a DAI. This included a score that combined

fecal blood presence, severity, and stool consistency (138). The animals' weights were

normalized to their baseline weight on Day 0 of the trial, as at this age, mice are still in a growth

phase and likely to gain weight even when under duress (139); and this was the rationale for

excluding weight loss as part of the DAI criteria. The presence of fecal blood was assessed

using Hemoccult Single Slide testing slides (Beckman Coulter, Brea, CA, U.S.).

Assessment of Lipocalin and Proinflammatory Cytokines

Plasma samples collected immediately after euthanasia, as well as fecal samples at

various time points, were frozen for evaluation of fecal lipocalin (LCN2); a neutrophil protein that

binds bacterial siderophores and serves as a biomarker for intestinal inflammation (71). Frozen

fecal samples were weighed up to 20 mg and reconstituted in phosphate buffered solution

(PBS) with 0.2 mL of 0.1% Tween 20, thawed, and vortexed to create a homogenous

suspension. The samples were centrifuged for 10 minutes at 12,000 rpm at 4 °C. Clear

supernatant was collected and stored at -20 °C until analysis. LCN2 concentration was

measured by a mouse Lipocalin-2/NGAL DuoSet ELISA kit (R & D Biosystems, USA) following

the manufacturer's instructions. The readings at wavelengths of 540 nm and 450 nm were

measured by a Thermo Scientific Varioskan LUX Multimode Microplate Reader. The readings at

540 nm were subtracted from the readings at 450 nm to correct for the discoloration of the

solution in serum.

We analyzed several pro-inflammatory cytokines in the mouse plasma samples which

were collected at the end of the study: CCL4 /macrophage inflammatory protein 1β (MIP1β),

which regulates the migration and activation of immune cells during inflammation (140); and IL-

1β, IL-6, and tumor necrosis factor alpha (TNF-α), which play several roles regulating immune

responses (141–143). The concentrations of mouse CCL4/MIP-1β, IL-1β/IL1F2, IL6, and TNF-a

were analyzed using the Simple Plex Ella Automated Immunoassay System (Ella) from R&D

Biosystems, USA. Mouse serum samples were diluted 10-fold using the kit-specific reagent

(SPCKA-MP-007374, Protein Simple, Bio-Techne), and the concentrations were measured

following the manufacturer's instructions on the Ella system. Mean values were calculated for

each cytokine and analyzed using ANOVA.

Quantification of Plasma SFN

Plasma collected after euthanasia was also used for SFN measurement using LC/MS

(38). To extract SFN, 50 µl of plasma homogenate was mixed with 3-5 fold of acetonitrile and

centrifuged at 14,000 rpm, 4 °C for 15 min, and the supernatant stored at -20°C. An AB SCIEX

QTRAP 4500 with a TurboV electrospray ionization source mass spectrometer (Applied

Biosystems, Carlsbad, CA) coupled to an Agilent 1200 Series HPLC system (Agilent

Technologies, Santa Clara, CA) was used for the quantification. HPLC separation was

performed on a Waters XBridge® C18 3.5 µm 5 cm x 2.1 mm column (Waters Corporation,

Milford, MA). Mobile phase A (water containing 0.1% formic acid) was first kept at 90% for 0.5

min, and then decreased to 5% over 1 min and maintained at 5% for 2 min, and then returned to

10% mobile phase B (ACN containing 0.1% formic acid) and maintained for 3 min. The flow rate

was 400 L/min. Positive ion MS/MS was conducted to detect SFN with the following conditions:

source temperature, 400 °C; curtain gas (CUR), 30 psi; ionspray voltage (IS), 4500 V;

desolvation gas temperature (TEM), 500 °C; ion source gas 1 (GS1), 60 psi; ion source gas 2

(GS2), 40 psi; collision gas (CAD), high; entrance potential (EP), 4 eV; collision energy (CE), 15

eV. The MS/MS transition of 178 > 114 was used to detect SFN, and a dwell time of 50 ms was

used for the transition. Data acquisition and quantitation were performed using Analyst software

(Applied Biosystems, Carlsbad, CA).

Histological Analysis of Tissues

After euthanasia, tissue segments (1 cm, collected in duplicate) from the ileum, proximal

colon and distal colon were collected and fixed in 4% paraformaldehyde overnight for

histological evaluation. Tissues were immediately rinsed with phosphate buffer solution (PBS),

placed in 2% paraformaldehyde/0.2% picric acid as a preservative, and stored at 4□ until

transport to the University of Maine Electron Microscopy Laboratory (Orono, Maine) for

processing. All processing protocols took place in a biosafety cabinet using aseptic techniques

to reduce contamination. A pipette was used to gently remove the previous PBS that the tissue

was submerged in without disturbing tissue while leaving a little bit of PBS behind to prevent the

tissue drying out. The sample tubes were refilled with fresh PBS, and the wash step was

repeated four times throughout the day every three hours with samples stored at 4□ between

washes.

After the final wash step, the tissue samples were transferred into embedding baskets.

The samples were dehydrated by immersing the samples in a graded series of ethanol at 4□

(50%, 70%, 80%, 95%, 100% ethanol), cleared in xylene, and then infiltrated and finally

embedded in Paraplast X-tra. During embedding, care was taken to orient the intestine samples

to give a cross section during sectioning. Slides were sectioned at 5µm and stained in

Hematoxylin and Eosin (144). Images were taken using an Olympus BX41 microscope with a

Zeiss ERc 5s digital camera.

The tissues were scored with 7 criteria (Figure S1) used to assess inflammation in the

ileum, and the proximal and distal colon tissues, performed according to parameters provided

by the Mawe Lab at the University of Vermont (145). Epithelial damage and architectural

changes were scored from 0 (no damage) to 2 (extensive damage). Similarly, infiltration of

mononuclear cells in the lamina propria was scored on a scale of 0 (no infiltration) to 2

(extensive infiltration). Infiltration of polymorphonuclear cells in both the lamina propria and the

epithelium were scored on a scale of 0-2, with 0 indicating no infiltration, 1 equating to sighting

of ≥1 cell in each viewing field, and 2 equating to >3 cells in each viewing field. Abscesses,

ulcers, erosion, and branched crypts were scored together with 0 being absence of these

damage indicators and 1 being presence. Finally, presence of granulomas was scored with 0

indicating no granulomas and 1 equating to ≥1 granulomas. Scores by treatment, anatomical

location, and trial were compared with analysis of variance.

<u>DNA extraction and 16S rRNA bacterial sequencing library preparation</u>

On Day 16, following euthanasia, intestinal tissue segments (2 cm in length) were

collected from the ileum, cecum, proximal colon, and distal colon; placed in RNAlater

preservative (Invitrogen, Waltham, MA, U.S.); and transported overnight on ice to the University

of Maine for DNA extraction. All DNA extraction processing steps took place in a biosafety

cabinet using aseptic techniques to reduce contamination. All tissues containing their resident

gut microbiota were gently homogenized with vortexing, then treated with propidium monoazide

(PMA; BioTium) following kit protocols at a final concentration of 25 μmol. PMA covalently binds

to relic/free DNA and DNA inside compromised/dead cell membranes, and prevents

amplification in downstream protocols to preclude dead DNA from the sequence data.

Following PMA treatment, bulk DNA was extracted from tissue-associated bacterial

communities (n = 80 samples), or no-template (water) control samples (n = 4, one for each

extraction batch) using commercially available kits optimized for fecal-based microbial

communities (Zymo fecal/soil kit), and some aliquots archived. DNA extract was roughly

quantified and purity-checked with a Nanodrop spectrophotometer. Samples underwent DNA

amplicon sequencing of the 16S rRNA gene V3-V4 region, using primers 341F (146) and 806R

(147) and protocols consistent with The Earth Microbiome Project (148), and sequenced on an

Illumina MiSeq platform using the 2 x 300-nt V3 kit (Molecular Research Labs, Clearwater, TX).

Nine samples failed the first sequencing run, and were extracted again for repeated sequencing.

Raw sequence data (fastq files and metadata) from both runs are publicly available from the

NCBI Sequence Read Archive (SRA) under BioProject Accession PRJNA909836.

16S rRNA bacterial community sequencing analysis

Amplicon sequence data was processed using previously curated workflows in the Ishaq

Lab (Supplemental Material) which used the DADA2 pipeline ver. 1.22 (149) in the R software

environment ver. 4.1.1 (150). The dataset started with 38,287,212 million raw reads from 80

samples and 4 negative controls. All samples were sequenced in an initial batch together, but 5

samples failed to meet all quality control standards and were resequenced. The two sequencing

batches were put through quality control steps separately in DADA2 and combined before

rarefaction. Trimming parameters were designated based on visual assessment of the

aggregated quality scores at each base from all samples (plotQualityProfile in DADA2): the first

and last 10 bases were trimmed, and sequences were discarded if they had ambiguous bases,

more than two errors, or matching the PhiX version 3 positive control (Illumina; FC-110-3001).

After filtering, 43,382,928 paired non-unique reads and 289 samples remained.

The DADA algorithm was used to estimate the error rates for the sequencing run,

dereplicate the reads, pick SVs which represent 'microbial individuals', and remove chimeric

artifacts from the sequence table. Taxonomy was assigned using the Silva taxonomic training

data version 138.1 (151) and reads matching chloroplasts and mitochondria taxa were removed

using the dplyr package (152). No-template control samples were used to remove

contaminating sequences from the samples by extraction batch (153). The sequence table,

taxonomy, and metadata were combined using the phyloseq package (154) to facilitate

statistical analysis and visualization, representing 80 samples and 22,805 taxa. Due to the

variability in sequences per sample which passed quality assurance parameters (range 12,060 -

157,040 sequences/sample), the data were rarefied (155, 156) to 12,060 sequences/sample.

Normality was checked using a Shapiro-Wilks test on alpha diversity metrics generated

from rarefied data; observed richness (W = 0.86662, p-value = 1.186e-06); evenness (W =

0.89141, p-value = 9.775e-06); and Shannon diversity were not normally distributed (W =

0.95012, p-value = 0.005061). Linear models compared alpha diversity metrics (lme4 package

(157)), in which anatomical location, diet treatment, and trial (either 1 or 2) were used as factors.

Jaccard unweighted similarity was used to calculate sample similarity based on

community membership (species presence/absence), and non-parametric multidimensional

scaling (Trial 1 run 20 stress = 0.1940131; Trial 2 run 20 stress = 0.1748438) and tested with

permutational analysis of variance (permANOVA) by using the vegan package (158). Random

forest feature prediction with permutation was used to identify differentially abundant SVs based

on factorial conditions (159). Plots were made using the ggplot2 (160), ggpubr (161), and

phyloseq packages. Source Tracker algorithms which had been modified for the R platform

(162, 163) were used to identify source: sink effects based on anatomical location. This was

used to determine if the cecum could be the source for population sinks in the colon, as a proxy

for the model's applicability to the human gut anatomical features and microbial communities

(Figure S6).

<u>Quantitative PCR performed on genes which metabolize glucoraphanin</u>

The ability of gut bacteria to perform glucoraphanin conversion to sulforaphane was

assessed using gene sequences and primers associated with the established pathway in

*Bacteroides thetaiotaomicron* VPI-5482 (73) (see Table S1). Gene sequences of interest from

*B. thetaiotaomicron* VPI-5482 were evaluated through the ApE-A plasmid Editor v3.1.3 to

confirm optimal melting temperature and GC content of primers, and assessment of primers

was performed using NCBI primer blast (https://www.ncbi.nlm.nih.gov/tools/primer-blast/) and

IDT's OligoAnalyzer (www.idtdna.com). Quantitative polymerase chain reaction (qPCR) was

completed on extracted DNA to determine copy numbers of glucoraphanin-metabolizing genes

previously established in *B. thetaiotaomicron* using a Applied Biosystems QuantStudio 6 Flex

Real-Time PCR system (Applied Biosystems, Foster City, CA, USA), primer sets BT2156-

BT2160, and Luna® Universal qPCR Master Mix were employed. After diluting primers to 10 µM

working concentration, the primer concentration in each well was 0.25 µM. PCR conditions

consisted of one cycle of 50°C for 2 min, one cycle of 95°C for 1 min, and forty cycles of 95°C

for 15 sec, 60°C for 30 sec and 72°C at either 20, 25 or 30 sec based on the primer set used

(Table S1). Standard curves were generated for each gene by serially diluting geneblocks (IDT)

six times. All standards and samples were tested in triplicate, with standards and negative

controls present on both plates. Using the standard curve produced in the Quantstudio analysis

software, sample gene copy numbers were quantified. R software was used to run an analysis

of variance, with corrections applied for multiple comparisons through Tukey's HSD.

**Acknowledgements**

All authors have read and approved the final manuscript. This project was supported by

the USDA National Institute of Food and Agriculture through the Maine Agricultural & Forest

Experiment Station: Hatch Project Numbers ME022102 and ME022329 (Ishaq) and ME022303

(Li); the USDA-NIFA-AFRI Foundational Program [Li and Chen; USDA/NIFA 2018-67017-

27520/2018-67017-36797]; and the National Institute of Health [Li and Ishaq; NIH/NIDDK

1R15DK133826-01] which supported Marissa Kinney, Timothy Hunt, and Benjamin Hunt.

Johanna Holman was supported by ME0-22303 (Li), and Lola Holcomb was supported by US

National Science Foundation One Health and the Environment (OG&E): Convergence of Social

and Biological Sciences NRT program grant DGE-1922560, and the UMaine Graduate School

of Biomedical Science and Engineering.

809
810
811 **Author Contributions**

812 Conceptualization, S.L.I., Y.L., T.Z., G.M., P.M., G.C.; Methodology, S.L.I., Y.L., T.Z., G.M.,
813 P.M., J.H., L.H., M.H., B.L., EP; Software, S.L.I.; Formal Analysis, J.H., L.H., S.L.I.; M.H., A.S.,
814 T.H., B.H., M.K.; Investigation, M.H., B.L.; J.H., L.H., L.C.; A.S., J.P.; Resources, G.M., S.L.I.,

Y.L.; Data Curation, S.L.I., J.H., L.H.; Writing - Original Draft, J.H., L.H., S.L.I., Y.L.; Writing - Review and Editing; S.L.I., Y.L., T.Z., G.M., P.M., J.H., L.H., M.H., B.L., G.C.,T.H., B.H., E.P., M.K., J.P.; Visualization, J.H., L.H., S.L.I., A.S.,T.H., B.H., M.K.; Supervision, S.L.I., Y.L., T.Z.; G.M.; Project Administration, S.L.I., Y.L., G.M.; Funding Acquisition, S.L.I., Y.L., G.M. G.C.

## References

1. GBD 2016 DALYs and HALE Collaborators. 2017. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet 390:1260–1344.

2. Sulz MC, Burri E, Michetti P, Rogler G, Peyrin-Biroulet L, Seibold F, on behalf of the Swiss IBDnet, an official working group of the Swiss Society of Gastroenterology. 2020. Treatment Algorithms for Crohn's Disease. Digestion 101 Suppl 1:43–57.

3. Lichtenstein GR, Shahabi A, Seabury SA, Lakdawalla DN, Espinosa OD, Green S, Brauer M, Baldassano RN. 2020. Lifetime Economic Burden of Crohn's Disease and Ulcerative Colitis by Age at Diagnosis. Clin Gastroenterol Hepatol 18:889–897.e10.

4. Gao N, Qiao Z, Yan S, Zhu L. 2022. Evaluation of health-related quality of life and influencing factors in patients with Crohn disease. J Int Med Res 50:3000605221098868.

5. Mitropoulou M-A, Fradelos EC, Lee KY, Malli F, Tsaras K, Christodoulou NG, Papathanasiou IV. 2022. Quality of Life in Patients With Inflammatory Bowel Disease: Importance of Psychological Symptoms. Cureus 14:e28502.

6. Roda G, Chien Ng S, Kotze PG, Argollo M, Panaccione R, Spinelli A, Kaser A, Peyrin-Biroulet L, Danese S. 2020. Crohn's disease. Nat Rev Dis Primers 6:22.

7. Spehlmann ME, Begun AZ, Burghardt J, Lepage P, Raedler A, Schreiber S. 2008. Epidemiology of inflammatory bowel disease in a German twin cohort: results of a

nationwide study. Inflamm Bowel Dis 14:968–976.

8.  Lindoso L, Mondal K, Venkateswaran S, Somineni HK, Ballengee C, Walters TD, Griffiths

A, Noe JD, Crandall W, Snapper S, Rabizadeh S, Rosh JR, LeLeiko N, Guthery S, Mack D,

Kellermayer R, Gulati AS, Pfefferkorn MD, Moulton DE, Keljo D, Cohen S, Oliva-Hemker M,

Heyman MB, Otley A, Baker SS, Evans JS, Kirschner BS, Patel AS, Ziring D, Stephens

MC, Baldassano R, Dubinsky MC, Markowitz J, Denson LA, Hyams J, Kugathasan S,

Ananthakrishnan AN. 2018. The Effect of Early-Life Environmental Exposures on Disease

Phenotype and Clinical Course of Crohn's Disease in Children. Am J Gastroenterol

113:1524–1529.

9.  Örtqvist AK, Lundholm C, Halfvarson J, Ludvigsson JF, Almqvist C. 2019. Fetal and early

life antibiotics exposure and very early onset inflammatory bowel disease: a population-

based study. Gut 68:218–225.

10. Segal AW. 2019. Studies on patients establish Crohn's disease as a manifestation of

impaired innate immunity. J Intern Med 286:373–388.

11. Hazel K, O'Connor A. 2020. Emerging treatments for inflammatory bowel disease. Ther

Adv Chronic Dis 11:2040622319899297.

12. Gunasekera DC, Ma J, Vacharathit V, Shah P, Ramakrishnan A, Uprety P, Shen Z, Sheh

A, Brayton CF, Whary MT, Fox JG, Bream JH. 2020. The development of colitis in Il10-/-

mice is dependent on IL-22. Mucosal Immunol 13:493–506.

13. Jiang Y, Wu S-H, Shu X-O, Xiang Y-B, Ji B-T, Milne GL, Cai Q, Zhang X, Gao Y-T, Zheng

W, Yang G. 2014. Cruciferous vegetable intake is inversely correlated with circulating levels

of proinflammatory markers in women. J Acad Nutr Diet 114:700–8.e2.

14. Holman J, Hurd M, Moses P, Mawe G, Zhang T, Ishaq SL, Li Y. 2023. Interplay of

Broccoli/Broccoli Sprout Bioactives with Gut Microbiota in Reducing Inflammation in

Inflammatory Bowel Diseases. J Nutr Biochem 113:109238.

15. Gibbons SM, Kearney SM, Smillie CS, Alm EJ. 2017. Two dynamic regimes in the human

gut microbiome. PLoS Comput Biol 13:e1005364.

16. Derrien M, Alvarez A-S, de Vos WM. 2019. The Gut Microbiota in the First Decade of Life.

Trends Microbiol 27:997–1010.

17. Ostrem Loss E, Thompson J, Cheung PLK, Qian Y, Venturelli OS. 2023. Carbohydrate

complexity limits microbial growth and reduces the sensitivity of human gut communities to

perturbations. Nat Ecol Evol 7:127–142.

18. Swidsinski A, Ladhoff A, Pernthaler A, Swidsinski S, Loening-Baucke V, Ortner M, Weber J,

Hoffmann U, Schreiber S, Dietel M, Lochs H. 2002. Mucosal flora in inflammatory bowel

disease. Gastroenterology 122:44–54.

19. Duvallet C, Gibbons SM, Gurry T, Irizarry RA, Alm EJ. 2017. Meta-analysis of gut

microbiome studies identifies disease-specific and shared responses. Nat Commun 8:1784.

20. Shalon D, Culver RN, Grembi JA, Folz J, Treit PV, Shi H, Rosenberger FA, Dethlefsen L,

Meng X, Yaffe E, Aranda-Díaz A, Geyer PE, Mueller-Reif JB, Spencer S, Patterson AD,

Triadafilopoulos G, Holmes SP, Mann M, Fiehn O, Relman DA, Huang KC. 2023. Profiling

the human intestinal environment under physiological conditions. Nature

https://doi.org/10.1038/s41586-023-05989-7.

21. Khan I, Bai Y, Ullah N, Liu G, Rajoka MSR, Zhang C. 2022. Differential Susceptibility of the

Gut Microbiota to DSS Treatment Interferes in the Conserved Microbiome Association in

Mouse Models of Colitis and Is Related to the Initial Gut Microbiota Difference. Advanced

Gut & Microbiome Research 2022:7813278.

22. Leshem A, Segal E, Elinav E. 2020. The Gut Microbiome and Individual-Specific

Responses to Diet. mSystems 5:e00665–20.

23. Gibbons SM, Gurry T, Lampe JW, Chakrabarti A, Dam V, Everard A, Goas A, Gross G,

Kleerebezem M, Lane J, Maukonen J, Penna ALB, Pot B, Valdes AM, Walton G, Weiss A,

Zanzer YC, Venlet NV, Miani M. 2022. Perspective: Leveraging the Gut Microbiota to

Predict Personalized Responses to Dietary, Prebiotic, and Probiotic Interventions. Adv Nutr

13:1450–1461.

24. Lewis JD, Abreu MT. 2017. Diet as a Trigger or Therapy for Inflammatory Bowel Diseases.

Gastroenterology 152:398–414.e6.

25. Limdi JK. 2018. Dietary practices and inflammatory bowel disease. Indian J Gastroenterol

37:284–292.

26. Gutiérrez A, Scharl M, Sempere L, Holler E, Zapater P, Almenta I, González-Navajas JM,

Such J, Wiest R, Rogler G, Francés R. 2014. Genetic susceptibility to increased bacterial

translocation influences the response to biological therapy in patients with Crohn's disease.

Gut 63:272–280.

27. Buisine MP, Desreumaux P, Debailleul V, Gambiez L, Geboes K, Ectors N, Delescaut MP,

Degand P, Aubert JP, Colombel JF, Porchet N. 1999. Abnormalities in mucin gene

expression in Crohn's disease. Inflamm Bowel Dis 5:24–32.

28. Söderholm JD, Olaison G, Peterson KH, Franzén LE, Lindmark T, Wirén M, Tagesson C,

Sjödahl R. 2002. Augmented increase in tight junction permeability by luminal stimuli in the

non-inflamed ileum of Crohn's disease. Gut 50:307–313.

29. Farrugia A, Arasaradnam R. 2021. Bile acid diarrhoea: pathophysiology, diagnosis and

management. Frontline Gastroenterol 12:500–507.

30.  Eckburg PB, Relman DA. 2007. The role of microbes in Crohn's disease. Clin Infect Dis

44:256–262.

31.  Opipari A, Franchi L. 2015. Role of inflammasomes in intestinal inflammation and Crohn's

disease. Inflamm Bowel Dis 21:173–181.

32.  Weaver CT, Hatton RD. 2009. Interplay between the TH17 and TReg cell lineages: a (co-

)evolutionary perspective. Nat Rev Immunol 9:883–889.

33.  Nieminen U, Jussila A, Nordling S, Mustonen H, Färkkilä MA. 2014. Inflammation and

disease duration have a cumulative effect on the risk of dysplasia and carcinoma in IBD: A

case-control observational study based on registry data. International Journal of Cancer

134:189–196.

34.  Kaakoush NO, Day AS, Huinao KD, Leach ST, Lemberg DA, Dowd SE, Mitchell HM. 2012.

Microbial dysbiosis in pediatric patients with Crohn's disease. J Clin Microbiol 50:3258–

3266.

35.  Zhuang X, Liu C, Zhan S, Tian Z, Li N, Mao R, Zeng Z, Chen M. 2021. Gut Microbiota

Profile in Pediatric Patients With Inflammatory Bowel Disease: A Systematic Review. Front

Pediatr 9:626232.

36.  Kowalska-Duplaga K, Gosiewski T, Kapusta P, Sroka-Oleksiak A, Wędrychowicz A,

Pieczarkowski S, Ludwig-Słomczyńska AH, Wołkow PP, Fyderek K. 2019. Differences in

the intestinal microbiome of healthy children and patients with newly diagnosed Crohn's

disease. Sci Rep 9:1–11.

37.  Hansen T, Duerksen DR. 2018. Enteral Nutrition in the Management of Pediatric and Adult

Crohn's Disease. Nutrients 10.

38. Zhang T, Holman J, McKinstry D, Trindade BC, Eaton KA, Mendoza-Castrejon J, Ho S,

Wells E, Yuan H, Wen B, Sun D, Chen GY, Li Y. 2022. A steamed broccoli sprout diet

preparation that reduces colitis via the gut microbiota. J Nutr Biochem 112:109215.

39. Narbad A, Rossiter JT. 2018. Gut Glucosinolate Metabolism and Isothiocyanate Production.

Mol Nutr Food Res 62:e1700991.

40. Shapiro TA, Fahey JW, Wade KL, Stephenson KK, Talalay P. 2001. Chemoprotective

glucosinolates and isothiocyanates of broccoli sprouts: metabolism and excretion in

humans. Cancer Epidemiol Biomarkers Prev 10:501–508.

41. Wei L-Y, Zhang J-K, Zheng L, Chen Y. 2022. The functional role of sulforaphane in

intestinal inflammation: a review. Food Funct 13:514–529.

42. Mueller K, Blum NM, Mueller AS. 2013. Examination of the Anti-Inflammatory, Antioxidant,

and Xenobiotic-Inducing Potential of Broccoli Extract and Various Essential Oils during a

Mild DSS-Induced Colitis in Rats. ISRN Gastroenterology

https://doi.org/10.1155/2013/710856.

43. Liu T, Zhang L, Joo D, Sun S-C. 2017. NF-κB signaling in inflammation. Signal Transduct

Target Ther 2:17023.

44. Burnett JP, Lim G, Li Y, Shah RB, Lim R, Paholak HJ, McDermott SP, Sun L, Tsume Y, Bai

S, Wicha MS, Sun D, Zhang T. 2017. Sulforaphane enhances the anticancer activity of

taxanes against triple negative breast cancer by killing cancer stem cells. Cancer Lett

394:52–64.

45. Holman J, Colucci L, Baudewyns D, Balkan J, Hunt T, Hunt B, Kinney M, Holcomb L, Chen

G, Moses PL, Mawe GM, Zhang T, Li Y, Ishaq SL. 2023. Steamed broccoli sprouts alleviate

DSS-induced inflammation and retain gut microbial biogeography in mice.

46. Li Y, Zhang T, Holman J, McKinstry D, Trindade BC, Mendoza-Castrejon J, Ho S, Sun D,

Wells E, Chen G. 2022. A Specific Broccoli Sprout Preparation Reduces Chemically-

Induced Colitis Via Gut Microbiota. Curr Dev Nutr 6:307–307.

47. Zhang Y, Tan L, Li C, Wu H, Ran D, Zhang Z. 2020. Sulforaphane alter the microbiota and

mitigate colitis severity on mice ulcerative colitis induced by DSS. AMB Express 10:119.

48. Fahey JW, Haristoy X, Dolan PM, Kensler TW, Scholtus I, Stephenson KK, Talalay P,

Lozniewski A. 2002. Sulforaphane inhibits extracellular, intracellular, and antibiotic-resistant

strains of *Helicobacter pylori* and prevents benzo[*a*]pyrene-induced stomach tumors. Proc

Natl Acad Sci U S A 99:7610–7615.

49. Abukhabta S, Khalil Ghawi S, Karatzas KA, Charalampopoulos D, McDougall G, Allwood

JW, Verrall S, Lavery S, Latimer C, Pourshahidi LK, Lawther R, O'Connor G, Rowland I,

Gill CIR. 2021. Sulforaphane-enriched extracts from glucoraphanin-rich broccoli exert

antimicrobial activity against gut pathogens in vitro and innovative cooking methods

increase in vivo intestinal delivery of sulforaphane. Eur J Nutr 60:1263–1276.

50. Yuanfeng W, Chengzhi L, Ligen Z, Juan S, Xinjie S, Yao Z, Jianwei M. 2021. Approaches

for enhancing the stability and formation of sulforaphane. Food Chem 345:128771.

51. Yagishita Y, Fahey JW, Dinkova-Kostova AT, Kensler TW. 2019. Broccoli or Sulforaphane:

Is It the Source or Dose That Matters? Molecules 24:3593.

52. Eichele DD, Kharbanda KK. 2017. Dextran sodium sulfate colitis murine model: An

indispensable tool for advancing our understanding of inflammatory bowel diseases

pathogenesis. World J Gastroenterol 23:6016–6029.

53. Wirtz S, Popp V, Kindermann M, Gerlach K, Weigmann B, Fichtner-Feigl S, Neurath MF.

2017. Chemically induced mouse models of acute and chronic intestinal inflammation. Nat

Protoc 12:1295–1309.

54. Hou JK, Lee D, Lewis J. 2014. Diet and inflammatory bowel disease: review of patient-

targeted recommendations. Clin Gastroenterol Hepatol 12:1592–1600.

55. Walker A, Schmitt-Kopplin P. 2021. The role of fecal sulfur metabolome in inflammatory

bowel diseases. Int J Med Microbiol 311:151513.

56. Wright EK, Kamm MA, Teo SM, Inouye M, Wagner J, Kirkwood CD. 2015. Recent

advances in characterizing the gastrointestinal microbiome in Crohn's disease: a

systematic review. Inflamm Bowel Dis 21:1219–1228.

57. Chiodini RJ, Dowd SE, Chamberlin WM, Galandiuk S, Davis B, Glassing A. 2015. Microbial

Population Differentials between Mucosal and Submucosal Intestinal Tissues in Advanced

Crohn's Disease of the Ileum. PLoS One 10:e0134382.

58. Pascal V, Pozuelo M, Borruel N, Casellas F, Campos D, Santiago A, Martinez X, Varela E,

Sarrabayrouse G, Machiels K, Vermeire S, Sokol H, Guarner F, Manichanh C. 2017. A

microbial signature for Crohn's disease. Gut 66:813–822.

59. Ghazi LJ. 2023. Crohn Disease. Medscape 172940.

60. Rennick DM, Fort MM. 2000. XII. IL-10-deficient (IL-10–/–) mice and intestinal

inflammation. Am J Physiol Gastrointest Liver Physiol 278:G829–G833.

61. Keubler LM, Buettner M, Häger C, Bleich A. 2015. A Multihit Model: Colitis Lessons from

the Interleukin-10-deficient Mouse. Inflamm Bowel Dis 21:1967–1975.

62. Duell BL, Tan CK, Carey AJ, Wu F, Cripps AW, Ulett GC. 2012. Recent insights into

microbial triggers of interleukin-10 production in the host and the impact on infectious

disease pathogenesis. FEMS Immunol Med Microbiol 64:295–313.

63. Glocker E-O, Kotlarz D, Klein C, Shah N, Grimbacher B. 2011. IL-10 and IL-10 receptor

defects in humans. Ann N Y Acad Sci 1246:102–107.

64. Burich A, Hershberg R, Waggie K, Zeng W, Brabb T, Westrich G, Viney JL, Maggio-Price L.

2001. *Helicobacter*-induced inflammatory bowel disease in IL-10- and T cell-deficient mice.

Am J Physiol Gastrointest Liver Physiol 281:G764–78.

65. Kühn R, Löhler J, Rennick D, Rajewsky K, Müller W. 1993. Interleukin-10-deficient mice

develop chronic enterocolitis. Cell 75:263–274.

66. Kennedy RJ, Hoper M, Deodhar K, Erwin PJ, Kirk SJ, Gardiner KR. 2000. Interleukin 10-

deficient colitis: new similarities to human inflammatory bowel disease. Br J Surg 87:1346–

1351.

67. Mähler M, Leiter EH. 2002. Genetic and environmental context determines the course of

colitis developing in IL-10-deficient mice. Inflamm Bowel Dis 8:347–355.

68. Knoop KA, Gustafsson JK, McDonald KG, Kulkarni DH, Coughlin PE, McCrate S, Kim D,

Hsieh C-S, Hogan SP, Elson CO, Tarr PI, Newberry RD. 2017. Microbial antigen encounter

during a preweaning interval is critical for tolerance to gut bacteria. Sci Immunol

2:eaao1314.

69. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, Ross MC, Lloyd

RE, Doddapaneni H, Metcalf GA, Muzny D, Gibbs RA, Vatanen T, Huttenhower C, Xavier

RJ, Rewers M, Hagopian W, Toppari J, Ziegler A-G, She J-X, Akolkar B, Lernmark A, Hyoty

H, Vehik K, Krischer JP, Petrosino JF. 2018. Temporal development of the gut microbiome

in early childhood from the TEDDY study. Nature 562:583–588.

70. Fahey JW, Zhang Y, Talalay P. 1997. Broccoli sprouts: an exceptionally rich source of

inducers of enzymes that protect against chemical carcinogens. Proc Natl Acad Sci U S A

94:10367–10372.

71. Chassaing B, Srinivasan G, Delgado MA, Young AN, Gewirtz AT, Vijay-Kumar M. 2012.

Fecal Lipocalin 2, a Sensitive and Broadly Dynamic Non-Invasive Biomarker for Intestinal

Inflammation. PLoS One 7:e44328.

72. Nowicki D, Maciąg-Dorszyńska M, Bogucka K, Szalewska-Pałasz A, Herman-Antosiewicz

A. 2019. Various modes of action of dietary phytochemicals, sulforaphane and phenethyl

isothiocyanate, on pathogenic bacteria. Sci Rep 9:13677.

73. Liou CS, Sirk SJ, Diaz CAC, Klein AP, Fischer CR, Higginbottom SK, Erez A, Donia MS,

Sonnenburg JL, Sattely ES. 2020. A Metabolic Pathway for Activation of Dietary

Glucosinolates by a Human Gut Symbiont. Cell 180:717–728.e19.

74. Baumgart DC, Sandborn WJ. 2012. Crohn's disease. Lancet 380:1590–1605.

75. Knoch B, Nones K, Barnett MPG, McNabb WC, Roy NC. 2010. Diversity of caecal bacteria

is altered in interleukin-10 gene-deficient mice before and after colitis onset and when fed

polyunsaturated fatty acids. Microbiology 156:3306–3316.

76. Overstreet A-MC, Ramer-Tait AE, Suchodolski JS, Hostetter JM, Wang C, Jergens AE,

Phillips GJ, Wannemuehler MJ. 2020. Temporal Dynamics of Chronic Inflammation on the

Cecal Microbiota in IL-10-/- Mice. Front Immunol 11:585431.

77. Berding K, Holscher HD, Arthur AE, Donovan SM. 2018. Fecal microbiome composition

and stability in 4- to 8-year old children is associated with dietary patterns and nutrient

intake. J Nutr Biochem 56:165–174.

78. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, Almeida M, Arumugam M,

Batto J-M, Kennedy S, Leonard P, Li J, Burgdorf K, Grarup N, Jørgensen T, Brandslund I,

Nielsen HB, Juncker AS, Bertalan M, Levenez F, Pons N, Rasmussen S, Sunagawa S, Tap J, Tims S, Zoetendal EG, Brunak S, Clément K, Doré J, Kleerebezem M, Kristiansen K, Renault P, Sicheritz-Ponten T, de Vos WM, Zucker J-D, Raes J, Hansen T, Bork P, Wang J, Ehrlich SD, Pedersen O. 2013. Richness of human gut microbiome correlates with metabolic markers. Nature 500:541–546.

79. Park JH, Kwon JG, Kim SJ, Song DK, Lee SG, Kim ES, Cho KB, Jang BI, Kim DH, Sin J-I, Kim TW, Song IH, Park KS. 2015. Alterations of colonic contractility in an interleukin-10 knockout mouse model of inflammatory bowel disease. J Neurogastroenterol Motil 21:51–61.

80. Cohen SL, Moore AM, Ward WE. 2004. Interleukin-10 knockout mouse: a model for studying bone metabolism during intestinal inflammation. Inflamm Bowel Dis 10:557–563.

81. Veeranki OL, Bhattacharya A, Marshall JR, Zhang Y. 2013. Organ-specific exposure and response to sulforaphane, a key chemopreventive ingredient in broccoli: implications for cancer prevention. Br J Nutr 109:25–32.

82. Okayasu I, Hatakeyama S, Yamada M, Ohkusa T, Inagaki Y, Nakaya R. 1990. A novel method in the induction of reliable experimental acute and chronic ulcerative colitis in mice. Gastroenterology 98:694–702.

83. Poritz LS, Garver KI, Green C, Fitzpatrick L, Ruggiero F, Koltun WA. 2007. Loss of the tight junction protein ZO-1 in dextran sulfate sodium induced colitis. J Surg Res 140:12–19.

84. Chassaing B, Aitken JD, Malleshappa M, Vijay-Kumar M. 2014. Dextran sulfate sodium (DSS)-induced colitis in mice. Curr Protoc Immunol 104:15.25.1–15.25.14.

85. Araki A, Kanai T, Ishikura T, Makita S, Uraushihara K, Iiyama R, Totsuka T, Takeda K, Akira S, Watanabe M. 2005. MyD88-deficient mice develop severe intestinal inflammation

in dextran sodium sulfate colitis. J Gastroenterol 40:16–23.

86.   Dieleman LA, Ridwan BU, Tennyson GS, Beagley KW, Bucy RP, Elson CO. 1994. Dextran

sulfate sodium-induced colitis occurs in severe combined immunodeficient mice.

Gastroenterology 107:1643–1652.

87.   Bhattacharyya S, Dudeja PK, Tobacman JK. 2009. ROS, Hsp27, and IKKbeta mediate

dextran sodium sulfate (DSS) activation of IkappaBa, NFkappaB, and IL-8. Inflamm Bowel

Dis 15:673–683.

88.   Yang Y, Jia H, Lyu W, Furukawa K, Li X, Hasebe Y, Kato H. 2022. Dietary Eggshell

Membrane Powder Improves Survival Rate and Ameliorates Gut Dysbiosis in Interleukin-10

Knockout Mice. Front Nutr 9:895665.

89.   Wang D, Jin H, Sheng J, Cheng L, Lin Q, Lazerev M, Jin P, Li X. 2022. A high salt diet

protects interleukin 10-deficient mice against chronic colitis by improving the mucosal

barrier function. Mol Immunol 150:39–46.

90.   Zhang Z, Hyun JE, Thiesen A, Park H, Hotte N, Watanabe H, Higashiyama T, Madsen KL.

2020. Sex-Specific Differences in the Gut Microbiome in Response to Dietary Fiber

Supplementation in IL-10-Deficient Mice. Nutrients 12:2088.

91.   Kondo H, Abe I, Gotoh K, Fukui A, Takanari H, Ishii Y, Ikebe Y, Kira S, Oniki T, Saito S,

Aoki K, Tanino T, Mitarai K, Kawano K, Miyoshi M, Fujinami M, Yoshimura S, Ayabe R,

Okada N, Nagano Y, Akioka H, Shinohara T, Akiyoshi K, Masaki T, Teshima Y, Yufu K,

Nakagawa M, Takahashi N. 2018. Interleukin 10 Treatment Ameliorates High-Fat Diet-

Induced Inflammatory Atrial Remodeling and Fibrillation. Circ Arrhythm Electrophysiol

11:e006040.

92.   Kowalski GM, Nicholls HT, Risis S, Watson NK, Kanellakis P, Bruce CR, Bobik A,

Lancaster GI, Febbraio MA. 2011. Deficiency of haematopoietic-cell-derived IL-10 does not

exacerbate high-fat-diet-induced inflammation or insulin resistance in mice. Diabetologia

54:888–899.

93.  Martínez I, Maldonado-Gomez MX, Gomes-Neto JC, Kittana H, Ding H, Schmaltz R,

Joglekar P, Cardona RJ, Marsteller NL, Kembel SW, Benson AK, Peterson DA, Ramer-Tait

AE, Walter J. 2018. Experimental evaluation of the importance of colonization history in

early-life gut microbiota assembly. Elife 7:e36521.

94.  Kemp KM, Colson J, Lorenz RG, Maynard CL, Pollock JS. 2021. Early life stress in mice

alters gut microbiota independent of maternal microbiota inheritance. Am J Physiol Regul

Integr Comp Physiol 320:R663–R674.

95.  Knoop KA, McDonald KG, Coughlin PE, Kulkarni DH, Gustafsson JK, Rusconi B, John V,

Ndao IM, Beigelman A, Good M, Warner BB, Elson CO, Hsieh C-S, Hogan SP, Tarr PI,

Newberry RD. 2020. Synchronization of mothers and offspring promotes tolerance and

limits allergy. JCI Insight 5:e137943.

96.  Schloss PD, Schubert AM, Zackular JP, Iverson KD, Young VB, Petrosino JF. 2012.

Stabilization of the murine gut microbiome following weaning. Gut Microbes 3:383–393.

97.  Delage C, Taib T, Mamma C, Lerouet D, Besson VC. 2021. Traumatic Brain Injury: An Age-

Dependent View of Post-Traumatic Neuroinflammation and Its Treatment. Pharmaceutics

13:1624.

98.  Bäckhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, Li Y, Xia Y, Xie H,

Zhong H, Khan MT, Zhang J, Li J, Xiao L, Al-Aama J, Zhang D, Lee YS, Kotowska D,

Colding C, Tremaroli V, Yin Y, Bergman S, Xu X, Madsen L, Kristiansen K, Dahlgren J,

Wang J. 2015. Dynamics and Stabilization of the Human Gut Microbiome during the First

Year of Life. Cell Host Microbe 17:852.

99.  Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, Clemente

JC, Knight R, Heath AC, Leibel RL, Rosenbaum M, Gordon JI. 2013. The Long-Term

Stability of the Human Gut Microbiota. Science 341:1237439.

100.  Tropini C, Earle KA, Huang KC, Sonnenburg JL. 2017. The Gut Microbiome: Connecting

Spatial Organization to Function. Cell Host Microbe 21:433–442.

101.  Kadriu H, Yeoman C. 2022. The Unseen Minority: Biogeographical Investigations of the

Ruminant Gastrointestinal Microbiome Highlight the Importance of Frequently Ignored

Anatomical Regions, p. 179–191. *In* Kogut, MH, Zhang, G (eds.), Gut Microbiota, Immunity,

and Health in Production Animals. Springer International Publishing, Cham, Switzerland.

102.  Suzuki TA, Nachman MW. 2016. Spatial Heterogeneity of Gut Microbial Composition

along the Gastrointestinal Tract in Natural Populations of House Mice. PLoS One

11:e0163720.

103.  Yeoman CJ, Ishaq SL, Bichi E, Olivo SK, Lowe J, Aldridge BM. 2018. Biogeographical

differences in the influence of maternal microbial sources on the early successional

development of the bovine neonatal gastrointestinal tract. Sci Rep 8:3197.

104.  Hildebrand F, Nguyen TLA, Brinkman B, Yunta RG, Cauwe B, Vandenabeele P, Liston

A, Raes J. 2013. Inflammation-associated enterotypes, host genotype, cage and inter-

individual effects drive gut microbiota variation in common laboratory mice. Genome Biol

14:R4.

105.  Snipes RL. 1981. Anatomy of the cecum of the laboratory mouse and rat. Anat Embryol

162:455–474.

106.    Gordon JH, Dubos R. 1970. The anaerobic bacterial flora of the mouse cecum. J Exp

Med 132:251–260.

107.    Boger-May A, Reed T, LaTorre D, Ruley-Haase K, Hoffman H, English L, Roncagli C,

Overstreet A-M, Boone D. 2022. Altered microbial biogeography in an innate model of

colitis. Gut Microbes 14:2123677.

108.    Kozik AJ, Nakatsu CH, Chun H, Jones-Hall YL. 2019. Comparison of the fecal, cecal,

and mucus microbiome in male and female mice after TNBS-induced colitis. PLoS One

14:e0225079.

109.    Hubbard TD, Murray IA, Nichols RG, Cassel K, Podolsky M, Kuzu G, Tian Y, Smith P,

Kennett MJ, Patterson AD, Perdew GH. 2017. Dietary Broccoli Impacts Microbial

Community Structure and Attenuates Chemically Induced Colitis in Mice in an Ah receptor

dependent manner. J Funct Foods 37:685–698.

110.    Sokol H, Pigneur B, Watterlot L, Lakhdari O, Bermúdez-Humarán LG, Gratadoux J-J,

Blugeon S, Bridonneau C, Furet J-P, Corthier G, Grangette C, Vasquez N, Pochart P,

Trugnan G, Thomas G, Blottière HM, Doré J, Marteau P, Seksik P, Langella P. 2008.

*Faecalibacterium prausnitzii* is an anti-inflammatory commensal bacterium identified by gut

microbiota analysis of Crohn disease patients. Proc Natl Acad Sci U S A 105:16731–16736.

111.    Ridlon JM, Devendran S, Alves JM, Doden H, Wolf PG, Pereira GV, Ly L, Volland A,

Takei H, Nittono H, Murai T, Kurosawa T, Chlipala GE, Green SJ, Hernandez AG, Fields

CJ, Wright CL, Kakiyama G, Cann I, Kashyap P, McCracken V, Gaskins HR. 2020. The "in

vivo lifestyle" of bile acid 7α-dehydroxylating bacteria: comparative genomics,

metatranscriptomic, and bile acid metabolomics analysis of a defined microbial community

in gnotobiotic mice. Gut Microbes 11:381–404.

112.    Sonnenburg JL, Xu J, Leip DD, Chen C-H, Westover BP, Weatherford J, Buhler JD,

1159            Gordon JI. 2005. Glycan foraging in vivo by an intestine-adapted bacterial symbiont.

1160            Science 307:1955–1959.

1161    113.    Büchler G, Wos-Oxley ML, Smoczek A, Zschemisch N-H, Neumann D, Pieper DH,

1162            Hedrich HJ, Bleich A. 2012. Strain-specific colitis susceptibility in IL10-deficient mice

1163            depends on complex gut microbiota-host interactions. Inflamm Bowel Dis 18:943–954.

1164    114.    Wang R, Halimulati M, Huang X, Ma Y, Li L, Zhang Z. 2022. Sulforaphane-driven

1165            reprogramming of gut microbiome and metabolome ameliorates the progression of

1166            hyperuricemia. J Advert Res In Press.

1167    115.    Sikorska-Zimny K, Beneduce L. 2021. The Metabolism of Glucosinolates by Gut

1168            Microbiota. Nutrients 13:2750.

1169    116.    Luang-In V, Albaser AA, Nueno-Palop C, Bennett MH, Narbad A, Rossiter JT. 2016.

1170            Glucosinolate and Desulfo-glucosinolate Metabolism by a Selection of Human Gut Bacteria.

1171            Curr Microbiol 73:442–451.

1172    117.    Laiz L, Miller AZ, Jurado V, Akatova E, Sanchez-Moral S, Gonzalez JM, Dionísio A,

1173            Macedo MF, Saiz-Jimenez C. 2009. Isolation of five Rubrobacter strains from

1174            biodeteriorated monuments. Naturwissenschaften 96:71–79.

1175    118.    Axelrad JE, Cadwell KH, Colombel J-F, Shah SC. 2021. The role of gastrointestinal

1176            pathogens in inflammatory bowel disease: a systematic review. Therap Adv Gastroenterol

1177            14:17562848211004493.

1178    119.    Hedin CR, McCarthy NE, Louis P, Farquharson FM, McCartney S, Taylor K, Prescott

1179            NJ, Murrells T, Stagg AJ, Whelan K, Lindsay JO. 2014. Altered intestinal microbiota and

1180            blood T cell phenotype are shared by patients with Crohn's disease and their unaffected

siblings. Gut 63:1578–1586.

120.   Schwab M, Reynders V, Loitsch S, Steinhilber D, Schröder O, Stein J. 2008. The dietary

histone deacetylase inhibitor sulforaphane induces human beta-defensin-2 in intestinal

epithelial cells. Immunology 125:241–251.

121.   Liu T-C, Gurram B, Baldridge MT, Head R, Lam V, Luo C, Cao Y, Simpson P, Hayward

M, Holtz ML, Bousounis P, Noe J, Lerner D, Cabrera J, Biank V, Stephens M, Huttenhower

C, McGovern DP, Xavier RJ, Stappenbeck TS, Salzman NH. 2016. Paneth cell defects in

Crohn's disease patients promote dysbiosis. JCI Insight 1:e86907.

122.   Ericsson AC, Franklin CL. 2021. The gut microbiome of laboratory mice: considerations

and best practices for translational research. Mamm Genome 32:239–250.

123.   Chichlowski M, Sharp JM, Vanderford DA, Myles MH, Hale LP. 2008. *Helicobacter*

*typhlonius* and *Helicobacter rodentium* differentially affect the severity of colon inflammation

and inflammation-associated neoplasia in IL10-deficient mice. Comp Med 58:534–541.

124.   Makki K, Deehan EC, Walter J, Bäckhed F. 2018. The Impact of Dietary Fiber on Gut

Microbiota in Host Health and Disease. Cell Host Microbe 23:705–715.

125.   Brotherton CS, Martin CA, Long MD, Kappelman MD, Sandler RS. 2016. Avoidance of

Fiber Is Associated With Greater Risk of Crohn's Disease Flare in a 6-Month Period. Clin

Gastroenterol Hepatol 14:1130–1136.

126.   Armstrong HK, Bording-Jorgensen M, Santer DM, Zhang Z, Valcheva R, Rieger AM,

Sung-Ho Kim J, Dijk SI, Mahmood R, Ogungbola O, Jovel J, Moreau F, Gorman H, Dickner

R, Jerasi J, Mander IK, Lafleur D, Cheng C, Petrova A, Jeanson T-L, Mason A, Sergi CM,

Levine A, Chadee K, Armstrong D, Rauscher S, Bernstein CN, Carroll MW, Huynh HQ,

Walter J, Madsen KL, Dieleman LA, Wine E. 2022. Unfermented β-fructan Fibers Fuel

Inflammation in Select Inflammatory Bowel Disease Patients. Gastroenterology

https://doi.org/10.1053/j.gastro.2022.09.034.

127.    He C, Huang L, Lei P, Liu X, Li B, Shan Y. 2018. Sulforaphane Normalizes Intestinal

Flora and Enhances Gut Barrier in Mice with BBN-Induced Bladder Cancer. Mol Nutr Food

Res 62:e1800427.

128.    Hodgkinson K, El Abbar F, Dobranowski P, Manoogian J, Butcher J, Figeys D, Mack D,

Stintzi A. 2023. Butyrate's role in human health and the current progress towards its clinical

application to treat gastrointestinal disease. Clin Nutr 42:61–75.

129.    Hallert C, Björck I, Nyman M, Pousette A, Grännö C, Svensson H. 2003. Increasing fecal

butyrate in ulcerative colitis patients by diet: controlled pilot study. Inflamm Bowel Dis

9:116–121.

130.    O'Flaherty S, Briner Crawley A, Theriot CM, Barrangou R. 2018. The Lactobacillus Bile

Salt Hydrolase Repertoire Reveals Niche-Specific Adaptation. mSphere 3.

131.    Wlodarska M, Luo C, Kolde R, d'Hennezel E, Annand JW, Heim CE, Krastel P, Schmitt

EK, Omar AS, Creasey EA, Garner AL, Mohammadi S, O'Connell DJ, Abubucker S, Arthur

TD, Franzosa EA, Huttenhower C, Murphy LO, Haiser HJ, Vlamakis H, Porter JA, Xavier

RJ. 2017. Indoleacrylic Acid Produced by Commensal Peptostreptococcus Species

Suppresses Inflammation. Cell Host Microbe 22:25–37.e6.

132.    Seifert HA, Gerstner G, Kent G, Vandenbark AA, Offner H. 2019. Estrogen-induced

compensatory mechanisms protect IL-10-deficient mice from developing EAE. J

Neuroinflammation 16:195.

133.    Bábíčková J, Tóthová Ľ, Lengyelová E, Bartoňová A, Hodosy J, Gardlík R, Celec P.

2015. Sex Differences in Experimentally Induced Colitis in Mice: a Role for Estrogens.

Inflammation 38:1996–2006.

134.    Jašarević E, Howerton CL, Howard CD, Bale TL. 2015. Alterations in the Vaginal

Microbiome by Maternal Stress Are Associated With Metabolic Reprogramming of the

Offspring Gut and Brain. Endocrinology 156:3265–3276.

135.    Park KT, Ehrlich OG, Allen JI, Meadows P, Szigethy EM, Henrichsen K, Kim SC, Lawton

RC, Murphy SM, Regueiro M, Rubin DT, Engel-Nitz NM, Heller CA. 2020. The Cost of

Inflammatory Bowel Disease: An Initiative From the Crohn's & Colitis Foundation. Inflamm

Bowel Dis 26:1–10.

136.    Singh S, Qian AS, Nguyen NH, Ho SKM, Luo J, Jairath V, Sandborn WJ, Ma C. 2022.

Trends in U.S. Health Care Spending on Inflammatory Bowel Diseases, 1996-2016.

Inflamm Bowel Dis 28:364–372.

137.    Peery AF, Crockett SD, Murphy CC, Lund JL, Dellon ES, Williams JL, Jensen ET,

Shaheen NJ, Barritt AS, Lieber SR, Kochar B, Barnes EL, Fan YC, Pate V, Galanko J,

Baron TH, Sandler RS. 2019. Burden and Cost of Gastrointestinal, Liver, and Pancreatic

Diseases in the United States: Update 2018. Gastroenterology 156:254–272.e11.

138.    Dieleman LA, Peña AS, Meuwissen SG, van Rees EP. 1997. Role of animal models for

the pathogenesis and treatment of inflammatory bowel disease. Scand J Gastroenterol

Suppl 223:99–104.

139.    Roy N, Barnett M, Knoch B, Dommels Y, McNabb W. 2007. Nutrigenomics applied to an

animal model of Inflammatory Bowel Diseases: transcriptomic analysis of the effects of

eicosapentaenoic acid- and arachidonic acid-enriched diets. Mutat Res 622:103–116.

140.    Gong W, Yu J, Zheng T, Liu P, Zhao F, Liu J, Hong Z, Ren H, Gu G, Wang G, Wu X,

Zhao Y, Ren J. 2021. CCL4-mediated targeting of spleen tyrosine kinase (Syk) inhibitor

using nanoparticles alleviates inflammatory bowel disease. Clin Transl Med 11:e339.

141.   Yu W, Li Q, Shao C, Zhang Y, Kang C, Zheng Y, Liu X, Liu X, Yan J. 2022. The Cao-

Xiang-Wei-Kang formula attenuates the progression of experimental colitis by restoring the

homeostasis of the microbiome and suppressing inflammation. Front Pharmacol

13:946065.

142.   Prakash T, Janadri S. 2022. Anti-inflammatory effect of wedelolactone on DSS induced

colitis in rats: IL-6/STAT3 signaling pathway. J Ayurveda Integr Med 100544.

143.   Gobert AP, Al-Greene NT, Singh K, Coburn LA, Sierra JC, Verriere TG, Luis PB,

Schneider C, Asim M, Allaman MM, Barry DP, Cleveland JL, Destefano Shields CE,

Casero RA Jr, Washington MK, Piazuelo MB, Wilson KT. 2018. Distinct Immunomodulatory

Effects of Spermine Oxidase in Colitis Induced by Epithelial Injury or Infection. Front

Immunol 9:1242.

144.   Cardiff RD, Miller CH, Munn RJ. 2014. Manual hematoxylin and eosin staining of mouse

tissue sections. Cold Spring Harb Protoc 2014:655–658.

145.   Spohn SN, Bianco F, Scott RB, Keenan CM, Linton AA, O'Neill CH, Bonora E, Dicay M,

Lavoie B, Wilcox RL, MacNaughton WK, De Giorgio R, Sharkey KA, Mawe GM. 2016.

Protective Actions of Epithelial 5-Hydroxytryptamine 4 Receptors in Normal and Inflamed

Colon. Gastroenterology 151:933–944.e3.

146.   Fadrosh DW, Ma B, Gajer P, Sengamalay N, Ott S, Brotman RM, Ravel J. 2014. An

improved dual-indexing approach for multiplexed 16S rRNA gene sequencing on the

Illumina MiSeq platform. Microbiome 2:6.

147.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ,

Fierer N, Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of

sequences per sample. Proc Natl Acad Sci U S A 108:4516–4522.

148.   2011. The Earth Microbiome Project. The Earth Microbiome Project.

http://www.earthmicrobiome.org.

149.   Callahan B. 2022. DADA2: Fast and accurate sample inference from amplicon data with

single-nucleotide resolution (1.16). https://benjjneb.github.io/dada2/tutorial.html.

150.   RCoreTeam. 2022. R: a language and environment for statistical computing (4.2.2). R

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/.

151.   Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007.

SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA

sequence data compatible with ARB. Nucleic Acids Res 35:7188–7196.

152.   Wickham H, Francois R, Henry L, Müller K. 2015. dplyr: A Grammar of Data

Manipulation. R package version 0.4. 3. R Found Stat Comput , Vienna https://CRAN R-

project org/package= dplyr.

153.   Ishaq SL. 2017. Phyloseq_dealing_with_neg_controls_ Ishaq_example.R. Github.

https://github.com/SueIshaq/Examples-DADA2-Phyloseq. Retrieved 5 March 2019.

154.   McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLoS One 8:e61217.

155.   Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, Lozupone C, Zaneveld

JR, Vázquez-Baeza Y, Birmingham A, Hyde ER, Knight R. 2017. Normalization and

microbial differential abundance strategies depend upon data characteristics. Microbiome

5:27.

156.   Cameron ES, Schmidt PJ, Tremblay BJ-M, Emelko MB, Müller KM. 2021. Enhancing

diversity analysis by repeatedly rarefying next generation sequencing data describing

microbial communities. Sci Rep 11:22302.

157.    Bates D, Mächler M, Zurich E, Bolker BM, Walker SC. 2015. Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software, Articles 67:1–48.

158.    Oksanen J, Guillaume Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin

PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2020.

Vegan: Community Ecology Package (2.5-7). https://cran.r-

project.org/web/packages/vegan/vignettes/diversity-vegan.pdf.

159.    Archer E. 2022. rfpermute: Estimate Permutation p-Values for Random Forest

Importance Metrics (2.5.1). https://github.com/EricArcher/rfPermute.

160.    Wickham H. 2016. ggplot2: Elegant graphics for Data Analysis. Springer Publishing

Company, New York. https://dl.acm.org/citation.cfm?id=1795559.

161.    Kassambara A. 2022. ggpubr:"ggplot2" based publication ready plots (0.1.7).

https://rpkgs.datanovia.com/ggpubr/.

162.    Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, Bushman

FD, Knight R, Kelley ST. 2011. Bayesian community-wide culture-independent microbial

source tracking. Nat Methods 8:761–763.

163.    Galey M. 2021. Mi-Seq Analysis. GitHub. https://mgaley-004.github.io/MiSeq-

Analysis/Tutorials/SourceSink.html, https://github.com/mgaley-004/MiSeq-Analysis.

Retrieved 7 December 2022.

1315

**Figure Legends**

**Figure 1. Experimental design for a dietary prevention model of Crohn's Disease in Interleukin-10 knockout mice beginning at age 4 or 7 weeks.** Mice were raised in a barrier facility and fed a control or 10% raw broccoli sprout diet for 7 days, then for an additional 16 days as they were moved to a conventional room and co-housed with a *Helicobacter hepaticus* positive mouse to induce immune-mediated enterocolitis.

**Figure 2. A 10% raw broccoli sprout diet significantly attenuated the development of colitis in IL-10-KO mice that were A) 4 - 6 weeks old during trial or B) 7 - 9 weeks old during the trial.** Brown lines indicate control diet (n = 4, trial 1; n = 6 trial 2) and green lines indicate sprout diet (n = 5, trial 1; n = 5, trial 2). All mice were in a growth phase, so body weights and time scale were normalized to the day mice were exposed to *H. hepaticus*-positive mice, set at 100% starting weight and Day 0, respectively. DAI scores are calculated by weight loss intensity score, fecal blood, and fecal consistency. Overall model significance is included on the chart, and single timepoint comparisons significance was designated as $p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$ by two-way ANOVA, and post multiple comparisons by Sidak.

**Figure 3**. **Levels of (A) fecal lipocalin during the diet trial and plasma (B) lipocalin, (C) cytokines, and (D) sulforaphane on the last day, in IL-10-KO mice with enterocolitis during consuming 10% (w/w) raw broccoli sprouts or control diets.** Stool samples were collected from mice (n=10/group across both trials) on Days 0 (conventionalization), 10 (onset of symptoms), and 15/16 of experiment for Lipocalin-2 (Lpn/Lcn) concentration determination. Plasma was collected on Days 15/16 for lipocalin, cytokines, and sulforaphane concentration. Data were similar across both trials and combined. For fecal lipocalin, data was normalized by the weight of feces, and significance set as * $p < 0.05$.

**Figure 4. Observed bacterial richness within the gastrointestinal tracts IL-10-KO mice fed control diets or raw broccoli sprout diets from 4-6 weeks of age (Trial 1) and 7-9 weeks of age (Trial 2).** Panel A shows comparison of broccoli-fed and control mice in Trial 1 and Trial 2 mice across the whole gastrointestinal tract ($***p < 0.001$); Panel B shows comparisons of broccoli-fed and control mice in which were 4 - 6 weeks old for the duration of the first trial and 7 - 9 weeks old in the second trial, in each anatomical location scraping site ($*p < 0.05$). Graphics made using phyloseq and ggplot2 packages in R. Significance added by Wilcox tests from the ggsignif package in R.

**Figure 5. Principal coordinates analysis of bacterial community similarity within the gastrointestinal tracts of 4- or 7-week-old IL-10-KO mice fed control diets or broccoli sprout diets.** Calculations using weighted Bray-Curtis dissimilarity show differences in the taxonomic structure of trial 1 mice in Panel A, and trial 2 mice in Panel B. Graphic made using phyloseq, and vegan packages in R.

**Figure 6. Bacterial community dissimilarity within a specific anatomical location along the gastrointestinal tract of IL-10-KO mice fed control diets or broccoli sprout diets beginning at 4 or 7 weeks of age.** Calculations using weighted Bray-Curtis dissimilarity show differences in the taxonomic structure of (A) trial 1 mice sampled at age 6 weeks , and (B) trial 2 mice sampled at age 9 weeks, after each had been consuming diets for 3 weeks. Principal

coordinates analysis graphic made using phyloseq, and vegan packages in R.

**Figure 7. Bacterial SVs of genera which have putative pathogenicity.** Trial 1 (left): 36 gut
samples and 107 SVs; Trial 2 (right): 39 gut samples and 103 SVs. Graphic made using
phyloseq and ggplot2 packages in R.

**Figure 8. Read counts of bacterial SVs of genera which have putative capacity to convert**
**glucoraphanin to sulforaphane (GSL).** Strains of bacteria in these genera have been
demonstrated to perform myrosinase-like activity in the digestive tract, as reviewed in (115).
Trial 1 (left): 36 gut samples and 241 SVs; Trial 2 (right): 39 gut samples and 353 SVs. Graphic
made using phyloseq and ggplot2 packages in R.

**Figure 9. Biogeographic frequency of species from putative GSL-converting genera in**
**the treatment groups.** The Silva Database identified *Bacteroides* species *acidifaciens*,
*caecimuris*, and *sartorii*, *Enterococcus cecorum*, *Staphylococcus aureus*, and *Lactobacillus*
*intestinalis*. NCBI BLASTN identified additional *B. acidifaciens*, *B. caecimuris*, and *B. sartorii.*
BLASTN uniquely identified *Enterococcus* species *casseliflavus* and *faecalis*, and *Lactobacillus*
*gasseri*. *Bacteroides thetaiotaomicron* was linked to two (2) SVs based on genome mapping. *E.*
*cecorum* (frequency 12), *S. aureus* (2), and *B. thetaiotaomicron* (5) were not included in the
figure as their frequencies were minimal.

**Figure 10: Prevalence of the operon BT2159-BT2156 in *Bacteroides thetaiotaomicron***
**(VPI-5482) in 4-6 week-old (Trial 1) and 7-9 week-old (Trial 2) IL-10-KO mice fed either a**
**control or raw broccoli sprout diet.** A. Percentages indicate the number of samples with a
max operon mean copy count in each category (red low <100, purple medium 100-10,000, blue
high >10,000). B. Average high indication across all anatomical sites by trial and diet.
1381

1382

1383 **Tables**

1384

**Table 1. Statistical comparison of principal coordinates analysis of bacterial community**
**similarity within the gastrointestinal tracts of 4- or 7-week-old IL-10-KO mice fed control**
**diets or broccoli sprout diets.** Comparisons were done using dietary treatment and
anatomical locations as factors, and subsetting by trial, as well as by anatomical location, as
noted. Unweighted Jaccard Similarity and weighted Bray-Curtis metrics were used to calculate
beta diversity. Number of permutations = 9999.

1391

| Permanova Tests of Dissimilarity | Df | Test type | Trial 1, aged 4-6 weeks | | Trial 2, aged 7-9 weeks | |
|---|---|---|---|---|---|---|
| | | | F | P-value | F | P-value |
| Diet | 1 | Jac | 2.7779 | 0.0001*** | 1.1898 | 0.0258* |
| | | BC | 7.5867 | 0.0001*** | 2.4498 | 0.0022** |
| Anatomical Location | 3 | Jac | 1.2027 | 0.0291* | 1.0955 | 0.0359* |
| | | BC | 1.5991 | 0.0205* | 1.4958 | 0.0139* |
| Diet: Anatomical Location | 3 | Jac | 1.0704 | 0.1780 | 0.9559 | 0.8477 |
| | | BC | 1.0482 | 0.3504 | 0.8272 | 0.8300 |
| Diet effect within ileum | 1 | Jac | 1.1319 | 0.1242 | 1.0154 | 0.3133 |
| | | BC | 1.4445 | 0.1220 | 1.3366 | 0.1699 |
| Diet effect within cecum | 1 | Jac | 1.9352 | 0.0074** | 1.0393 | 0.2663 |
| | | BC | 4.4036 | 0.0081** | 1.6667 | 0.0818 |
| Diet effect within proximal colon | 1 | Jac | 1.3499 | 0.0554 | 0.9949 | 0.3998 |
| | | BC | 2.0148 | 0.0247* | 0.8776 | 0.6080 |
| Diet effect within distal colon | 1 | Jac | 1.7018 | 0.0261* | 1.0058 | 0.4183 |
| | | BC | 4.1923 | 0.0158* | 0.9797 | 0.4679 |

1392 Significance codes: p ≤ 0.001 *** | p ≤ 0.01 ** | p ≤ 0.05 *
1393 Jaccard dissimilarity = "Jac" | Bray-Curtis dissimilarity = "BC"
1394

1395 **Supplemental Material Legends**

1396 **Supplemental Figures and Table.** Additional data visualizations are included to demonstrate
1397 minor points or null results.

1398 **IL10 mouse R code.** The R code that was used to clean, visualize, and statistically analyze the

1399    bacterial community sequencing data presented.

**A** PCoA of Weighted Bray-Curtis Dissimilarity for Trial 1

**B** PCoA of Weighted Bray-Curtis Dissimilarity for Trial 2

Diet
broccoli
control

**A**    Trial 1 genera with putative pathogenicity

**B**    Trial 2 genera with putative pathogenicity

Genus

Bacillus    Enterococcus    Helicobacter

Clostridioides    Escherichia-Shigella    Staphylococcus

**A** Trial 1 genera with putative capacity for GSL conversion

**B** Trial 2 genera with putative capacity for GSL conversion

Genus
- Bacillus
- Bacteroides
- Enterococcus
- Lactobacillus
- Lactococcus
- Pseudomonas
- Staphylococcus

# A

**C57BL/6 IL10 – Helicobacter**

| Age 4-6 weeks | Ileum contents | Cecum contents | Prox. colon contents | Dist. colon contents |
|---|---|---|---|---|
| Control | 25% | 0% | 50% | 0% |
| | 25% | 80% | 50% | 75% |
| | 50% | 20% | 0% | 25% |
| Broccoli | 20% | 0% | 20% | 0% |
| | 40% | 25% | 40% | 0% |
| | 40% | 75% | 40% | 100% |

| Age 7-9 weeks | Ileum contents | Cecum contents | Prox. colon contents | Dist. colon contents |
|---|---|---|---|---|
| Control | 17% | 33% | 17% | 17% |
| | 17% | 33% | 67% | 83% |
| | 67% | 33% | 17% | 0% |
| Broccoli | 0% | 20% | 20% | 0% |
| | 80% | 60% | 60% | 60% |
| | 20% | 20% | 20% | 40% |

**BT2159-BT2156 operon**
Mean copy number

| 0% | 0% | 0% |
|---|---|---|
| 10% | 10% | 10% |
| 20% | 20% | 20% |
| 30% | 30% | 30% |
| 40% | 40% | 40% |
| 50% | 50% | 50% |
| 60% | 60% | 60% |
| 70% | 70% | 70% |
| 80% | 80% | 80% |
| 90% | 90% | 90% |
| 100% | 100% | 100% |

Low <100     High >10,000

# B

## High Operon Indication ( > 10,000) - Average Across All Anatomical Sites

High Operon Indication (%)

- 4-6 week-old: Broccoli 64%, Control 24%
- 7-9 week-old: Broccoli 25%, Control 29%