**Elevated glucocorticoid alters the developmental dynamics of hypothalamic neurogenesis in**

**zebrafish**

Helen Eachus[1], Min-Kyeung. Choi[1], Anna Tochwin[1], Johanna Kaspareit[2], May Ho[3], Soojin Ryu*[1,2]

* Corresponding author

[1] Living Systems Institute, University of Exeter, Stocker Road, Exeter, EX4 4QD. United Kingdom.

[2] University Medical Center, Johannes Gutenberg University Mainz**,** Duesbergweg 6**,** 55128

Mainz, Germany.

[3] Nanyang Technological University, 50 Nanyang Avenue, Singapore 639798.

**Abstract**

Exposure to excess glucocorticoid (GC) during early development is implicated in adult

dysfunctions. Reduced adult hippocampal neurogenesis is a well-known consequence of exposure

to early life stress or elevated GC, however the effects on neurogenesis during development and

effects on other brain regions are not well understood. Using an optogenetic zebrafish model, here

we analysed the effects of GC exposure on neurogenesis during development in the whole brain.

We identify that the hypothalamus is a highly GC-sensitive region where elevated GC causes

precocious development. This is followed by failed maturation and early decline accompanied by

impaired feeding, growth, and longevity.  In GC-exposed animals, the developmental trajectory of

hypothalamic progenitor cells is strikingly altered, potentially mediated by direct regulation of

transcription factors such as *rx3* by GC. Our data provide cellular and molecular level insight into

GC-induced alteration of the hypothalamic development trajectory, a process crucial for health

across the life-course.

**Introduction**

Glucocorticoids (GC) are the key effectors of the stress response and have pleiotropic effects on the

body, acting to restore homeostasis and thus allow an animal to respond adaptively to a threat[1].

Since the developing brain is plastic, GC exposure during early life has the potential to alter the

developmental trajectory of the brain. Indeed, alteration of the development trajectory may be an

adaptive mechanism employed by animals exposed to stress during early life. For example,

antenatal GC treatment has well documented effects on lung maturation, as well as other organs, in

preterm babies[2]. However, such treatment has also been associated with negative outcomes in later

life, such as increased incidence of mental health issues during childhood[3], impacted development

of fronto-parietal brain functions during adolescence[4], and cortical thinning in children with

associated affective disorders[5]. Early life GC exposure leading to adverse consequences in later life

is an example of the concept known as early life programming of adult disease[6,7,8,9].

One of the most well-documented effects of GC exposure during early life or early life stress (ELS)

is reduced adult neurogenesis in the hippocampus, including reduced cell proliferation[10,11,12,13,14].

Indeed adrenalectomy is sufficient to prevent reduced proliferation in chronically stressed mice,

suggesting that GC is the main driving force behind this phenotype[15]. In contrast, only a few studies

have looked at the early life effects of stress or GC exposure on neurogenesis during development,

but there are some indications that its effects might be developmentally dynamic. Specifically, it is

possible that GC-induced reduction of cell proliferation is a delayed consequence of a GC-induced

change to the development trajectory that began in early life. Indeed, rats exposed to ELS exhibited

enhanced hippocampal cell proliferation and improved stress-associated behavioural performance

as young adults, whilst these effects were reversed by middle age[10]. A potential mechanistic

explanation for the ELS-induced reduction in adult hippocampal neurogenesis is that enhanced cell

proliferation at an earlier time-point depletes the stem cell pool over time[16].

Despite the well-documented effects of GC exposure on hippocampal neurogenesis, the effects of

GC exposure on other brain regions are unclear. Recently, a first study of the effects of ELS on

neurogenesis in the hypothalamus revealed that cell proliferation and numbers of hypothalamic

stem cells known as tanycytes were reduced in the adult mouse[17]. Tanycytes are unique radial glial-

like cells that line the walls of the 3rd ventricle, have NSPC (neural stem/progenitor cell) properties

that persist into adulthood, send projections into neighbouring hypothalamic nuclei and are able to

sense glucose levels in the CSF[18]. Whilst most previous studies have focused on effects of stress

and GC exposure on neurogenesis in the hippocampus, the aforementioned study supports that the

hippocampus is not uniquely affected by stress/GC. However, whether and how GC affects

hypothalamic neurogenesis during development is completely unknown.

Here we used an optogenetic zebrafish model in which endogenous cortisol level is elevated to test

the hypothesis that GC exposure alters the developmental trajectory of the brain. The model utilises

optogenetic manipulation of steroidogenic interrenal cells to increase the endogenous GC level

during larval and juvenile stages[19]. Whilst the mammalian adrenal gland, counterpart of the fish

interrenal[20], is known to produce GCs, mineralocorticoids (MCs), and androgens[21]; MCs and

androgens are not produced by the larval zebrafish interrenal gland[22,23,24,25]. Therefore, the

phenotypes observed in our model are primarily due to the effects of GC and its downstream

effectors, rather than MC or androgen over-exposure.

We found that GC exposure affected cell proliferation in a region-specific manner and that these

effects were dynamic across the life-course. In particular, we found that the hypothalamus is a

highly GC-sensitive brain region. In early life, we observed precocious development of the

hypothalamus and hypothalamus-associated feeding behaviour in GC-exposed animals. This was followed by a rapid decline, indicated by failed hypothalamic maturation and suppressed feeding and growth. Our data uncover a tissue-specific time-dependent plasticity of the hypothalamus in response to GC, and provide cellular and molecular insights for GC-induced alteration of the hypothalamic development trajectory.

**Results**

**Neurogenesis is altered in an optogenetic zebrafish model of elevated GC**

Our lab previously generated an optogenetic zebrafish model, *Tg(star:bPAC-2A-tdTomato)*[uex300], in which the endogenous cortisol level can be increased specifically in response to blue light exposure using an optogenetic protein *beggiatoa* Photoactivated Adenylate Cylase (bPAC) (Fig.1a)[19,26,27]. In order to determine the effects of elevated GC on the brain across development, we raised transgenic fish under standard aquarium lighting conditions, where the blue light component of the white light is sufficient to activate bPAC, which led to significantly elevated cortisol levels in *Tg(star:bPAC-2A-tdTomato)*[uex300] compared with wild types from hatching through until late juvenile/early adult stage (Fig.1b)[19]. *Tg(star:bPAC-2A-tdTomato)*[uex300] raised under standard light conditions will be referred to throughout as star:bPAC+ or bPAC+.

To investigate the effects of elevated GC on brain development we performed bulk RNA-sequencing (RNA-seq) of whole brain samples from star:bPAC+ and wild-type fish at different developmental stages[19]. Gene Ontology (GO) analysis of differentially expressed genes (DEGs) that were down-regulated in star:bPAC+ brains compared with wild types were enriched for a number of biological processes, including Neurogenesis at all developmental stages (GO:0022008, adjusted p-value 2.47E-20 at 6 days post fertilization (dpf), 5.45E-08 at 13 dpf, 1.11E-17 at 120 dpf). Up-regulated genes were not enriched for GO term neurogenesis. To investigate the effects of GC exposure on neurogenesis during development, we focused downstream analysis on the early life time-points: 5/6 dpf and 13 dpf. 225 genes associated with neurogenesis were down-regulated in star:bPAC+ brains at either 6 dpf or 13 dpf (Fig.1c). To confirm our results and identify specific features of neurogenesis that are altered by elevated GC, we performed qPCR on independent samples of whole brains from 5 dpf star:bPAC+ and wild types for genes associated with neurogenesis processes or distinct cell types. Of the 14 categories of genes analysed, only one category was associated with consistent differential expression in star:bPAC+. Expression of *pcna*, *mki67* and *mcm5*, all associated with cell proliferation, showed increased expression in star:bPAC+ compared with wild types (Supplementary Fig.1). These data support that neurogenesis, and especially cell proliferation, is altered by elevated GC.

To identify whether the effect of GC on neurogenesis is developmentally dynamic, we performed
unsupervised clustering analysis of neurogenesis DEGs at 6 dpf and 13 dpf and identified 6
different clusters. Cluster 2 and cluster 5 genes showed a relatively stronger down-regulation at 6
dpf compared to 13 dpf, meanwhile cluster 1 genes showed a relatively stronger downregulation at
13 dpf (Fig.1d). Cluster 3 genes were strongly downregulated at both time-points (Fig.1d).
Interestingly, cluster 4 and cluster 6 genes showed a timepoint-specific down-regulation. Cluster 4
genes were downregulated in star:bPAC+ at 6 dpf but were not differentially expressed at 13 dpf
(Fig.1d). Meanwhile cluster 6 genes were down-regulated in star:bPAC+ at 13 dpf but not at 6 dpf
(Fig.1d). These results suggest that some effects of elevated GC on neurogenesis are temporarily
dynamic during early life. GO analysis of genes within the clusters indicated that all clusters
contained genes associated with neuron differentiation (Fig.1e). Gliogenesis and glial cell
development were enriched in multiple clusters. There were also some terms which were restricted
to single or small numbers of clusters. For example, cluster 4 genes were associated with neuron
fate commitment, cluster 3 with neuron migration, and cluster 2 with neuronal stem cell population
maintenance (Fig.1e). These results suggest that specific aspects of neurogenesis exhibit
temporally dynamic differential expression during early life as a result of GC exposure.
128

**Precocious hypothalamic development following elevated GC**
In order to determine whether the effects of elevated GC on cell proliferation were brain-wide or
region specific we performed immunohistochemistry (IHC) for mitosis marker phospho-histoneH3
(pH3) on brains of 5 dpf star:bPAC+ and wild-type larvae (Fig.2a). Cell counting of proliferating
pH3+ cells across brain regions indicated a trend towards an increase in number of pH3+ cells in
the valvula cerebelli (ce-v) of the hindbrain, however, the only significant increase was restricted to
the hypothalamus (Fig.2b). We next sought to determine which cell types were associated with the
increased proliferative capacity in the hypothalamus of 5 dpf star:bPAC+. To this end, we performed
Fluorescent *in situ* hybridization (FISH) for *rx3* (*retinal homeobox gene 3),* combined with IHC for
Pcna and pH3. *rx3* is a transcription factor known to be expressed in hypothalamic progenitor cells
in zebrafish where it plays a critical role in hypothalamic development[28], and its mammalian
orthologue *Rax* is known to be expressed in hypothalamic radial glia[29]. Pcna and pH3 are markers
for proliferating cells during S-phase and M-phase respectively. Pcna+ cells were situated around
the 3[rd] ventricle of the hypothalamus, especially around the lateral recess, in both star:bPAC+ and
wild-type larvae, with a small number of pH3+ cells scattered throughout this region (Fig.2c). We
distinguish the Pcna+ region around the hypothalamic ventricles as the 'proliferative zone' and
observed co-localisation of *rx3* with Pcna and pH3 within the proliferative zone (Fig.2c, white
arrowheads).
147

Quantitative analysis of confocal stacks revealed that the hypothalamus was significantly larger
(Fig.2f), and we found a significant increase in *rx3* and pcna signal intensity, and significantly more
pH3+ cells in the hypothalamus of 5 dpf star:bPAC+ compared with wild types (Fig.2g-i). To
determine whether the observed increase in hypothalamic proliferation in the hypothalamus of 5 dpf
star:bPAC+ larvae is a direct effect of developmental GC exposure, we treated star:bPAC+ larvae
with the GR antagonist RU-486 from 2 dpf until 5 dpf. Treatment with RU-486 did not significantly
affect hypothalamic volume and elicited only a trend reduction in hypothalamic Pcna, however it
was sufficient to significantly reduce the number of pH3+ mitotic cells in the hypothalamus
(Supplementary Fig.2), suggesting that GR signalling is a significant driver of the increased
hypothalamic proliferation observed in star:bPAC+ larvae. Further, we observed that the excess
hypothalamic pH3+ cells were *rx3*+, meanwhile *rx3*-/pH3+ cells were not different (Fig.2i). These
results suggest that excess proliferating hypothalamic cells in star:bPAC+ are *rx3*-expressing radial
glia. Interestingly, we identified two Glucocorticoid Response Elements (GREs) within the zebrafish
*rx3* gene (Fig.2l). GRE1 is in the 5' UTR, which is most likely a regulatory region. ChIP-qPCR
analysis revealed enrichment of GR antibody (compared with IgG) at both GREs in larval brain
samples, supporting that GR does indeed bind to regulatory regions of the *rx3* gene (Fig.2m).
Further, enrichment of GR was higher in dexamethasone-treated wild-type samples, compared with
wild-type controls, and higher still in star:bPAC+ samples. This experiment shows that GC exposure
directly regulates GR-mediated control of *rx3* gene expression in the brain.
167
Whilst *rx3* appeared to be co-expressed with Pcna in proliferating radial glia, we also analysed Blbp
(brain lipid binding protein, fabp7a) localisation, which is known to be expressed in predominantly
quiescent radial glia in the zebrafish brain[30]. Consistent with this, we observed strong expression of
Blbp in cells lining the third ventricle in the Pcna-negative rostral domains of the hypothalamus,
whilst some Blbp signal was also observed in Pcna-positive cells of the lateral recess (Fig.2d). In 5
dpf star:bPAC+ larvae, we observed increased Blbp signal, especially in the arcuate region of the
hypothalamus (Fig.2d,j), suggesting that increased numbers of glial cells might include both
proliferative and quiescent subtypes. Finally we observed increased signal for Otpa (orthopedia
homeobox a) in the hypothalamus of 5 dpf star:bPAC+ larvae (Fig.2e,k), which is predominantly
expressed in early post-mitotic neuronal precursors[31], suggesting that the excess GC-exposed
hypothalamic proliferative radial glia do exit the cell cycle, leading to increased numbers of neuronal
precursors. Together these data suggest that elevated GC drives precocious development of the
hypothalamus in zebrafish larvae.
181
**Failed hypothalamic maturation following elevated GC**
To test whether the effects of GC exposure on hypothalamic neurogenesis might be temporally
dynamic, we investigated hypothalamic cell proliferation at 13 dpf. In the 13 dpf hypothalamus, *rx3*

was expressed in a subset of Pcna+ cells around the lateral recess, and Blbp was expressed mainly

along the midline 3$^{rd}$ ventricle anterior to the proliferative zone, in a similar manner to at 5 dpf

(Fig.3a). In 13 dpf star:bPAC+ larvae, we observed that the size of the hypothalamus was no longer

larger, which was previously observed at 5 dpf (Fig.3b). Expression of hypothalamic *rx3* was low at

this developmental stage, and showed a trend towards a reduction in star:bPAC+ compared with

wild type (Fig.3a,c). In stark contrast to the 5 dpf hypothalamus, we now observed a dramatic

reduction in expression of Pcna in 13 dpf star:bPAC+ hypothalamus (Fig.3a,d). Meanwhile, Blbp

signal intensity was not different between 13 dpf wild-type and star:bPAC+ larvae (Fig.3a,e).

Strikingly, when we performed co-expression analysis, we observed that whilst only a small

proportion of Blbp+ cells were proliferative in 13 dpf wild-type hypothalamus (~10%), almost all

Blbp+ cells were quiescent in star:bPAC+ (Fig.3f). Further, we observed that in wild types most

hypothalamic *rx3*+ cells (~70%) were proliferative and expressed Pcna, however in star:bPAC+

most *rx3*+ cells (~85%) were quiescent and didn't express Pcna (Fig.3g). These data support that

proliferative radial glia are lost in the hypothalamus of 13 dpf star:bPAC+ larvae. Next, we sought to

identify an explanation for the loss of proliferative radial glia in the hypothalamus. We first analysed

cell death in the hypothalamus of 13 dpf larvae using a TUNEL assay. We did not observe a

significant difference in the number of apoptotic cells in the hypothalamus of 13 dpf star:bPAC+

larvae compared to wild types (Supplementary Fig.3a), suggesting that loss of hypothalamic radial

glia is not due to increased cell death. We reasoned that the loss of proliferation in the star:bPAC+

hypothalamus might be due to senescence of the formerly highly proliferative radial glia. In support

of this hypothesis, we observed that one of the five significantly enriched KEGG pathways in our

whole brain RNA-seq data set in the 13 dpf star:bPAC+ was cellular senescence, a pathway which

was not enriched in the 6 dpf star:bPAC+ samples (Supplementary Fig.3b,c). In the RNA-seq

analysis, 102 of 150 genes associated with the KEGG pathway cellular senescence were

differentially expressed in 13 dpf star:bPAC+ brain samples, meanwhile at 6 dpf only 60 of those

genes were differentially expressed, and they generally showed a lesser degree of fold change

(Supplementary Fig.3d). This suggested that proliferative radial glia might have been lost in the

hypothalamus of 13 dpf star:bPAC+ larvae due to cellular senescence. Additionally, we confirmed

that the loss of hypothalamic proliferation is maintained into later life, since we observed that in

juvenile star:bPAC+ fish, the hypothalamus is smaller and contained fewer mitotic cells

(Supplementary Fig.4a-c). This suggests that the proliferative hypothalamic cells lost at 13 dpf do

not re-enter the cell cycle later in development, suggesting that they have become senescent, rather

than quiescent.

Since we observed a dramatic reduction in cell proliferation in the hypothalamus of 13 dpf

star:bPAC+ larvae, we postulated that downstream neurogenesis and neuronal differentiation might

also be affected. Since one of the hypothalamus-regulated functions is feeding behaviour and food

intake, we sought to determine whether specific neurons that regulate feeding were altered in the

hypothalamus of star:bPAC+ larvae at 13 dpf. *agouti-related peptide* (*agrp*) is expressed in a cluster

of cells in the arcuate nucleus of the hypothalamus, and we observed significantly fewer numbers of

these cells in star:bPAC+ larvae (Fig.3h,i). A small number of cells expressing *cart4* (*cocaine- and*

*amphetamine-regulated transcript*) were observed near to the medial lateral hypothalamus (mLH)

and the total number of these cells were also reduced in star:bPAC+ larvae (Fig.3h,j). Orexin+ (*hcrt*,

*hypocretin*) cells were observed in the dorso-rostral hypothalamus and we observed a trend

reduction in cell count in star:bPAC+ larvae (Fig.3h,k). We observed *npy* (*neuropeptide y*) cells near

the mLH and these cells were not affected in star:bPAC+ larvae (Supplementary Fig.5a,b). Finally,

*pomca* (*proopiomelanocortin*) is also expressed in the arcuate hypothalamic nucleus, and these

cells were not significantly reduced in star:bPAC+ larvae (Supplementary Fig.5a,c). Of the

neuromodulators where we observed reduced numbers in star:bPAC+, *agrp*[32] and *hcrt*[33] are both

known to be appetite-stimulators, meanwhile the effects of *cart* on appetite are more complex[34].

These data indicate that neurogenesis of specific neuronal subtypes that stimulate feeding are

reduced in 13 dpf star:bPAC+ larvae.

**Progenitor cells fail to differentiate in the hypothalamus of star:bPAC+ larvae**

Since we observed that the effects of GC on hypothalamic proliferation changed dramatically

between 5 dpf and 13 dpf, we sought to track the proliferating radial glia of the hypothalamus

between these timepoints to determine their fate. To do this we exposed 4 dpf larvae to BrDU

(Bromodeoxyuridine) overnight for 17 hours until 5 dpf and then fixed larvae at 5, 8, 10 or 13 dpf for

double-IHC analysis of BrDU and Pcna (Fig.4a). At 5 dpf we observed successful incorporation of

BrDU into proliferating cells, as observed by co-expression with Pcna, in the hypothalamus of both

star:bPAC+ and wild-type larvae (Fig.4b). By 8, 10 and 13 dpf, BrDU+ cells in the wild-type

hypothalamus have begun exiting the cell cycle and are now mostly observed around the perimeter

of the Pcna-expressing proliferative zone (Fig.4c, Supplementary Fig.6). Indeed, we observed that

the percentage of hypothalamic BrDU+ cells that co-express Pcna drops from around 75% at 5 dpf

to 50-55% at 8-13 dpf (Fig.4d), suggesting that some of the BrDU+ cells that were proliferating at 5

dpf have lost their progenitor cell status and differentiated.

In stark contrast, hypothalamic BrDU+ cells in star:bPAC+ larvae are still predominantly observed

within the Pcna+ proliferative zone after 5 dpf (Fig.4c, Supplementary Fig.6). In fact, the percentage

of hypothalamic BrDU+ cells that co-express Pcna remains at around 100% from 5 dpf until 13 dpf

in star:bPAC+ larvae (Fig.4d). This indicates that in star:bPAC+ larvae, hypothalamic progenitor

cells remain proliferative between 5 dpf and 13 dpf and do not differentiate. Further, we observed an

increase in the number of hypothalamic BrDU+ cells between 5 dpf and 8 dpf in wild types,

supporting that these cells are dividing and therefore increase in number, however no such increase

was observed in star:bPAC+ larvae (Fig.4e). Similarly, the number of hypothalamic Pcna+ cells
increased in wild types between 5 dpf and 10 dpf, indicating that the proliferative zone continues to
expand as cells divide during this period (Fig.4f). No increase in the number of hypothalamic Pcna+
cells was observed in star:bPAC+ larvae during this developmental period (Fig.4f), further
supporting that neurogenesis appears to stall during this developmental window in star:bPAC+
larvae. Together, these data support that in star:bPAC+ larvae, hypothalamic progenitor cells fail to
differentiate, and neurogenesis is stalled during the 5-13 dpf time window.
266

**Functional consequences of elevated GC: precocious feeding followed by physical decline**
The increased size and proliferative capacity of the hypothalamus in 5 dpf star:bPAC+ larvae
suggested that development of the hypothalamus is accelerated in these animals. Therefore, we
hypothesised that behaviours associated with the hypothalamus that develop around this
developmental stage might also be altered and thus we analysed feeding in 5 dpf larvae. Feeding
usually emerges at around 120 hpf (5 dpf) in lab-raised zebrafish larvae, and larvae are capable of
eating small live food. We exposed 5 dpf larvae to live rotifers and saw that star:bPAC+ larvae
consumed significantly more rotifers during the trial (Fig.5a), suggesting that GC-induced
precocious hypothalamic development is accompanied by early emergence of an associated
behaviour, feeding.
277

In light of the reduction of cells expressing feeding-related neuromodulators in the hypothalamus of
13 dpf star:bPAC larvae, we sought to determine whether feeding was also affected at 13 dpf. In
stark contrast to 5 dpf, at 13 dpf star:bPAC+ larvae fed less than wild types (Fig.5b). Interestingly, in
13 dpf wild-type larvae we observed that food intake positively correlated with body size, in that
larger larvae ate more, whilst in star:bPAC+ larvae no significant trend was apparent. (Fig.5c).
Hence, star:bPAC+ larvae consumed less, regardless of their size. Whilst at 5 dpf and 13 dpf we did
not observe any difference in larval body size, at 28 dpf and 120 dpf star:bPAC+ fish were
significantly smaller than wild types (Fig.5d). This suggests that the reduction in food intake in
star:bPAC+ animals from 13 dpf ultimately leads to impaired growth. Finally, we also observed
reduced fertilisation rates and reduced long-term survival of star:bPAC+ fish (Fig.5e,f). Together,
these results point towards potential functional consequences of failed hypothalamic development
after 5 dpf, including impaired behaviour and an early physical decline. Further work is required to
directly link altered hypothalamic neurogenesis with the observed functional alterations in GC-
exposed animals.
292

**Discussion**
This work shows for the first time that elevated glucocorticoid alters the trajectory of hypothalamic
development and function. We identify that the hypothalamus is a highly GC-sensitive region where

elevated GC causes precocious development followed by failed maturation and early decline
accompanied by impaired feeding, growth, and longevity.

Whilst many studies report GC-induced reduction in cell proliferation[35], our data supports the
hypothesis that, at least in some cases, the reduced cell proliferation following GC exposure might
be a consequence of a prior increase in proliferation. Indeed the concept of GC-inducible stem cells
is not novel[36,37], and there are many reports of a GC-induced increase in cell proliferation of various
cell types including NSPCs[38,39,40,41]. The initial GC-induced increase in cell proliferation observed
here might be in part mediated by mechanisms akin to those observed during brain injury, whereby
formerly quiescent radial glia enter the cell cycle in response to trauma[30,42]. In the published
literature, there are very few previous studies of effects of GC on cell proliferation across
development. Of those, Noorlander *et al.*, observed that antenatal GC treatment in mice led to an
initial reduction in cell proliferation in the embryonic hippocampus, followed by increased cell
proliferation during the postnatal period and a subsequent reduction in proliferation during
adulthood[43]. There are also examples in mouse models where ELS induced an initial increase in cell
proliferation in the hippocampus during early life which was again followed by a subsequent
reduction in cell proliferation[10] or hippocampal volume[12] in later life. Further, a meta-analysis of age-
dependent effects of ELS on cell proliferation shows a negative correlation between age and
change in cell proliferation, with the ELS-exposed animals analysed earliest in development
exhibiting increased cell proliferation and older ELS-exposed animals exhibiting a reduction in cell
proliferation[44]. It is possible that our interpretation of the results presented here also applies to other
studies, in that GC drives precocious proliferation leading to failed maturation and a rapid decline.

In our model, we observed profound effects on a population of NSPCs within the proliferative zone
of the developing hypothalamus. Indeed this population of hypothalamic radial glia might be akin to
tanycyte cells observed in mammals, which also express *Rax/rx3* and *Blbp*[18], and were recently
shown to be sensitive to ELS exposure in adult mice[17]. We postulate that these cells might be
especially sensitive to the effects of elevated GC, since they are adjacent to the 3$^{rd}$ ventricle and
thus might detect elevated GC levels within the CSF. Indeed, neurogenesis of tanycyte cells is even
responsive to dietary cues[45]. Another potential driver of increased sensitivity of the hypothalamus to
GC would be increased levels of GR, however, it is not currently clear whether the hypothalamus
expresses higher levels of GR than other brain regions in zebrafish, or indeed in other vertebrates.
As such, an explanation for the specificity of the proliferation phenotype observed in our study
remains elusive. We cannot rule-out the possibility that the hypothalamus is not uniquely affected in
our model, but rather that the effects of GC on different brain regions might emerge across different
time-scales.

We observed significant effects on feeding behaviour of developmental GC-exposed star:bPAC+
larvae, whereby 5 dpf larvae consumed more and 13 dpf larvae consumed less than controls. The
observed reduction in feeding persists into adulthood in this model[19]. These changes correlated with
increased proliferative hypothalamic radial glia at 5 dpf, which were reduced by 13 dpf, alongside a
reduction in numbers of feeding-regulating neurons that express *agrp*, *cart* and *hcrt at* 13 dpf. The
role of these hypothalamic neuropeptides in food consumption in mammals is well established[46,47,48].
In larval zebrafish, hypothalamic *agrp* neurons are known to stimulate feeding[32], and orexin,
produced by *hcrt* neurons, is also known to stimulate food intake in zebrafish[33]. As such, it is likely
that the observed reduction in numbers of *agrp* and *hcrt* neurons in 13 dpf star:bPAC+ larvae
underlie the observed reduction in food consumption at the same developmental stage. Another
hypothesis regarding the basis of the altered food consumption in the star:bPAC+ larvae relates to
the observed changes in hypothalamic radial glia. The population of hypothalamic radial glia altered
by developmental GC exposure in our model display similarities to mammalian tanycyte cells, based
on location, morphology, and marker expression[18]. Recent work demonstrated that hypothalamic
tanycytes can regulate food intake[49]. Hypothalamic tanycytes are known to project to neighbouring
orexigenic *npy*/*agrp* and anorexigenic *pomc* neurons of the arcuate nucleus, and tanycyte activation
was shown to depolarise these neurons *in vitro*, and lead to hyperphagia *in vivo*. It is interesting to
postulate that if similar functionality occurs in the hypothalamic radial glia of zebrafish, then the
observed increase in these cells at 5 dpf might contribute to the increased feeding behaviour
observed, meanwhile the subsequent reduction of these cells might contribute to the reduction of
feeding behaviour observed in 13 dpf star:bPAC+ larvae. Further work is required to characterise
the development and function of the zebrafish hypothalamic radial glia, and to determine the
mechanisms underlying the altered feeding observed in star:bPAC+ larvae.

We hypothesise that the precocious hypothalamic development observed here may represent an
example of GC-induced adaptive plasticity (Fig.6), as observed in other animals. For example in red
squirrels, females exposed to high-density cues (mimicking increased competition) increase their
levels of endogenous GC which drives them to produce offspring that grow faster than controls[50].
This so-called adaptive plasticity is acknowledged to have short-term fitness benefits, but the
investment is often associated with costs in later life[51]. The costs of accelerated development can
include reduced longevity, which is known as the growth-rate lifespan trade-off[52], as well as
reductions in breeding success[53]. Interestingly, there is evidence directly linking divergence from the
normal growth trajectory with longevity, since sticklebacks that are forced to grow faster have a
reduced lifespan, whilst those in which growth was slowed down lived longer than controls[52]. In our
model, chronic GC exposure ultimately appears to lead to allostatic overload, a maladaptive state
whereby an organism can no longer adapt to its conditions[54]. Whilst GC-induced acceleration of
growth-rate leading to adverse phenotypes later is widely observed at the organism level, the

underlying cellular and molecular mechanisms are poorly understood. Our work provides new
insight into these processes in the developing brain.

In the context of the early-late life trade-off, the subsequent reduction in breeding success and/or
longevity is thought to be related to oxidative stress[55,56]. Exposure to elevated GC is known to
increase oxidative stress[57], a phenomenon in which an over-accumulation of reactive oxygen
species (ROS) occurs in cells and tissues[58]. ROS is known to cause cellular damage, especially to
DNA, and can lead to accelerated telomere shortening[59]. In a previous study of amphibian larvae,
the relationship between stress, development, growth and aging depended on the type of stressor
exposure[60]. Predator exposure initiated faster development and enhanced growth of survivors, but
they showed signs of oxidative stress, had shorter telomeres and reduced long-term survival[60]. This
indicates that stress exposure may initiate faster development in early life, but at a cost to long-term
health and longevity.

Telomeres shorten with every round of cell division and when a critical size is reached this imposes
a functional limit on cell replication, leading to replicative senescence, which can contribute to age-
related diseases[59]. One possibility in our model is that the excess proliferation of hypothalamic
progenitors observed in 5 dpf star:bPAC+ larvae leads to subsequent senescence of those cells. In
contrast to quiescent, apoptotic, or terminally differentiated cells, senescent cells have incurred an
irreversible cell cycle arrest, yet remain viable, have alterations in metabolic activity and undergo
dramatic changes in gene expression[61]. In support of our hypothesis, a previous study of liver
progenitor cells reported that GC exposure induced cell proliferation leading to long-term replicative
senescence/stemness exhaustion[62], and GC is also reported to induce senescence of NSPCs *in*
*vitro*[63]. The presence of senescent cells in the proliferative zone of the GC-exposed hypothalamus
at 13 dpf might explain why we observed reduced expression of Pcna and *rx3*, and a reduced
proliferative capacity of the remaining glial cells in the proliferative zone of star:bPAC+ larvae.
Telomere shortening and senescence are associated with aging and age-related diseases; and
models of accelerated aging present with phenotypes such as reduced longevity and physical
decline[64], as observed in our model. Additionally, chronic exposure to GC is associated with signs of
accelerated aging at a cellular level[65]. Further analysis of oxidative stress and telomere homeostasis
would provide further insight into the mechanisms underlying the phenotypes observed here.

We conclude that the phenotypes observed in our optogenetic model are the result of exposure to
elevated GC produced in the interrenal, rather than to other interrenal steroid hormones, which are
not detectable in larval zebrafish[19,20,66]. We propose that GC-induced alteration of the hypothalamic
development trajectory observed in our model is mediated by GR-regulated gene transcription and
potentially epigenetic mechanisms. In support of this, we showed that treatment with GR antagonist

RU-486 reduced the number of pH3+ mitotic cells in the hypothalamus of star:bPAC+ larvae. We
also provide evidence for regulation of hypothalamic *rx3* by GR. Indeed, GR signalling is known to
regulate cell cycle progression in NSPCs by inducing a specific pattern of DNA methylation during
aging[67]. In a model of replicative senescence, GC exposure lead to altered methylation of GR target
gene *fkbp5*, which was exacerbated by age, and the subsequently increased *fkbp5* expression was
associated with inflammation and myocardial infarction in humans, suggesting that GC is linked with
age-related disease via a mechanism involving epigenetic regulation of *fkbp5*[68]. Meanwhile in
cancer cells, GC exposure can induce a cell dormancy state which is in part senescence and in part
quiescence, mediated by GR target gene *CDNK1C* (*cyclin-dependent kinase inhibitor 1C)*[69], further
supporting that GC exposure might regulate cell proliferation in our model via GR-mediated
regulation of cell cycle.

We speculate that the effects of elevated GC on the hypothalamic development trajectory reported
here might be an example of allostatic overload. Allostasis is the process of activating adaptive
mechanisms in response to external or internal changes, however prolonged exposure to stressors
can lead to accumulation of so-called allostatic load, and subsequently a maladaptive state of
allostatic overload[8], which is associated with age-related diseases and reduced longevity. In support
of this, it has been proposed that effects of stress or GC on NSPCs in young individuals may affect
their renewal potential in the long-term, predisposing to adult disease[36]. We hypothesise that whilst
precocious hypothalamic development induced by GC might be an example of adaptive plasticity in
the short-term, the cost of this may be paid in later life through subsequent replication senescence
of those cells, leading to failed hypothalamic maturation, subsequent impaired feeding behaviour
and ultimately physical decline (Fig.6). Further understanding of how stress and GC exposure can
alter development trajectories at the molecular and cellular level is of critical importance to reduce
the burden of mental and physical ill health across the life-course.

**Methods**

**Zebrafish Husbandry and Maintenance**
Zebrafish experiments were performed at the University of Exeter Aquatic Resource Centre and at
the Johannes Gutenberg University of Mainz in compliance with local and national animal welfare
laws, guidelines, and policies and approved by local government (Landesuntersuchugsamt
Rheinland-Pfalz, Germany –  23 177-07/G20-1-033 and Home Office PPL number: PEF291C4D).
*Tg(2kbStAR:bPAC-2A-tdTomato)[uex300]* were bred with wild-type TU zebrafish strain and were
screened at 4 dpf on a fluorescent stereomicroscope. Adult zebrafish were maintained under
standard zebrafish husbandry conditions, under a 12:12 light/dark cycle[70].

**RNA sequencing**

The detailed protocol for sample collection, RNA preparation, mRNA sequencing, and bioinformatic analysis is available in an online methods repository[71].

**Gene Ontology analysis**

The functional enrichment analysis was performed using g:Profiler (version *e107_eg54_p17_bf42210)* with an adjusted p-value of 0.05. For GO analysis of neurogenesis gene clusters (Fig1e), significant GO terms for each cluster were manually slimmed according to QuickGO ancestor charts. Heatmaps were generated in R using the heatmap.2 function.

**qPCR**

Fish were immobilised using ice-cold water and larvae or dissected juvenile/adult brains were stored in RNA later solution. Sample collection was carried out between 08:00 and 10:00. Larvae were subsequently dissected, such that each replicate consisted of 15 larval heads (eyes and jaw removed) or 3 juvenile/adult brains. RNA was extracted using TRIzol™ Reagent (Invitrogen™, 15596026), as previously described[72], and cDNA was synthesised using High-Capacity RNA-to-cDNA™ Kit (Applied Biosystems, 4387406). Approximately 100 ng cDNA was used in a 10 μl qPCR reaction with PowerUp™ SYBR™ Green Master Mix (A25778, Applied Biosystems). Primer sequences can be found in Supplementary table 1. The reactions were run in Hard-Shell 96-Well PCR Plates (HSP9601, BioRad) on a CFX96 Real Time PCR machine (Bio-Rad) using a standard protocol. Relative expression was calculated using the $2^{-\Delta\Delta Ct}$ method, with *18s* as a reference gene.

**Fluorescent in situ hybridisation (FISH) and immunohistochemistry (IHC)**

The FISH and IHC methods were based on a protocol by Jakob von Trotha (2016) for whole zebrafish embryos and larvae. Larvae or juvenile fish were immobilised in ice-cold water and then whole larvae or dissected juvenile fish brains were fixed in 4% PFA overnight at 4 °C, dehydrated in methanol and subsequently stored at -20C. After rehydration, larvae or brains were permeabilised using proteinase K (10 μg/ml) treatment at 37 °C and then re-fixed in PFA. DIG-conjugated mRNA probes were hybridised overnight at 65 °C. *In situ* hybridisation probes for *rx3*, *agrp*, *cart4, npy* were synthesised using primers listed in Supplementary Table 1. *pomc* probe was synthesised from a plasmid, as previously described[73]. After removal of probe and washing, samples were blocked for 2 hours in blocking solution (1% blocking reagent; (Roche, 11096176001) in malic acid buffer (0.15M maleic acid, 0.15M NaCl, PH 7.5)) and then incubated overnight at 4 °C in anti-DIG POD antibody (Roche, 11207733910) diluted 1:300 in blocking solution. After more washing, samples were incubated for 40 minutes in the dark with 1:200 FITC-tyramide (synthesised from product 46410, Thermo Scientific), 0.003% H2O2, 2% dextran sulphate in PBST. For IHC antigen retrieval was

performed according to published methods[74]. Samples were blocked in 10% Normal goat serum in

PBST for 2 hours before incubation in primary antibody overnight at 4 °C. For IHC of BrDU-treated

samples, samples were treated with 1N HCl for 30 minutes prior to blocking. Primary antibodies

used were: pH3 (Merck/Millipore, 06-570), pcna (Sigma, MABE288), BLBP (Sigma, ABN14), otpa[75],

BrDU (abcam, ab6326). After washing all day, samples were incubated overnight in 1:1000

secondary antibody with 1:200 hoechst (H3570, Invitrogen). Secondary antibodies used were Goat

anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488 (A-11008,

Invitrogen), Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 633 (A-

21070, Invitrogen), Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor

568 (A-11004, Invitrogen), Goat anti-Rat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa

Fluor 488 (A-11006, Invitrogen). After staining all samples were washed further then cleared

overnight in glycerol. Samples were imaged on a Zeiss LSM 880 confocal microscope using a 25x

or 10x objective.

**Image processing**

Image processing was performed in Fiji (ImageJ2, version 2.9.0). For whole brain counting of cells,

z-stacks were taken across the brain from dorsal to ventral and regions were defined using the

hoechst staining with the Z-Brain viewer[76]. Images were thresholded and PH3 cells in each region

were manually counted in a blinded manner. For counting of cells within the hypothalamus, cell

counts were performed manually from z-stacks, in a blinded manner. Similarly, hypothalamic

volume was calculated based on hoechst staining in z-stacks, using Z-Brain viewer. Corrected Total

Cell Fluorescence (CTCF) was calculated as Integrated Density of cells – (Area of cells X Mean

fluorescence of background reading). For CTCF calculations, labelled cells were identified first by

thresholding each image and creating ROIs of the labelled cells using the Analyze Particles

function. The total Integrated density and total area of all labelled cells was then used to calculate

CTCF for each sample. For co-expression analysis, ROIs of labelled cells were generated for each

marker as described above. Percentage overlap was calculated using the Analyze Particles function

with ROIs of marker 1 cells against a mask of marker 2 cells. Total area of cells was calculated from

the thresholded mask/ROIs of labelled cells. pH3 cells were manually scored as *rx3-* or *rx3+* using

the ROIs/masks as described above, in a blinded manner.

**Feeding behaviour**

Prior to behavioural analysis, larvae were transferred to 24 well cell culture plates (83.3922,

Sarstedt) with 1 larva per well in 1.5 ml aquarium water. Larvae were then left to acclimate to the

behavioural testing room (maintained at 28 °C) for 2 hours. 5 dpf larvae were fed 1 ml of diluted live

rotifers (approximately 20), meanwhile 13 dpf larvae were fed 5 live artemia. Larvae were recorded

for 10 minutes using Basler Video Recording Software. The camera used was a Basler (Germany)

acA1300-200um USB 3.0 camera with ON Semiconductor PYTHON 1300 CMOS sensor (203

frames per second at 1.3 MP resolution) with a Computar Zoom Lens 18-108/2.5 (Japan). Number

of successful and unsuccessful prey captures and latency to first hunting attempt were analysed

manually from video recordings in a blinded manner.

**BrDU experiment**

523    Larvae were transferred to a 12 well dish with 10 larvae per well. Larvae were incubated overnight

524    from 4 dpf in 3 ml 10 mM 5-Bromo-2′-deoxyuridine (B5002, Sigma) with 1% DMSO for 17 hours. On

525    the morning of 5 dpf, BrDU was removed by washing 3-times with aquarium water. Larvae were

526    subsequently fixed in PFA or raised until a later stage for fixation.

**Drug treatments**

529    Larvae were incubated from 6 hpf until 120 hpf with 50 µM dexamethasone (Sigma-Aldrich D2915)

530    or from 48 hpf until 120 hpf with 2 µM RU-486 (Sigma-Aldrich, M8046). DMSO solution was used as

531    a control. Solutions were changed daily.

**ChIP-qPCR**

We analysed the sequence for the *rx3* gene (ENSDARG00000052893,

chromosome:GRCz11:21:10755554:10759823:1). Primers for *rx3* GRE1 and GRE2 are described

in supplementary table 1 and were designed using Primer3. GR bounded chromatin was prepared

following the protocol from Idilli *et al.*, 2020[77], with minor modifications. For each sample, 100

dissected 5 dpf larval heads (without eyes) were incubated for 10 minutes at room temperature in

1% formaldehyde in PBS with protease inhibitors (A32955, Thermo Scientific) for cross-linking. We

used a glucocorticoid receptor polyclonal antibody (24050-1-AP, Proteintech) and Rabbit IgG

Isotype Control (10500C, Invitrogen) as primary antibodies.

**TUNEL assay**

544    Apoptotic cells were labelled using the In Situ Cell Death Detection Kit, Fluorescein (11684795910,

545    Roche). Larvae were first fixed, dehydrated, rehydrated and permeabilised as per the FISH

546    protocol. Larvae were incubated in TUNEL reaction mix at 37 °C for 2 hours and subsequently,

547    washed and cleared as per the FISH protocol.

**Statistics**

550    All analyses and graphs were generated in R (RStudio 2022.07.2). Prior to testing for statistically

551    significant differences between groups, data were tested for normality and variance. Where data did

552    not fit the assumptions for t-testing or ANOVA, non-parametric alternatives were used. Where
553    appropriate, Bonferroni correction for multiple testing was used. In all bar graphs mean +/- standard
554    error is presented for each group.
555
556

**Figures & Figure Legends**

**Figure 1. Neurogenesis is altered in an optogenetic zebrafish model of elevated GC.**

**a**. In star:bPAC+ larvae, blue light activates cAMP signalling in the interrenal gland, which leads to

release of cortisol. **b.** Wild-type and star:bPAC+ larvae were raised under a standard 12:12

light/dark cycle under white light. In these conditions, basal cortisol levels are elevated in

star:bPAC+ larvae from 2 dpf until approximately 2 months of age. **c.** Heatmap showing fold change

for 225 genes associated with GO:0022008 (neurogenesis) in star:bPAC+ whole-brain samples that

were downregulated at either 6 dpf or 13 dpf (or both), relative to wild type, from an RNA-

sequencing analysis. **d.** Scaled expression (cpm) of neurogenesis-associated genes in star:bPAC+

and wild-type whole-brain samples at 6 dpf and 13 dpf, according to the 6 clusters shown in c.

Boxes show the median and quartiles, whiskers are inter-quartile range and gray shading shows the

smoothed conditional means. **e.** Significant slimmed neurogenesis child GO terms at p<0.05,

identified using g:Profiler, are shown for each cluster. Size of the dot indicates the ratio of

differentially expressed genes from each GO term, whilst the colour shows the adjusted p-value. **a-b**
taken from Choi *et al.*,[19].

**Figure 2. Precocious hypothalamic development following elevated GC.**

**a.** Whole-brain IHC for proliferation marker pH3 as seen in a maximum intensity projection in 5 dpf
wild-type and star:bPAC+ larvae. Scale bar is 60 μm. **b.** Cell counting of pH3 across brain regions

revealed an increase in the hypothalamus (H) of 5 dpf star:bPAC+ larvae, compared with wild type

(Wilcox Exact test with Bonferroni correction (N=14, p= 0.0013195). Brain regions are labelled

according to the ZFIN anatomy database. **c**. Co-expression of *rx3*, pcna and pH3 in the

hypothalamus of 5 dpf wild-type and star:bPAC+ larvae, observed in a single plane image of FISH-

IHC. Scale bar 40 μm. Dashed line indicates the ventricles and arrowheads indicate excess *rx3+*

pH3+ cells in star:bPAC+ hypothalamus. **d.** IHC for blbp in the arcuate nucleus of 5 dpf wild-type

and star:bPAC+ larvae, observed in a single plane image. Scale bar 40 μm. **e.** IHC for otpa in the

hypothalamus of 5 dpf wild-type and star:bPAC+ larvae, as observed in a maximum intensity

projection. Scale bar 40 μm. **f-h**. The hypothalamus of 5 dpf star:bPAC+ larvae is larger (f, t= -

2.6193, df= 11.207, p-value = 0.02353) and has increased intensity of *rx3* (g, N=7, t= -2.3011, df=

8.5909, p-value = 0.04823) and pcna (h, N=7, t =-2.4409, df= 11.766, p-value = 0.03147). **i.** Excess

proliferating pH3+ cells in the hypothalamus of 5 dpf star:bPAC+ larvae are *rx3+* radial glia (N=7,

Two-way ANOVA with Tukey's test: total pH3+ cells (p= 0.0293550), *rx3+* pH3+ cells (p=

0.0373125)). **j-k.** Intensity of blbp (j, N=9, t= -2.2734, df= 14.903, p-value= 0.03823) and otpa (k,

N=12, t= -2.44, df= 16.227, p-value= 0.02652) was increased in the hypothalamus of 5 dpf

star:bPAC+ larvae. **l.** The zebrafish *rx3* gene contains two GREs, one in exon one and another in

intron one. The region shown is ENSDARG00000052893, GRCz11:21:10755554:10759823:1

(including flanking sequence). **m.** ChIP qPCR analysis of two GREs in the *rx3* gene shows

enrichment for GR compared with IgG control in 5 dpf wild-type brains. Enrichment of GR was

higher in wild types exposed to dexamethasone, and higher still in star:bPAC+ brains for both

GREs.

601
**Figure 3. Failed hypothalamic maturation following elevated GC.**

**a**. Expression of *rx3*, pcna and blbp in the hypothalamus of 13 dpf WT and star:bPAC larvae,
observed via FISH-IHC in single plane confocal images through the hypothalamus. Scale bar is 40
μm. **b-e.** Quantitative analysis of confocal images shown in (a), N=8. Volume of the hypothalamus is
not different (b, t = 0.31771, df = 10.899, p-value = 0.7567), meanwhile, *rx3* CTCF shows a trend
towards reduction (c, t = 1.8613, df = 8.3108, p-value = 0.09835), pcna CTCF is significantly
reduced (d, t = 4.0207, df = 7.7651, p-value = 0.00408), and blbp CTCF is not significantly different
(e, t = -1.2827, df = 13.092, p-value = 0.2219) in the hypothalamus of 13 dpf star:bPAC+ larvae
compared with wild types. **f-g.** Quantitative analysis of co-expression of pcna in hypothalamic radial
glia, from the images shown in (a). blbp cells (f, t = -2.5888, df = 7.5126, p-value = 0.0339) and *rx3*
cells (g, t = 4.7867, df = 13.444, p-value = 0.0003238) have a significant reduction in pcna co-
expression in the hypothalamus of 13 dpf star:bPAC+ larvae compared with wild types. **h.**
Expression of genes involved in feeding regulation in the hypothalamus of 13 dpf star:bPAC+ and
wild-type larvae as shown in maximum intensity projections of confocal stacks. Scale bar is 40 μm.
**i-k.** Quantitative analysis of cell counts in the stacks shown in (h). Number of *agrp* cells is
significantly lower (i, N=6-9, t = 2.2891, df = 11.255, p-value = 0.04234), number of *cart4* cells is
significantly reduced (j, N=20-21, t = 2.8544, df = 34.862, p-value = 0.007212), number of *hcrt* cells

shows a trend reduction (k, N=15-18, t = 1.9553, df = 26.571, p-value = 0.06115). All graphs show

the mean with standard error. #, $p<0.1$,  *, $p<0.05$, **, $p<0.01$, ***, $p<0.001$.

**Figure 4. Progenitor cells fail to differentiate in the hypothalamus following elevated GC.**

**a**. Schematic showing the experimental paradigm. Wild-type and star:bPAC+ larvae were treated with BrDU overnight from 4 dpf until 5 dpf for 17 hours. Following removal of BrDU compound, larvae were collected at 5 dpf, 8 dpf, 10 dpf and 13 dpf for lineage tracing analysis. **b,c.** Confocal images of BrDU lineage tracing analysis at 5 dpf (b) and 13 dpf (c). In each panel wild types are shown in the upper row and star:bPAC+ larvae in the bottom row. The left-most column (Merge)

shows the overlap between BrDU-labelled cells with endogenous pcna, whilst single channel
images for BrDU and pcna are adjacent. All images are a single plane from a confocal z-stack. In all
cases the scale bar indicates 40 μm. **d-f.** Quantitative analysis of BrDU cell fate. Whilst the
percentage of BrDU cells expressing pcna decreases over time in wild types, this remains high in
star:bPAC+ (d). Whilst BrDU+ cells increase between 5 dpf and 8 dpf in wild types, this does not
occur in star:bPAC+ (e). Similarly, pcna+ cells increase between 5 dpf and 10 dpf in wild types, but
not in star:bPAC+ larvae (f). Wild types are shown in blue, star:bPAC+ in pink. All graphs show the
mean with standard error. N=5-10 per group. *, $p<0.05$, **, $p<0.01$ for t-tests with Bonferroni
correction.
639
640

**Figure 5. Functional consequences of elevated GC: precocious feeding followed by physical decline**.

**a.** Analysis of feeding behaviour reveals that 5 dpf star:bPAC+ larvae eat more (N=12, t = -2.1694, df = 21.996, p-value = 0.04113). **b.** Analysis of feeding behaviour in 13 dpf wild-type and star:bPAC+ larvae reveals that food intake is significantly reduced (N=14, t = 2.5626, df = 24.299, p-value = 0.017). **c.** In 13 dpf wild-type larvae, food intake correlates positively with body size (linear regression, R-squared: 0.3619, F: 6.805 on 1 and 12 DF, p-value: 0.02286), however in star:bPAC+, there is no significant correlation (R-squared: 0.1877, F: 2.773 on 1 and 12 DF, p-value: 0.1218). **d.** star:bPAC+ larvae are significantly smaller than wild types at 28 dpf and 120 dpf, but not at 5 dpf or 13 dpf (Two-way ANOVA with Tukey's test, *** p<0.001 for 28 and 120 dpf). **e.** Percentage of embryos that were fertilised by adult male star:bPAC+ fish is significantly lower compared to wild types (N=16,19 pairs, t = 2.7181, df = 31.072, p-value = 0.01064). **f.** Percentage of star:bPAC+ fish that survived to 2 months is significantly reduced, compared with wild types (Wilcoxon test, N=9, W = 65.5, p-value = 0.004315).

Figure 6. Model proposing how elevated GC alters the trajectory of hypothalamic development and function.

In our model of elevated GC, we suggest that in the short-term, GC exposure drives adaptive plasticity, whereby we observed precocious hypothalamic development. In the long-term, chronic GC exposure leads to an accumulation of allostatic load, leading to allostatic overload. We observed this via failed hypothalamic maturation during the larval stage, and subsequent early physical decline of exposed fish.

**Acknowledgements**

This project is supported by an award from The Dennis and Mireille Gillings Foundation to SR and the German Federal Office for Education Research (grant number 01GQ1404) to SR. HE received support from the Society for Endocrinology and Wellcome Trust Institutional Strategic Support Fund 3 scheme (ISSF3) to Translational Research Exchange @ Exeter. MH received support from the CN Yang Scholars Programme at Nanyang Technological University. We would like to acknowledge the support of Ms. Kathrin Domdera at the University of Mainz and the Exeter Aquatic Resources Centre staff at the University of Exeter for expert zebrafish care as well as Dr Corin Liddle at Exeter Bioimaging Centre for microscopy. We are grateful to Dr Elina Jacobs, Dr Kate Ellacott and Professor Gil Levkowitz for feedback on the manuscript.

**Data Availability**

All sequenced reads for RNA-seq were deposited in European Nucleotide Archive (ENA, PRJEB53713).

**Author contributions**

HE designed the study, performed the experiments, analysed the data, provided funding and wrote the manuscript. MK designed, performed and analysed the RNA-seq experiments, and performed the GR-ChIP. AT, JK, and MH performed some of the experiments. SR provided funding and contributed to study design and manuscript writing.

**Materials & Correspondence**

For materials and correspondence please contact Soojin Ryu.

**Competing interests**

SR holds a patent, European patent number 2928288 and US patent number 10,080,355: "A novel inducible model of stress.". The remaining authors declare no known competing interests.

**References**

1.  Chrousos GP. Stress and disorders of the stress system. *Nature Reviews Endocrinology* **5**, 374-381 (2009).

2.  McKinlay CJD, Dalziel SR, Harding JE. Antenatal glucocorticoids: where are we after forty years? *J Dev Orig Health Dis* **6**, 127-142 (2015).

3.  Wolford E*, et al.* Associations of antenatal glucocorticoid exposure with mental health in children. *Psychological Medicine* **50**, 247-257 (2020).

704     4.     Ilg L, Klados M, Alexander N, Kirschbaum C, Li S-C. Long-term impacts of prenatal synthetic
705             glucocorticoids exposure on functional brain correlates of cognitive monitoring in
706             adolescence. *Scientific Reports* **8**, 7715 (2018).
707
708     5.     Davis EP, Sandman CA, Buss C, Wing DA, Head K. Fetal glucocorticoid exposure is
709             associated with preadolescent brain development. *Biol Psychiatry* **74**, 647-655 (2013).
710
711     6.     Bale TL*, et al.* Early Life Programming and Neurodevelopmental Disorders. *Biol Psychiatry*
712             **68**, 314-319 (2010).
713
714     7.     Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on
715             the brain, behaviour and cognition. *Nat Rev Neurosci* **10**, 434-445 (2009).
716
717     8.     Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-
718             related disease. *Physiol Behav* **106**, 29-39 (2012).
719
720     9.     Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming part 1: outcomes.
721             *Nature Reviews Endocrinology* **10**, 391-402 (2014).
722
723     10.     Suri D*, et al.* Early stress evokes age-dependent biphasic changes in hippocampal
724             neurogenesis, BDNF expression, and cognition. *Biol Psychiatry* **73**, 658-666 (2013).
725
726     11.     Mirescu C, Peters JD, Gould E. Early life experience alters response of adult neurogenesis
727             to stress. *Nat Neurosci* **7**, 841-846 (2004).
728
729     12.     Naninck EF*, et al.* Chronic early life stress alters developmental and adult neurogenesis and
730             impairs cognitive function in mice. *Hippocampus* **25**, 309-328 (2015).
731
732     13.     Lemaire V, Koehl M, Le Moal M, Abrous D. Prenatal stress produces learning deficits
733             associated with an inhibition of neurogenesis in the hippocampus. *Proceedings of the*
734             *National Academy of Sciences* **97**, 11032-11037 (2000).
735
736     14.     Sawamoto A*, et al.* 3,5,6,7,8,3',4'-Heptamethoxyflavone, a Citrus Flavonoid, Ameliorates
737             Corticosterone-Induced Depression-like Behavior and Restores Brain-Derived Neurotrophic
738             Factor Expression, Neurogenesis, and Neuroplasticity in the Hippocampus. *Molecules* **21**,
739             541 (2016).
740
741     15.     Lehmann ML, Brachman RA, Martinowich K, Schloesser RJ, Herkenham M. Glucocorticoids
742             Orchestrate Divergent Effects on Mood through Adult Neurogenesis. *The Journal of*
743             *Neuroscience* **33**, 2961-2972 (2013).
744
745     16.     Youssef M, Atsak P, Cardenas J, Kosmidis S, Leonardo ED, Dranovsky A. Early life stress
746             delays hippocampal development and diminishes the adult stem cell pool in mice. *Sci Rep* **9**,
747             4120 (2019).
748
749     17.     Bielefeld P*, et al.* Early life stress decreases cell proliferation and the number of putative
750             adult neural stem cells in the adult hypothalamus. *Stress* **24**, 189-195 (2021).
751
752     18.     Goodman T, Hajihosseini MK. Hypothalamic tanycytes-masters and servants of metabolic,
753             neuroendocrine, and neurogenic functions. *Front Neurosci* **9**, 387 (2015).
754
755     19.     Choi M-K*, et al.* Developmental Glucocorticoid exposure in zebrafish primes expression of
756             genes linked to human psychiatric disorders. *bioRxiv*, 2023.2002.2013.528363 (2023).
757

20.   Bacila I, Cunliffe VT, Krone NP. Interrenal development and function in zebrafish. *Molecular*
*and Cellular Endocrinology* **535**, 111372 (2021).

21.   Santulli G. *Adrenal Glands: From Pathophysiology to Clinical Evidence*. Nova Biomedical
(2015).

22.   Eachus H*, et al.* Genetic Disruption of 21-Hydroxylase in Zebrafish Causes Interrenal
Hyperplasia. *Endocrinology* **158**, 4165-4173 (2017).

23.   Oakes JA*, et al.* Ferredoxin 1b Deficiency Leads to Testis Disorganization, Impaired
Spermatogenesis, and Feminization in Zebrafish. *Endocrinology* **160**, 2401-2416 (2019).

24.   Li N, Oakes JA, Storbeck K-H, Cunliffe VT, Krone NP. The P450 side-chain cleavage
enzyme Cyp11a2 facilitates steroidogenesis in zebrafish. *Journal of Endocrinology* **244**, 309-
321 (2020).

25.   Oakes JA, Barnard L, Storbeck K-H, Cunliffe VT, Krone NP. 11β-Hydroxylase loss disrupts
steroidogenesis and reproductive function in zebrafish. *Journal of Endocrinology* **247**, 197-
212 (2020).

26.   Nagpal J, Eachus H, Lityagina O, Ryu S. Optogenetic induction of chronic glucocorticoid
exposure in early-life impairs stress-response in larval zebrafish. *bioRxiv*,
2022.2009.2009.507267 (2022).

27.   Gutierrez-Triana JA*, et al.* Manipulation of Interrenal Cell Function in Developing Zebrafish
Using Genetically Targeted Ablation and an Optogenetic Tool. *Endocrinology* **156**, 3394-
3401 (2015).

28.   Muthu V, Eachus H, Ellis P, Brown S, Placzek M. Rx3 and Shh direct anisotropic growth and
specification in the zebrafish tuberal/anterior hypothalamus. *Development (Cambridge,*
*England)* **143**, 2651-2663 (2016).

29.   Pak T, Yoo S, Miranda-Angulo AL, Wang H, Blackshaw S. Rax-CreERT2 knock-in mice: a
tool for selective and conditional gene deletion in progenitor cells and radial glia of the retina
and hypothalamus. *PLoS One* **9**, e90381 (2014).

30.   Kutsia M, Takeuchi Y, Ranawat N, Masai I. Single-cell transcriptome analysis reveals
heterogeneity and a dynamic regenerative response of quiescent radial glia in adult
zebrafish brain. *bioRxiv*, 2022.2007.2027.501663 (2022).

31.   Ryu S*, et al.* Orthopedia homeodomain protein is essential for diencephalic dopaminergic
neuron development. *Curr Biol* **17**, 873-880 (2007).

32.   Shainer I*, et al.* Agouti-Related Protein 2 Is a New Player in the Teleost Stress Response
System. *Curr Biol* **29**, 2009-2019 e2007 (2019).

33.   Yokobori E*, et al.* Stimulatory effect of intracerebroventricular administration of orexin A on
food intake in the zebrafish, Danio rerio. *Peptides* **32**, 1357-1362 (2011).

34.   Smith KL*, et al.* Overexpression of CART in the PVN Increases Food Intake and Weight
Gain in Rats. *Obesity* **16**, 2239-2244 (2008).

35.   Sundberg M, Savola S, Hienola A, Korhonen L, Lindholm D. Glucocorticoid hormones
decrease proliferation of embryonic neural stem cells through ubiquitin-mediated
degradation of cyclin D1. *J Neurosci* **26**, 5402-5410 (2006).

813
814    36.    Bornstein SR*, et al.* Stress-inducible-stem cells: a new view on endocrine, metabolic and
815           mental disease? *Molecular Psychiatry* **24**, 2-9 (2019).
816
817    37.    Bornstein SR, Berger I, Steenblock C. Are Nestin-positive cells responsive to stress? *Stress*
818           **23**, 662-666 (2020).
819
820    38.    Ninomiya E, Hattori T, Toyoda M, Umezawa A, Hamazaki T, Shintaku H. Glucocorticoids
821           promote neural progenitor cell proliferation derived from human induced pluripotent stem
822           cells. *Springerplus* **3**, 527 (2014).
823
824    39.    Heidt T*, et al.* Chronic variable stress activates hematopoietic stem cells. *Nat Med* **20**, 754-
825           758 (2014).
826
827    40.    Leung CS, Kosyk O, Welter EM, Dietrich N, Archer TK, Zannas AS. Chronic stress-driven
828           glucocorticoid receptor activation programs key cell phenotypes and functional epigenomic
829           patterns in human fibroblasts. *iScience* **25**, 104960 (2022).
830
831    41.    Nürnberg E, Horschitz S, Schloss P, Meyer-Lindenberg A. Basal glucocorticoid receptor
832           activation induces proliferation and inhibits neuronal differentiation of human induced
833           pluripotent stem cell-derived neuronal precursor cells. *The Journal of Steroid Biochemistry*
834           *and Molecular Biology* **182**, 119-126 (2018).
835
836    42.    Yu S, He J. Stochastic cell-cycle entry and cell-state-dependent fate outputs of injury-
837           reactivated tectal radial glia in zebrafish. *eLife* **8**, e48660 (2019).
838
839    43.    Noorlander CW*, et al.* Antenatal glucocorticoid treatment affects hippocampal development
840           in mice. *PLoS One* **9**, e85671 (2014).
841
842    44.    Loi M, Koricka S, Lucassen PJ, Joels M. Age- and sex-dependent effects of early life stress
843           on hippocampal neurogenesis. *Front Endocrinol (Lausanne)* **5**, 13 (2014).
844
845    45.    Lee DA*, et al.* Tanycytes of the hypothalamic median eminence form a diet-responsive
846           neurogenic niche. *Nat Neurosci* **15**, 700-702 (2012).
847
848    46.    Coll AP, Farooqi IS, O'Rahilly S. The Hormonal Control of Food Intake. *Cell* **129**, 251-262
849           (2007).
850
851    47.    Timper K, Brüning JC. Hypothalamic circuits regulating appetite and energy homeostasis:
852           pathways to obesity. *Dis Model Mech* **10**, 679-689 (2017).
853
854    48.    Zeltser LM. Feeding circuit development and early-life influences on future feeding
855           behaviour. *Nat Rev Neurosci* **19**, 302-316 (2018).
856
857    49.    Bolborea M, Pollatzek E, Benford H, Sotelo-Hitschfeld T, Dale N. Hypothalamic tanycytes
858           generate acute hyperphagia through activation of the arcuate neuronal network. *Proc Natl*
859           *Acad Sci U S A* **117**, 14473-14481 (2020).
860
861    50.    Dantzer B*, et al.* Density Triggers Maternal Hormones That Increase Adaptive Offspring
862           Growth in a Wild Mammal. *Science* **340**, 1215-1217 (2013).
863
864    51.    Metcalfe NB, Monaghan P. Compensation for a bad start: grow now, pay later? *Trends in*
865           *Ecology & Evolution* **16**, 254-260 (2001).
866

52.  Lee W-S, Monaghan P, Metcalfe NB. Experimental demonstration of the growth rate–
lifespan trade-off. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122370
(2013).

53.  Lemaître J-F*, et al.* Early-late life trade-offs and the evolution of ageing in the wild.
*Proceedings of the Royal Society B: Biological Sciences* **282**, 20150209 (2015).

54.  McEwen BS, Wingfield JC. The concept of allostasis in biology and biomedicine. *Horm
Behav* **43**, 2-15 (2003).

55.  Monaghan P. Organismal stress, telomeres and life histories. *J Exp Biol* **217**, 57-66 (2014).

56.  Jennings BJ, Ozanne SE, Hales CN. Nutrition, oxidative damage, telomere shortening, and
cellular senescence: individual or connected agents of aging? *Mol Genet Metab* **71**, 32-42
(2000).

57.  Costantini D, Marasco V, Møller AP. A meta-analysis of glucocorticoids as modulators of
oxidative stress in vertebrates. *Journal of Comparative Physiology B* **181**, 447-456 (2011).

58.  Pizzino G*, et al.* Oxidative Stress: Harms and Benefits for Human Health. *Oxid Med Cell
Longev* **2017**, 8416763 (2017).

59.  Barnes RP, Fouquerel E, Opresko PL. The impact of oxidative DNA damage and stress on
telomere homeostasis. *Mech Ageing Dev* **177**, 37-45 (2019).

60.  Burraco P, Díaz-Paniagua C, Gomez-Mestre I. Different effects of accelerated development
and enhanced growth on oxidative stress and telomere shortening in amphibian larvae.
*Scientific Reports* **7**, 7494 (2017).

61.  Kumari R, Jat P. Mechanisms of Cellular Senescence: Cell Cycle Arrest and Senescence
Associated Secretory Phenotype. *Frontiers in Cell and Developmental Biology* **9**,  (2021).

62.  Yang F*, et al.* Secretory galectin-3 induced by glucocorticoid stress triggers stemness
exhaustion of hepatic progenitor cells. *J Biol Chem* **295**, 16852-16862 (2020).

63.  Bose R, Moors M, Tofighi R, Cascante A, Hermanson O, Ceccatelli S. Glucocorticoids
induce long-lasting effects in neural stem cells resulting in senescence-related alterations.
*Cell Death Dis* **1**, e92 (2010).

64.  Anchelin M, Alcaraz-Pérez F, Martínez CM, Bernabé-García M, Mulero V, Cayuela ML.
Premature aging in telomerase-deficient zebrafish. *Dis Model Mech* **6**, 1101-1112 (2013).

65.  Bobba-Alves N*, et al.* Chronic Glucocorticoid Stress Reveals Increased Energy Expenditure
and Accelerated Aging as Cellular Features of Allostatic Load. *bioRxiv*,
2022.2002.2022.481548 (2022).

66.  Eachus H*, et al.* Genetic Disruption of 21-Hydroxylase in Zebrafish Causes Interrenal
Hyperplasia. *Endocrinology* **158**, 4165-4173 (2017).

67.  Schouten M*, et al.* Circadian glucocorticoid oscillations preserve a population of adult
hippocampal neural stem cells in the aging brain. *Molecular Psychiatry* **25**, 1382-1405
(2020).
919

920 68.    Zannas AS*, et al.* Epigenetic upregulation of FKBP5 by aging and stress contributes to NF-
921        κB–driven inflammation and cardiovascular risk. *Proceedings of the National Academy of*
922        *Sciences* **116**, 11370-11379 (2019).
923
924 69.    Prekovic S*, et al.* Glucocorticoid receptor triggers a reversible drug-tolerant dormancy state
925        with acquired therapeutic vulnerabilities in lung cancer. *Nature Communications* **12**, 4360
926        (2021).
927
928 70.    Westerfield M. *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio)*,
929        4th Edition edn. University of Oregon Press (2000).
930
931 71.    Choi M-K, Cook A, Eachus H, Tochwin A, Kuntz S, Ryu S. Comprehensive analysis methods
932        for developmental GC exposed zebrafish. Preprint at
933        http://dx.doi.org/10.17504/protocols.io.kxygx9ooog8j/v1 (2023).
934
935 72.    Eachus H, Bright C, Cunliffe VT, Placzek M, Wood JD, Watt PJ. Disrupted-in-Schizophrenia-
936        1 is essential for normal hypothalamic-pituitary-interrenal (HPI) axis function. *Hum Mol*
937        *Genet* **26**, 1992-2005 (2017).
938
939 73.    Herzog W*, et al.* Adenohypophysis formation in the zebrafish and its dependence on sonic
940        hedgehog. *Dev Biol* **254**, 36-49 (2003).
941
942 74.    Inoue D, Wittbrodt J. One for all--a highly efficient and versatile method for fluorescent
943        immunostaining in fish embryos. *PLoS One* **6**, e19713 (2011).
944
945 75.    Wolf A, Ryu S. Specification of posterior hypothalamic neurons requires coordinated
946        activities of Fezf2, Otp, Sim1a and Foxb1.*2*. *Development* **140**, 1762-1773 (2013).
947
948 76.    Randlett O*, et al.* Whole-brain activity mapping onto a zebrafish brain atlas. *Nat Methods* **12**,
949        1039-1046 (2015).
950
951 77.    Idilli AI*, et al.* Expression of tert Prevents ALT in Zebrafish Brain Tumors. *Frontiers in Cell*
952        *and Developmental Biology* **8**,  (2020).
953
954