Elevated Glucocorticoid alters the trajectory of hypothalamic development and function

Helen Eachus[1], Min-Kyeung. Choi[1], Anna Tochwin[1], Johanna Kaspareit[2], May Ho[3], Soojin Ryu*[1,2]

* Corresponding author

[1] Living Systems Institute, University of Exeter, Stocker Road, Exeter, EX4 4QD. United Kingdom.

[2] University Medical Center, Johannes Gutenberg University Mainz**,** Duesbergweg 6**,** 55128

Mainz, Germany.

[3] Nanyang Technological University, 50 Nanyang Avenue, Singapore 639798.

Abstract

Exposure to excess glucocorticoid (GC) during early development is implicated in disease risk in

later life. GC-induced acceleration of growth-rate leading to adverse phenotypes later is widely

observed at the organism level, but cellular and molecular understanding of this process is currently

lacking. Using an optogenetic zebrafish model, here we analysed the effects of GC exposure on

neurogenesis during development in the whole brain. We identify that the hypothalamus is a highly

GC-sensitive region where elevated GC causes precocious development followed by failed

maturation and early decline accompanied by impaired feeding, growth, and longevity. In GC-

exposed animals, the developmental trajectory of hypothalamic progenitor cells is strikingly altered,

potentially mediated by direct regulation of transcription factors such as *rx3* by GC. Our data provide

cellular and molecular level insight to GC-induced adaptive plasticity leading to allostatic overload in

a developing brain, a process crucial for health across the life-course.

Introduction

Glucocorticoids (GC) are the key effectors of the stress response and have pleiotropic effects on the

body, acting to restore homeostasis and thus allow an animal to respond adaptively to a threat[1].

Since the developing brain is plastic, GC exposure during early life has the potential to alter the

developmental trajectory of the brain. Indeed, alteration of the development trajectory may be an

adaptive mechanism employed by animals exposed to stress during early life. In red squirrels,

females exposed to high-density cues (mimicking increased competition) increase their levels of

endogenous GC and produce offspring which grow faster than controls[2]. This so-called adaptive

plasticity is acknowledged to have short-term fitness benefits, but the investment is often associated

with costs in later life[3]. Such costs of accelerated development include reduced longevity, which is

known as the growth-rate lifespan trade-off[4], as well as reductions in breeding success[5]. Meanwhile

in humans, exposure to GC in early life is associated with development of various adverse

phenotypes in later life, a concept known as early life programming of adult disease[6,7,8,9]. For

example, antenatal GC treatment has well documented effects on lung maturation, as well as other

organs, in preterm babies[10]. However, such treatment has also been associated with negative outcomes in later life, such as increased incidence of mental health issues during childhood[11], impacted development of fronto-parietal brain functions during adolescence[12], and cortical thinning in children with associated affective disorders[13].

One of the most well-documented effects of GC exposure during early life or early life stress (ELS) is reduced adult neurogenesis in the hippocampus, including reduced cell proliferation[14,15,16,17,18]. Indeed adrenalectomy is sufficient to prevent reduced proliferation in chronically stressed mice, suggesting that GC is the main driving force behind this phenotype[19]. In contrast, only a few studies have looked at the early life effects of stress or GC exposure on neurogenesis, but there are some indications that its effects might be developmentally dynamic. Specifically, it is possible that GC-induced reduction of cell proliferation is a delayed consequence of a GC-induced change to the development trajectory that began in early life. Indeed, rats exposed to early life stress (ELS) exhibited enhanced hippocampal cell proliferation and improved stress-associated behavioural performance as young adults, whilst these effects were reversed by middle age[14]. A potential mechanistic explanation for the ELS-induced reduction in adult hippocampal neurogenesis is that enhanced cell proliferation at an earlier time-point depletes the stem cell pool over time[20].

Despite the well-documented effects of GC exposure on hippocampal neurogenesis, the effects of GC exposure on other brain regions are unclear. Recently, a first study of the effects of ELS on neurogenesis in the hypothalamus revealed that cell proliferation and numbers of hypothalamic stem cells known as tanycytes were reduced in the adult mouse[21]. Tanycytes are unique radial glial-like cells that line the walls of the 3rd ventricle, have NSPC (neural stem/progenitor cell) properties that persist into adulthood, send projections into neighbouring hypothalamic nuclei and are able to sense glucose levels in the CSF[22]. Whilst most previous studies have focused on effects of stress and GC exposure on neurogenesis in the hippocampus, the aforementioned study supports that the hippocampus is not uniquely affected by stress/GC. However, whether and how GC affects hypothalamic neurogenesis during development is completely unknown.

Here we used an optogenetic zebrafish model in which endogenous cortisol level is elevated to test the hypothesis that GC exposure alters the developmental trajectory of the brain. We found that GC exposure affected cell proliferation in a region-specific manner and that these effects were dynamic across the life-course. In particular, we found that the hypothalamus is a highly GC-sensitive brain region. In early life, we observed precocious development of the hypothalamus and hypothalamus-associated feeding behaviour in GC-exposed animals. This was followed by a rapid decline, indicated by failed hypothalamic maturation and suppressed feeding and growth. Our data uncover a tissue-specific time-dependent plasticity of the hypothalamus in response to GC, and provide

75 cellular and molecular insights for GC-induced adaptive plasticity leading to allostatic overload in the
76 developing brain.
77
78 **Results**
79
Neurogenesis is altered in an optogenetic zebrafish model of elevated GC
Our lab previously generated an optogenetic zebrafish model, *Tg(star:bPAC-2A-tdTomato)^{uex300}*, in
which the endogenous cortisol level can be increased specifically in response to acute blue light
exposure using an optogenetic protein *beggiatoa* Photoactivated Adenylate Cylase (bPAC)
(Fig.1a)[23,24,25]. In order to determine the effects of elevated GC on the brain across development, we
raised transgenic fish under standard aquarium lighting conditions, where the blue light component
of the white light is sufficient to activate bPAC, which led to significantly elevated cortisol levels in
Tg(star:bPAC-2A-tdTomato)^{uex300} compared with wild types from hatching through until late
juvenile/early adult stage (Fig.1b)[25]. *Tg(star:bPAC-2A-tdTomato)^{uex300}* raised under standard light
condition will be referred to throughout as star:bPAC+ or bPAC+.
90
To investigate the effects of elevated GC on brain development we performed bulk RNA-
sequencing (RNA-seq) of whole brain samples from star:bPAC+ and wild-type fish at different
developmental stages[25]. Gene Ontology (GO) analysis of differentially expressed genes (DEGs) that
were down-regulated in star:bPAC+ brains compared with wild types were enriched for a number of
biological processes, including Neurogenesis at all developmental stages (GO:0022008, adjusted p-
value 2.47E-20 at 6 days post fertilization (dpf), 5.45E-08 at 13 dpf, 1.11E-17 at 120 dpf). Up-
regulated genes were not enriched for GO term neurogenesis. To investigate the effects of GC
exposure on neurogenesis during development, we focused downstream analysis on the early life
time-points: 5/6 dpf and 13 dpf. 225 genes associated with neurogenesis were down-regulated in
star:bPAC+ brains at either 6 dpf or 13 dpf (Fig.1c). To confirm our results and identify specific
features of neurogenesis that are altered by elevated GC, we performed qPCR on independent
samples of whole brains from 5 dpf star:bPAC+ and wild types for genes associated with
neurogenesis processes or distinct cell types. Of the 14 categories of genes analysed, only one
category was associated with consistent differential expression in star:bPAC+. Expression of *pcna*,
mki67 and *mcm5*, all associated with cell proliferation, showed increased expression in star:bPAC+
compared with wild types (Supplementary Fig.1). These data support that neurogenesis, and
especially cell proliferation, is altered by elevated GC.
108
To identify whether the effect of GC on neurogenesis is developmentally dynamic, we performed
unsupervised clustering analysis of neurogenesis DEGs at 6 dpf and 13 dpf and identified 6
different clusters. Cluster 2 and cluster 5 genes showed a relatively stronger down-regulation at 6

dpf compared to 13 dpf, meanwhile cluster 1 genes showed a relatively stronger downregulation at
13 dpf (Fig.1d). Cluster 3 genes were strongly downregulated at both time-points (Fig.1d).
Interestingly, cluster 4 and cluster 6 genes showed a timepoint-specific down-regulation. Cluster 4
genes were downregulated in star:bPAC+ at 6 dpf but were not differentially expressed at 13 dpf
(Fig.1d). Meanwhile cluster 6 genes were down-regulated in star:bPAC+ at 13 dpf but not at 13 dpf
(Fig.1d). These results suggest that some effects of elevated GC on neurogenesis are temporarily
dynamic during early life. GO analysis of genes within the clusters indicated that all clusters
contained genes associated with neuron differentiation (Fig.1e). Gliogenesis and glial cell
development were enriched in multiple clusters. There were also some terms which were restricted
to single or small numbers of clusters. For example, cluster 4 genes were associated with neuron
fate commitment, cluster 3 with neuron migration, and cluster 2 with neuronal stem cell population
maintenance (Fig.1e). These results suggest that specific aspects of neurogenesis exhibit
temporally dynamic differential expression during early life as a result of GC exposure.
125

Precocious hypothalamic development following elevated GC
In order to determine whether the effects of elevated GC on cell proliferation were brain-wide or
region specific we performed immunohistochemistry (IHC) for mitosis marker phospho-histoneH3
(pH3) on brains of 5 dpf star:bPAC+ and wild-type larvae (Fig.2a). Cell counting of proliferating
pH3+ cells across brain regions indicated a trend towards an increase in number of pH3+ cells in
the valvula cerebelli (ce-v) of the hindbrain, however, the only significant increase was restricted to
the hypothalamus (Fig.2b). We next sought to determine which cell types were associated with the
increased proliferative capacity in the hypothalamus of 5 dpf star:bPAC+. To this end, we performed
Fluorescent *in situ* hybridization (FISH) for *rx3* (*retinal homeobox gene 3),* combined with IHC for
pcna and pH3. *rx3* is a transcription factor known to be expressed in hypothalamic progenitor cells
in zebrafish where it plays a critical role in hypothalamic development[26], and its mammalian
orthologue *Rax* is known to be expressed in hypothalamic radial glia[27]. pcna and pH3 are markers
for proliferating cells during S-phase and M-phase respectively. pcna+ cells were situated around
the 3rd ventricle of the hypothalamus, especially around the lateral recess, in both star:bPAC+ and
wild-type larvae, with a small number of pH3+ cells scattered throughout this region (Fig.2c). We
distinguish the pcna+ region around the hypothalamic ventricles as the 'proliferative zone' and
observed co-localisation of *rx3* with pcna and pH3 within the proliferative zone (Fig.2c, white
arrowheads).
144

Quantitative analysis of confocal stacks revealed that the hypothalamus was significantly larger
(Fig.2f), and we found a significant increase in *rx3* and pcna signal intensity in the hypothalamus of
5 dpf star:bPAC+ compared with wild types (Fig.2g-h). Further, we observed that the excess
hypothalamic pH3+ cells were *rx3*+, meanwhile *rx3*-/pH3+ cells were not different (Fig.2i). These

results suggest that excess proliferating hypothalamic cells in star:bPAC+ are *rx3*-expressing radial

glia. Interestingly, we identified two Glucocorticoid Response Elements (GREs) within the zebrafish

rx3 gene (Fig.2l). GRE1 is in the 5' UTR, which is most likely a regulatory region. ChIP-qPCR

analysis revealed enrichment of GR antibody (compared with IgG) at both GREs in larval brain

samples, supporting that GR does indeed bind to regulatory regions of the *rx3* gene (Fig.2m).

Further, enrichment of GR was higher in dexamethasone-treated wild-type samples, compared with

wild-type controls, and higher still in star:bPAC+ samples. This experiment shows that GC exposure

directly regulates GR-mediated control of *rx3* gene expression in the brain.

Whilst *rx3* appeared to be co-expressed with pcna in proliferating radial glia, we also analysed blbp

(brain lipid binding protein, fabp7a) localisation, which is known to be expressed in predominantly

quiescent radial glia in the zebrafish brain[28]. Consistent with this, we observed strong expression of

blbp in cells lining the third ventricle in the pcna-negative rostral domains of the hypothalamus,

whilst some blbp signal was also observed in pcna-positive cells of the lateral recess (Fig.2d). In 5

dpf star:bPAC+ larvae, we observed increased blbp signal, especially in the arcuate region of the

hypothalamus (Fig.2d,j), suggesting that increased numbers of glial cells might include both

proliferative and quiescent subtypes. Finally we observed increased signal for otpa (orthopedia

homeobox a) in the hypothalamus of 5 dpf star:bPAC+ larvae (Fig.2e,k), which is predominantly

expressed in early post-mitotic neuronal precursors[29], suggesting that the excess GC-exposed

hypothalamic proliferative radial glia do exit the cell cycle, leading to increased numbers of neuronal

precursors. Together these data suggest that elevated GC drives precocious development of the

hypothalamus in zebrafish larvae.

Failed hypothalamic maturation following elevated GC

To test whether the effects of GC exposure on hypothalamic neurogenesis might be temporally

dynamic, we investigated hypothalamic cell proliferation at 13 dpf. In the 13 dpf hypothalamus, *rx3*

was expressed in a subset of pcna+ cells around the lateral recess, and blbp was expressed mainly

along the midline 3[rd] ventricle anterior to the proliferative zone, in a similar manner to at 5 dpf

(Fig.3a). In 13 dpf star:bPAC+ larvae, we observed that the size of the hypothalamus was no longer

larger, as was previously observed at 5 dpf (Fig.3b). Expression of hypothalamic *rx3* was very low at

this developmental stage, and showed a trend towards a reduction in star:bPAC+ compared with

wild type (Fig.3a,c). In stark contrast to the 5 dpf hypothalamus, we now observed a dramatic

reduction in expression of pcna in 13 dpf star:bPAC+ hypothalamus (Fig.3a,d). Meanwhile, blbp

signal intensity was not different between 13 dpf wild-type and star:bPAC+ larvae (Fig.3a,e).

Strikingly, when we performed co-expression analysis, we observed that whilst only a small

proportion of blbp+ cells were proliferative in 13 dpf wild-type hypothalamus (~10%), almost all

blbp+ cells were quiescent in star:bPAC+ (Fig.3f). Further, we observed that in wild types most
hypothalamic *rx3*+ cells (~70%) were proliferative and expressed pcna, however in star:bPAC+
most *rx3*+ cells (~85%) were quiescent and didn't express pcna (Fig.3g). These data support that
proliferative radial glia are lost in the hypothalamus of 13 dpf star:bPAC+ larvae. Next, we sought to
identify an explanation for the loss of proliferative radial glia in the hypothalamus. We first analysed
cell death in the hypothalamus of 13 dpf larvae using a TUNEL assay. We did not observe a
significant difference in the number of apoptotic cells in the hypothalamus of 13 dpf star:bPAC+
larvae compared to wild types (Supplementary Fig.2a), suggesting that loss of hypothalamic radial
glia is not due to increased cell death. We reasoned that the loss of proliferation in the star:bPAC+
hypothalamus might be due to senescence of the formerly highly proliferative radial glia. In support
of this hypothesis, we observed that one of the five significantly enriched KEGG pathways in our
whole brain RNA-seq data set in the 13 dpf star:bPAC+ was cellular senescence, a pathway which
was not enriched in the 6 dpf star:bPAC+ samples (Supplementary Fig.2b,c). In the RNA-seq
analysis, 102 of 150 genes associated with the KEGG pathway cellular senescence were
differentially expressed in 13 dpf star:bPAC+ brain samples, meanwhile at 6 dpf only 60 of those
genes were differentially expressed, and they generally showed a lesser degree of fold change
(Supplementary Fig.2d). This suggested that proliferative radial glia might have been lost in the
hypothalamus of 13 dpf star:bPAC+ larvae due to cellular senescence. Additionally, we confirmed
that the loss of hypothalamic proliferation is maintained into later life, since we observed that in
juvenile star:bPAC+ fish, the hypothalamus is smaller and contained fewer mitotic cells
(Supplementary Fig.3a-c). This suggests that the proliferative hypothalamic cells lost at 13 dpf do
not re-enter the cell cycle later in development, indicating that they have become senescent, rather
than quiescent.

Since we observed a dramatic reduction in cell proliferation in the hypothalamus of 13 dpf
star:bPAC+ larvae, we postulated that downstream neurogenesis and neuronal differentiation might
also be affected. Since one of the hypothalamus-regulated functions is feeding behaviour and food
intake, we sought to determine whether specific neurons that regulate feeding were altered in the
hypothalamus of star:bPAC+ larvae at 13 dpf. *agouti-related peptide* (*agrp*) is expressed in a cluster
of cells in the arcuate nucleus of the hypothalamus, and we observed significantly fewer numbers of
these cells in star:bPAC+ larvae (Fig.3h,i). A small number of cells expressing *cart4* (*cocaine- and
amphetamine-regulated transcript*) were observed near to the medial lateral hypothalamus (mLH)
and the total number of these cells were also reduced in star:bPAC+ larvae (Fig.3h,j). Orexin+ (*hcrt*,
hypocretin) cells were observed in the dorso-rostral hypothalamus and we observed a trend
reduction in cell count in star:bPAC+ larvae (Fig.3h,k). We observed *npy* (*neuropeptide y*) cells near
the mLH and these cells were not affected in star:bPAC+ larvae (Supplementary Fig.4a,b). Finally,
pomca (*proopiomelanocortin*) is also expressed in the arcuate hypothalamic nucleus, and these

cells were not significantly reduced in star:bPAC+ larvae (Supplementary Fig.4a,c). Of the
neuromodulators where we observed reduced numbers in star:bPAC+, *agrp*[30] and *hcrt*[31] are both
known to be appetite-stimulators, meanwhile the effects of *cart* on appetite are more complex[32].
These data indicate that neurogenesis of specific neuronal subtypes that stimulate feeding are
reduced in 13 dpf star:bPAC+ larvae.
228
Progenitor cells fail to differentiate in the hypothalamus of star:bPAC+ larvae
Since we observed that the effects of GC on hypothalamic proliferation changed dramatically
between 5 dpf and 13 dpf, we sought to track the proliferating radial glia of the hypothalamus
between these timepoints to determine their fate. To do this we exposed 4 dpf larvae to BrDU
(Bromodeoxyuridine) overnight for 17 hours until 5 dpf and then fixed larvae at 5, 8, 10 or 13 dpf for
double-IHC analysis of BrDU and pcna (Fig.4a). At 5 dpf we observed successful incorporation of
BrDU into proliferating cells, as observed by co-expression with pcna, in the hypothalamus of both
star:bPAC+ and wild-type larvae (Fig.4b). By 8, 10 and 13 dpf, BrDU+ cells in the wild-type
hypothalamus have begun exiting the cell cycle and are now mostly observed around the perimeter
of the pcna-expressing proliferative zone (Fig.4c, Supplementary Fig.5). Indeed, we observed that
the percentage of hypothalamic BrDU+ cells that co-express pcna drops from around 75% at 5 dpf
to 50-55% at 8-13 dpf (Fig.4d), suggesting that some of the BrDU+ cells that were proliferating at 5
dpf have lost their progenitor cell status and differentiated.
In stark contrast, hypothalamic BrDU+ cells in star:bPAC+ larvae are still predominantly observed
within the pcna+ proliferative zone after 5 dpf (Fig.4c, Supplementary Fig.5). In fact, the percentage
of hypothalamic BrDU+ cells that co-express pcna remains at around 100% from 5 dpf until 13 dpf
in star:bPAC+ larvae (Fig.4d). This indicates that in star:bPAC+ larvae, hypothalamic progenitor
cells remain proliferative between 5 dpf and 13 dpf and do not differentiate. Further, we observed an
increase in the number of hypothalamic BrDU+ cells between 5 dpf and 8 dpf in wild types,
supporting that these cells are dividing and therefore increase in number, however no such increase
was observed in star:bPAC+ larvae (Fig.4e). Similarly, the number of hypothalamic pcna+ cells
increased in wild types between 5 dpf and 10 dpf, indicating that the proliferative zone continues to
expand as cells divide during this period (Fig.4f). No increase in the number of hypothalamic pcna+
cells was observed in star:bPAC+ larvae during this developmental period (Fig.4f), further
supporting that neurogenesis appears to stall during this developmental window in star:bPAC+
larvae. Together, these data support that in star:bPAC+ larvae, hypothalamic progenitor cells fail to
differentiate, and neurogenesis is stalled during the 5-13 dpf time window.
256
Functional consequences of elevated GC: precocious feeding followed by physical decline
The increased size and proliferative capacity of the hypothalamus in 5 dpf star:bPAC+ larvae
suggested that development of the hypothalamus is accelerated in these animals. Therefore, we

hypothesised that behaviours associated with the hypothalamus that develop around this
developmental stage might also be altered and thus we analysed feeding in 5 dpf larvae. Feeding
usually emerges at around 120 hpf (5 dpf) in lab-raised zebrafish larvae, and larvae are capable of
eating small live food. We exposed 5 dpf larvae to live rotifers and saw that star:bPAC+ larvae
consumed significantly more rotifers during the trial (Fig.5a), suggesting that GC-induced
precocious hypothalamic development is accompanied by early emergence of an associated
behaviour, feeding.

In light of the reduction of cells expressing feeding-related neuromodulators in the hypothalamus of
13 dpf star:bPAC larvae, we sought to determine whether feeding was also affected at 13 dpf. In
stark contrast to 5 dpf, at 13 dpf star:bPAC+ larvae fed less than wild types (Fig.5b). Interestingly, in
13 dpf wild-type larvae we observed that food intake positively correlated with body size, in that
larger larvae ate more, whilst in star:bPAC+ larvae no significant trend was apparent. (Fig.5c).
Hence, star:bPAC+ larvae consumed less, regardless of their size. Whilst at 5 dpf and 13 dpf we did
not observe any difference in larval body size, at 28 dpf and 120 dpf star:bPAC+ fish were
significantly smaller than wild types (Fig.5d). This suggests that the reduction in food intake in
star:bPAC+ animals from 13 dpf ultimately leads to impaired growth. Finally, we also observed
reduced fertilisation rates and reduced long-term survival of star:bPAC+ fish (Fig.5e,f). Together,
these results point towards functional consequences of failed hypothalamic development after 5 dpf,
including impaired behaviour and an early physical decline.

Discussion
This work shows for the first time that elevated glucocorticoid alters the trajectory of hypothalamic
development and function. We identify that the hypothalamus is a highly GC-sensitive region where
elevated GC causes precocious development followed by failed maturation and early decline
accompanied by impaired feeding, growth, and longevity. We propose that the precocious
hypothalamic development observed here is an example of GC-induced adaptive plasticity (Fig.6).
However, chronic GC exposure ultimately leads to allostatic overload. Whilst GC-induced
acceleration of growth-rate leading to adverse phenotypes later is widely observed at the organism
level, the underlying cellular and molecular mechanisms are poorly understood. Our work provides
new insight into these processes in the developing brain.

In the fields of ecology and evolutionary biology, it is acknowledged that a trade-off may exist
between early life and later life fitness[3,4,5]. For example, animals which invest more in reproduction
during early life, such as via accelerated growth and maturation, might pay a cost for this in later life
in terms of reduced fitness and longevity. Indeed, an increased investment in early life success
likely represents an evolutionary advantage when exposed to adverse conditions during early life, in

that reproduction should be achieved before it's too late. In this context, the precocious
hypothalamic development observed in our model might represent a striking example of an adaptive
cellular response to the adverse conditions, in which GC-exposed individuals attempt to maximise
their breeding success by growing faster. On the other hand, it has also been observed that severe
exposure to stress or GC can impair growth temporarily and that this is followed by a period of so-
called catch-up growth, where growth rate is temporarily increased until normal size is reached[4,33].
In light of this, an alternative explanation is that in our model excess GC initially causes reduced
growth (prior to 5 dpf), and that our observation of increased size and proliferation of the
hypothalamus at 5 dpf represents a period of catch-up growth. However, this seems unlikely, since
catch-up growth would usually only occur after removal of the excess GC, which is not the case
here.
308
In the context of the early-late life trade-off, the subsequent reduction in breeding success and/or
longevity is thought to be related to oxidative stress[34,35]. Exposure to elevated GC is known to
increase oxidative stress[36], a phenomenon in which an over-accumulation of oxygen reactive
species (ROS) occurs in cells and tissues[37]. ROS is known to cause cellular damage, especially to
DNA, and can lead to accelerated telomere shortening[38]. In a previous study of amphibian larvae,
the relationship between stress, development, growth and aging depended on the type of stressor
exposure[39]. Predator exposure initiated faster development and enhanced growth of survivors, but
they showed signs of oxidative stress, had shorter telomeres and reduced long-term survival[39]. This
indicates that stress exposure may initiate faster development in early life, but at a cost to long-term
health and longevity.
319
Telomeres shorten with every round of cell division and when a critical size is reached this imposes
a functional limit on cell replication, leading to replicative senescence, which can contribute to age-
related diseases[38]. One possibility in our model is that the excess proliferation of hypothalamic
progenitors observed in 5 dpf star:bPAC+ larvae leads to subsequent senescence of those cells. In
contrast to quiescent, apoptotic, or terminally differentiated cells, senescent cells have incurred an
irreversible cell cycle arrest, yet remain viable, have alterations in metabolic activity and undergo
dramatic changes in gene expression[40]. In support of our hypothesis, a previous study of liver
progenitor cells reported that GC exposure induced cell proliferation leading to long-term replicative
senescence/stemness exhaustion[41], and GC is also reported to induce senescence of NSPCs
(neural stem/progenitor cells) *in vitro*[42]. The presence of senescent cells in the proliferative zone of
the GC-exposed hypothalamus at 13 dpf might explain why we observed reduced expression of
pcna and *rx3*, and a reduced proliferative capacity of the remaining glial cells in the proliferative
zone of star:bPAC+ larvae. Telomere shortening and senescence are associated with aging and
age-related diseases; and models of accelerated aging present with phenotypes such as reduced

longevity and physical decline[43], as observed in our model. Additionally, chronic exposure to GC is
associated with signs of accelerated aging at a cellular level[44]. Further analysis of oxidative stress
and telomere homeostasis would provide further insight into the mechanisms underlying the
phenotypes observed here.

In our model, we observed profound effects on a population of NSPCs within the proliferative zone
of the developing hypothalamus. Indeed this population of hypothalamic radial glia might be akin to
tanycyte cells observed in mammals, which also express *Rax*/*rx3* and *blbp*[22], and were recently
shown to be sensitive to ELS exposure in adult mice[21]. We postulate that these cells might be
especially sensitive to the effects of elevated GC, since they are adjacent to the 3^{rd} ventricle and
thus might detect elevated GC levels within the CSF. Indeed, neurogenesis of tanycyte cells is even
responsive to dietary cues[45]. Another potential driver of increased sensitivity of the hypothalamus to
GC would be increased levels of GR, however, it is not currently clear whether the hypothalamus
expresses higher levels of GR than other brain regions in zebrafish, or indeed in other vertebrates.
As such, an explanation for the specificity of the proliferation phenotype observed in our study
remains elusive. We cannot rule-out the possibility that the hypothalamus is not in fact uniquely
affected in our model, but rather that the effects of GC on different brain regions might emerge
across different time-scales.

Whilst many studies report GC-induced reduction in cell proliferation[46], our data supports the
hypothesis that, at least in some cases, the reduced cell proliferation following GC exposure might
be a consequence of a prior increase in proliferation. Indeed the concept of GC-inducible stem cells
is not novel[47,48], and there are many reports of a GC-induced increase in cell proliferation of various
cell types including NSPCs[49,50,51,52]. The initial GC-induced increase in cell proliferation observed
here might be in part mediated by mechanisms akin to those observed during brain injury, whereby
formerly quiescent radial glia enter the cell cycle in response to trauma[28,53]. In the published
literature, there are very few previous studies of effects of GC on cell proliferation across
development. Of those, Noorlander *et al.*, observed that antenatal GC treatment in mice led to an
initial reduction in cell proliferation in the embryonic hippocampus, followed by increased cell
proliferation during the postnatal period and a subsequent reduction in proliferation during
adulthood[54]. There are also examples in mouse models where ELS induced an initial increase in cell
proliferation in the hippocampus during early life which was again followed by a subsequent
reduction in cell proliferation[14] or hippocampal volume[16] in later life. Further, a meta-analysis of age-
dependent effects of ELS on cell proliferation shows a negative correlation between age and
change in cell proliferation, with the ELS-exposed animals analysed earliest in development
exhibiting increased cell proliferation and older ELS-exposed animals exhibiting a reduction in cell
proliferation[55]. It is possible that our interpretation of the results presented here also applies to these

studies, in that GC drives precocious proliferation leading to failed maturation and a rapid decline.

Interestingly, there is evidence directly linking divergence from the normal growth trajectory with

longevity, since sticklebacks that are forced to grow faster have a reduced lifespan, whilst those in

which growth was slowed down lived longer than controls[4].

We conclude that the phenotypes observed in our optogenetic model are the result of exposure to

elevated GC produced in the interrenal, rather than to other interrenal steroid hormones, since

cortisol is the primary output of the interrenal gland in zebrafish[25,56,57]. It is likely that the GC-induced

alteration of the hypothalamic development trajectory observed in our model is mediated by GR-

regulated gene transcription and potentially epigenetic mechanisms. Indeed, GR signalling is known

to regulate cell cycle progression in NSPCs by inducing a specific pattern of DNA methylation during

aging[58]. In a model of replicative senescence, GC exposure lead to altered methylation of GR target

gene *fkbp5*, which was exacerbated by age, and the subsequently increased *fkbp5* expression was

associated with inflammation and myocardial infarction in humans, suggesting that GC is linked with

age-related disease via a mechanism involving epigenetic regulation of *fkbp5*[59]. Meanwhile in

cancer cells, GC exposure can induce a cell dormancy state which is in part senescence and in part

quiescence, mediated by GR target gene *CDNK1C* (*cyclin-dependent kinase inhibitor 1C)[60]*, further

supporting that GC exposure might regulate cell proliferation in our model via GR-mediated

regulation of cell cycle.

We speculate that the effects of elevated GC on the hypothalamic development trajectory reported

here might be an example of allostatic overload. Allostasis is the process of activating adaptive

mechanisms in response to external or internal changes, however prolonged exposure to stressors

can lead to accumulation of so-called allostatic load, and subsequently a maladaptive state of

allostatic overload[8], which is associated with age-related diseases and reduced longevity. In support

of this, it has been proposed that effects of stress or GC on NSPCs in young individuals may affect

their renewal potential in the long-term, predisposing to adult disease[47]. Whilst precocious

hypothalamic development induced by GC might be an example of adaptive plasticity in the short-

term, the cost of this may be paid in later life through subsequent replication senescence of those

cells, leading to failed hypothalamic maturation, subsequent impaired feeding behaviour and

ultimately physical decline (Fig.6). Further understanding of how stress and GC exposure can alter

development trajectories at the molecular and cellular level is of critical importance to reduce the

burden of mental and physical ill health across the life-course.

Methods

Zebrafish Husbandry and Maintenance

Zebrafish experiments were performed at the University of Exeter Aquatic Resource Centre and at
the Johannes Gutenberg University of Mainz in compliance with local and national animal welfare
laws, guidelines, and policies and approved by local government (Landesuntersuchugsamt
Rheinland-Pfalz, Germany – 23 177-07/G20-1-033 and Home Office PPL number: PEF291C4D).
Tg(2kbStAR:bPAC-2A-tdTomato)[uex300] were bred with wild-type TU zebrafish strain and were
screened at 4 dpf on a fluorescent stereomicroscope. Adult zebrafish were maintained under
standard zebrafish husbandry conditions, under a 12:12 light/dark cycle[61].
415
416 **RNA sequencing**
417 The detailed protocol for sample collection, RNA preparation, mRNA sequencing, and bioinformatic
418 analysis is available in an online methods repository[62].
419
420 **Gene Ontology analysis**
421 The   functional   enrichment   analysis   was   performed   using   g:Profiler   (version
422 *e107_eg54_p17_bf42210)* with an adjusted p-value of 0.05. For GO analysis of neurogenesis gene
423 clusters (Fig1e), significant GO terms for each cluster were manually slimmed according to
424 QuickGO ancestor charts. Heatmaps were generated in R using the heatmap.2 function.
425
qPCR
Fish were immobilised using ice-cold water and larvae or dissected juvenile/adult brains were stored
in RNA later solution. Sample collection was carried out between 08:00 and 10:00. Larvae were
subsequently dissected, such that each replicate consisted of 15 larval heads (eyes and jaw
removed) or 3 juvenile/adult brains. RNA was extracted using TRIzol™ Reagent (Invitrogen™,
15596026), as previously described[63], and cDNA was synthesised using High-Capacity RNA-to-
cDNA™ Kit (Applied Biosystems, 4387406). Approximately 100 ng cDNA was used in a 10 µl qPCR
reaction with PowerUp™ SYBR™ Green Master Mix (A25778, Applied Biosystems). Primer
sequences can be found in Supplementary table 1. The reactions were run in Hard-Shell 96-Well
PCR Plates (HSP9601, BioRad) on a CFX96 Real Time PCR machine (Bio-Rad) using a standard
protocol. Relative expression was calculated using the $2^{-\Delta\Delta Ct}$ method, with *18s* as a reference gene.
437
Fluorescent in situ hybridisation (FISH) and immunohistochemistry (IHC)
The FISH and IHC methods were based on a protocol by Jakob von Trotha (2016) for whole
zebrafish embryos and larvae. Larvae or juvenile fish were immobilised in ice-cold water and then
whole larvae or dissected juvenile fish brains were fixed in 4% PFA overnight at 4 °C, dehydrated in
methanol and subsequently stored at -20C. After rehydration, larvae or brains were permeabilised
using proteinase K (10 µg/ml) treatment at 37 °C and then re-fixed in PFA. DIG-conjugated mRNA

probes were hybridised overnight at 65 °C. *In situ* hybridisation probes for *rx3*, *agrp*, *cart4, npy* were

synthesised using primers listed in Supplementary Table 1. *pomc* probe was synthesised from a

plasmid, as previously described[64]. After removal of probe and washing, samples were blocked for 2

hours in blocking solution (1% blocking reagent; (Roche, 11096176001) in malic acid buffer (0.15M

maleic acid, 0.15M NaCl, PH 7.5)) and then incubated overnight at 4 °C in anti-DIG POD antibody

(Roche, 11207733910) diluted 1:300 in blocking solution. After more washing, samples were

incubated for 40 minutes in the dark with 1:200 FITC-tyramide (synthesised from product 46410,

Thermo Scientific), 0.003% H_2O_2, 2% dextran sulphate in PBST. For IHC antigen retrieval was

performed according to published methods[65]. Samples were blocked in 10% Normal goat serum in

PBST for 2 hours before incubation in primary antibody overnight at 4 °C. For IHC of BrDU-treated

samples, samples were treated with 1N HCl for 30 minutes prior to blocking. Primary antibodies

used were: pH3 (Merck/Millipore, 06-570), pcna (Sigma, MABE288), BLBP (Sigma, ABN14), otpa[66],

BrDU (abcam, ab6326). After washing all day, samples were incubated overnight in 1:1000

secondary antibody with 1:200 hoechst (H3570, Invitrogen). Secondary antibodies used were Goat

anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488 (A-11008,

Invitrogen), Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 633 (A-

21070, Invitrogen), Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor

568 (A-11004, Invitrogen), Goat anti-Rat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa

Fluor 488 (A-11006, Invitrogen). After staining all samples were washed further then cleared

overnight in glycerol. Samples were imaged on a Zeiss LSM 880 confocal microscope using a 25x

or 10x objective.

Image processing

Image processing was performed in Fiji (ImageJ2, version 2.9.0). For whole brain counting of cells,

z-stacks were taken across the brain from dorsal to ventral and regions were defined using the

hoechst staining with the Z-Brain viewer[67]. Images were thresholded and PH3 cells in each region

were manually counted in a blinded manner. For counting of cells within the hypothalamus, cell

counts were performed manually from z-stacks, in a blinded manner. Similarly, hypothalamic

volume was calculated based on hoechst staining in z-stacks, using Z-Brain viewer. Corrected Total

Cell Fluorescence (CTCF) was calculated as Integrated Density of cells – (Area of cells X Mean

fluorescence of background reading). For CTCF calculations, labelled cells were identified first by

thresholding each image and creating ROIs of the labelled cells using the Analyze Particles

function. The total Integrated density and total area of all labelled cells was then used to calculate

CTCF for each sample. For co-expression analysis, ROIs of labelled cells were generated for each

marker as described above. Percentage overlap was calculated using the Analyze Particles function

with ROIs of marker 1 cells against a mask of marker 2 cells. Total area of cells was calculated from

the thresholded mask/ROIs of labelled cells. pH3 cells were manually scored as *rx3-* or *rx3*+ using

the ROIs/masks as described above, in a blinded manner.

Feeding behaviour

Prior to behavioural analysis, larvae were transferred to 24 well cell culture plates (83.3922,

Sarstedt) with 1 larva per well in 1.5 ml aquarium water. Larvae were then left to acclimate to the

behavioural testing room (maintained at 28 °C) for 2 hours. 5 dpf larvae were fed 1 ml of diluted live

rotifers (approximately 20), meanwhile 13 dpf larvae were fed 5 live artemia. Larvae were recorded

for 10 minutes using Basler Video Recording Software. The camera used was a Basler (Germany)

acA1300-200um USB 3.0 camera with ON Semiconductor PYTHON 1300 CMOS sensor (203

frames per second at 1.3 MP resolution) with a Computar Zoom Lens 18-108/2.5 (Japan). Number

of successful and unsuccessful prey captures and latency to first hunting attempt were analysed

manually from video recordings in a blinded manner.

BrDU experiment

495   Larvae were transferred to a 12 well dish with 10 larvae per well. Larvae were incubated overnight

496   from 4 dpf in 3 ml 10 mM 5-Bromo-2′-deoxyuridine (B5002, Sigma) with 1% DMSO for 17 hours. On

497   the morning of 5 dpf, BrDU was removed by washing 3-times with aquarium water. Larvae were

498   subsequently fixed in PFA or raised until a later stage for fixation.

Dexamethasone treatment

501   Larvae were incubated from 6 hpf to 5 dpf with 50 μM dexamethasone (Sigma-Aldrich D2915). The

502   solution was changed daily.

ChIP-qPCR

We analysed the sequence for the *rx3* gene (ENSDARG00000052893,

chromosome:GRCz11:21:10755554:10759823:1). Primers for *rx3* GRE1 and GRE2 are described

in supplementary table 1 and were designed using Primer3. GR bounded chromatin was prepared

following the protocol from Idilli *et al.*, 2020[68], with minor modifications. For each sample, 100

dissected 5 dpf larval heads (without eyes) were incubated for 10 minutes at room temperature in

1% formaldehyde in PBS with protease inhibitors (A32955, Thermo Scientific) for cross-linking. We

used a glucocorticoid receptor polyclonal antibody (24050-1-AP, Proteintech) and Rabbit IgG

Isotype Control (10500C, Invitrogen) as primary antibodies.

TUNEL assay

Apoptotic cells were labelled using the In Situ Cell Death Detection Kit, Fluorescein (11684795910,

Roche). Larvae were first fixed, dehydrated, rehydrated and permeabilised as per the FISH

protocol. Larvae were incubated in TUNEL reaction mix at 37 °C for 2 hours and subsequently,

washed and cleared as per the FISH protocol.

Statistics

All analyses and graphs were generated in R (RStudio 2022.07.2). Prior to testing for statistically

significant differences between groups, data were tested for normality and variance. Where data did

not fit the assumptions for t-testing or ANOVA, non-parametric alternatives were used. Where

appropriate, Bonferroni correction for multiple testing was used. In all bar graphs mean +/- standard

error is presented for each group.

 Figures & Figure Legends

Figure 1. Neurogenesis is altered in an optogenetic zebrafish model of elevated GC.
a. In star:bPAC+ larvae, blue light activates cAMP signalling in the interrenal gland, which leads to release of cortisol. **b.** Wild-type and star:bPAC+ larvae were raised under a standard 12:12 light/dark cycle under white light. In these conditions, basal cortisol levels are elevated in star:bPAC+ larvae from 2 dpf until approximately 2 months of age. **c.** Heatmap showing fold change for 225 genes associated with GO:0022008 (neurogenesis) in star:bPAC+ whole-brain samples that were downregulated at either 6 dpf or 13 dpf (or both), relative to wild type, from an RNA-sequencing analysis. **d.** Scaled expression (cpm) of neurogenesis-associated genes in star:bPAC+ and wild-type whole-brain samples at 6 dpf and 13 dpf, according to the 6 clusters shown in c. Boxes show the median and quartiles, whiskers are inter-quartile range and gray shading shows the smoothed conditional means. **e**. Significant slimmed neurogenesis child GO terms at p<0.05, identified using g:Profiler, are shown for each cluster. Size of the dot indicates the ratio of

543   differentially expressed genes from each GO term, whilst the colour shows the adjusted p-value. **a-b**

544   taken from Choi *et al.*,[25].

546   **Figure 2. Precocious hypothalamic development following elevated GC.**

547   **a.** Whole-brain IHC for proliferation marker pH3 as seen in a maximum intensity projection in 5 dpf

548   wild-type and star:bPAC+ larvae. Scale bar is 60 μm. **b.** Cell counting of pH3 across brain regions

revealed an increase in the hypothalamus (H) of 5 dpf star:bPAC+ larvae, compared with wild type (Wilcox Exact test with Bonferroni correction (N=14, p= 0.0013195). Brain regions are labelled according to the ZFIN anatomy database. **c**. Co-expression of *rx3*, pcna and pH3 in the hypothalamus of 5 dpf wild-type and star:bPAC+ larvae, observed in a single plane image of FISH-IHC. Scale bar 40 μm. Dashed line indicates the ventricles and arrowheads indicate excess *rx3*+ pH3+ cells in star:bPAC+ hypothalamus. **d.** IHC for blbp in the arcuate nucleus of 5 dpf wild-type and star:bPAC+ larvae, observed in a single plane image. Scale bar 40 μm. **e.** IHC for otpa in the hypothalamus of 5 dpf wild-type and star:bPAC+ larvae, as observed in a maximum intensity projection. Scale bar 40 μm. **f-h**. The hypothalamus of 5 dpf star:bPAC+ larvae is larger (f, t= -2.6193, df= 11.207, p-value = 0.02353) and has increased intensity of *rx3* (g, N=7, t= -2.3011, df= 8.5909, p-value = 0.04823) and pcna (h, N=7, t =-2.4409, df= 11.766, p-value = 0.03147). **i.** Excess proliferating pH3+ cells in the hypothalamus of 5 dpf star:bPAC+ larvae are *rx3*+ radial glia (N=7, Two-way ANOVA with Tukey's test: total pH3+ cells (p= 0.0293550), *rx3*+ pH3+ cells (p= 0.0373125)). **j-k.** Intensity of blbp (j, N=9, t= -2.2734, df= 14.903, p-value= 0.03823) and otpa (k, N=12, t= -2.44, df= 16.227, p-value= 0.02652) was increased in the hypothalamus of 5 dpf star:bPAC+ larvae. **l.** The zebrafish *rx3* gene contains two GREs, one in exon one and another in intron one. The region shown is ENSDARG00000052893, GRCz11:21:10755554:10759823:1 (including flanking sequence). **m.** ChIP qPCR analysis of two GREs in the *rx3* gene shows enrichment for GR compared with IgG control in 5 dpf wild-type brains. Enrichment of GR was higher in wild types exposed to dexamethasone, and higher still in star:bPAC+ brains for both GREs.

572
573 **Figure 3. Failed hypothalamic maturation following elevated GC.**

a. Expression of *rx3*, pcna and blbp in the hypothalamus of 13 dpf WT and star:bPAC larvae, observed via FISH-IHC in single plane confocal images through the hypothalamus. Scale bar is 40 μm. **b-e.** Quantitative analysis of confocal images shown in (a), N=8. Volume of the hypothalamus is not different (b, t = 0.31771, df = 10.899, p-value = 0.7567), meanwhile, *rx3* CTCF shows a trend towards reduction (c, t = 1.8613, df = 8.3108, p-value = 0.09835), pcna CTCF is significantly reduced (d, t = 4.0207, df = 7.7651, p-value = 0.00408), and blbp CTCF is not significantly different (e, t = -1.2827, df = 13.092, p-value = 0.2219) in the hypothalamus of 13 dpf star:bPAC+ larvae compared with wild types. **f-g.** Quantitative analysis of co-expression of pcna in hypothalamic radial glia, from the images shown in (a). blbp cells (f, t = -2.5888, df = 7.5126, p-value = 0.0339) and *rx3* cells (g, t = 4.7867, df = 13.444, p-value = 0.0003238) have a significant reduction in pcna co-expression in the hypothalamus of 13 dpf star:bPAC+ larvae compared with wild types. **h.** Expression of genes involved in feeding regulation in the hypothalamus of 13 dpf star:bPAC+ and wild-type larvae as shown in maximum intensity projections of confocal stacks. Scale bar is 40 μm. **i-k.** Quantitative analysis of cell counts in the stacks shown in (h). Number of *agrp* cells is significantly lower (i, N=6-9, t = 2.2891, df = 11.255, p-value = 0.04234), number of *cart4* cells is significantly reduced (j, N=20-21, t = 2.8544, df = 34.862, p-value = 0.007212), number of *hcrt* cells

590   shows a trend reduction (k, N=15-18, t = 1.9553, df = 26.571, p-value = 0.06115). All graphs show

591   the mean with standard error. #, $p<0.1$, *, $p<0.05$, **, $p<0.01$, ***, $p<0.001$.

Figure 4. Progenitor cells fail to differentiate in the hypothalamus following elevated GC.

a. Schematic showing the experimental paradigm. Wild-type and star:bPAC+ larvae were treated with BrDU overnight from 4 dpf until 5 dpf for 17 hours. Following removal of BrDU compound, larvae were collected at 5 dpf, 8 dpf, 10 dpf and 13 dpf for lineage tracing analysis. **b,c.** Confocal images of BrDU lineage tracing analysis at 5 dpf (b) and 13 dpf (c). In each panel wild types are shown in the upper row and star:bPAC+ larvae in the bottom row. The left-most column (Merge)

shows the overlap between BrDU-labelled cells with endogenous pcna, whilst single channel
images for BrDU and pcna are adjacent. All images are a single plane from a confocal z-stack. In all
cases the scale bar indicates 40 μm. **d-f.** Quantitative analysis of BrDU cell fate. Whilst the
percentage of BrDU cells expressing pcna decreases over time in wild types, this remains high in
star:bPAC+ (d). Whilst BrDU+ cells increase between 5 dpf and 8 dpf in wild types, this does not
occur in star:bPAC+ (e). Similarly, pcna+ cells increase between 5 dpf and 10 dpf in wild types, but
not in star:bPAC+ larvae (f). Wild types are shown in blue, star:bPAC+ in pink. All graphs show the
mean with standard error. N=5-10 per group. *, $p<0.05$, **, $p<0.01$ for t-tests with Bonferroni
correction.
610
611

Figure 5. Functional consequences of elevated GC: precocious feeding followed by physical decline.

a. Analysis of feeding behaviour reveals that 5 dpf star:bPAC+ larvae eat more (N=12, t = -2.1694, df = 21.996, p-value = 0.04113). **b.** Analysis of feeding behaviour in 13 dpf wild-type and star:bPAC+ larvae reveals that food intake is significantly reduced (N=14, t = 2.5626, df = 24.299, p-value = 0.017). **c.** In 13 dpf wild-type larvae, food intake correlates positively with body size (linear regression, R-squared: 0.3619, F: 6.805 on 1 and 12 DF, p-value: 0.02286), however in star:bPAC+, there is no significant correlation (R-squared: 0.1877, F: 2.773 on 1 and 12 DF, p-value: 0.1218). **d.** star:bPAC+ larvae are significantly smaller than wild types at 28 dpf and 120 dpf, but not at 5 dpf or 13 dpf (Two-way ANOVA with Tukey's test, *** p<0.001 for 28 and 120 dpf). **e.** Percentage of embryos that were fertilised by adult male star:bPAC+ fish is significantly lower compared to wild types (N=16,19 pairs, t = 2.7181, df = 31.072, p-value = 0.01064). **f.** Percentage of star:bPAC+ fish that survived to 2 months is significantly reduced, compared with wild types (Wilcoxon test, N=9, W = 65.5, p-value = 0.004315).

Figure 6. Model describing how elevated GC alters the trajectory of hypothalamic development and function.

In our model of elevated GC, we suggest that in the short-term, GC exposure drives adaptive plasticity, whereby we observed precocious hypothalamic development. In the long-term, chronic GC exposure leads to an accumulation of allostatic load, leading to allostatic overload. We observed this via failed hypothalamic maturation during the larval stage, and subsequent early physical decline of exposed fish.

Acknowledgements

This project is supported by an award from The Dennis and Mireille Gillings Foundation to SR and the German Federal Office for Education Research (grant number 01GQ1404) to SR. HE received support from the Society for Endocrinology and Wellcome Trust Institutional Strategic Support Fund 3 scheme (ISSF3) to Translational Research Exchange @ Exeter. MH received support from the CN Yang Scholars Programme at Nanyang Technological University. We would like to acknowledge the support of Ms. Kathrin Domdera at the University of Mainz and the Exeter Aquatic Resources Centre staff at the University of Exeter for expert zebrafish care as well as Dr Corin Liddle at Exeter Bioimaging Centre for microscopy, and Dr Elina Jacobs and Dr Kate Ellacott for feedback on the manuscript.

Data Availability

All sequenced reads for RNA-seq were deposited in European Nucleotide Archive (ENA, PRJEB53713).

Author contributions

HE designed the study, performed the experiments, analysed the data, provided funding and wrote the manuscript. MK designed, performed and analysed the RNA-seq experiments, and performed the GR-ChIP. AT, JK, and MH performed some of the experiments. SR provided funding and contributed to study design and manuscript writing.

Materials & Correspondence

For materials and correspondence please contact Soojin Ryu.

Competing interests

SR holds a patent, European patent number 2928288 and US patent number 10,080,355: "A novel inducible model of stress.". The remaining authors declare no known competing interests.

References

1.   Chrousos GP. Stress and disorders of the stress system. *Nature Reviews Endocrinology* **5**, 374-381 (2009).

2.   Dantzer B*, et al.* Density Triggers Maternal Hormones That Increase Adaptive Offspring Growth in a Wild Mammal. *Science* **340**, 1215-1217 (2013).

3.   Metcalfe NB, Monaghan P. Compensation for a bad start: grow now, pay later? *Trends in Ecology & Evolution* **16**, 254-260 (2001).

675 4.   Lee W-S, Monaghan P, Metcalfe NB. Experimental demonstration of the growth rate–
676   lifespan trade-off. *Proceedings of the Royal Society B: Biological Sciences* **280**, 20122370
677   (2013).
678
679 5.   Lemaître J-F*, et al.* Early-late life trade-offs and the evolution of ageing in the wild.
680   *Proceedings of the Royal Society B: Biological Sciences* **282**, 20150209 (2015).
681
682 6.   Bale TL*, et al.* Early Life Programming and Neurodevelopmental Disorders. *Biol Psychiatry*
683   **68**, 314-319 (2010).
684
685 7.   Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on
686   the brain, behaviour and cognition. *Nat Rev Neurosci* **10**, 434-445 (2009).
687
688 8.   Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-
689   related disease. *Physiol Behav* **106**, 29-39 (2012).
690
691 9.   Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming part 1: outcomes.
692   *Nature Reviews Endocrinology* **10**, 391-402 (2014).
693
694 10.   McKinlay CJD, Dalziel SR, Harding JE. Antenatal glucocorticoids: where are we after forty
695   years? *J Dev Orig Health Dis* **6**, 127-142 (2015).
696
697 11.   Wolford E*, et al.* Associations of antenatal glucocorticoid exposure with mental health in
698   children. *Psychological Medicine* **50**, 247-257 (2020).
699
700 12.   Ilg L, Klados M, Alexander N, Kirschbaum C, Li S-C. Long-term impacts of prenatal synthetic
701   glucocorticoids exposure on functional brain correlates of cognitive monitoring in
702   adolescence. *Scientific Reports* **8**, 7715 (2018).
703
704 13.   Davis EP, Sandman CA, Buss C, Wing DA, Head K. Fetal glucocorticoid exposure is
705   associated with preadolescent brain development. *Biol Psychiatry* **74**, 647-655 (2013).
706
707 14.   Suri D*, et al.* Early stress evokes age-dependent biphasic changes in hippocampal
708   neurogenesis, BDNF expression, and cognition. *Biol Psychiatry* **73**, 658-666 (2013).
709
710 15.   Mirescu C, Peters JD, Gould E. Early life experience alters response of adult neurogenesis
711   to stress. *Nat Neurosci* **7**, 841-846 (2004).
712
713 16.   Naninck EF*, et al.* Chronic early life stress alters developmental and adult neurogenesis and
714   impairs cognitive function in mice. *Hippocampus* **25**, 309-328 (2015).
715
716 17.   Lemaire V, Koehl M, Le Moal M, Abrous D. Prenatal stress produces learning deficits
717   associated with an inhibition of neurogenesis in the hippocampus. *Proceedings of the
718   National Academy of Sciences* **97**, 11032-11037 (2000).
719
720 18.   Sawamoto A*, et al.* 3,5,6,7,8,3′,4′-Heptamethoxyflavone, a Citrus Flavonoid, Ameliorates
721   Corticosterone-Induced Depression-like Behavior and Restores Brain-Derived Neurotrophic
722   Factor Expression, Neurogenesis, and Neuroplasticity in the Hippocampus. *Molecules* **21**,
723   541 (2016).
724
725 19.   Lehmann ML, Brachman RA, Martinowich K, Schloesser RJ, Herkenham M. Glucocorticoids
726   Orchestrate Divergent Effects on Mood through Adult Neurogenesis. *The Journal of
727   Neuroscience* **33**, 2961-2972 (2013).
728

729 20. Youssef M, Atsak P, Cardenas J, Kosmidis S, Leonardo ED, Dranovsky A. Early life stress
730   delays hippocampal development and diminishes the adult stem cell pool in mice. *Sci Rep* **9**,
731   4120 (2019).
732
733 21. Bielefeld P*, et al.* Early life stress decreases cell proliferation and the number of putative
734   adult neural stem cells in the adult hypothalamus. *Stress* **24**, 189-195 (2021).
735
736 22. Goodman T, Hajihosseini MK. Hypothalamic tanycytes-masters and servants of metabolic,
737   neuroendocrine, and neurogenic functions. *Front Neurosci* **9**, 387 (2015).
738
739 23. Nagpal J, Eachus H, Lityagina O, Ryu S. Optogenetic induction of chronic glucocorticoid
740   exposure in early-life impairs stress-response in larval zebrafish. *bioRxiv*,
741   2022.2009.2009.507267 (2022).
742
743 24. Gutierrez-Triana JA*, et al.* Manipulation of Interrenal Cell Function in Developing Zebrafish
744   Using Genetically Targeted Ablation and an Optogenetic Tool. *Endocrinology* **156**, 3394-
745   3401 (2015).
746
747 25. Choi M-K*, et al.* Developmental Glucocorticoid exposure in zebrafish primes expression of
748   genes linked to human psychiatric disorders. *bioRxiv*, 2023.2002.2013.528363 (2023).
749
750 26. Muthu V, Eachus H, Ellis P, Brown S, Placzek M. Rx3 and Shh direct anisotropic growth and
751   specification in the zebrafish tuberal/anterior hypothalamus. *Development (Cambridge,*
752   *England)* **143**, 2651-2663 (2016).
753
754 27. Pak T, Yoo S, Miranda-Angulo AL, Wang H, Blackshaw S. Rax-CreERT2 knock-in mice: a
755   tool for selective and conditional gene deletion in progenitor cells and radial glia of the retina
756   and hypothalamus. *PLoS One* **9**, e90381 (2014).
757
758 28. Kutsia M, Takeuchi Y, Ranawat N, Masai I. Single-cell transcriptome analysis reveals
759   heterogeneity and a dynamic regenerative response of quiescent radial glia in adult
760   zebrafish brain. *bioRxiv*, 2022.2007.2027.501663 (2022).
761
762 29. Ryu S*, et al.* Orthopedia homeodomain protein is essential for diencephalic dopaminergic
763   neuron development. *Curr Biol* **17**, 873-880 (2007).
764
765 30. Shainer I*, et al.* Agouti-Related Protein 2 Is a New Player in the Teleost Stress Response
766   System. *Curr Biol* **29**, 2009-2019 e2007 (2019).
767
768 31. Yokobori E*, et al.* Stimulatory effect of intracerebroventricular administration of orexin A on
769   food intake in the zebrafish, Danio rerio. *Peptides* **32**, 1357-1362 (2011).
770
771 32. Smith KL*, et al.* Overexpression of CART in the PVN Increases Food Intake and Weight
772   Gain in Rats. *Obesity* **16**, 2239-2244 (2008).
773
774 33. Baron J*, et al.* Catch-up growth after glucocorticoid excess: a mechanism intrinsic to the
775   growth plate. *Endocrinology* **135**, 1367-1371 (1994).
776
777 34. Monaghan P. Organismal stress, telomeres and life histories. *J Exp Biol* **217**, 57-66 (2014).
778
779 35. Jennings BJ, Ozanne SE, Hales CN. Nutrition, oxidative damage, telomere shortening, and
780   cellular senescence: individual or connected agents of aging? *Mol Genet Metab* **71**, 32-42
781   (2000).
782

783   36.   Costantini D, Marasco V, Møller AP. A meta-analysis of glucocorticoids as modulators of
784   oxidative stress in vertebrates. *Journal of Comparative Physiology B* **181**, 447-456 (2011).
785

786   37.   Pizzino G*, et al.* Oxidative Stress: Harms and Benefits for Human Health. *Oxid Med Cell*
787   *Longev* **2017**, 8416763 (2017).
788

789   38.   Barnes RP, Fouquerel E, Opresko PL. The impact of oxidative DNA damage and stress on
790   telomere homeostasis. *Mech Ageing Dev* **177**, 37-45 (2019).
791

792   39.   Burraco P, Díaz-Paniagua C, Gomez-Mestre I. Different effects of accelerated development
793   and enhanced growth on oxidative stress and telomere shortening in amphibian larvae.
794   *Scientific Reports* **7**, 7494 (2017).
795

796   40.   Kumari R, Jat P. Mechanisms of Cellular Senescence: Cell Cycle Arrest and Senescence
797   Associated Secretory Phenotype. *Frontiers in Cell and Developmental Biology* **9**, (2021).
798

799   41.   Yang F*, et al.* Secretory galectin-3 induced by glucocorticoid stress triggers stemness
800   exhaustion of hepatic progenitor cells. *J Biol Chem* **295**, 16852-16862 (2020).
801

802   42.   Bose R, Moors M, Tofighi R, Cascante A, Hermanson O, Ceccatelli S. Glucocorticoids
803   induce long-lasting effects in neural stem cells resulting in senescence-related alterations.
804   *Cell Death Dis* **1**, e92 (2010).
805

806   43.   Anchelin M, Alcaraz-Pérez F, Martínez CM, Bernabé-García M, Mulero V, Cayuela ML.
807   Premature aging in telomerase-deficient zebrafish. *Dis Model Mech* **6**, 1101-1112 (2013).
808

809   44.   Bobba-Alves N*, et al.* Chronic Glucocorticoid Stress Reveals Increased Energy Expenditure
810   and Accelerated Aging as Cellular Features of Allostatic Load. *bioRxiv*,
811   2022.2002.2022.481548 (2022).
812

813   45.   Lee DA*, et al.* Tanycytes of the hypothalamic median eminence form a diet-responsive
814   neurogenic niche. *Nat Neurosci* **15**, 700-702 (2012).
815

816   46.   Sundberg M, Savola S, Hienola A, Korhonen L, Lindholm D. Glucocorticoid hormones
817   decrease proliferation of embryonic neural stem cells through ubiquitin-mediated
818   degradation of cyclin D1. *J Neurosci* **26**, 5402-5410 (2006).
819

820   47.   Bornstein SR*, et al.* Stress-inducible-stem cells: a new view on endocrine, metabolic and
821   mental disease? *Molecular Psychiatry* **24**, 2-9 (2019).
822

823   48.   Bornstein SR, Berger I, Steenblock C. Are Nestin-positive cells responsive to stress? *Stress*
824   **23**, 662-666 (2020).
825

826   49.   Ninomiya E, Hattori T, Toyoda M, Umezawa A, Hamazaki T, Shintaku H. Glucocorticoids
827   promote neural progenitor cell proliferation derived from human induced pluripotent stem
828   cells. *Springerplus* **3**, 527 (2014).
829

830   50.   Heidt T*, et al.* Chronic variable stress activates hematopoietic stem cells. *Nat Med* **20**, 754-
831   758 (2014).
832

833   51.   Leung CS, Kosyk O, Welter EM, Dietrich N, Archer TK, Zannas AS. Chronic stress-driven
834   glucocorticoid receptor activation programs key cell phenotypes and functional epigenomic
835   patterns in human fibroblasts. *iScience* **25**, 104960 (2022).
836

52. Nürnberg E, Horschitz S, Schloss P, Meyer-Lindenberg A. Basal glucocorticoid receptor
activation induces proliferation and inhibits neuronal differentiation of human induced
pluripotent stem cell-derived neuronal precursor cells. *The Journal of Steroid Biochemistry*
and Molecular Biology **182**, 119-126 (2018).

53. Yu S, He J. Stochastic cell-cycle entry and cell-state-dependent fate outputs of injury-
reactivated tectal radial glia in zebrafish. *eLife* **8**, e48660 (2019).

54. Noorlander CW*, et al.* Antenatal glucocorticoid treatment affects hippocampal development
in mice. *PLoS One* **9**, e85671 (2014).

55. Loi M, Koricka S, Lucassen PJ, Joels M. Age- and sex-dependent effects of early life stress
on hippocampal neurogenesis. *Front Endocrinol (Lausanne)* **5**, 13 (2014).

56. Bacila I, Cunliffe VT, Krone NP. Interrenal development and function in zebrafish. *Molecular*
and Cellular Endocrinology **535**, 111372 (2021).

57. Eachus H*, et al.* Genetic Disruption of 21-Hydroxylase in Zebrafish Causes Interrenal
Hyperplasia. *Endocrinology* **158**, 4165-4173 (2017).

58. Schouten M*, et al.* Circadian glucocorticoid oscillations preserve a population of adult
hippocampal neural stem cells in the aging brain. *Molecular Psychiatry* **25**, 1382-1405
(2020).

59. Zannas AS*, et al.* Epigenetic upregulation of FKBP5 by aging and stress contributes to NF-
κB–driven inflammation and cardiovascular risk. *Proceedings of the National Academy of*
Sciences **116**, 11370-11379 (2019).

60. Prekovic S*, et al.* Glucocorticoid receptor triggers a reversible drug-tolerant dormancy state
with acquired therapeutic vulnerabilities in lung cancer. *Nature Communications* **12**, 4360
(2021).

61. Westerfield M. *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio)*,
4th Edition edn. University of Oregon Press (2000).

62. Choi M-K, Cook A, Eachus H, Tochwin A, Kuntz S, Ryu S. Comprehensive analysis methods
for developmental GC exposed zebrafish. Preprint at
http://dx.doi.org/10.17504/protocols.io.kxygx9ooog8j/v1 (2023).

63. Eachus H, Bright C, Cunliffe VT, Placzek M, Wood JD, Watt PJ. Disrupted-in-Schizophrenia-
1 is essential for normal hypothalamic-pituitary-interrenal (HPI) axis function. *Hum Mol*
Genet **26**, 1992-2005 (2017).

64. Herzog W*, et al.* Adenohypophysis formation in the zebrafish and its dependence on sonic
hedgehog. *Dev Biol* **254**, 36-49 (2003).

65. Inoue D, Wittbrodt J. One for all--a highly efficient and versatile method for fluorescent
immunostaining in fish embryos. *PLoS One* **6**, e19713 (2011).

66. Wolf A, Ryu S. Specification of posterior hypothalamic neurons requires coordinated
activities of Fezf2, Otp, Sim1a and Foxb1.2. *Development* **140**, 1762-1773 (2013).

67. Randlett O*, et al.* Whole-brain activity mapping onto a zebrafish brain atlas. *Nat Methods* **12**,
1039-1046 (2015).
891

892   68.   Idilli AI, *et al.* Expression of tert Prevents ALT in Zebrafish Brain Tumors. *Frontiers in Cell*
893   *and Developmental Biology* **8**, (2020).
894
895