Revisit the Inhibitory Effects of Glucocorticoids on Immunocytes

Shuting Wu[1, 2], Shushu Zhao[1 †], Yiwei Zhong[1, 2&3], and Bin Wang[1, 2 &3 *]

[1]Key Laboratory of Medical Molecular Virology (MOE/NHC/CAMS), School of Basic Medical Sciences, Fudan University, Shanghai, China.

[2]Fudan-Advaccine Join-Lab for Vaccine Research, Fudan University, Shanghai, China.

[3]Shanghai Institute of Infectious Disease and Biosecurity, Fudan University, Shanghai, China.

†The Wistar Institute, Philadelphia 3601 Spruce Street, PA 19104, USA

*Corresponding author: Bin Wang, email: bwang3@fudan.edu.cn. Address: 131 Dong An Road, 409 Fuxing Building, Shanghai 200032, People's Republic of China.

Running head: INHIBITORY EFFECTS OF DEX ON IMMUNOCYTES

Abstract

Glucocorticoids (GCs) are efficacious agents for reducing inflammation and suppressing immune

responses, exerting various effects on immune cells through the intracellular glucocorticoid receptor

(GR), and impacting both innate and adaptive immunity. In the context of COVID-19, glucocorticoids

are often used to treat severe cases of patients by reducing inflammation, suppressing immune responses,

and ameliorating the severity of COVID-19. However, the precise inhibitory effects on immune cells

have yet to be comprehensively delineated. In this study, we extensively examined the inhibitory effects

of treating Balb/c mice with dexamethasone (DEX) on lymphoid and myeloid cells. We observed that

high doses of DEX treatment resulted in a reduction in the number of immunocytes and an attenuation

of their activity. Particularly noteworthy, macrophages, DC cells, and monocytes were diminished by

approximately 90% following high doses of DEX, while B cells experienced a reduction of about 70%

and CD3 T cells were less affected. Furthermore, our findings demonstrated that DEX induces the

inhibition of immune cells by engaging in high-affinity binding to GR. Consequently, we conclude that

DEX treatments affect a broad range of immune cells, encompassing both lymphoid and myeloid cells,

through depletion or the down-regulation of immune function, potentially acting via the GR signaling

pathway. These findings may enhance the clinical applicability of DEX in achieving transient immune

deficiency.

KEYWORDS: Glucocorticoids; Dexamethasone; Glucocorticoid Receptor; Immune Suppression;

Lymphoid Cell; Myeloid Cell.

Abbreviations

    DCs   dendritic cells

DEX	dexamethasone
DMSO	dimethyl sulfoxide
FCM	flow cytometry
GC	Glucocorticoid
GR	glucocorticoid receptor
HPA	hypothalamic-pituitary–adrenal
ip	intraperitoneal
MHC	major histocompatibility complex
PBMC	peripheral blood mononuclear cells
PBS	phosphate-buffered saline
PR	progesterone receptor
RU486	Mifepristone
TCR	T-cell receptor

Introduction

Glucocorticoids (GCs) are known inhibitors of the immune system. The synthesis and release of GCs are

under dynamic circadian and ultradian regulation by the hypothalamic-pituitary–

adrenal (HPA) axis[1]. Multiple studies have shown that GCs have potent anti-inflammatory and

immunosuppressive properties[2-5]. Natural and synthetic glucocorticoids remain at the forefront of anti-

inflammatory and immunosuppressive therapies used in the treatment of rheumatoid arthritis, multiple

sclerosis, psoriasis, and eczema. Additionally, they are used in the treatment of certain leukemias and to

combat the side effects of some chemotherapies and immunosuppressive regimens following organ

transplant[4, 6, 7]. GCs induce large-scale lymphodepletion that differs from specific depletion mediated

by antibodies in that it occurs predominantly through binding to the glucocorticoid receptor (GR) of

immunocytes. GCs form a complex with the intracellular GR, which induces a conformational change in

the receptor that allows translocation to the nucleus, where it binds to chromatin. This binding to

chromatin leads to the regulation of gene expression, resulting in various anti-inflammatory and

immunosuppressive effects [8-10]. In addition to genomic mechanisms, GCs elicit rapid effects by

interacting directly with enzymes and other cell proteins, including the regulation of signaling

pathways[11]. These rapid effects of GCs make them a promising treatment option for COVID-19, as

they can help reduce the hyperactive immune response that often leads to severe lung inflammation.

Dexamethasone (DEX), a widely used GC, has been shown to effectively achieve immune suppression

and alleviate over-reactive inflammation in the lungs of COVID-19 patients. This treatment approach

can be particularly beneficial for severe cases of the disease, where an excessive immune response plays

a major role in worsening the patient's condition[12]. However, the precise mechanism by which DEX

interacts with immune cells and exerts its anti-inflammatory and immunomodulatory effects in this

specific context is not fully understood. It is believed that DEX acts by inhibiting the production of pro-inflammatory molecules such as cytokines and chemokines, thereby reducing the recruitment and activation of immune cells in the lungs. Additionally, DEX may also interfere with the signaling pathways involved in immune cell activation, further suppressing the hyperactive immune response. Furthermore, the administration of DEX to DC results in a diminished level of major histocompatibility complex (MHC) class II and costimulatory molecules, consequently diminishing their capacity to activate T cells[13], while concurrently inhibiting the signal transmission pathway mediated by the T-cell receptor (TCR)[14]. Some reports have documented that GCs modulate apoptosis by repressing the expression of the Bcl-2 protein, which is encoded by the B-cell lymphoma/leukemia-2 gene[15]. Initial studies on the effects of GCs on B cells have yielded conflicting results, particularly with regards to the dosage relationship between GCs and B cell activity as well as antibody synthesis or levels[16]. GCs have also been found to suppress innate immune cells by regulating monocyte-macrophages. GCs can attenuate the exaggerated activitiy of immune cells that can arise during inflammation[17]. Intriguingly, glucocorticoids have been shown to enhance the accumulation and survival of neutrophils by inducing the expression of anti-apoptotic proteins such as members of the Bcl-2 family [18] , which has contradictory effects on B cells. The regulatory effects of GCs or DEX on the immune system are intricate and still needed to fully understand their comprehensive impact on their effects on innate immune cells such as dendritic cells and monocyte macrophages, as well as their role in adaptive immune cells such as B and T cells. By examining all these aspects, a better understanding of these mechanisms will be crucial in optimizing the use of DEX as a treatment for severe cases of COVID-19 and other virally induced lung inflammations. In this study, we investigated the effects of DEX on the myeloid and lymphoid systems to reveal if the effects of DEX varied across different immune cell types. The effects on

67   monocytes, DCs, macrophages, and NKTs were contingent on the dosage and duration, although CD3 T

68   cells were less affected by the DEX treatments, whereas an enhanced level of neutrophils was evidence.

69   These inhibitions by DEX exhibit a strong affinity for GR in the affected immune cells, suggesting a

70   mechanistic basis for its immunosuppressive properties.

Materials and Methods

Mice and cells lines

Female Balb/c mice were housed and treated according to the guidelines of the Institutional Animal Care and Use Committee of the Fudan University in China. All mice were 6–8 weeks old and age-matched. All experiments were approved by the Committee of Experimental Animals of SHMC with the reference number 202012037S.

Reagents

Dexamethasone (DEX) was purchased from Xi'an Reyphon Pharmaceutical CO (Xi'an, China). Mifepristone (RU486) was purchased from MCE (NJ, USA). Fluorochrome labeled antibodies against CD45(30-F11), CD4(GK1.5), CD8(53-6.7), CD25(3C7), CD11b(M1/70), F4/80, Ly6C(HK1.4), CD49b(DX5), B220(RA3-6B2) were purchased from BioLegend (CA, USA), CD11c(HL3), Ly6G(1A8) were purchased from BD Bioscience (CA, USA) and CD3 (145-2C11), Foxp3 (FJK-16S), fixable viability dye eFluor 780, Foxp3 transcription factor staining buffer kit were from eBioscience (OR, USA).

Treatments

The mice were treated with 100 μg DEX in phosphate-buffered saline (PBS)/animal or vehicle by intraperitoneal injections for 3 consecutive days. Peripheral blood mononuclear cells (PBMC) were collected 24 h after the last administration of DEX. When determining the effect of dexamethasone dosage on immune cells, the doses per mouse were adjusted to 5 mg/kg for high-dose and 1.25 mg/kg for low-dose. Eight micrograms of RU486 were dissolved in 20 μl dimethyl sulfoxide (DMSO), added in 200 μl corn oil (MCE, NJ, USA), and vortexed. Mice were given 400 mg/kg body weight RU486 in corn oil intraperitoneally every 12 h before each DEX administration, and PBMC were collected 72 h after those treatments for FCM analysis (Supplementary 1A).

Flow cytometry

For flow cytometry (FCM), whole blood with anticoagulant was treated with red cell lysis buffer. Single-

cell suspensions were stained with viability dye eflour780 in PBS for 15 minutes at room temperature

and then washed twice with PBS supplemented with 2% FBS. For detection of cell surface antigens, cells

were stained with fluorochrome-tagged antibodies for 15 minutes at room temperature; for detection of

Foxp3, cells were permeabilized and fixed using a commercial transcription factor staining buffer set

(eBioscience, OR, USA). All stained samples were analyzed on an LSR Fortessa (BD, CA, USA) with

FlowJo software (BD, CA, USA).

Statistical analysis

The student's t-test was used to compare continuous variables. Data were analyzed using one-way

ANOVA for single-factor analysis among multiple groups and two-way ANOVA for multiple-

dimensional comparisons. Data are shown as mean $\pm$ SEM or as percentages. P values reporting

statistical significance were calculated in Prism 9 (GraphPad Software). P values less than 0.05 were

deemed significant.

Results

Immunosuppressive Effects of Dexamethasone on Lymphoid and Myeloid Cells

In order to assess the impact of DEX on myeloid-derived innate immune cells, Balb/c mice were

subjected to daily treatment with either 5 mg/kg DEX or vehicle (PBS) for a duration 3 days via the

intraperitoneal (*ip*) route (Fig. 1A). Flow cytometry analysis of PBMC collected on the 4th day (sFig. 1A

and sFig. 2A) demonstrated a significant reduction in the levels of these cells, except for the neutrophils

(Figs. 1B, 1C, and sFig. 1B). Further examination revealed that DEX induced the most substantial

reductions in the proportions of macrophages (95%), DCs (90%), while demonstrating a comparatively

smaller reduction in NK and NKT cells (Fig. 1D). Interestingly, the level of neutrophils was found to be

increased, suggesting the involvement of a distinct regulatory mechanism. Given the relatively short half-

life of DEX in circulation (2-3 days), we sought to explore whether its effects could persist after cessation

of treatment. After 21 days of daily administration, the administrations were halted, and PBMC were

analyzed. The data demonstrated that the proportions of the aforementioned affected myeloid-derived

innate immune cells returned to normal levels 5 days after discontinuation of the drug (Figs. 2A-2F).

Immunosuppressive Effects of Dexamethasone on Lymphoid-Derived Cells

The subsequent investigation aimed to analyze the immunosuppressive effects of DEX on lymphoid-

derived cells, with a particular focus on T and B cells. The same treatment schedule as shown in Figure

1A was followed on Balb/c mice, and flow cytometry analysis was conducted to evaluate any alterations

in $CD3^+$, $CD4^+$, $CD8^+$, $CD4^+CD25^+$ T cells, and $B220^+$ B cells (sFigs 1A and 2A). The results revealed

that all analyzed cells were suppressed by DEX (Fig. 3A and sFig. 2B), with reductions of 72%, 40%,

45%, 40%, and 60%, respectively (Fig. 3B). Furthermore, the duration of the inhibitory effect on each of

these cell types following drug withdrawal was assessed. Unlike myeloid-derived cells, lymphoid-

derived cells exhibited a longer recovery period, taking up to 30 days to regain normal levels after

cessation of treatment (Figs. 4A-4E).

Immunosuppressive Effects Are Related to Dose of Dexamethasone

DEX is used in a wide range of doses to reduce the side effects of chemotherapy in patients with advanced

cancer[7]. To gain insight into the immunoregulatory effects of DEX on the immune system, we selected

high (5 mg/kg) and low (1.25 mg/kg) doses for study in Balb/c mice. We observed that quantities of the

immune cells were reduced less by the lower dose of DEX (Figs. 5A, 5D, and sFig. 3A). However,

macrophages and NKT cells were increased under the low dose of DEX (Figs. 5B-5D, sFigs. 3A, and

3B).

Dexamethasone Mediates Immunosuppression by Binding GR With a High Affinity

Since glucocorticoid-mediated immune modulation is attributed to glucocorticoid receptor-induced

alterations in gene expressions, we investigated whether the immunoregulatory effects of DEX were

related to GR signaling. Mifepristone (RU486), a progesterone receptor (PR) and GR antagonist that

binds to GR with a low affinity, was given intraperitoneally every 12 h before each DEX administration.

PBMC were collected 72 h after those treatments (sFigs. 3A-3B and 4A). We observed that RU486, as a

GR-blocker, abolished the immune regulation by low-dose DEX. The proportion of T and NK cells

returned to normal level (Fig. 6A). When a higher dose of DEX at 5 mg/kg was given, a more robust

immunosuppressive function was obtained and there was less effect of RU486 (Fig. 6B). These data

indicated that DEX was competitively bound to GR with RU486 and a stronger affinity was evident when

given at higher dose.

Discussion

Our results demonstrated DEX's immunoregulatory effects in down-regulating a wide range of immune cells in PBMC, both lymphoid and myeloid cell subsets. The doses of DEX in our study correspond to the doses used in clinical practice [19], suggesting that these findings may be relevant for understanding the immunosuppressive mechanisms of DEX in patients. Additionally, our data showed that the immunosuppressive function of DEX was dose-dependent, with higher doses leading to a stronger effect. This highlights the importance of carefully titrating the dosage of DEX in clinical settings to achieve the desired immunoregulatory outcomes. DEX, a synthetic glucocorticoid, had dramatic suppressive effects on DCs, monocytes, macrophages, NK, T, and B cells, but not on neutrophils and NKs.

Glucocorticoids signal through the glucocorticoid receptor (GR), a member of the superfamily of nuclear receptors that modulates both genomic and non-genomic regulatory pathways in almost every tissue in the bodies of mammals. Observed reductions of immune cells by the DEX treatments could be due to the fact that genomic regulation is receptor-dependent to suppress relevant gene expressions, and maybe also due to a mechanism of non-genomic regulation by inducing apoptosis [20, 21]. It has been reported that glucocorticoids could induce differential biological effects in dendritic cells based on GR isoforms to attenuate DC activity and maturation in one way [22] and in another to promote the differentiation of 'tolerogenic' DCs. Tolerance follows from down-regulation of the expression of MHC class II, co-stimulatory molecules (for example, CD80 and CD86) and pro-inflammatory cytokines (such as IL-12 and TNF) resulting in Treg induction [23]. Glucocorticoids also dampen T cell activation by interfering with TCR signaling and inhibition of interleukin (IL)-2 synthesis via binding to GR inside T cell cytoplasm has been reported[24]. Neutrophils are the earliest responding cell of the innate immune response to infections, particularly bacterial infections, and mount the initial inflammatory response.

They are also involved in the initiation of the inflammatory response by releasing pro-inflammatory

cytokines and chemokines. Interestingly, we found that the proportion of neutrophils was increased by

the DEX treatment, in direct contrast to the other immune cells. Observed enhanced neutrophils by the

DEX treatments may achieve this by promoting the differentiation and activation of neutrophil progenitor

cells in the bone marrow, leading to an increased production and release of mature neutrophils into

circulation. The enhanced effect on the neutrophils could also be explained by recent studies in which

neutrophils were found to be the cells more weakly responding to GCs and more resistant to DEX-

induced apoptosis by mechanisms that included upregulation of anti-apoptotic proteins such as Mcl-1

and XIAP[25]. Alternatively, several studies have shown that neutrophils can also have

immunosuppressive functions. They can suppress T cell proliferation and cytokine production, as well

as promote the differentiation of Tregs[26, 27]. This dual role of neutrophils highlights their importance

in maintaining immune homeostasis and preventing excessive inflammation.

Regarding B cells, immature B cells are more sensitive to glucocorticoid-induced apoptosis than mature

B cells[15, 28], suggesting a regulatory role for glucocorticoids in B cell development. Additionally,

glucocorticoids can also modulate B cell activation, proliferation, and antibody production. This

highlights the complex interplay between neutrophils, B cells, and glucocorticoids in immune regulation

and the delicate balance required for maintaining a properly functioning immune system, but more details

of the effects of GCs on humoral immunity are needed.

Although there have been many studies and clinical reports of glucocorticoid effects on immunity during

the preceding 60 years, this study has focused on the landscape of immunosuppression as contributed by

key immune cell types, particularly those with roles in adaptive immunity. To our surprise, we observed

that low DEX doses could increase the ratio of macrophages, while a reduction of macrophages was

achieved when DEX was at a high dose. This double-edged sword effect of GCs on macrophages had

not been reported previously and suggests a complex regulation via the GR signaling: at a low

concentration, GC was reported to positively regulate NLRP3, a component of the inflammasome

complex in macrophages, and thereby augment the pro-inflammatory response[29]. An alternative

explanation might be that low GC concentrations, in the nanomolar range, exert an immunostimulatory

effect on a range of macrophage functions, including adhesion, chemotaxis, phagocytosis, and cytokine

production[30]. The macrophage is a crucial bridging arm from innate to adaptive immune activation

and expresses pattern-recognition receptors, including Toll-like receptors that sense infectious agents and

harmful signals, hence activating the inflammasome complex and secretion of inflammatory

cytokines[31]. This cascade of reactivities serves to elicit adaptive immune responses.

Our findings confirm that DEX could depress immune cells via a high-affinity binding of GR but also

show that this could be reversed to a normal level when using RU486 to block the GR.

Treating severe cases of COVID-19 by duration with low doses of Dexamethasone (DEX) has displayed

promising outcomes in terms of reducing inflammation and enhancing patient outcomes. Nonetheless, it

is crucial to exercise caution when employing DEX due to its potential to exert immunosuppressive

effects, which could potentially impede the body's capability to mount an efficient immune response

against the virus and secondary infections. Consequently, further investigation is warranted to ascertain

the most optimal dosage and duration of DEX treatment in COVID-19 patients, with the objective of

striking a delicate balance between curbing excessive inflammation and preserving an active immune

response.

In conclusion, our study shows that treatment with a high dose of DEX negatively affects the numbers

of both myeloid and lymphoid cells, except for neutrophils and NKT. Although myeloid cells returned to

normal within 5 days of withdrawing the DEX treatment, the treatment affected lymphoid cells

profoundly, and they took longer to recover. These findings will facilitate further understanding of the

complexity of GCs effects on the immune responses and lead to a more economical use of the drug to

achieve broad immune suppression.

Statements and Declarations

Funding

The project leading to these results has received fundings from the Chinese National Natural Science

Foundation (81991492 and 82041039) to B. Wang.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author contributions

Bin Wang and Shuting Wu conceived, directed the experiments with input from all authors. Material

preparation, data collection and analysis were performed by Shuting Wu. Shuting Wu, Shushu Zhao, Gan

Zhao and Yiwei Zhong performed data analysis and interpretation. The first draft of the manuscript was

written by Shuting Wu and all authors commented on previous versions of the manuscript. All authors

read and approved the final manuscript.

Data Availability

The data generated in this study are available upon request from the corresponding author.

Ethics approval

This study was performed in line with the guidelines of the Institutional Animal Care and Use Committee

of the Fudan University in China. All experiments were approved by the Committee of Experimental

Animals of SHMC with the reference number 202012037S.

References

Biddie SC, Conway-Campbell BL, Lightman SL (2012) Dynamic regulation of glucocorticoid signalling in health and disease. *Rheumatology (Oxford)* 51: 403-412

Kizaki T, Oh-ishi S, Ookawara T, Yamamoto M, Izawa T, Ohno H (1996) Glucocorticoid-mediated generation of suppressor macrophages with high density Fc gamma RII during acute cold stress. *Endocrinology* 137: 4260-4267

Reed JC, Abidi AH, Alpers JD, Hoover RG, Robb RJ, Nowell PC (1986) Effect of cyclosporin A and dexamethasone on interleukin 2 receptor gene expression. *J Immunol* 137: 150-154

Giles AJ, Hutchinson MND, Sonnemann HM, Jung J, Fecci PE, Ratnam NM, Zhang W, Song H, Bailey R, Davis D et al (2018) Dexamethasone-induced immunosuppression: mechanisms and implications for immunotherapy. *J Immunother Cancer* 6: 51

Unger WW, Laban S, Kleijwegt FS, van der Slik AR, Roep BO (2009) Induction of Treg by monocyte-derived DC modulated by vitamin D3 or dexamethasone: differential role for PD-L1. *Eur J Immunol* 39: 3147-3159

Ronchetti S, Migliorati G, Bruscoli S, Riccardi C (2018) Defining the role of glucocorticoids in inflammation. *Clin Sci (Lond)* 132: 1529-1543

Shih A, Jackson KC, 2nd (2007) Role of corticosteroids in palliative care. *J Pain Palliat Care Pharmacother* 21: 69-76

Lu NZ, Cidlowski JA (2006) Glucocorticoid receptor isoforms generate transcription specificity. *Trends Cell Biol* 16: 301-307

Stahn C, Buttgereit F (2008) Genomic and nongenomic effects of glucocorticoids. *Nat Clin Pract Rheumatol* 4: 525-533

Grad I, Picard D (2007) The glucocorticoid responses are shaped by molecular chaperones. *Mol Cell Endocrinol* 275: 2-12

Oppong E, Cato AC (2015) Effects of Glucocorticoids in the Immune System. *Adv Exp Med Biol* 872: 217-233

Ledford H (2020) Coronavirus breakthrough: dexamethasone is first drug shown to save lives. *Nature* 582: 469

Vanderheyde N, Verhasselt V, Goldman M, Willems F (1999) Inhibition of human dendritic cell functions by methylprednisolone. *Transplantation* 67: 1342-1347

Lowenberg M, Verhaar AP, Bilderbeek J, Marle J, Buttgereit F, Peppelenbosch MP, van Deventer SJ, Hommes DW (2006) Glucocorticoids cause rapid dissociation of a T-cell-receptor-associated protein complex containing LCK and FYN. *EMBO Rep* 7: 1023-1029

Akiyama M, Yamada O, Agawa M, Yuza Y, Yanagisawa T, Eto Y, Yamada H (2008) Effects of prednisolone on specifically expressed genes in pediatric acute B-lymphoblastic leukemia. *J Pediatr Hematol Oncol* 30: 313-316

Alnemri ES, Fernandes TF, Haldar S, Croce CM, Litwack G (1992) Involvement of BCL-2 in glucocorticoid-induced apoptosis of human pre-B-leukemias. *Cancer Res* 52: 491-495

Cain DW, Cidlowski JA (2017) Immune regulation by glucocorticoids. *Nat Rev Immunol* 17: 233-247

Saffar AS, Ashdown H, Gounni AS (2011) The molecular mechanisms of glucocorticoids-mediated neutrophil survival. *Curr Drug Targets* 12: 556-562

Czock D, Keller F, Rasche FM, Haussler U (2005) Pharmacokinetics and pharmacodynamics of systemically administered glucocorticoids. *Clin Pharmacokinet* 44: 61-98

Boldizsar F, Talaber G, Szabo M, Bartis D, Palinkas L, Nemeth P, Berki T (2010) Emerging pathways of non-genomic glucocorticoid (GC) signalling in T cells. *Immunobiology* 215: 521-526

Biddie SC, John S, Hager GL (2010) Genome-wide mechanisms of nuclear receptor action. *Trends Endocrinol Metab* 21: 3-9

Cao Y, Bender IK, Konstantinidis AK, Shin SC, Jewell CM, Cidlowski JA, Schleimer RP, Lu NZ (2013) Glucocorticoid receptor translational isoforms underlie maturational stage-specific glucocorticoid sensitivities of dendritic cells in mice and humans. *Blood* 121: 1553-1562

Chamorro S, Garcia-Vallejo JJ, Unger WW, Fernandes RJ, Bruijns SC, Laban S, Roep BO, t Hart BA, van Kooyk Y (2009) TLR triggering on tolerogenic dendritic cells results in TLR2 up-regulation and a reduced proinflammatory immune program. *J Immunol* 183: 2984-2994

Northrop JP, Crabtree GR, Mattila PS (1992) Negative regulation of interleukin 2 transcription by the glucocorticoid receptor. *J Exp Med* 175: 1235-1245

Ronchetti S, Ricci E, Migliorati G, Gentili M, Riccardi C (2018) How Glucocorticoids Affect the Neutrophil Life. *International Journal of Molecular Sciences* 19

Langereis JD, Oudijk EJ, Schweizer RC, Lammers JW, Koenderman L, Ulfman LH (2011) Steroids induce a disequilibrium of secreted interleukin-1 receptor antagonist and interleukin-1beta synthesis by human neutrophils. *Eur Respir J* 37: 406-415

Leliefeld PH, Wessels CM, Leenen LP, Koenderman L, Pillay J (2016) The role of neutrophils in immune dysfunction during severe inflammation. *Crit Care* 20: 73

Garvy BA, King LE, Telford WG, Morford LA, Fraker PJ (1993) Chronic elevation of plasma corticosterone causes reductions in the number of cycling cells of the B lineage in murine bone marrow and induces apoptosis. *Immunology* 80: 587-592

Hermoso MA, Matsuguchi T, Smoak K, Cidlowski JA (2004) Glucocorticoids and tumor necrosis factor alpha cooperatively regulate toll-like receptor 2 gene expression. *Mol Cell Biol* 24: 4743-4756

Zhou JY, Zhong HJ, Yang C, Yan J, Wang HY, Jiang JX (2010) Corticosterone exerts immunostimulatory effects on macrophages via endoplasmic reticulum stress. *Br J Surg* 97: 281-293

Busillo JM, Azzam KM, Cidlowski JA (2011) Glucocorticoids sensitize the innate immune system through regulation of the NLRP3 inflammasome. *J Biol Chem* 286: 38703-38713

Figure & Figure Legends

Figure 1

Fig. 1 DEX's immunoregulatory effects on myeloid-derived innate immune cells

(A) Diagram of DEX treatment of mice. PBMC were isolated at 96 h after treatment with DEX and analyzed by flow cytometry (**B, C**). Antibodies to Ly6G, CD11c, F4/80, Ly6C, and CD49b were used to identify neutrophils, DC, macrophages, monocytes, and NK cells. Antibodies to CD3 and CD49b were used to identify NKT cells. The results are presented in percentages of CD11b+ cells (**B**) and CD45+ cells (**C**); (**D**) shows the proportion of the decline in each innate immune cell type. Error bars indicate ± SEM, **** P < 0.0001, n = 5. Statistical significance was determined with an unpaired two-tailed T-test. Data are representative of two independent experiments.

Figure 2

Fig. 2 Recovery curves of innate immune cells after drug withdrawal

PBMC were isolated on days 5, 9, 15, 20, and 30 after DEX withdrawal and analyzed by flow cytometry.

Antibodies to Ly6G, CD11c, F4/80, Ly6C, and CD49b were used to identify neutrophils, DC cells,

macrophages, monocytes, and NK cells; antibodies to CD3 and CD49b were used to identify NKT cells.

Results from the DEX treatment group are shown in blue and from PBS vehicle group in red. Shown are

the percentages of CD11b$^+$ neutrophil cells (**A**), CD11b$^+$ DC cells (**B**), CD11b$^+$ macrophages (**C**),

CD11b$^+$ monocytes (**D**), CD45$^+$ NK cells (**E**), and CD45$^+$ NKT cells (**F**) Error bars indicate ± SEM, ****

P < 0.0001. Statistical significance was determined with an unpaired two-tailed T-test. Data are

representative of two independent experiments.

Figure 3

Fig. 3 DEX's immunoregulatory effects on lymphatic cells

PBMC were isolated 96 h after treatment with 5 mg/kg DEX daily and analyzed by flow cytometry.

Antibody to B220 was used to identify B cells. The results are presented as percentages of CD45$^+$ cells

(**A**) and the percentage of the decline in each lymphatic immune cell type (**B**). Error bars indicate ± SEM,

**** $P < 0.0001$, n = 5. Statistical significance was determined with an unpaired two-tailed T-test. Data

are representative of two independent experiments.

Figure 4

Fig. 4 Recovery curves of lymphatic cells after drug withdrawal

PBMC were isolated on days 9, 20, and 30 after DEX withdrawal and T cell subsets in CD45$^+$ cells analyzed by flow cytometry. Antibody to B220 was used to identify B cells. Results from the DEX treatment group are shown in blue and from PBS vehicle group in red. Shown are the percentages of CD3$^+$T cells (**A**), CD4$^+$T cells (**B**), CD4$^+$CD25$^+$T cells(**C**), CD8$^+$T cells(**D**), B220$^+$B cells (**E**). The results are presented as percentages in CD45$^+$ cells. Error bars indicate ± SEM, **** $P < 0.0001$. Statistical significance was determined with an unpaired two-tailed T-test. Data are representative of two independent experiments.

Figure 5

Fig. 5 Immunosuppressive effects under different doses of DEX

PBMC were isolated on day 8 after treatment with different doses of DEX and analyzed by flow cytometry. Antibodies to Ly6G, CD11c, F4/80, Ly6C, CD49b, CD3, Foxp3, and B220 were used to identify neutrophils, DC cells, macrophages, monocytes, NK, T, Treg, and B cells, respectively. Antibodies to CD3 and CD49b were used to identify NKT cells. The results are presented as percentages in CD11b$^+$ cells (**A**) and CD45$^+$ cells (**B, C**) and the proportion of the decline in each immune cell type (**D**). Error bars indicate ± SEM, **** P < 0.0001. Statistical significance was determined with an unpaired one-way ANOVA test.

Figure 6

Fig. 6 The high-affinity binding of DEX to GR reduces immune cells

PBMC were isolated and analyzed by flow cytometry 3 days after treatment with 1.25 mg/kg DEX (**A**)

or 5 mg/kg DEX (**B**). Antibodies to B220, Foxp3, and CD49b were used to identify B, Treg, and NK

cells. Antibodies to CD3 and CD49b were used to identify NKT cells. The results are presented as

percentages in CD45$^+$ cells. Error bars indicate ± SEM, **** P < 0.0001. Statistical significance was

determined with an unpaired one-way ANOVA test.