Trophic eggs affect caste determination in the ant *Pogonomyrmex rugosus*

E. Genzoni[1], T. Schwander[1] and L. Keller[1]

[1] Department of Ecology and Evolution, Biophore Building, University of Lausanne, 1015 Lausanne Switzerland

Author contributions: EG, TS and LK designed the study. EG performed the experiment and analysed the data. EG and LK wrote the manuscript with input from TS.

Abstract

Understanding how a single genome creates distinct phenotypes remains a fundamental challenge for biologists. Social insects provide a striking example of polyphenism, with queen and worker castes exhibiting morphological, behavioural, and reproductive differences. Here we show that trophic eggs, which do not contain an embryo and are primarily regarded as a source of food, play a role in the process of caste determination in the harvester ant *Pogonomyrmex rugosus*. When first instar larvae were given access to trophic eggs, they mostly developed into workers. By contrast, larvae without access to trophic eggs developed into queens. We found that trophic eggs differ in many ways from reproductive eggs, including texture, morphology and their contents of protein, triglycerides, glycogen, sugar and small RNAs. Moreover, comparison of miRNA fragment size distributions suggests differences in the composition of miRNAs between the two egg types. This is the first demonstration of trophic eggs playing a role in caste determination in social insects.

Keyword: Trophic eggs, *Pogonomyrmex rugosus* ant, caste determination, miRNA, triglycerides, glycogen, proteins

Introduction

Many species of insects, spiders, amphibians, marine invertebrates and sharks produce trophic eggs, a special type of eggs that do not contain an embryo (Levin and Bridges 1995; Blake and Arnofsky 1999; Collin 2004; Kudo and Nakahira 2004; Perry and Roitberg 2006; Strathmann and Strathmann 2006; Gibson *et al.* 2012; López-Ortega and Williams 2018). It is generally assumed that these non-developing eggs are either a by-product of failed reproduction or that they serve as nutrition for offspring (Perry and Roitberg 2006). However, the suggestion that trophic eggs solely provide a nutritional function is based on surprisingly little evidence. We here report a direct function of trophic eggs in the determination of alternative phenotypes in ants.

Trophic eggs have been reported in many ant species (Figure 1 and Supplementary Table 1), where they are also thought to mostly or only serve as food for the offspring (Crespi 1977). Although they are believed to be generally laid by workers, they can also be laid by the queens (see Figure 1 and Supplementary Table 1). While conducting egg cross fostering experiments in the ant *Pogonomyrmex rugosus* to study worker size variation, we observed a sudden increase in the frequency of females developing into queens. During these experiments, we only cross fostered reproductive eggs, raising the possibility that the absence of trophic eggs influenced the process of caste determination. Similarly, changes in the ratio of trophic and reproductive eggs has been proposed as a possible explanation for changes in worker size in another *Pogonomyrmex* species following treatments with a JH analogue (Helms Cahan *et al.* 2011). Because these treatments resulted in an increased proportion of trophic eggs, the authors proposed that the presence of trophic eggs might have altered the nutritional environment of developing larvae, thereby leading to increased worker size. These observations prompted us to investigate whether trophic eggs play a role in caste determination in *P. rugosus*. Our experiments show that the presence of trophic eggs reduces the probability that female larvae develop into reproductive individuals. Metabolomic analyses also revealed profound differences between reproductive and trophic eggs, including in the composition of miRNAs and content of protein, triglycerides, glycogen, and sugar.

	Queen	Worker	Queen & worker	None
Myrmicinae	4	11	9	
Ectatomminae		5		
Heteroponerinae				
Formicinae		6	1?	
Dolichoderinae		2	1	
Aneuretinae				
Pseudomyrmecinae				
Myrmeciinae		1	1	
Dorylinae				
Amblyoponinae		2		1
Apomyrminae				
Proceratiinae				
Agroecomyrmecinae				
Paraponerinae				
Ponerinae		4		
Leptanillinae				
Martialinae				
Polistes				

Figure 1. Trophic egg production is widespread in ants. Simplified phylogenetic tree of ant families redrawn after Romiguier *et al.* (2022). The number of species with documented trophic egg production by queens, workers or both castes, as well as absence of trophic eggs, is indicated for each family. Details on the species and related references can be found in Supplementary Table 1.

Materials and methods

Pogonomyrmex rugosus colonies were initiated by queens collected after mating flights in 2008 (Bowie, Arizona, USA) or 2013 (Bowie and Florence, Arizona, USA). The colonies were maintained at 28°C and 60% humidity, with a 12-h/12-h light:dark cycle and fed once a week with grass seeds, flies and 20% honey water. Eggs were collected in October 2020 for the experiment investigating the effect of trophic eggs on larval caste fate, in November 2021 for the percentage of trophic eggs and from February to December 2021 for the egg content analysis.

Trophic and reproductive egg production

To verify that workers do not lay trophic eggs, as previously shown for other *Pogonomyrmex* species (Supplementary Table 1), we created 12 queenless colonies and waited approximately three weeks until workers started laying eggs. From each of these colonies, we isolated two groups of five workers for 12 hours every two days for two weeks in November 2020 to obtain eggs. Collected eggs were then placed for 10 days in a petri dish containing a water reservoir to study their development and distinguish whether they were trophic or reproductive.

To determine whether queens lay variable percentages of trophic eggs over time, we isolated 43 *P. rugosus* queens for 8 hours every day for 2 weeks, before and after hibernation, and counted the number of trophic and reproductive eggs laid (see results for how to discriminate the two types of eggs). The percentage of trophic eggs was compared using a linear mixed effect model within Rstudio (RStudio Team 2015), with before vs after hibernation as the explanatory variable and colony as a random factor.

To inform on the mechanisms of oogenesis, we assessed whether reproductive and trophic eggs were laid in a
random order, or whether eggs of a given type were laid in clusters. To this end, we isolated 11 queens for 10
hours, eight times over three weeks, and collected every hour the eggs laid. To determine whether reproductive
and trophic eggs were laid in a random order, we used Rstudio (package *snpar* v.1.0; RStudio Team 2015) to
perform a Wald–Wolfowitz runs test for each queen's egg laying sequence (this non-parametric test calculates the
likelihood that a binomial data sequence is random).

Trophic egg influence on the larval caste fate

To determine whether trophic eggs influence the process of caste determination, we compared the development
of freshly hatched (first instar) larvae placed in small recipient colonies with and without trophic eggs. From each
of 22 donor colonies, we obtained approximately 30 freshly hatched larvae by isolating the queens for 16 hours
(from 2pm to 6am) every day for three weeks (in October 2020), with a 24-hour break every three days. Eggs were
collected every eight hours and placed during 10 days in a petri-dish with a water reservoir ensuring a high humidity
until they hatched. After hatching, half of the larvae were then transferred into a recipient colony containing 20
workers, while the other half of the larvae were placed in identical recipient colonies, which received in addition
three 0-4 hours-old trophic eggs. There was no cross-fostering between colonies, so that larvae were always
placed in recipient colonies containing workers from the same donor colony. The recipient colonies were
maintained at 28°C and 60% humidity, with a 12-h/12-h light:dark cycle and fed twice a week with grass seeds,
flies and 20% honey water. The caste of each individual was recorded at the pupal stage. To compare the
proportion of queen pupae produced between recipient colonies with and without trophic eggs, we used Rstudio
(RStudio Team 2015) and the package *lme4* (Bates *et al.* 2015) to perform a binomial generalized linear mixed
effects analysis (GLMM) fit by maximum likelihood, with caste as response variable (binary categorical factor) and
presence/absence of trophic eggs as an explanatory variable. Donor colony was included as a random effect. To
test whether the presence of trophic eggs affects survival, we performed a linear mixed effect analysis with
mortality as a response variable, presence/absence of trophic eggs as explanatory variable, and colonies as
random effects. As we found a significantly higher survival of larvae in recipient colonies with trophic eggs than
recipient colonies without trophic eggs (see results), we tested whether the percentage of larvae developing into
queens was correlated with survival by performing a linear mixed effects analysis with the percentage of queen
pupae as response variable, the survival as an explanatory variable and colonies as a random factor.

Volume and content of trophic and reproductive eggs

The volumes of trophic (n=11) and reproductive eggs (n=14) were estimated by using the volume of an ellipse
$(\frac{4}{3} \times \pi \times egg\ lenght \times \left(\frac{egg\ width}{2}\right)^2)$, with egg length and width estimated on images under 10x magnification using
ZEN Microscopy Software (v. 1.1.2.0).

To determine the nutritional content of reproductive and trophic eggs, we quantified the proteins, triglycerides,
glycogen, and glucose in both types of eggs. We also quantified long and small RNAs (including miRNAs) as these

compounds have been shown to be involved in caste determination in other eusocial species (Suzzoni *et al.* 1979;

Schwander *et al.* 2008a; Helms Cahan *et al.* 2011; Cameron *et al.* 2013; Guo *et al.* 2013; Libbrecht *et al.* 2013;

Søvik *et al.* 2015; Collins *et al.* 2017). To obtain the two types of eggs, we isolated 12 queens for 10 hours (7am

to 5pm; from March to October 2021) in a dark petri-dish with three workers and a water supply. Eggs were

collected every hour (so all eggs were a maximum of one hour old), and trophic and reproductive eggs were flash-

frozen separately in liquid nitrogen. Twenty eggs were pooled for triglycerides-sugar-protein analyses and six eggs

for RNA analyses. They were kept at -80°C until the extractions were performed. After the 10 hours of isolation,

queens and workers were returned to their colony until the next isolation session. For each of the 12 colonies, we

obtained two replicates of reproductive and trophic egg pools (i.e., 24 replicates in total).

Triglycerides, glycogen and glucose were quantified as described in Tennessen *et al.* (2014), and protein levels

were measured using a Bradford assay (Bradford 1976). The 20 one-hour old eggs per sample were homogenized

with beads in 200µl of PBS buffer in a Precellys Evolution tissue homogenizer coupled with a Cryolys Evolution

(Bertin Technologies SAS).

For the Bradford assay, 10μl of the homogenate were put in a clear-bottom 96-well plate with 300μl of Coomassie

Plus Reagent (Thermo Scientific: 23200) and incubated for 10 minutes at room temperature. Protein standard

(Sigma: P5369) was used as standard (ranging from 0-0.5mg/ml) and protein absorbance was read at 595nm on

a Hidex Sense Microplate Reader.

For the triglycerides assay, 90μl of homogenate were heat treated at 70°C for 10 minutes, then 40μl were mixed

with 40μl of Triglyceride Reagent (Sigma: T2449) for digestion and 40μl were mixed with PBS buffer for free

glycerol measurement. After 30 minutes incubation at 37°C, 30μl of each sample and standards were transferred

to clear-bottom 96-well plate. 100μl of Free Glycerol Reagent (Sigma: F6428) was added to each sample, mixed

well by pipetting, and incubated five minutes at 37°C. Glycerol standard solution (Sigma: G7793) was used as

standard (ranging from 0-1.0mg/ml TAG) and absorbance was read at 540nm on a Hidex Sense Microplate

Reader. The triglycerides concentration in each sample was determined by subtracting the absorbance of free

glycerol in the corresponding sample.

Glucose and glycogen were quantified as in Tennessen *et al.* (2014). A 90μl aliquot was heat treated at 70°C for

10min and then diluted 1:2 with PBS. The standard curves for glucose (Sigma, G6918) and glycogen (Sigma:

G0885) were made by diluting stocks to 160μg/ml, making 1:1 serial dilution for 160, 80, 40, 20 and 10μg/ml. 40μl

of each sample was pipetted in duplicates of a clear microplate, and 30μl of each glucose or glycogen standard

was pipetted in duplicates. Amyloglucosidase enzyme (Sigma, A1602) was diluted 3μl into 2000μl of PBS, and

40μl diluted enzyme was pipetted to the glycogen standards and to one well of the sample (for total glucose

determination), 40μl PBS was pipetted to the glucose standards and to the other sample well (for free glucose

determination). The plate was incubated at 37°C for 60 minutes. 30μl of each standard and samples (in duplicates)

were transferred to a UV 96-well plate and 100μl Glucose Assay Reagent (G3293) was pipetted to each well. The

plate was incubated at room temperature for 15 minutes and the absorbance was read at 340nm on a Hidex Sense

Microplate Reader. The glycogen concentration was quantified by subtracting the free glucose absorbance from

the total glycogen + glucose absorbance.

Concentrations of each compound (protein, triglycerides, glycogen, and glucose) were compared between

reproductive and trophic eggs with Rstudio (RStudio Team 2015) and the package *lme4* (Bates *et al.* 2015) using

a linear mixed effects analysis (LMER), with the concentration as response variable and egg type as explanatory

variable. Colony and extraction batch were added as random effects in the model.

Total and small RNA, and DNA

RNA (>200 nt) and small RNA were isolated using the miRNeasy Mini Kit (Qiagen, cat. no. 217004) and RNeasy®

MinElute® Cleanup Kit (Qiagen, cat. no. 74204), respectively, following manufacturer instructions. RNA (>200 nt)

and small RNA concentrations were measured with a QuantiFluor® RNA System (Promega). RNA (>200 nt)

integrity was examined with an Agilent Fragment Analyzer (at the Lausanne Genomic Technologies Facility)

using a High Sensitivity Assay and small RNA were examined using the small RNA kit (at the Gene Expression

Core Facility at EPFL).

The miRNA and RNA (>200nt) concentrations were compared between reproductive and trophic eggs with paired-

t-tests in Rstudio (RStudio Team 2015; for each type of eggs we used the average of the two replicates per colony).

We also compared the fragment size distributions from 18 to 24 nucleotides for miRNAs (Sohel 2016) with a Mantel

test.

DNA was extracted from pools of six eggs using TRIzol (Life Technologies). DNA concentration was measured

with a Nanodrop 3300 (ThermoFisher), and DNA integrity was examined with an Agilent Fragment Analyzer (at

the Lausanne Genomic Technologies Facility) using a High Sensitivity Assay. DNA concentrations were compared

between reproductive and trophic eggs using paired-t-tests (sample size is 5 for both types of eggs, each sample

being a pool of 6 eggs).

Results

Trophic and reproductive egg characteristics

P. rugosus queens lay two types of eggs that are morphologically different. Reproductive eggs are white with a

bright surface and have a distinct oval shape, a homogenous content as well as a solid chorion (Figure 2A), while

trophic eggs are rounder, have a smooth surface and a granular looking content as well as a fragile chorion (Figure

2D). Trophic eggs had a significantly larger volume (94.3±4.3nL; n=11) than reproductive eggs (n=14; 63.3±1.6nL;

two-sample t-test, t(23) = -9.54, p = 1.8×10^{-09}). *P. rugosus* workers only laid reproductive eggs. They started to lay

eggs approximately three weeks after queen removal (n=12 queenless recipient colonies) and approximately 90%

of the eggs successfully hatched. However, only approximately 5% successfully developed into pupae which were

all males.

Figure 2. Morphology and development of eggs laid by *P. rugosus* queens. Reproductive egg general morphology (**A**), with embryonic development at approximately 25 hours (**B**) and 65 hours (**C**). In trophic eggs (**D**), there is no embryonic development at 25 hours (**E**) nor at 65 hours (**F**). Panels B, C, E, F represent fluorescence images with DAPI-counterstained nuclei.

The percentage of eggs that were trophic was higher before hibernation (61.6$\pm$1.4% mean$\pm$ SE; n=43 colonies) than after (50.3$\pm$2.0%; LMER, t(86)=5.04, p=9x10^{-6}). This higher percentage was due to a reduced number of reproductive eggs, the number of trophic eggs laid remained stable. The production of the two types of eggs was not random (Wald-Wolfowitz runs tests, p-values for the 11 queens in Table 1). Instead, each of the 11 queens tended to lay relatively long sequences of either reproductive (6.1$\pm$0.7; mean number per sequence $\pm$SE) or trophic eggs (6.0$\pm$0.5; Figure 3).

Figure 3. Egg laying sequences from eleven *P. rugosus* queens. Every row shows the sequence of reproductive R and trophic T eggs laid by a given queen (queen ID in the orange cell). Each egg laying session lasted 10 hours. Each yellow square separates two egg laying sessions and represents an interval of minimum 16 hours to several days.

Table 1. Wald-Wolfowitz runs tests on the queen's egg sequence. Significant p-values (corrected for multiple testing) indicate that queens do not lay reproductive and trophic eggs in a random sequence.

Queen ID	P-value for egg sequence	P-value of random sequence	Number of eggs per sequence
338	$4.1*10^{-11}$	0.419	94
117	$4.3*10^{-07}$	0.567	63
173	$1.7*10^{-03}$	0.755	92
303	$4.8*10^{-13}$	0.765	110
215	$9.8*10^{-11}$	0.292	70
120	$1.4*10^{-05}$	0.518	75
12B	$1.9*10^{-03}$	0.298	38
316	$3.4*10^{-09}$	0.737	93
193	$4.3*10^{-12}$	0.655	101
150	$1.4*10^{-10}$	0.630	62
125	$1.5*10^{-05}$	0.404	58

185 The concentrations of protein, triglycerides, glycogen, and glucose were significantly higher in reproductive than

186 trophic eggs (LMER, protein: t = -13.11, p <0.0001; triglycerides: t = -11.66, p <0.0001; glycogen: t = -11.98,

187 p <0.0001; glucose: t = -18.60, p <0.0001; Figure 4).

Figure 4. Concentration of protein (**A**), triglycerides (**B**), glycogen (**C**) and glucose (**D**) in reproductive and trophic eggs. Each dot represents the average of the two replicates per colony.

The amount of small RNA (<200 nt, including miRNA and tRNA; Nagano and Fraser 2011) was significantly higher in reproductive eggs (44.3±1.4ng, mean ± SE) than in trophic eggs (22.3±1.1ng; paired-t-test, $t_{(23)}$ = 15.9, p = $6.5*10^{-14}$). The same was true for longer RNAs (>200 nt; reproductive eggs: 7.6±0.6ng, mean ± SE; trophic eggs: 3.6±0.3ng; paired-t-test, $t_{(23)}$ = 7.2, p = $2.7*10^{-7}$).

The DNA quantification showed that the amount of DNA was about twice higher in reproductive (15.9$\pm$1.9ng/μl)
than trophic eggs (8.8$\pm$1.9ng/μl; t-test, $t_{(4.7)}$ = 2.7, p = 0.045).

There was a significant difference in the miRNA fragment size distribution between reproductive and trophic eggs
(Mantel test, r_M = 0.26, p<.0001), as shown on the PCA (Figure 5A). There was no difference in the tRNA fragment
size distribution between the two types of eggs (Mantel test, r_M = 0.01, p=0.30, Figure 5B).

Figure 5. First two principal components (PC1 and PC2) explaining size distribution variation for (**A**) miRNA and (**B**)
tRNA across egg samples, with reproductive eggs in grey and trophic eggs in black. There is a separation of the samples
by egg types for miRNAs, but not for tRNA. Ellipses enclose each of the egg type groups.

Trophic eggs influence caste fate of larvae

The percentage of larvae that developed into queens was significantly lower in recipient colonies that received trophic eggs (27±9% mean±SE; n=22) than in recipient colonies without trophic eggs (83±10%; n=22; binomial GLMM, z = 4.25, p = 2x10⁻⁵; Figure 6A and Supplementary Table 2). The survival of larvae until the pupal stage was also significantly lower in the colonies without trophic eggs (16.9±3.8%; n=22; LMER, z = 2.66, p = 0.008) than in colonies with trophic eggs (30.2±6.7%; mean±SE; n=22) , but the 1.8 fold survival decrease cannot fully account for the 3 fold difference in queen percentage between the two treatments. Furthermore, there was no significant correlation between larval mortality and the percentage of larvae developing into queens (n=44 recipient colonies; LMER, z = 0.97, p = 0.34; Figure 6B). These analyses allow us to exclude differential survival between castes as an explanation for the higher percentage of queens developing in the recipient colonies without trophic eggs.

Figure 6. (**A**) Percentage of larvae which developed into queens in recipient colonies without (grey) or with (black) trophic eggs. (**B**) Scatterplot of the relationship between the percentage of larvae which developed into queens and the percentage of survival from larvae to pupae.

Discussion

Our study reveals that *P. rugosus* queens lay a very high proportion (0.6) of trophic eggs. These eggs differ in many ways from reproductive eggs. First, trophic eggs are larger, rounder, have a smoother surface, a more granular looking content as well as a more fragile chorion than reproductive eggs. Similar differences between trophic and reproductive eggs have been reported in other ant species (Wilson 1976; Wardlaw and Elmes 1995; Gobin *et al.* 1998; Dietemann and Peeters 2000; Dietemann *et al.* 2002; Perry and Roitberg 2006; Lee *et al.* 2017).

Our analyses also showed that trophic eggs are solely laid by queens; *P. rugosus* workers are able to produce
reproductive eggs which occasionally develop into males, but they do not lay trophic eggs. Moreover, trophic eggs
have a reduced DNA content.

Importantly, our experiments showed that the presence of trophic eggs influences the process of caste
determination. First instar female larvae fed with trophic eggs were significantly more likely to develop into workers
than larvae without access to trophic eggs. This was somewhat surprising because trophic eggs are generally
thought to be an important source of nutrients to the colony and, everything else being equal, one would think that
eating such eggs should increase the likelihood of females to develop into queens (which are usually larger than
workers). Indeed, a study in the Argentine ant *Linepithema humile* showed that the presence of queens in colonies
was associated with a drastic decrease in the number of trophic eggs (laid by workers) fed to the larvae as well as
a decrease in the proportion of larvae developing into queens (Bartels 1988). Bartels thus proposed that the
deprivation of trophic eggs may have an inhibitory effect on the probability of larvae to develop into queens (Bartels
1988), but no experiment was performed to show a causal effect of trophic eggs. In some lineages of *P. barbatus*,
the experimental increase of maternal juvenile hormone resulted in a 50% increase in worker body size, as well
as a sharp reduction in total number of progeny reared and a higher proportion of trophic eggs laid by queens
(Helms Cahan *et al.* 2011). This was interpreted as an effect of trophic egg availability or brood/worker ratio on
the nutritional environment. Importantly, in these two studies the consumption of trophic eggs was suggested to
either increase the size of the individuals produced or the likelihood to develop into queens (which are larger than
workers). By contrast, our study reveals that the consumption of trophic eggs reduces the likelihood of developing
into queens.

Our analyses revealed that trophic eggs have a lower content of protein, triglycerides, glycogen, and glucose than
reproductive eggs. A reduced protein content of trophic as compared to reproductive eggs has also been
documented in *Pheidole pallidula* (Lorber and Passera 1981). These findings are in line with the view that trophic
eggs do not simply have a nutritive function as it might then be expected that they should at least contain as much
nutrients as reproductive eggs. Interestingly, our analyses also revealed important differences in RNA and miRNA
content between the two egg types. miRNAs have already been suggested to influence larval caste determination
in the honeybee (Guo *et al.* 2013) with worker jelly being enriched in miRNAs compared to royal jelly (Guo *et al.*
2013; Zhu *et al.* 2017). These studies suggest that it is not the royal jelly that stimulates larval differentiation into
queen, but rather the worker jelly which stimulates the development of larva into worker. Similarly, our study
reveals that compounds found in trophic eggs, perhaps miRNAs, influence larval development towards the worker
phenotype. Interestingly, it has also been recently shown that trophallactic fluid in the ant *Camponotus floridanus*
contains non-digestion related proteins, microRNAs and juvenile hormone (LeBoeuf *et al.* 2016). Moreover,
comparison of trophallactic fluid proteins across social insect species revealed that many are regulators of growth,
development and behavioral maturation (Meurville and LeBoeuf 2021). Finally, a recent study showed that pupae
of several ant species produce secretions that play an important role for early larval nutrition with young larvae
exhibiting stunted growth and decreased survival without access to the fluid (Snir et al. 2022). This raises the

possibility that chemicals delivered in trophic eggs, trophallactic fluids and pupae secretions play previously

unsuspected roles in communication and caste development. Given that some ants do not perform trophallaxis, it

would be interesting to determine whether there are differences in the content of trophic eggs of species performing

trophallaxis and species which do not.

Maternal effects on the process of caste determination have been demonstrated in several social insect species,

including *P. rugosus*, either by queen behaviour or content of the eggs being produced (De Menten *et al.* 2005;

Linksvayer 2006; Schwander *et al.* 2008b; Libbrecht *et al.* 2013; Wei *et al.* 2019). This is, to our knowledge the

first experimental demonstrations that provisioning of trophic eggs influences caste fate. Since only queens

produce trophic eggs in *P. rugosus,* trophic egg provisioning could be the main mechanism underlying the

previously documented maternal effects on the process of caste determination. In species where workers produce

trophic eggs (Supplementary Table 1), the same mechanism could allow workers to influence colony level caste

ratios.

Finally, our analyses also revealed seasonal differences in the proportion of reproductive and trophic eggs, with a

higher ratio of trophic eggs before hibernation than after. In *Pogonomyrmex* the production of new queens occurs

after hibernation (Smith and Tschinkel 2006) or when the queen dies or is removed from the colony (pers. obs).

Thus, new queens are typically produced when there are fewer trophic eggs. Our results predict that under natural

conditions, a decrease in the proportion of trophic eggs should lead to an increase in the larvae developing into

queens. The same logic applies to species where trophic eggs are laid only by the workers in queenright colonies

(Supplementary Table 1). After the queen's death, workers start producing their own male offspring and lay mostly

(if not only) reproductive eggs (*Temnothorax recedens*, Dejean and Passera 1974; *Plagiolepis pygmaea*, Passera

1980; *Myrmecia gulosa*, Dietemann *et al.* 2002), which again leads to a decrease, or cessation, in trophic egg

production. A decrease in trophic egg production and the development of queens were observed simultaneously

in freshly orphaned colonies of *Temnothorax recedens* (Dejean and Passera 1974), *Plagiolepis pygmaea* (Passera

1980) and *Myrmecia gulosa* (Dietemann *et al.* 2002). These examples are consistent with the view that trophic

eggs may also play a role in the process of caste determination in other ant species.

In conclusion, this study provides a new striking example of how females can influence the developmental fate of

their offspring. Because many ants produce trophic eggs, it is possible that this mechanism of parental

manipulation is widespread and play an important role in the general process of caste determination.

Acknowledgments

292   We thank Dr. C. Berney for technical assistance to develop wet lab protocols. We are grateful to S. McGregor and

293   Prof. M. Chapuisat for their helpful comments on the manuscript. This work was supported by an ERC grant and

294   the Swiss NSF (LK).

References

References

Amor, F., P. Ortega, R. Boulay, and X. Cerdá. 2017. Frequent colony orphaning triggers the production of
replacement queens via worker thelytoky in a desert-dwelling ant. *Insectes Soc.* **64**:373–378.
doi.org/10.1007/s00040-017-0556-9

Augustin, J. O., J. F. Santos, and S. L. Elliot. 2011. A behavioral repertoire of *Atta sexdens* (Hymenoptera,
Formicidae) queens during the claustral founding and ergonomic stages. *Insectes Soc.* **58**:197–206.
doi.org/10.1007/s00040-010-0137-7

Azevedo, D. O., J. C. Zanuncio, J. H. C. Delabie, and J. E. Serrão. 2011. Temporal variation of vitellogenin
synthesis in *Ectatomma tuberculatum* (Formicidae: Ectatomminae) workers. *J. Insect Physiol.* **57**:972–977.
doi.org/10.1016/j.jinsphys.2011.04.015

Baroni Urbani, C. B. 1991. Indiscriminate oophagy by ant larvae: an explanation for brood serial organization?
Insectes Soc. **38**:229–239. doi.org/10.1007/BF01314909

Bartels, P. J. 1988. Reproductive caste inhibition by Argentine ant queens: New mechanisms of queen control.
Insectes Soc. **35**:70–81. doi.org/10.1007/BF02224139

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat.*
Softw. **67**:1–48. doi.org/doi:10.18637/jss.v067.i01

Blake, J. A., and P. L. Arnofsky. 1999. Reproduction and larval development of the spioniform Polychaeta with
application to systematics and phylogeny. *Hydrobiologia.* **402**:57-106. doi.org/10.1023/A

Bradford, M. M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing
the principle of protein–dye binding. *Anal. Biochem.* **72**:248–254. doi.org/10.1016/0003-2697(76)90527-3

Brian, M. V, Rigby, C. 1978. The Trophic Eggs of Myrmica Rubra L. *Insect Molecular Biology.* **25**(1):89-110.
CamargoMathias, M., and F. Caetano. 1995. Trophic eggs in workers of *Neoponera villosa* ants
(Hymenoptera:Ponerinae). *J. Adv. Zool.* **16**:62–66.

Cameron, R. C., E. J. Duncan, and P. K. Dearden. 2013. Biased gene expression in early honeybee larval
development. *BMC Genomics.* **14**:1–12. BioMed Central. doi.org/10.1186/1471-2164-14-903

Cassill, D. 2002. Brood care strategies by newly mated monogyne *Solenopsis invicta* (Hymenoptera: Formicidae)
queens during colony founding. *Ann. Entomol. Soc. Am.* **95**:208–212. doi.org/10.1603/0013-
8746(2002)095[0208:BCSBNM]2.0.CO;2

Choe, J. C. 1988. Worker reproduction and social evolution in ants (Hymenoptera: Formicidae). Advances in
myrmecology. 163-187

Collin, R. 2004. Phylogenetic effects, the loss of complex characters, and the evolution of development in
calyptraeid gastropods. *Evolution.* **58**:1488–1502. doi.org/10.1111/j.0014-3820.2004.tb01729.x

Collins, D. H., I. Mohorianu, M. Beckers, V. Moulton, T. Dalmay, and Bourke A. F. 2017. MicroRNAs Associated
with Caste Determination and Differentiation in a Primitively Eusocial Insect. *Sci. Rep.* **7**:1–9. Nature
Publishing Group. doi.org/10.1038/srep45674

Crespi, B. J. 1977. Cannibalism and trophic eggs in subsocial and eusocial insects. Pp. 176–213 in M. A. Elgar
and B. J. Crespi, eds. Cannibalism: Ecology and Evolution Among Diverse Taxa.

Dejean, A., and L. Passera. 1974. Ponte des ouvrières et inhibition royale chez la Fourmi *Temnothorax recedens*
(Nyl.) (Formicidae, Myrmicinae). *Insectes Soc.* **21**:343–355. doi.org/10.1007/BF02331564

Della Lucia, T. M. C., E. F. Vilela, D. D. O. Moreira, J. M. S. Bento, and N. Dos Anjos. 1990. Egg-laying in Atta
sexdens rubropilosa, under laboratory conditions. Pp. 173–179 in R. K. Vander Meer, K. Jaffé, and A.
Cedeno, eds. Applied Myrmecology: A World Perspective. Westview Press, New York.

Dietemann, V., B. Hölldobler, and C. Peeters. 2002. Caste specialization and differentiation in reproductive
potential in the phylogenetically primitive ant *Myrmecia gulosa. Insectes Soc.* **49**:289–298.
doi.org/10.1007/s00040-002-8316-9

Dietemann, V., and C. Peeters. 2000. Queen influence on the shift from trophic to reproductive eggs laid by
workers of the ponerine ant *Pachycondyla apicalis. Insectes Soc.* **47**:223–228.
doi.org/10.1007/PL00001707

Dijkstra, M. B., D. R. Nash, and J. J. Boomsma. 2005. Self-restraint and sterility in workers of *Acromyrmex* and
Atta leafcutter ants. *Insectes Soc.* **52**:67–76. doi.org/10.1007/s00040-004-0775-8

Fletcher, D. J. C., and K. G. Ross. 1985. Regulation of reproduction in Eusocial hymenoptera. *Annu. Rev. Entomol.*
30:319–343.

Freeland, J. 1958. Biological and social patterns in the Australian bulldog ants of the genus *Myrmecia. Aust. J.*
Zool. **6**:1–18. doi.org/https://doi.org/10.1071/ZO9580001

Gibson, G., C. Hart, C. Coulter, and H. Xu. 2012. Nurse eggs form through an active process of apoptosis in the
spionid *Polydora cornuta* (Annelida). *Integr. Comp. Biol.* **52**:151–160. doi.org/10.1093/icb/ics054

Gobin, B., J. Billen, and C. Peeters. 1999. Policing behaviour towards virgin egg layers in a polygynous ponerine
ant. *Anim. Behav.* **58**:1117–1122. doi.org/10.1006/anbe.1999.1245

Gobin, B., and F. Ito. 2000. Queens and major workers of *Acanthomyrmex ferox* redistribute nutrients with trophic
eggs. *Naturwissenschaften* **87**:323–326. doi.org/10.1007/s001140050731

Gobin, B., C. Peeters, and J. Billen. 1998. Production of trophic eggs by virgin workers in the ponerine ant
Gnamptogenys menadensis. Physiol. Entomol. **23**:329–336. doi.org/10.1046/j.1365-3032.1998.00102.x

Guo, X., S. Su, G. Skogerboe, S. Dai, W. Li, Z. Li, F. Liu, R. Ni, Y. Guo, S. Chen, S. Zhang, and R. Chen. 2013.
Recipe for a busy bee: MicroRNAs in honey bee caste determination. *PLoS One* **8**:1–10.
doi.org/10.1371/journal.pone.0081661

Haskins, C.P., and Whelden R.M. 1965. "Queenlessness," Worker Sibship, and Colony Versus Population
Structure in the Formicid Genus Rhytidoponera'. *Psyche: A Journal of Entomology.* **72**(ID 040465): 26
pages. doi.org/10.1155/1965/40465

Heinze, J., S. Cover, and B. Hölldobler. 1995. Neither worker, nor queen: an ant caste specialized in the production
of unfertilized eggs. *Psyche A J. Entomol.* **102**:173–185. doi.org/10.1155/1995/65249

Heinze, J., S. Foitzik, B. Oberstadt, O. Rüppell, and B. Hölldobler. 1999. A female caste specialized for the
production of unfertilized eggs in the ant *Crematogaster smithi. Naturwissenschaften.* **86**:93–95.
doi.org/10.1007/s001140050579

Helms Cahan, S., C. J. Graves, and C. S. Brent. 2011. Intergenerational effect of juvenile hormone on offspring in
Pogonomyrmex harvester ants. *J. Comp. Physiol. B*. **181**:991–999. doi.org/10.1007/s00360-011-0587-x

Hölldobler, B., and N. F. Carlin. 1989. Colony founding, queen control and worker reproduction in the ant
Aphaenogaster (=Novomessor) cockerelli (Hymenoptera: Formicidae). *Psyche (Stuttg).* **96**:131–152.

Hölldobler, B., and E. O. Wilson. 1983. Queen Control in Colonies of Weaver Ants (Hymenoptera: Formicidae).
Ann. Entomol. Soc. Am. **76**:235–238. doi.org/10.1093/aesa/76.2.235

Hora, R. R., C. Poteaux, C. Doums, D. Fresneau, and R. Fénéron. 2007. Egg cannibalism in a facultative
polygynous ant: Conflict for reproduction or strategy to survive? *Ethology.* **113**:909-916.
doi.org/10.1111/j.1439-0310.2007.01391.x

Hung, A. C. F. 1973. Reproductive biology in dulotic ants: preliminary report (Hymenoptera: Formicidae). *Entomol.*
News. **84**:253–259.

Ito, F. 1991. Preliminary report on queenless reproduction in a primitive ponerine ant *Amblyopone sp.* (reclinata
group) in west Java, indonesia. *Psyche (Stuttg).* **98**:319–322.

Ito, F. 1993. Social organization in a primitive ponerine ant: Queenless reproduction, dominance hierarchy and
functional polygyny in *Amblyopone sp*. (reclinata group) (hymenoptera: Formicidae: Ponerinae*). J. Nat. Hist.*
27:1315–1324. doi.org/10.1080/00222939300770751

Ito, F. 2005. Mechanisms regulating functional monogyny in a Japanese population of *Leptothorax acervorum*
(Hymenoptera, Formicidae): Dominance hierarchy and preferential egg cannibalism. *Belgian J. Zool.* **135**:3-
8.

Khila, A., and E. Abouheif. 2008. Reproductive constraint is a developmental mechanism that maintains social
harmony in advanced ant societies. *Proc. Natl. Acad. Sci. U.S.A.* **105**:17884-17889.
doi.org/10.1073/pnas.0807351105

Khila, A., and E. Abouheif. 2010. Evaluating the role of reproductive constraints in ant social evolution. *Philos.*
Trans. R. Soc. B. **365**:617–630. doi.org/10.1098/rstb.2009.0257

Klowden, M. J. 2013. Developmental Systems. Pp. 149–196 in M. J. Klowden, ed. Physiological Systems in Insects
(Third Edition). Academic Press.

Kudo, S., and T. Nakahira. 2004. Effects of Trophic-Eggs on Offspring Performance and Rivalry in a Sub-Social
Bug. *Oikos.* **107**:28–35.

LeBoeuf, A. C., P. Waridel, C. S. Brent, A. N. Gonçalves, L. Menin, D. Ortiz, O. Riba-Grognuz, A. Koto, Z. G.
Soares, E. Privman, E. A. Miska, R. Benton, and L. Keller. 2016. Oral transfer of chemical cues, growth
proteins and hormones in social insects. *Elife.* **5:e20375**:1-27. doi.org/10.7554/eLife.20375

Lee, C. C., H. Nakao, S. P. Tseng, H. W. Hsu, G. L. Lin, J. W. Tay, J. Billen, F. Ito, C. Y. Lee, C. C. Lin, and C. C.
S. Yang. 2017. Worker reproduction of the invasive yellow crazy ant *Anoplolepis gracilipes. Front. Zool.*
14(1):1-12. doi.org/10.1186/s12983-017-0210-4

Levin, L. A., and T. S. Bridges. 1995. Pattern and diversity in reproduction and development. Pp. 1–48 in L.
McEdward, ed. Ecology of marine invertebrate larvae. CRC Press. doi.org/10.1201/9780138758950-1

Libbrecht, R., M. Corona, F. Wende, D. O. Azevedo, J. E. Serrão, and L. Keller. 2013. Interplay between insulin
signaling, juvenile hormone, and vitellogenin regulates maternal effects on polyphenism in ants. *Proc. Natl.*
Acad. Sci. U.S.A. **110**:11050–11055. doi.org/10.1073/pnas.1221781110

Linksvayer, T.A. 2006. Direct, Maternal, and Sibsocial Genetic Effects on Individual and Colony Traits in an Ant.
Evolution. **60**(12): 2552. doi.org/10.1554/06-011.1

López-Ortega, M., and T. Williams. 2018. Natural enemy defense, provisioning and oviposition site selection as
maternal strategies to enhance offspring survival in a sub-social bug. *PLoS One* **13**:1–18.
doi.org/10.1371/journal.pone.0195665

Lorber, B., and L. Passera. 1981. Étude comparative des protéines solubles des œufs de la fourmi *Pheidole*
pallidula Nyl. Bull Intérieur SF-UIEIS 97–99.

Masuko, K. 2003. Larval oophagy in the ant *Amblyopone silvestrii* (Hymenoptera, Formicidae). *Insectes Soc*.
50:317–322. doi.org/10.1007/s00040-003-0688-y

De Menten, L., Fournier, D., Brent, C., Passera, L., Vargo, E.L., and Aron. S. 2005. Dual Mechanism of Queen
Influence over Sex Ratio in the Ant *Pheidole Pallidula. Behavioral Ecology and Sociobiology.* **58**(6): 527-33.
doi.org/10.1007/s00265-005-0964-0.

Meurville, M.-P., and LeBoeuf, A.C. 2021. Trophallaxis: The Functions and Evolution of Social Fluid Exchange in
Ant Colonies (Hymeno Ptera: Formicidae). *Myrmecological News.* **31**:1-20. doi.org/doi:
10.25849/myrmecol.news_031:001 13

Nagano, T., and Fraser, P. 2011. No-Nonsense Functions for Long Noncoding RNAs. *Cell* **145**:178-181.
doi.org/10.1016/J.CELL.2011.03.014

Passera, L. 1978. Une nouvelle catégorie d'œufs alimentaires: les œufs alimentaires émis par les reines vierges
de *Pheidole pallidula* (Nyl.) (Formicidae, Myrmicinae). *Insectes Soc*. **25**:117–126.

Passera, L. 1980. La fonction inhibitrice des reines de la fourmi *Plagiolepis pygmaea* Latr.: Rôle des phéromones.
Insectes Soc. **27**:212–225. doi.org/10.1007/BF02223665

Peeters, C. 2017. Independent colony foundation in *Paraponera clavata* (hymenoptera: Formicidae): First workers
lay trophic eggs to feed queen's larvae. *Sociobiology* **64**:417–422.
doi.org/10.13102/sociobiology.v64i4.2092

Perry, J. C., and B. D. Roitberg. 2006. Trophic egg laying: Hypotheses and tests. *Oikos* **112**:706–714.
doi.org/10.1111/j.0030-1299.2006.14498.x

Romiguier, J., Borowiec, M. L., Weyna, A., Helleu, Q., Loire, E., La Mendola, C., Rabeling, C., Fisher, B. L., Ward,
P. S., Keller, L. 2022. Ant phylogenomics reveals a natural selection hotspot preceding the origin of complex
eusociality. *Current Biology* **32**(13):2942-2947.e4. doi.org/10.1016/j.cub.2022.05.001

RStudio Team. 2015. RStudio: Integrated Development Environment for R. Boston, MA.

Schwander, T., J. Y. Humbert, C. S. Brent, S. H. Cahan, L. Chapuis, E. Renai, and L. Keller. 2008. Maternal Effect
on Female Caste Determination in a Social Insect. *Curr. Biol.* **18**:265–269.
doi.org/10.1016/j.cub.2008.01.024

Smeeton, L. 1981. The Source of Males in *Myrmica Rubra* L. (Hym. Formicidae). *Insectes Sociaux.* **28**(3):263-78.
doi.org/10.1007/BF02223628.

Smith, A. A., B. Hölldobler, and J. Liebig. 2008. Hydrocarbon signals explain the pattern of worker and egg policing
in the ant *Aphaenogaster cockerelli. J. Chem. Ecol.* **34**:1275–1282. doi.org/10.1007/s10886-008-9529-9

Smith, C. R., C. Schoenick, K. E. Anderson, J. Gadau, and A. V Suarez. 2007. Potential and realized reproduction
by different worker castes in queen-less and queen-right colonies of *Pogonomyrmex badius. Insect. Soc.*
54:260-267. doi.org/10.1007/s00040-007-0940-y

Smith, C. R., and W. R. Tschinkel. 2006. The sociometry and sociogenesis of reproduction in the Florida harvester
ant, *Pogonomyrmex badius. J. Insect Sci.* **6**:1–11. doi.org/10.1673/2006_06_32.1

Snir, O., Alwaseem, H., Heissel, S. Sharma, A., Valdés-Rodríguez, S., Carroll, T.S., Jiang, C.S., Razzauti, J.,
Kronauer, D.J.C. 2022. The pupal moulting fluid has evolved social functions in ants. *Nature.*
612(7940):488-494. doi.org/10.1038/s41586-022-05480-9

Sohel, M. H. 2016. Extracellular/Circulating MicroRNAs: Release Mechanisms, Functions and Challenges. *Achiev.*
Life Sci. **10**:175–186. doi.org/10.1016/j.als.2016.11.007

Søvik, E., G. Bloch, and Y. Ben-Shahar. 2015. Function and evolution of microRNAs in eusocial Hymenoptera.
Front. Genet. **6**:1–11. doi.org/10.3389/fgene.2015.00193

Strathmann, M. F., and R. R. Strathmann. 2006. A vermetid gastropod with complex intracapsular cannibalism of
nurse eggs and sibling larvae and a high potential for invasion. *Pacific Sci.* **60**:97–108.
doi.org/10.1353/psc.2005.0062

Suzzoni, J. P., L. Passera, and A. Strambi. 1979. Taux des ecdystéroides et déterminisme des castes de la Fourmi
Pheidole pallidula (Hym. Formicidae).

Taylor, R.W. 1978. *Nothomyrmecia Macrops*: A Living-Fossil Ant Rediscovered. *Science.* **201**(4360):979-85.
doi.org/10.1126/science.201.4360.979

Tennessen, J. aso. M., W. E. Barry, J. Cox, and C. S. Thummel. 2014. Methods for studying metabolism in
Drosophila. Methods **68**:105–115. doi.org/10.1016/j.ymeth.2014.02.034

Torossian, C. 1968. Recherches Sur La Biologie et l'éthologie De *Dolichoderus Quadripunctatus* (L) (Hym. Form.
Dolichoderidæ). *Insectes Sociaux.* **15**(4):375-88

Volny, V. P., M. J. Greene, D. M. Gordon, V. P. Volny, M. J. Greene, and D. M. Gordon. 2006. Brood Production
and Lineage Discrimination in the Red Harvester Ant (*Pogonomyrmex barbatus*). *Ecology.* **87**:2194–2200.

Voss, S. H., J. F. McDonald, and C. H. Keith. 1988. Production and abortive development of fire ant trophic eggs.
Advances in myrmecology. Pp. 517-534. ed. J. C. Trager. New York : E. J. Brill.

Wardlaw, J. C., and G. W. Elmes. 1995. Trophic eggs laid by fertile *Myrmica* queens (Hymenoptera: Formicidae).
Insectes Soc. **42**:303–308. doi.org/10.1007/BF0124042

Wardlaw, J. C., and G. W. Elmes. 1998. Variability in oviposition by workers of six species of *Myrmica*
(hymenoptera, formicidae). *Insectes Soc.* **45**:369–384. doi.org/10.1007/s000400050096

Wei, H., He, X.J., Liao, C.H., Wu, X.B., Jiang, W.J., Zhang, B., Zhou, L.B., Zhang, L.Z., Barron, A.B., Zeng, Z.J.
2019. A Maternal Effect on Queen Production in Honeybees. *Current Biology.* **29**(13):2208-2213.e3.
doi.org/10.1016/j.cub.2019.05.059.

Wheeler, W. M. 1910. Ants: Their Structure, Development, and Behavior. The Columbia University Press, New
York. doi.org/10.1155/1922/80849

Wilson, E. O. 1976. A social ethogram of the neotropical arboreal ant *Zacryptocerus varians* (Fr. Smith). *Anim.*
Behav. **24**:354–363. doi.org/10.1016/S0003-3472(76)80043-7

Yamada, A., F. Ito, R. Hashim, and K. Eguchi. 2018. Queen polymorphism in Acanthomyrmex careoscrobis
Moffett, 1986 in Peninsular Malaysia (Hymenoptera: Formicidae: Myrmicinae), with descriptions of hitherto
unknown female castes and males. *Asian Myrmecology.* **10**(e010009):1-19. doi.org/10.20362/am.010009
Yamauchi, K., T. Furukawa, K. Kinomura, H. Takamine, and K. Tsuji. 1991. Secondary Polygyny by Inbred
Wingless Sexuals in the Dolichoderine Ant *Technomyrmex albipes. Behav. Ecol. Sociobiol.* **29**:313–319.
Zhu, K., M. Liu, Z. Fu, Z. Zhou, Y. Kong, H. Liang, Z. Lin, J. Luo, H. Zheng, P. Wan, J. Zhang, K. Zen, J. Chen, F.
Hu, C. Y. Zhang, J. Ren, and X. Chen. 2017. Plant microRNAs in larval food regulate honeybee caste
development. PLoS Genet. **13**:1–23. doi.org/10.1371/journal.pgen.1006946