Bacterial vaginosis is associated with increased amylolytic activity

Rosanne Hertzberger[1,2], Lisa Himschoot[3], Sylvia Bruisten[4], Leon Steenbergen[1], Warren Lewis[2¥], Piet Cools[3], Remco Kort [1,5 *].

[1] Amsterdam Institute for Life and Environment (A-LIFE), Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[2] Washington University School of Medicine, St Louis, MO, USA

[3] Department of Diagnostic Sciences, Faculty of Medicine and Health Sciences, Ghent University, Belgium

[4] Public Health Laboratory, Department of Infectious diseases, GGD, Amsterdam, The Netherlands

[5] ARTIS-Micropia, Amsterdam, The Netherlands

¥Current address

Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California San Diego, CA, USA.

*corresponding author; E-mail r.kort@vu.nl

Abstract

**Glycogen is the most abundant vaginal carbohydrate in reproductive aged women. Reduced vaginal glycogen is associated with lower levels of *Lactobacillus crispatus*, overgrowth of fastidious anaerobes such as *Gardnerella vaginalis* and increased risk of adverse reproductive and sexual health outcomes. Here we show that *Gardnerella vaginalis*, *Lactobacillus iners* and *Lactobacillus crispatus* can autonomously utilize glycogen as a source for growth. Using an ungelatinized and labeled form of raw amylose, a more degradation-resistant α-1,4-glucan, we were able to discriminate between the alpha-glucosidase activities of common vaginal bacterial species. The bacteria *Lactobacillus crispatus*, *Lactobacillus iners*, *Gardnerella piotii* as well as several other common vaginal species were not capable of raw amylose degradation, while *Gardnerella vaginalis*, *Gardnerella swidsinskii* and *Gardnerella leopoldii* were, with the latter two having the highest degradation rates. In contrast to the glycogen-degrading activity we previously identified in *Lactobacillus crispatus*, this *Gardnerella* alpha-glucosidase activity was not cell-bound and not repressed in the presence of glucose. Raw amylose degradation activity in vaginal swabs was strongly associated with bacterial vaginosis as assessed by Nugent scoring. Overall, our results show that the dysbiotic microbiota of bacterial vaginosis is associated with increased amylolytic activity, which is also found in pure cultures of *Gardnerella* species, but not in other common vaginal bacteria.*

Introduction

The vaginal mucosa of reproductive aged women is rich in glycogen (1, 2). Through shedding and lysis of the superficial vaginal epithelial cells, glycogen is released in the vaginal lumen where it is the primary carbon and energy source for bacteria colonizing the vagina. Glycogen levels were found to strongly depend on the vaginal microbiota composition. The more prevalent *Lactobacillus*-dominated vaginal microbiota is associated with higher levels of glycogen compared to the more diverse *Lactobacillus*-deprived microbiota with higher levels of fastidious anaerobes such as *Gardnerella*, *Fannyhessea* and *Prevotella*. This microbial state is also referred to as bacterial vaginosis (BV) and is a general risk factor for adverse reproductive and sexual health outcomes, such as HIV infection (3), HPV infection (progression) (4), subfertility (5), pelvic inflammatory disease (6), pregnancy loss (7) and preterm birth (8). Therefore, it is important to develop simple, low-cost screening options for bacterial vaginosis.

Gardnerella has been identified as a hallmark genus in bacterial vaginosis. Although this microbial state has a much higher bacterial diversity compared to the *Lactobacillus* dominated microbiota, *Gardnerella* alone can cause many of the (clinical) manifestations of bacterial vaginosis, including exfoliation of the epithelial wall lining the vagina and thinning of the mucus leading to abnormal secretions (9–11).

Alpha-glucosidases are enzymes that catalyze the hydrolysis of α1-4 or α1-6 linkages between glucose units of alpha-glucans. Alpha-glucosidases of vaginal bacteria are hypothesized to take a central role in carbon and energy metabolism by extracellular degradation of host-associated glycogen followed by transport and metabolism of the resulting smaller dextrins. Previously, we have elucidated the genetic basis, regulation and prevalence of one of these alpha-glucosidases: the *L. crispatus* amylopullulanase (12–14). Strikingly, we found a relatively high number of *L. crispatus* strains to be deficient for this amylopullulanase and identified frequently occurring mutations in its gene in the genomes of these strains as well as in

databases of the vaginal metagenome (14, 15). Insight in the ability of other common vaginal

microbes to digest alpha-glucans such as vaginal glycogen remains forthcoming.

Most studies characterizing the alpha-glucosidases of vaginal bacteria have used soluble

starch or glycogen (16–18). Yet, for the gut microbiome it was shown that the structure of the

alpha-glucan has an important role in distinguishing between enzymes of different bacterial

taxa. Animals fed with alpha-glucans, that are more resistant to degradation due to a denser

structure and a relatively lower degree of branching, show an increase of amylolytic species

such as *Bifidobacteria* (19, 20). These bacteria dispose of a type II amylase-pullulanase fusion

protein capable of hydrolyzing resistant starch (21, 22). A heterologously expressed

homologue of this enzyme in *Gardnerella* was found to degrade amylose, pullulan, glycogen

and starch (23). Recent research showed that 14 out of 15 tested *Gardnerella* strains carried

a full copy of this gene (17) and that isolates of four different *Gardnerella* species can transport

and metabolize the small maltodextrins derived from glycogen breakdown (24).

Here we characterized the ability of common vaginal bacteria *L. crispatus*, *L. iners* and

Gardnerella species to degrade a more resistant alpha-glucan: raw amylose. We show that

Gardnerella species can be distinguished from common vaginal lactobacilli by their ability to

break down raw labeled amylose. We further show that the degradation is significantly

associated with vaginal dysbiosis.

Results

83

Gardnerella and *Lactobacillus iners* can use glycogen as the sole carbon source for growth.

84
85

86 The optical density of *L. crispatus*, *L. iners* and *G. vaginalis* after anaerobic growth in NYCIII
87 medium with glucose or glycogen is shown in Figure 1. As expected, and previously shown,
88 the amylopullulanase-deficient strain of *L. crispatus* did not grow on glycogen, whereas the
89 amylopullulanase-sufficient strain grew to a comparable optical density on glycogen as on
90 glucose. The *L. iners* strain showed an equal optical density on glycogen compared to glucose,
91 while the two *G. vaginalis* strains showed a higher optical density on glycogen compared to
92 glucose. No increase in optical density was observed compared to the cell-free control when
93 growing isolates of either species in NYCIII medium without glucose or glycogen (results not
94 shown).

95

96 Figure 1: Optical density after 48 hours of growth in liquid NYCIII medium with 5 g/L glucose (white bars) or 5 g/L
97 glycogen (black bars). All measurements were performed in 3 biological replicates in separate experiments on
98 separate days except for *G. vaginalis* CCUG72423 where data from two biological replicates is used. Data represent
99 average and error bars are the standard deviation. Paired t-test was used to compare averages between glycogen
100 and glucose. * = p < 0,05.

Multi-well assay with undissolved labeled amylose to measure enzymatic rate of alpha-

glucosidases.

Next, we further characterized alpha-glucosidase activity of *L. crispatus*, *L. iners* and

Gardnerella vaginalis using a quantitative amylase assay with raw amylose labelled with a

covalently linked azure cross-linked (AZCL) chromogenic substrate (25) in a microtiter plate.

This assay is conventionally used to study amylase activity using agarose plates or in dissolved

form but is applied here using undissolved granules in a xanthan gum buffer to keep the

substrate dispersed.

The release rate of blue dye is dependent on the enzymatic digestion rate of the labelled glycan

(see a video of the dye diffusion in Supplementary Video S2 and calibration curves in

Supplementary Figure S3). No pretreatment, purification or dialysis of the sample is required,

and the assay is suitable for use in multi-well format in a plate reader as well as for visual

detection.

Figure 2: The AZCL-amylose substrate in xanthan gum buffer with buffer (negative control, left) or after

mixing with saliva (positive control, right). Still image from the video of the enzymatic assay

Supplementary Video S2.

***Gardnerella vaginalis* shows extracellular amylose-degrading activity**

In Figure 3, the enzymatic activity in pellets and supernatants as determined by the amylose

degradation assay is shown. All cultures had a minimal OD600 of 1.0. The amylose

degradation assay showed that the *G. vaginalis* strains were capable of degrading the raw

amylose substrate, while in *L. crispatus* and *L. iners* cultures, no increase of optical density

due to labeled amylose digestion was observed. The activity was mostly found in supernatants

of *G. vaginalis* and to a limited extent associated with the cell pellet. The activity was also

present in *G. vaginalis* cultures grown on medium with glucose, indicating that *G. vaginalis*

alpha-glucosidase activity is not subject to carbon-catabolite repression, as observed for the

glycogen-degrading activity of *L. crispatus*.

Figure 3: AZCL-amylose degradation rate expressed as the increase of optical density at 600 nm per

hour in resuspended cell pellets (white bars) and supernatants (grey bars) from *Lactobacillus crispatus*

RL10 (*pulA*[+]), *Lactobacillus iners* ACS 049-v-Sch2 and *Gardnerella vaginalis* strains CCUG 72422 and

CCUG 72423. Cultures were grown anaerobically in NYCIII medium with glycogen (left panel) or glucose

(right panel) for 48 hours. Bars represent the mean and error bars represent standard deviation of three

independent biological replicates, except for *G. vaginalis* CCUG72423 on glucose (two biological

replicates) and *G. vaginalis* CCUG72422 on glycogen (four biological replicates).

Gardnerella species show variation in amylolytic activity

Figure 4: AZCL-amylose degradation rate by optical density increase at 600 nanometer of cultures (cells in growth

medium) of various human vaginal isolates grown for 48 hours on NYC III medium with glucose. Data points

represent biological replicates that were corrected for a medium control.

AZCL-amylose degradation activity of different _Gardnerella_ species is shown in Figure 4.

Activity varied across _Gardnerella_ species: none of the three _G. piotii_ isolates showed

degradation activity in the assay whereas _G. swidsinskii_ and _G. leopoldii_ showed increased

activity compared to _G. vaginalis_. When all biological replicates of the three _G. vaginalis_ strains

(6 measurements) were compared to all biological replicates of the _G. piotii_ strains taken

together (6 measurements) a significant difference was observed between the degradation of

strains belonging to these species (Mann Whitney $p<0.05$). No amylase activity was observed

in any of the lactobacilli or other vaginal genera such as _Fannyhessea_, _Prevotella_,

Bifidobacterium and _Candida_ (Supplementary Figure 4).

AZCL-amylose degradation rate is associated with Nugent score but not with

abundance of specific _Gardnerella_ species.

The substantial differences in raw amylose degradation activity between three of the most

common vaginal species prompted us to study the presence of this enzymatic activity in vaginal

samples. We analyzed amylose degradation in 96 vaginal samples from a collection of swabs

obtained from women visiting an outpatient Sexually Transmitted Disease clinic in Amsterdam,

as described previously (12). The degradation rate in clinical samples across Nugent

categorization is shown in Figure 5.

Degradation of amylose was strongly associated with Nugent group. Namely, samples from

women with Nugent score between 7-10 (bacterial vaginosis) showed a mean amylose

degradation rate of 0,41 OD600/h +/- 0,27 compared to 0,22 OD600/h +/- 0,18 (mean +/-

standard deviation) in women with Nugent score 0-3 (indicating a *Lactobacillus* dominated

microbiome) (Mann-Whitney, p=0,0001). Women with an intermediate Nugent score (4-6)

showed a mean degradation rate of 0,33 +/- 0,33 OD600/h, which was also significantly lower

compared to women with bacterial vaginosis (*p*-value 0,039).

Figure 5: AZCL-amylose degradation rate of 96 vaginal samples with *Lactobacillus* (LAB)-dominant

microbiome (Nugent score 0-3), intermediate microbiome (Nugent-score 4-6) or bacterial vaginosis

(Nugent score 7-10). *p*-values were calculated by Mann-Whitney test and displayed as * *p* < 0.05 and

*** *p* < 0.0005. Horizontal bars indicate the mean.

In growth experiments, a higher amylose degradation rate was observed in cultures of

Gardnerella species compared to *Lactobacillus* species. Therefore, we hypothesized that the

observed difference in samples from women with high Nugent scores (bacterial vaginosis)

compared to women with low Nugent scores may be due to higher *Gardnerella* abundance in

the high Nugent score group. We therefore assessed whether the measured amylose

degradation rate was associated with *Gardnerella* abundance and found a positive correlation

between AZCL-amylose degradation rate and levels of *Gardnerella* and *Fannyhessea* (r of

0,24 and 0,23, *p*-value of 0,016 and 0,028, respectively) that was nevertheless non-significant

after Bonferroni correction for multiple testing (4 tests). Since *Gardnerella* showed strong

interspecies differences in amylolytic activity *in vitro*, we continued to quantify the various

Gardnerella species using species-specific qPCR. Given the high amylose degradation rate of

G. leopoldii and *G. swidsinskii* strains and the low rate in strains of *G. piotii* (Figure 4), we

expected to find a high abundance of *G. leopoldii* and *G. swidsinskii* in vaginal samples with

high Nugent scores and high AZCL-amylose degradation rate, and samples with high Nugent

score but low degradation rates to have less *G. leopoldii* and *swidsinskii*. However, none of

the *Gardnerella* spp. concentrations showed a significant correlation with AZCL-amylose

degradation rate of the vaginal samples, using a *p*-value threshold that was corrected using

Bonferroni for multiple testing (Table 1).

Table 1: correlation between raw AZCL-amylose degradation rate and bacterial abundance in vaginal

samples measured by two different qPCR protocols. Spearman's rank correlation coefficient (R_s) for

log10 transformed genome equivalents and AZCL-amylose degradation rate with approximate *p*-value

(*p*-value of < 0,013 are considered significant after Bonferroni correction for multiple testing).

Species	Assay	R_s (*p*-value)
Gardnerella	BV diagnostic PCR	0.245 (0.016)
Fannyhessea	BV diagnostic PCR	0.224 (0.028)
Lactobacillus	BV diagnostic PCR	-0.035 (0.734)
Bacteria	BV diagnostic PCR	0.211 (0.084)
Gardnerella leopoldii	*Gardnerella* species-specific qPCR	0.084 (0.417)
Gardnerella piotii	*Gardnerella* species-specific qPCR	0.096 (0.352)
Gardnerella swidsinskii	*Gardnerella* species-specific qPCR	0.192 (0.060)
Gardnerella vaginalis	*Gardnerella* species-specific qPCR	0.238 (0.020)

 Discussion

In this study we set out to characterize extracellular alpha-glucosidases of key member species of the vaginal microbiota. These enzymes catalyze the degradation of α1-4 and/or α1-6 linkages between glucose units of alpha-glucans such as amylopectin, glycogen and amylose. The female reproductive tract between puberty and menopause is rich in glycogen (26). Vaginal bacteria that express and secrete glycogen-degrading enzymes gain access to these carbon and energy stores in a general nutrient-poor environment.

Most *L. crispatus* strains express an amylopullulanase that is capable of degrading starch and allows growth on glycogen as a single carbon source (13–15). Despite the ability to degrade starch, here we showed that *L. crispatus* is unable to break down a more degradation resistant alpha-glucan: raw amylose. Amylose consists of α1-4 linked glucose units and has a denser structure compared to α-glucans such as amylopectin, pullulan and glycogen due to the absence of branches (that are introduced by α1-6 linkages). Vaginal lactobacilli as well as other common vaginal species, such as *Fannyhessea vaginae*, *Prevotella bivia* and *Candida albicans* were all unable to degrade raw labeled amylose. Furthermore, we saw profound differences in the raw amylolytic activity of various *Gardnerella* species, which went previously undetected using glycogen as a substrate (17).

These findings are contrasting previously published results on vaginal bacterial alpha-glucosidases. A characterization of a variety of amylopullulanases from vaginal species showed that alpha-glucosidases of *L. crispatus, L. iners, G. vaginalis* and *Prevotella bivia* were capable of amylose degradation in addition to glycogen and pullulan, when heterologously expressed in *Escherichia coli* (23). The difference may stem from the type of amylose used. Woolston *et al.* used amylose dissolved in DMSO which may alter the crystal structure and therefore the physical accessibility, whereas in our study we have used ungelatinized labeled amylose kernels. A second difference may be caused by the cleavage of the signal peptide in their study and the heterologous expression in *E. coli* which may lead to posttranscriptional modification of these enzymes.

Previously, resistant starches (such as high amylose corn starch) were found to modulate the

gut microbiome. High amylose corn starch is degraded by specific gut genera, one of which is

Bifidobacterium that is closely related to the genus *Gardnerella*. The amylopullulanase of

bifidobacteria has the same carbohydrate-binding, amylase and pullulanase domains, N-

terminal signal peptide and C-terminal transmembrane domain as the *Gardnerella* gene that

is likely responsible for the observed raw amylose degradation activity (17, 23).

Using raw amylose we found strong differences in amylolytic activity between *Gardnerella*

species and other vaginal genera. Besides substrate degradation we also found differences

between the regulation. Alpha-glucosidase activity of *G. vaginalis* was in this study also

observed in cultures grown on glucose whereas we previously found *L. crispatus*

amylopullulanase to be repressed in the presence of the products of glycogen degradation

such as glucose, maltose or maltotriose (14).

Apart from the absence of catabolite repression, and the ability to degrade raw amylose, we

identified a third characteristic that was found to be distinctive: unlike to what we previously

found for *L. crispatus* (14), the *Gardnerella* amylolytic activity was mostly found in the

extracellular environment and only to a minor extent cell-bound. This is not consistent with the

presence of a C-terminal transmembrane domain in the candidate amylopullulanase gene. We

do consider that this could be the result of the linkages in our chemically inaccessible form of

the labeled substrate and that there may be in fact some cell-associated activity that is

incapable of accessing the linkages embedded in the granules. However, this phenomenon

has been described previously for the *Gardnerella* sialidase gene *nanH*. The *nanH* gene also

has a C-terminal transmembrane domain, yet the activity of the gene product is exclusively

found in spent medium (27). Further experiments are required to understand the exact location

of these enzymes and potential cleavage of the transmembrane domain.

Although the association between AZCL-amylose degradation rate and Nugent categorization

was high, we did not find evidence that the association was related to the abundance of any

specific *Gardnerella* species. For example, samples that had high abundance of *G. swidsinskii*

were equally likely to show high or low amylose degradation activity compared to samples that

were high in *G. piotii* even though the former *Gardnerella* species had much higher activity in

culture compared to the latter. Some samples with no detectable *G. swidsinskii* or *G. leopoldii*

were amongst the top 10% of samples in terms of amylase degradation.

Many have speculated that there is a significant role of the host in vaginal glycogen hydrolysis

and demonstrated the presence of human amylase in vaginal samples with different microbial

community states (16, 18, 28). Apart from microbial amylases, human amylase may contribute

to the observed amylose degradation activity in the vagina. Likely, human amylase activity, if

any, is not present at biologically relevant levels in *Lactobacillus*-dominated vaginal

environments, given the very low levels of AZCL-amylose breakdown in most of the samples

with low Nugent scores (0-3). We cannot exclude the presence of human amylase activity in

samples with higher Nugent scores. Since *in vitro* we did not observe any raw amylose

degradation in several other vaginal bacterial strains commonly associated with bacterial

vaginosis, such as *Fannyhessae, Prevotella* or vaginal *Bifidobacteria* the origin of this activity

remains elusive.

To further identify the contribution of *Gardnerella* in glycogen degradation in the vaginal

environment it is now crucial to analyze the genetic basis for this activity. We especially

focused on the amylopullulanase gene that was confirmed to encode a 200 kDa pullulanase-

amylase enzyme by heterologous expression in two previous studies (Woolston and

Bhandari). The amylopullulanase gene sequence of the strains used in this study show

extensive genetic variation but no apparent genetic disruptions coinciding with the (in)ability of

a *Gardnerella* strain to degrade raw amylose (not shown). One of the *G. piotii* strains (UGent

18.01) had a full copy of the amylopullulanase gene whereas another strain did not (UGent

21.28). Both strains were not capable of AZCL-amylose degradation. *G. leopoldii* and *G.*

swidsinskii strains had a smaller gene, confirmed to be an amylase (17), upstream of the

amylopullulanase gene which may explain the higher amylase activity observed in these

strains. However, we also detected one *G. leopoldii* and one *G. swidsinskii* strain had

inconsistent high amylose activity. These strains were thus not described here due to high

variability in degradation rates between experiments. Future experiments should focus on the

identification, expression and regulation of the genetic basis of the *Gardnerella* amylase

activity observed in this study, as well as the factors that impact its activity in different strains

to elucidate the underlying mechanism.

Taken together, the characteristics of the *Gardnerella* amyloglucosidase could explain the low

glycogen levels in samples with bacterial vaginosis (26). Lactobacilli may in contrast only

'graze' the accessible linkages of the glycogen molecules and cease amylopullulanase

expression as soon as the smaller maltodextrins start to accumulate (14). Overgrowth of

Gardnerella typically seen in BV in combination with the metabolic differences may add to the

reduced glycogen concentrations in BV.

Lastly, we consider that the secreted nature of the glycogen-degrading activity may mean that

these enzymes do not remain confined to the vaginal space but may travel to the tissues of

the higher reproductive tract. The enzymes could hypothetically migrate through the cervix to

access the endometrium lining of the uterus. Glycogen stores of the endometrium and fallopian

tubes function as an early carbohydrate source for the implanted blastocyst in many animals

(29). *Gardnerella* enzymes degrading endometrial glycogen may impact the chances of

successful implantation.

Previously, *Gardnerella* sialidase and cytolysin were found to be linked to clinical

characteristics of bacterial vaginosis, namely mucus thinning and epithelial exfoliation (9, 11).

Here we consider *Gardnerella*'s glycogen metabolism in the vagina as a third factor that links

colonization and outgrowth of this specific bacterium to a BV-characteristic: low luminal

glycogen levels. Furthermore, glycogen hydrolysis in addition with epithelial exfoliation and

mucus degradation may explain the extensive mucosal barrier degradation that is associated

with *Gardnerella* colonization.

Acknowledgments

VU Amsterdam: Alexander Woudstra, Ritesh Panchoe for experiments setting up anaerobic cultivation. Jurgen Haanstra for supervision of students. Douwe Molenaar for help with data analysis and statistics. Martin Braster for help with the anaerobic chamber. Sara Morselli for help with additional calibration curves.

GGD Amsterdam: Mirjam van Dierdorp for isolating the DNA from vaginal samples.

Ghent University: Mario Vaneechoutte for providing different vaginal strains. Leen van Simaey and Aliona Rosca for their assistance in AZCL-amylose assays of different member species of the vaginal microbiota.

Megazyme kindly donated 5 gram of fine AZCL-amylose for this study.

Author contributions

Conceptualization, R.H., R.K. and W.L.; Methodology and data analysis, R.H., L. H., L. van S., W.L., R.K., P.C., S.B.; Experiments: R.H., L. H., L. van S.; Data curation: R.H., L.H., L. van S.; Writing—original draft preparation, R.H.; Writing review and editing, R.H., L.H., S.B., P.C., R.K.; Supervision: S.B., P.C., R.K. All authors have read and agreed to the published version of the manuscript.

Methods

Bacterial cultivation

For this study we selected various vaginal strains isolated from vaginal samples. The isolation and characterization of the *Lactobacillus crispatus* amylopullulanase-sufficient strain RL10 and amylopullulanase-deficient strain RL09 were previously described (12–14). *Lactobacillus iners* ACS 049-v-Sch2 was isolated from a vaginal sample and identified using MALDI-TOF. Various *Gardnerella* strains whose identification and genome sequences were previously published (30) were included as well as several other members of the vaginal ecology who were also identified using MALDI-TOF. An overview of all strains can be found in Supplementary Table S1.

Cultivation of bacteria was carried out in two laboratories (Ghent University and Vrije Universiteit (VU) Amsterdam) with different equipment. The results from the VU Amsterdam laboratory are depicted in Figure 1, 2, 3, 5, S1, S2 and S3. The results from experiments by Ghent University are depicted in Figure 4 and S4. All strains were anaerobically cultivated from -80 °C glycerol stocks (VU Amsterdam: NYCIII medium + 20% glycerol, Ghent: TSB + 15% glycerol) on agar plates under conditions indicated in Table S1. From these plates colonies were picked to inoculate liquid New York City (NYC)-III broth with 10% horse serum as described previously (14) or YEPG broth for *Candida albicans*. All cultures were handled under anaerobic conditions (VU Amsterdam: N_2 + 10% CO_2, Ghent University: 10% H_2, 10% CO_2, 80% N_2) in an anaerobic chamber (VU Amsterdam: PLAS Labs 855 NB; Ghent university: Bugbox Plus). For the three most common vaginal bacterial species -*Gardnerella vaginalis, Lactobacillus iners* and *Lactobacillus crispatus*- we studied growth on glycogen by comparing optical density after anaerobic incubation in NYCIII medium with glucose or glycogen. To verify whether any other medium components resulted in growth we inoculated these species also in NYCIII medium without glucose or glycogen supplemented.

In Amsterdam the optical density of cultures after 48 hours of growth was determined by

diluting in saline and measuring the turbidity at 600 nm with a 1 mL cuvette using a UV-VIS

spectrophotometer (Cary 50 Bio, Varian), subtracting the cell-free medium control. In Ghent

the optical density was measured by diluting the culture in phosphate-buffered saline in a 96-

well microtiter plate and measuring the turbidity in a plate reader at 600 nm (GloMax® Explorer

Multimode Microplate Reader, Promega).

AZCL-amylose assay

AZCL-amylose was used in a raw ungelatinized form and suspended in an amylase buffer with

xanthan gum to disperse the granules. After mixing with the sample (either vaginal sample of

bacterial culture, spent medium or resuspended cell pellet) the increase in optical density in a

microtiter plate was measured at 600 nm. The lack of heating or other pretreatment results in

the substrate retaining a coarse, granular, undissolved structure. Pelleting is prevented in this

assay by suspension in a viscous buffer containing xanthan gum (see Supplementary Figure

S2). Xanthan gum (Sigma Life Sciences) was dissolved at 5 mg/mL in amylase buffer (100

mM sodium acetate buffer containing 5 mM $CaCl_2$, pH adjusted with HCl to 5,3). This solution

was autoclaved at 121 °C for 20 minutes and kept at room temperature. AZCL-amylose (O-

AZAMYF) is commercially available in two forms: regular and fine. The granules in the 'fine'

product are small enough to pipet with regular pipet tips to dose and mix with the sample. This

solution was stable without sedimentation of the raw amylose granules for up to 48 hours.

AZCL-amylose was dispersed in the xanthan gum/amylase buffer prior to each experiment by

shortly vortexing.

The AZCL-amylose assay was also carried out at both Ghent University and VU Amsterdam,

with different equipment. In both protocols the assay was carried out in a flat bottom 96-well

plate covered with adhesive film. A total of 190 μL reagent solution was added to 10 μL sample

and mixed by carefully pipetting up and down. The plate was incubated in a plate reader

(Amsterdam: Multiskan microplate photometer, Thermo Fisher; Ghent University: GloMax®

Explorer Multimode Microplate Reader, Promega) at 37 °C and absorption was measured at

600 nm. Maximum rate was taken as the maximum slope of a three-hour time frame (including 19 datapoints). The enzyme assay was validated using *Aspergillus oryzae* amylase (Sigma-Aldrich) solution in amylase buffer with 150 units/μL where one unit corresponds to the amount of enzyme needed to cleave 1 μmol of maltose per minute at pH 6.0 and 25 °C.

AZCL-amylose assay video

The video of the assay shown in Supplementary materials S2 was made by prewarming the substrate suspension and microtiter plate to 40 °C after which 10 μL of saliva or amylase buffer (negative control) was added and filmed at room temperature on the bench using a Samsung smartphone. The footage was clipped to include only the two relevant wells and accelerated 50 times.

AZCL-amylose assay on bacterial cultures

To study the prevalence of raw amylose degradation activity amongst *Gardnerella* species, we screened a set of 14 human vaginal *Gardnerella* strains, including *Gardnerella leopoldii, G. swidsinskii, G. vaginalis* and *G. piotii* and *Gardnerella* genomospecies 3. We compared these to isolates of other common species colonizing the human vagina including *Candida albicans*, *Fannyhessea vaginae* (previously known as *Atopobium vaginae*), *Prevotella bivia,* *Bifidobacterium* and several *Lactobacillus* species.

For this enzymatic assay with bacterial cultures we only included cultures with sufficient growth with a minimal OD600 of 1,0 (VU Amsterdam) or 0,2 (Ghent University). Cultures were either used without centrifugation or centrifuged for 1 minute at 20,000 rcf at room temperature to separate cells. Spent medium was aspirated and mixed with the substrate and cell pellets were resuspended in sterile phosphate-buffered saline and mixed with the substrate. In experiments with full cultures (shown in figure 4 and S4) a correction with a NYCIII medium control was included. For one biological replicate this medium control was not available so instead the *L. iners* measurement was taken as the baseline since this species was shown in all experiments to have no activity compared to the medium control.

In one experiment the optical density and AZCL-amylose degradation rates of *Gardnerella*

cultures on NYCIII with glucose were omitted because of lack of sufficient growth. One other

biological replicate was omitted (optical densities as well as AZCL-degradation rate) since the

optical densities were more than twice the usual optical densities. We ruled out contamination

but the reason of the unusually high optical density remained unexplained and was not

observed afterwards. Therefore, we decided to leave the data out.

One *G. leopoldii* (4042406) and one *G. swidsinskii* strain (GS10234) showed highly discrepant

results between different experiments and were omitted for this reason.

Vaginal samples

Vaginal swabs were collected from patients visiting the outpatient clinic for Sexually

Transmitted Disease of the Public Health Service in Amsterdam (The Netherlands) as

described previously (12). Ethics approval was obtained by the AMC ethics review board

(document reference number W12_086 # 12.17.0104). The procedure for consent to

participate for use of vaginal swabs for scientific research has been approved (reference

number W15_159 # 15.0193). The swabs were eluted in Amies buffer and 15% glycerol and

cysteine were added for cryoprotection to preserve live bacteria and enzymes for subsequent

isolation and analysis. The samples were stored at -80 °C.

Samples for this study were selected based on Nugent scores choosing one third of samples

with scores between 0-3 (*Lactobacillus* dominated), one third with scores between 4-6

(intermediate) and one third with a Nugent score between 7-10 (bacterial vaginosis).

AZCL-amylose assay on vaginal samples

For the amylose assay samples were thawed, vortexed and pipetted but not homogenized or

centrifuged, which means that mucus and other solid material was still present in the sample

when pipetted. 10 µL of the liquid fraction was transferred to a microtiter plate to which 190 µL

of the AZCL-amylose in the amylase buffer with xanthan gum was added. The assay and rate

calculation was carried out as described above.

DNA extraction and bacterial quantification in vaginal samples

The commercially available multiplex PCR for BV diagnosis (ATRiDa, The Netherlands) was performed according to instructions of the manufacturer (31, 32) on isolated DNA from vaginal samples targeting total *Lactobacillus*, total *Gardnerella*, total *Fannyhessea* and total bacterial load. Both the DNA extraction as well as the qPCR reaction were performed as described previously (33). This same extracted DNA was utilized for *G. vaginalis*, *G. piotii*, *G. leopoldii* and *G. swidsinskii* specific qPCR assays (34). To account for the multiple testing, we adjusted the *p*-value threshold by Bonferroni-correction leading to a required *p*-value of 0,013 (p=,05 divided by 4 outcomes) for the BV diagnostic PCR and the *Gardnerella* species-specific qPCR (also 4 outcomes). As expected, *Gardnerella, Fannyhessea* and total bacterial abundance were strongly associated with Nugent score, and these species had a higher abundance in samples with Nugent score 7-10 compared to samples with a Nugent score of 0-3, while lactobacilli were more abundant in women with low Nugent scores compared to women with high Nugent scores. All correlations between the genera and Nugent score group are shown in Table S3.

We used the log transformed data to find correlations between AZCL-amylose degradation rate and abundance of any of the targets. All data processing and statistical analysis were performed in Microsoft Excel and GraphPad Prism 9.5.0.

References

1.　Cruickshank, R.1934. The Conversion of the Glycogen of the Vagina Into Lactic Acid. J.
Pathol. Bacteriol. 39: 213–219.

2.　Mirmonsef P, Hotton AL, Gilbert D, Gioia CJ, Maric D, Hope TJ, Landay AL, Spear GT.
2016. Glycogen Levels in Undiluted Genital Fluid and Their Relationship to Vaginal pH,
Estrogen, and Progesterone. PloS One 11:e0153553.

3.　Gosmann C, Anahtar MN, Handley SA, Farcasanu M, Abu-Ali G, Bowman BA,
Padavattan N, Desai C, Droit L, Moodley A, Dong M, Chen Y, Ismail N, Ndung'u T,
Ghebremichael MS, Wesemann DR, Mitchell C, Dong KL, Huttenhower C, Walker BD,
Virgin HW, Kwon DS. 2017. Lactobacillus-Deficient Cervicovaginal Bacterial
Communities Are Associated with Increased HIV Acquisition in Young South African
Women. Immunity 46:29–37.

4.　Lin D, Kouzy R, Abi Jaoude J, Noticewala SS, Delgado Medrano AY, Klopp AH,
Taniguchi CM, Colbert LE. 2020. Microbiome factors in HPV-driven carcinogenesis and
cancers. PLoS Pathog 16:e1008524.

5.　Vergaro P, Tiscornia G, Barragán M, García D, Rodriguez A, Santaló J, Vassena R.
2019. Vaginal microbiota profile at the time of embryo transfer does not affect live birth
rate in IVF cycles with donated oocytes. Reprod Biomed Online 38:883–891.

6.　Haggerty CL, Hillier SL, Bass DC, Ness RB, PID Evaluation and Clinical Health study
investigators. 2004. Bacterial vaginosis and anaerobic bacteria are associated with
endometritis. Clin Infect Dis Off Publ Infect Dis Soc Am 39:990–995.

7.　Brown RG, Al-Memar M, Marchesi JR, Lee YS, Smith A, Chan D, Lewis H, Kindinger L,
Terzidou V, Bourne T, Bennett PR, MacIntyre DA. 2019. Establishment of vaginal
microbiota composition in early pregnancy and its association with subsequent preterm
prelabor rupture of the fetal membranes. Transl Res. 207:30-43.

8.　Gudnadottir U, Debelius JW, Du J, Hugerth LW, Danielsson H, Schuppe-Koistinen I,
Fransson E, Brusselaers N. 2022. The vaginal microbiome and the risk of preterm birth:
a systematic review and network meta-analysis. Sci Rep 12:7926.

9.　Gilbert NM, Lewis WG, Lewis AL. 2013. Clinical features of bacterial vaginosis in a
murine model of vaginal infection with *Gardnerella vaginalis*. PloS One 8:e59539.

10.　Gelber SE, Aguilar JL, Lewis KLT, Ratner AJ. 2008. Functional and phylogenetic
characterization of Vaginolysin, the human-specific cytolysin from *Gardnerella vaginalis*.
J Bacteriol 190:3896–3903.

11.　Lewis WG, Robinson LS, Gilbert NM, Perry JC, Lewis AL. 2013. Degradation, foraging,
and depletion of mucus sialoglycans by the vagina-adapted Actinobacterium *Gardnerella*
vaginalis. J Biol Chem 288:12067–12079.

12.　Dols JAM, Molenaar D, van der Helm JJ, Caspers MPM, de Kat Angelino-Bart A,
Schuren FHJ, Speksnijder AGCL, Westerhoff HV, Richardus JH, Boon ME, Reid G, de
Vries HJC, Kort R. 2016. Molecular assessment of bacterial vaginosis by *Lactobacillus*
abundance and species diversity. BMC Infect Dis. 16:180.

13.　van der Veer C, Hertzberger RY, Bruisten SM, Tytgat HLP, Swanenburg J, de Kat
Angelino-Bart A, Schuren F, Molenaar D, Reid G, de Vries H, Kort R. 2019. Comparative

genomics of human *Lactobacillus crispatus* isolates reveals genes for glycosylation and
glycogen degradation: implications for in vivo dominance of the vaginal microbiota.
Microbiome 7:49.

14. Hertzberger R, May A, Kramer G, van Vondelen I, Molenaar D, Kort R. 2022. Genetic
Elements Orchestrating *Lactobacillus crispatus* Glycogen Metabolism in the Vagina. 10.
Int J Mol Sci 23:5590.

15. Sycuro L, Lithgow K, Cochinamogulos A, Muirhead K, Konschuh S, Oluoch L, Mugo N,
Roxby A. 2023. Glycogen-Degrading pullulanase a is variably present and correlated
with lactic acid in vaginal samples from young african women. Am J Obstet Gynecol
228:S788.

16. Nunn Kenetta L., Clair Geremy C., Adkins Joshua N., Engbrecht Kristin, Fillmore
Thomas, Forney Larry J., Young Vincent B. 2020. Amylases in the Human Vagina.
mSphere 5:e00943-20.

17. Bhandari P, Tingley J, Abbott DW, Hill JE. 2022. Glycogen degrading activities of
catalytic domains of α-amylase and α-amylase-pullulanase enzymes conserved in
Gardnerella spp. from the vaginal microbiome. bioRxiv10.19.512974

18. Spear GT, French AL, Gilbert D, Zariffard MR, Mirmonsef P, Sullivan TH, Spear WW,
Landay A, Micci S, Lee BH, Hamaker BR. 2014. Human alpha-amylase present in lower-
genital-tract mucosal fluid processes glycogen to support vaginal colonization by
Lactobacillus. J Infect Dis 210:1019–1028.

19. Brown I, Warhurst M, Arcot J, Playne M, Illman RJ, Topping DL. 1997. Fecal Numbers of
Bifidobacteria Are Higher in Pigs Fed *Bifidobacterium longum* with a High Amylose
Cornstarch Than with a Low Amylose Cornstarch. J Nutr 127:1822–1827.

20. Sybille T, June Z, Michael K, Roy M, Maria L. M. 2013. The intestinal microbiota in aged
mice is modulated by dietary resistant starch and correlated with improvements in host
responses. FEMS Microbiol Ecol 83:299–309.

21. Jung D-H, Seo D-H, Kim Y-J, Chung W-H, Nam Y-D, Park C-S. 2020. The presence of
resistant starch-degrading amylases in *Bifidobacterium adolescentis* of the human gut.
Int J Biol Macromol 161:389–397.

22. Maier TV, Lucio M, Lee LH, VerBerkmoes NC, Brislawn CJ, Bernhardt J, Lamendella R,
McDermott JE, Bergeron N, Heinzmann SS, Morton JT, González A, Ackermann G,
Knight R, Riedel K, Krauss RM, Schmitt-Kopplin P, Jansson JK. 2017. Impact of Dietary
Resistant Starch on the Human Gut Microbiome, Metaproteome, and Metabolome. mBio
8:e01343-17.

23. Woolston BM, Jenkins DJ, Hood-Pishchany MI, Nahoum SR, Balskus EP. 2021.
Characterization of vaginal microbial enzymes identifies amylopullulanases that support
growth of *Lactobacillus crispatus* on glycogen. bioRxiv 07.19.452977

24. Bhandari P, Hill JE. 2022. Transport and utilization of glycogen breakdown products by
Gardnerella spp. from the human vaginal microbiome. bioRxiv 11.01.514706

25. McCleary BV. 1980. New chromogenic substrates for the assay of alpha-amylase and
(1→4)-β-d-glucanase. Carbohydr Res 86:97–104.

523 26. Mirmonsef P, Hotton AL, Gilbert D, Burgad D, Landay A, Weber KM, Cohen M, Ravel J,
524    Spear GT. 2014. Free glycogen in vaginal fluids is associated with *Lactobacillus*
525    colonization and low vaginal pH. PloS One 9:e102467.

526 27. Robinson LS, Schwebke J, Lewis WG, Lewis AL. 2019. Identification and
527    characterization of NanH2 and NanH3, enzymes responsible for sialidase activity in the
528    vaginal bacterium *Gardnerella vaginalis*. J Biol Chem 294:5230–5245.

529 28. Nasioudis D, Beghini J, Bongiovanni AM, Giraldo PC, Linhares IM, Witkin SS. 2015.
530    alpha-Amylase in Vaginal Fluid: Association With Conditions Favorable to Dominance of
531    *Lactobacillus*. Reprod Sci 22:1393-8.

532 29. Dean M. 2019. Glycogen in the uterus and fallopian tubes is an important source of
533    glucose during early pregnancydagger. Biol Reprod 101:297–305.

534 30. Vaneechoutte M, Guschin A, Van Simaey L, Gansemans Y, Van Nieuwerburgh F, Cools
535    P 2019. Emended description of *Gardnerella vaginalis* and description of *Gardnerella*
536    *leopoldii* sp. nov., *Gardnerella piotii* sp. nov. and *Gardnerella swidsinskii* sp. nov., with
537    delineation of 13 genomic species within the genus *Gardnerella*. Int J Syst Evol Microbiol
538    69:679–687.

539 31. Shipitsyna E, Khusnutdinova T, Budilovskaya O, Krysanova A, Shalepo K, Savicheva A,
540    Unemo M. 2020. Bacterial vaginosis-associated vaginal microbiota is an age-
541    independent risk factor for *Chlamydia trachomatis*, *Mycoplasma genitalium* and
542    *Trichomonas vaginalis* infections in low-risk women, St. Petersburg, Russia. Eur J Clin
543    Microbiol Infect Dis. 39:1221-1230.

544 32. Rumyantseva T, Shipitsyna E, Guschin A, Unemo M. 2016. Evaluation and subsequent
545    optimizations of the quantitative AmpliSens Florocenosis/Bacterial vaginosis-FRT
546    multiplex real-time PCR assay for diagnosis of bacterial vaginosis. APMIS Acta Pathol
547    Microbiol Immunol Scand 124:1099–1108.

548 33. van der Veer C, van Houdt R, van Dam A, de Vries H, Bruisten S. 2018. Accuracy of a
549    commercial multiplex PCR for the diagnosis of bacterial vaginosis. J Med Microbiol
550    67:1265-1270.

551 34. Latka A, Van Simaey L, Reynders M, Cools P, Rogier T, Lebbe B, Corsini L, Landlinger
552    C, Vaneechoutte M. 2022. Optimization of propidium monoazide qPCR (viability-qPCR)
553    to quantify the killing by the *Gardnerella*-Specific endolysin PM-477, directly in vaginal
554    samples from women with bacterial vaginosis. Antibiotics 11:111.

555