Identification of levoglucosan degradation pathways in bacteria and sequence similarity

network analysis

Arashdeep Kaur,[1,2] Nichollas E. Scott,[3] Marion Herisse,[3] Ethan D. Goddard-Borger,[4,5] Sacha

Pidot,[3] Spencer J. Williams*[1,2]

[1] School of Chemistry, University of Melbourne, Parkville, Victoria 3010, Australia.

[2] Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville,

Victoria 3010, Australia

[3] Department of Microbiology and Immunology, University of Melbourne, at the Peter

Doherty Institute for Infection and Immunity, Victoria, 3000, Australia

[4] Department of Medical Biology, University of Melbourne, Parkville, Victoria 3010,

Australia

[5] ACRF Chemical Biology Division, The Walter and Eliza Hall Institute of Medical

Research, Parkville, Victoria 3010, Australia

E-mail: sjwill@unimelb.edu.au; ORCID: 0000-0001-6341-4364

Keywords: anhydrosugar; carbohydrate; metabolism; sequence similarity network analysis

Abstract

Levoglucosan is produced in the pyrolysis of cellulose and starch, including from bushfires or

the burning of biofuels, and is deposited from the atmosphere across the surface of the earth.

We describe two levoglucosan degrading *Paenarthrobacter* spp. (*Paenarthrobacter*

nitrojuajacolis LG01 and *Paenarthrobacter histidinolovorans* LG02) that were isolated by

metabolic enrichment on levoglucosan as sole carbon source. Genome sequencing and

proteomics analysis revealed expression of a series of gene clusters encoding known

levoglucosan degrading enzymes, levoglucosan dehydrogenase (LGDH, LgdA), 3-keto-

levoglucosan β-eliminase (LgdB1) and glucose 3-dehydrogenase (LgdC), along with an ABC

transporter cassette and associated solute binding protein. However, no homologues of 3-

ketoglucose dehydratase (LgdB2) were evident, while the expressed gene clusters contained a

range of putative sugar phosphate isomerase/xylose isomerases with weak similarity to LgdB2.

Sequence similarity network analysis of genome neighbors of LgdA revealed that homologues

of LgdB1 and LgdC are generally conserved in a range of bacteria in the phyla Firmicutes,

Actinobacteria and Proteobacteria. One sugar phosphate isomerase/xylose isomerase cluster

(LgdB3) was identified with limited distribution mutually exclusive with LgdB2 that may fulfil

a similar function. LgdB1, LgdB2 and LgdB3 adopt similar predicted 3D folds suggesting

overlapping function in processing intermediates in LG metabolism. Our findings highlight the

diversity within the LGDH pathway through which bacteria utilize levoglucosan as a nutrient

source.

Introduction

Levoglucosan (LG; 1,6-anhydro-β-D-glucose) is a bicyclic anhydride of D-glucose (McGill and Williams 2008). It is formed by a range of chemical processes including pyrolysis of glucose (Sugisawa and Edo 1966), starch and cellulose (Lakshmanan et al. 1969; Itabaiana Junior et al. 2020), and is produced by the burning of biomass (Suciu et al. 2019). Emissions of LG into the atmosphere arise from burning of plant matter in open fires (1.7 million tonnes per annum) and biofuels (2.1 million tonnes per annum) and contributes to aerosols in the troposphere that affect air quality (Li et al. 2021). LG is formed from the burning of cellulose above 300 ℃ (Simoneit et al. 1999), and consequently, LG can be used as a molecular marker for biomass burning (Li et al. 2021), as a measure of air quality (Li et al. 2021), and has been used to trace ancient fire histories (paleo-fire records) in glacier snow and ice (You and Xu 2018). While LG degrades with a lifetime of 1.8 days in the atmosphere, an estimated 0.7 million tonnes of LG are deposited annually through dry deposition and in rain: 'sugar from the heavens' (Li et al. 2021). Consequently, soils across the earth likely experience continuous, low levels of LG deposition, irrespective of proximity to biomass combustion processes or precipitation.

Microorganisms that can grow on LG as sole carbon source are readily recovered from soils (Kitamura et al. 1991; Yasui et al. 1991; Nakahara et al. 1994; Iwazaki et al. 2018) including those associated with wood burning (Arya et al. 2022), as well as plant leaves (Iwazaki et al. 2018), and pond (Iwazaki et al. 2018) and waste water (Arya et al. 2022), suggesting that LG degrading organisms are widespread in Nature. While humans and mice do not appear to metabolize LG (Migliaccio et al. 2009; Bergauff et al. 2010), pathways for its degradation have been elucidated in fungi and bacteria. Fungi appear to exclusively use a pathway involving LG kinase, an ATP-dependent enzyme that converts LG in one step to glucose-6-phosphate (Kitamura and Yasui 1991; Bacik et al. 2015). On the other hand, bacteria use a multistep pathway (Yasui et al. 1991) that was first completely elucidated in *Bacillus smithii* S-720M (Kuritani et al. 2020) (**Figure 1**). In this organism, LG metabolism is initiated by the action of NADH-dependent LG dehydrogenase (LGDH, also known as LgdA), which forms 3-keto-LG. LGDH from *Pseudarthobacter phenanthenivorans* Sphe3 has been studied extensively, including determination of its 3D structure, identification of its active site, and the binding mode of LG (Sugiura et al. 2018). The 1,6-anhydro ring of 3-keto-LG is cleaved to give 2-hydroxy-3-keto-D-glucal, catalyzed by 3-keto-LG β-eliminase (LgdB1), and this is hydrated to give 3-keto-D-glucose, catalysed by 3-ketoglucose dehydratase (LgdB2). Finally,

3-keto-D-glucose is reduced to D-glucose, catalyzed by NADH-dependent glucose 3-

dehydrogenase (LgdC). LgdB2 appears to be a bifunctional enzyme that can also catalyze the

conversion of 3-keto-LG to 2-hydroxy-3-keto-D-glucal. Overall, this complex pathway

involves 3-4 enzymes that achieve the formal hydrolysis of LG. This pathway bears some

similarity to glycoside hydrolases of carbohydrate active enzyme family GH4 (Yip et al. 2004)

and GH109 (Liu et al. 2007), which in a single-step catalyze glycoside hydrolysis in an NADH-

dependent manner through a redox mechanism involving a 3-keto sugar intermediate (Yip and

Withers 2006). Possibly, the requirement for multiple enzymes for the net hydrolysis of LG

relates to the dramatic conformational change that occurs when the 1,6-anhydro ring of LG is

cleaved, meaning that different enzymes are required to recognize the substrate and

intermediates in the pathway.

All LG-degrading bacteria studied at a genetic level contain close homologs of *lgdA*,

and previous work has sought to identify genes involved in LG transport and metabolism by

analysis of the surrounding genomic regions. In the case of *Bacillus smithii* S-270M, the gene

cluster containing *lgdA-lgdB1-lgdB2-lgdC* also contains genes encoding ABC transporter

subunits and an ABC transporter substrate binding protein (Kuritani et al. 2020). However,

other LG degrading organisms such as *P. phenanthenivorans* Sphe3, *Paenibacillus athensensis*

MEC069, *Shinella sumterensis* MEC087, *Microbacterium* MEC084, and *Klebsiella*

pneumoniae MEC097 contain *lgdA-lgdB1* and genes encoding ABC transporter subunits and

substrate binding proteins, but homologs of *lgdB2* and *lgdC* were not identified (Arya et al.

2022), raising questions as to whether other enzymes may be used for the final steps of LG

metabolism.

Here, we isolate two LG degrading bacteria from soil and report their draft genome

sequences. Both organisms contain close homologs of *lgdA*. We perform comparative

proteomics to identify the operons that are upregulated upon growth on LG, and use sequence

similarity network analysis to study the genome neighborhoods surrounding *lgdA* to provide

insight into conserved genes surrounding this gene in a range of bacteria.

Materials and methods

Bacterial growth media

LG growth media was prepared using M9 salt medium, trace metal solution and vitamin

mixture and 5 mM levoglucosan as sole carbon source. M9 minimal medium consisted of 12.8

g/L Na_2HPO_4, 3 g/L KH_2PO_4, 1 g/L NH_4Cl, 0.5 g/L NaCl, 0.6 g/L $MgSO_4$, 0.22 g/L

$CaCl_2.2H_2O$ salts and trace metal solution consisted of 14.4 mg/L $FeCl_3$, 4.0 mg/L $CoCl_2.6H_2O$,

2.9 mg/L $CuCl_2.2H_2O$, 3.0 mg/L $MnCl_2.4H_2O$, 15.0 mg/L $ZnCl_2$, 0.2 mg/L $NiCl_2$, 3 mg/L

$Na_2MoO_4.6H_2O$ and 6 mg/L H_3BO_3. Vitamin mixture consisted of 2 µg/L biotin, 5 µg/L

calcium pantothenate, 10 µg/L thiamine hydrochloride, 10 µg/L p-aminobenzoic acid, 20 µg/L

nicotinic acid, 50 µg/L pyridoxamine dihydrochloride, and 4 µg/L B12 (cyanocobalamin).

Isolation of levoglucosan degrading species: *Paenarthrobacter nitrojuajacolis* LG01 and

Paenarthrobacter histidinolovorans LG02 were isolated from enriched culture, obtained from

soil samples collected from Royal Park (Melbourne, Australia).

Approximately 1 g of soil samples were suspended in 5 mL of sterilized LG growth media

containing 5 mM of LG as sole carbon source. The culture was incubated at 28 °C for 5 days

with continuous shaking at 250 rpm. A subsample of 50 µL was supplemented into freshly

prepared levoglucosan media and grown for a further 3 days. This step was repeated 5 times

and after the 5[th] transfer, cells were plated on LB agar plates consisting of 10 g/L tryptone, 5

g/L NaCl, 5 g/L yeast, 15 g/L agar and incubated overnight at 28 °C in the dark. Single colonies

were picked and used to inoculate fresh LG growth media and incubated at 28 °C with

continuous shaking at 250 rpm. Then, cells were plated and re-inoculated into fresh liquid LG

growth media 2 more times. Stock cultures were prepared by addition of 10% glycerol and

stored at -80 °C. Genomic DNA of LG degrading species was extracted using the GenElute

DNA extraction kit (Sigma) with inclusion of lysozyme and RNAase.

Phenotypic assays

Strains LG01 and LG02 were grown to visible turbidity in LG growth media containing 20

mM of levoglucosan at 28 °C for 3 days with continuous shaking at 250 rpm. Cultures were

centrifuged at 10000 rpm for 10 min (using Sigma laborzentrifugan model 1-15, rotor 12124)

and the supernatant liquid was diluted with 50% D_2O. ^{13}C-NMR spectra of diluted supernatant

liquid was recorded using a 500 MHz instrument.

Genome sequence, assembly, and annotation

Genomic DNA was sequenced using an Illumina MiSeq at the Peter Doherty Institute for

Infection and Immunity, Parkville, Victoria, Australia. DNA was prepared for sequencing on

the Illumina MiSeq platform using the Nextera XT DNA preparation kit (Illumina) with ×300

bp paired end chemistry and with a targeted sequencing depth of >50×. Draft genomes were

assembled using Shovill v1.1.0 (https://github.com/tseemann/shovill) and annotated using

Prokka v1.14.5 (Seemann 2014). GC percentage and ANI calculations were performed using

ANI calculator (https://www.ezbiocloud.net/tools/ani) (Yoon et al. 2017). Assembled genomes

have been deposited at the NCBI (Bioproject accession: LG01, PRJNA925556; LG02,

PRJNA925557). General features for isolated bacteria are reported in Table 1.

Sample preparation for proteomic analysis: Cells were grown to mid log phase and washed

3 times with phosphate buffered saline (PBS), collected by centrifugation at 10000 rpm, and

then snap frozen in liquid nitrogen and stored at -80 °C. Pelleted snap frozen bacterial cells

were solubilized in 4% SDS, 100 mM HEPES by boiling for 10 min at 95 °C. Protein

concentrations were assessed using the bicinchoninic acid protein assay (Thermo Fisher

Scientific) and 100 μg of each independent replicate prepared for digestion using Micro S-traps

(Protifi, USA) according to the manufacturer's instructions. Samples were reduced with 10

mM DTT for 10 mins at 95 °C and then alkylated with 40 mM iodoacetamide in the dark for 1

h. Samples were acidified to 1.2% phosphoric acid and diluted with seven volumes of S-trap

wash buffer (90% methanol, 100 mM tetraethylammonium bromide, pH 7.1), then were loaded

onto S-traps and washed 3 times with 400 μl of S-trap wash buffer. Samples were then digested

with 3 μg of trypsin (a 1:33 protease/protein ratio) in 100 mM tetraethylammonium bromide

for 4 h at 37 °C, then were collected by centrifugation with washes of 100 mM

tetraethylammonium bromide, followed by 0.2% formic acid followed by 0.2% formic acid /

50% acetonitrile. Samples were dried and further cleaned up using C18 Stagetips to ensure the

removal of particulate matter (Rappsilber et al. 2003; Rappsilber et al. 2007).

Reverse phase liquid chromatography–mass spectrometry for proteomics: Prepared

purified peptides from each sample were re-suspended in Buffer A* (2% acetonitrile, 0.01%

trifluoroacetic acid) and separated using a two-column chromatography setup composed of a

PepMap100 C_{18} 20-mm by 75-μm trap and a PepMap C_{18} 500-mm by 75-μm analytical column

(Thermo Fisher Scientific). Samples were concentrated onto the trap column at 5 μl/min for 5

min with Buffer A (0.1% formic acid, 2% DMSO) and then infused into an Orbitrap Eclipse

Mass Spectrometer (Thermo Fisher Scientific) or a Orbitrap Fusion Lumos equipped with a

FAIMS Pro interface both at 300 nl/minute via the analytical columns using a Dionex Ultimate

3000 UPLC (Thermo Fisher Scientific). For samples infused into the Orbitrap Fusion Lumos,

125-minute analytical runs were undertaken by altering the buffer composition from 2% Buffer

B (0.1% formic acid, 77.9% acetonitrile, 2% DMSO) to 22% B over 95 min, then from 22% B

to 40% B over 10 min, then from 40% B to 80% B over 5 min. The composition was held at

80% B for 5 min, and then dropped to 2% B over 2 min, then held at 2% B for another 8 min.

For samples infused into the Orbitrap Eclipse 95-minute analytical runs were undertaken by

altering the buffer composition from 2% Buffer B (0.1% formic acid, 77.9% acetonitrile, 2% DMSO) to 22% B over 65 min, then from 22% B to 40% B over 10 min, then from 40% B to 80% B over 5 min. The composition was held at 80% B for 5 min, and then dropped to 2% B over 2 min, then held at 2% B for another 8 min. The Orbitrap Eclipse mass spectrometer was operated in a data-dependent mode automatically switching between the acquisition of a single Orbitrap MS scan (375-1500 m/z and a resolution of 120k) and 3 seconds of Orbitrap MS/MS HCD scans of precursors (NCE of 30%, a maximal injection time of 22 ms, and a resolution of 15k). The Fusion Lumos Mass Spectrometer was operated in a stepped FAIMS data-dependent mode at two different FAIMS CVs -40 and -60. For each FAIMS CV a single Orbitrap MS scan (300-1600 m/z and a resolution of 60k) was acquired every 1.5 seconds followed by Orbitrap MS/MS HCD scans of precursors (Stepped NCE 28,32,37%, maximal injection time of 100 ms and a resolution of 30k).

Proteomic data analysis: Identification and LFQ analysis were accomplished using MaxQuant (v2.0.2.0) (Cox and Mann 2008) using the in house generated proteomes of LG01 and LG02 allowing for oxidation on Methionine. Prior to MaxQuant analysis datasets acquired on the Fusion Lumos were separated into individual FAIMS fractions using the FAIMS MzXML Generator (Hebert et al. 2018). The LFQ and "Match Between Run" options were enabled to allow comparison between samples. The resulting data files were processed using Perseus (v1.4.0.6) (Tyanova et al. 2016) to compare the growth conditions using Student's t-tests as well as Pearson correlation analyses.

Bioinformatic analysis: The protein sequences of LG degrading enzymes from *Pseudarthrobacter phenanthrenivorans* Sphe3 and *B. smithii* S-270M were used as queries to search the genomes of LG01 and LG02 using BLASTp.

The LGDH sequence from isolated strain LG01 was submitted as query in phmmer web server (https://www.ebi.ac.uk/Tools/hmmer/search/phmmer) (Potter et al. 2018). Standard parameters were used except the cut-off for E-value was set to 10^{-130}. The resulting sequences were used to build a hidden Markov model (HMM), which was used as query for *hmmsearch* in hmmer web server (https://www.ebi.ac.uk/Tools/hmmer/search/hmmsearch). The results were filtered and protein sequences with E-value $\leq$ -200 (**see Supplementary material file**) were used to generate a sequence similarity network (SSN) using the Enzyme Function Initiative-Enzyme Similarity Tool (https://efi.igb.illinois.edu/efi-est/) with minimum alignment score 120 (corresponding to $\geq$60% identity) and 170 (corresponding to $\geq$70%

identity) (Zallot et al. 2019). The SSN generated using the alignment score 120 was used to generate genome neighborhood diagrams (GND) with open reading frame (ORF) ± 10 neighbors using the Enzyme Function Initiative-Genome Neighborhood Tool (https://efi.igb.illinois.edu/efi-gnt/). A script was used to extract the accession codes of the retrieved neighbors (**Figure S2**). The LGDH neighbors were used to generate an SSN of neighbors (SSNN). The resulting SSNN was plotted using Cytoscape v3.9, with minimum alignment score of 50 (**Fig 5**).

The individual nodes of the LGDH SSN were manually coloured according to the occurrence of *lgdB1*, *lgdB2*, *lgdB3* and *lgdC* gene homologues in their neighborhood (ORF ± 10 neighbors). Initially, the EFI tool was used to generate a cluster number for each unique isofunctional cluster within the LGDH SSNN. The cluster number of the neighborhood genes from the SSNN for *lgdB1*, *lgdB2*, *lgdB3* and *lgdC* were mapped onto the LGDH SSN (alignment score of 170).

Results and discussion

Isolation and characterization of sulfoglycolytic bacteria

Bacteria able to grow on LG were selected by using soil samples (collected from the South-East corner of Royal Park, Melbourne) to inoculate a vitamin-enriched minimal media containing LG as sole carbon source. Sequential subculturing into fresh LG-minimal media, followed by repeated plating onto LB agar and then regrowth in LG-minimal media, led to isolation of strains LG01 and LG02 (**Table S1**). Strains LG01 and LG02 grew robustly on LG with peak growth rates of 0.073 and 0.288 A_{600}/min and achieved stationary phase after approximately 82 and 18 h, respectively (**Figure S1**). The absorbance at stationary phase for cultures grown on LG were approximately the same for cultures grown on equimolar glucose. ^{13}C-NMR analysis of spent culture medium of LG01 and LG02 grown on LG (20 mM) media did not reveal any carbon containing metabolic products.

To investigate the genetic basis for LG consumption by LG01 and LG02, genomic DNA extracted from these bacteria were sequenced using the Illumina NextSeq platform. **Table 1** shows the key features of the two draft genomes. Based on 16S rRNA gene sequence analysis the organisms were assigned as *Paenarthrobacter nitrojuajacolis* LG01 and *Paenarthrobacter histidinolovorans* LG02.

Identification of genetic loci expressed upon growth on LG

To identify possible proteins involved in LG metabolism we examined changes in the proteome of LG01 and LG02 cultivated on glucose versus LG at mid-log phase (**Figure 2**). For LG01, the largest and most significant changes we observed were an increase in the abundance of proteins encoded in two gene clusters in the genome of LG01 that comprise 19 genes, of which two are transcription factors (**Figure 3a**). For LG02, the largest and most significant changes were observed in proteins encoded in three gene clusters in LG02 that comprise 12 genes (**Figure 3b**). Both the identified clusters for LG01 and LG02 encode close homologs of *P. phenanthenivorans* Sphe3 and *B. smithii* S-270M LGDH proteins with 96.7% and 70.1% identity (for LG01) and 97.9% and 70.7% (for LG02), respectively. Both organisms also encode homologs of LgdB1 and LgdC, but with lower percentage identity (**Tables S3 and S4**). Collectively, the presence of genes encoding LgdA, LgdB1 and LgdC provides strong evidence for canonical LGDH pathways. Neither organism contains genes encoding homologs of LgdB2. However, the first cluster identified in LG02 is situated at the start of a contig, suggesting it is incomplete. Given that the genomic region encoding LG metabolism in LG02 appears incomplete, we focused our attention on LG01.

SSN analysis of LG metabolizing genetic loci

Previous work demonstrated that *B. smithii* contains four genes *lgdA-lgdB1-lgdB2-lgdC* that contribute to degradation of LG (Kuritani et al. 2020), whereas Arya and coworkers identified several organisms, such as *P. athensensis* MEC069 and *Microbacterium* MEC084 that appeared to contain only *lgdA-lgdB1* (Arya et al. 2022). On the other hand, LG01 contains *lgdA-lgdB1-lgdC*. LgdA and LgdC belong to the Gfo/Idh/MocA superfamily, which contains NAD(P)-dependent oxidoreductases that act on a wide range of hydroxylated substrates (Taberman et al. 2016). On the other hand, LgdB1 and LgdB2 belong to protein family PF01261 within the PFAM database, a large family of proteins that contain diverse activities including inositol catabolism (Fry et al. 2001).

To gain insight into the structure and composition of LgdA-containing gene clusters, we used SSN analysis. SSN analysis uses all-by-all pairwise sequence alignments to generate networks that show sequence relationships as a graph (Atkinson et al. 2009). Initially, the LgdA sequence from *P. nitrojuajacolis* LG01 was used as a query for a BLAST search on the phmmer webserver, which retrieved 379 sequences with *E*-value < 130. These were used to build a hidden Markov model (HMM) that was used as a query in the hmmsearch server. This retrieved 631 sequences with *E*-value < 130, which were filtered to *E*-value < 200 (percentage identity

> 60%) to give 265 sequences. These sequences were used to build an SSN using the Enzyme Function Initiative tools (https://efi.igb.illinois.edu/). At an alignment score of >170, this SSN breaks into subclusters that are closely aligned with taxonomy (**Figure 4a**). The sequences were used to retrieve the genome neighborhoods (± 10 genes around *lgdA*), providing 4877 neighbors. Some examples of genome neighborhood diagrams for bacteria from different phyla are provided in **Figure 4b**. These gene clusters reveal conservation of *lgdB1*, but not of *lgdB2* or *lgdC* (*vide infra*).

The gene neighbors were used to build an SSN of neighbors (SSNN). The SSNN contains isofunctional protein clusters that allow analysis of the common neighbors (**Figure 5**). The SSNN reveals three large protein clusters that contain components of ABC transporters: ATP binding protein, substrate binding protein and ABC transporter permease, along with several smaller clusters containing putative multifunction transporters (MFS), which presumably facilitate binding and import of LG. Multiple clusters are present that correspond to ROK, LytTR, PucR and LacI type transcription factors. Numerous oxidoreductase clusters are present. Three clusters, including the largest oxidoreductase cluster, belong to PF01408 and PF02894, and correspond to the nucleotide binding and C-terminal domains of LgdC/LgdA. The LgdC sequences from *B. smithii*, and LG01 and LG02 are all located in the largest cluster. Six clusters belong to PF01261, annotated as sugar phosphate isomerase or xylose isomerase/epimerase. LgdB1 from *B. smithii*, LG01 and LG02 are present within the largest of these clusters, which contains proteins of around 275 residues. LgdB2 from *B. smithii* is located within a smaller cluster, which contains proteins of around 325 residues. The remaining four clusters contain representatives that belong to the same PFAM family, but do not contain characterized members. The genomes of LG01 and LG02 encode proteins that fall into these orphan clusters, which we have named LgdB3 (71 sequences in cluster) and LgdB4 (4 sequences in cluster).

To understand the organization and composition of the LG degrading clusters we coloured the SSN of LgdA homologues according to the presence of *lgdB1*, *lgdB2*, *lgdB3* and *lgdC* in their genome neighborhood (**Figure 6a-d**). Almost all LgdA homologs (253 out of 265) have LgdB1 encoded in their gene cluster, while only a minority have LgdB2 (57 out of 265) or LgdB3 (71 out of 265) encoded in their gene cluster. A Venn diagram reveals that: (1) 126 organisms have only LgdB1; (2) LgdB2 always co-occurs with LgdB1 but never with LgdB3; and (3) LgdB3, with one exception, co-occurs with LgdB1 (**Figure 6e**). LgdB1 is widely distributed, occurring in Actinobacteria, Firmicutes, Proteobacteria, Deinococcus and Bacteroidetes. LgdB2 is more restricted, occurring mainly in Firmicutes and one Deinococcus.

LgdB3 occurs actinobacteria and proteobacteria. Collectively, these data suggest that LgdB2 and LgdB3 may have overlapping function, but that they are not essential. The rarer LgdB4 sugar phosphate isomerase is encoded within just 4 Actinobacteria, and co-occurs with only LgdA and LgdB1.

Approximately half (151 out of 265) of LgdA-containing organisms have genes for LgdC in their gene cluster. It has been suggested that LgdA may function as an oxidoreductase capable of reducing 3-keto-glucose, which could explain the absence of LgdC is some bacteria (Arya et al. 2022). LgdC is distributed across Actinobacteria, Firmicutes, Proteobacteria and Deinococcus.

Structural analysis of LG degrading proteins

Only a single example of an LG-degrading protein in the LGDH pathway has been experimentally determined, LGDH from *P. phenanthrenivorans* Sphe3 (PDB 6A3I) (Sugiura et al. 2018). Comparison of the AlphaFold2(Jumper et al. 2021; Varadi et al. 2022) model for LgdA from LG01 shows almost identical 3D fold (RMSD 0.252 Å; **Figure 7a**), and the presence of nucleotide cofactor and LG binding sites (**Figure 7b**).

Comparison of the AlphaFold2 models for *B. smithii* and LG01 LgdB1 proteins shows similar 3D folds (RMSD 0.433 Å) (**Figure 7c**), suggestive of similar functions. The peptide chain of LgdB1 is much shorter than LgdB2, LgdB3 and LgdB4, and comparison of AlphaFold models were poor, with high RMSD values (**Table S4b**). An overlay of *B. smithii* LgdB2, LG01 LgdB3 (RMSD 1.345 Å versus *B. smithii* LgdB2), and LgdB4 (RMSD 3.288 Å versus *B. smithii* LgdB2) shows broadly similar $(\alpha\beta)_8$ barrel folds, but weaker structural similarity (**Figure 7d**). The weaker structural similarity is difficult to interpret in the absence of structural studies to determine the molecular basis of substrate recognition and catalysis, but suggests that these proteins could have similar functions, or may act on modified intermediates in the pathway (eg 6-phospho-2-hydroxy-3-keto-D-glucal, formed by hexokinase), or may not be involved in the pathway.

Comparison of the AlphaFold2 models of *B. smithii* and LG01 LgdC reveals similar 3D folds (RMSD 0.641 Å) (**Figure 7e**). Overlay of LG01 LgdC and the *P. phenantrenivorans* LgdA in complex with NADH allowed a model to be built that showed LgdC could accommodate NADH, consistent with its demonstrated function as an NADH-dependent glucose 3-dehydrogenase (**Figure 7f**).

Conclusions

Two LG degrading soil bacteria belonging to the *Paenarthrobacter* genus (strains LG01 and LG02) were isolated by enrichment culture involving growth on LG from soil. These organisms can effect the complete degradation of LG, and give a stationary phase optical density on LG that is similar to that on glucose. Comparative proteomics revealed the presence of key genes encoding homologues of proteins involved in LG degradation, namely LGDH (LgdA), 3-keto-LG β-eliminase (LgdB1), and glucose 3-dehydrogenase (LgdC). However, the upregulated proteins included several additional members of the PFAM PF01261 sugar phosphate isomerase/xylose isomerase family that exhibit low homology to LgdB1. Structural modelling shows that these adopt similar folds to LgdB1 and their occurrence within the genomic regions of a wide range of bacteria encoding *lgdA-lgdB1-lgdC* gene clusters suggest that they may assist in the processing of intermediates in the LG-degradation pathway.

Abbreviations

ABC: ATP-binding cassette

DNA: Deoxyribonucleic acid

GND: Genome neighborhood diagram

HMM: Hidden Markov Model

LG: Levoglucosan

LGDH: Levoglucosan dehydrogenase

NADH: reduced nicotinamide adenine dinucleotide

ORF: open reading frame

SSN: Sequence similarity network

SSNN: Sequence similarity network of neighbors

Authors' contributions

SJW conceived the study with input from AK. AK performed enrichment culture and microbial growth and characterization. NS conducted proteomics. MH and SP performed genome sequencing, assembly and analysis. AK and EGB performed bioinformatic analysis. AK and SJW wrote the paper with input and approval of the final draft from the authors.

Funding

S.J.W is supported by the Australian Research Council (DP210100233, DP210100235). N.E.S

is supported by the Australian Research Council (FT200100270, DP210100362). E.D.G-B.

acknowledges support from: The Walter and Eliza Hall Institute of Medical Research; National

Health and Medical Research Council of Australia (NHMRC) project grant GNT2000517; the

Australian Cancer Research Fund; and the Brian M. Davis Charitable Foundation Centenary

Fellowship. We thank the Melbourne Mass Spectrometry and Proteomics Facility of The Bio21

Molecular Science and Biotechnology Institute for access to MS instrumentation.

Availability of data and material

The genome datasets generated and/or analysed during the current study are available in the

NCBI database with BioProject accession codes PRJNA925556 (strain LG01) and

PRJNA925557 (strain LG02). The mass spectrometry proteomics data has been deposited in

the Proteome Xchange Consortium via the PRIDE partner repository with the data set

identifier: PXD039086.

Code availability

Not applicable.

Declarations

Conflict of interest

The authors declare that they have no conflict of interest

Ethics approval

Not applicable

Consent to participate

Not applicable

Consent for publication

Not applicable

References

Arya AS, Hang MTH, Eiteman MA (2022) Isolation and Characterization of Levoglucosan-
Metabolizing Bacteria. Appl. Environ. Microbiol. 88:e0186821

Atkinson HJ, Morris JH, Ferrin TE, Babbitt PC (2009) Using sequence similarity networks
for visualization of relationships across diverse protein superfamilies. PLoS One
4:e4345
Bacik JP et al. (2015) Producing glucose 6-phosphate from cellulosic biomass: structural
insights into levoglucosan bioconversion. J. Biol. Chem. 290:26638-26648
Bergauff MA et al. (2010) Urinary levoglucosan as a biomarker of wood smoke: results of
human exposure studies. J. Expo. Sci. Environ. Epidemiol. 20:385-392
Cox J, Mann M (2008) MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat.
Biotechnol. 26:1367-1372
Fry J, Wood M, Poole PS (2001) Investigation of myo-inositol catabolism in *Rhizobium*
leguminosarum bv. viciae and its effect on nodulation competitiveness. Mol. Plant
Microbe Interact. 14:1016-1025
Hebert AS et al. (2018) Comprehensive single-shot proteomics with FAIMS on a hybrid
Orbitrap mass spectrometer. Anal. Chem. 90:9529-9537
Itabaiana Junior I, Avelar do Nascimento M, de Souza ROMA, Dufour A, Wojcieszak R
(2020) Levoglucosan: a promising platform molecule? Green Chem. 22:5859-5880
Iwazaki S, Hirai H, Hamaguchi N, Yoshida N (2018) Isolation of levoglucosan-utilizing
thermophilic bacteria. Sci. Rep. 8:4066
Jumper J et al. (2021) Highly accurate protein structure prediction with AlphaFold. Nature
596:583-589
Kitamura Y, Abe Y, Yasui T (1991) Metabolism of Levoglucosan (1,6-Anhydro-α-d-
glucopyranose) in Microorganisms. Agric. Biol. Chem. 55:515-521
Kitamura Y, Yasui T (1991) Purification and Some Properties of Levoglucosan (1,6-
Anhydro-β-d-glucopyranose) Kinase from the Yeast *Sporobolomyces salmonicolor*.
Agric. Biol. Chem. 55:523-529
Kuritani Y et al. (2020) Conversion of levoglucosan into glucose by the coordination of four
enzymes through oxidation, elimination, hydration, and reduction. Sci. Rep. 10:20066
Lakshmanan CM, Gal-or B, Hoelscher HE (1969) Production of levoglucosan by pyrolysis of
carbohydrates. Ind. Eng. Chem. Res. Dev. 8:261-267
Li Y et al. (2021) Impacts of chemical degradation on the global budget of atmospheric
levoglucosan and its use as a biomass burning tracer. Environ. Sci. Technol. 55:5525-
5536
Liu QP et al. (2007) Bacterial glycosidases for the production of universal red blood cells.
Nat. Biotechnol. 25:454-464
McGill N, Williams SJ (2008) Anomeric anhydro sugars. In: Fraser-Reid BO et al. (eds)
Glycoscience: Chemistry and Chemical Biology, 2 edn. Springer, Heidelberg,
Germany, pp 737-753
Migliaccio CT, Bergauff MA, Palmer CP, Jessop F, Noonan CW, Ward TJ (2009) Urinary
levoglucosan as a biomarker of wood smoke exposure: observations in a mouse model
and in children. Environ. Health Perspect. 117:74-79
Nakahara K, Kitamura Y, Yamagishi Y, Shoun H, Yasui T (1994) Levoglucosan
dehydrogenase involved in the assimilation of levoglucosan in Arthrobacter sp. I-552.
Biosci. Biotechnol. Biochem. 58:2193-2196
Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD (2018) HMMER web server:
2018 update. Nucleic Acids Res. 46:W200-W204
Rappsilber J, Ishihama Y, Mann M (2003) Stop and Go Extraction Tips for Matrix-Assisted
Laser Desorption/Ionization, Nanoelectrospray, and LC/MS Sample Pretreatment in
Proteomics. Anal. Chem. 75:663-670

Rappsilber J, Mann M, Ishihama Y (2007) Protocol for micro-purification, enrichment, pre-
fractionation and storage of peptides for proteomics using StageTips. Nat. Protoc.
2:1896-1906

Seemann T (2014) Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068-
2069

Simoneit BRT et al. (1999) Levoglucosan, a tracer for cellulose in biomass burning and
atmospheric particles. Atmospheric Environ. 33:173-182

Suciu LG, Masiello CA, Griffin RJ (2019) Anhydrosugars as tracers in the Earth system.
Biogeochemistry 146:209-256

Sugisawa H, Edo H (1966) The thermal degradation of sugars I. Thermal polymerization of
glucose. J. Food. Sci. 31:561-565

Sugiura M, Nakahara M, Yamada C, Arakawa T, Kitaoka M, Fushinobu S (2018)
Identification, functional characterization, and crystal structure determination of
bacterial levoglucosan dehydrogenase. J. Biol. Chem. 293:17375-17386

Taberman H, Parkkinen T, Rouvinen J (2016) Structural and functional features of the
NAD(P) dependent Gfo/Idh/MocA protein family oxidoreductases. Protein Sci.
25:778-786

Tyanova S et al. (2016) The Perseus computational platform for comprehensive analysis of
(prote)omics data. Nat. Methods 13:731

Varadi M et al. (2022) AlphaFold Protein Structure Database: massively expanding the
structural coverage of protein-sequence space with high-accuracy models. Nucleic
Acids Res. 50:D439-D444

Yasui T, Kitamura Y, Nakahara K, Abe Y (1991) Metabolism of Levoglucosan (1,6-
Anhydro-β-d-glucopyranose) in Bacteria. Agric. Biol. Chem. 55:1927-1929

Yip VL et al. (2004) An unusual mechanism of glycoside hydrolysis involving redox and
elimination steps by a family 4 β-glycosidase from *Thermotoga maritima*. J. Am.
Chem. Soc. 126:8354-8355

Yip VL, Withers SG (2006) Breakdown of oligosaccharides by the process of elimination.
Curr. Opin. Chem. Biol. 10:147-155

Yoon SH, Ha SM, Lim J, Kwon S, Chun J (2017) A large-scale evaluation of algorithms to
calculate average nucleotide identity. Antonie Van Leeuwenhoek 110:1281-1286

You C, Xu C (2018) Review of levoglucosan in glacier snow and ice studies: Recent progress
and future perspectives. Sci. Total Environ. 616-617:1533-1539

Zallot R, Oberg N, Gerlt JA (2019) The EFI web resource for genomic enzymology tools:
leveraging protein, genome, and metagenome databases to discover novel enzymes
and metabolic pathways. Biochemistry 58:4169-4182

Table 1. Key features of the LG01 (BioProject accession: PRJNA925556) and LG02

(BioProject accession: PRJNA925557) draft genome assemblies.

	LG01	LG02
Genome size	4777618	4332355
Number of contigs	86	862
GC %	62.86	62.92
Number of ORFs	4455	4403
Number of putative genes	4519	4462
Number of putative tRNA	56	55
Number of putative rRNA	7	3
Number of putative tmRNA	1	1
Number of genes assigned a function (%)	2419 (53%)	2094 (47%)

Fig. 1. Pathway for levoglucosan (LG) metabolism in *Bacillus smithii* S-2701M. NADH-dependent LG dehydrogenase (LgdA, also known as LGDH) oxidizes LG to 3-keto-LG. 3-Keto-LG β-eliminase (LgdB1) cleaves the 1,6-anhydro ring to give 2-hydroxy-3-keto-D-glucal. 3-Keto-glucose dehydratase (LgdB2) catalyses the hydration of 2-hydroxy-3-keto-D-glucal. Glucose 3-dehydrogenase (LgdC) catalyzes the reduction of 3-keto-D-glucose to D-glucose. LgdB2 is reported to be a bifunctional enzyme that can also catalyze the conversion of 3-keto-LG to 2-hydroxy-3-keto-D-glucal.

Fig. 2. Manhattan plot of comparative proteomics data for a) *Paenarthrobacter nitrojuajacolis* LG01 and b) *Paenarthobacter histidinolovorans* LG02, grown on LG versus glucose. The relative fold change, denoted by the colour, and the -log10(p-value), denoted by on the *y*-axis, of the observed differences are plotted according to gene order, *x*-axis, revealing two gene clusters of up-regulated proteins in LG01 and three gene clusters in LG02.

500 **Fig. 3.** Gene clusters identified by comparative proteomics analysis. a) Up-regulated gene
501 clusters identified in *Paenarthrobacter nitrojuajacolis* LG01. b) Up-regulated gene clusters
502 identified in *Paenarthrobacter histidinolovorans* LG02.

Fig. 4. a) Sequence similarity network of LgdA orthologs with alignment score 170 (i.e., 70% identity). Each node is an LgdA ortholog belonging to PF01408 and PF0289, and nodes are colored based on phylum Firmicutes (Persian green), Actinobacteria (purple), Proteobacteria (Humbrol blue), Deinococcus (light blue), Bacteroidetes (light purple) and unknown (grey). Yellow diamonds indicate proteins functionally characterized: *Pseudarthrobacter phenanthrenivorans* Sphe3 and *Bacillus smithii* S-2701M, and in this study: *Paenarthrobacter nitrojuajacolis* LG01 and *Paenarthrobacter histidinolovorans* LG02. **b)** Genome neighborhood diagrams (GND) for different species that contains LgdA (red), LgdB1 (bright pink), LgdB2 (magenta) and LgdC (blue). Species names are colored according to phylum as for the SSN.

516   w

Fig. 5. Sequence similarity network of neighbors (SSNN) of LgdA. SSNN was generated with an alignment score of 50. Isofunctional clusters are colored accordingly to their assigned function. Three large clusters are for ABC transporters (orange, yellow and light yellow). LgdB1 and LgdB2 map onto two clusters colored bright pink and magenta. LgdC maps onto a blue cluster. Additional clusters annotated as inosose dehydratase/sugar phosphate isomerase (light pink) and sugar phosphate isomerase/epimerase (new one) contain proteins annotated in this work as LgdB3 and LgdB4. Yellow diamonds denote proteins that have been studied previously or in this study. From *Bacillus smithii* S-2701M: *Bs*LgdB1, *Bs*LgdB2, *Bs*LgdC; from *Paenarthrobacter nitrojuajacolis* LG01: *Pn*LgdB1, *Pn*LgdB3, *Pn*LgdC; from *Paenarthrobacter histidinolovorans* LG02: *Ph*LgdB1, *Ph*LgdB3, *Ph*LgdC. Clusters smaller than 7 members are not shown, except for the cluster for LgdB4.

Fig. 6. Sequence similarity network (SNN) of LgdA orthologues with alignment score 170 (i.e. 70% identity). Each node is an LgdA ortholog belong to PF01408 and PF02894. (**a-d**), nodes are colored according to presence of specified LG degrading pathway enzymes located within an ORF window ± 10 of LgdA (each LG degrading gene belongs to a specific cluster in SSNN, **Figure 5**). Nodes are colored if they possess the following within their genome neighborhood: a) LgdB1 (bright pink), b) LgdB2 (magenta); c) LgdB3 (light pink); d) LgdC (blue). **e)** Venn diagram shows co-occurrence of LgdB1, LgdB2 and LgdB3 in each genome neighborhood surrounding LgdA.

Figure 7. LG01 Alphafold2 models and overlays with experimentally determined structure of LgdA. (a) An overlay of the model for LgdA from LG01 (cyan) with levoglucosan dehydrogenase from *Pseudarthrobacter phenanthrenivorans* Sphe3 (green, PDB ID: 6A3I). (b) Close-up view of the putative active site of LgdA from LG01 (cyan) with nucleotide cofactor and levoglucosan substrate overlaid from 6A3I, and the proposed catalytic residues E103 and H193 shown in stick form. (c) An overlay of the models for LG01 and *B. smithii* LgdB1 (cyan and green, respectively). (d) An overlay of the models for *B. smithii* LgdB2 (green), LG01 LgdB3 (cyan) and LG01 LgdB4 (magenta). (e) An overlay of the models for *B.*

548   *smithii* and LG01 LgdC (cyan and green, respectively). (f) A close-up view of the putative
549   active site of LG01 LgdC, with nucleotide cofactor overlaid from 6A3G, and the putative
550   catalytic residues E95 and H185 shown in stick form.