The post-translational modification profile of TAR DNA-Binding Protein
(TDP-43) in platelets of patients with Alzheimer's disease: An exploratory
study for blood-based biomarker development.

Qwynton Johnson[1], Mahan Hadjian[1], Alpha Bah[1], Sean Smith[1], Edina Kosa[1], Abdulbaki
Agbas[1,2]*

[1] College of Osteopathic Medicine, Department of Basic Sciences, Kansas City University, Kansas
City, MO, USA

[2] Heartland Center for Mitochondrial Medicine, Kansas City, MO, USA

* Correspondence:
Abdulbaki Agbas
aagbas@kansascity.edu

Keywords: TDP-43, post-translational modification, platelets, Alzheimer's disease, biomarkers

Abstract

The assignment of blood-based biomarkers for neurodegenerative diseases is of great clinical value.
Well-developed and validated blood-based biomarkers can serve in early diagnosis and prognosis as
well as aid in patient screening when recruiting for clinical trials. We attempted to establish a portfolio
for post-translationally modified TAR DNA/RNA-binding protein (TDP-43), a regulator of nuclear
transcription, in platelet cytosol obtained from patients with Alzheimer's disease (AD) comparing to
age-matched healthy subjects and a disease control cohort. We aimed to identify the most prominent
post-translational modifications of TDP-43 as an AD-relevant biomarker and to demonstrate that such
an assessment can be performed in peripheral blood. We have isolated TDP-43 protein from human
platelet cytosol utilizing an Immunoaffinity chromatography. The eluates were immunoprobed with a
series of antibodies raised against post-translationally modified proteins. We employed a capillary
electrophoretic immunoassay (CEI) to assess the phosphorylated TDP-43 profile. We observed that
SUMOylation, phosphorylation, ubiquitination, and cysteine oxidation of TDP-43 are more prominent
in platelet cytosol of AD patients as compared to control subjects. These studies will pave the way for
identifying disease-specific TDP-43 derivatives that can be potential biomarkers for early diagnosis
and the development of therapeutics.

1 Introduction

Alzheimer's disease (AD) is the most common cause of dementia and is currently estimated to affect
more than 5 million people in the United States, with an expected increase to 13.8 million by the year
2060. The typical clinical presentation is progressive loss of memory and cognitive function, ultimately
leading to a loss of independence, inflicting a heavy personal toll on the patient and the family. The costs
of care of patients with AD in 2010 were estimated at more than \$172 billion in the United States
alone. The annual cost is projected to inflate to one trillion dollars by 2050 unless disease-modifying
treatments are developed (1). Attempts are being made to fill this demand, with the global market for
AD treatment expected to grow from \$4.8 billion in 2020 to an estimated \$6.3 billion by 2025. That
said, such attempts have been met with challenges; clinical trials of Alzheimer's drugs ultimately

failed more than 99% of the time between 2002 and 2012 (2). While treatment is multifaceted, a
deterrent to developing disease-modifying therapies is the lack of biomarkers for early detection and
evaluation. Therefore, well-validated and well-defined biomarkers of Alzheimer's disease are
needed. These biomarkers will improve the design of clinical trials, allow for development of more
effective therapeutics, and offer the opportunity for prevention trials.

TAR DNA-binding protein (TDP-43) is a pivotal protein in the regulation of transcription and, by
extension, cell homeostasis (3). There are several studies demonstrating that post-translational
modifications of TDP-43 occur in a wide variety of neurodegenerative states in the central nervous
system (CNS) where TDP-43 is associated with plaque formation (4-6). These plaques are insoluble
by nature and can be histologically observed only in post-mortem tissues. Both overexpression and
downregulation of TDP-43 is toxic for cells (7). Cells are unpredictable to process post-
translationally modified TDP-43 and that of aggregated forms. This feature makes the therapeutic
targeting of TDP-43 a challenging task. Identification of disease-specific post-translationally
modified (PTM) TDP-43 profiles is necessary so that therapeutic intervention may be logically
designed; hence, to establish chemically modified TDP-43 as a disease-specific and a validated
blood-based biomarker is of importance. The PTM profile of TDP-43 may help to discriminate
neurodegenerative diseases based on the progress stage. This will enhance our understanding of the
pathological TDP-43 occurrence in neurodegenerative diseases.

We are working on developing a blood-based biomarker for neurodegenerative diseases such as AD,
amyotrophic lateral sclerosis (ALS), and inclusion body myositis (IBM). Peripheral blood sampling
provides easy access for obtaining repeated specimens using less invasive methods. Blood-derived
platelets are a particularly attractive platform, providing an encapsulated environment for the
cytosolic form of TDP-43 protein protected from the actions of several degradative enzymes such as
proteases and phosphatases in blood serum/plasma (8). We have demonstrated the occurrence of
hyperphosphorylation of TDP-43 in platelets (8-10) and others showed that plasma TDP-43 is largely
or wholly restricted to platelets (11). Other potential PTMs with the capacity to produce toxic gain of
function in the cell, such as cleavage, aggregation, acetylation, ubiquitination, SUMOylation, and
cysteine-oxidation, need to be documented since neurodegenerative diseases vary in their repertoire
of PTMs of target proteins (12). Blood-derived platelets provide a surrogate venue with a less
invasive access to the central nervous system to evaluate PTMs of TDP-43 during the course
diseases.

In this explorative study, we will compile a PTM profile of TDP-43 in platelets derived from patients
with AD. This approach, if successful, can be applied to other neurodegenerative diseases such as
amyotrophic lateral sclerosis Parkinson's disease, Niemen-Pick C, and motor neuron disease.

2   Materials and Methods

2.1 *Platelet cytosol preparation:* Alzheimer's disease patients were recruited through University of
Kansas Alzheimer's Disease Research Center (KUADRC). Blood samples were obtained from
KUADRC under an approved IRB protocol (IRB#299644-3). All patients underwent annual Uniform
Data Set clinical evaluations, which includes a battery of cognitive tests. The subjects included in this
study were all males. The preparation of platelets was described in a previous manuscript (10).
Briefly, approximately 8-9 mL of venous blood were collected over a vacutainer collection tube
containing acid citrate dextrose (ACD) anti-coagulant reagent and brought to the laboratory within 1
hr. Platelet-rich plasma (PRP) was carefully separated from the buffy coat after centrifuging at 200 x
g for 20 minutes. Prostaglandin I_2 (PGI_2) was added to PRP to prevent platelet aggregation (1 µL of

1mg /mL PGI$_2$ per mL of PRP), gently mixed, and subjected to centrifugation (1,200 x g for 15
minutes) to pellet the platelets. The platelet pellet was washed with 1.0 mL of citrate wash buffer and
subjected to an additional centrifugation (1,200 x g for 15 minutes) to eliminate possible red blood
cell contamination. The pellet was then ruptured in 0.6 mL of rupture buffer and further subjected to
sonication. The resulting platelet homogenate was centrifuged at 20,000 x g for 30 min at 4 °C to
remove platelet membranes. The supernatant was aliquoted and stored at -80 °C freezer until use.

2.2 High Performance Immunoprecipitation (HPIP):

2.2.1 Cross-linking anti-TDP-43 antibody to Protein-A-conjugated affinity media: Approximately
10 µg commercial anti-TDP-43 rabbit antibody (Proteintech Cat #10782-2-AP) was chemically
cross-linked to Protein-A-conjugated affinity media pre-packed in a pipette tip (PhyTip Cat #PTR-
92-20-01) (tip-affinity column). Cross-linking chemistry was performed in the presence of
disuccinimidyl suberate (DSS), a non-cleavable cross-linking agent, according to an established
method in our laboratory. The procedure of cross-linking the primary antibody to affinity media
eliminates the co-elution of IgG antibodies with the target protein; hence, it increases the stability of
antibody binding (13) and allows for reuse of the tip-affinity column up to two more times without
losing antibody binding efficiency. We employed a fully automated multi-channel pipette equipped
with software (PureSpeed, RAININ Cat #17013549) for the cross-linking process. A second tip-
affinity column cross-linked with 10 µg home-prepared non-immune rabbit IgG antibody was used as
a negative control.

2.2.2 Immunoprecipitation of TDP-43: Platelet cytosols obtained from Alzheimer's patients and
age-matched controls were added to a deep-well plate and the automated multi-channel pipette was
placed into position over the plate. Pre-programmed pipetting was initiated. The HPIP procedure
included steps for equilibration, capture, wash, and elution. The TDP-43-enriched eluate was then
neutralized with a small amount of 1M Tris-Buffer (pH 8.0). This eluate proceeded to SDS-PAGE or
was stored at -20 °C for future use. Detailed information on the HPIP program utilized with the
electronic pipette is provided in **Supplemental Table-1.**

2.3 Immunoblot Analysis: One hundred micrograms of platelet cytosol protein from the
immunoprecipitation eluate (see step 2.2.2.) were loaded into a home-prepared one-well analytical
SDS-PAGE gradient (4-20%) gel. Proteins were resolved in the electrophoretic field (100 V; 1hr)
under reducing conditions and subsequently transferred onto PVDF membrane (116 mA; 1hr). Total
protein staining was performed according to manufacturer instructions (Revert 700, Total Protein
Staining, LI-COR Cat #926-11021) to assess the protein transfer efficiency and to normalize protein
loading after completing the immunoblotting procedure. The membrane was destained using
commercial reagent (Revert 700, Reversal Solution, LI-COR Cat #926-11013) and then blocked in a
fish serum blocking buffer (Thermo Fisher, Cat #37527) for 1 hr at RT. The membrane was dissected
into 10 equal strips and stored at 4 °C until use. Each strip was individually incubated with a relevant
primary antibody (i.e., anti-SUMO1 and -SUMO 2/3, anti-ubiquitin, anti-cysteine, and anti-acetyl-
lysine antibodies) overnight at 4 °C. The following day, the individual membrane strips were
thoroughly washed with Tris-Buffered Saline, 0.1% Tween-20 (TBST) buffer and incubated with IR-
conjugated secondary antibodies (IRDye 800CW, Cat #926-32211 [rabbit], #926-32210 [mouse]) for
1 hr at RT. The membrane strips were washed with TBST for 3x5 min and 1x5 min with TBS to
eliminate the nonspecific fluorescence generated by Tween-20. The membranes were then scanned
by an imager (LI-COR Odyssey) and the images were analyzed by Image Studio[TM] Lite
Quantification software (Ver.3.1.4).

2.4 Capillary Electrophoretic Immunoassay (CEI): Anti-phospho-(S409/410)-TDP-43 antibody
(Proteintech Cat #80007-1-RR) was used to determine the phosphorylated TDP-43 profile in both
AD and ALS platelet cytosols. The assay was performed according to a well-established
methodology developed in our lab (10). The samples were prepared according to manufacturer
instructions (Proteinsimple kit Cat #SM-W004). The fully automated CEI analyzer (Proteinsimple,
Cat#004-600. This product is currently discontinued. Updated versions are available) is capable of
measuring the phosphorylated TDP-43 levels in very small volume (0.4 µL), precisely analyzed the
results, and created the final graph. Due to this features, CEI became a preferred method in our
laboratory for high throughput screening for protein-based analysis in platelet cytosols obtained from
patients with neurodegenerative diseases.

2.5 Sequence based prediction analysis: The TDP-43 amino acid sequence (UniProt/Swiss-Prot :
Q13148) was explored via several proteomic databases for all predicted post-translational
modifications that can occur (14). We showed only the predicted PTM sites in human TDP-43,
rendered by PhosphositePlus (v6.6.0.4)

3. Results

We chemically cross-linked the polyclonal anti-TDP-43 rabbit antibody and non-immune rabbit IgG
on Protein-A-conjugated affinity material (**Fig.1**) and checked for any IgG cleavage during the
washing and elution process. We did not observe IgG molecules in the eluates (**Supplemental Fig-
1A**). One hundred micrograms of platelet cytosolic protein mixture was applied to the HPIP tip-
affinity column and antibody-bound proteins were eluted. The immunoblots of the eluates obtained
from the platelet cytosol of AD patients and age-matched control subjects showed different levels of
PTMs of TDP-43. The signal intensities were normalized to total protein staining according to the
manufacturer protocol and the data were graphically illustrated (**Fig.2A**). Ubiquitinated and
SUMOylated TDP-43 appeared to be the most notable PTMs, expressing the greatest signal intensity.
We have observed reduced cysteine oxidation of TDP-43 in platelet cytosol samples obtained from
patients with AD (**Fig.2B**) as compared to age-matched controls (**Fig.2C**). We were not able to detect
acetyl-lysine modification. Due to low protein concentration (1-1.5 mg/mL) of individual platelet
cytosol, we pooled both AD (n=10) and age-matched control cohort (n=10) samples and performed
HPIP only once as a proof-of-concept. For this reason, statistical analysis was not performed at this
time (**Fig.2A**). Data obtained from the CEI assay revealed two phosphorylated TDP-43 (pTDP-43)
species observed in platelet cytosol samples obtained from AD patients and from that of ALS
patients (the disease control). There was a notable elevation of the pTDP-43 band localized at
approximately 60 kDa in the AD platelet cytosol (green peak) as opposed to the healthy control (blue
peak) and the ALS disease control (red peak). Note that the same anti-phospho-(S409/410)-TDP-43
antibody recognized different phosphorylated TDP-43 protein species in different neurodegenerative
diseases (**Fig.3**)**.** Although we have utilized a number of publicly available programs, the most
popular PTM studies were revealed using the PhosphositePlus (Ver.6.6.0.4) online prediction
program with a number of references (14). Peak heights on y- axis refers to how many publications
are available in particular PTMs of TDP-43 (**Fig.4**). The non-immune rabbit IgG tip-column,
assigned as a negative control, did not pull down TDP-43 from the platelet cytosol samples
(**Supplemental Fig-1C**).

4.   Discussion

The profile of PTMs of TDP-43 could be a useful indicator for the early diagnosis and prognosis of
neurodegenerative diseases as well as discriminating such diseases from one another. Platelet-derived

TDP-43 has been considered a potential candidate as a surrogate blood-based biomarker for
neurodegenerative disease (8, 10, 15, 16). In this study, we analyzed the most popular PTMs of TDP-
43 in platelet cytosols obtained from patients with AD as well as a healthy age-matched control and
ALs as disease control group.

We chose platelet cytosol as the venue for measuring the TDP-43 profile. Platelets are anucleated,
cell fragments derived from megakaryocytes (17-19). TDP-43 is a nuclear protein that has the ability
to shuttle back and forth between the nucleus and cytosol (20). When TDP-43 proteins undergo
PTMs due to disease, as in neurodegeneration, or physical trauma ,such as repeated concussions in
contact sports (21, 22), TDP-43 translocates to the cytoplasm. The cytoplasmic accumulation of
TDP-43 is sufficient to induce aggregation through protein-protein interaction and phosphorylation
(23, 24). Although remains largely unexplored, new studies support the notion that phosphorylation
of TDP-43 occurs after aggregate formation in cytosol (reviewed in (25). Therefore, platelets are an
appropriate system of study, reflecting a cytosolic environment where PTMs of TDP-43 will be
abundant. In a separate pilot study, it has been shown that approximately 90 % of plasma TDP-43
was found in platelet enriched plasma as compared to 10% TDP-43 in platelet poor plasma (11) (R.
Bowser, personal communication, October 12, 2022).

There are several publicly available PTM prediction programs that can be employed for forecasting
possible PTMs of TDP-43. We have checked the majority of the sites provided in a review paper
(14); that said, many of links were either inactive or under maintenance at the time of writing this
manuscript. Of all the proteomic databases and tools that we tried, only phosphosite.org
(https://www.phosphosite.org/homeAction.action) generated a comprehensive output. Therefore, we
presented the data output of PhosphositePlus v6.6.0.4, which provided all popular PTMs along with
references for each (**Fig.4**). Prediction analysis of TDP-43 for natural disorder regions (PONDR®)
revealed that most of the phosphorylation events occur between amino acids 359-410 (**Supplemental**
Figure-2).

We have isolated platelets from whole blood within a one hour timeframe by employing a two-step
centrifugation as described in a previous manuscript (10). After lysis and removal of membranous
fragments by centrifugation, the platelet cytosol samples were stored at -80 °C until use. We
immunoprecipitated TDP-43 proteins from platelet cytosol obtained from AD subjects ,as well as
age-matched but otherwise healthy control subjects, employing HPIP technology as illustrated in
Fig.1. Platelet cytosol samples obtained from patients with ALS were included as a disease control
cohort for the CEI assay so as to compare the pTDP-43 profile in both diseases (i.e., AD and ALS).
Traditional immunoblotting was employed to demonstrate that TDP-43 undergoes posttranslational
modification in AD as compared to control subjects. One notable pattern observed in AD patients
was a modest increase in TDP-43 SUMOylation as compared to healthy patients (**Fig.2**). The
regulatory role of SUMOylation has been implicated in fundamental processes, with several reports
indicating that the aging process could be influenced by SUMOylation (26-30). Based on our
exploratory results (**Fig.2**), we may propose that the SUMOylation characteristics of TDP-43 could
be considered as a disease-specific hallmark and possible target in AD. We recognize that we must
also study other neurodegenerative diseases such as ALS, Parkinson's disease, Huntington's disease,
multiple system atrophy, and progressive supranuclear palsy and demonstrate the SUMOylation
pattern (or lack thereof) of TDP-43 in such diseases (31-33).

We have noticed that there is a pattern of increase in TDP-43 ubiquitination in parallel to increasing
SUMOylation. These two PTMs vary in terms of function. Ubiquitination is used to tag proteins for
degradation in proteasome machinery, whereas SUMOylation alters gene expression, chromatin

structure, signal transduction, and maintenance of the genome (34). These two patterns (i.e.,
SUMOylation and ubiquitination) may suggest that TDP-43 ubiquitination is a response to aberrant
protein expression caused by increased TDP-43 SUMOylation ,an attempt to remove the aberrant
protein via the proteasome apparatus. More work needs to be done concerning the SUMOylation
characteristics of TDP-43 during AD progression into the more advanced stage. The other relevant
biological question would be whether SUMOylation of TDP-43 induces proteasome activity so that
ubiquitinated TDP-43 can be eliminated from cell.

The most obvious post-translationally modified TDP-43 species is pTDP-43, with 64 potential
phosphorylation sites (41 Ser, 15 Thr, and 8 Tyr residue) available on the protein (**Supplemental
Fig. 2**) (8). However, **d**isorder **e**nhanced **p**hosphorylation **p**redictor (DEPP) revealed that 20 Ser, 4
Thr, and 4 Tyr residues are statistically more probable sites for phosphorylation. Most of these sites
are located in the C-terminus region of the sequence. We employed CEI technology, and observed
that commercial anti-phospho-(S409/410)-TDP-43 antibody recognized two separate protein
populations ,at MW 47 kDa and MW 60 kDa, in platelets obtained from patients with ALS and
patients with AD, respectively (**Fig.3**). This pattern has held for several ongoing clinical trials that we
are involved in. It is currently unknown whether TDP-43 undergoes a hyper-phosphorylation process
that prompts this presence in AD platelets. If that is the case, we may consider the ~60 kDa pTDP-43
species to be an AD-specific signature protein discriminating from ALS, which instead expresses an
alternate but equally specific pTDP-43 species at ~47 kDa. More work needs to be performed in
terms of reproducibility, specificity, and sensitivity of pTDP-43 measurements.

We have also observed that anti-phospho-(S409/410)-TDP-43 antibody recognizes cross-linked
pTDP-43 in ALS at MW ~230 kDa. (**Supplemental Figure-3**). It is possible that as a super-
phosphorylated TDP-43 conglomerate, this TDP-43 species do not completely enter the separation
matrix packed in the capillary column of CEI. High-stringency solubilization and reducing conditions
may resolve some of the cross-linked pTDP-43 conglomerates, and we are currently working on this
issue and testing whether AD platelet cytosol shows a similar pattern (i.e., ~230 kDa protein
conglomerate).

Analysis using CEI has precisely provided accurate molecular weight information for target proteins
in every assay we have performed using commercial molecular weight markers. We did not use the
other antibodies utilized during immunoblotting studies (i.e., anti- SUMO1, anti-SUMO2-3, anti-
cysteine sulfonation, anti-acetylated Lys, anti-ubiquitin antibodies) in the CEI assay system. Not
every antibody works well in the CEI platform and optimization steps (concerning kinetics of binding
and minimizing background) must be taken for each antibody prior to use in CEI assays. We
routinely measure pTDP-43 levels in human platelet cytosol for several clinical trials (16); hence the
anti-phospho-(S409/410)-TDP-43 antibody was well optimized for CEI assays. We have included
CEI data for only the pTDP-43 profile in this manuscript CEI studies utilizing other commercial
antibodies may be done once the binding kinetics of those antibodies have been optimized in the
future.

Acetylation of proteins can occur as a co-translational or post-translational modification (35). Two
main sites (Lys145 and Lys192) of acetylation in TDP-43 were identified by generating acetylation
mimics (36). However, in a contradictory study, mass spectrometry analysis revealed that residues
Lys145 and Lys192 were not acetylated in TDP-43 inclusions found in brain samples of ALS
patients (37). We were also unable to assess acetylation of TDP-43 on Lys residues by
HPIP/immunoblotting methods (**Fig.2**). Either the anti-acetyl lysine antibody that we used for this
study (Cytoskeleton, Inc., Cat# AAC01-S) did not work in our hands or there was no recognizable

acetylation of TDP-43 in AD platelet cytosol. In the future, we will use antibodies from other
vendors ,paying special attention to other potential acetylation sites of TDP-43, such as Lys82 (37),
to clarify this ambiguity regarding whether Lys acetylation in TDP-43 is disease-specific.

The other peculiar trend we noticed was that all forms of TDP-43 expressing oxidized cysteine
moieties with an oxidized thiol group (i.e., sulfenic RSOH, sulfinic RSO_2H, and sulfonic RSO_3H)
were reduced in AD patients as compared to age-matched controls. It seems that there was less
cysteine modification in AD than in age-matched controls (**Fig.2**), which is contrary to what we
expected (i.e., more sulfonated cysteine residues in TDP-43). It has been hypothesized that reactive
oxygen species may trigger some TDP-43 pathology (38). Cohen et al., proposed that there are at
least six cysteine residues (Cyst39, Cys50, Cys173, Cys175, Cys198, and Cys244) in the TDP-43
protein sequence that are targets for cysteine oxidation (38). We have demonstrated in a previous
manuscript that functional protein undergoes aggregation via -SH groups in the aging brain (39).
Since a disulfide bridge (S-S) forms between two cysteine amino acids, we were expecting that
cysteine modification in TDP-43 would be elevated. It is possible that TDP-43 takes on a more
soluble form in platelet cytosol; hence, TDP-43 aggregation is less. This conjecture needs to be
experimentally tested by first pre-treating isolated platelets with dimedone (5.5-Dimethyl-1, 3-
cyclohexanedione) ,which will help identify cysteine residues expressing sulfenic acid groups from
other cysteine oxoforms (40), then immunoprecipitating TDP-43, and immunoprobing with an anti-
cysteine (sulfonate) antibody.

In conclusion, several post-translational modifications of TDP-43 can be analyzed in human platelet
cytosol, mediated by the compromise of the blood-brain barrier in neurodegenerative diseases (41,
42). Many CNS biomolecules can be observed in peripheral blood tissue; therefore, the profile of
platelet TDP-43 and its derivative species could serve as a surrogate biomarker for neurodegenerative
diseases.

3   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

4   Author Contributions

QJ, MH, AB, and SS are medical students. All were involved in experimental procedure, data
analysis, and graphic creations. EK helped the students perform the experiments. MH was involved
in editing the manuscript. AA conceived the idea, planned the experiments, and wrote the
manuscript.

5   Funding

This project is funded by an intramural grant awarded to AA.

6   Acknowledgments

The authors acknowledge the Office of Research and Sponsored Programs at Kansas City University
for awarding summer research fellowships for QJ, MH, and AB.

References

1.   Brookmeyer R, Evans DA, Hebert L, Langa KM, Heeringa SG, Plassman BL, et al. National
estimates of the prevalence of Alzheimer's disease in the United States. Alzheimers Dement.
2011;7(1):61-73.

2.   Bredesen DE, John V. Next generation therapeutics for Alzheimer's disease. EMBO Mol
Med. 2013;5(6):795-8.

3.   Zuo-Shang X. Does a loss of TDP-43 function cause neurodegeneration? MolNeurodeg.
2012;7:27.

4.   DeTure MA, Dickson DW. The neuropathological diagnosis of Alzheimer's disease.
Molecular neurodegeneration. 2019;14(1):32.

5.   Davis SA, Gan KA, Dowell JA, Cairns NJ, Gitcho MA. TDP-43 expression influences
amyloidbeta plaque deposition and tau aggregation. Neurobiology of disease. 2017;103:154-62.

6.   Lloyd GM, Trejo-Lopez JA, Xia Y, McFarland KN, Lincoln SJ, Ertekin-Taner N, et al.
Prominent amyloid plaque pathology and cerebral amyloid angiopathy in APP V717I (London)
carrier - phenotypic variability in autosomal dominant Alzheimer's disease. Acta Neuropathol
Commun. 2020;8(1):31.

7.   Suzuki H, Lee K, Matsuoka M. TDP-43-induced death is associated with altered regulation of
BIM and Bcl-xL and attenuated by caspase-mediated TDP-43 cleavage. The Journal of biological
chemistry. 2011;286(15):13171-83.

8.   Wilhite R, Sage JM, Bouzid A, Primavera T, Agbas A. Platelet phosphorylated TDP-43: an
exploratory study for a peripheral surrogate biomarker development for Alzheimer's disease. Future
Sci OA. 2017;3(4):FSO238.

9.   Statland JM, Barohn RJ, McVey AL, Katz JS, Dimachkie MM. Patterns of Weakness,
Classification of Motor Neuron Disease, and Clinical Diagnosis of Sporadic Amyotrophic Lateral
Sclerosis. Neurol Clin. 2015;33(4):735-48.

10.   Sage J, Hall L, McVey A, Barohn RJ, Statland JM, Jawdat O, et al. Use of Capillary
Electrophoresis Immunoassay to Serach for Potential Biomarkers of Amyotrophic Lateral Sclerosis
in Human Platelets. JVisExp. 2020;156(e60638):1-9.

11.   Luthi-Carter R, Tentiller N, Le Roux-Bourdieu M, Font L, Agbas A, Pfiefer A, et al., editors.
Compartmentalization of plasma TDP-43 in platelets: Implications for TDP-43 related biomarker
development. International Conference on Alzheimer's and Parkinson's Diseases and related
neurological disorders; 2023 3/30/2023; Gothenburg, Sweden. Gothenburf, Sweden2023.

12.   Buratti E. TDP-43 post-translational modifications in health and disease. Expert Opin Ther
Targets. 2018;22(3):279-93.

13.   Hermanson GT, Mallia AK, Smith PK. Immobilized affinity ligand techniques. USA:
Academic Press, Inc.; 1992 1992. 453 p.

14.   Kamath KS, Vasavada MS, Srivastava S. Proteomic databases and tools to decipher post-
translational modifications. J Proteomics. 2011;75(1):127-44.

15.   Wilkins HM, Dimachkie MM, Agbas A. Blood-based Biomarkers for Amyotrophic Lateral
Sclerosis. In: Araki T, editor. Amyotrophic Lateral Sclerosis. Brisbane (AU)2021.

16.   Statland JM, Moore D, Wang Y, Walsh M, Mozaffar T, Elman L, et al. Rasagiline for
amyotrophic lateral sclerosis: A randomized, controlled trial. Muscle Nerve. 2019;59(2):201-7.

17.   Schulze H, Korpal M, Hurov J, Kim SW, Zhang J, Cantley LC, et al. Characterization of the
megakaryocyte demarcation membrane system and its role in thrombopoiesis. Blood.
2006;107(10):3868-75.

18.   Thon JN, Italiano JE. Platelets: production, morphology and ultrastructure. Handb Exp
Pharmacol. 2012(210):3-22.

19.   Junt T, Schulze H, Chen Z, Massberg S, Goerge T, Krueger A, et al. Dynamic visualization of
thrombopoiesis within bone marrow. Science. 2007;317(5845):1767-70.

20.   Suk TR, Rousseaux MWC. The role of TDP-43 mislocalization in amyotrophic lateral
sclerosis. Molecular neurodegeneration. 2020;15(1):45.

21.   Tripodis Y, Alosco ML, Zirogiannis N, Gavett BE, Chaisson C, Martin B, et al. The Effect of
Traumatic Brain Injury History with Loss of Consciousness on Rate of Cognitive Decline Among
Older Adults with Normal Cognition and Alzheimer's Disease Dementia. Journal of Alzheimer's
disease : JAD. 2017;59(1):251-63.

22.   Turner RC, Lucke-Wold BP, Robson MJ, Omalu BI, Petraglia AL, Bailes JE. Repetitive
traumatic brain injury and development of chronic traumatic encephalopathy: a potential role for
biomarkers in diagnosis, prognosis, and treatment? Front Neurol. 2012;3:186.

23.   Neumann M, Kwong LK, Lee EB, Kremmer E, Flatley A, Xu Y, et al. Phosphorylation of
S409/410 of TDP-43 is a consistent feature in all sporadic and familial forms of TDP-43
proteinopathies. Acta neuropathologica. 2009;117(2):137-49.

24.   Sasaguri H, Chew J, Xu YF, Gendron TF, Garrett A, Lee CW, et al. The extreme N-terminus
of TDP-43 mediates the cytoplasmic aggregation of TDP-43 and associated toxicity in vivo. Brain
research. 2016;1647:57-64.

25.   Eck RJ, Kraemer BC, Liachko NF. Regulation of TDP-43 phosphorylation in aging and
disease. Geroscience. 2021;43(4):1605-14.

26.   Princz A, Tavernarakis N. SUMOylation in Neurodegenerative Diseases. Gerontology.
2020;66(2):122-30.

27.   Vijayakumaran S, Pountney DL. SUMOylation, aging and autophagy in neurodegeneration.
Neurotoxicology. 2018;66:53-7.

28.   Ficulle E, Sufian MDS, Tinelli C, Corbo M, Feligioni M. Aging-related SUMOylation pattern
in the cortex and blood plasma of wild type mice. Neuroscience letters. 2018;668:48-54.

29.   Gong L, Sun Q, Li DW. Sumoylation in Cellular Senescence and Aging. Curr Mol Med.
2017;16(10):871-6.

30.   Li F, Zhang L, Craddock J, Bruce-Keller AJ, Dasuri K, Nguyen A, et al. Aging and dietary
restriction effects on ubiquitination, sumoylation, and the proteasome in the heart. Mech Ageing Dev.
2008;129(9):515-21.

31.   Krumova P, Weishaupt JH. Sumoylation in neurodegenerative diseases. Cell Mol Life Sci.
2013;70(12):2123-38.

32.   Mandel N, Agarwal N. Role of SUMOylation in Neurodegenerative Diseases. Cells.
2022;11(21).

33.   Chen X, Zhang Y, Wang Q, Qin Y, Yang X, Xing Z, et al. The function of SUMOylation and
its crucial roles in the development of neurological diseases. FASEB journal : official publication of
the Federation of American Societies for Experimental Biology. 2021;35(4):e21510.

34.   Gill G. SUMO and ubiquitin in the nucleus: different functions, similar mechanisms? Genes
Dev. 2004;18(17):2046-59.

35.   Polevoda B, Sherman F. The diversity of acetylated proteins. Genome Biol.
2002;3(5):reviews0006.

36.   Cohen TJ, Hwang AW, Restrepo CR, Yuan CX, Trojanowski JQ, Lee VM. An acetylation
switch controls TDP-43 function and aggregation propensity. Nat Commun. 2015;6:5845.

37.   Kametani F, Obi T, Shishido T, Akatsu H, Murayama S, Saito Y, et al. Mass spectrometric
analysis of accumulated TDP-43 in amyotrophic lateral sclerosis brains. Sci Rep. 2016;6:23281.

38.   Cohen TJ, Hwang AW, Unger T, Trojanowski JQ, Lee VM. Redox signalling directly
regulates TDP-43 via cysteine oxidation and disulphide cross-linking. The EMBO journal.
2012;31(5):1241-52.

39.   Agbas A, Zaidi A, Michaelis EK. Decreased activity and increased aggregation of brain
calcineurin during aging. Brain research. 2005;1059(1):59-71.

40.   Seo YH, Carroll KS. Profiling protein thiol oxidation in tumor cells using sulfenic acid-
specific antibodies. Proceedings of the National Academy of Sciences of the United States of
America. 2009;106(38):16163-8.

41.   Tayler H, Miners JS, Guzel O, MacLachlan R, Love S. Mediators of cerebral hypoperfusion
and blood-brain barrier leakiness in Alzheimer's disease, vascular dementia and mixed dementia.
Brain Pathol. 2021;31(4):e12935.

42.   Jacob A, Alexander JJ. Complement and blood-brain barrier integrity. Mol Immunol.
2014;61(2):149-52.

FIGURE LEGENDS

FIGURE-1: Illustration of the workflow for high performance immunoprecipitation (HPIP).
Tip-columns were prepared by cross-linking a-TDP-43 antibody (steps-1,2) and used for high
efficiency immuno-isolation of TDP-43 (steps-3,4)from platelet cytosol. The eluate were analyzed by
classical immunoblotting assay (step-5). Created with BioRender.com

FIGURE-2: Post-translational modifications profile of TDP-43. (A) The bar graphs illustrate
comparison between AD patients and age-matched control cohort in terms of PTMs of TDP-43
profile in platelet cytosol. The more noticeable elevations (blue bar) were observed in SUMOylation
and ubiquitination detected by anti-SUMO-1/anti-SUMO2,3 and anti-ubiquitination antibodies,
respectively. Since pooled platelet cytosol samples from AD patients (n=10) and age-matched control
cohort (n=10) were used for HPIP, the assay performed only once; hence, no statistical analysis were
performed. **(B)** Shows the immunoblot analysis of AD patients and **(C)** shows that of age-matched
control cohort of PTM TDP-43 profile with various antibodies.

FIGURE-3: pTDP-43 electropherogram of capillary electrophoretic immunoassay (CEI). Anti-
p(S409/410)TDP-43 antibody was used in this assay. The same antibody recognized TDP-43 species
with different molecular weight. The prominent pTDP-43 peak appeared at ~60 kDa (green color
peak) for AD patients whereas the most prominent pTDP-43 peak appeared at ~47 kDa (red color
peak) for ALS patients. Control group pTDP-43 peaks (blue color peak) did appeared in both
location with the same intensity.

FIGURE-4: PhosphositePlus predictions fort post-translational modifications of TDP-43. TDP-
43 amino acid sequence (Uniport/Swiss-Prot: Q13148) was uploaded in PhosphositePlus
(Ver.6.6.0.4) program and several PTMs were predicted. The most notable PTMS were
phosphorylation, acetylation, and ubiquitination. The height of circles represent the total number of
published papers. The most phosphorylation sites were appeared to be in C-terminal region of TDP-
43. The most ubiquitination sites were located on of the RNA-binding site of TDP-43 (RRM-1)

Supplemental Figure-1: Anti-TDP-43 antibody cross-linking onto tip-affinity column. (A) After
cross-linking anti-TDP-43 antibody (rabbit), a HPIP procedure was performed with PBS buffer (pH
7.4) as input sample, Flow-through and eluate was collected and analyzed by 4-20% gradient
SDS/PAGE. The gel was stained by silver staining method. No IgG leaching was observed, meaning
we achieved to cross-link the antibody to tip-affinity column. **(B)** Anti-TDP-43 antibody was
chemically cross-linked onto a tip-affinity column (PhyTip Cat# PTR-92-20-01). HPIP was
performed . Input consisted of platelet cytosol obtained from AD patients. Input and eluate samples
were analyzed by 4-20% gradient SDS/PAGE followed by silver staining. TDP-43 was pull down
and no IgG leach was observed. **(C)** Two tip-affinity columns were prepared using non-immune
rabbit IgG (left lane) and anti-TDP-43 antibody (rabbit) (right lane) were cross-linked. The tip-
affinity columns were used for immunoisolating TDP-43from platelet cytosols obtained from AD
patients. No TDP-43 signal was obtained from non-immune IGG column whilst anti-TDP-43
antibody column pulled down TDP-43 proteins. Non-immune IgG column served as negative control.

Supplemental Figure-2: Disorder enhanced phosphorylation predictor (DEPP). DEEP predictor
revealed that the most of the phosphorylation events occur between amino acids 359-410 (framed
region).

Supplemental Figure-3: Super phosphorylated TDP-43. Approximately 230 kDa protein peak
(right-pink color) in platelet cytosols obtained from ALS patients. Anti- phosphorylated (S409/410)
TDP-43 antibody was used for CEI assay. We did not observed the same peak when we performed
CEI assay wit pan anti-TDP-43 antibody.

Supplemental Table-1: Programming steps for fully automated multi-channel pipette. These
programming steps were optimized for anti-TDP-43 antibody. Users be advised that each antibody
has different binding kinetics onto affinity matrix; therefore, antibody of interest need to be
optimized.

Figure-1

Figure-2

Figure-3

Figure-4