***PerturbSci-Kinetics*: Dissecting key regulators of transcriptome kinetics through scalable single-cell**

RNA profiling of pooled CRISPR screens

Zihan Xu[1,2], Andras Sziraki[1,2], Jasper Lee[1], Wei Zhou[1,3], Junyue Cao[1,3,4]

Affiliations:

[1]Laboratory of Single Cell Genomics and Population Dynamics, The Rockefeller University, New York,

NY, USA

[2]The David Rockefeller Graduate Program in Bioscience, The Rockefeller University, New York, NY,

USA

[3]Senior author

[4]Lead Contact

*Correspondence: wzhou@rockefeller.edu (W.Z.), jcao@rockefeller.edu (J.C.)

Abstract

Here we described *PerturbSci-Kinetics*, a novel combinatorial indexing method for capturing three-layer

single-cell readout (*i.e.,* whole transcriptome, nascent transcriptome, sgRNA identities) across hundreds

of genetic perturbations. Through *PerturbSci-Kinetics* profiling of pooled CRISPR screens targeting a

variety of biological processes, we were able to decipher the complexity of RNA regulations at multiple

levels (*e.g.,* synthesis, processing, degradation), and revealed key regulators involved in miRNA and

mitochondrial RNA processing pathways. Our technique opens up the possibility of systematically

decoding the genome-wide regulatory network underlying RNA temporal dynamics at scale and cost-

effectively.

Main

Cellular functions are determined by the expression of millions of RNA molecules, which are tightly regulated across several critical steps, including but not limited to RNA synthesis, splicing, and degradation. Dysregulated transcriptome kinetics have been linked to various diseases, including cancer[1], intellectual disability[2], and neurodegenerative disorders[3]. However, our knowledge regarding how critical molecular regulators affect genome-wide RNA kinetics is still scarce, partly due to the lack of scalable tools. For example, while single-cell transcriptome analysis coupled with pooled CRISPR screens have recently yielded fundamental insight into the gene regulatory mechanisms[4–9], the readout of these methods only provides a snapshot of gene expression programs, thus is insufficient to decipher the complexity of RNA dynamics (*e.g.,* synthesis, splicing, and degradation). To resolve this challenge, we developed *PerturbSci-Kinetics*, by integrating CRISPR-based pooled genetic screens, highly scalable single-cell RNA-seq by combinatorial indexing, and metabolic labeling to recover single-cell transcriptome dynamics across hundreds of genetic perturbations.

The key features of the new method include: (i) A novel combinatorial indexing strategy (referred to as '*PerturbSci')* was developed for targeted enrichment and amplification of the sgRNA region that carries the same cellular barcode with the single-cell whole transcriptome (**Fig 1a**). A modified CROP-seq vector system[8] was adopted in *PerturbSci*, enabling the direct capture of sgRNA sequences[59] (**Extended Data Fig 1**). With extensive optimizations on primer designs and reaction conditions (**Extended Data Fig 2**), *PerturbSci* yields a high capture rate of sgRNA (*i.e.,* up to 99.7%), comparable to previous approaches for single-cell profiling of pooled CRISPR screens[4–9]. Furthermore, built on an extensively improved single-cell RNA-seq by three-level combinatorial indexing (*i.e.,* EasySci-RNA[10]), *PerturbSci* substantially reduced the library preparation costs for single-cell RNA profiling of pooled CRISPR screens (**Fig 1b, Supplementary file 3**). In addition, to maximize the gene knockdown efficacy, we used a multimeric fusion protein *dCas9-KRAB-MeCP2[11]*, a highly potent transcriptional repressor that outperforms conventional dCas9 repressors. (ii) By integrating *PerturbSci* with 4-thiouridine (4sU) labeling method, *PerturbSci-Kinetics* exhibited an order of magnitude higher throughput than the previous single-cell metabolic profiling approaches (*e.g.*, scEU-seq, sci-fate, scNT-seq)[12–15](**Fig 1a**). Of note, we extensively optimized the cell fixation condition to reduce the cell loss rate during permeabilization and *in-situ* thiol (SH)-linked alkylation reaction[16–22] (referred to as 'chemical conversion') (**Extended Data Fig 3**). Following 4sU labeling and chemical conversion, the nascent transcriptome and the whole

transcriptome from the same cell can be distinguished by T to C conversion in reads mapping to mRNAs[14].

The kinetic rate of mRNA dynamics (*e.g.,* synthesis and degradation) were then calculated as a multi-

layer readout for each genetic perturbation (**Fig 1a**, **Methods**). We further optimized the computational

pipeline for nascent reads calling based on the established pipeline of sci-fate[14], enabling the separation

of single cell nascent transcriptomes with high accuracy (**Extended Data Fig 4**).

As a proof-of-concept, we first tested our approach in a mouse 3T3-L1-CRISPRi cell line transduced with

a non-target control (NTC) sgRNA or sgRNA targeting a *Fto* gene (encoding an RNA demethylase). We

found that sgRNA expression was detected in over 99% of all cells, with a median of 284 sgRNA UMI

detected per cell in our optimal condition (*i.e.,* 1uM gRNA primer + 50uM dT primer in reverse

transcription) (**Extended Data Fig 2f**). We then generated a human HEK293 cell line with the inducible

expression of dCas9-KRAB-MeCP2[11] (HEK293-idCas9) and tested the sgRNA capture efficiency using

an NTC sgRNA and a sgRNA targeting the *IGF1R* gene (encoding insulin-like growth factor 1 receptor).

The transductions of the NTC and target sgRNAs were performed independently, such that each cell

received a unique perturbation. We then carried out a *PerturbSci* experiment on a 1:1 mixture of cells

from these two conditions. We recovered the target sgRNA expression in 96.7% of cells, of which 95.2%

were annotated as sgRNA singlets with a median of 81 sgRNA UMIs detected per cell (**Fig 1c**). Single-

cell gene expression analysis confirmed the induction of dCas9 after Dox treatment, as well as the

significantly decreased *IGF1R* expression in cells transduced with the target sgRNA (**Fig 1d**). Strongly

reduced *IGF1R* mRNA and protein levels were further validated by RT-qPCR and flow cytometry

(**Extended Data Fig 5**), validating the high knockdown efficiency of the system.

We next sought to validate the *PerturbSci-Kinetics* for capturing three-layer readout (*i.e.,* whole

transcriptome, nascent transcriptome, sgRNA identities) at the single-cell level. Following 4sU labeling

(200uM for two hours), we mixed HEK293-idCas9 cells transduced with NTC or IGF1R sgRNA at a 1:1

ratio for fixation and chemical conversion. We observed a significant enrichment of T to C mismatches in

mapped reads of the chemical conversion group, similar to our previous study[14](**Fig 1e**). A median of

22.1% of newly synthesized reads was recovered in labeled and chemically converted cells, compared to

only 0.8% in control groups (**Fig 1f**). Reassuringly, the proportion of reads mapped to exonic regions was

significantly lower in newly synthesized reads compared with pre-existing reads (p-value < 1e-20,

Tukey's test after ANOVA) (**Fig 1g**). Indeed, genes with a higher fraction of nascent reads were

significantly enriched in highly dynamic biological processes such as transcription coregulator activity

(FDR = 5.7e-12) and protein kinase activity (FDR = 2.6e-08)[23] **(Fig 1h)**. By contrast, genes with a lower

fraction of nascent reads were strongly enriched for processes essential for cell vitality, such as the

structural constituent of ribosome (FDR = 1.5e-42), unfolded protein binding (FDR = 4.5e-11), and

translation regulator activity (FDR = 8.2e-10) **(Fig 1i)**. Notably, the chemical conversion step is fully

compatible with sgRNA detection at single-cell resolution: we recovered sgRNAs from 97% of chemically

converted cells (a median of 62 sgRNA UMIs/cell), 92.6% of which were annotated as sgRNA singlets

(Fig 1j-k). These analyses demonstrate the capacity of *PerturbSci-Kinetics* to profile both transcriptome

dynamics and the associated perturbation identity at the single-cell level.

Fig. 1. *PerturbSci-Kinetics* **enables joint profiling of transcriptome dynamics and high-throughput gene perturbations by pooled CRISPR screens. a**. Scheme of the experimental and computational strategy for *PerturbSci-Kinetics*. The dot plot on the upper right shows the number of cells profiled in this study for comparison with the published single-cell metabolic profiling datasets[14,15,24]. Scale, the highest number of cells profiled in a single experiment of each technique. IAA, iodoacetamide. *4sU, chemically modified 4sU. R, steady-state RNA level. α, RNA synthesis rate. β, RNA degradation rate. Exp, steady-state expression. Synth, synthesis rate. Deg, degradation rate. **b**. Bar plot showing the estimated library preparation cost for *PeturbSci-Kinetics* and other published techniques[25,26] for single-cell transcriptome analysis coupled with CRISPR screens. **c**. Scatter plot showing the number of unique sgRNA transcripts detected per cell in the *PerturbSci* experiment for profiling cells transduced with sgNTC or sgIGF1R. **d**. The left box plot shows the normalized expression of dCas9-KRAB-MeCP2 in untreated or Dox-induced HEK293-idCas9 cells. The right box plot shows the normalized expression of IGF1R in Dox-induced HEK293-idCas9 cells transduced with sgNTC or sgIGF1R. Gene counts of each single cell were normalized by the total gene count, multiplied by 1e4 and then log-transformed. **e.** Bar plot showing the normalized percentage of all possible single base mismatches in reads from sci-fate (blue), and *PerturbSci-*

Kinetics on chemically converted (green) or unconverted cells (orange). Normalized mismatch rates, the
percentage of each type of mismatch in all sequencing bases. **f**. Box plot showing the fraction of recovered
nascent reads in single-cell transcriptomes across conditions: no 4sU labeling + no chemical conversion,
4sU labeling + no chemical conversion, and 4sU labeling + chemical conversion. **g.** Box plot showing the
ratio of reads mapped to exonic regions of the genome in nascent reads, pre-existing reads, and reads of
the whole transcriptomes across single cells. **h-i.** Bar plots showing the significantly enriched Gene
Ontology (GO) terms in the list of genes with low (h) or high (i) nascent reads ratio (**Methods**). **j.** Box
plot showing the number of unique sgRNA transcripts detected per cell in cells with or without the
chemical conversion. **k.** We performed *PerturbSci-Kinetics* experiment using converted/unconverted
HEK293-idCas9 cells transduced with sgNTC/sgIGF1R. Stacked bar plot showing the fraction of
converted/unconverted cells identified as sgNTC/sgIGF1R singlets, doublets, and cells with no sgRNA
detected.
126

To dissect the impact of key genetic regulators on transcriptome kinetics, we performed a *PerturbSci-Kinetics* experiment on HEK293-idCas9 cells transduced with a library of 699 sgRNAs, containing 15 NTC sgRNAs and sgRNAs targeting 228 genes involved in a variety of biological processes including mRNA transcription, processing, degradation, and others (**Fig 2a, Supplementary Table 1**). The cloning and lentiviral packaging were carried out in a pooled fashion similar to the previous report[27] (**Methods**). We then infected the HEK293-idCas9 cell line with the sgRNA lentiviral library at a low multiplicity of infection (MOI) (2 repeats at MOI = 0.1 and 2 repeats at MOI = 0.2) to ensure most cells received only one sgRNA. After a 5-day puromycin selection to remove non-infected cells, we harvested a fraction of cells for bulk library preparation ('day 0' samples). The rest of the cells were treated with Doxycycline (Dox) to induce the dCas9-KRAB-MeCP2 expression for an additional seven days. We then introduced 4sU labeling (200uM for two hours) and harvested samples for both bulk and single-cell *PerturbSci-Kinetics* library preparation ('day 7' samples). The time window for the screening period was chosen to minimize the effect of population dropout[28] (**Methods**).

As expected, the induction of CRISPRi significantly changed the abundance of sgRNAs in the cell population, which is consistent between replicates and the previous study[29] (**Extended Data Fig 6a-b, Supplementary Table 2, 3**). For example, the sgRNAs targeting genes involved in essential biological functions, such as DNA replication, ribosome assembly, and rRNA processing, were strongly depleted in the screen (**Extended Data Fig 6c**). Reassuringly, the sgRNA abundance recovered by *PerturbSci-kinetics* strongly correlated with the bulk library (Pearson correlation r = 0.988, p-value < 2.2e-16) (**Fig 2b**). After filtering out low-quality cells, we recovered 161,966 labeled cells, 88.1% of which had matched sgRNAs. 78% of these matched cells were annotated as sgRNA singlets (**Extended Data Fig 7a**). Despite the relatively low (17.9%) duplication rate of sequencing, we obtained a median of 2,155 UMIs per cell. Most (698 out of 699) sgRNAs were recovered, with a median of 28 sgRNA UMIs detected per cell. We further filtered out sgRNAs with low knockdown efficiencies (<= 40% expression reduction of target genes compared with NTC) (**Extended Data Fig 7b-e**). Finally, 98,315 cells were retained for downstream analysis, corresponding to a median of 484 cells per gene perturbation with a median of 67.7% knockdown efficiency of target genes (**Fig 2c**). To further validate the impact of perturbations, we aggregated single-cell transcriptomes and generated a 'pseudo-cell' for each targeted gene, followed by PCA dimension reduction and UMAP visualization[30]. Indeed, perturbations targeting paralogous genes

(*e.g., EXOSC5* and *EXOSC6*; *CNOT2* and *CNOT3*) or related biological processes (*e.g.,* RNA degradation,

RNA splicing, oxidative phosphorylation (OXPHOS) and energy metabolism) were readily clustered

together in the low dimension space (**Fig 2d**).

Taking advantage of *PerturbSci-Kinetics* for uniquely capturing multiple layers of information, we

performed differentially-expressed gene (DEG) analysis (**Supplementary Table 4**) and quantified gene-

specific synthesis and degradation rates of DEGs in each perturbation based on an ordinary differential

equation[31] (**Methods**). As a quality control, we first examined the kinetics of genes targeted by CRISPRi,

which were known to function through transcriptional repression[32,33]. Indeed, these genes exhibited

strongly reduced synthesis rates while their degradation rates were only mildly affected (**Fig 2c**). We then

investigated the impact of genetic perturbations on the global transcriptome dynamics (*i.e.,* synthesis,

splicing and degradation) (**Methods, Supplementary Table 5, 6**). As expected, the knockdown of genes

involved in transcription initiation (*e.g., GTF2E1*, *TAF2*, *MED21,* and *MNAT1*), mRNA synthesis (*e.g.,*

POLR2B and *POLR2K*), and chromatin remodeling (*e.g., SMC3, RAD21, CTCF, ARID1A*) significantly

downregulated the global synthesis rates but not the degradation rates (**Fig 2e-f**). In contrast, perturbations

targeting components of critical biological processes such as DNA replication (*e.g.*, *POLA2*, *POLD1*),

ribosome synthesis and rRNA processing (*e.g.*, *POLR1A, POLR1B, RPL11, RPS15A*), mRNA and protein

processing (*e.g.*, *CNOT2, CNOT3, CCT3, CCT4*) substantially reduced both RNA synthesis and

degradation globally, indicating a compensatory mechanism for maintaining overall transcriptome

homeostasis (**Fig 2e-f, Extended Data Fig 8a, b**). Furthermore, we observed significantly reduced

fractions of exonic reads in nascent transcripts, an indicator of dysregulated splicing dynamics, following

perturbations of genes involved in the main steps of RNA processing, including 5' capping (*e.g., NCBP1*),

RNA splicing (*e.g., LSM2, LSM4, PRPF38B, HNRNPK*), and 3' cleavage/polyadenylation (*e.g., CPSF2,*

CPSF6, NUDT21, CSTF3) (**Fig 2g, Supplementary Table 7**). In addition, the knockdown of genes

involved in OXPHOS & energy metabolism (*e.g., GAPDH, NDUFS2, ACO2*) also significantly reduced

the exonic reads ratio in nascent reads (**Fig 2g, Extended Data Fig 8c**), potentially due to the fact that the

mRNA processing is highly energy-dependent[34,35,36].

We next sought to investigate the regulators of mitochondrial RNA dynamics by quantifying the ratio of

nascent/total read counts (referred to as "turnover rate") mapped to mitochondrial genes (**Methods**).

Notably, we observed a significantly downregulated turnover rate of mitochondrial-specific RNA

following the perturbation of multiple metabolism-related genes (*e.g., GAPDH, FH, PKM* involved in glycolysis, *ACO2,* and *IDH3A* involved in the TCA cycle, *NDUFS2* and *COX6B1* involved in oxidative phosphorylation) **(Fig 2h, Extended Data Fig 8d)**. Furthermore, the knockdown of *LRPPRC* introduced the most substantial defect in the mitochondrial turnover and the expression levels of all mitochondrial protein-coding genes **(Fig 2h, Extended Data Fig 9a)**. Intriguingly, 5 of 13 mitochondrial protein-coding genes, including *MT-CO1, MT-ATP8, MT-ND4, MT-CYB, and MT-ATP6*, were regulated by both decreased transcription and increased degradation (**Extended Data Fig 9a, Supplementary Table 9**). This result was supported by a previous study[37] (**Extended Data Fig 9b**) and was also consistent with the known functions of *LRPPRC* in regulating the life cycles of mitochondrial RNA from synthesis to degradation[38–40]. For comparison, the nuclear-encoded differentially expressed genes (DEGs) following *LRPPRC* knockdown were significantly changed mostly at the transcription level (39 out of 48 genes, **Extended Data Fig 9c**). Upon closer inspection of promoter regions of these synthesis-regulated genes, we observed a strong enrichment of *ATF4* and *CEBPG* binding motifs, suggesting their potential roles as downstream transcriptional regulators of *LRPPRC*. Indeed, *ATF4* and *CEGPG* have been reported as core transcriptional activators involved in stress sensing[41],and both genes were substantially upregulated in *LRPPRC* knockdown cells (**Extended Data Fig 9d-e**).

Extending the above analysis, we examined the gene-specific synthesis and degradation regulation across all perturbations (**Supplementary Table 10**). Among all 14,618 perturbation-DEG pairs identified in the study, 22.9% of them exhibited rate changes, in which 15.1% showed significant synthesis rate changes only, 3.6% showed degradation rate changes only, and 4.2% showed both changes, suggesting complex mechanisms regulating gene expression upon perturbations[42] (**Extended Data Fig 10**). As expected, most degradation-regulated DEGs were associated with perturbations on mRNA surveillance/processing (*e.g., UPF1, UPF2, SMG5, SMG7* in nonsense-mediated mRNA decay pathway; *EXOSC2, EXOSC5, EXOSC6* in RNA exosome; *CSTF3, CPSF2, CPSF6, NUDT21, XRN2* for 3' polyadenylation; *RNMT, NCBP1* related to 5' RNA capping) **(Fig 2i-j)**. For example, the knockdown of two critical regulators in the microRNA (miRNA) pathway[43] (*i.e., DROSHA* and *DICER1*[44,45], **Extended Data Fig 11a**) resulted in a group of highly overlapped DEGs(**Extended Data Fig 11b**). These DEGs were upregulated through decreased degradation (*e.g.,* miRNA-mediated silencing complex (RISC) components: *TNRC6A and TNRC6B*[46]) or increased transcription (*e.g.,* miRNA host genes: *MIR181A1HG*[47], *FTX*[48]; genes involved in miRNA biogenesis: *DDX3X*[49]) **(Fig 2k-m, Extended Data Fig 11c, Supplementary Table 11)**. To

explore the underlying regulatory mechanisms, we examined the gene-specific binding patterns of *Ago2*,
one of the core components in RISC for targeted mRNA binding and degradation[50]. Indeed, *Ago2* binding
was strongly enriched in the 5' and 3' untranslated regions (UTR) of the genes with reduced degradation,
but not in genes with upregulated synthesis **(Fig 2n)**, consistent with prior reports that miRNA induces
targeted RNA degradation and translation repression mainly through binding to the UTR[44,51]. The analysis
further demonstrates the unique capacity of *PerturbSci-Kinetics* for deciphering the regulatory
mechanisms (degradation vs. transcription) involved in gene expression changes upon genetic
perturbations.
228
Lastly, to our knowledge, the studies described here provided the first method to quantitatively
characterize the genome-wide mRNA kinetic rates (*e.g.,* synthesis and degradation rates) across hundreds
of genetic perturbations in a single experiment. We included the step-by-step protocols and the data
processing pipeline as supplementary files (**Supplementary file 1-4**) to facilitate the broad applications
of the technique. Our analysis illustrates the advantages of *PerturbSci-Kinetics* over conventional assays
that solely profile gene expression changes. By capturing three layers of readout (*e.g.,* whole, nascent
transcriptome, and sgRNA identify) at the single-cell resolution, *PerturbSci-Kinetics* uniquely enables us
to dissect the critical regulators of gene-specific transcription, processing, and degradation in a massive-
parallel manner. Finally, *PerturbSci-Kinetics* is built on the recently developed *EasySci-RNA*[10] and can be
readily scaled up to profiling genome-wide perturbations (*e.g.,* 10,000s genes or cis-regulatory elements)
across millions of single cells, thus enabling the systematic characterization of cell-type-specific gene
regulatory network at unprecedented scale and resolution.

Fig 2. Characterizing the impact of genetic perturbations on gene-specific transcriptional and degradation dynamics with *PerturbSci-Kinetics*. a. Scheme of the experimental design of the *PerturbSci-Kinetics* screen. The main steps are described in the text. **b.** The scatter plot shows the correlation between perturbation-associated cell count (PerturbSci-Kinetics) and sgRNA read counts (bulk screen). **c.** Box plot showing the log2 transformed fold change of gene expression, synthesis rates, and degradation rates of target genes across perturbations in comparison with the NTC cells. **d.** UMAP visualization of genetic perturbations profiled by *PerturbSci-Kinetics*. We aggregated single-cell transcriptomes in each perturbation, followed by dimension reduction using PCA and visualization using

UMAP. Population classes, the functional categories of genes targeted in different perturbations. **e-h.** Scatter plots showing the extent and the significance of changes on the distributions of global synthesis (e), degradation (f), nascent exonic reads ratio (g), and mitochondrial transcriptome turnover (h) upon perturbations compared to NTC cells. The fold changes were calculated by dividing the median values of each perturbation with that of NTC cells and were log2 transformed. **i.** Scatter plot showing the number of synthesis/degradation-regulated DEGs from different perturbations. nDEGs: the number of DEGs. **j.** Top20 perturbations ordered by the number of degradation-regulated DEGs. Synthesis only, DEGs with significant changes in synthesis rates. Degradation only, DEGs with significant changes in degradation rates. Synthesis+degradation, DEGs with significant changes in both synthesis and degradation rates. **k-l.** Venn diagrams showing the number of merged DEGs with significantly enhanced synthesis (k) or impaired degradation (l) between *DROSHA* and *DICER1*. Based on statistical test results, merged DEGs of *DROSHA* and *DICER1* perturbations were classified into synthesis-regulated genes (*i.e.,* the upregulation of these genes was mainly driven by increased synthesis rates) and degradation-regulated genes (*i.e.,* the upregulation of these genes were mainly driven by reduced degradation rates). Merged DEGs with p-value <= 0.05 on synthesis increase/degradation decrease in at least one perturbation were included in the diagram, in which genes with p-value < 0.1 on synthesis increase/degradation decrease in both perturbations were regarded as shared hits between two perturbations. **m.** Heatmaps showing the steady-state expression, synthesis and degradation rate changes of genes sharing the same regulatory mechanism upon *DROSHA* and *DICER1* knockdown as shown in k, l. Tiles of each row were colored by fold changes of values of perturbations relative to NTC. **n.** Line plot showing the *Ago2* binding patterns on the transcript regions of protein-coding genes in Figure 2n and 2o. The transcript regions of genes were assembled by merging all exons, and were divided into 5'UTR, coding sequence (CDS), and 3'UTR based on coordinates of the 5' most start codon and the 3' most stop codon. Single-base coverage of *Ago2* eCLIP on each gene was calculated, binned, and scaled to 0-1. After merging and averaging scaled binned coverage of genes in the same group together, the lowest coverage value in the CDS was used to scale the averaged merged coverage again to visualize the *Ago2*/RISC binding pattern.

Endnotes

Acknowledgments: We thank all members of the Cao lab for helpful discussions and feedback. We thank Dr. R. Satija (New York Genome Center) for insightful feedback related to this work. We thank the Tissue Culture facility of the University of California, Berkeley for the 3T3 cell line, and the Scott Keeney Lab at Memorial Sloan Kettering Cancer Center for the HEK293 cell line. We thank members of the Rockefeller University Flow Cytometry Resource Center and the Rockefeller University Genomics Resource Center for their extensive help with FACS sorting and sequencing experiments. We also thank members of the Information Technology and High-Performance Computing team at Rockefeller University, especially J. Banfelder and B. Jayaraman for the great support. We acknowledge that the research resulting in this publication was supported, in part, by The G. Harold and Leila Y. Mathers Charitable Foundation.

Funding: This work was funded by grants from the NIH (1DP2HG012522, 1R01AG076932 and RM1HG011014) and the Mathers Foundation to J.C..

Author contributions: J.C. and W.Z. conceptualized and supervised the project. Z.X. performed experiments, including technique development and optimization, with input from J.L.. Z.X. performed computational analyses, with input from A.S.. J.C., W.Z., and Z.X. wrote the manuscript with input and biological insight from all co-authors.

Competing interests statement: J.C., W.Z., and Z.X. are inventors on pending patent applications related to *PerturbSci-Kinetics*. Other authors declare no competing interests.

Supplementary Figures:

Extended Data Fig. 1. Scheme of plasmids and experiment procedures of *PerturbSci*. a. The vector

system used in *PerturbSci* for dCas9 and sgRNA expression. The expression of the enhanced CRISPRi

silencer dCas9-KRAB-MeCP2[11] was controlled by the tetracycline responsive (Tet-on) promoter. A GFP

sequence was added to the original CROP-seq-opti plasmid[9] as an indicator of successful sgRNA

transduction and for the lentivirus titer measurement. **b**. The library preparation scheme and the final

library structures of *PerturbSci*, including a scalable combinatorial indexing strategy with direct sgRNA

capture and enrichment that reduced the library preparation cost, enhanced the sensitivity of the sgRNA

capture compared to the original CROP-seq[8], and avoided the extensive barcodes swapping detected in

Perturb-seq[9].

Extended Data Fig. 2. Representative optimizations on sgRNA enrichment of *PerturbSci*. a. Multiple
RT primers targeting different sgRNA scaffold regions were mixed with polyT primers respectively and
were used in our test experiment for targeted enrichment of sgRNA (RT primer 2-4 were modified from
primers used in Direct-capture Perturb-seq[52], CRISPR-sciATAC[53], and ECCITE-seq[54]). CB, cell
barcode. P_R1, partial TruSeq read1 sequence. **b-c.** A 96-well plate was divided into 4 parts and RT was performed

using different combinations of sgRNA capture primers and shortdT primers. After ligation, cells were
mixed and redistributed for SSS. We tested the capture efficiency of sgRNA by different RT primers in
PerturbSci using "Direct PCR" and tested the efficiency of by-product removal by "sgRNA-only PCR"
(Scheme shown in b) followed by gel electrophoresis for analyzing the PCR product (c). Crosses in b,
potential Tn5 tagmentation sites. As shown in c, sgRNA primer 2 and 3 yielded strongest amplification
signals following PCR, while primer1 and 4 recovered weak signals. In addition, tagmentation removed
large by-products generated potentially from polyT priming (as shown in b). **d.** We tested different
conditions in post-multiplex PCR purification to obtain the input for the sgRNA enrichment PCR that
could maximize the recovery of the sgRNA library. Left lane: 0.7x-1.5x double-size AMPURE beads
purification followed by the sgRNA enrichment PCR reaction. Middle lane: 0.8x-1.2x AMPURE beads
purification followed by the sgRNA enrichment PCR reaction. Right lane: Gel extraction on multiplex
PCR product within 175-275 bp range followed by the sgRNA enrichment PCR reaction. The recovered
sgRNA libraries generated from gRNA primer2 and 3 were marked on the gel image. Based on the result,
the sgRNA primer2 and the 0.8-1.2x AMPURE beads purification condition yielded the best performance.
e. A representative gel image of the final libraries of *PerturbSci*, including the sgRNA library (Lane 1)
and the whole transcriptome library (Lane 2). **f-i.** We tested different concentrations of sgRNA RT primers
in the PerturbSci experiment using 3T3-L1-CRISPRi cells transduced with either sgFto and sgNTC. The
box plots show the number of unique sgRNA transcripts (f) or mRNA transcripts (g) detected per cell, the
cell recovery rate (h) and sgRNA capture purity (i) across different sgRNA RT primer concentrations. **j-**
k. We performed *PerturbSci* experiment with 3T3-L1-CRISPRi cells transduced with sgFto and sgNTC
in a pooled or separate manner. The box plots show the number of unique sgRNA transcripts detected per
cell (j) and sgRNA capture purity (k) across the two conditions. **l.** Scatter plot showing the correlation
between log2-transformed aggregated gene expression profiled by *PerturbSci* and *EasySci*[10] in the mouse
3T3-L1-CRISPRi cell line.

Extended Data Fig. 3. Representative optimizations on fixation conditions of *PerturbSci-Kinetics*. We aimed to search for an optimal fixation condition that can i) minimize the cell loss during the fixation and chemical conversion, ii) reduce the RNA cross-contamination, iii) be compatible with *in-situ* combinatorial indexing of cellular transcriptomes. **a-c.** We tested different cell fixation conditions on HEK293-idCas9 cells followed by *PerturbSci* profiling and quantified the fraction of cells that were assigned to different groups (a), the number of unique sgRNA (b) and mRNAs (c) detected per cell. PFA fixation conditions at the room temperature (RT) were too strong to recover sufficient signals. FA fixation at 4°C yielded higher total UMI counts but showed stronger cross-contamination, indicating that under 4°C it was a milder fixative compared to 4% PFA. **d.** Scatter plot showing the number of unique mRNA transcripts recovered from human HEK293-idCas9 cells and mouse 3T3 cells in a *PerturbSci* experiment. The human and mouse cell mixture was fixed by 4°C PFA+BS3 condition. Reads were aligned to a combined human-mouse reference genome and the species origins of single cells were identified by the fraction of species-specific read counts. The clear separation of cells from two species indicated the good

compatibility of this fixation condition with *PerturbSci*. **e-f.** Dot plots showing the relative recovery rate (with standard error of the mean) of HEK293-idCas9 cells in different fixation conditions (n = 4) following HCl permeabilization (d) and chemical conversion (e). All values were normalized by the standard condition used in sci-fate (PFA fixation)[14]. **g.** Box plot showing the number of unique transcripts detected per cell with or without chemical conversion. Fixation conditions included in the plots: 4°C PFA+BS3: cells were fixed with 4% PFA in PBS for 15 minutes, and were further fixed by 2mM BS3 during and after Triton-X100 permeabilization (**Methods**). 4°C FA+BS3: cells were fixed with 1% Formaldehyde (FA) in PBS for 10 minutes, and were further fixed by 2mM BS3 during and after Triton-X100 permeabilization. 4°C FA: cells were only fixed once with 1% Formaldehyde (FA) in PBS for 10 minutes. 4°C PFA: cells were only fixed once with 4% PFA in PBS for 15 minutes as sci-fate[14].

Extended Data Fig. 4. Optimization of the computational pipeline for nascent reads calling. a-c. Bar plots showing the normalized mismatch rates of all 12 mismatch types detected in unconverted cells (a), converted cells (b), and the original sci-fate A549 dataset[14] (c) at different positions of the reads using the original sci-fate mutation calling pipeline[14]. **d-f.** Bar plots showing the normalized mismatch rates of all 12 mismatch types detected in unconverted cells (d), converted cells (e), and the original sci-fate A549 dataset[14] (f) at different positions of the reads using the updated mutation calling pipeline. Considering the different sequencing lengths between the present dataset and sci-fate, the Read2 from sci-fate were trimmed to the same length as the present dataset before processing. Compared to the original pipeline, the updated pipeline further filtered the mismatch based on the CIGAR string and only mismatches with "CIGAR = M" were kept. As shown in the result, this optimized pipeline efficiently removed the unaligned mismatches enriched at the 5' and 3' end of reads. Normalized mismatch rates in each bin, the percentage of each type of mismatch in all sequencing bases within the bin.

Extended Data Fig. 5. Validation of the CRISPRi performance. Strongly reduced *IGF-1R* mRNA and protein levels in HEK293-idCas9 cells after Dox induction were further validated by **a.** RT-qPCR (n=4. ****, p-value < 1e-4, Tukey's test after ANOVA) and **b.** flow cytometry. Isotype, isotype control. αIGFIR, anti-IGF1R.

Extended Data Fig. 6. The changes in sgRNA abundance are consistent between replicates and previously published data. a. Heatmap showing the overall Pearson correlations of normalized sgRNA read counts between the plasmid library and bulk screen replicates at different sampling times. For each library, read counts of sgRNAs were normalized first by the sum of total counts and then by the counts of sgNTC. **b**. Box plot showing the reproducible trends of deletion upon CRISPRi between the present study and a prior report[29]. We calculated the fraction changes (After vs. before the CRISPRi induction) of sgRNAs for each gene, followed by log2 transformation. **c**. Bar plot showing the different extent of deletion of cells receiving sgRNAs targeting genes in different categories in the bulk screen. The knockdown on genes with higher essentiality caused stronger cell growth arrest.

Extended Data Fig. 7. Quality control and sgRNA filtering on the *PerturbSci-Kinetics* library. a. We

filtered out cells assigned to multiple gRNAs based on two criteria: the cell is defined as a sgRNA singlet

if the most abundant sgRNA in the cell took >= 60% of total sgRNA counts and was at least 3-fold of the

second most abundant sgRNA. The histogram shows the fraction distribution of the most abundant sgRNA

in assigned singlets (78%) and doublet cells (22%). **b-e**. Dotplots showing the expression fold changes of

target genes upon CRISPRi induction compared to NTC. Each dot represents a sgRNA. Fold change <

0.6 was used for sgRNA filtering, and target genes with 3, 2, 1, 0 on-target sgRNA(s) were shown in b-e,

respectively. FC, fold change.

Extended Data Fig. 8. A systematic view of the effects of perturbations on global synthesis rates, global degradation rates, exonic reads ratio, and mitochondrial turnover rates. For each gene category, we calculated the fraction of genetic perturbations associated with significant changes in global synthesis rates (a), global degradation rates (b), fractions of exonic reads in the nascent transcriptome (c), and mitochondrial RNA turnover rates (d). Overall global transcription could be affected by more genes than degradation. Perturbation on essential genes, such as DNA replication genes, could affect both global synthesis and degradation. Perturbations on chromatin remodelers only specifically impaired the global synthesis rates but not the degradation rates, supporting the established theory that gene expression is

regulated by chromatin folding. In addition to the enrichment of genes in transcription, spliceosome and

mRNA surveillance, perturbation on OXPHOS genes and metabolism-related genes also affected the RNA

processing, consistent with the fact that 5' capping, 3' polyadenylation, and RNA splicing are highly

energy-dependent processes. That knockdown of OXPHOS genes and metabolism-related genes could

reduce the mitochondrial transcriptome dynamics and also supported the complex feedback mechanisms

between energy metabolism and mitochondrial transcription[55].

Extended Data Fig. 9. *PerturbSci-kinetics* **identified LRPPRC as the master regulator of mitochondrial RNA dynamics. a.** Heatmap showing the relative fold changes of gene expression, synthesis and degradation rates of mitochondrial protein-coding genes upon NDUFS2, CYC1, BCS1L and LRPPRC knockdown compared to NTC cells. Perturbation on genes encoding electron transport chain components resulted in stable steady-state expression with impaired turnover. However, LRPPRC knockdown significantly disrupted the mitochondrial transcriptome dynamics by inhibiting the synthesis of almost all mitochondrial protein-coding genes and promoting the degradation of multiple genes including *MT-ND6, MT-CO1, MT-ATP8, MT-ND4, MT-CYB* and *MT-ATP6*. **b.** The heatmap on the left showed the mitochondrial protein-coding gene expression changes between wild-type and *LRPPRC*-knockout mice heart tissue, as reported by Siira, S.J., et al.[37]. The heatmap on the right showed the extent of the mRNA secondary structure increase upon *Lrpprc* knockdown observed in the published study[37], which positively correlated with the elevated degradation rates of genes detected in our study (Pearson correlation r = 0.708, p-value = 6.8e-3). The result further validated the mRNA-stabilizing mechanism of *Lrpprc*. **c.** Bar plot showing the fraction of genes regulated by synthesis, degradation or both in mitochondrial-encoded and nuclear-encoded DEGs. **d.** Bar plot showing the enrichment of *ATF4/CEBPG*

motifs at promoter regions of DEGs with or without significant synthesis changes. Nc DEGs w/o synth changes, Nuclear-encoded differentially expressed genes without synthesis changes. Nc DEGs w/ synth changes, Nuclear-encoded differentially expressed genes with synthesis changes. A large part of synthesis-regulated nuclear-encoded DEGs showed motif enrichment, suggesting the activation of an integrated stress response transcriptional program mediated by ATF4/CEBPG upon LRPPRC knockdown[41]. 5kb regions around transcription start sites of input genes were used for motif scanning and enrichment calculation using RcisTarget[56]. We identified two transcription factors (*ATF4* and *CEBPG*) that were i) significantly upregulated upon *LRPPRC* knockdown ii) significantly over-represented in the surroundings of the transcription start site of the synthesis-regulated nuclear-encoded DEGs (Normalized motif enrichment score of 16 for *ATF4* and 16.6 for *CEBPG*). **e.** The transcriptional regulatory network in *LRPPRC* perturbation inferred from our analysis. Notably, it was consistent with the prior study[41] that *ATF4* was regulated at both transcriptional and post-transcriptional levels.

Extended Data Fig. 10. *PerturbSci-Kinetics* **captured the synthesis/degradation rates of DEGs upon**

perturbations. a-d. Box plots showing the log2 transformed fold changes of synthesis or degradation

rates between perturbations and NTC cells for DE genes in four categories: synth up (DEGs with

significantly increased synthesis rates), synth down (DEGs with significantly decreased synthesis rates),

deg up (DEGs with significantly increased degradation rates), deg down (DEGs with significantly

decreased degradation rates).

Extended Data Fig. 11. The overview of the miRNA biogenesis pathway and perturbations on pathway members. a. Illustration of the canonical miRNA biogenesis pathway. After the transcription of miRNA host genes, the primary miRNA (pri-miRNA) forms into a hairpin and is processed by *Drosha*. Processed precursor miRNA (pre-miRNA) is transported to the cytoplasm by Exportin-5. The stem loop is cleaved by *Dicer1*, and one strand of the double-stranded short RNA is selected and loaded into the RISC for targeting mRNA[44]. **b.** Venn diagram showing the overlap of upregulated DEGs across perturbations on four genes encoding main members of the miRNA pathway. The knockdown of two critical RNases in this pathway (*i.e., DROSHA* and *DICER1*) resulted in significantly overlapped DEGs (p-value = 2.2e-16, Fisher's exact test). In contrast, *AGO2* knockdown resulted in more unique transcriptome features, and only 1 DEG (*PRTG*, identified to be mainly regulated by degradation and has been reported as a miRNA target[57]) overlapped with DEGs from *DROSHA* and *DICER1* knockdown, indicating the RNAi-independent roles of *AGO2*. Interestingly, *XPO5* knockdown showed no upregulated DEGs, which is consistent with a previous report in which *XPO5* silencing only minimally perturbed the miRNA biogenesis, indicating the existence of an alternative miRNA transportation pathway[45]. **c.** Bar plot showing the fraction of upregulated DEGs driven by synthesis changes and degradation changes upon *DROSHA*, *DICER1*, and *AGO2* perturbations. While *DROSHA* and *DICER1* knockdown resulted in increased synthesis and reduced degradation, *AGO2* knockdown only affected gene expression transcriptionally, which was consistent with the previous finding that *AGO2* knockdown resulted in a global increase of synthesis rates **(Fig 2e)**, and further supported its roles in nuclear transcription regulation[58–60]. As *Drosha* is upstream of *Dicer1* in the pathway, we indeed observed stronger effects of *DROSHA* knockdown than *DICER1* knockdown, which was supported by the previous study[45].

Materials and Methods:

Cell culture

The 3T3-L1-CRISPRi cell line was obtained from the Tissue Culture facility at the University of California, Berkeley. The HEK293 cell line was a gift from the Scott Keeney Lab at Memorial Sloan Kettering Cancer Center. The HEK293T cell line was obtained from ATCC (CRL-3216). All cells were maintained at 37 °C and 5% CO2 in high glucose DMEM medium supplemented with L-Glutamine and Sodium Pyruvate (Gibco 11995065) and 10% Fetal Bovine Serum (FBS; Sigma F4135). When generating a monoclonal cell line, the medium was supplemented with 1% Penicillin-Streptomycin (Gibco 15140163). In the screening experiment, sgRNA-transduced HEK293-idCas9 cells were cultured in high glucose DMEM medium supplemented with L-Glutamine (Gibco 11965092) and 10% FBS, following the induction of dCas9-KRAB-MeCP2 expression by 1ug/ml Dox (Sigma D5207),

Generation of monoclonal HEK293-idCas9 cell line

To generate HEK293 with Dox-inducible dCas9-KRAB-MeCP2 expression, the lentiviral plasmid Lenti-idCas9-KRAB-MeCP2-T2A-mCherry-Neo was constructed. A dCas9-KRAB-MeCP2-T2A insert was amplified from dCas9-KRAB-MeCP2 (Addgene #110821). A T2A-mCherry Gblock was synthesized by IDT. Gibson Assembly reaction (NEB E2611S) was performed at 50 °C with a mixture of Bsp119I-digested Lenti-Neo-iCas9 (Thermo FD0124; Addgene #85400), dCas9-KRAB-MeCP2-T2A amplicon, T2A-mCherry Gblock for 60 minutes to construct a dCas9-KRAB-MeCP2-T2A-mCherry plasmid. The reaction product was transformed into NEBstable competent cells (NEB C3040H), and colonies were inoculated and amplified in LB medium (Gibco 10855001) with 50ug/ml Sodium Ampicillin (Sigma A8351) at 37 °C overnight.

After plasmid extraction (QIAGEN No.27106) and sequencing validation, the plasmid was co-transfected with psPAX2 (Addgene #12260) and pMD2.G (Addgene #12259) into low-passage HEK293T cells in a 10cm dish using Polyjet (SignaGen SL100688) for 24 hours. Cells were gently washed twice with PBS, then cultured in a medium with 10mM Sodium Butyrate (Sigma TR-1008-G) for another 24 hours. The supernatant was collected, and cell debris was cleared by spinning down (5min, 1000xg) and passed through a 0.45 μm filter. The lentivirus was concentrated 10x by the Lenti-X concentrator (TaKaRa 631231), and the virus suspension was flash frozen by Liquid Nitrogen and was stored at -80 °C.

The lentivirus titer was determined by examining the ratio of mCherry+ cells after 24 hours of transduction

and 48 hours of Dox induction. Polybrene (Sigma TR-1003) at a final concentration of 8ug/ml was used

to enhance the transduction efficiency. Then HEK293 cells were counted and transduced with lentivirus

at MOI = 0.2 for 48 hours. Cells were treated with Dox for 48 hours, and the top 10% of cells with the

strongest mCherry fluorescence were sorted to each well of a 96-well plate containing 100ul medium.

After a 3-week expansion, monoclonal cells that survived were transferred to larger dishes for further

expansion. We picked the clone with inducible homogeneous strong mCherry expression and normal

morphology for the following experiment.

Gene Knockdown and efficacy examination

To simplify the lentiviral titer measurement, CROP-seq-opti-Puro-T2A-GFP was assembled by adding a

T2A-GFP downstream of Puromycin resistant protein coding sequence on the CROP-seq-opti plasmid

(Addgene #106280). Flanking MluI and CsiI digestion sites were added to the GFP Gblock (IDT) by PCR.

Both amplicon and CROP-seq-opti vector were digested using MluI (Thermo, FD0564) and CsiI (Thermo,

FD2114) at 37 °C for 30 minutes, and were ligated at room temperature for 20 minutes using the Blunt/TA

Ligase Master Mix (NEB M0367S). Transformation, clone amplification, and sequencing validation were

done as stated above.

Oligos corresponding to individual guides for ligation were ordered as standard DNA oligos from IDT

with the following design:

Plus strand: 5'-CACCG[20bp sgRNA plus strand sequence]-3'

Minus strand: 5'-AAAC[20bp sgRNA minus strand sequence]C-3'

Oligos were reconstituted into 100uM and were mixed and phosphorylated using T4 PNK (NEB M0201S)

by incubating at 37 °C for 30 minutes. The reaction was heated at 95 °C for 5 minutes and then ramped

down to 25 °C by -0.1 °C/second to anneal oligos into a double-stranded duplex. The CROP-seq-opti-

Puro-T2A-GFP was digested by Esp3I (NEB R0734L) at 37 °C for 30 minutes, then the linearized

backbone and the annealed duplex were ligated at room temperature for 20 minutes using the Blunt/TA

Ligase Master Mix (NEB M0367S). Transformation, clone amplification, sequencing validation,

lentivirus generation, and titer measurement were done as stated above.

For the mouse 3T3-L1-CRISPRi cells, they were counted and incubated with lentivirus inserted with either non-target control (NTC) sgRNA or sgRNA targeting a *Fto* gene, and 8ug/ml of Polybrene. For the human HEK293-idCas9 cells, they were counted and incubated with NTC sgRNA or sgRNA targeting an *IGF1R* gene, and 8ug/ml of Polybrene. Transduction was then performed at MOI = 0.2 for 48 hours. Based on the results of our puromycin titration experiments, sgRNA-transduced 3T3-L1-CRISPRi cells were selected by 2.5ug/ml Puromycin for 2 days and 2ug/ml Puromycin for 3 days, and sgRNA-transduced HEK293-idCas9 cells were selected by 1.5ug/ml Puromycin for 3 days and 1ug/ml Puromycin for 2 days.

As dCas9-BFP-KRAB was constitutively expressed in 3T3-L1-CRISPRi cells, the target gene started being silenced once sgRNA lentivirus was introduced. For HEK293-idCas9 cells, Dox treatment for a minimum of 72 hours was required before examining the knockdown effect.

For RT-qPCR validation, primers targeting IGF1R were selected from PrimerBank (https://pga.mgh.harvard.edu/primerbank/) and were synthesized from IDT. Total RNA in 1e6 cells of each sample was extracted using the RNeasy Mini kit (QIAGEN 74104) and the concentration was measured by Nanodrop. 1ug total RNA was then reverse-transcribed into the first strand cDNA by SuperScript VILO Master Mix (Thermo 11755050). PowerTrack SYBR Green Master Mix (Thermo A46109) was used for RT-qPCR following the manufacturer's instructions.

For flow cytometry validation, 1e6 cells of each sample were harvested and resuspended in 100ul of PBS-0.1% sodium azide-2% FBS. BV421 Mouse Anti-Human CD221 (BD 565966) and BV421 Mouse IgG1 k Isotype Control (BD 562438) at the final concentration of 10 ug/ml were added, and reactions were incubated at 4 °C in the dark with rotation for 30 minutes. Cells were then washed twice using PBS-0.1% sodium azide-2% FBS, and fluorescence signals were recorded.

Construction of pooled sgRNA library

Genes of interest were selected manually, considering their functions and expression levels in HEK293 cells. The sgRNA sequences targeting genes of interest with the best performances were obtained from an established optimized sgRNA library (only sgRNA set A is considered)[29]. Finally, 684 sgRNAs targeting 228 genes (3 sgRNAs/gene) and 15 non-targeting controls were included in the present study.

The single-stranded sgRNA library was synthesized in a pooled manner by IDT in the following format:

5'-GGCTTTATATATCTTGTGGAAAGGACGAAACACCG[20bp   sgRNA   plus   strand

sequence]GTTTAAGAGCTATGCTGGAAACAGCATAGCAAGTT-3'

100ng of oligo pool was amplified by PCR using primers targeting 5' homology arm (HA) and 3' HA

with limited cycles (x12) to avoid introducing amplification biases. The PCR product was purified, and

double-stranded library amplicons were extracted by DNA electrophoresis and gel extraction. Then the

insert was cloned into Esp3I-digested CROP-seq-opti-Puro-T2A-GFP by Gibson Assembly (50 °C for 60

minutes). In parallel, a control Gibson Assembly reaction containing only the backbone was set. Both

reactions were cleaned up by 0.75x AMPURE beads (Beckman Coulter A63882) and eluted in 5uL EB

buffer (QIAGEN 19086), then were transformed into Endura Electrocompetent Cells (Lucigen, 602422)

by electroporation (Gene Pulser Xcell Electroporation System, Bio-Rad, 1652662). After 1 hour of

recovery at 250rpm, 37 °C, each reaction was spread onto an in-house 245 mm Square agarose plate

(Corning, 431111) with 100ug/ml of Carbenicillin (Thermo, 10177012) and was then grown at 32 °C for

13 hours to minimize potential recombination and growth biases. All colonies from each reaction were

scraped from the plate and the CROP-seq-opti-Puro-T2A-GFP-sgRNA plasmid library was extracted

using ZymoPURE II Plasmid Midiprep Kit (Zymo, D4200). The lentiviral library was generated as stated

above with extended virus production time. The step-by-step protocol is included in the supplementary

materials.

The pooled *PerturbSci-Kinetics* screen experiment

For each replicate, 7e6 uninduced HEK293-idCas9 cells were seeded. After 12 hours, two replicates were

transduced at MOI=0.1 (1000x coverage/sgRNA) and another two replicates were transduced at MOI=0.2

(2000x coverage/sgRNA) with 8ug/ml of Polybrene for 24 hours. Then we replaced the culture medium

with the virus-free medium and culture cells for another 24 hours. Transduced cells were selected by

1.5ug/ml of Puromycin for 3 days and 1ug/ml of Puromycin for 2 days. During the selection, we passed

cells every 2 or 3 days to ensure at least 1000x coverage. At the end of the drug selection, we harvested

1.4e6 cells in each replicate (2000x coverage/sgRNA) as day0 samples of the bulk screen and pellet down

at 500xg, 4 °C for 5 minutes. Cell pellets were stored at -80 °C for genomic DNA extraction later. Then

the dCas9-KRAB-MeCP2 expression was induced by adding Dox at the final concentration of 1ug/ml,

and L-glutamine+, sodium pyruvate-, high glucose DMEM was used to sensitize cells to perturbations on

energy metabolism genes. Cells were cultured in this condition for additional 7 days and were passed

every other day with 4000x coverage/sgRNA. On day7, 6ml of the original media from each plate was

mixed with 6uL of 200mM 4sU (Sigma T4509-25MG) dissolved in DMSO (VWR 97063-136) and was

put back for nascent RNA metabolic labeling. After 2 hours of treatment, 1.4e6 cells in each replicate

were harvested as day7 samples of the bulk screen, and the rest of the cells were fixed and stored for

single-cell *PerturbSci-Kinetics* profiling (see the next section).

Genomic DNA of bulk screen samples was extracted using Quick-DNA Miniprep Plus Kit (Zymo,

D4068T) following the manufacturer's instructions and quantified by Nanodrop. All genomic DNA was

used for PCR to ensure coverage. The primer targeting the U6 promoter region with P5-i5-Read1 overhang

and the primer targeting the sgRNA scaffold region with P7-i7-Read2 overhang was used for generating

the bulk screen libraries for sequencing.

Library preparation for the *PerturbSci-Kinetics*

After trypsinization, cells in each 10cm dish were collected into a 15ml falcon tube and kept on ice. Cells

were spun down at 300xg for 5 minutes (4 °C) and washed once in 3ml ice-cold PBS. Cells were fixed

with 5ml ice-cold 4% Paraformaldehyde (PFA) in PBS (Santa Cruz Biotechnology sc-281692) for 15

minutes on ice. PFA was then quenched by adding 250ul 2.5M Glycine (Sigma 50046-50G), and cells

were pelleted at 500xg for 5 minutes (4 °C). Fixed cells were washed once with 1ml PBSR (PBS, 0.%

SUPERase In (Thermo AM2696), and 10mM dithiothreitol (DTT; Thermo R0861)), and were then

resuspended, permeabilized, and further fixed in 1ml PBSR-triton-BS3 (PBS, 0.1% SUPERase In, 0.2%

Triton-X100 (Sigma X100-500ML), 2mM bis(sulfosuccinimidyl)suberate (BS3; Thermo, PG82083),

10mM DTT) for 5 minutes. Additional 4ml of PBS-BS3 (PBS, 2mM BS3, 10mM DTT) was then added

to dilute Triton-X100 while keeping the concentration of BS3, and cells were incubated on ice for 15

minutes. Cells were pelleted at 500xg, 4 °C for 5 minutes and resuspended in 500ul nuclease-free water

(Corning 46-000-CM) supplemented with 0.1% SUPERase In and 10mM DTT. 3ml of 0.05N HCl (Fisher

Chemical, SA54-1) was added for further permeabilization. After 3 minutes of incubation on ice, 3.5ml

Tris-HCl, pH 8.0 (Thermo 15568025), and 35ul of 10% Triton X-100 were added to each tube to neutralize

the HCl. After spinning down at 4 °C, 500xg for 5 minutes, cells were finally resuspended in 400ul PSB-

DTT at the concentration of ~2e6 cells/100ul (PBS, 1% SUPERase In, 1% BSA (NEB B90000S), 1mM

DTT), mixed with 10% DMSO, and were slow-frozen and stored in -80 °C.

The chemical conversion was performed before the library preparation. Cells were thawed with shaking

in the 37 °C water bath and spun down, then were washed once with 400ul PSB without DTT. Next, cells

were resuspended in 100ul PSB, mixed with 40ul Sodium Phosphate buffer (PH 8.0, 500mM), 40ul IAA

(100mM, Sigma I1149-5G), 20ul nuclease-free water, and 200ul DMSO with the order. The reaction was

incubated at 50 °C for 15 minutes and was quenched by adding 8ul 1M DTT. Then cells were washed

with PBS and were filtered through a 20um strainer (Pluriselect 43-10020-60). Cells were finally

resuspended in 100ul PSB.

For library preparation, a step-by-step protocol is included as a supplementary file.

Reads processing

For bulk screen libraries, bcl files were demultiplexed into fastq files based on index 7 barcodes. Reads

for each sample were further extracted by index 5 barcode matching. Then every read pair was matched

against two constant sequences (Read1: 11-25bp, Read2: 11-25bp) to remove reads generated from the

PCR by-product. For all matching steps, a maximum of 1 mismatch was allowed. Finally, sgRNA

sequences were extracted from filtered read pairs (at 26-45bp of R1), assigned to sgRNA identities with

no mismatch allowed, and read counts matrices at sgRNA and gene levels were quantified.

For *PerturbSci-Kinetics* transcriptome reads processing and whole-transcriptome/nascent transcriptome

gene counting, the pipeline was developed based on *EasySci*[10] and *Sci-fate*[14] with minor modifications.

After demultiplexing on index 7, Read1 were matched against a constant sequence on the sgRNA capture

primer to remove unspecific priming, and cell barcodes and UMI sequences sequenced in Read1 were

added to the headers of the fastq files of Read2, which were retained for further processing. After potential

polyA sequences and low-quality bases were trimmed from Read2 by Trim Galore[61], reads were aligned

to a customized reference genome consisting of a complete hg38 reference genome and the dCas9-KRAB-

MeCP2 sequence from Lenti-idCas9-KRAB-MECP2-T2A-mCherry-Neo using STAR[62]. Unmapped

reads and reads with mapping score < 30 were filtered by samtools[63]. Then deduplication at the single-

cell level was performed based on the UMI sequences and the alignment location, and retained reads were

split into SAM files per cell. These single-cell sam files were converted into alignment tsv files using the sam2tsv function in jvarkit[64]. Only reads with FLAG values of 0 or 16 and high-quality mismatches with QUAL scores > 45 and CIGAR of M in them were maintained. Mutations were further filtered against background SNPs called by VarScan using our in-house *EasySci* data on HEK293 cells. Reads in which at least 30% of mutations were T to C mismatches were identified as nascent reads, and the list of reads were extracted from single-cell whole transcriptome sam files by Picard[65]. Finally, single-cell whole transcriptome gene x cell count matrix and nascent transcriptome gene x cell count matrix were constructed by assigning reads to genes if the aligned coordinates overlapped with the gene locations on the genome. At the same time, single cell exonic/intronic read numbers were also counted by checking whether reads were mapped to the exonic or the intronic regions of genes. To quantify dCas9-KRAB-MECP2 expression, a customized gtf file consisting of the complete hg38 genomic annotations and additional annotations for dCas9 was used in this step.

Read1 and read2 of *PerturbSci-Kinetics* sgRNA libraries were matched against constant sequences respectively, allowing a maximum of 1 mismatch. For each filtered read pair, cell barcode, sgRNA sequence, and UMI were extracted from designed positions. Extracted sgRNA sequences with a maximum of 1 mismatch from the sgRNA library were accepted and corrected, and the corresponding UMI was used for deduplication. De-duplication was performed by collapsing identical UMI sequences of each individual corrected sgRNA under a unique cell barcode. Cells with overall sgRNA UMI counts higher than 10 were maintained and the sgRNA x cell count matrix was constructed.

Bulk screen sgRNA counts analysis

For each bulk screen library, read counts of sgRNAs were normalized first by the sum of total counts to remove the biases from sequencing depth, and then the abundance of each sgRNA relative to the sum of sgNTC was calculated, assuming the NTC cells had no selection pressure during the screen. The Pearson correlations across replicates were calculated based on the relative abundances. Then the fraction changes (After vs. before the CRISPRi induction) of sgRNAs were calculated within each replicate, and the mean fold changes across replicates were log2 transformed. The raw counts of another external bulk CRISPRi screen dataset[29] was processed as stated above and the log2 mean relative abundance was compared to the current study.

sgRNA singlets identification and off-target sgRNA removal

In the cell mixture experiments, cells with at least 200 whole transcriptome UMIs and 200 genes detected, and unannotated reads ratio < 40% were kept. If the count of the most abundant sgRNA was at least 3-fold of the second most abundant sgRNA within this single cell, then this cell was identified as a sgRNA singlet.

In the screen dataset, cells with at least 300 whole transcriptome UMIs and 200 genes detected, and unannotated reads ratio < 40% were kept. sgRNA identities of cells were assigned and doublets were removed based on the following criteria: the cell is assigned to a single sgRNA if the most abundant sgRNA in the cell took >= 60% of total sgRNA counts and was at least 3-fold of the second most abundant sgRNA. Then whole transcriptomes and sgRNA profiles of single cells were integrated with the matched nascent transcriptomes.

Target genes with the number of cells perturbed >= 50 were kept for further filtering. The knockdown efficiency was calculated at the individual sgRNA level to remove potential off-target or inefficient sgRNAs: whole transcriptome counts of all cells receiving the same sgRNA were merged, normalized by the total counts, and scaled using 1e6 as the scale factor, then the fold changes of the target gene expressions were calculated by comparing the normalized expression levels between corresponding perturbations and NTC. sgRNAs with >= 40% of target gene expression reduction relative to NTC were regarded as "effective sgRNAs", and singlets receiving these sgRNAs were kept as "on-target cells". Downstream analyses were done at the target gene level by analyzing all cells receiving different sgRNAs targeting the same gene together.

Gene Ontology analysis of genes with high or low nascent reads ratio

To validate the specificity of 4sU labeling and the computational identification of nascent reads, we identified features of gene groups with different turnover rates. Single cells were split into nascent transcriptomes and pre-existing transcriptomes, and were loaded into Seurat[30]. Nascent transcriptomes and pre-existing transcriptomes were normalized, scaled independently, and DEGs between the two groups were identified by FindMarkers function[30] with default parameters. Then GO enrichment analyses were performed using ClusterProfiler[66] on upregulated genes (genes with significantly higher fraction of

738 nascent counts, FDR of 0.05) and downregulated genes (genes with significantly lower fraction of nascent
739 counts, FDR of 0.05) respectively.
740

UMAP embedding on pseudo-cells
The count matrix of the "on-target" cells described above was loaded into Seurat[30], and DEGs of each
perturbation (compared to NTC) were retrieved by FindMarkers function[30] with default parameters. Cells
from perturbations with over one DEGs (by FindMarkers function[30]) were selected. We also included cells
from genetic perturbations involved in similar pathways of the top perturbations. The fold changes of the
normalized gene expression between perturbations and NTC were calculated, and were binned based on
the gene-specific expression levels in NTC. The top 3% of genes showing the highest fold changes within
each bin were selected and merged as features for Principal Component Analysis (PCA). The top 9 PCs
were used as input for Uniform Manifold Approximation and Projection (UMAP) embedding (min.dist =
0.3, n.neighbors = 10).
751

Differential expression analysis
Pairwise differential expression analyses between each perturbation and NTC cells were performed by the
differentialGeneTest() function of Monocle 2[67]. To identify DEGs with rate changes, we selected
significant hits (FDR of 5%, likelihood) with a >= 1.5-fold expression difference and counts per million
(CPM) >= 5 in at least one of the tested cell pairs. To showcase LRPPRC and miRNA pathway
perturbations, more stringent criteria were used to obtain DEGs with high confidence: significant hits
(FDR of 5%, likelihood) with a >= 1.5-fold expression difference and CPM >= 50 in at least one of the
tested cell pairs were kept.
760

761 **Synthesis and degradation rates calculation**
762 After the induction of CRISPRi for 7 days, we assumed new transcriptomic steady states had been
763 established at the perturbation level before the 4sU labeling, and the labeling didn't disturb these new
764 transcriptomic steady states. The following RNA dynamics differential equation is used for synthesis and
765 degradation rates calculation similar to the previous study[31]:

$$\frac{d(R)}{d(t)} = \alpha - R \cdot \beta$$

767 In which R is the mRNA abundance of each gene, α is the synthesis rate of this gene, and β is the

768 degradation rate of this gene. Since the RNA synthesis follows the zero-order kinetics and RNA

769 degradation follows the first-order kinetics in cells, $\frac{d(R)}{d(t)}$ is determined by α and $R \cdot \beta$.

770 As steady states had been established, the mRNA level of each gene didn't change. We can get:

$$\frac{d(R)}{d(t)} = 0$$

$$R = \frac{\alpha}{\beta}$$

773 Under the assumption that the labeling efficiency was 100%, all nascent RNA were labeled during the

774 4sU incubation, and pre-existing RNA would only degrade. So, for nascent RNA (R_n), $R_n(t = 0) = 0$

775 and $\alpha_n = \alpha$. For pre-existing RNA (R_p), $R_p(t = 0) = R = \alpha/\beta$ and $\alpha_p = 0$. Based on these boundary

776 conditions, we could further solve the differential equation above on nascent RNA and pre-existing RNA

777 of each gene.

$$\begin{cases} R_n = \frac{\alpha}{\beta}(1 - e^{-\beta \cdot t}) \\ R_p = \frac{\alpha}{\beta}e^{-\beta \cdot t} \end{cases}$$

As *PerturbSci-Kinetics* directly measured whole transcriptome gene expression levels and nascent

transcriptome gene expression levels, pre-existing gene expression levels could be obtained by subtracting

nascent transcriptome expressions from the whole transcriptome expressions. As cells were labeled by

4sU for 2 hours ($t = 2$), α and β of each gene could be calculated based on the equations above.

Due to the shallow sequencing and the sparsity of the single cell expression data, synthesis and degradation

rates of DEGs were calculated at the pseudo-cell level. We aggregated the expression profiles of all cells

with the same target gene knockdown, normalized the expressions of genes by the sum of gene counts,

and scaled the size of the total counts to 1e6. Synthesis and degradation rates of DEGs in the corresponding

perturbed pseudo-cell were calculated as stated above. DEGs with only nascent counts or degradation

counts were excluded from further examination since their rates couldn't be estimated.

To examine the significance of synthesis and degradation rate changes upon perturbation, regarding the

different cell sizes across different perturbations and NTC, which could affect the robustness of rate

calculation, randomization tests were adopted. Only perturbations with cell number $>= 50$ were examined.

For each DEG belonging to each perturbation, background distributions of the synthesis and degradation

rate were generated: a subset of cells with the same size as the corresponding perturbed cells was randomly

sampled from a mixed pool consisting of corresponding perturbed cells and NTC cells, then these cells

were aggregated into a background pseudo-cell, and synthesis and degradation rates of the gene for testing

were calculated as stated above, and the process was repeated for 500 times. Rates = 0 were assigned if

only nascent counts or degradation counts were sampled during the process (referred to as invalid

samplings), but only genes with less than 50 (10%) "invalid samplings" were kept for p-value calculation.

The two-sided empirical p-values for the synthesis and degradation rate changes were calculated

respectively by examining the occurrence of extreme values in background distributions compared to the

rates from perturbed pseudo-cell. Rate changes with p-value $<= 0.05$ were regarded as significant, and the

directions of the rate changes were determined by comparing the rates from the perturbed pseudo-cell with

the background mean values. The fold changes of rates for each significant gene were calculated as

follows: only NTC cells were sampled at the same size as perturbed cells and aggregated, and the

background rates were calculated at the pseudo-cell level. After resampling for 200 times, these gene-

specific rates were averaged. Fold changes of the rates = rates in perturbed pseudo-cell / mean rates from

the NTC-only background.

Global changes of key statistics upon perturbations

For global synthesis and degradation rate changes, considering the noise from lowly-expressed genes, we

selected top1000 highly-expressed genes from NTC cells, then calculated their synthesis rates and

degradation rates in NTC cells and all perturbations with cell number $>= 50$. KS tests were performed to

compare rate distributions between each perturbation and NTC cells.

During the reads processing, the number of reads aligned to exonic/intronic regions were counted at the

single cell level. Then the distributions of exonic reads percentage in nascent reads from single cells with

the same target gene knockdown and NTC cells were compared using the KS tests to identify genes

affecting RNA processing.

The ratio of nascent mitochondrial read counts to total mitochondrial read counts was calculated in each
single cell, and the distributions of the ratio from single cells with the same target gene knockdown and
NTC cells were compared using the KS tests to identify the master regulator of mitochondrial mRNA
dynamics.

In all global statistics examinations, the p-values were corrected from multiple comparisons, and
comparisons with FDR <= 0.05 were considered as significant. The median value from each perturbation
and NTC cells were compared to determine the direction of significant changes.

Ago2 **eCLIP coverage analysis**

To identify the potential different RISC binding patterns between synthesis/degradation-regulated DEGs
in *DROSHA* and *DICER1* perturbations, we reprocessed the raw data of *Ago2* eCLIP obtained from Hela
cells (two replicates, SRR7240709 and SRR7240710) from Zhang, K et, al[68]. Potential adapters at 3' ends
of reads were trimmed by Cutadapt[69], and the first 6-base UMI were extracted and attached to headers of
the reads. After STAR alignment[62] and samtools filtering[63], only uniquely aligned reads were kept and
deduplication was performed based on the UMI and mapping coordinates using UMI-tools[70]. Then bam
files were transformed to the single-base coverage by BEDtools[71]. The transcript regions of genes-of-
interest were reconstructed based on the hg38 genome annotation gtf file from GENCODE. Briefly, for
each gene, the exonic regions were extracted and were redivided into 5'UTR, CDS, and 3'UTR by the
5'most start codon and the 3'most stop codon annotated in the gtf. The *Ago2* binding coverages of these
designated regions were obtained by intersection and were binned. A small background (0.1/base) was
added for smoothing. The gene-specific signal in each bin was normalized by the number of bases in each
bin, and the binned coverage of each gene was scaled to be within 0-1. After aggregating scaled coverages
of synthesis/degradation-regulated genes respectively, the second scaling was performed to visualize the
relative enrichment of *Ago2* binding at UTR compared to the CDS: fold changes of the scaled binned
coverage relative to the lowest coverage value in the CDS along the aggregated transcript were calculated.

Data Availability

The data generated by this study can be downloaded in raw and processed forms from the NCBI Gene
Expression Omnibus (GSE218566, reviewers' token: itqlgacczrgxpmb).

Code Availability

The computation scripts for processing *PerturbSci-Kinetics* were included as supplementary files.

Supplementary Tables (provided as Microsoft Excel files)

Supplementary Table 1: Genes and sgRNAs included in the study. Each gene ("gene_symbol") has 3 sgRNAs, and they were named in the format "Gene_number" ("names"). sgRNA sequences were included in "sgRNA_seq". The "gene_class" is the functional category of each gene.

Supplementary Table 2: Raw sgRNA counts of the bulk screen samples collected at different time points. Read counts of each sgRNA ("sgRNA_name") from 4 replicates at day 0 and day 7 were included.

Supplementary Table 3: Relative sgRNA abundance fold changes between day 7 and day 0. The "Day7_vs_Day0_repX" is the fold changes of relative sgRNA abundance at the gene level (**Methods**).

Supplementary Table 4: Filtered differentially expressed genes between perturbations with cell number >= 50 and NTC. For each gene ("Gene_symbol"), the "perturbation" is the target gene in perturbed cells. The "DEGs_direction" is the direction of gene expression changes comparing perturbed cells to the NTC cells, and the "DEGs_FC" is the fold change of the gene expression changes comparing perturbed cells to the NTC cells. The "max.CPM.between.KD.NTC" and "min.CPM.between.KD.NTC" are the pseudobulk expression levels of the gene that showed higher and lower expression in perturbed cells or the NTC cells. The expression level was quantified by counts per million. The "qval" is the false discovery rate (one-sided likelihood ratio test with adjustment for multiple comparisons).

Supplementary Table 5: Information about perturbations that showed significant global synthesis rate changes. The "adj.p" is the false discovery rate adjusted for multiple comparisons. The "direction" is the direction of the changes on the global synthesis rates distributions comparing perturbed cells to the NTC cells, and the "KD_median/NTC_median" is the quantitative measurement of the changes. The "gene_class" is the functional category of target genes ("Perturbations").

Supplementary Table 6: Information about perturbations that showed significant global degradation rate changes. The "adj.p" is the false discovery rate adjusted for multiple comparisons. The "direction" is the direction of the changes on the global degradation rates distributions comparing perturbed cells to the

NTC cells, and the "KD_median/NTC_median" is the quantitative measurement of the changes. The

"gene_class" is the functional category of target genes ("Perturbations").

Supplementary Table 7: Information about perturbations that showed significant nascent exonic reads

ratio changes. The "adj.p" is the false discovery rate adjusted for multiple comparisons. The "direction"

is the direction of the changes on the nascent exonic reads ratio distributions comparing perturbed cells to

the NTC cells, and the "KD_median/NTC_median" is the quantitative measurement of the changes. The

"gene_class" is the functional category of target genes ("Perturbations").

Supplementary Table 8: Information about perturbations that showed significant mitochondrial RNA

turnover changes. The "adj.p" is the false discovery rate adjusted for multiple comparisons. The

"direction" is the direction of the changes in the distributions of mitochondrial nascent/total reads ratio

comparing perturbed cells to the NTC cells, and the "KD_median/NTC_median" is the quantitative

measurement of the changes. The "gene_class" is the functional category of target genes ("Perturbations").

Supplementary Table 9: Steady-state expression and synthesis/degradation dynamics of mitochondrial

genes upon *LRPPRC*, *NDUFS2*, *CYC1*, *BCS1L* perturbations. The "synth_rate", "synth_FC",

"synth_pval", "synth_direction" are the synthesis rate of the gene in the perturbed cells, the fold change

of the synthesis rate of the gene in the perturbed cells compared to the NTC cells, the significance of the

synthesis rate change, and the direction of the synthesis rate changes. The "deg_rate", "deg_FC",

"deg_pval", "deg_direction" are the degradation rate of the gene in the perturbed cells, the fold change of

the degradation rate of the gene in the perturbed cells compared to the NTC cells, the significance of the

degradation rate change, and the direction of the degradation rate changes. The "DEG_qval" and

"DEG_fold.change" are the multiple comparison-corrected FDR and the fold change of the steady-state

gene expression change in perturbed cells compared to the NTC cells.

Supplementary Table 10: Differentially expressed genes with significant synthesis and/or degradation

changes. The "perturbations" is the target gene of the perturbed cells, and the "Gene_symbols" is the

symbols of DEGs with significant synthesis and/or degradation rate changes in corresponding

perturbations. The type of significant rate change of each gene is included in the "Regulation_type". The

"Synth_deg_FC", the "Synth_deg_direction", and the "Synth_deg_pval" reflect the fold change, the

direction of the change, and the randomization test p-value of the rate indicated in the "Regulation_type".

"DEGs_FC", "DEGs_direction", and "max.expr.between.KD.NTC" are the fold changes of gene
expression, the direction of the change, and the maximum pseudobulk CPM between the corresponding
perturbation and the NTC cells.

Supplementary Table 11: Steady-state expression and synthesis/degradation dynamics of merged DEGs
upon *DROSHA* and *DICER1* perturbations. The "synth_rate", "synth_FC", "synth_pval",
"synth_direction" are the synthesis rate of the gene in the perturbed cells, the fold change of the synthesis
rate of the gene in the perturbed cells compared to the NTC cells, the significance of the synthesis rate
change, and the direction of the synthesis rate changes. The "deg_rate", "deg_FC", "deg_pval",
"deg_direction" are the degradation rate of the gene in the perturbed cells, the fold change of the
degradation rate of the gene in the perturbed cells compared to the NTC cells, the significance of the
degradation rate change, and the direction of the degradation rate changes. The "DEG_fold.change" and
"DEG_qval" are the fold change of the steady-state gene expression change in perturbed cells compared
to the NTC cells and the multiple comparison-corrected FDR.

Supplementary files

Supplementary file 1: Detailed experiment protocols for *PerturbSci-Kinetics*, including all materials and
equipment needed, step-by-step descriptions, and representative gel images.

Supplementary file 2: Primer sequences used in the *PerturbSci-Kinetics* experiment. The design
principles and sequences of the oligo pool library, bulk screen sequencing primer, shortdT RT primers,
sgRNA capture primers, ligation primers, sgRNA inner i7 primers, and P5/P7 primers were included. The
columns indicate the positions on the 96-well plate (Well positions), an identifier of the sequence (Names),
the full primer sequence (Sequences), and the barcode sequence (Barcodes).

Supplementary file 3: The overall costs for *PerturbSci-Kinetics* library preparation. Reagents used in
each step were included, and the costs were calculated based on the scale of the real experiment.

Supplementary file 4: Computational pipeline scripts and notes for processing *PerturbSci-Kinetics* data,
from sequencer-generated files to single-cell gene count matrix.

Reference:

1. Huang, H. *et al.* Recognition of RNA N6-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. *Nat. Cell Biol.* **20**, 285–295 (2018).

2. Kurosaki, T., Popp, M. W. & Maquat, L. E. Quality and quantity control of gene expression by nonsense-mediated mRNA decay. *Nat. Rev. Mol. Cell Biol.* **20**, 406–420 (2019).

3. Weskamp, K. & Barmada, S. J. RNA Degradation in Neurodegenerative Disease. *Adv Neurobiol* **20**, 103–142 (2018).

4. Jaitin, D. A. *et al.* Dissecting Immune Circuits by Linking CRISPR-Pooled Screens with Single-Cell RNA-Seq. *Cell* **167**, 1883–1896.e15 (2016).

5. Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867–1882.e21 (2016).

6. Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853–1866.e17 (2016).

7. Xie, S., Duan, J., Li, B., Zhou, P. & Hon, G. C. Multiplexed Engineering and Analysis of Combinatorial Enhancer Activity in Single Cells. *Mol. Cell* **66**, 285–299.e5 (2017).

8. Datlinger, P. *et al.* Pooled CRISPR screening with single-cell transcriptome readout. *Nat. Methods* **14**, 297–301 (2017).

9. Hill, A. J. *et al.* On the design of CRISPR-based single-cell molecular screens. *Nat. Methods* **15**, 271–274 (2018).

10. Sziraki, A. *et al.* A global view of aging and Alzheimer's pathogenesis-associated cell population dynamics and molecular signatures in the human and mouse brains. Preprint at https://doi.org/10.1101/2022.09.28.509825.

11. Yeo, N. C. *et al.* An enhanced CRISPR repressor for targeted mammalian gene regulation. *Nat. Methods* **15**, 611–616 (2018).

12. Erhard, F. *et al.* scSLAM-seq reveals core features of transcription dynamics in single cells. *Nature* **571**, 419–423 (2019).

13. Hendriks, G.-J. *et al.* NASC-seq monitors RNA synthesis in single cells. *Nat. Commun.* **10**, 3138 (2019).

14. Cao, J., Zhou, W., Steemers, F., Trapnell, C. & Shendure, J. Sci-fate characterizes the dynamics of gene expression in single cells. *Nat. Biotechnol.* **38**, 980–988 (2020).

15. Qiu, Q. *et al.* Massively parallel and time-resolved RNA sequencing in single cells with

scNT-seq. *Nat. Methods* **17**, 991–1001 (2020).

16.   Cleary, M. D., Meiering, C. D., Jan, E., Guymon, R. & Boothroyd, J. C. Biosynthetic labeling of RNA with uracil phosphoribosyltransferase allows cell-specific microarray analysis of mRNA synthesis and decay. *Nat. Biotechnol.* **23**, 232–237 (2005).

17.   Dolken, L. *et al.* High-resolution gene expression profiling for simultaneous kinetic parameter analysis of RNA synthesis and decay. *RNA* **14**, 1959–1972 (2008).

18.   Miller, C. *et al.* Dynamic transcriptome analysis measures rates of mRNA synthesis and decay in yeast. *Mol. Syst. Biol.* **7**, 458–458 (2014).

19.   Duffy, E. E. *et al.* Tracking Distinct RNA Populations Using Efficient and Reversible Covalent Chemistry. *Mol. Cell* **59**, 858–866 (2015).

20.   Schwalb, B. *et al.* TT-seq maps the human transient transcriptome. *Science* **352**, 1225–1228 (2016).

21.   Rabani, M. *et al.* Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nat. Biotechnol.* **29**, 436–442 (2011).

22.   Miller, M. R., Robinson, K. J., Cleary, M. D. & Doe, C. Q. TU-tagging: cell type–specific RNA isolation from intact complex tissues. *Nat. Methods* **6**, 439–441 (2009).

23.   Kawata, K. *et al.* Metabolic labeling of RNA using multiple ribonucleoside analogs enables the simultaneous evaluation of RNA synthesis and degradation rates. *Genome Res.* **30**, 1481–1491 (2020).

24.   Battich, N. *et al.* Sequencing metabolically labeled transcripts in single cells reveals mRNA turnover strategies. *Science* **367**, 1151–1156 (2020).

25.   Ziegenhain, C. *et al.* Comparative Analysis of Single-Cell RNA Sequencing Methods. *Mol. Cell* **65**, 631–643.e4 (2017).

26.   Ding, J. *et al.* Systematic comparison of single-cell and single-nucleus RNA-sequencing methods. *Nat. Biotechnol.* **38**, 737–746 (2020).

27.   Joung, J. *et al.* Genome-scale CRISPR-Cas9 knockout and transcriptional activation screening. *Nat. Protoc.* **12**, 828–863 (2017).

28.   Replogle, J. M. *et al.* Mapping information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. *Cell* **185**, 2559–2575.e28 (2022).

29.   Sanson, K. R. *et al.* Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities. *Nat. Commun.* **9**, 5416 (2018).

30.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888–1902.e21 (2019).

31.   Qiu, X. *et al.* Mapping transcriptomic vector fields of single cells. *Cell* **185**, 690–711.e45 (2022).

32.   Jones, P. L. *et al.* Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat. Genet.* **19**, 187–191 (1998).

33.   Dominguez, A. A., Lim, W. A. & Qi, L. S. Beyond editing: repurposing CRISPR–Cas9 for precision genome regulation and interrogation. *Nature Reviews Molecular Cell Biology* vol. 17 5–15 Preprint at https://doi.org/10.1038/nrm.2015.2 (2016).

34.   Kim, S. H. & Lin, R. J. Pre-mRNA splicing within an assembled yeast spliceosome requires an RNA-dependent ATPase and ATP hydrolysis. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 888–892 (1993).

35.   Colgan, D. F. & Manley, J. L. Mechanism and regulation of mRNA polyadenylation. *Genes & Development* vol. 11 2755–2766 Preprint at https://doi.org/10.1101/gad.11.21.2755 (1997).

36.   Kikkawa, S. *et al.* Conversion of GDP into GTP by nucleoside diphosphate kinase on the GTP-binding proteins. *J. Biol. Chem.* **265**, 21536–21540 (1990).

37.   Siira, S. J. *et al.* LRPPRC-mediated folding of the mitochondrial transcriptome. *Nat. Commun.* **8**, 1532 (2017).

38.   Ruzzenente, B. *et al.* LRPPRC is necessary for polyadenylation and coordination of translation of mitochondrial mRNAs. *EMBO J.* **31**, 443–456 (2012).

39.   Liu, L. *et al.* LRP130 protein remodels mitochondria and stimulates fatty acid oxidation. *J. Biol. Chem.* **286**, 41253–41264 (2011).

40.   Pajak, A. *et al.* Defects of mitochondrial RNA turnover lead to the accumulation of double-stranded RNA in vivo. *PLoS Genet.* **15**, e1008240 (2019).

41.   Pakos-Zebrucka, K. *et al.* The integrated stress response. *EMBO Rep.* **17**, 1374–1395 (2016).

42.   Buccitelli, C. & Selbach, M. mRNAs, proteins and the emerging principles of gene expression control. *Nat. Rev. Genet.* **21**, 630–644 (2020).

43.   Chipman, L. B. & Pasquinelli, A. E. miRNA Targeting: Growing beyond the Seed. *Trends Genet.* **35**, 215–222 (2019).

44.   Treiber, T., Treiber, N. & Meister, G. Regulation of microRNA biogenesis and its crosstalk

with other cellular pathways. *Nat. Rev. Mol. Cell Biol.* **20**, 5–20 (2019).

45. Kim, Y.-K., Kim, B. & Kim, V. N. Re-evaluation of the roles of DROSHA, Export in 5, and DICER in microRNA biogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E1881–9 (2016).

46. Park, M. S. *et al.* Multidomain Convergence of Argonaute during RISC Assembly Correlates with the Formation of Internal Water Clusters. *Mol. Cell* **75**, 725–740.e6 (2019).

47. Liu, B., Shyr, Y., Cai, J. & Liu, Q. Interplay between miRNAs and host genes and their role in cancer. *Brief. Funct. Genomics* **18**, 255–266 (2018).

48. Chureau, C. *et al.* Ftx is a non-coding RNA which affects Xist expression and chromatin structure within the X-inactivation center region. *Hum. Mol. Genet.* **20**, 705–718 (2011).

49. Zhao, L., Mao, Y., Zhao, Y. & He, Y. DDX3X promotes the biogenesis of a subset of miRNAs and the potential roles they played in cancer development. *Scientific Reports* vol. 6 Preprint at https://doi.org/10.1038/srep32739 (2016).

50. Heinrichs, A. A slice of the action. *Nat. Rev. Mol. Cell Biol.* **5**, 677–677 (2004).

51. Lytle, J. R., Yario, T. A. & Steitz, J. A. Target mRNAs are repressed as efficiently by microRNA-binding sites in the 5' UTR as in the 3' UTR. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 9667–9672 (2007).

52. Replogle, J. M. *et al.* Combinatorial single-cell CRISPR screens by direct guide RNA capture and targeted sequencing. *Nat. Biotechnol.* **38**, 954–961 (2020).

53. Liscovitch-Brauer, N. *et al.* Profiling the genetic determinants of chromatin accessibility with scalable single-cell CRISPR screens. *Nat. Biotechnol.* **39**, 1270–1277 (2021).

54. Mimitou, E. P. *et al.* Multiplexed detection of proteins, transcriptomes, clonotypes and CRISPR perturbations in single cells. *Nat. Methods* **16**, 409–412 (2019).

55. Scarpulla, R. C., Vega, R. B. & Kelly, D. P. Transcriptional integration of mitochondrial biogenesis. *Trends Endocrinol. Metab.* **23**, 459–466 (2012).

56. Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat. Methods* **14**, 1083–1086 (2017).

57. La Torre, A., Georgi, S. & Reh, T. A. Conserved microRNA pathway regulates developmental timing of retinal neurogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E2362–70 (2013).

58. Janowski, B. A. *et al.* Involvement of AGO1 and AGO2 in mammalian transcriptional silencing. *Nat. Struct. Mol. Biol.* **13**, 787–792 (2006).

59. Griffin, K. N. *et al.* Widespread association of the Argonaute protein AGO2 with meiotic

chromatin suggests a distinct nuclear function in mammalian male reproduction. *Genome Res.* **32**, 1655–1668 (2022).

60.   Moshkovich, N. *et al.* RNAi-independent role for Argonaute2 in CTCF/CP190 chromatin insulator function. *Genes Dev.* **25**, 1686–1701 (2011).

61.   Krueger, F. A wrapper around Cutadapt and FastQC to consistently apply adapter and quality trimming to FastQ files, with extra functionality for RRBS data. *TrimGalore (accessed on 27 August 2019).*

62.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

63.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

64.   Lindenbaum, P. JVarkit: java-based utilities for Bioinformatics. (2015) doi:10.6084/m9.figshare.1425030.v1.

65.   Picard. https://broadinstitute.github.io/picard/.

66.   Guangchuang Yu [aut, cre], Li-Gen Wang [ctb], GiovanniDall'Olio [ctb] (formula interface of compareCluster). *clusterProfiler*. (Bioconductor, 2017). doi:10.18129/B9.BIOC.CLUSTERPROFILER.

67.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods* **14**, 979–982 (2017).

68.   Zhang, K. *et al.* A novel class of microRNA-recognition elements that function only within open reading frames. *Nat. Struct. Mol. Biol.* **25**, 1019–1027 (2018).

69.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

70.   Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Research* vol. 27 491–499 Preprint at https://doi.org/10.1101/gr.209601.116 (2017).

71.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).