Colour patterns are less variable in highly unpalatable nudibranchs

Cedric P. van den Berg[1†], John A. Endler[2], Leon Drummond[1], Bethany R. Dawson[1], Carl Santiago[1], Nathalie Weber[3], Karen L. Cheney[1]

[1]Visual Ecology Laboratory, School of Biological Sciences, The University of Queensland, 4072 Brisbane, Australia

[2]Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Geelong, Victoria 3216, Australia

[3]Faculty of Biology and Medicine, School of Biological Sciences, The University of Lausanne, 1015 Lausanne, Switzerland

Corresponding author: c.vandenberg@bristol.ac.uk

† Current address: School of Biological Sciences, The University of Bristol, BS81TQ, United Kingdom

ORICD ID:

CPvdB 0000-0001-6422-7237

JAE 0000-0002-7557-7627

LD 0000-0003-2090-7876

BRD 0000-0001-6677-4453

CS 0000-0002-3812-2535

NW 0000-0001-6709-9787

KLC 0000-0001-5622-9494

Abstract

Predators learn and memorise the association between conspicuous colour patterns of

aposematic prey and their underlying chemical or secondary defences. Consequently, variation

in signal design within a species should be selected against, because it can decrease the rate of

predator learning and enhance the rates of predator errors. However, quantitative assessments

have not been made on whether the strength of chemical defences influences colour pattern

variation. We examined this by quantifying colour pattern variation using Quantitative Colour

Pattern Analysis (QCPA) in 12 Dorid nudibranch species (Infraorder: Doridoidei) that varied

in their unpalatability. We accounted for the physiological limitations of a potential predator's

visual system (a triggerfish, *Rhinecanthus aculeatus*), modelling animal appearance along an

escalating predation sequence. We found that various colour pattern statistics were less variable

in highly unpalatable species, with pattern statistics being up to 72% less variable than those of

more palatable species. No correlations indicating the opposite were found across 157 colour

pattern statistics. However, the strength and number of correlations depended on viewing

distance. Our results suggest that low colour pattern variability could be favourable for

aposematic signalling in Dorid nudibranchs. We provide evidence for distance-dependent

signalling facilitating context-specific feature selection in multicomponent visual signals.

Keywords: Signal honesty, aposematism, crypsis, predator psychology, purifying selection,

defensive animal colouration

1. Introduction

Many animals use aposematic colour patterns to warn potential predators of underlying

defences [1]. Once aposematism has evolved, purifying selection is expected to constrain

colour pattern diversity within a species and/or populations as predators learn to associate a

visual signal with unprofitability [2–7]. A consistent appearance across aposematic individuals

may facilitate predator learning and memorisation, whereas variation in signal design may

cause predators to make errors when attacking prey and decrease rates of predator learning

[4,8–10]. Yet some degree of colour pattern diversity within aposematic species is ubiquitous

and is thought to be driven by countervailing evolutionary factors such as genetic drift, gene

flow, sexual selection, or variation in resource abundance [11–13].

Colour pattern variation in aposematic species depends on the absolute and relative
abundance of individual species in a given environment and the strength of their secondary
defences [10,14,15]. For strongly defended and abundant species, anti-apostatic selection is
expected to be common [15]. This is because, in strongly defended species, rare and unfamiliar
morphs, proportional to the strength of associated secondary defences, are increasingly
predated on [3]. Such correlations between reduced colour pattern variability and increased
strength of secondary defences may further be accelerated in aposematic species with
aggregated occurrence as inclusive fitness through kin selection is argued to enhance anti-
apostatic selection [16,17].

Cumulatively, such mechanisms explain variable colour patterns in aposematic species
across heterogeneous environments while explaining reduced variability within aposematic
species at a local scale (e.g. [18,19]). Specifically, aposematism in a homogeneous and stable
environment is expected to coincide with selection towards reduced colour pattern variability
within a population [20,21]. However, whether the strength of chemical defences influences
pattern variability is unclear, and quantitative assessments are lacking (see [21,22] for review).
Highly defended species should exhibit more uniform colour patterns to strengthen predator
learning of the aposematic signal. However, attacking well-defended prey is costly; therefore,
predators may generalise more broadly between colour patterns of previously attacked prey and
prey they subsequently encounter (e.g. [23–25]). This could relax selection pressures on
uniform colour patterns within a population resulting in higher variability. Selection for or
against colour pattern variability could also only occur on individual colour pattern elements,
depending on which elements of the signal predators learn or pay attention to [26].

When interpreting the ecological relevance of phenotypic variation, it is important to
consider how the appearance of an organism changes as a function of observer acuity and
viewing distance [27]. For example, colour patterns may be cryptic when viewed from a
distance but may become aposematic as a predator approaches [27,28]. Consequently, predator
learning of associations between primary and secondary prey defences, or the subsequent
retrieval of formed associations from memory upon encounters with aposematic prey, might
happen at a specific range of viewing distances. Animals also detect objects and make decisions
on their identity and quality based on complex combinations of spatiochromatic features [29–
32]. However, empirical evidence on the ecological significance of colour pattern variability in
aposematic animals remains largely restricted to investigations of colour alone (e.g. [33,34]).

In this study, we examined whether highly defended aposematic nudibranch species exhibited less variable colour patterns than less defended species. We further hypothesized that distance-dependent signalling might also affect any correlations between colour pattern variability and level of defence. To do this, we modelled the visual appearance of 12 sympatric Dorid nudibranch species, for which we have previously published information on their levels and type of chemical defences [35]. We did this across viewing distances corresponding to an escalating predation sequence [8,36]. We quantified the perception of within-species colour pattern variability using the Quantitative Colour pattern Analysis (QCPA) [37], allowing for the consideration of colour, luminance and spatial vision of triggerfish (*Rhinecanthus aculeatus*). We then correlated the perceived variability of colour patterns with our previously published unpalatability data for each species [35].

2. Materials and Methods

(a) Study species

We took digital photographs of 353 Dorid nudibranchs using a calibrated Olympus EPL-5 with a 60mm macro lens. These individuals belonged to 12 species: *Goniobranchus collingwoodi* (N=20); *Phyllidiella pustulosa* (N=40); *Glossodoris vespa* (N=32); *Goniobranchus splendidus* (N=37); F) *Doriprismatica atromarginata* (N=35); *Hypselodoris bennetti* (N=13); *Chromodoris kuiteri* (N=49); *Chromodoris magnifica* (N=14); *Phyllidia ocellata* (N=32); *Chromodoris elisabethina* (N=31); *Discodoris* sp. (N=19); *Aphelodoris varia* (N=31) (Fig. 1) from five locations on the east coast of Australia: Mackay (QLD), Sunshine Coast (SE Queensland, QLD), Gold Coast (SE QLD), Cook Island (New South Wales, NSW) and Nelson Bay (NSW) between March 2016 and February 2021. Two individuals of *Chromodoris magnifica* were provided by an aquarium supplier (Cairns Marine, Pty Ltd, Cairns, QLD). A sub-sample of these images were previously used to investigate background habitat variability of nudibranch molluscs [38] and distance-dependent signalling in terms of colour pattern detectability and boldness [39]. These species were selected as they were relatively abundant at our study sites and cover a broad range of visual appearance and strength of chemical defences. Specifically, we have previously provided a detailed investigation into the strength and identity of chemical defences in these species sampled from the same locations as individuals from this study [35].

Two out of 12 species (*Doriprismatica atromarginata, Goniobranchus splendidus*) were sampled across sites in QLD and NSW in high numbers, whereas the other species were

only sampled in either NSW or QLD or with highly uneven numbers between sites (Table S2).

Nudibranchs were either photographed underwater against their natural habitat (n = 216) with

the camera in an Olympus PT-EP10 underwater housing and using white LED illumination

from a combination of VK6r and PV62 Scubalamp video lights. Alternatively, nudibranchs (n

= 137) were collected, taken back to the laboratory, submerged in water in a petri dish and

photographed against a white background with the same camera. In the laboratory, nudibranchs

were illuminated with 300nm-700nm full-spectrum white LED lights. Details on collection

sites and dates are provided in the Supplementary Information, camera and illumination spectra

are provided in [37]. Nudibranchs were collected under the Queensland General Fisheries

Permits 183990, 205961 and NSW Scientific Collection Permit P16/0052-1.0.

(b) Image analysis

We used ImageJ [40] and the MICA toolbox [41], to manually segment the images into

regions of interest (ROI), selecting the animal from its background and defining a size standard.

All nudibranch were aligned vertically prior to analysis using QCPA [37]. To analyse the

nudibranch colour patterns, we used the visual system parameters of a trichromatic triggerfish,

Rhinecanthus aculatus, a common shallow reef inhabitant found throughout the Indo-Pacific,

which feeds on invertebrates, algae, and detritus [42]. Their visual system has been well

described [43–48].

We analysed colour patterns for viewing distances of 2cm, 5cm, 10cm and 30cm

assuming the spatial acuity of the triggerfish was 3 cycles per degree [43,47]. Following acuity

modelling, the images were processed with a Receptor Noise Limited (RNL) ranked filter

(falloff: 3, radius: 5, repetition: 5) and clustered using RNL clustering with a colour threshold

of $2\,\Delta S$ [48,49] and a luminance contrast threshold of $4\,\Delta S$ [50] for all analyses except the local

edge intensity analysis (LEIA) which does not require image segmentation but is recommended

to be subjected to RNL ranked filtering. We calculated Weber fractions based on a relative

photoreceptor abundance of 1:2:2:2 (sw:mw:lw:dbl) and photoreceptor noise of 0.05, resulting

in 0.07:0.05:0.05:0.05.

QCPA analysis was achieved using a custom batch script [51] running on high

performance computing (HPC) infrastructure. We analysed each animal's values for: 1) colour

adjacency analysis (CAA), which describes pattern geometry in a segmented image; 2) visual

contrast analysis (VCA), which describes pattern boldness based on chromatic and spatial

pattern element properties in a clustered image; 3) boundary strength analysis (BSA), which

describes the colour and luminance contrast of boundaries between pattern elements at the scale

of an animal in an unclustered image; and 4) local edge intensity analysis (LEIA) which

describes the strength of colour and luminance contrast at the scale of an edge-detecting

receptive field in an unclustered image. This resulted in a highly descriptive array of 157 colour

pattern statistics for each animal. A full detailed description of each pattern statistic is found in

[37]. Here we use CAA, VCA, BSA, and LEIA as prefixes for each type of analysis. All pattern

analyses used a segmented image and measured transitions between pixels along vertical (along

body axis) and horizontal (perpendicular to body axis) sampling transects in a transition matrix.

Statistics ending with 'vrt' or 'hrz' are the vertical and horizontal version (analysing the

respective transition matrix only) of their respective statistic (analysing the full transition

matrix) and are able to represent differential directionality sensitivity in the visual system of an

observer and directionality in patterns such as stripes [52–54]. LEIA does not use a transition

matrix, but equally discriminates between horizontal and vertical edge contrast [37].

Figure 1. Representative photographs of the 12 species used in this study, ordered from most
unpalatable to least unpalatable with the strength of chemical defences reported as $1\text{-}ED_{50}$ [35].
A) *Goniobranchus collingwoodi;* B) *Phyllidiella pustulosa;* C) *Glossodoris vesp*a; D)
Goniobranchus splendidus; E) *Doriprismatica atromarginata;* F) *Hypselodoris bennetti;* G)
Chromodoris kuiteri; H) *Phyllidia ocellata*; I) *Chromodoris magnifica*; J) *Chromodoris
elisabethina;* K) *Discodoris sp.*; L) *Aphelodoris varia.* The level of chemical defence ($1\text{-}ED_{50}$)
is indicated underneath each species.

To determine the level of chemical defences for each species (i.e. [7]), we used

previously published data on the deterrent properties from feeding rejection assays with

rockpool shrimp (*Palaemon serenus*) that are similar to assays performed with triggerfish and

toadfish [35]. Here, we use unpalatability as opposed to toxicity, as the former is more likely

reflective of a repellent property of chemical defences [35,55]. Assays were conducted by

adding extracted nudibranch compounds at increasing concentrations to food pellets made from

squid mantle. Effective dose (ED_{50}) values were calculated based on the concentration that

elicited a rejection response in at least 50% of the shrimp. For the purpose of this study, ED_{50}

values from [35] were averaged when multiple extracts from the same species were tested. We

then subtracted these values from 1 so that values close to 0 were the least defended/most

palatable and values close to 1 highly defended/least palatable) (Table S2). Importantly,

specimens in [35] were collected from the same locations as in this study. Although

Chromodoris magnifica was not included in [35], [56] demonstrated that this species also stores

latrunculin A as the sole defensive compound in the mantle rim. We have also recently shown

that this is at concentrations in between those found in *Chromodoris elisabethina* and

Chromodoris kuiteri [57]. We, therefore, set unpalatable ED_{50} values as the average from these

two sister species. Lastly, assay data for *Glossodoris vespa* is presented in [58].

(d) Statistical analysis

To quantify colour pattern variability, we calculated the inter-quartile range (IQR) for

each parameter of each species. All pattern statistics were then scaled from 0 to 1 using the

'caret' package [59], with 1 being most variable. We assessed whether there were colour pattern

differences between sites for those species that were sampled roughly equally across locations

(*G. splendidus, D. atromarginata*); however, we found no significant differences between

locations in the IQR across all 157 parameters (lmer: $F_{17} = 2.12$, $p = 0.16$). Due to the small

number of species and variable colour pattern statistics, it was not possible to estimate the

phylogenetic signal reliably using phylogenetic least square regression (pgls), with lambda

estimates being inconclusive. Therefore, in line with previous studies [60,61], we assumed a

phylogenetic signal of 0 for all viewing distances when analysing the median IQR for each

species across all 157 pattern statistics using a Pearson product moment correlation test in the

'stats' package [62]. Prior to analysis, the IQR values were normalised using the

'bestNormalize' package [63]. As all colour pattern statistics essentially describe the same three

core properties (colour, luminance and pattern), we applied a conservative Bonferroni

correction of n = 3 to avoid an excessive type II error (for discussion see [64]), resulting in an

adjusted significance threshold of $p = 0.017$.

We present the results from the largest viewing distance to the smallest to replicate an

escalating predation sequence [8]. Detailed reporting of test statistics for all 157 pattern

statistics at each viewing distance are presented in the supplementary material.

3. Results

The species-specific interquartile range (IQR) of various colour pattern statistics was

negatively correlated with increasing unpalatability levels at each viewing distance (Pearson

correlation: $r < -0.63$, $p < 0.017$, Fig. 2), indicating that highly unpalatable species showed less

less among-individual variation in visual appearance than more palatable ones. Importantly, we

did not find any significant positive correlations in our data (Tables S3) nor any positive

correlations of moderate strength ($r > 0.58$, $p < 0.05$) which would have indicated the opposite.

At 30 cm, the colour pattern regularity measured as the Shannon diversity of the

synonymous transitions in the transition matrix (*CAA.Hc*) is 71% less variable in highly

defended species ($r = -0.84$, $p = 0.001$). This is the case for both, looking at the horizontal

(*CAA.Hc.Hrz*, $r = -0.85$, $p < 0.001$) and the vertical axis (CAA.Hc.Vrt, $r = -0.84$, $p = 0.001$) of

colour patterns with variability (IQR) in highly defended species being 72% (Hrz) and 71%

(Vrt) more reduced (Fig. 2A, Table 1). Additionally, nine other pattern statistics showed

moderately strong negative correlations between chemical defences and colour pattern

variability ($r < -0.58$, $p < 0.05$, Table S2).

At 10 cm, six out of 157 pattern statistics were significantly negatively correlated with

unpalatability (Fig. 2 B, Table 1). The more unpalatable a species was, the less variable the

colour pattern boldness elicited by the mean abundance weighted saturation contrast between

colour pattern elements. This was the case for both, saturation measured as the distance from

the achromatic point in the RNL colour space [65], as well as maximum possible pairwise

contrast between photoreceptors irrespective of relative abundances and receptor noise [66].

These are denoted as *Ssat* and *Dmax* respectively (see [37] for details). The mean abundance-

weighted saturation contrast measured as *Dmax* saturation (*VCA.MDmax*) between colour

pattern elements was 52% ($r = -0.72$, $p = 0.008$) less variable in highly defended species than

it was in undefended ones, irrespective of the horizontal or vertical axis (Fig. 2 B, Table 1).

When measuring saturation as RNL saturation (*Ssat*), the mean abundance-weighted saturation

contrast (*VCA.MSsat*) was 50% ($r = -0.71$, $p = 0.010$) less variable in highly defended species

than in undefended ones.

At a closer viewing distance of 5 cm, three out of 157 pattern statistics were significantly
negatively correlated with unpalatability (Fig. 2 C, Table 1). The mean saturation contrast
(*VCA.MDmax*, $r = -0.70$, $p = 0.011$) was 51% less variable in highly defended species as
opposed to undefended ones. This was true for both, the vertical (*VCA.MDmax.Vrt*, $r = -0.71$,
$p = 0.010$) and horizontal axis (*VCA.MDmax.Vrt*, $r = -0.71$, $p = 0.010$). Additionally, we found
six colour pattern statistics showing moderately strong negative correlations ($r < -0.58$, $p < 0.05$,
Table S2).

Lastly, at 2 cm, the same three out of 157 statistics were significantly negatively
correlated with unpalatability (Fig. 2 D, Table 1). This distance most likely reflects immediate
contact between predator and prey. The more unpalatable a species was, the mean abundance
weighted saturation contrast between colour pattern elements was 49% less variable
(*VCA.MDmax*, $r = -0.70$, $p = 0.011$). However, saturation contrast variation along the horizontal
(*VCA.MDmax.Hrz*, $r = -0.75$, $p = 0.005$) and vertical axis (*VCA.MDmax.Vrt*, $r = -0.72$, $p =$
0.008) was slightly more reduced in highly defended species when looked at separately. We
also found three other pattern statistics to show a moderately strong negative correlation ($r < -$
0.58, $p < 0.05$, Table S2).

Figure 2. All significant correlations between colour pattern variability (interquartile range, IQR) and level of unpalatability (ED_{50}) out of 157 assessed colour pattern statistics across all 12 species (Pearson correlation: $r < -0.5$, $p < 0.017$) for each viewing distance: A) 30 cm, B) 10 cm, C) 5 cm, D) 2cm. Grey shading indicates 95% confidence intervals of the linear regressions. The most likely corresponding step of the predation sequence is indicated with schematics: exclamation mark signifies encounter and detection; the question mark signifies identification and approach, and finally subjugation and consumption. The x-axis shows the normalised 1-ED_{50} from palatability assays with Palaemon shrimp where 0 refers to the most palatable species and 1 refers to the most unpalatable species. The y-axis shows the variability of colour pattern statistics expressed as the normalised IQR (interquartile range) for each species where 0 refers to the least variable species and 1 to the most variable species.

Table 1. All significantly correlated pattern statistics for each viewing distance with a Bonferroni adjusted Pearson correlation significance of $p < 0.017$. Statistics are ranked according to their significance at each viewing distance. Pearson correlations (r) are also expressed as a percentage.

Viewing distance	Stage in Predation Sequence	Pattern statistic	Pearson Correlation [r] / % [r²*100]	P value
30cm	Detection / Identification	CAA.Hc.Hrz	-0.85 / 72	<0.001
		CAA.Hc	-0.84 / 71	<0.001
		CAA.Hc.Vrt	-0.84 / 71	<0.001
10cm	Identification / Approach	VCA.MDmax	-0.72 / 52	0.008
		VCA.MDmax.Hrz	-0.72 / 52	0.008
		VCA.MDmax.Vrt	-0.72 / 52	0.008
		VCA.MSsat	-0.71 / 50	0.010
		VCA.MSsat.Hrz	-0.71 / 50	0.010
		VCA.MSsat.Vrt	-0.71 / 50	0.010
5cm	Approach / Subjugation	VCA.MDmax.Hrz	-0.71 / 51	0.010
		VCA.MDmax.Vrt	-0.71 / 51	0.010
		VCA.MDmax	-0.70 / 49	0.011
2cm	Subjugation / Consumption	VCA.MDmax.Hrz	-0.75 / 56	0.005
		VCA.MDmax.Vrt	-0.72 / 52	0.008
		VCA.MDmax	-0.70 / 49	0.011

Figure 3. Pattern variability measured as the variability of the Shannon colour diversity index (*CAA.HC*) at 30 cm viewing distance (Pearson correlation: $r = -0.84$, $p = < 0.001$). *CAA.HC* calculates the Shannon diversity index of the synonymous (off-diagonal) transitions in the transition matrix, *i.e.* the evenness of colour pattern elements in a pattern (electronic Supplement in [37]).

4. Discussion

Our results indicate that highly defended species exhibit less variable colour patterns than moderately or undefended species. We found that different measures of colour pattern variation, specifically parameters measuring chromatic boldness (*VCA.MSsat & VCA.MDmax*) and colour pattern regularity (*CAA.Hc*), were correlated with unpalatability at different viewing distances. This is intriguing, as chromatic contrast and colour pattern regularity are generally regarded as indicators of potent aposematic signalling [67,68]. Our data does not show any differential importance of the pattern information contained in the horizontal or vertical dimension along the body axis at any viewing distance (Table 1). This suggests that selective pressure imposed on colour pattern variability by differential orientation sensitivity of predatory visual systems (e.g. [53,54]) may be negligible. In fact, the strongest correlations between colour pattern variability and the strength of chemical defences were found at 30cm where colour pattern regularity (*CAA.Hc*). *CAA.Hc* was 72% less variable in species with strong chemical defences.

The second most significant correlation occurred at a viewing distance of 10cm, where prey identification and discrimination are likely possible. Indeed, colour pattern boldness through internal colour contrast, such as that measured by *VCA.MSsat* and *VCA.MDmax* is a crucial component of the effectiveness of aposematic signalling [28,69]. Given that variability in colour, not luminance, contrast is significantly correlated with the strength of secondary defences further aligns with the importance of colour for object identification and assessment [70], a crucial aspect of aposematic signalling. Interestingly, the maximum possible colour contrast between colour pattern elements (*Dmax*) by a triggerfish (*Rhincenathus aculeatus*) almost identically matches the colour contrast calculated using the RNL model (*Ssat*). This suggests that the visual system of *R. aculeatus* is indeed very well adapted to perceive colour pattern contrast in nudibranch molluscs.

We also highlight how distance-dependent signalling in Dorid nudibranchs can facilitate selection on specific aspects of the design and function of aposematic colour patterns [27,32,71,72]. Aposematic colour patterns in nudibranchs might have evolved to be cryptic at a distance but bold when viewed up close or bold upon successful detection [20,71]. For example, we found the largest number of correlations between pattern variation and unpalatability at 10 cm, a distance where fine colour pattern details become apparent (Fig. 2 & 3). Our findings align with the idea of purifying selection shaping the regularity-specific colour

pattern elements (as opposed to the entirety of an animal's visual appearance) as context-
specific aposematic signals in Dorid nudibranchs (e.g. [26]). Specifically, we show negative
correlations between colour pattern variability and the strength of secondary defences to occur
for separate colour pattern properties at distinct viewing distances. Thus, our results suggest
colour patterns of nudibranch molluscs to comprise true multicomponent signals [73] with
distance-dependent functionality.

Our study is unable to assess how and if the observed correlations between traits are
heritable or the consequence of phenotypic plasticity, due to the absence of reliable estimates
of signal heritability for each of the measured traits. Without such estimates, conclusions of
natural selection are complicated as selection affects the phenotypes, not the genotypes of
species in a given community [3,74,75]. Therefore, how and if the observed phenotypes are
indeed confounded by the evolutionary history between species remains unknown. However,
high degrees of phenotypic plasticity and rapid radiation of species morphology in Dorid
nudibranchs are common (e.g. [76]) and the general presence of gene flow across populations
in the study area is likely [26]. For example, *Phyllidia ocellata* is known to be highly variable
in appearance across its range [77], but is found to be very uniform in appearance within our
study area (Fig. 3), thus further supporting our assumptions of significant connectivity between
sampling locations. Furthermore, it is reasonable to assume similar visual and cognitive
abilities of the predator community across the studied area [78], as well as the overall similarity
of the visual appearance of rocky reef habitats across the area [38]. Together, these factors
indicate our study to quantify variability between species with low levels of phylogenetic
confound, consisting of well-connected meta-populations in a comparably homogeneous
landscape of selective pressures across the study area.

Existing literature predicts highly defended species to display uniform colour patterns, as
consistent appearance across individuals should enhance learning and memorisation of
aposematic signals by predators. However, it is not easy to explain why moderately defended
species have moderate levels of pattern variation. Moderately defended nudibranchs may have
different predators compared to highly defended species. These predators (e.g. crabs) may not
be as reliant on visual signals when attacking prey, so selection for uniform patterns may be
relaxed for these species (e.g. [79]). Alternatively, attacking moderately defended species is less
costly for predators and may occur more frequently, particularly in areas or times where
alternative food sources are unavailable, which could weaken predator learning. Finally,

moderately defended species could be found on particularly complex backgrounds, making a
uniform colour pattern less effective, again weakening selection on uniformity.

However, reasoning for specific causes underlying the observed correlation is difficult.
For example, unlike in other organisms considered in the study of aposematic colouration [e.g.
19,77] predator communities and their interactions with local populations of nudibranchs
beyond anecdotal observations remain poorly understood. Furthermore, many predictions on
the direction of selection for or against colour pattern variability in aposematic signals [3,4]
rely on assumptions on the genetic makeup and thus mechanisms underlying the traits under
selection. However, it remains unknown which genes or how many of them are involved in the
expression of colour pattern morphology in nudibranchs; a complex trait under multiple
selective pressures (e.g. [73,81]). Indeed, genetic mechanisms underlying adaptive processes
in defensive colouration remain largely elusive, even for individual components of colour
patterns such as spot size or colour variability of a distinct pattern element [but see 80].
Furthermore, how and if environmental factors, such as turbidity, temperature and variability
in spatial and seasonal processes (e.g. [83–86]) interact with such selection pressures and thus
variability in phenotype, remains unknown as well. Lastly, the defensive properties of
secondary metabolites can be highly context and predator specific [35,87,88]. Such context-
dependent or predator specific insights do not exist for the study of secondary metabolites in
nudibranchs, despite the growing number of identified compounds and generalisable assays.

Therefore, drawing conclusions the specific mechanisms underlying our observation is
difficult, especially given the complicated and poorly resolved phylogenetic relations between
many nudibranch species discussed previously. Importantly, our understanding of both, human
and animal visual perception and cognition are constantly evolving, and with it, our
understanding of the ecological significance of signal variation in the context of aposematism.
For example, Raviv *et al.* [89] recently showed that, in humans, variability in visual stimuli is
actually likely to enhance learning under certain conditions. This is a highly relevant finding
for our understanding of selective pressures shaping diversity in defensive animal colouration
which is currently not considered in the literature. In fact, perceiving variability as a factor
enhancing (rather than inhibiting) animal learning diametrically opposes the general
understanding in the existing literature. Importantly, such differentiated views on the
complexity of visual perception and cognition, in combination with the known complexity of
factors underlying phenotypic variation in aposematic animals, make it rather unsurprising that

the strength of secondary defences and colour pattern morphology share a continuous rather
than a discrete relationship.

Nudibranchs only have rudimentary visual abilities enabling basic phototaxis [90] and
the species observed in this study live in thermally comparably stable aquatic habitats (i.e. no
species were collected in rockpools or at shallow depths <2m) experiencing only gradual
seasonal fluctuations. Given the multitude of alternative selective pressures such as
intraspecific signalling or thermoregulation acting on colour pattern morphology (and thus
variability) in traditionally considered study organisms such as poison dart frogs [e.g. 89],
makes our findings particularly valuable for understanding the potential effects of visual
predation on colour pattern variability in aposematic species.

Our study has important implications for quantifying colour pattern variation in
complex natural environments by addressing the importance of ecologically relevant observers
and their perception of complex, multicomponent visual signals. Colour pattern variability can
be observed and quantified in several ways; however, metrics that are perceptually relevant and
contributing to the cognitive processes should be used to help inform our knowledge regarding
the ecology and evolution of animal colouration [27,92,93]. To date, few studies investigating
colour pattern variability make this distinction beyond the modality of colour (e.g. [50,94,95]).
In fact, our study is the first to incorporate both, spatial acuity as well as colour and luminance
contrast perception of an ecologically relevant observer in the assessment of aposematic signal
variability. In the context of a steadily growing array of tools available for the study of animal
colour patterns, we emphasize the need for careful consideration of the physiological limits of
animal visual systems and the ecologically relevant perceptual context studies investigating
quantitative and qualitative aspects of the ecology and evolution of aposematism and defensive
animal colouration more broadly. Importantly, we test each colour pattern statistic individually.
However, considering colour pattern statistics cumulatively, allowing for interactions between
multiple descriptors, is of great interest in future studies (see [32] for discussion).

In summary, we showed that colour pattern variability in aposematic nudibranchs is
negatively correlated with the strength of their secondary defences, thus being indicative of
quantitative and qualitative signal honesty [10,21,22,96] in local populations of Dorid
nudibranchs belonging to substantially different genera and families. Our findings align with
broader indications of quantitative signalling honesty in nudibranchs from existing literature
[61]. In fact, the strength of the association between reduced intraspecific colour pattern

variability and increased secondary defences is among the strongest indications of quantitative
signalling honesty in any aposematic organism observed to date (see [22] for review).

Data accessibility

Authors' contributions

CPvdB: Conceptualisation, data curation, formal analysis, funding acquisition, investigation,
methodology, project administration, resources, software, validation, visualisation, writing –
original draft, writing – review & editing. JAE: project funding acquisition, writing - review &
editing. BRD, CS, NW: investigation. KLC: Funding acquisition, project administration,
resources, validation, writing - review & editing.

Competing interests

We declare we have no competing interests.

Acknowledgements

We would like to thank Dr Nicholas Condon for his assistance with the creation of automated
QCPA scripts; various volunteers for assistance with image analysis; the High-Performance
Computing (HPC) infrastructure at UQ (Wiener & Awoonga) and infrastructure provided by
Simone Blomberg which contributed to the computing of image statistics.

Funding

This work was funded by the Australian Research Council (FT190199313 awarded to KLC and
DP180102363 awarded to JAE), Holsworth Wildlife Research Endowment (two grants
awarded to CPvdB) and the Australasian Society for the Study of Animal Behaviour (research
grant awarded to CPvdB).

References

1. Poulton EB. 1890 *The Colours of Animals*. Kegan Paul, Trench & Trubner.

2. Borer M, Van Noort T, Rahier M, Naisbit RE. 2010 Positive frequency-dependent selection on warning color in alpine leaf beetles. *Evolution (N. Y).* **64**, 3629–3633. (doi:10.1111/j.1558-5646.2010.01137.x)

3. Endler JA, Greenwood JJD. 1988 Frequency-dependent predation, crypsis and aposematic coloration. *Philos. Trans. R. Soc. London. Ser. B Biol. Sci.* **319**, 505–523. (doi:10.1098/rstb.1988.0062)

4. Endler JA, Mappes J. 2004 Predator Mixes and the Conspicuousness of Aposematic Signals. *Am. Nat.* **163**, 532–547. (doi:10.1086/382662)

5. Mallet JLB, Barton NH. 1989 Strong Natural Selection in a Warning-Color Hybrid Zone. *Evolution (N. Y).* **43**, 421. (doi:10.2307/2409217)

6. Sherratt TN, Speed MP, Ruxton GD. 2004 Natural selection on unpalatable species imposed by state-dependent foraging behaviour. *J. Theor. Biol.* **228**, 217–226. (doi:10.1016/j.jtbi.2003.12.009)

7. Brower LP, Ryerson WN, Coppinger LL, Glazier SC. 1968 Ecological Chemistry and the Palatability Spectrum. *Science.* **161**, 1349–1350. (doi:10.1126/science.161.3848.1349)

8. Endler JA. 1991 Interactions between predators and prey. In *Behavioural Ecology* (eds JR Krebs, NB Davies), pp. 169–196. Oxford: Blackwell Scientific.

9. Karpestam E, Merilaita S, Forsman A. 2014 Natural levels of colour polymorphism reduce performance of visual predators searching for camouflaged prey. *Biol. J. Linn. Soc.* **112**, 546–555. (doi:10.1111/bij.12276)

10. Mappes J, Marples N, Endler JA. 2005 The complex business of survival by aposematism. *Trends Ecol. Evol.* **20**, 598–603. (doi:10.1016/j.tree.2005.07.011)

11. Briolat ES, Burdfield-Steel ER, Paul SC, Katja HR, Seymoure BM, Stankowich T, Stuckert AMM. 2019 Diversity in warning coloration: selective paradox or the norm? *Biol. Rev.* **94**, 388–414. (doi:10.1111/brv.12460)

12. Speed MP, Turner JRG. 1999 Learning and memory in mimicry: II. Do we understand the mimicry spectrum? *Biol. J. Linn. Soc.* **67**, 281–312. (doi:10.1006/bijl.1998.0310)

13. Turner JRG. 1981 Adaptation and Evolution in Heliconius: A Defense of NeoDarwinism. *Annu. Rev. Ecol. Syst.* **12**, 99–121. (doi:10.1146/annurev.es.12.110181.000531)

14. Clarke B, O'Donald P. 1964 Frequency-dependent selection. *Heredity (Edinb).* **19**, 201–206. (doi:10.1038/hdy.1964.25)

15. Greenwood JJD. 1985 Frequency-Dependent Selection by Seed-Predators. *Oikos* **44**, 195.

485   (doi:10.2307/3544062)

16.   Fisher RA. 1930 *The genetical theory of natural selection.* Oxford: Clarendon Press.
(doi:10.5962/bhl.title.27468)

17.   Guilford T. 1985 Is Kin Selection Involved in the Evolution of Warning Coloration? *Oikos*
45, 31. (doi:10.2307/3565219)

18.   Barnett JB, Michalis C, Scott-Samuel NE, Cuthill IC. 2021 Colour pattern variation forms
local background matching camouflage in a leaf-mimicking toad. *J. Evol. Biol.* **34**, 1531–
1540. (doi:10.1111/jeb.13923)

19.   Rönkä K, Valkonen J, Nokelainen O, Rojas B, Gordon S, Burdfield-Steel E, Mappes J.
2020 Geographic mosaic of selection by avian predators on hindwing warning colour in
a polymorphic aposematic moth. *Ecol. Lett.* (doi:10.1101/2020.04.08.032078)

20.   Sherratt TN. 2002 The coevolution of warning signals. *Proc. R. Soc. B Biol. Sci.* **269**,
741–746. (doi:10.1098/rspb.2001.1944)

21.   Summers K, Speed MP, Blount JD, Stuckert AMM. 2015 Are aposematic signals honest?
A review. *J. Evol. Biol.* **28**, 1583–1599. (doi:10.1111/jeb.12676)

22.   White TE, Umbers KDL. 2021 Meta-analytic evidence for quantitative honesty in
aposematic signals. *Proc. R. Soc. B Biol. Sci.* **288**. (doi:10.1098/rspb.2021.0679)

23.   Banich MT, Caccamise D. 2010 *Generalization of knowledge : multidisciplinary*
perspectives. New York: Psychology Press.

24.   Gamberale G, Tullberg BS. 1996 Evidence for a peak-shift in predator generalization
among aposematic prey. *Proc. Biol. Sci.* **263**, 1329–1334. (doi:10.1098/rspb.1996.0195)

25.   Rowe C. 1999 Receiver psychology and the evolution of multicomponent signals. *Anim.*
Behav. **58**, 921–931. (doi:10.1006/anbe.1999.1242)

26.   Winters AE, Green NF, Wilson NG, How MJ, Garson MJ, Marshall NJ, Cheney KL. 2017
Stabilizing selection on individual pattern elements of aposematic signals. *Proc. R. Soc.*
B Sci. **284**.

27.   Endler JA. 1978 A predator's view of animal color patterns. *Evol. Biol.* **11**, 320–364.
(doi:10.1007/978-1-4615-6956-5_5)

28.   Barnett JB, Scott-Samuel NE, Cuthill IC. 2016 Aposematism: balancing salience and
camouflage. *Biol. Lett.* **12**, 20160335. (doi:10.1098/rsbl.2016.0335)

29.   Endler JA, Houde AE. 1995 Geographic variation in female preferences for male traits in
Poecilia reticulata. Evolution (N. Y). **49**, 456. (doi:10.2307/2410270)

30.   Rosselli FB, Alemi A, Ansuini A, Zoccolan D. 2015 Object similarity affects the
perceptual strategy underlying invariant visual object recognition in rats. *Front. Neural*

519   *Circuits* **9**, 1–22. (doi:10.3389/fncir.2015.00010)

520 31. Sibeaux A, Cole GL, Endler JA. 2019 The relative importance of local and global visual

521   contrast in mate choice. *Anim. Behav.* **154**, 143–159. (doi:10.1016/j.anbehav.2019.06.020)

522 32. van den Berg CP, Endler JA, Papinczak DEJ, Cheney KL. 2022 Using colour pattern edge

523   contrast statistics to predict detection speed and success in triggerfish (*Rhinecanthus*

524   *aculeatus*). *J. Exp. Biol.* (doi:10.1242/jeb.244677)

525 33. Rönkä K, De Pasqual C, Mappes J, Gordon S, Rojas B. 2018 Colour alone matters: no

526   predator generalization among morphs of an aposematic moth. *Anim. Behav.* **135**, 153–

527   163. (doi:10.1016/j.anbehav.2017.11.015)

528 34. Nokelainen O, Galarza JA, Kirvesoja J, Suisto K, Mappes J. 2022 Genetic colour variation

529   visible for predators and conspecifics is concealed from humans in a polymorphic moth.

530   *J. Evol. Biol.* **35**, 467–478. (doi:10.1111/jeb.13994)

531 35. Winters AE, Chan W, White AM, Berg CP, Garson MJ, Cheney KL. 2022 Weapons or

532   deterrents? Nudibranch molluscs use distinct ecological modes of chemical defence

533   against predators. *J. Anim. Ecol.* **91**, 831–844. (doi:10.1111/1365-2656.13643)

534 36. Endler JA. 1986 Defense Against Predators. In *Predator-prey relationships, Perspectives*

535   *and Approaches from the Study of Lower Vertebrates*, pp. 109–134. Chicago: University

536   of Chicago Press.

537 37. van den Berg CP, Troscianko J, Endler JA, Marshall NJ, Cheney KL. 2020 Quantitative

538   Colour Pattern Analysis (QCPA): A comprehensive framework for the analysis of colour

539   patterns in nature. *Methods Ecol. Evol.* **11**, 316–332. (doi:10.1111/2041-210X.13328)

540 38. van den Berg CP, Endler JA, Cheney KL. 2022 Highly defended nudibranchs do not

541   'escape' to visually diverse background habitats. *bioRxiv*

542   (doi:https://doi.org/10.1101/2022.12.13.520332)

543 39. van den Berg CP, Endler JA, Cheney KL. 2022 Signal detectability and boldness are not

544   the same: the function of defensive colouration in nudibranchs is distance-dependent.

545   *bioRxiv* (doi:https://doi.org/10.1101/2022.12.20.521213)

546 40. Schneider CA, Rasband WS, Eliceiri KW. 2012 NIH Image to ImageJ: 25 years of image

547   analysis. *Nat. Methods* **9**, 671–675. (doi:10.1038/nmeth.2089)

548 41. Troscianko J, Stevens M. 2015 Image calibration and analysis toolbox – a free software

549   suite for objectively measuring reflectance, colour and pattern. *Methods Ecol. Evol.* **6**,

550   1320–1331. (doi:10.1111/2041-210X.12439)

551 42. Randall JE, Allen GR, Steene RC. 1997 *Fishes of the Great Barrier Reef and Coral Sea*.

552   Crawford House Publishing.

43. Champ CM, Wallis G, Vorobyev M, Siebeck U, Marshall J. 2014 Visual acuity in a
species of coral reef fish: *Rhinecanthus aculeatus*. *Brain. Behav. Evol.* **83**, 31–42.
(doi:10.1159/000356977)

44. Champ CM, Vorobyev M, Marshall NJ. 2016 Colour thresholds in a coral reef fish. *R.*
Soc. Open Sci. **3**, 160399. (doi:10.1098/rsos.160399)

45. Cheney KL, Newport C, McClure EC, Marshall NJ. 2013 Colour vision and response bias
in a coral reef fish. *J. Exp. Biol.* **216**, 2967–2973. (doi:10.1242/jeb.087932)

46. Pignatelli V, Champ CM, Marshall J, Vorobyev M. 2010 Double cones are used for colour
discrimination in the reef fish, *Rhinecanthus aculeatus*. *Biol. Lett.* **6**, 537–539.
(doi:10.1098/rsbl.2009.1010)

47. Cheney KL, Hudson J, de Busserolles F, Luehrmann M, Shaughnessy A, van den Berg C,
Green NF, Marshall NJ, Cortesi F. 2022 Seeing Picasso: an investigation into the visual
system of the triggerfish *Rhinecanthus aculeatus*. *J. Exp. Biol.* **225**, jeb243907.
(doi:10.1242/jeb.243907)

48. Green NF, Guevara E, Osorio DC, Endler JA, Marshall NJ, Vorobyev M, Cheney KL.
2022 Colour discrimination thresholds vary throughout colour space in a reef fish
(*Rhinecanthus aculeatus*). *J. Exp. Biol.* **225**, jeb243533. (doi:10.1242/jeb.243533)

49. Cheney KL, Green NF, Vibert AP, Vorobyev M, Marshall NJ, Osorio DC, Endler JA.
2019 An Ishihara-style test of animal colour vision. *J. Exp. Biol.* **222**, jeb189787.
(doi:10.1242/jeb.189787)

50. van den Berg CP, Hollenkamp M, Mitchell LJ, Watson EJ, Green NF, Marshall NJ,
Cheney KL. 2020 More than noise: Context-dependant luminance contrast discrimination
in a coral reef fish (*Rhinecanthus aculeatus*). *J. Exp. Biol.* **223**, jeb.232090.
(doi:10.1242/jeb.232090)

51. van den Berg CP, Condon ND, Conradsen CI, White TE, Cheney KL. 2023 Automated
workflows using Quantitative Colour Pattern Analysis (QCPA): A guide to batch
processing and downstream data analysis. *bioRxiv*
(doi:https://doi.org/10.1101/2023.02.02.526788)

52. Endler JA. 2012 A framework for analysing colour pattern geometry: Adjacent colours.
Biol. J. Linn. Soc. **107**, 233–253. (doi:10.1111/j.1095-8312.2012.01937.x)

53. Endler JA, Cole GL, Kranz AM. 2018 Boundary strength analysis: Combining colour
pattern geometry and coloured patch visual properties for use in predicting behaviour and
fitness. *Methods Ecol. Evol.* **9**, 2334–2348. (doi:10.1111/2041-210X.13073)

54. Hubel DH, Wiesel TN. 1962 Receptive fields, binocular interaction and functional

architecture in the cat's visual cortex. *J. Physiol.* **160**, 106–154.
(doi:10.1113/jphysiol.1962.sp006837)

55. Chan W, Shaughnessy AEP, van den Berg CP, Garson MJ, Cheney KL. 2021 The Validity
of Brine Shrimp (Artemia Sp.) Toxicity Assays to Assess the Ecological Function of
Marine Natural Products. *J. Chem. Ecol.* **47**, 834–846. (doi:10.1007/s10886-021-01264-
z)

56. Cheney KL, White A, Mudianta IW, Winters AE, Quezada M, Capon RJ, Mollo E, Garson
MJ. 2016 Choose your weaponry: Selective storage of a single toxic compound,
latrunculin a, by closely related nudibranch molluscs. *PLoS One* **11**, 1–16.
(doi:10.1371/journal.pone.0145134)

57. Chan W. 2022 The Chemical Ecology of Nudibranchs.

58. Winters AE, White AM, Dewi AS, Mudianta IW, Wilson NG, Forster LC, Garson MJ,
Cheney KL. 2018 Distribution of Defensive Metabolites in Nudibranch Molluscs. *J.*
Chem. Ecol. **44**, 384–396. (doi:10.1007/s10886-018-0941-5)

59. Kuhn M. 2021 caret: Classification and Regression Training. R package version 6.0-90.
https://CRAN.R-project.org/package=caret.

60. Cheney KL, Cortesi F, How MJ, Wilson NG, Blomberg SP, Winters AE, Umanzör S,
Marshall NJ. 2014 Conspicuous visual signals do not coevolve with increased body size
in marine sea slugs. *J. Evol. Biol.* **27**, 676–87. (doi:10.1111/jeb.12348)

61. Cortesi F, Cheney KL. 2010 Conspicuousness is correlated with toxicity in marine
opisthobranchs. *J. Evol. Biol.* **23**, 1509–18. (doi:10.1111/j.1420-9101.2010.02018.x)

62. R Core Team. 2021 R: A Language and Environment for Statistical Computing.

63. Peterson RA, Cavanaugh JE. 2020 Ordered quantile normalization: a semiparametric
transformation built for the cross-validation era. *J. Appl. Stat.* **47**, 2312–2327.
(doi:10.1080/02664763.2019.1630372)

64. Moran MD. 2003 Arguments for rejecting the sequential Bonferroni in ecological studies.
Oikos **100**, 403–405. (doi:10.1034/j.1600-0706.2003.12010.x)

65. Hempel de Ibarra N, Giurfa M, Vorobyev M. 2001 Detection of coloured patterns by
honeybees through chromatic and achromatic cues. *J. Comp. Physiol. A Sensory, Neural,*
Behav. Physiol. **187**, 215–224. (doi:10.1007/s003590100192)

66. Endler JA, Mielke PW. 2005 Comparing entire colour patterns as birds see them. *Biol. J.*
Linn. Soc. **86**, 405–431. (doi:10.1111/j.1095-8312.2005.00540.x)

67. Ruxton GD, Allen WL, Sherratt TN, Speed MP. 2018 *Avoiding Attack*. New York: Oxford
University Press. (doi:10.1093/oso/9780199688678.001.0001)

68. Kenward B, Wachtmeister CA, Ghirlanda S, Enquist M. 2004 Spots and stripes: The evolution of repetition in visual signal form. *J. Theor. Biol.* **230**, 407–419. (doi:10.1016/j.jtbi.2004.06.008)

69. Aronsson M, Gamberale-Stille G. 2009 Importance of internal pattern contrast and contrast against the background in aposematic signals. *Behav. Ecol.* **20**, 1356–1362. (doi:10.1093/beheco/arp141)

70. Osorio D, Vorobyev M. 2008 A review of the evolution of animal colour vision and visual communication signals. *Vision Res.* **48**, 2042–2051. (doi:10.1016/j.visres.2008.06.018)

71. Barnett JB, Cuthill IC. 2015 Distance-dependent defensive coloration. *Curr. Biol.* **24**, R1157–R1158. (doi:10.1016/j.cub.2014.11.015)

72. Marshall NJ. 2000 Communication and camouflage with the same 'bright' colours in reef fishes. *Philos. Trans. R. Soc. London. Ser. B Biol. Sci.* **355**, 1243–1248. (doi:10.1098/rstb.2000.0676)

73. Hebets EA, Papaj DR. 2005 Complex signal function: Developing a framework of testable hypotheses. *Behav. Ecol. Sociobiol.* **57**, 197–214. (doi:10.1007/s00265-004-0865-7)

74. Falconer DS, Mackay TFC. 1996 *Introduction to quantitative genetics.* 4th ed. /. Burnt Mill, England: Longman.

75. Lande R, Arnold SJ. 1983 The Measurement of Selection on Correlated Characters. *Evolution (N. Y).* **37**, 1210. (doi:10.2307/2408842)

76. Layton KKS, Gosliner TM, Wilson NG. 2018 Flexible colour patterns obscure identification and mimicry in Indo-Pacific Chromodoris nudibranchs (Gastropoda: Chromodorididae). *Mol. Phylogenet. Evol.* **124**, 27–36. (doi:10.1016/j.ympev.2018.02.008)

77. Debelius H, Kuiter RH. 2007 *Nudibranchs of the world*. Frankfurt, Germany: IKAN-Unterwasserarchiv.

78. Losey GS, McFarland WN, Loew ER, Zamzow JP, Nelson P a., Marshall NJ. 2003 Visual biology of Hawaiian coral reef fishes. I. Ocular transmission and visual pigments. *Copeia* **2003**, 433–454. (doi:10.1643/01-053)

79. Stevens M. 2007 Predator perception and the interrelation between different forms of protective coloration. *Proc. R. Soc. B Biol. Sci.* **274**, 1457–64. (doi:10.1098/rspb.2007.0220)

80. Nokelainen O, Valkonen J, Lindstedt C, Mappes J. 2014 Changes in predator community structure shifts the efficacy of two warning signals in Arctiid moths. *J. Anim. Ecol.* **83**, 598–605. (doi:10.1111/1365-2656.12169)

81. Postema EG, Lippey MK, Armstrong-Ingram T. 2022 Color under pressure: how multiple factors shape defensive coloration. *Behav. Ecol.* , 1–13. (doi:10.1093/beheco/arac056)

82. De Pasqual C, Suisto K, Kirvesoja J, Gordon S, Ketola T, Mappes J. 2022 Heterozygote advantage and pleiotropy contribute to intraspecific color trait variability. *Evolution (N. Y).* , 2389–2403. (doi:10.1111/evo.14597)

83. Cadena V, Rankin K, Smith KR, Endler JA, Stuart-Fox D. 2018 Temperature-induced colour change varies seasonally in bearded dragon lizards. *Biol. J. Linn. Soc.* **123**, 422–430. (doi:10.1093/biolinnean/blx152)

84. Larkin MF, Smith SDA, Willan RC, Davis TR. 2018 Diel and seasonal variation in heterobranch sea slug assemblages within an embayment in temperate eastern Australia. *Mar. Biodivers.* **48**, 1541–1550. (doi:10.1007/s12526-017-0700-9)

85. Schubert J, Smith SDA. 2020 Sea slugs—"rare in space and time"—but not always. *Diversity* **12**, 1–14. (doi:10.3390/d12110423)

86. Smith SDA, Davis TR. 2019 Slugging it out for science: volunteers provide valuable data on the diversity and distribution of heterobranch sea slugs. *Molluscan Res.* **39**, 214–223. (doi:10.1080/13235818.2019.1594600)

87. Barnett CA, Bateson M, Rowe C. 2007 State-dependent decision making: Educated predators strategically trade off the costs and benefits of consuming aposematic prey. *Behav. Ecol.* **18**, 645–651. (doi:10.1093/beheco/arm027)

88. Rojas B, Burdfield-Steel E, Pakkanen H, Suisto K, Maczka M, Schulz S, Mappes J. 2017 How to fight multiple enemies: Target-specific chemical defences in an aposematic moth. *Proc. R. Soc. B Biol. Sci.* **284**. (doi:10.1098/rspb.2017.1424)

89. Raviv L, Lupyan G, Green SC. 2022 How variability shapes learning and generalization. *Trends Cogn. Sci.* **26**, 462–483. (doi:10.1016/j.tics.2022.03.007)

90. Eakin RM, Westfall JA, Dennis MJ. 1967 Fine Structure of the Eye of a Nudibranch Mollusc, *Hermissenda Crassicornis. J. Cell Sci.* **2**, 349–358. (doi:10.1242/jcs.2.3.349)

91. Yeager J, Barnett JB. 2022 Continuous Variation in an Aposematic Pattern Affects Background Contrast, but Is Not Associated With Differences in Microhabitat Use. *Front. Ecol. Evol.* **10**, 1–10. (doi:10.3389/fevo.2022.803996)

92. Endler JA, Mappes J. 2017 The current and future state of animal coloration research. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, 20160352. (doi:10.1098/rstb.2016.0352)

93. Kemp DJ, Herberstein ME, Fleishman LJ, Endler JA, Bennett ATD, Dyer AG, Hart NS, Marshall NJ, Whiting MJ. 2015 An integrative framework for the appraisal of coloration in nature. *Am. Nat.* **185**, 705–724. (doi:10.1086/681021)

689   94.   Olsson P, Lind O, Kelber A. 2018 Chromatic and achromatic vision: parameter choice
690   and limitations for reliable model predictions. *Behav. Ecol.* **29**, 273–282.
691   (doi:10.1093/beheco/arx133)

692   95.   Silvasti SA, Valkonen JK, Nokelainen O. 2021 Behavioural thresholds of blue tit colour
693   vision and the effect of background chromatic complexity. *Vision Res.* **182**, 46–57.
694   (doi:10.1016/j.visres.2020.11.013)

695   96.   Blount JD, Speed MP, Ruxton GD, Stephens P a. 2009 Warning displays may function as
696   honest signals of toxicity. *Proc. Biol. Sci.* **276**, 871–877. (doi:10.1098/rspb.2008.1407)

697