DLK-dependent mitochondrial fission drives axon degeneration and neuronal cell death

Jorge Gómez-Deza J [1], Matthew Nebiyou[1], Mor R. Alkaslasi[1], Preethi Somasundaran[2], Anastasia
L. Slavutsky[1], Michael E. Ward[3], Trent A. Watkins[2], Claire E. Le Pichon[1]

[1] *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National
Institutes of Health, Bethesda, United States
[2] Departments of Neurosurgery and Neuroscience, Baylor College of Medicine, Houston, Texas
[3] National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda,
United States

Abstract

Currently there are no effective treatments for an array of neurodegenerative disorders to a large
part because cell-based models fail to recapitulate disease. Here we developed a robust human
iPSCbased model where laser axotomy causes retrograde axon degeneration leading to neuronal
cell death. Time-lapse confocal imaging revealed that damage triggers a wave of mitochondrial
fission proceeding from the site of injury to the soma. We demonstrated that mitochondrial
fission and resultant cell death is entirely dependent on phosphorylation of dynamin related
protein 1 (DRP1) by dual leucine zipper kinase (DLK). Importantly, we show that CRISPR mediated
Drp1 depletion protected mouse retinal ganglion neurons from mitochondrial fission and
degeneration after optic nerve crush. Our results provide a powerful platform for studying
degeneration of human neurons, pinpoint key early events in damage related neural death and
new focus for therapeutic intervention.

Introduction

Neuronal death and axon loss are universal hallmarks of neurodegeneration, relevant to Alzheimer's disease, ALS and other neurodegenerative diseases. However, our understanding of the mechanisms controlling axon degeneration and neuron death is incomplete. Advances in this area have enormous potential to benefit all types of neurodegenerative conditions including those arising from nerve injury and traumatic brain injury.

Separate mechanisms regulate axon degeneration and neuronal apoptosis[1]. For example, deletion of BAX (BCL-2-associated X protein), a central regulator of apoptosis, does not protect axons from Wallerian degeneration. On the other hand, BAX deletion protects neurons following trophic factor withdrawal, a developmental model of axon degeneration and death[2]. SARM1 is the central executioner of Wallerian axon degeneration[3,4]. However, it is unclear how universal this pathway is for axon degeneration since some neurodegenerative models are not rescued by loss of SARM1[5,6]. Furthermore, while most neurodegenerative diseases involve loss of axons and cell bodies, Wallerian axon degeneration does not regulate neuronal cell death.

Mitochondria are a central hub of axon health regulation, at the nexus of axonal energy production, oxidative stress, and calcium buffering[7,8]. They also play critical roles in degeneration and regeneration[9,10]. Healthy maintenance of the mitochondrial network is mediated by a balance of mitochondrial fission and fusion regulated by specialized machinery[11]. Interestingly, mutations in some of these genes cause neurodegenerative conditions such as motor and sensory neuropathy (eg. Mfn1 in Charcot-Marie-Tooth[12]). A key player in the mitochondrial fission process is dynamin related protein 1 or DRP1, a GTPase that oligomerizes around the outer mitochondrial membrane to cause mitochondrial fission[13]. DRP1 also regulates apoptotis[11,14–16]. Under conditions that drive apoptosis, DRP1 promotes a type of apoptotic mitochondrial fission in which BAX permeabilizes the outer mitochondrial membrane, allowing the release of pro-apoptotic factors such as cytochrome c[17,18].

Much of what we know about mechanisms regulating neuron degeneration and death comes from a wealth of studies of apoptosis in cell lines (but not neurons), or from animal models (neurons, but not human). How translatable this work is and how applicable to neurons remains to be determined. One well established model of retrograde axon degeneration is a trophic factor deprivation model, in which embryonic mouse primary neurons are deprived of NGF (nerve growth factor), which leads to their degeneration[19,20]. Recent advances in iPSC-derived neuron techniques now enable the generation of a diverse range of human neuron subtypes. These efforts have already generated exciting findings in the neurodegeneration field[21–23]. We therefore set out to study axon injury using a widely-adopted glutamatergic neuron model (i3Neurons[24]).

We and others have shown that the dual leucine zipper kinase (DLK, or MAP3K12), is a key regulator of axon degeneration and a driver of neuron cell death[25–29]. The canonical DLK pathway involves a phosphorylation cascade downstream of DLK homodimerization and a transcriptional

71 stress response, including phosphorylation of the transcription factor cJun[29]. Despite its
72 consideration as a drug target for several neurological conditions, relatively little is known about
73 DLK function in human neurons.
74
Here, we investigate the behavior of i3Neurons following laser axotomy, and the role of DLK in
the injury response of human neurons. Following axotomy, we observe progressive retrograde
axon degeneration from the injury site back towards the cell body, causing subsequent neuronal
death, establishing a unique human model of neurodegeneration. This contrasts with the
behavior of cultured mouse neurons, which survive and regenerate after this type of injury[30–32],
but resembles the pattern of neurodegeneration observed in optic nerve injuries in living mice[26].
81
In axons proximal to the injury, mitochondria shrink and undergo DRP1-dependent fission in a
retrograde wave, leading to apoptosis of the cell body. DLK regulates mitochondrial fission via
direct phosphorylation of DRP1 at serine 616. Such DRP1 phosphorylation by DLK is necessary for
localization of BAX to axonal mitochondria, which causes axon degeneration. Blocking DLK or
DRP1 protects axons from degeneration and neurons from cell death both in human i3Neurons
in vitro and *in vivo* in the mouse. This interaction between DLK and DRP1 identifies a new
substrate for DLK, a new kinase for DRP1, and a novel signaling pathway for DLK. Collectively, our
findings reveal an apoptotic cascade that is initiated in the axon that both underlies non-
Wallerian axon degeneration and drives neuronal cell death.

Results

Axotomy in human neurons causes both proximal and distal axon degeneration

Axotomy has often been used as a method to study axon degeneration. After axotomy, the distal portion of the axon undergoes Wallerian degeneration, a phenomenon controlled by the NADase SARM1, the key executioner of this type of axon degeneration[3,4]. To study the response of human neurons to axotomy, we performed laser axotomies in i3Neurons that were sparsely transduced with a fluorescent protein to visualize individual neurons (Fig. 1A). The distal portion of these axons underwent Wallerian degeneration as expected (Fig. 1B, C). To our surprise, we observed that the portion of the axon proximal to the injury and still connected to the cell body underwent progressive retrograde axon degeneration (Fig. 1B-C). Because SARM1-dependent degeneration relies on the loss of the axon survival factor NMNAT2, which occurs when the axon is severed from the cell body, we hypothesized that the proximal axon degeneration we observed might not rely on SARM1 and might provide an opportunity to elucidate novel pathways of axon degeneration.

We tested the requirement for SARM1 in the degeneration of axons both proximal and distal to the injury site. Using CRISPR/Cas9, we knocked out SARM1 in i3Neurons [33]. We then axotomized these neurons and control neurons and imaged the axon on both sides of the injury. As expected, SARM1 was necessary for distal axon degeneration, but not proximal axon degeneration (Fig. 1B-C). We therefore decided to investigate the mechanisms underlying proximal axon degeneration in these neurons. For the remainder of the study, we focus specifically on the axon segment proximal to the injury site.

Axotomy in human neurons triggers a wave of DRP1-dependent mitochondrial fission

Mitochondria play pivotal roles in axon degeneration, due to their importance in energy production, calcium homeostasis, production of reactive oxygen species and signaling[34,35]. We transduced i3Neurons with a mitochondrial-targeted GFP and plated them in the center of wells to allow axons to project outwards and visualize individual axons. After axotomy, mitochondria underwent rapid shrinkage (Fig. 1D-G; Video S1) and increased in number (Fig. 1H). Using live imaging, we determined that the increase in mitochondrial number was due to fission events rather than mitochondrial transport and localization to the site of injury. Mitochondrial fission was most evident from close analysis of live imaging (Video S1 and Fig. 1D-E).

Mitochondrial fission is a dynamic process that is regulated by proteins localized in the inner and outer mitochondrial membranes[13]. A critical regulator of fission is dynamin related protein 1 (DRP1), a cytoplasmic GTPase. DRP1 is recruited to the outer mitochondrial membrane and helically oligomerizes around the mitochondrion to constrict it and bring about scission of the membrane. To test whether the fission we observed was dependent on DRP1, we knocked down DRP1 using CRISPR interference by expressing a previously validated guide RNA to DRP1 in i3Neurons expressing a catalytically dead Cas9 (dCAS9)[21,33]. DRP1 knockdown prevented the increase in mitochondrial particle number (Fig. 1H). We confirmed this result by inhibiting DRP1 using P110, a peptide that blocks the interaction between DRP1 and FIS1 (mitochondrial fission

135   protein 1), another protein that is essential for fission (Fig. 1I). In line with previous reports[36,37],
136   these events were dependent on calcium influx into the injured axon which, when buffered using
137   BAPTA-AM, also prevented mitochondrial fission (Fig. S1A-D; Videos S2-5).
138
139
Mitochondrial fission precedes proximal axon degeneration
To characterize the spatial and temporal dynamics of the axonal response to injury, we analyzed
mitochondrial fission and axon degeneration in 100 μm segments of the axon relative to the
injury site every 10 min following axotomy. This analysis revealed that mitochondrial fission
began seconds after injury and progressed in a retrograde manner back towards the cell body.
One hour after axotomy, mitochondria 500 μm away from the injury had undergone fission (Fig.
1J-K Video S6). This fission preceded a slower retrograde wave of axon degeneration back
towards the cell body (Fig. 1L-M, Video S7).
148
149   Time lapse imaging of the cell bodies of axotomized neurons revealed that the retrograde wave
150   of axon degeneration was followed by fragmentation of mitochondria in the soma and cell death
151   (Fig. 1N-O; Videos S8-9). Axotomy led to death of 84% of cells analyzed. We have thus uncovered
152   an exciting new human neuron model in which axon injury initiates progressive dying back of the
153   axon and ultimately results in neuronal cell death.
154
155
DLK regulates axotomy-induced mitochondrial fission
To understand mechanisms governing neurodegeneration in this model, we started by examining
a pathway involved in the cellular response to axon injury. The dual leucine zipper kinase (DLK;
MAP3K12) is a key regulator of the axon damage response and neuronal cell death [29,38]. Because
mitochondria are critical regulators of axon degeneration, we were curious whether DLK protein
might localize to axonal mitochondria. In i3Neurons, we observed association of DLK with 77% of
axonal mitochondria (Fig S2 A-C). We further corroborated this association by immuno-EM (Fig.
S2D), and through observation of DLK co-trafficking with mitochondria (Fig. S2E).
164
To investigate how DLK responds to axon damage in human neurons, we performed laser
axotomies and live imaging of DLK and mitochondria. We observed an intriguing presence of DLK
puncta at 45% of sites of mitochondrial fission (Fig. 2A-B, Video S10). To determine whether DLK
influences mitochondrial fission, we generated DLK KO neurons using CRISPR/Cas9 (Fig. S2F). The
knockout neurons were validated by immunofluorescent staining and western blotting (Fig. S2G-
H). In the absence of injury, we observed no mitochondrial phenotypes in these cells including
number, size, movement, and oxygen consumption rate (Fig. S2I-P). Strikingly, DLK KO neurons
did not exhibit the axonal mitochondrial fission we had observed in response to axotomy in wild
type neurons (Fig. 2C-D; Videos S11-12). Corroborating this result, pharmacological inhibition of
DLK using GNE3511 prevented mitochondrial fission after injury (Fig. 2E, Videos S13-14).
175
To better understand the mechanism by which DLK regulates mitochondrial fission, DLK KO
neurons were transduced with wild type DLK or 2 mutant versions of DLK. DLK C127S cannot be
palmitoylated, hindering its ability to properly signal[39] . Kinase dead DLK (S302A) cannot be trans-

phosphorylated and is catalytically inactive [27,40]. Only the expression of wild type DLK rescued the
fission phenotype in DLK KO neurons (Fig. 2F; Videos S15-17), indicating that DLK kinase activity
is critical for this process. We thus uncover a role for DLK in regulating mitochondrial fission after
injury.
183
184
DLK directly phosphorylates DRP1 to induce mitochondrial fission
DRP1 phosphorylation state regulates its function state, with phosphorylation at serine 616
promoting mitochondrial fission[41]. We speculated that DLK might phosphorylate DRP1 at this
residue. In order to more easily study this signaling mechanism, we turned to a heterologous
system, HEK293 cells, that do not express DLK. Overexpression of DLK in these cells can be used
to study DLK signaling since its homodimerization is sufficient to drive pathway activity [40,42].
191
Expression of wild type DLK, but not DLK C127S (palmitoyl-site mutant) or DLK S302A (kinase
dead), resulted in DLK signaling as reflected by the increased phosphorylation of its downstream
target c-Jun (Fig. 3 A-B). Interestingly, when we co-expressed DLK and DRP1, phosphorylation at
S616 was increased in this model of DLK activation (Fig. 3 A, C), but not with mutant DLK, showing
that DLK activity is sufficient to cause DRP1 S616 phosphorylation. Levels of total DRP1 remained
unchanged, as did levels of S637 pDRP1, a site whose phosphorylation is thought to impair DRP1
GTPase activity [43,44] (Fig. 3 A, D-E).
199
To further validate that DLK signaling causes DRP1 phosphorylation, we purified GST-DRP1 from
HEK293 cells expressing wild type DLK or DLK S302A, and performed targeted mass spectrometry
to quantify DRP1 phosphorylation (Fig. 3F). The relative abundance of phospho-DRP1 S616
peptides was significantly increased only in the HEK293 cells expressing wild type but not kinase
dead DLK (Fig. 3G). These results demonstrate that DLK kinase activity can promote DRP1
phosphorylation at S616 either directly or indirectly.
206
207   Next, to determine whether DLK can directly phosphorylate DRP1, we performed an *in vitro*
208   kinase assay using proteins separately purified from HEK293 cells. Remarkably, incubation of
209   DRP1 with wild type DLK increased pDRP1 S616, whereas incubation with kinase dead DLK did
210   not (Fig. 3H,I). We thus demonstrate that DLK can directly phosphorylate DRP1 at serine 616.
211
To link this phosphorylation of DRP1 to a potential effect on mitochondria, we analyzed
mitochondrial morphology in the HEK293 cells. Similar to what we had observed in injured axons,
DLK signaling resulted in an increase in mitochondrial fragmentation (Fig. 3 J,K). DLK-induced
mitochondrial fragmentation was prevented by knocking down DRP1 (Fig. 3 K-N). These data
identify DLK as a novel kinase of DRP1 whose role is to drive mitochondrial fission via direct
phosphorylation of DRP1 at serine 616.
218
219
Axotomy causes DLK-dependent phosphorylation of DRP1
To validate these results in the i3Neurons, DLK KO or WT neurons were center-plated such that
axon fractions could be harvested to examine axonal DRP1 phosphorylation after injury (Fig. 4A).

Two hours after axotomy, phospho-DRP1 S616 (pDRP1 S616) was increased in axons but not cell
bodies of wild type neurons. This did not occur in DLK KO neurons (Fig. 4B-E). The increase in
pDRP1 S616 in wild type but not DLK KO axons was further validated by immunostaining (Fig. 4F-
G).
227
To examine whether DRP1 phosphorylation at S616 also occurs *in vivo*, we performed optic nerve
crush in mice. Increased pDRP1 S616 relative to total DRP1 was detected in optic nerve proximal
to the crush site 24 hours after injury (Fig. 4H-J). Levels of total DRP1 in injured nerves remained
unchanged (Fig. 4I,K). We next asked whether DLK also regulates the phosphorylation of DRP1 *in*
vivo after injury. Optic nerve crush was performed in mice treated with vehicle or DLK inhibitor
GNE-3511. We observed a reduction in the levels of pDRP1 S616 relative to total DRP1 in mice
treated with GNE-3511 compared to the vehicle controls, demonstrating that axon injury triggers
DLK-dependent DRP1 phosphorylation *in vivo* as well as *in vitro.*
236
237
Inhibiting DRP1 delays axon degeneration and neuron death
Since axotomy leads to progressive degeneration of the axon proximal to the injury and
eventually to cell death, we asked to what extent DLK and DRP1 control the degeneration of each
compartment. DLK has been shown to contribute to axon degeneration and to neuron death in
many contexts [25–29,45]. We hypothesized that DLK would promote axon degeneration in
axotomized human neurons. Whereas wild type axons had degenerated as early as 4 hours after
injury, DLK KO axons were still present after 24 hours (Fig. 5A-C). Interestingly, in neurons with
CRISPRi knockdown of DRP1, axons were also protected from degeneration (Fig. 5D-F).
246
247 To determine whether loss of DLK or DRP1 also protected neurons from death, we followed the
248 fate of cell bodies of axotomized neurons. In control neurons we observed cell death in about
249 70% of cells after axotomy whereas only 18% of DLK KO cells and 33% of DRP1 knockdown cells
250 died (Fig 5G-L). These results demonstrate that DRP1 and DLK are key regulators of neuronal cell
251 death after axotomy.
252
253
Loss of DRP1 prevents neuron death after axotomy *in vivo*
We then asked how universal this pathway is, or whether it was unique to the i3Neuron model.
It has been established that silencing DLK blocks retinal ganglion cell (RGC) death after optic nerve
crush [26,27,45,46]. However, DRP1 has not previously been implicated downstream of this DLK
signaling. To knock down DRP1, we used an AAV delivery strategy to express a small Cas9 (sa-
Cas9) together with a guide RNA to DRP1 [47]. AAVs were injected into the intravitreal space to
transduce RGCs. Optic nerve crush was performed two weeks later, and retinas were harvested
3 days post injury, a time point at which injured RGCs express cleaved caspase 3 (Fig. 5M). The
average number of cleaved caspase 3-positive RGCs with saCas9 and a control gRNA was similar
to a control virus expressing just EGFP (55 ± 7.83 and 61 ± 6.24, respectively (Fig 5O). However,
knocking down DRP1 significantly reduced the caspase positive cell count to 20 ± 8.93 (Fig. 5N-
P), demonstrating that this DRP1-dependent pathway is essential for neuronal cell death after
axon injury *in vivo*.

BAX mediates cell death after axotomy

To better understand this type of axon degeneration and cell death, we asked whether they
occurred via canonical pathways of apoptosis, and in particular whether they involved BAX. In
models of apoptosis, DRP1 contributes to BAX recruitment to mitochondria [48], and DRP1 and BAX
interact to drive mitochondrial depolarization and fragmentation [18,49]. Jenner et al. showed a
close interaction between DRP1 and BAX in HEK cells undergoing apoptosis using a fluorescence
complementation biosensor (ddRFP [50]). This system produces enhanced red fluorescence upon
dimerization of A_1 and B_1 proteins. Using A_1-DRP1 and B_1-BAX fusion constructs, we tested
whether DRP1 and BAX closely interact in the axon after injury. One hour after axotomy, we
observed a significant increase in red fluorescence. The interaction between DRP1 and BAX
occurred at mitochondria as shown by co-localization with mito-GFP (Fig. 6A). However, DLK
inhibitor GNE3511 reduced the interaction (Fig. 6A-B). To test whether the interaction between
DRP1 and BAX was dependent on the phosphorylation of DRP1 at S616, we mutated the B_1-DRP
construct to carry a S616A point mutation, preventing its phosphorylation at this site. A_1-BAX and
B_1-DRP1 S616A did not show the increase in red fluorescence, demonstrating that the interaction
between BAX and DRP1 is more likely to occur if DRP1 is phosphorylated at S616 (Fig. 6C), which
is mediated by DLK (Fig. 3).

We next tested whether inhibiting BAX would prevent cell death and axon degeneration in
i3Neurons using BAI, a highly selective small molecule inhibitor of BAX[51]. BAX inhibition
significantly preserved cell bodies of axotomized neurons (Fig. 6D-E). We also found that BAX
inhibition significantly reduced axon degeneration measured 24 hours after injury (Fig. 6F),
indicating that BAX is a key mediator of axon degeneration and cell death after axotomy.

Finally, we asked whether local activation of BAX in the axon was sufficient to cause axons to
degenerate. Other studies have shown stimulation of Bax with a Bcl-x antagonist causes axon
degeneration[19,52] . We grew i3Neurons in microfluidic chambers, which physically isolate axons
from cell bodies and dendrites, and added BAX activator (BAX-A), which binds to the N-terminus
of BAX to promote its oligomerization[53], to the axon compartment. We measured axon integrity
24 hours later. We also tested this in the presence of DLK inhibitor GNE3511 (Fig. 6H). Local
activation of BAX caused axons to degenerate, independently of DLK inhibition (Fig. 6H-I). These
data show that axotomy-induced degeneration relies on BAX as well as DRP1 and DLK, and reveal
that these proteins interact in a signaling cascade. Upon axotomy, DLK directly phosphorylates
DRP1 at S616, which facilitates the interaction between DRP1 and BAX in axonal mitochondria
and leads to BAX-dependent apoptosis.

Discussion

307   In this work, we establish a novel and reproducible model of progressive axon degeneration and
308   neuronal death in human neurons and unravel a cascade of events governing neurodegeneration
309   after axon injury. We show that DLK directly phosphorylates DRP1 to drive mitochondrial fission
310   and BAX-dependent apoptosis that originates in the axon and leads to degeneration of the axon

followed by the cell body. We demonstrate the relevance of this pathway *in vivo*, extending from cultured human neurons to retinal ganglion cells of mice with optic nerve crush injuries. We hypothesize this pathway is broadly implicated in neuronal degeneration caused by axon damage, such as spinal cord injury and traumatic brain injury.

While BAX and DLK have separately been implicated in neuronal apoptosis of retinal ganglion cells after optic nerve crush[26,45,54,55], an interaction between the two has not previously been reported, nor with DRP1. One study reported that DRP1-dependent mitochondrial fission occurs in axons after damage[56]. Here, we show how DLK, DRP1, and BAX cooperate to cause neuronal apoptosis after axon damage, and in particular, how the same apoptotic mechanism can follow axon loss[57] .

Until now, it was also unclear whether pathways distinct from SARM1 can mediate axon degeneration. Many clinical descriptions of axon degeneration refer to "dying back" as a pattern in which distal axons are lost before cell bodies[58,59]. It has been suggested that dying back axon degeneration overlaps with Wallerian degeneration[60], but this remains unclear. Although axon degeneration is rescued by SARM1 deletion in some models (e.g. chemotherapy-induced peripheral neuropathy[61]; TDP43-linked ALS[62]), it is not in others (SOD1 mouse model of ALS[5]; and optic nerve crush[6]). We have shown that a type of non-Wallerian axon degeneration that is more akin to apoptosis can occur after axon damage, and that it spreads retrogradely, similar to a dying back. It is possible that additional axon degeneration pathways also exist.

Therapeutically targeting this pathway offers multiple options. There is precedence for testing some of these strategies, such as blocking BAX[54,55], or targeting mitochondrial dynamics[63], including promoting mitochondrial fusion[30] and blocking mitochondrial fission[64–67]. Blocking DLK signaling using small molecules has been considered[68–70], although a Phase I clinical trial in ALS patients was recently halted due to safety concerns[71]. However, specifically targeting the interaction between DLK and DRP1 has never been attempted.

In addition to protecting axons as we have shown, targeting DRP1 has the potential to protect dendrites[72] . However, it also has possible issues of toxicity as mitochondrial fission is critical for regeneration and motor neuron survival after sciatic nerve injury[73]. A more complete understanding of DRP1 regulation in homeostatic and injury conditions will be helpful. For example, future studies may identify other targetable DRP1 phosphorylation sites in addition to serine 616 as well as additional kinases. To date, kinases of DRP1 that have been identified include CDK5[74,75], TBK1[76], ERK1/2[77,78] CAMKIα[79] and CAMK2[80]; here, we highlight DLK as a novel DRP1 kinase.

As a cytoplasmic facing kinase, DLK has the potential to regulate many substrates in addition to MKK4 and MKK7[81] which had been described for canonical DLK signaling, and DRP1 which we now identify. Based in its localization, DLK likely has many substrates and regulators, and these will be of great interest to uncover as well as the functions of these interactions. DLK is reported to associate with the plasma membrane, Rab3, Rab6, Rab7, Rab11, Golgi-derived, and LAMP1-positive vesicles[39,82]. Localization of DLK to mitochondria has also just recently been described in

cultured mouse embryonic DRG neurons[82], corroborating our observation of DLK-mitochondrial
association.

Our study highlights a novel signaling pathway downstream of DLK. It has generally been thought
that DLK-dependent transcriptional events govern neuronal death [26]. Future studies will
determine whether dying-back and/or apoptosis are independent of the transcriptional response
or whether one or both of these processes requires both the DLK/DRP1/fission cascade and the
DLK-mediated transcriptional response. It will be interesting to examine how the local
mitochondrial mechanism we have uncovered for DLK is coordinated with transcriptional
regulation of cell death factors by DLK, for example BID and PUMA[19].

DRP1-dependent apoptosis has been described in many cell types[15]. The real novelty here lies in
the activation of DRP1 by DLK, and how axonal DLK can trigger this apoptotic pathway in neurons.
Our results call to mind new questions, such as how is DLK localized to the site of mitochondrial
fission so it can regulate DRP1 phosphorylation? Does it associate with factors like MFF, which is
an essential factor in recruiting DRP1 to mitochondria[83]? And what is the mechanism of activation
of DLK that would link it to mitochondrial fission? In the future, it will be of great interest to
explore under which conditions this pathway is activated, as well as the specificity of this pathway
to different neuronal subtypes.

Methods

I3Neuron differentiation

I3Neurons were differentiated as previously described. Briefly, i^3 iPSCs were dissociated using Accutase (Life Technologies, cat. no. A1110501). Cells were plated in Matrigel-coated (1:100 Corning) plates in Neuronal induction media on day 0 (Knockout Dulbecco's modified Eagle's medium (DMEM)/F12 medium; Life Technologies Corporation, cat. no. 12660012), 1X N2 supplement (Life Technologies, cat. no. 17502048), 1× GlutaMAX (Thermofisher Scientific, cat. no. 35050061), 1× MEM nonessential amino acids (NEAA) (Thermofisher Scientific, cat. no. 11140050), 10 μM ROCK inhibitor (Y-27632; Selleckchem, cat. no. S1049), and 2 μg/ml doxycycline (Clontech, cat. no. 631311). Neuronal induction media was changed once a day for 2 more days. On day 3 of induction, cells were dissociated using Accutase and plated in dishes coated with poly-L-ornithine (PLO; 0.1 mg/ml; Sigma, cat. no. P3655-10MG). Cells were plated in neuronal maturation media (BrainPhys medium (STEMCELL Technologies, cat. no. 05790), 1× B27 Plus Supplement (ThermoFisher Scientific, cat. no. A3582801), 10 ng/ml BDNF (PeproTech, cat. no. 450-02), 10 ng/ml NT-3 (PeproTech, cat. no. 450-03), 1 mg/ml mouse laminin (Invitrogen, cat. no. 23017015), and 2 μg/ml doxycycline). Half of the neuronal maturation media was removed and replenished with fresh media every 2-3 days.

DRP1 knockdown using dead Cas9

For DRP1 knockdown, we used CRISPRi-i^3 iPSCs containing a CAG promoter-driven dCas9-BFP-KRAB cassette inserted into the CLYBL safe harbor locus (Addgene #127968). iPSCs were transduced with control gRNA or a previously validated gRNA targeting DRP1[84] and differentiated as described above.

Laser Axotomies

Laser Axotomies were performed on 10-14 day old neurons using a Zeiss LSM 880 Upright 2-Photon microscope equipped with an ablate laser. Temperature and CO_2 levels were maintained using a Pecon microscope incubator. Ablation laser was set to 800 nm, 100% power and 100 iterations. Neurons were cultured in μ-Dish 35 mm dishes (Ibidi, Cat. No. 81156 or 81166 depending on the application) in Hybernate E (Gibco, Cat. No. A1247601), 1× B27 Plus Supplement (ThermoFisher Scientific, cat. no. A3582801), 10 ng/ml BDNF (PeproTech, cat. no. 450-02), 10 ng/ml NT-3 (PeproTech, cat. no. 450-03), 1 mg/ml mouse laminin (Invitrogen, cat. no. 23017015), and 2 μg/ml doxycycline.

Mitochondrial fission after axotomy

iPSCs were differentiated as described above. On day 1 of differentiation, cells were transduced with the appropriate lentivirus. Neuronal induction media was changed 24 hours later. On day 3 of differentiation, 50,000 cells were plated in a 5 μL drop of neuronal maturation media in the middle of a 35mm dish (ibidi, Cat. No. 81156) for the axons to grow out from the center. Dishes were incubated for 20 minutes at 37°C to allow cells to attach, and 2mL of neuronal maturation media was then added to each well. Cells were fed with half-media changes every two days until the day of the experiment. Laser axotomies were performed as described above. Single axons were randomly selected. Five images were taken before axotomy every 5 seconds and axons

were imaged for 5 minutes after axotomy every 5 seconds. The number of mitochondrial particles
was quantified using imageJ imaging software and normalized to the number of mitochondria
pre-axotomy. When treating neurons with different chemical compounds, neurons were
pretreated for one hour prior to performing axotomies. The following compounds were used
GNE3511 (500 nM, Sigma, Cat. No. 5331680001), P110 (1 µM, Tocris, Cat. No. 6897), BAPTA-AM
(10mM Tocris, Cat. No. 2787), BSTA1 (BAX activator, 10 µM, Sigma, Cat. No. SML2243) and BAI1
(BAX inhibitor, 10 µM, Medchemexpress, Cat. No. HY-103269).
426
Axon separation and axotomies
For axon separation experiments after axotomy, 1.5 million cells were plated in 150 µl of
maturation media in a strip in the center of a 6 cm dish. Cells were allowed to attach for 15
minutes at 37°C, after which 8 ml of media was added to the dish. Neurons were cultured for 1
month to allow for the axons to project outwards. Axotomy was performed using a sharp blade.
Axons and cell bodies were separated and processed for western blotting.
433
Immunostaining
Cells were fixed in cold 4% PFA for 5-10 minutes, permeabilized using 0.1% TritonX in PBS for 5
mins and blocked in 5% normal donkey serum (NDS) in PBS at room temperature for 1 hour. Cells
were immunostained with the selected primary antibody in 2.5% NDS overnight at 4°C. Cells were
washed three times with PBS and stained with the appropriate secondary antibody at a
concentration of 1:500 in 2.5% NDS for 1 hour at room temperature. Cells were washed three
times with PBS and DAPI was used as a nuclear counterstain. Antibodies used: βIII tubulin
(Thermo Fischer, cat. no. 2G10, 1:500), anti-pDRP1 S637 (CST, cat. no. 4867S, 1:1000), DLK (CST,
cat. no. GF-32-1, 1:500), TOMM20 (Santa Cruz, cat. no. SC-17764, 1:500).
443
Calcium imaging
Neurons were plated as described above in the center of a 35mm dish, and their axons were
allowed to grow outwards. 10-14 days post induction, neurons were treated with Fluo-4 AM cell
permeable calcium dye (ThermoFischer, Cat. No. F14201) following the manufacturers
indications. Axotomies were performed on single axons as described above and imaged every 2
seconds for 2 minutes after axotomy. Fluorescence intensities were calculated using ImageJ
software and normalized to pre-axotomy levels.
451
HEK cell culture and transfection
HEK-293T cells were cultured in DMEM (Gibco, cat. no 11995065) supplemented with 10% FBS
(Gibco; cat. no. 10437028). For transfection, cells were dissociated using Trypsin-EDTA (0.25%)
(ThermoFischer, Cat. No. 25200056) and plated in DMEM, 10%FBS. The day after plating cells
were transfected with 1 µg DNA plasmid, 8 µl lipofectamine 2000 (Invitrogen, Cat. No. 11668030)
for 24 hours. For western blotting, cells were harvested in ice cold RIPA buffer (Invitrogen, cat.
no. 89901) supplemented with protease inhibitors (Sigma, 11836153001) and PhosStop (Sigma
cat. no. 4906845001). For immunostaining, cells were fixed in 4% PFA for 10 minutes.
460
Western blotting

Protein concentrations were calculated using a BCA assay (ThermoFisher, cat. no. 23225). 10-20
µg of total protein lysate was loaded into a 4–20% Mini-PROTEAN® TGX™ Precast Protein Gels
(Biorad, cat. no. 4561095). Gels were transferred into PVDF membranes and blocked for 1 hour
at room temperature in 5% BSA. Membranes were incubated with the appropriate primary
antibody in 2.5% BSA solution in a cold room, rocking, overnight. Membranes were washed 3
times with PBS 0.1% Tween solution. Secondary antibody was incubated for 1 hour at room
temperature at a concentration of 1:8000 in 2.5% BSA solution. Membranes were washed 3 times
with PBS 0.1% Tween solution prior to developing. Western blots were developed using Clarity
Western ECL Substrate (Biorad, cat. no. 1705061) and imaged using a ChemiDoc MP Imaging
system (Biorad). Band intensity was quantified using FIJI imaging analysis software. The following
antibodies were used at the concentrations stated: anti-DRP1 (Santa Cruz Biotechnology, cat. no.
sc-101270, 1:1000), anti-pDRP1 S616 (CST, cat. no. 3455S, 1:1000), anti-ß Actin (Sigma, cat. no.
A1978, 1:5000), anti-p-cJun S63 (CST, cat. no. 9261, 1:1000), anti-pDRP1 S637 (CST, cat. no.
4867S, 1:1000).
476

Axon Degeneration index and neuron survival
Induction of iPSCs into neurons was performed as described above. On day 1 of differentiation,
neurons were transduced with a cytoplasmic mApple (cyto mApple)-expressing lentivirus, and
the media was changed 24 hours later. On day 3 of differentiation, transduced neurons were
mixed with mixed with untransduced neurons at a ratio of 1:8000 to achieve sparse labelling of
neurons, and re-plated on a 35mm gridded dish (Ibidi Cat. No. 81166). On day 14 of
differentiation, the positions of selected axons or neuron somas was recorded and laser
axotomies were performed as described above. Dished were stored in a temperature and CO_2
controlled incubator until imaged next. Injured axons and somas were re-located at different
time points after axotomy using the gridded dish and imaged. Axons were injured <1mm away
from the soma. The axon degeneration index was calculated as previously described[33].
488

Airyscan imaging
Neurons were plated in 1mm German Glass Coverslips thickness #1.5 (Electron Microscopy
Sciences, Cat. No. 72290-04). 10-day old neurons were fixed with 4% PFA for 10 minutes at
room temperature. Cells were permeabilized with 0.1% Triton-X for 5 minutes and blocked in
5% normal donkey serum (NDS) at room temperature for an hour. Primary antibody incubation
was carried out in 2.5% NDS overnight at 4°C. Cells were washed three times with PBS and
stained with the appropriate secondary antibody at a concentration of 1:500 in 2.5% NDS for 1
hour at room temperature. Cells were washed three times with PBS and DAPI was used as a
nuclear counterstain. Coverslips were mounted using ProLong Glass antifade (Invitrogen, Cat.
No. P36980).
499

Seahorse analysis
For Seahorse analysis, 50,000 cells per well were plated after 3 days of induction in a specialized
Seahorse analyzer 96-well plate (Agilent, cat. no. 103774-100). Cells were washed in PBS once
and incubated in a hypoxic chamber following the manufacturer's procedure. Seahorse analysis
was performed using a Seahorse XFe 96 Analyzer (Agilent). The average value for three wells was
used for analysis.

Protein purification and *in vitro* kinase assay

DLK and DRP1 were cloned into a pcDNA3-C-terminal GST vector. 12 million HEK293 cells were plated in 10 cm dished coated with Matrigel. The day after, cells were transfected with 10 µg plasmids and 40 µl lipofectamine. 24 hours post transfection, cells were washed twice with PBS and harvested in 880 µl ice cold RIPA buffer (Invitrogen, cat. no. 89901) supplemented with protease inhibitors (Sigma, 11836153001) and PhosStop (Sigma cat. no. 4906845001). Lysates were sonicated and spun at 18000g 10 minutes at 4°C and the pellet was discarded. 500 µl Glutathione-Sepharose beads (Sigma Cat. No. GE17-0756-01) were washed twice in 1ml in lysis buffer. Washed beads to 1 ml lysate and incubate end-over-end shaking in cold room overnight. The next morning. Beads were washed 4 times with lysis buffer + 0.4M NaCl. The protein-containing beads were pelleted by spinning 1min 8000rpm at 4°C, the supernatant was discarded, and the beads were resuspended in fresh 200 mM glutathione 1X PBS solution for elution of the desired protein. The beds were incubated for 20 minutes at room temperature shaking. The eluted protein and the beads were separated by briefly spinning in a SpinX column tubes (Sigma Cat. No. CLS8162-24EA). The purity of purified protein was by running 5 µg of protein on an SDS PAGE gel and performing Comassie stain. The purified protein was stored at -80°C. For *in vitro* kinase assay, purified GST-DLK or GST-DLK-S302A was mixed with purified GST-DRP1 at a 1:1 ratio in 50 µl kinase buffer (50mM TrisHCL pH 7.5, 150mM NaCl, 10mM MgCl, 10mM MnCl$_2$, 1.8mM ATP) incubated for 60 mins at 37 °C. The reaction was terminated by adding SDS gel loading buffer and the sample was subjected to western blotting.

Phospho-mass spectrometry

DRP1 bands, with ~0.5 µg protein on each band, were cut from SDS-PAGE. In-gel samples were reduced with 10 mM Tris(2-carboxyethyl)phosphine hydrochloride, alkylated with N-Ethylmaleimide, and digested with trypsin at 37°C for 18 hours. Peptides were extracted, desalted and injected into a nano-LC-MS/MS system where an Ultimate 3000 HPLC was coupled to an Orbitrap Lumos mass spectrometer (Thermo Scientific) via an Easy-Spray ion source (Thermo Scientific). Peptides were separated on an ES902 Easy-Spray column (75-µm inner diameter, 25 cm length, 3 µm C18 beads; Thermo Scientific) with mobile phase B (0.1% formic acid in LC-MS grade acetonitrile) increased from 3% to 24% in 60 min. The flow rate was maintained at 300 nl/min. MS and MS/MS data were acquired on Thermo Scientific Orbitrap Lumos mass spectrometer. MS1 scans were acquired in orbitrap at of a resolution 120k with a mass range of m/z 400-1500. MS2 scans were acquired in ion trap with ETciD method at normal scan rate. The isolation width was 1.6 m/z. Ions were excluded after 1 acquisition and the exclusion duration is 9 sec. MS1 scans were performed every 3 sec. As many MS2 scans were acquired within each MS1 scan cycle.

Proteome Discoverer software version 2.4 was used for protein identification and quantitation. Database search was performed against Sprot Human database using Mascot search engine. Mass tolerances for MS1 and MS2 scans were set to 5 ppm and 0.4 Da, respectively. Up to 2 missed cleavage was allowed for trypsin digestion. NEM on cysteines was set as fixed modification. Oxidation (M) and Phosphorylation (STY) were searched as variable modifications. Spectra of phosphopeptides matched by database search were manually checked. The search results were filtered by a false discovery rate of 1% at the protein level. Proteins detected with 1-2 peptides were further filtered out. Protein abundance values were calculated by summing

550   the abundance of unique and razor peptides matched to that protein. Each sample group
551   contains 4 replicates. The ratios were calculated using protein abundance without normalization.
552
Electron microscopy
Neurons were plated in 1mm German Glass Coverslips thickness #1.5 (Electron Microscopy
Sciences, Cat. No. 72290-04). 10 Day old neurons were fixed with in 4% PFA, 1% Glutaraldehyde
for 40 minutes. Fixed cells were placed in phosphate buffer containing 0.1% sodium borohydride
to inactivate residual aldehyde groups. Cells were then washed with phosphate buffer several
times until the solution was clear of bubbles. To improve reagent penetration, the cells were then
treated with phosphate buffer containing 0.05% Triton X-100. To prevent nonspecific binding of
the immunoreagents, cells were incubated in Aurion blocking solution for Goat gold conjugates
(Electron Microscopy Sciences, Hatfield, PA., containing phosphate buffer, pH 7.4, normal goat
serum (NGS), bovine serum albumin (BSA), and cold water fish skin gelatin (CWFG)). After
blocking, cells were incubated in the primary antibody, Rabbit anti-DLK (1:100), (Sigma-Aldrich,
St. Louis, MO) diluted with incubation buffer (phosphate buffer containing 0.2% Aurion
acetylated bovine serum albumin (BSA-c), Electron Microscopy Sciences, Hatfield, PA., 5% CWFG
and 5% NGS, pH 7.4) for two hours at room temperature. After washes with incubation buffer,
cells were incubated in the secondary antibody Aurion ultra-small gold-conjugated
F(ab')$_2$ fragments of Goat anti-Rabbit IgG, (Electron Microscopy Sciences, Hatfield, PA.) diluted
1:100 with incubation buffer overnight at 4°C. To remove unbound secondary antibody, cells
were washed thoroughly with incubation buffer and then with PBS. After washes, cells were
prepared for silver enhancement.
572
Silver Enhancement and Cell processing for Electron Microscopy
Cells were washed with MilliQ water and then transferred to Aurion R-Gent SE-EM silver
enhancement solution, Electron Microscopy Sciences, Hatfield, PA. and incubated at room
temperature for 35 minutes. The enhancement was stopped by washing cells several times in
MilliQ water. The following processing steps were carried out using the variable wattage Pelco
BioWave Pro microwave oven (Ted Pella, Inc., Redding, CA.): cells were rinsed in MilliQ water,
post-fixed in 1% osmium tetroxide made in MilliQ water, ethanol dehydration series up to 100%
ethanol, followed by a Embed-812 resin (Electron Microscopy Sciences, Hatfield, PA.) infiltration
series up to 100% resin. The epoxy resin was polymerized for 20 hours in an oven set at 60°C.
Ultra-thin sections (90nm) were prepared on a Leica EM UC7 ultramicrotome. Ultra-thin sections
were picked up and placed on 200-mesh cooper grids (Electron Microscopy Sciences, Hatfield,
PA) and post-stained with uranyl acetate and lead citrate. Imaging was performed on a JEOL-1400
Transmission Electron Microscope operating at 80kV and images were acquired on an AMT
BioSprint 29 camera.
587
Generation of DLK KO iPSCs
WT iPSCs were transfected with Cas9-GFP, and two gRNA containing plasmids targeting exons 3
and 5. gRNA expressing plasmids were purchased from Sigma Aldrich (U6-gRNA:hPGK-puro-2A-
tBFP, gRNA1: CCCAGGCTCCCTGCTACTGCAT, gRNA 2: TCCTTTGGCGTGGTGCTATGGG). 1 day after
transfection, expression of Cas9 and gRNAs was validated by the expression of GFP and BFP
respectively and iPSCs were treated for 3 days with 10 µM Puromycin (Invitrogen Cat. No.

A11113803) and allowed to recover for 3 days. Cells were then dissociated with accutase for 10
minutes and dissociated my pipetting forcefully 10 times. A 1:2 serial dilution was then
performed in two 6 well plates. iPSCs were fed every 3 days until individual colonies were visible
(usually 10-14 days). Individually colonies were picked using a P1000 pipette and transferred into
individual wells in a 24-well plate. Individual clones were allowed to expand, passaged and their
DNA was extracted for DLK KO validation using PCR (Forward primer:
TCAGGTGAATGCTGAGCCAGCT, reverse primer: TGGAGACTGTTGCTTCCCACAC). A shorter band
from ~1200bp to ~200bp was expected if the KO was successful. Potential clones were
sequenced. Further validation was performed by differentiating potential clones into neurons
and knock out of DLK was validated by western blotting.

Animals
All animal care and experimental procedures were performed in accordance with animal study
proposals approved by the National Institute of Child Health and Human Disease Animal Care and
Use Committee (animal protocol number 20-003, Le Pichon lab) and by the Baylor College of
Medicine ACUC (animal protocol #AN-7208; Watkins lab). C57Bl/6J (Jax#000664) mice of both
sexes, aged between 6-10 weeks old, were used in this study.

Mouse tissue preparation
The proximal segment of the injured nerve was harvested, and 4 nerves were pooled to obtain
enough protein. Nerves were mechanically lysed in 40 μl ice cold RIPA buffer (Invitrogen, cat. no.
89901) supplemented with protease inhibitors (Sigma, 11836153001) and PhosStop (Sigma cat.
no. 4906845001) for 2 minutes using a micro pestle. Nerves were then sonicated for 15 minutes
at 4°C. Protein lysates were centrifuged at 18 000g, 4°C for 10 minutes and the pellet was
discarded. The samples were subjected to the western blotting procedure described above.
Abundance of phopho-DRP1/ total DRP1 was obtained running two western blots.
Phosphorylated and unphosphorylated protein was divided by the total protein and then divided
to each other to obtain the relative levels of phosphorylated protein.

Adeno-associated virus (AAV)
Adeno-associated viral vectors carrying human synapsin-1 (*hSyn1*) promoter driving expression
of saCas9-U6-control, saCas9-U6-DRP1-gRNA, or eGFP were generated by Epoch Life Sciences Inc.
The plasmids were then packaged into AAV2 capsids at the Optogenetics and Viral Vector Core
at Duncan Neurological Research Institute, Houston.

Intravitreal injections
Mice to be injected were anesthetized with 3% isoflurane. The eyes were sterilized by 3 repeated
applications of 5% ophthalmic betadine (Henry Schein #6900250), Opti-clear ophthalmic eye
wash (Akorn #NDC 17478-620-04) and a dry wipe. A topical anesthetic 0.5% proparacaine HCl
ophthalmic solution (Henry Schein #1365345) was applied. The eyeball was punctured using a 5-
μl Hamilton syringe loaded with a custom 33-gauge needle (Hamilton #7803-5) and some of the
intraocular pressure was relieved. The same puncture site was used to reinsert the Hamilton
needle and 2 μl of AAV in titers ranging from 10^{12} -10^{13} vg/ml was delivered per eye.

Intra-orbital optic nerve crush (ONC)

Two weeks after intravitreal injections, animals underwent optic nerve crush surgeries. Mice undergoing surgery were dosed with 1mg/kg buprenorphine sustained release formulation 1 hour before surgery. Right before surgery, mice were anesthetized with 3% isoflurane. The non-surgical eye received artificial tears ointment (Covetrus #11695-6832-1. The surgical eye was sterilized as described in the intravitreal injections section above. Topical anesthetic 0.5% proparacaine HCl ophthalmic solution (Henry Schein #1365345) was applied on the surgical eye (left). A pair of Vannas scissors (World precision instruments #501777) were used to make incisions in the conjunctival layers. The optic nerve section in the intra-orbital space was exposed by using two pairs of suture-tying forceps (Fine science tools #1106307) to gently clear the soft tissue behind the eye until the optic nerve was visible. The optic nerve was manually crushed for 5s by using a pair of Dumont forceps (Fine science tools #1125325). The eyeball was gently placed back into the orbit. Animals were post-surgically monitored in their home cages until sternal recumbency was observed.

Immunolabeling of retinae

At the experimental endpoint (3 days after optic nerve crush) mice were euthanized by an overdose of isoflurane and followed by decapitation. The eyes were enucleated and drop-fixed in 4% paraformaldehyde for 1 hour. The retinae were dissected out in 1X phosphate-buffered saline (PBS) and then blocked with blocking buffer (5% goat serum, 0.5% Triton X-100 and 0.025% sodium azide in 1X PBS) for 30min. The retinae were then immersed in primary antibody solution prepared by dilution in blocking buffer solution and incubated for 5 days in 4°C. They were then washed three times in 1X PBS with 0.5% TritonX-100 for 30 min each, and then moved to appropriate secondary antibody solution prepared in blocking buffer and incubated overnight in 4°C. After secondary antibody incubation, they were washed again three times in 1X PBS with 0.5% TritonX-100 for 30 min each. Retinae were then mounted in Drop-n-Stain EverBrite™ Mounting Medium (Biotium #23008) onto slides and imaged using Zeiss Axio Imager Z1 fluorescence microscope.

DLK inhibitor treatment

GNE-3511 solution was synthesized as previously described[68]. Briefly, GNE-3511 was dissolved in a solution of 0.5% w/v USP Grade Methyl Cellulose and 0.2% v/v Tween 80 in water up to 48 hours before dosing. Mice were dosed by oral gavage every 12 hours at 75 mg/kg with 10 ml/kg body weight of a 7.5 mg/ml solution of GNE-3511 or vehicle alone.

Funding

This work was supported by NIH intramural funding NICHD ZIA-HD008966 (CLP), by a Milton-Safenowitz postdoctoral fellowship from the ALS Association (JG-D), by NIH grant R01NS112691 (TAW) and grant #021-107 from Mission Connect, a project of the TIRR Foundation (TAW), and NIH intramural funding NINDS 1ZIANS003155-03 (MEW).

Acknowledgements
We would like to thank Drs. Vincent Schram and Chip Dye and the NICHD Microscopy core, Dr.
Yan Li and the NINDS proteomics core, the NHLBI FACS core facility staff, and Dr. Gareth
Thomas (Temple University) for reagents. We would also like to thank members of Dr. Michael
Ward's lab for advice, and Dr. Nicholas Ryba, Dr. Richard Youle, Dr. David Simon, Dr. Alexander
Chesler, and Dr. Joseph Lewcock for helpful discussion and comments on the manuscript.
688
689

Author contributions
Conceptualization: JGD and CLP.
Methodology and Investigation: all authors.
Formal Analysis: MN and JGD.
Visualization: JGD and CLP.
Writing: JGD and CLP.
Supervision: CLP, TW, MEW.
Funding acquisition: CLP, TW, MEW.

References

1. Whitmore, A.V., Lindsten, T., Raff, M.C., and Thompson, C.B. (2003). The proapoptotic
proteins Bax and Bak are not involved in Wallerian degeneration. Cell Death Differ *10*, 260–261.
10.1038/sj.cdd.4401147.

2. Deckwerth, T.L., Elliott, J.L., Knudson, C.M., Johnson, E.M., Snider, W.D., and Korsmeyer, S.J.
(1996). BAX Is Required for Neuronal Death after Trophic Factor Deprivation and during
Development. Neuron *17*, 401–411. 10.1016/s0896-6273(00)80173-7.

3. Osterloh, J.M., Yang, J., Rooney, T.M., Fox, A.N., Adalbert, R., Powell, E.H., Sheehan, A.E.,
Avery, M.A., Hackett, R., Logan, M.A., et al. (2012). dSarm/Sarm1 Is Required for Activation of
an Injury-Induced Axon Death Pathway. Science *337*, 481–484. 10.1126/science.1223899.

4. Gerdts, J., Brace, E.J., Sasaki, Y., DiAntonio, A., and Milbrandt, J. (2015). SARM1 activation
triggers axon degeneration locally via NAD$^+$ destruction. Science *348*, 453–457.
10.1126/science.1258366.

5. Peters, O.M., Lewis, E.A., Osterloh, J.M., Weiss, A., Salameh, J.S., Metterville, J., Brown, R.H.,
and Freeman, M.R. (2018). Loss of Sarm1 does not suppress motor neuron degeneration in the
SOD1G93A mouse model of amyotrophic lateral sclerosis. Hum Mol Genet.
10.1093/hmg/ddy260.

6. Fernandes, K.A., Mitchell, K.L., Patel, A., Marola, O.J., Shrager, P., Zack, D.J., Libby, R.T., and
Welsbie, D.S. (2018). Role of SARM1 and DR6 in retinal ganglion cell axonal and somal
degeneration following axonal injury. Exp Eye Res *171*, 54–61. 10.1016/j.exer.2018.03.007.

7. Mandal, A., and Drerup, C.M. (2019). Axonal Transport and Mitochondrial Function in
Neurons. Front Cell Neurosci *13*, 373. 10.3389/fncel.2019.00373.

8. Smith, G.M., and Gallo, G. (2018). The role of mitochondria in axon development and
regeneration. Dev Neurobiol *78*, 221–237. 10.1002/dneu.22546.

9. Han, S.M., Baig, H.S., and Hammarlund, M. (2016). Mitochondria Localize to Injured Axons to
Support Regeneration. Neuron *92*, 1308–1323. 10.1016/j.neuron.2016.11.025.

10. Lin, M.-Y., Cheng, X.-T., Tammineni, P., Xie, Y., Zhou, B., Cai, Q., and Sheng, Z.-H. (2017).
Releasing Syntaphilin Removes Stressed Mitochondria from Axons Independent of Mitophagy
under Pathophysiological Conditions. Neuron *94*, 595-610.e6. 10.1016/j.neuron.2017.04.004.

11. Karbowski, M., and Youle, R.J. (2003). Dynamics of mitochondrial morphology in healthy
cells and during apoptosis. Cell Death Differ *10*, 870–880. 10.1038/sj.cdd.4401260.

12. Palau, F., Estela, A., Pla-Martín, D., and Sánchez-Piris, M. (2009). Inherited Neuromuscular
Diseases, Translation from Pathomechanisms to Therapies. Adv Exp Med Biol *652*, 129–137.
10.1007/978-90-481-2813-6_9.

13. Tilokani, L., Nagashima, S., Paupe, V., and Prudent, J. (2018). Mitochondrial dynamics:
overview of molecular mechanisms. Essays Biochem *62*, 341–360. 10.1042/ebc20170104.

14. Youle, R.J., and Bliek, A.M. van der (2012). Mitochondrial Fission, Fusion, and Stress. Science
337, 1062–1065. 10.1126/science.1219855.

15. Lee, Y., Jeong, S.-Y., Karbowski, M., Smith, C.L., and Youle, R.J. (2004). Roles of the
Mammalian Mitochondrial Fission and Fusion Mediators Fis1, Drp1, and Opa1 in Apoptosis. Mol
Biol Cell *15*, 5001–5011. 10.1091/mbc.e04-04-0294.

16. Frank, S., Gaume, B., Bergmann-Leitner, E.S., Leitner, W.W., Robert, E.G., Catez, F., Smith,
C.L., and Youle, R.J. (2001). The Role of Dynamin-Related Protein 1, a Mediator of Mitochondrial
Fission, in Apoptosis. Dev Cell *1*, 515–525. 10.1016/s1534-5807(01)00055-7.

17. Milani, M., Beckett, A.J., Al-Zebeeby, A., Luo, X., Prior, I.A., Cohen, G.M., and Varadarajan, S.
(2019). DRP-1 functions independently of mitochondrial structural perturbations to facilitate
BH3 mimetic-mediated apoptosis. Cell Death Discov *5*, 117. 10.1038/s41420-019-0199-x.

18. Jenner, A., Peña-Blanco, A., Salvador-Gallego, R., Ugarte-Uribe, B., Zollo, C., Ganief, T.,
Bierlmeier, J., Mund, M., Lee, J.E., Ries, J., et al. (2022). DRP1 interacts directly with BAX to
induce its activation and apoptosis. Embo J *41*, e108587. 10.15252/embj.2021108587.

19. Simon, D.J., Pitts, J., Hertz, N.T., Yang, J., Yamagishi, Y., Olsen, O., Tešić Mark, M., Molina, H.,
and Tessier-Lavigne, M. (2016). Axon Degeneration Gated by Retrograde Activation of Somatic
Pro-apoptotic Signaling. Cell *164*, 1031–1045. 10.1016/j.cell.2016.01.032.

20. Simon, D.J., Weimer, R.M., McLaughlin, T., Kallop, D., Stanger, K., Yang, J., O'Leary, D.D.M.,
Hannoush, R.N., and Tessier-Lavigne, M. (2012). A Caspase Cascade Regulating Developmental
Axon Degeneration. J Neurosci *32*, 17540–17553. 10.1523/jneurosci.3012-12.2012.

21. Tian, R., Gachechiladze, M.A., Ludwig, C.H., Laurie, M.T., Hong, J.Y., Nathaniel, D., Prabhu,
A.V., Fernandopulle, M.S., Patel, R., Abshari, M., et al. (2019). CRISPR Interference-Based
Platform for Multimodal Genetic Screens in Human iPSC-Derived Neurons. Neuron *104*, 239-
255.e12. 10.1016/j.neuron.2019.07.014.

22. Liao, Y.-C., Fernandopulle, M.S., Wang, G., Choi, H., Hao, L., Drerup, C.M., Patel, R., Qamar,
S., Nixon-Abell, J., Shen, Y., et al. (2019). RNA Granules Hitchhike on Lysosomes for Long-
Distance Transport, Using Annexin A11 as a Molecular Tether. Cell *179*, 147-164.e20.
10.1016/j.cell.2019.08.050.

23. Goldsmith, J., Ordureau, A., Harper, J.W., and Holzbaur, E.L.F. (2022). Brain-derived
autophagosome profiling reveals the engulfment of nucleoid-enriched mitochondrial fragments
by basal autophagy in neurons. Neuron *110*, 967-976.e8. 10.1016/j.neuron.2021.12.029.

24. Fernandopulle, M.S., Prestil, R., Grunseich, C., Wang, C., Gan, L., and Ward, M.E. (2018).
Transcription Factor–Mediated Differentiation of Human iPSCs into Neurons. Curr Protoc Cell
Biology *79*, e51. 10.1002/cpcb.51.

25. Miller, B.R., Press, C., Daniels, R.W., Sasaki, Y., Milbrandt, J., and DiAntonio, A. (2009). A dual
leucine kinase–dependent axon self-destruction program promotes Wallerian degeneration.
Nat Neurosci *12*, 387–389. 10.1038/nn.2290.

26. Watkins, T.A., Wang, B., Huntwork-Rodriguez, S., Yang, J., Jiang, Z., Eastham-Anderson, J.,
Modrusan, Z., Kaminker, J.S., Tessier-Lavigne, M., and Lewcock, J.W. (2013). DLK initiates a
transcriptional program that couples apoptotic and regenerative responses to axonal injury.
Proc National Acad Sci *110*, 4039–4044. 10.1073/pnas.1211074110.

27. Huntwork-Rodriguez, S., Wang, B., Watkins, T., Ghosh, A.S., Pozniak, C.D., Bustos, D.,
Newton, K., Kirkpatrick, D.S., and Lewcock, J.W. (2013). JNK-mediated phosphorylation of DLK
suppresses its ubiquitination to promote neuronal apoptosis. J Cell Biology *202*, 747–763.
10.1083/jcb.201303066.

28. Pichon, C.E.L., Meilandt, W.J., Dominguez, S., Solanoy, H., Lin, H., Ngu, H., Gogineni, A.,
Ghosh, A.S., Jiang, Z., Lee, S.-H., et al. (2017). Loss of dual leucine zipper kinase signaling is
protective in animal models of neurodegenerative disease. Sci Transl Med *9*, eaag0394.
10.1126/scitranslmed.aag0394.

29. Adib, E.A., Smithson, L.J., and Collins, C.A. (2018). An axonal stress response pathway:
degenerative and regenerative signaling by DLK. Curr Opin Neurobiol *53*, 110–119.
10.1016/j.conb.2018.07.002.

30. Franco, A., Dang, X., Zhang, L., Molinoff, P.B., and Dorn, G.W. (2022). Mitochondrial
dysfunction and pharmacodynamics of mitofusin activation in murine Charcot-Marie-Tooth
disease type 2A. J Pharmacol Exp Ther *383*, JPET-AR-2022-001332. 10.1124/jpet.122.001332.

31. Taylor, A.M., Blurton-Jones, M., Rhee, S.W., Cribbs, D.H., Cotman, C.W., and Jeon, N.L.
(2005). A microfluidic culture platform for CNS axonal injury, regeneration and transport. Nat
Methods *2*, 599–605. 10.1038/nmeth777.

32. Tong, Z., Segura-Feliu, M., Seira, O., Homs-Corbera, A., Río, J.A.D., and Samitier, J. (2015). A
microfluidic neuronal platform for neuron axotomy and controlled regenerative studies. Rsc
Adv *5*, 73457–73466. 10.1039/c5ra11522a.

33. Gómez-Deza, J., Slavutsky, A.L., Nebiyou, M., and Pichon, C.E.L. (2022). Local production of
reactive oxygen species drives vincristine-induced axon degeneration. Biorxiv,
2022.07.30.501173. 10.1101/2022.07.30.501173.

34. Court, F.A., and Coleman, M.P. (2012). Mitochondria as a central sensor for axonal
degenerative stimuli. Trends Neurosci *35*, 364–372. 10.1016/j.tins.2012.04.001.

35. Wang, B., Huang, M., Shang, D., Yan, X., Zhao, B., and Zhang, X. (2021). Mitochondrial
Behavior in Axon Degeneration and Regeneration. Front Aging Neurosci *13*, 650038.
10.3389/fnagi.2021.650038.

36. George, E., Glass, J., and Griffin, J. (1995). Axotomy-induced axonal degeneration is
mediated by calcium influx through ion-specific channels. J Neurosci *15*, 6445–6452.
10.1523/jneurosci.15-10-06445.1995.

37. Villegas, R., Martinez, N.W., Lillo, J., Pihan, P., Hernandez, D., Twiss, J.L., and Court, F.A.
(2014). Calcium Release from Intra-Axonal Endoplasmic Reticulum Leads to Axon Degeneration
through Mitochondrial Dysfunction. J Neurosci *34*, 7179–7189. 10.1523/jneurosci.4784-
13.2014.

38. Farley, M.M., and Watkins, T.A. (2018). Intrinsic Neuronal Stress Response Pathways in
Injury and Disease. Annu Rev Pathology Mech Dis *13*, 93–116. 10.1146/annurev-pathol-012414-
040354.

39. Holland, S.M., Collura, K.M., Ketschek, A., Noma, K., Ferguson, T.A., Jin, Y., Gallo, G., and
Thomas, G.M. (2016). Palmitoylation controls DLK localization, interactions and activity to
ensure effective axonal injury signaling. Proc National Acad Sci *113*, 763–768.
10.1073/pnas.1514123113.

40. Nihalani, D., Merritt, S., and Holzman, L.B. (2000). Identification of Structural and Functional
Domains in Mixed Lineage Kinase Dual Leucine Zipper-bearing Kinase Required for Complex
Formation and Stress-activated Protein Kinase Activation*. J Biol Chem *275*, 7273–7279.
10.1074/jbc.275.10.7273.

41. Longo, D.L., and Archer, S.L. (2013). Mitochondrial Dynamics — Mitochondrial Fission and
Fusion in Human Diseases. New Engl J Medicine *369*, 2236–2251. 10.1056/nejmra1215233.

42. Ghosh, A.S., Wang, B., Pozniak, C.D., Chen, M., Watts, R.J., and Lewcock, J.W. (2011). DLK
induces developmental neuronal degeneration via selective regulation of proapoptotic JNK
activity. J Cell Biology *194*, 751–764. 10.1083/jcb.201103153.

43. Yu, R., Liu, T., Ning, C., Tan, F., Jin, S.-B., Lendahl, U., Zhao, J., and Nistér, M. (2019). The
phosphorylation status of Ser-637 in dynamin-related protein 1 (Drp1) does not determine Drp1
recruitment to mitochondria. J Biol Chem *294*, 17262–17277. 10.1074/jbc.ra119.008202.

44. Chang, C.-R., and Blackstone, C. (2007). Cyclic AMP-dependent Protein Kinase
Phosphorylation of Drp1 Regulates Its GTPase Activity and Mitochondrial Morphology*. J Biol
Chem *282*, 21583–21587. 10.1074/jbc.c700083200.

45. Welsbie, D.S., Mitchell, K.L., Jaskula-Ranga, V., Sluch, V.M., Yang, Z., Kim, J., Buehler, E.,
Patel, A., Martin, S.E., Zhang, P.-W., et al. (2017). Enhanced Functional Genomic Screening
Identifies Novel Mediators of Dual Leucine Zipper Kinase-Dependent Injury Signaling in
Neurons. Neuron *94*, 1142-1154.e6. 10.1016/j.neuron.2017.06.008.

46. Larhammar, M., Huntwork-Rodriguez, S., Jiang, Z., Solanoy, H., Ghosh, A.S., Wang, B.,
Kaminker, J.S., Huang, K., Eastham-Anderson, J., Siu, M., et al. (2017). Dual leucine zipper
kinase-dependent PERK activation contributes to neuronal degeneration following insult. Elife
6, e20725. 10.7554/elife.20725.

47. Hunker, A.C., Soden, M.E., Krayushkina, D., Heymann, G., Awatramani, R., and Zweifel, L.S.
(2020). Conditional Single Vector CRISPR/SaCas9 Viruses for Efficient Mutagenesis in the Adult
Mouse Nervous System. Cell Reports *30*, 4303-4316.e6. 10.1016/j.celrep.2020.02.092.

48. Robitaille, K., Daviau, A., Lachance, G., Couture, J.-P., and Blouin, R. (2008). Calphostin C-
induced apoptosis is mediated by a tissue transglutaminase-dependent mechanism involving
the DLK/JNK signaling pathway. Cell Death Differ *15*, 1522–1531. 10.1038/cdd.2008.77.

49. Karbowski, M., Lee, Y.-J., Gaume, B., Jeong, S.-Y., Frank, S., Nechushtan, A., Santel, A., Fuller,
M., Smith, C.L., and Youle, R.J. (2002). Spatial and temporal association of Bax with
mitochondrial fission sites, Drp1, and Mfn2 during apoptosis. J Cell Biology *159*, 931–938.
10.1083/jcb.200209124.

50. Alford, S.C., Abdelfattah, A.S., Ding, Y., and Campbell, R.E. (2012). A Fluorogenic Red
Fluorescent Protein Heterodimer. Chem Biol *19*, 353–360. 10.1016/j.chembiol.2012.01.006.

51. Garner, T.P., Amgalan, D., Reyna, D.E., Li, S., Kitsis, R.N., and Gavathiotis, E. (2019). Small
Molecule Allosteric Inhibitors of BAX. Nat Chem Biol *15*, 322–330. 10.1038/s41589-018-0223-0.

52. Cosker, K.E., Pazyra-Murphy, M.F., Fenstermacher, S.J., and Segal, R.A. (2013). Target-
Derived Neurotrophins Coordinate Transcription and Transport of Bclw to Prevent Axonal
Degeneration. J Neurosci *33*, 5195–5207. 10.1523/jneurosci.3862-12.2013.

53. Reyna, D.E., Garner, T.P., Lopez, A., Kopp, F., Choudhary, G.S., Sridharan, A., Narayanagari,
S.-R., Mitchell, K., Dong, B., Bartholdy, B.A., et al. (2017). Direct Activation of BAX by BTSA1
Overcomes Apoptosis Resistance in Acute Myeloid Leukemia. Cancer Cell *32*, 490-505.e10.
10.1016/j.ccell.2017.09.001.

54. Donahue, R.J., Maes, M.E., Grosser, J.A., and Nickells, R.W. (2020). BAX-Depleted Retinal
Ganglion Cells Survive and Become Quiescent Following Optic Nerve Damage. Mol Neurobiol
57, 1070–1084. 10.1007/s12035-019-01783-7.

55. Goldenberg-Cohen, N., Dratviman-Storobinsky, O., El, S.D.B., Cheporko, Y., and Hochhauser,
E. (2011). Protective Effect of Bax Ablation Against Cell Loss in the Retinal Ganglion Layer
Induced by Optic Nerve Crush in Transgenic Mice. J Neuro-ophthalmol *31*, 331–338.
10.1097/wno.0b013e318227e4fb.

56. Kedra, J., Lin, S., Pacheco, A., Gallo, G., and Smith, G.M. (2021). Axotomy Induces Drp1-
Dependent Fragmentation of Axonal Mitochondria. Front Mol Neurosci *14*, 668670.
10.3389/fnmol.2021.668670.

57. Barrientos, S.A., Martinez, N.W., Yoo, S., Jara, J.S., Zamorano, S., Hetz, C., Twiss, J.L., Alvarez,
J., and Court, F.A. (2011). Axonal Degeneration Is Mediated by the Mitochondrial Permeability
Transition Pore. J Neurosci *31*, 966–978. 10.1523/jneurosci.4065-10.2011.

58. Cavanagh, J.B. (1964). The significance of the "dying back" process in experimental and
human neurological disease. Int Rev Exp Pathol *3*, 219–267.

59. Yaron, A., and Schuldiner, O. (2016). Common and Divergent Mechanisms in Developmental
Neuronal Remodeling and Dying Back Neurodegeneration. Curr Biol *26*, R628–R639.
10.1016/j.cub.2016.05.025.

60. Coleman, M. (2005). Axon degeneration mechanisms: commonality amid diversity. Nat Rev
Neurosci *6*, 889–898. 10.1038/nrn1788.

61. Geisler, S., Doan, R.A., Cheng, G.C., Cetinkaya-Fisgin, A., Huang, S.X., Höke, A., Milbrandt, J.,
and DiAntonio, A. (2019). Vincristine and bortezomib use distinct upstream mechanisms to
activate a common SARM1-dependent axon degeneration program. Jci Insight *4*, e129920.
10.1172/jci.insight.129920.

62. White, M.A., Lin, Z., Kim, E., Henstridge, C.M., Altamira, E.P., Hunt, C.K., Burchill, E.,
Callaghan, I., Loreto, A., Brown-Wright, H., et al. (2019). Sarm1 deletion suppresses TDP-43-
linked motor neuron degeneration and cortical spine loss. Acta Neuropathologica Commun *7*,
166. 10.1186/s40478-019-0800-9.

63. Yang, J., Wu, Z., Renier, N., Simon, D.J., Uryu, K., Park, D.S., Greer, P.A., Tournier, C., Davis,
R.J., and Tessier-Lavigne, M. (2015). Pathological Axonal Death through a MAPK Cascade that
Triggers a Local Energy Deficit. Cell *160*, 161–176. 10.1016/j.cell.2014.11.053.

64. Guo, X., Disatnik, M.-H., Monbureau, M., Shamloo, M., Mochly-Rosen, D., and Qi, X. (2013).
Inhibition of mitochondrial fragmentation diminishes Huntington's disease–associated
neurodegeneration. J Clin Invest *123*, 5371–5388. 10.1172/jci70911.

65. Joshi, A.U., Saw, N.L., Vogel, H., Cunnigham, A.D., Shamloo, M., and Mochly-Rosen, D.
(2018). Inhibition of Drp1/Fis1 interaction slows progression of amyotrophic lateral sclerosis.
Embo Mol Med *10*. 10.15252/emmm.201708166.

66. Chen, Z., Chai, E., Mou, Y., Roda, R.H., Blackstone, C., and Li, X.-J. (2022). Inhibiting
mitochondrial fission rescues degeneration in hereditary spastic paraplegia neurons. Brain.
10.1093/brain/awab488.

67. Flippo, K.H., Lin, Z., Dickey, A.S., Zhou, X., Dhanesha, N.A., Walters, G.C., Liu, Y., Merrill, R.A.,
Meller, R., Simon, R.P., et al. (2020). Deletion of a Neuronal Drp1 Activator Protects against
Cerebral Ischemia. J Neurosci *40*, 3119–3129. 10.1523/jneurosci.1926-19.2020.

68. Patel, S., Cohen, F., Dean, B.J., Torre, K.D.L., Deshmukh, G., Estrada, A.A., Ghosh, A.S.,
Gibbons, P., Gustafson, A., Huestis, M.P., et al. (2015). Discovery of Dual Leucine Zipper Kinase
(DLK, MAP3K12) Inhibitors with Activity in Neurodegeneration Models. J Med Chem *58*, 401–
418. 10.1021/jm5013984.

69. Pichon, C.E.L., Meilandt, W.J., Dominguez, S., Solanoy, H., Lin, H., Ngu, H., Gogineni, A.,
Ghosh, A.S., Jiang, Z., Lee, S.-H., et al. (2017). Loss of dual leucine zipper kinase signaling is
protective in animal models of neurodegenerative disease. Sci Transl Med *9*, eaag0394.
10.1126/scitranslmed.aag0394.

70. Siu, M., Ghosh, A.S., and Lewcock, J.W. (2018). Dual Leucine Zipper Kinase Inhibitors for the
Treatment of Neurodegeneration. J Med Chem *61*, 8078–8087.
10.1021/acs.jmedchem.8b00370.

71. Katz, J.S., Rothstein, J.D., Cudkowicz, M.E., Genge, A., Oskarsson, B., Hains, A.B., Chen, C.,
Galanter, J., Burgess, B.L., Cho, W., et al. (2022). A Phase 1 study of GDC-0134, a dual leucine
zipper kinase inhibitor, in ALS. Ann Clin Transl Neur *9*, 50–66. 10.1002/acn3.51491.

72. Chen, L., Nye, D.M., Stone, M.C., Weiner, A.T., Gheres, K.W., Xiong, X., Collins, C.A., and
Rolls, M.M. (2016). Mitochondria and Caspases Tune Nmnat-Mediated Stabilization to Promote
Axon Regeneration. Plos Genet *12*, e1006503. 10.1371/journal.pgen.1006503.

73. Kiryu-Seo, S., Tamada, H., Kato, Y., Yasuda, K., Ishihara, N., Nomura, M., Mihara, K., and
Kiyama, H. (2016). Mitochondrial fission is an acute and adaptive response in injured motor
neurons. Sci Rep-uk *6*, 28331. 10.1038/srep28331.

74. Rong, R., Xia, X., Peng, H., Li, H., You, M., Liang, Z., Yao, F., Yao, X., Xiong, K., Huang, J., et al.
(2020). Cdk5-mediated Drp1 phosphorylation drives mitochondrial defects and neuronal
apoptosis in radiation-induced optic neuropathy. Cell Death Dis *11*, 720. 10.1038/s41419-020-
02922-y.

75. Jahani-Asl, A., Huang, E., Irrcher, I., Rashidian, J., Ishihara, N., Lagace, D.C., Slack, R.S., and
Park, D.S. (2015). CDK5 phosphorylates DRP1 and drives mitochondrial defects in NMDA-
induced neuronal death. Hum Mol Genet *24*, 4573–4583. 10.1093/hmg/ddv188.

76. Chen, S., Liu, S., Wang, J., Wu, Q., Wang, A., Guan, H., Zhang, Q., Zhang, D., Wang, X., Song,
H., et al. (2020). TBK1-Mediated DRP1 Targeting Confers Nucleic Acid Sensing to Reprogram
Mitochondrial Dynamics and Physiology. Mol Cell *80*, 810-827.e7.
10.1016/j.molcel.2020.10.018.

77. Gan, X., Huang, S., Wu, L., Wang, Y., Hu, G., Li, G., Zhang, H., Yu, H., Swerdlow, R.H., Chen,
J.X., et al. (2014). Inhibition of ERK-DLP1 signaling and mitochondrial division alleviates
mitochondrial dysfunction in Alzheimer's disease cybrid cell. Biochimica Et Biophysica Acta Bba
- Mol Basis Dis *1842*, 220–231. 10.1016/j.bbadis.2013.11.009.

78. Yu, T., Jhun, B.S., and Yoon, Y. (2011). High-Glucose Stimulation Increases Reactive Oxygen
Species Production Through the Calcium and Mitogen-Activated Protein Kinase-Mediated
Activation of Mitochondrial Fission. Antioxid Redox Sign *14*, 425–437. 10.1089/ars.2010.3284.

79. Han, X.-J., Lu, Y.-F., Li, S.-A., Kaitsuka, T., Sato, Y., Tomizawa, K., Nairn, A.C., Takei, K., Matsui,
H., and Matsushita, M. (2008). CaM kinase Iα–induced phosphorylation of Drp1 regulates
mitochondrial morphology. J Cell Biology *182*, 573–585. 10.1083/jcb.200802164.

80. Yang, D., Rong, R., Yang, R., You, M., Wang, M., Li, H., and Ji, D. (2021). CaMK II -induced
Drp1 phosphorylation contributes to blue light-induced AIF-mediated necroptosis in retinal
R28 cells. Biochem Bioph Res Co *559*, 113–120. 10.1016/j.bbrc.2021.04.082.

81. Tedeschi, A., and Bradke, F. (2013). The DLK signalling pathway—a double-edged sword in
neural development and regeneration. Embo Rep *14*, 605–614. 10.1038/embor.2013.64.

82. Tortosa, E., Ghosh, A.S., Li, Q., Wong, W.R., Hinkle, T., Sandoval, W., Rose, C.M., and
Hoogenraad, C.C. (2022). Stress-induced vesicular assemblies of dual leucine zipper kinase are
signaling hubs involved in kinase activation and neurodegeneration. Embo J *41*, e110155.
10.15252/embj.2021110155.

83. Otera, H., Ishihara, N., and Mihara, K. (2013). New insights into the function and regulation
of mitochondrial fission. Biochimica Et Biophysica Acta Bba - Mol Cell Res *1833*, 1256–1268.
10.1016/j.bbamcr.2013.02.002.

958
959

Figure 1: Axotomy triggers a wave of DRP1-dependent mitochondrial fission, axon degeneration and cell death

Figure 1 – Axotomy triggers a wave of DRP1-dependent mitochondrial fission, axon degeneration and cell death

A) Schematic representation of i3Neuron laser axotomy using Biorender. Dashed box indicates field of view. * Marks the site of axotomy

B) Representative images of WT and SARM1 KO axons transduced with cytoplasmic mApple (red) pre and 4 hours post axotomy (PA). * Marks the site of axotomy. (Scale bar = 100 μm).

C) Quantification of axon degeneration index (ADI) in WT and SARM1 KO neurons distal and proximal to the site of axon injury 4 hours post axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 38 axons. Two-way ANOVA, Bonferroni correction ($p \leq 0.01$ **, not significant, ns).

D) Representative images of i3Neuron axons transduced with mitoGFP (green) pre- and 5 minutes post axotomy (PA). * Marks the site of axotomy. (Scale bar = 20 μm). Time-lapse images of mitochondria shown in E.

E) Representative images of mitochondria undergoing fission after axon injury outlined in D.

F) Schematic representation of i3Neuron laser axotomy. Dashed box indicates field of view.

G) Quantification of mitochondrial length pre (green) and 5 minutes post axotomy (grey) (PA). Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 28 axotomized neurons. Two-way ANOVA, Bonferroni correction ($p \leq 0.05$ *).

H) Normalized number of mitochondrial particles post axotomy in control and DRP1 KD neurons. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 25 axotomized neurons. Two-way ANOVA, Bonferroni correction ($p \leq 0.05$ *).

I) Normalized number of mitochondrial particles 3 mins post axotomy in DMSO and P110 treated neurons. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 26 axotomized neurons. Two-way ANOVA, Bonferroni correction ($p \leq 0.05$ *).

J) Quantification of axotomy-induced mitochondrial fission wave. Normalized number of mitochondrial particles post axotomy 0-100, 200-300 and 400-500 μm from the site of axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 23 axotomized neurons. One-way ANOVA, Bonferroni correction (*$p \leq 0.05$, 0-100 μm vs 200-300 μm, # $p \leq 0.05$, 0-100 μm vs 400-500 μm).

K) Representative images of axotomy-induced mitochondrial fission wave, Pre axotomy (0), 10-, 30- and 120-minutes post axotomy. Neurons sparsely transduced with mitoGFP (green). (Scale bar = 25 μm).

L) Quantification of axotomy-induced progressive axon degeneration. Axon degeneration index (ADI) 0-100, 200-300 and 400-500μm from the site of axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 14 axotomized neurons. (*$p \leq 0.05$, 0-100 μm vs 200-300 μm, ## $p \leq 0.01$, ### $p \leq 0.005$, 0-100 μm vs 400-500 μm).

M) Representative images of axotomy-induced axon degeneration. Pre axotomy (0), 10-, 30- and 120-minutes post axotomy. Neurons sparsely transduced with cyto-mApple (red). * Marks the site of axotomy. (Scale bar = 25 μm).

N) Representative images of axotomy-induced neuron death. Pre and 12-hours post axotomy. Arrow indicates axotomized neuron. Neurons sparsely transduced with cyto- mApple (red) and mitoGFP (green). (Scale bar = 40 μm).

1003 O) Quantification of axotomy-induced neuron death. Percentage of dead neurons 12 hours post
1004   axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥30
1005   axotomized neurons. Unpaired t-test (p≤0.01 **).

Figure 2: DLK regulates axotomy-induced mitochondrial fission

1006 **Figure 2 – DLK regulates axotomy-induced mitochondrial fission**

1007

A) Representative images of DLK colocalization at the site of mitochondrial fission after
axotomy. DLK-GFP (green) and mitochondria (mitoRFP, magenta). (Scale bar = 10 μm).
B) Percentage mitochondrial fission events where DLK is localized at the site of fission. Results
are represented as mean ± SEM. N=3 independent differentiations, N≥30 axotomized
neurons.
C) Representative images of WT and DLK KO neurons transduced with mitoGFP (green) pre (Pre)
and 5 mins post axotomy (PA). * Marks the site of axotomy. (Scale bar = 25 μm).
D) Normalized number of mitochondrial particles post axotomy in WT (black) and DLK KO (green)
neurons. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 30
axotomized neurons. (Two-way ANOVA, Bonferroni correction ($p \leq 0.05$ *)
E) Normalized number of mitochondrial particles post axotomy in DMSO (black) and GNE3511
(green)-treated neurons. Results are represented as mean ± SEM. N=3 independent
differentiations, N≥ 25 axotomized neurons. (Two-way ANOVA, Bonferroni correction ($p \leq 0.05$
*)
F) Normalized number of mitochondrial particles post axotomy in DLK KO neurons transduced
with WT DLK-GFP (green), DLK-C127S-GFP (black) or DLK-S302A-GFP (purple). Results are
represented as mean ± SEM. N=3 independent differentiations, N≥ 23 axotomized neurons.
Two-way ANOVA, Bonferroni correction (*$p \leq 0.05$, WT DLK vs DLK C127S, # $p \leq 0.05$, WT DLK vs
S302A).

Figure 3: Activation of the DLK pathway by overexpression causes increased DRP1 phopshorylation

Figure 3 – Activation of the DLK pathway by overexpression causes increased DRP1 phosphorylation

A) Representative western blots of HEK293A cells transfected for 24 hours with WT, C127S and S302A DLK-GFP. Immunoblot for pS616-DRP1, pS637-DRP1, total DRP1, DLK, pS63-cJun and loading control ß actin.

B) Quantification of pS63-cJun/ ß actin levels after 24-hour expression of WT, C127S and S302A DLK-GFP in HEK293A cells. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (p≤0.05 *).

C) Quantification of pS616-DRP1/total DRP1 levels after 24-hour expression of WT, C127S and S302A DLK-GFP in HEK293A cells. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (p≤0.05 *).

D) Quantification of total DRP1/ ß actin levels after 24-hour expression of WT, C127S and S302A DLK-GFP in HEK293A cells. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (not significant (ns)).

E) Quantification of pS637 DRP1/ ß actin levels after 24-hour expression of WT, C127S and S302A DLK-GFP in HEK293A cells. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (not significant (ns)).

F) Schematic representation of GST-DRP1 and DLK-GFP co-expression, purification and phospho-mass spectrometry.

G) Phospho-mass spectrometry. pS616-DRP1 peptide / total DRP1 peptide levels from purified GST-DRP1 in HEK 293A cells expressing GST-DRP1, GST-DRP1 and DLK-GFP or GST-DRP1 and DLK-S302A-GFP. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (p≤0.01 **).

H) Schematic representation of the *in vitro* kinase assay using purified DLK, DLK S302A and DRP1 to test the direct phosphorylation of DRP1 by DLK.

I) *In vitro* kinase assay. Utilizing separately purified DRP1, WT DLK and kinase dead DLK S302A. Representative immunoblots for pS616-DRP1, total DRP1 and DLK. Results are represented as mean ± SEM. One-way ANOVA, Bonferroni correction (p≤0.01 **).

J) Representative images of HEK-293A cells expressing WT, C127S and S302A DLK-GFP (green) and mitoTracker (orange) for 24 hours. (Scale bar = 10 μm).

K) Quantification of mitochondrial morphology in HEK293A cells transfected for 24 hours with WT, C127S and S302A DLK-GFP. N=3 individual transfections. Results are represented as mean ± SEM. Two-way ANOVA, Bonferroni correction (WT vs C127S p≤0.0001 ****, WT vs S302A p≤0.0001 ****, C127S vs S302A not significant ns).

L) Representative western blots HEK-293A cells expressing dCAS9 transduced with control and DRP1 gRNAs. Immunoblot for total DRP1 and loading control ß actin.

M) Representative images of Control and DRP1 KD HEK-293A cells expressing DLK-HALO for 24 hours. NLS-GFP (green), DLK-HALO (white), mitoTracker (orange). (Scale bar = 10 μm).

N) Quantification of Control and DRP1 KD HEK-293A cells mitochondrial morphology. Cells transfected for 24 hours with DLK-HALO. N=3 individual transfections. Results are represented as mean ± SEM. (Two-way ANOVA, Bonferroni correction (p≤0.01 **)

Figure 4: Axotomy causes DLK-dependent phosphorylation of DRP1

1071 **Figure 4 – Axotomy causes DLK-dependent phosphorylation of DRP1**
1072

A) Schematic representation of i3Neuron center plating and axotomy for protein harvesting
after injury.
B) Representative western blots of WT and DLK KO neuron cell bodies untreated (UT), 2 and 4
hours post axotomy. Immunoblot for p-S616 DRP1, total DRP1 and loading control ß actin.
C) Quantification of pDRP1 (S616)/ total DRP1 levels in WT and DLK KO neuron cell bodies
untreated (UT), 2 and 4 hours post axotomy. Results normalized to UT. Results are
represented as mean ± SEM. N=3 independent differentiations. One-way ANOVA. No
significant changes observed.
D) Representative western blots of WT and DLK KO neuron axons untreated (UT), 2 and 4 hours
post axotomy. Immunoblot for p-S616 DRP1, total DRP1 and loading control ß actin.
E) Quantification of pDRP1 (S616)/ total DRP1 levels in WT and DLK KO neuron axons 0, 2 and 4
hours post axotomy. Results normalized to UT. Results are represented as mean ± SEM. N=3
independent differentiations. Two-way ANOVA. Bonferroni correction (p≤0.05*).
F) Representative images of WT and DLK KO axons stained for ßIII tubulin (red) and pDRP1 S616
(white) untreated (UT), 30 mins and 1 hour post axotomy. (Scale bar = 25 μm).
G) Quantification of pDRP1 S616 fluorescence in WT and DLK KO neuron axons after axotomy.
Results normalized to untreated axons. Results are represented as mean ± SEM. N=3
independent differentiations. Two-way ANOVA, Bonferroni correction (p≤0.05 *)
H) Schematic representation of optic nerve crush injury model. Bracket and * indicate the
proximal portion of the injured nerve harvested for western blotting.
I) Representative western blots of proximal portion of ipsilateral (I) injured and contralateral
(C) optic nerves 24 hours after optic nerve crush. Immunoblot for p-S616 DRP1, total DRP1
and loading control ß actin.
J) Quantification of pDRP1 (S616)/ total DRP1 levels after optic nerve crush. Results normalized
to contralateral side (C). Results are represented as mean ± SEM. Unpaired t-test (p≤0.05 *).
K) Quantification of total DRP1/ ß actin levels after optic nerve crush. Results normalized to
contralateral side (C). Results are represented as mean ± SEM. Unpaired t-test not significant
(ns).
L) Representative western blots of proximal portion of optic nerves 24 hours after optic nerve
crush in mice treated with vehicle (veh) or DLK inhibitor (GNE-3511). Immunoblot for p-S616
DRP1, total DRP1 and loading control ß actin.
M) Quantification of total pDRP1 (S616)/ total DRP1 levels after optic nerve crush in mice treated
with vehicle (veh) or DLK inhibitor (GNE-3511). Results are represented as mean ± SEM.

Figure 5: Inhibiting DRP1 delays axonal degeneration

Figure 5 – Blocking DRP1 or DLK delays axon degeneration

A) Schematic representation of i3Neuron laser axotomy. Dashed box indicates the field of view.

B) Representative images of WT and DLK KO neuron axons transduced with cyto mApple (red) proximal to the site of injury 0,4, 8 and 24 hours post axotomy (Scale bar = 25 μm).

C) Quantification of axon degeneration index (ADI) in WT and DLK KO neurons 0, 4, 8 and 24 hours post axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 33 axons. Two-way ANOVA, Bonferroni correction (p≤0.005 ***, p≤0.001 ****).

D) Schematic representation of i3Neuron laser axotomy. Dashed box indicates field of view.

E) Representative images of Control and DRP1 KD neuron axons transduced with cyto mApple (red) proximal to the site of injury 0, 4, 8 and 24 hours post axotomy (Scale bar = 25 μm).

F) Quantification of axon degeneration index (ADI) in WT and DLK KO neurons 0, 4, 8 and 24 hours post axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 33 axons. Two-way ANOVA, Bonferroni correction (p≤0.005 ***, p≤0.001 ****).

G) Schematic representation of i3Neuron laser axotomy. Dashed box indicates field of view.

H) Representative images of WT and DLK KO neuron cell bodies transduced with cyto mApple (red) pre and 18 hours post axotomy (PA). (Scale bar = 40 μm).

I) Percentage degenerated WT and DLK KO 18 post axotomy. Results are represented as mean ± SEM. N=3 independent differentiations. Unpaired t-test (p≤0.05 *).

J) Schematic representation of i3Neuron laser axotomy. Dashed box indicates field of view.

K) Representative images of Control and DRP1 KD neuron cell bodies transduced with cyto mApple (red) pre and 18 hours post axotomy (PA). (Scale bar = 40 μm).

L) Percentage degenerated Control and DRP1 KD 18 post axotomy. Results are represented as mean ± SEM. N=3 independent differentiations. Unpaired t-test (p≤0.05 *).

M) Schematic representation of RGC survival following optic nerve crush.

N) Western blots of Control and DRP1 gRNA transduced optic nerved. Immunoblot for DRP1 and loading control ß actin.

O) Representative images of cleaved caspase 3 positive cells in saCas9-Control and saCas9-DRP1 gRNA transduced RGCs 3 days after ONC. (Scale bar = 25 μm).

P) Average cleaved caspase 3 positive cells per animal in saCas9-Control, saCas9-DRP1 gRNA and EGFP transduced RGCs after nerve crush. Results are represented as mean ± SEM. N≥4 animals per condition. One-way ANOVA, Bonferroni correction. (Control gRNA vs EGFP, not significant, ns. Control gRNA vs DRP1 gRNA p≤0.05 *. DRP1 gRNA vs EGFP p≤0.01 **).

Fig 6: BAX mediates cell death afer axotomy

Figure 6 – BAX regulates cell death after axotomy

A) Representative images of WT neurons treated with DMSO or GNE3511 pre- and 1-hour post-axotomy (PA) expressing mitoGFP (green) and A_1-BAX/B_1-DRP1 complexes (red). (Scale bar = 5 μm).

B) Quantification of A_1-BAX/B_1-DRP1 fluorescence fold change in the mitochondria of DMSO or GNE3511 treated neurons 1-hour post-axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 34 axotomized neurons. Two-way ANOVA, Bonferroni correction. ($p \leq 0.05$ *).

C) Quantification of A_1-BAX/B_1- WT DRP1 and A_1-BAX/B_1- DRP1 S616A fluorescence fold change in the mitochondria 1-hour post-axotomy. Results are represented as mean ± SEM. N=3 independent differentiations, N≥ 34 axotomized neurons. Two-way ANOVA, Bonferroni correction. ($p \leq 0.05$ *).

D) Representative images of DMSO and BAXi treated neuron cell bodies transduced with cyto mApple (red) pre and 18 hours post axotomy (PA). (Scale bar = 40 μm).

E) Percent degenerated neurons 18 hours post axotomy treated with either DMSO or BAXi. Results are represented as mean ± SEM. N=3 independent differentiations. Unpaired t-test ($p \leq 0.05$ *).

F) Quantification of axon degeneration index (ADI) of DMSO and BAX-A-treated neurons 24 hours after axotomy. Results are represented as mean ± SEM. N=3 independent differentiations. One-way ANOVA, Bonferroni correction ($p \leq 0.001$ ***, not significant, ns).

G) Illustration of microfluidic devices used to separate axons from somas allowing for the local treatment of i3Neuron axons.

H) Representative images of WT axons separated using microfluidic chambers treated with DMSO, BAX-A or BAX-A + GNE3511 for 24 hours. (Scale bar = 25 μm).

I) Quantification of axon degeneration index (ADI) of isolated axons treated with DMSO BAX-A and BAX-A + GNE3511 for 24 hours. Results are represented as mean ± SEM. N=3 independent differentiations. One-way ANOVA, Bonferroni correction ($p \leq 0.001$ ****, not significant, ns).