# Soil legacies of extreme droughts enhance the performance of invading plants

Madhav P. Thakur[1,*], Maartje A. van der Sloot[2], Rutger A. Wilschut[3], S. Emilia Hannula[4], Freddy ten Hooven[5], Stefan Geisen[3], Casper W. Quist[6], Katja Steinauer[7], Wim H. van der Putten[3,5]

[1]*Institute of Ecology and Evolution and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland*

[2]*Plant Ecology and Nature Conservation Group, Wageningen University, Wageningen, The Netherlands*

[3]*Department of Nematology, Wageningen University and Research, 6708 PB, Wageningen, The Netherlands*

[4]*Institute of Environmental Sciences, Leiden University, Leiden*

[5]*Department of Terrestrial Ecology, Netherlands Institute of Ecology (NIOO- KNAW), Wageningen, The Netherlands*

[6]*Biosystematics Group, Wageningen University, 6708 PB Wageningen, the Netherlands*

[7]*Institute of Plant Sciences, University of Bern, Bern, Switzerland*

*Corresponding author: madhav.thakur@unibe.ch

**Abstract**

Extreme droughts can weaken the biotic resistance of native plant communities against the establishment of invading plants. However, we know little about the underlying mechanisms. Using a plant-soil feedback approach, we tested how an extreme drought event alters the soil-mediated biotic resistance of resident native plant communities against invading plant species from native and non-native ranges, namely non-resident natives, native range-expanders, and alien plants. We show that all three types of invading plants performed better in soils with a legacy of extreme drought independent of resident native plant diversity. Path models revealed that extreme drought effects on non-resident natives were mediated by the root biomass of resident native plants and endophytic fungal pathogens during drought, whereas alien plant performance was mediated only *via* the root biomass of resident native plants also during drought. Our results highlight that the performance of resident native plants during extreme drought and subsequent effects on soil fungi determine the performance of invading plants from native and non-native origins after extreme droughts.

**Keywords:** plant-soil feedback; diversity-invasibility relationships; climate extremes; biological invasion; soil microorganisms; fungal endophytes

**Introduction**

The success and failure of invading plants in new environments have intrigued ecologists for

a long time[1–4]. Several factors ranging from biotic to abiotic features of the new environment,

and from ecological to evolutionary characteristics of the invading plant determine its

success[5–9]. One of the classic ideas that have persisted over several decades is how the

diversity of native plant communities constrains or facilitates the establishment of invading

plants - the so-called diversity-invasibility relationship[2,10–12]. The contemporary understanding

of diversity-invasibility relationships is that greater diversity of native plants at local scales

(plot level) constrains the establishment of invading plants (e.g., particularly the alien plants

that come from other continents), whereas greater diversity of native plants at regional

scales (e.g., landscape level observational studies) correlates positively with the diversity of

non-native plants[13,14]. Yet, even at the local scale, studies have shown variable biotic

resistance of native plant diversity against invading plants, underlain by mechanisms that

remain little understood to date[15,16].

The key idea behind negative diversity-invasibility relationships is that greater plant diversity

can utilize resources (e.g., soil nutrients) more efficiently than low-diverse plant

communities, which in turn constrains the establishment of invading plants due to the lesser

availability of resources[2,17,18]. Recent advances in invasion biology further point to the

importance of soil microorganisms like fungi to determine the establishment success (or

failure) of invading plants[19–22]. For instance, pathogenic soil fungi in a new environment of

invading plants often weakly inhibit them compared to microorganisms from their previous

environments[23–25]. Diverse native plant communities could also influence the establishment

of invading plants through their ability to steer soil microorganisms[26]. For instance, a recent

experimental study showed that greater fungal diversity associated with native diverse plant

communities could constrain the performance of alien plants[27], as such diverse communities

are more likely to contain plant pathogens able to infect a new host (i.e., sampling effects).

Conversely, pathogenic soil fungi specialized on certain host plants may also become diluted

in diverse plant communities[28–30], which potentially increases the chance of the successful

establishment of invading plants given the reduced pathogenic load[31]. Taken together, the

performance of invading plants in diverse resident plant communities could very well depend

on the type of biotic interactions (e.g., pathogenic, or mutualistic) that emerge between the

invading plant and the resident soil microorganisms associated with resident plants.

Diverse native plant communities are further shown to be resistant to external disturbances,

such as drought[32–34], which could affect the success of invading plants[35,36]. With the

increasing magnitude and frequency of droughts across the biosphere, it is likely that biotic

resistance (or facilitation) of diverse native plant communities through soil microorganisms

against invading plants would change. This expectation is plausible as droughts shift the

structure and function of soil microbial communities[37–42], which could subsequently affect the

performance of incoming invading plants through drought legacies in the soil[43].

Here, we study how the performance of three types of invading plants- local native plants,

range-expanding (or intracontinental or neonative) plants and alien (or exotic or

intercontinental) plants[44,45] are affected by soil legacies of native plant diversity and extreme

drought (Figure 1). Using three broad categories of plant invaders facilitates our

understanding of how plant communities may change after extreme drought events, and

whether diversity-invasibility relationships vary for various kinds of invader plants. Most

diversity-invasibility plant studies have focused on alien plants[46], whereas local natives

(those not yet present in a native community but in near surroundings: hereafter non-

resident natives for brevity) and range-expanding plants have rarely been used to examine

their potential to invade a new environment that may have the history of native plant diversity

and extreme drought.

We use a plant-soil feedback (PSF) experimental approach to investigate soil legacy effects

on the performance of invading plants. The PSF approach is a robust way to test the effects

of soil microorganisms and soil nutrients in determining the performance of plants through

soil legacy or carry-over effects: PSF consists of a conditioning phase (to create a soil

legacy) and a feedback phase (to test soil legacy effects, Figure 1)[47,48]. In line with the

diversity-invasibility hypothesis, we hypothesize that native plant diversity will constrain the

establishment of all three types of invaders, whereas extreme drought will weaken the native

plant diversity effect. Moreover, we suspect that these native diversity and drought effects

will be mediated *via* soil microorganisms and soil nutrient availability. More specifically, we

investigate the role of various functional groups of soil fungi that could affect the

performance of invading plants. Given that climate change continues to alter local plant

community composition through both local diversity losses and gains, our study attempts to

provide insights into underlying mechanisms behind the success and failure of invading

plants by zooming in on the roles of soil fungi and soil nutrients.

**Results**

The performance of all invading plants was strongly affected by the soil legacy of extreme

droughts, whereas the effects of soil legacies of native plant diversity on all three types of

invader plants were weak (Table 1, Figures 2 and 3). More specifically, we found that all

three types of invading plants had greater shoot biomass in soils with a history of extreme

drought (Figure 2). Range-expanding plants in general had the least shoot biomass

compared to that of non-resident natives and alien plant invaders (Figure 2). The root

biomass of invading plants was significantly greater in soils with a history of high native plant

diversity contradicting the diversity-invasibility hypothesis (Table 1, Figure 3). Species-

specific shoot and root biomasses of all invading plants used during the feedback phase are

provided in supplementary figures 1 and 2.

To explore the underlying mechanisms that could have driven the effects of soil legacies of

extreme drought and native plant diversity on the performance of invading plants during the

feedback phase, we examined how abiotic (soil C, N and P responses) and biotic (native

plant root and soil fungal responses) variables differed in soils from the conditioning phase.

Extreme drought and native plant diversity consistently reduced native shoot and root

biomass, whereas soil PO4 was also lower in soils with higher native plant diversity during

the conditioning phase (Table 2). We quantified the response of 12 functional groups of soil

fungi, of which the majority was influenced by native plant species richness (Table 2).

However, only the relative abundance of total endophytic fungi (mainly the relative

abundance of *Rhizoctonia solani*) decreased in soils that experienced extreme drought

(Table 2). Multivariate analyses further showed significant compositional differences among

soil fungal communities across native plant diversity (F-value$_{1,175}$ = 29.36, p-value=0.01),

whereas neither extreme drought nor its interaction with plant species richness significantly

altered fungal community composition (p-value>0.05, Supplementary table 1).

We used path models to link the biotic and abiotic variables from the conditioning phase

soils to explain the performance of invading plants during the feedback phase. We ran path

models separately for the three types of invading plants, and only present those of native

and alien plants (Figure 4) due to the lack of goodness of fit of the path model of range-

expanding plants (details in method section). In none of the path models, we found soil N

and soil P (soil PO$_4$) from the conditioning phase explaining the performance of invader

plants during the feedback phase, whereas root biomass production during the conditioning

phase and the abundance of specific fungal groups did explain some of the variation in non-resident natives and alien plant performance. More specifically, the positive effects of extreme drought on invading non-resident native plants were mediated *via* lower root biomass production of resident native plants, which subsequently reduced the relative abundance of endophytic fungal pathogens (Figure 4 A). Given the negative relationship between endophytic fungal pathogens and invader performance, our path model results suggest that non-resident native plants are likely to experience weaker pathogenic pressure in soils that experienced extreme drought. Alien invader performance could also be partly explained through root biomass and fungal parasite (or mycoparasites: fungi that parasitize other fungi) responses during the conditioning phase (Figure 4 B). For instance, the positive soil legacy effect of native plant diversity on the performance of alien invaders (i.e., greater root biomass) was partly driven through the root biomass of native plants during the conditioning phase (Figure 4 B), whereas the performance of alien plants aboveground (i.e., shoot biomass) during the feedback phase was mediated through negative effects of native plant diversity on soil fungal parasites (Figure 4 B). Positive effects of extreme drought on alien invaders also occurred through decreased root biomass of resident native plants during the conditioning phase (Figure 4 B).

**Discussion**

There is a growing consensus that the increasing intensity of drought events across the biosphere will alter plant community composition with detrimental consequences for native plant biodiversity and ecosystem functions[49,50]. Extreme droughts can shift plant community composition by benefitting one set of species while harming the others[51,52], and thereby often enhancing plant turnover rates[53,54]. Many studies have suggested that alien plants have an advantage over native plants during and after extreme droughts[55–57], but factors that could lead to such an advantage remain poorly understood. Through a plant-soil feedback

experiment, we highlight that invading plants perform better in soils with an extreme drought

legacy. This seemed true for all three types (non-resident natives, range-expanders, and

alien plants) of invading plants irrespective of the soil legacy of native plant diversity. In fact,

there were no signs of biotic resistance in soils with the legacy of native diverse community,

but instead of a biotic facilitation through increments in root biomass production of invading

plants. We identified some specific biotic pathways through which invading plants exploit

soils of extreme drought or native plant diversity legacies. For instance, when extreme

drought reduced the root biomass of native plant communities, it benefitted the non-resident

native plants through a reduction in endophytic fungal pathogens. It is likely that lower

production of roots during extreme droughts can simultaneously reduce the amount of fungal

pathogens that reside inside the roots[58,59].

Soil fungal communities are known to determine the success (or failure) of invading plants[60–

62]. Endophytic fungi (those penetrating plant roots) in particular have received recent

attention for their positive roles (e.g., enhanced nutrient uptake and pathogen defense) in

driving the host performance including those of alien plants[60,63–65], but also most pathogenic

fungi that can inhibit plant performance are endophytic[66,67]. While we found that extreme

drought and plant diversity both reduced the relative abundance of endophytic fungi during

the conditioning phase (Table 2), it was the reduction of relative abundances of plant

pathogenic endophytic fungi due to drought-induced lower root biomass during the

conditioning phase through which extreme drought indirectly promoted the performance of

non-resident native plants (Figure 4 A). Based on ITS sequencing data, it was pathogens

like *Rhizoctonia solani* dominating the group of pathogenic endophytic fungi, which have a

wide range of host plants[68]. It is well known that the high abundance of *R. solani* can

severely damage their host plants[69,70]. A strong reduction of root biomass in our study during

extreme drought could also have reduced the abundance of such endophytic fungi[71,72], which

subsequently promoted the performance of invading plants[73]. A recent study further

suggested that fungal community responses to drought were closely related to root trait

responses to soil water scarcity although pathogenic fungal responses were rather weak to

drought[74]. We still know little about when and how (plant) host-associated pathogenic fungal

species would respond to extreme droughts. The present results help us speculate that if

some pathogens decline at the expense of lower resource availability (e.g., root biomass),

invading plants can exploit such environments to establish themselves. Whether this

mechanism further leads to a fast turnover of plant communities after extreme droughts

remains to be tested.

Despite the overall weak soil legacy effects of native plant diversity relative to extreme

droughts on the performance of all three types of invaders, our results still highlight biotic

facilitation in promoting plant performance in these soils, specifically through an increase in

root biomass of the invading plants. For the success of alien plants in native plant diversity

soils, diversity-dependent reduction in mycoparasites (composed of basidiomycete taxa like

*Chalciporus*, *Cystobasidium* and ascomycete taxa *Purpureocillium*) played some role (e.g.,

Figure 4 B). Mycoparasites are usually considered to regulate fungal pathogens and thereby

promote the growth of plants[75], but their decline could also be associated with increases of

some other beneficial fungi that could foster the establishment of invading alien plants.

Both positive soil legacy effects of extreme drought and native plant diversity were in part

driven by the reduction of root biomass production in resident native plants during the

conditioning phase. Many previous studies have pointed to the importance of resource

availability in the soil driving the success of invading plants[3,76]. Our results indeed suggest

that enhanced performance of invading plants to partly associated with both resource

availabilities in terms of root biomass and with the reduction of mycoparasites. The

remaining roots of resident native plants in soils could affect nutrient availability through root

decomposition, and further influence soil microorganisms that subsequently could determine

the success of a new invading plant. We accordingly suggest that if an extreme drought

lowers the root production in native plant communities and thereby alters soil fungal

communities, it could open up opportunities for invading plants to establish, at least for non-

resident native and alien plants used in our study. Indeed, our results are entirely based on

soil legacy effects, and it is likely that the presence of native resident plants (for instance

those that survive the extreme drought) could constrain the establishment of invading plants

after extreme droughts through competitive interactions.  Future studies can shed insight on

what other aspects of plant roots (e.g., morphological and/or chemical traits) of native plant

communities during extreme droughts and subsequent effects on soil fungal communities

can influence the performance of invading plants.

A central question of 21[st]-century ecology is how multiple global change factors determine

the dynamics of biodiversity change in ecosystems[77,78]. Extreme drought events are going to

be a key driver of biodiversity dynamics, particularly through altering the plant- and soil

community shifts in terrestrial ecosystems[79–82]. Our results confirm that the performance of

invading plants enhances in soils with the legacy of extreme drought, and these effects may

even exist independent of the background diversity of resident native plants. Moreover, we

highlight that such drought legacies in soil are mediated through how extreme drought would

alter the performance of previously present native plant communities and their associations

with soil fungi.

**Materials and Methods**

*Conditioning phase*

The experiment consisted of two stages, the conditioning phase, and the feedback phase

(Figure 1). The conditioning phase was performed to create the soil legacy of a native plant

diversity gradient crossed with ambient water and extreme drought conditions. We grew four

different native species to simulate different native plant diversity levels from 1 to 4 (Figure

1). These plant species were used based on their native status in Europe and their common

co-occurrence in grasslands. Plant seeds were obtained from a local nursery company that

collects seeds from the field (https://www.cruydthoeck.nl/). We added plants as seedling

(about 2-3 cm in height) to gamma-sterilized soils (also collected from the field before the

sterilization: Vlietberg, the Netherlands, 51° 86′ N, 5° 89' E), which were then grown as

monocultures, two species combination, three species combination, and four species

combination - each containing four individuals as per the substitutive design. Live soils for

the conditioning phase were collected where these species naturally grow from six different

sites near Nijmegen, the Netherlands (close to the site where soils were collected for

sterilization). These soils varied in their C, N and P content (Supplementary table 2), which is

why we used soils from specific sites (even when they were in close vicinity) as random

intercepts in mixed models (details below). We planted plant individuals in two-liter PVC pots

(~1800 g of sterilized soils mixed with 200 g of alive soil from the field, pot diameter 12.3 cm;

pot height 13.0 cm) during the conditioning phase with all diversity levels replicated 24 times

to create an adequate amount of soils for the feedback phase. Moreover, the field soil

inoculation (from six sites) to these pots was replicated four times which led to 96 pots for

native diversity treatments. We then doubled the diversity gradient pots (96) to create both

ambient and extreme drought treatments. Please note that given the substitutive design, we

were not able to replicate all compositions of plant pairs and three species combinations,

and our statistical analysis is based on the diversity levels (1, 2, 3 and 4), which were

adequately replicated during the conditioning phase.

Moisture levels during the conditioning phase were monitored regularly by weighing the pots

and correcting when large differences (e.g., exceeding a difference of ~10 g) within the

treatments occurred. The conditioning of the soil with a native plant diversity gradient took

place for ten weeks in total (eight weeks with a regular supply of water, and the last two

weeks with half of the pots receiving no water at all). This extreme drought treatment

resulted in a complete wilting of plants at the end of the conditioning phase (Figure 1). At the

end of the conditioning phase, the moisture percentage in the soil was calculated using the

gravimetric method to evaluate soil water differences between ambient and extreme drought

conditioning treatments. The extreme drought treatment resulted in four times less soil

moisture compared to the ambient conditions, which was sufficient to create wilting in plants

across diversity treatments (Figure 1 b, Supplementary figure 3).

At the end of the conditioning phase (week 10 of the plant growth), plant shoots were

clipped, roots were collected from half of the conditioned soil, and some of the conditioned

soils were kept for the measurement of soil abiotic and fungal analysis. The roots were

washed thoroughly to ensure no influence of residual soil on the root biomass. Both roots

and shoots were dried at 40 °C for several days and weighed afterwards for the estimation

of the shoot and root dry biomass. Soil samples (approximately ten grams) were taken from

the conditioned to measure soil C, N and P content. Amounts of carbon (C) and nitrogen (N)

were determined using an elemental analyzer (Flash EA 1112, Thermo Scientific) by

following the Micro-Dumas combustion method. We further determined the amount of

phosphorous (P) by extracting $NaHCO_3$ and determining the plant-available

orthophosphates in the soil following the Olsen P method [83].

*Fungal groups measurements*

We measured the relative abundance of various soil fungal groups from the conditioning

phase soils by amplicon sequencing using the Illumina Miseq PE300 platform at BGI (Hong

Kong). Total DNA was extracted from 250 mg of fresh soil using the DNeasy Powersoil kit

(Qiagen, Hilden, Germany) according to manufactures protocol. For multiplexing and re-

identification individual samples adapters and barcodes were added to samples by

polymerase chain reaction (PCR). We used the primers ITS4 and ITS9 targeting the ITS2

region[84]. PCRs were performed in 25-µl reaction mixtures and contained 2.5 µl dNTP (4005

µM), 0.15 µl of FastStart Expand High Fidelity polymerase (Roche Applied Sciences), 2.5 µl

10× PCR buffer with MgCl2, 1µl MgCl2 (25mM), 1.25 µl BSA (4mg/ml), 0.5 µl of each of the

two primers (10mM) and 1 µl soil DNA (~5–20 ng). The PCR conditions involved initial step

at 95°C for 5 min, followed by 35 cycles of 95°C for 45 seconds and 54°C for 60 seconds,

and then 72°C for 90 seconds. The final extension step was 72°C for 10 minutes. The final

PCR product was purified using Agencourt AMPure XP magnetic bead system (Beckman

Coulter Life Sciences, Indianapolis, Indiana, USA) with a volume ratio of PCR product to

beads of 1: 0.7.  After purification, we analyzed the PCR products in a Fragment Analyzer

using a Standard Sensitivity NGS Fragment Analysis kit (1bp-6000bp) and following

manufacturer's instructions (Advanced Analytical Technologies GmbH, Heidelberg,

Germany). The purified amplicons were diluted to the same concentration before sending for

sequencing using the Illumina MiSeq platform for 300 bp paired-end reads.

ITS sequences were analyzed using the PIPITS pipeline (version 2.7, standard settings,

Gweon et al. 2015) with VSEARCH implemented to pair sequences. The ITS2 region was

extracted using ITSx [86] and short reads (<100bp) were removed. Chimeric sequences were

removed by comparing with UNITE uchime database (version 8.2). Clustering of reads into

OTUs was performed at an identity threshold of 97%. Sequences were aligned and

classified using with RDP against the UNITE fungal database[87,88] to obtain a match with the

closest known species for each OTU. The resulting OTUs were parsed against the FunGuild

(v1.1) database to assign putative life strategies[89] for the OTUs with 'probable' or 'highly

probable' classification and the assignment was further curated using in-house databases[90].

We resolved 12 major fungal functional groups that were used in further analysis. These

groups were 1) plant pathogenic fungi, 2) total endophytic fungi, 3) endophytic fungal

pathogens, 4) arbuscular mycorrhizal fungi, 5) ectomycorrhizal fungi, 6) ericoid mycorrhizal

fungi, 7) animal pathogenic fungi, 8) epiphytic fungi, 9) fungal parasites, 10) total

saprotrophic fungi, 11) saprotrophic endophytic fungi, 12) wood saprotrophic fungi.

*Feedback phase*

To prepare for the feedback phase, the conditioned soils (300 grams) were mixed with

gamma-sterilized soils (300 grams)- these soils were the same that were collected from the

field from Vlietberg, the Netherlands (51° 86′ N, 5° 89' E) for the conditioning phase. One-

liter PVC pots (width 10 cm; length 10 cm; height 11 cm) were planted with an individual

plant (Figure 1). Seeds of feedback phase plants were also purchased from the same local

nursery company (https://www.cruydthoeck.nl/) where we purchased seeds for the

conditioning phase plants, except that *Ceratochloa sitchensis* (the alien Poaceae) and

*Senecio inaequidens* (the alien Asteraceae) were collected from the field in the Netherlands

(near Wageningen). List of all 12 feedback-phase plant species is provided in Figure 1.

These commonly found grassland plants were chosen in the feedback phase to create a

phylogenetically balanced design relative to the conditioning phase (Figure 1). Each of these

twelve species was planted after germination (about 2-3 cm in height) to the feedback phase

pots filled with conditioned soils with a history of different native plant diversity levels (one to

four species) and drought treatments.  We grew each of the 12 feedback species separately

to examine their performance in the conditioned soils. Each feedback species was replicated

six times per drought and diversity treatments, which were distributed evenly across six soil

origins from the conditioning phase. In this way, we examined the performance 576 plant

individuals during the feedback phase. The plants were grown for six weeks and were

watered regularly. The moisture difference between the extreme drought soils and ambient

soils was overcome by adding 36 ml extra water to extreme drought soils at the beginning of

the feedback phase. In this way, we excluded the effects of moisture difference already at

the start of the feedback phase. After six weeks, the shoots were harvested by clipping,

dried (again at 40 °C for several days) and weighed. Roots were harvested by washing them

out of the soil gently and carefully prior to drying (also at 40 °C for several days) and

weighing.

*Statistical analysis*

Both conditioning and feedback phase response variables were analyzed using mixed-

effects models. For the feedback phase, we explained variation in invader plant performance

(shoot and root biomass) using a three-way interaction model with native plant diversity,

drought, and invader type as three fixed effects, whereas soil inoculum was used as random

intercept with six different levels (i.e., six different soils). Additionally, we used specific plant

species as another random intercept in our feedback phase mixed-effect models (Table 1).

Given that we measured many variables during the conditioning phase to understand soil

biotic and abiotic characteristics as well as plant performance, we ran models separately for

all those variables (Table 2). For all these mixed-effect models, six soil inoculums were

again used as a random intercept, whereas native plant diversity and drought were the fixed

effects. The list of all the response variables during the conditioning phase is provided in

table 2, and could be categorized into plant community, soil abiotic variable and soil fungal

group responses. Linearity assumptions of the model were visually inspected using the

homogeneity of variance, and when not met, we either log-transformed the data or used

negative binomial error distribution instead of normal distribution (indicated in Table 2).

Finally, we examined the relationship between the soil legacy treatments and the

performance of invading plants in the feedback phase using path models to be able to

distinguish between direct and indirect effects from the conditioning phase treatments that

influence the feedback phase invading plants. Model structures in our path models were

similar to that of mixed-effect models for conditioning and feedback phase response

variables. However, since we ran path models separately for three types of invading plants,

we excluded specific plant species as a random term. Indeed, soil origin having six levels

compared to four levels of plant species as a random intercept within the invader type

provides a more robust parameter estimation in mixed models [91]. Moreover, two independent

random intercepts (soil type and plant species) in mixed-effect models prevented model

convergence, which is why we opted to use a single random term instead of two in all our

path models.  We included soil fungal and plant response data from the conditioning phase

and linked them to invading plant performance (shoot and root biomass) during the feedback

phase. For some variables, we tested both pathways to identify the causal links (Figure 4).

Model selection was used based on the inclusion of variables that resulted in non-saturated

path models e.g., model overfit) as per Shipley's test of d-separation which yields Fisher's C

statistic (Chi-square distributed)[92]. None of our path models showed a good fit (p-value

<0.05) when ran for range-expanding invaders with similar set of variables (e.g., soil fungal

groups and/or soil nutrients) used for local native and alien invasive plants. The inclusion of

soil nutrients in path models also failed to provide a good fit (p-value<0.05) for non-resident

native and alien plants. We eventually selected two path models-one for non-resident and

one for alien invaders with plant biomass responses and one of the fungal group responses

to associate with the respective performance (both shoot and root biomass) of invading

plants (Figure 4). Moreover, the path model for the alien plant was further reduced to obtain

a good fit (the direct paths from conditioning phase plant diversity to feedback phase plant

shoot and root biomass were removed), although for the sake of comparison between local

and alien invaders, we have shown the full model for alien plants as well (Figure 4).

All statistical analyses were carried out in R statistical software version 4.1.0 [93]. Mixed effect

models were run with the lme4 package for R, whereas we used the performance package

to check the linearity model assumption[94]. The F-value, p-value and degrees of freedom

were estimated using the lmerTest package[95]. Multi-variate analysis of fungal communities

were performed using the *manyglm* function from the mvabund package which incorporates

the framework of generalized linear models [96]. In path models, we used the nlme package[97]

to run the mixed-effect models. The path model analysis was carried out with the

piecewiseSEM package[98].

**Acknowledgements**

We thank Jake Alexander and Arjen Biere for their suggestions on the experimental design.

We are grateful to Ludovico Formenti for his help in drawing figure 1. MPT acknowledges the

funding from the Swiss State Secretariat for Education, Research and Innovation (SERI)

under contract number M822.00029. WHvdP acknowledges the support from ERC advanced

grant (ERC-ADV 323020 SPECIALS).

**Author contribution:** MPT and WHvdP conceived the study. MvdS, MPT, RAW, SEH, FtH,

SG, CQ, KS performed the experiment. MPT analysed the data with inputs from SEH and

wrote the manuscript together with WHvdP. All authors contributed to revisions.

**Competing Interest Statement:** The authors declare no conflict of interest.

**Data availability:** All data used for the analysis is uploaded as a supplementary data file.

**References**

1.  Darwin, C. *On the origin of species by means of natural selection*. (J. Murray, 1859).

2.  Elton, C. *The ecology of invasions by animals and plants*. (Methuen, 1958).

3.  Davis, M. A., Grime, J. P. & Thompson, K. Fluctuating resources in plant communities: A general theory of invasibility. *Journal of Ecology* **88**, 528–534 (2000).

4.  Pearson, D. E., Ortega, Y. K., Eren, Ö. & Hierro, J. L. Community Assembly Theory as a Framework for Biological Invasions. *Trends Ecol Evol* **33**, 313–325 (2018).

5.  Parepa, M., Fischer, M. & Bossdorf, O. Environmental variability promotes plant invasion. *Nat Commun* **4**, 1–4 (2013).

6.  Ricciardi, A. *et al.* Invasion Science: A Horizon Scan of Emerging Challenges and Opportunities. *Trends Ecol Evol* **32**, 464–474 (2017).

7.  Blackburn, T. M. *et al.* A proposed unified framework for biological invasions. *Trends Ecol Evol* **26**, 333–339 (2011).

8.  Bock, D. G. *et al.* What we still don't know about invasion genetics. *Mol Ecol* **24**, 2277–2297 (2015).

9.  van Kleunen, M., Bossdorf, O. & Dawson, W. The Ecology and Evolution of Alien Plants. *Annu Rev Ecol Evol Syst* **49**, 25–47 (2018).

10. Levine, J. M. & D'Antonio, C. M. Elton Revisited : A Review of Evidence Linking Diversity and Invasibility. *Oikos* **87**, 15–26 (1999).

11. Beaury, E. M., Finn, J. T., Corbin, J. D., Barr, V. & Bradley, B. A. Biotic resistance to invasion is ubiquitous across ecosystems of the United States. *Ecol Lett* **23**, 476–482 (2020).

12. Feng, Y., Fouqueray, T. D. & van Kleunen, M. Linking Darwin's naturalisation hypothesis and Elton's diversity–invasibility hypothesis in experimental grassland communities. *Journal of Ecology* **107**, 794–805 (2019).

13. Fridley, A. J. D. *et al.* The Invasion Paradox : Reconciling Pattern and Process in Species Invasions. *Ecology* **88**, 3–17 (2008).

14. Tomasetto, F., Duncan, R. P. & Hulme, P. E. Resolving the invasion paradox: pervasive scale and study dependence in the native-alien species richness relationship. *Ecol Lett* **22**, 1038–1046 (2019).

15. Smith, N. S. & Côté, I. M. Multiple drivers of contrasting diversity–invasibility relationships at fine spatial grains. *Ecology* **100**, 1–11 (2019).

16. Tanentzap, A. J., Bazely, D. R. & Lafortezza, R. Diversity-invasibility relationships across multiple scales in disturbed forest understoreys. *Biol Invasions* **12**, 2105–2116 (2010).

17. Herbold, B. & Moyle, P. Introduced Species and Vacant Niches. *Am Nat* **128**, 751–760 (1986).

18. Davis, M., Thomposon, K., Grime, J. & Liebhold, A. The Local Mechanism Driving Community Assembly and Species Diversity. *Ecography* **28**, 696–704 (2005).

19. Theoharides, K. A. & Dukes, J. S. Plant invasion across space and time: Factors affecting nonindigenous species success during four stages of invasion. *New Phytologist* **176**, 256–273 (2007).

20. Zhang, Z., Liu, Y., Brunel, C. & van Kleunen, M. Soil-microorganism-mediated invasional meltdown in plants. *Nat Ecol Evol* **4**, 1612–1621 (2020).

21. Wolfe, B. E. & Klironomos, J. N. Breaking new ground: Soil communities and exotic plant invasion. *Bioscience* **55**, 477–487 (2005).

22. Palmieri, D., Vitale, S., Lima, G., Di Pietro, A. & Turrà, D. A bacterial endophyte exploits chemotropism of a fungal pathogen for plant colonization. *Nat Commun* **11**, 1–11 (2020).

23. Callaway, R. M., Thelen, G. C., Rodriguez, A. & Holben, W. E. Soil biota and exotic plant invasion. *Nature* **427**, 731–733 (2004).

24. Van der Putten, W. H., Klironomos, J. N. & Wardle, D. A. Microbial ecology of biological invasions. *ISME J* **1**, 28–37 (2007).

25. Reinhart, K. O., Packer, A., Van Der Putten, W. H. & Clay, K. Plant-soil biota interactions and spatial distribution of black cherry in its native and invasive ranges. *Ecol Lett* **6**, 1046–1050 (2003).

26. Zhang, Z., Liu, Y., Brunel, C. & van Kleunen, M. Evidence for Elton's diversity–invasibility hypothesis from belowground. *Ecology* **101**, 1–10 (2020).

27. Shen, C., Wang, J., He, J. Z., Yu, F. H. & Ge, Y. Plant Diversity Enhances Soil Fungal Diversity and Microbial Resistance to Plant Invasion. *Appl Environ Microbiol* **87**, (2021).

28. Mommer, L. *et al.* Lost in diversity: the interactions between soil-borne fungi, biodiversity and plant productivity. *New Phytologist* **218**, 542–553 (2018).

29. Thakur, M. P. *et al.* Plant–Soil Feedbacks and Temporal Dynamics of Plant Diversity–Productivity Relationships. *Trends Ecol Evol* **36**, 651–661 (2021).

30. Maron, J. L., Marler, M., Klironomos, J. N. & Cleveland, C. C. Soil fungal pathogens and the relationship between plant diversity and productivity. *Ecol Lett* **14**, 36–41 (2011).

31. Zhang, Z., Liu, Y., Brunel, C. & van Kleunen, M. Evidence for Elton's diversity–invasibility hypothesis from belowground. *Ecology* **101**, 1–10 (2020).

32. Isbell, F. *et al.* Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature* **526**, 574–577 (2015).

33. Wagg, C. *et al.* Plant diversity maintains long-term ecosystem productivity under frequent drought by increasing short-term variation. *Ecology* **98**, 2952–2961 (2017).

34. Bennett, J. A. *et al.* Resistance of soil biota and plant growth to disturbance increases with plant diversity. *Ecol Lett* **23**, 119–128 (2020).

35. Orbán, I., Szitár, K., Kalapos, T. & Körel-Dulay, G. The role of disturbance in invasive plant establishment in a changing climate: insights from a drought experiment. *Biol Invasions* **23**, 1877–1890 (2021).

36.  Liu, Y. *et al.* Do invasive alien plants benefit more from global environmental change than
native plants? *Glob Chang Biol* **23**, 3363–3370 (2017).

37.  Hueso, S., García, C. & Hernández, T. Severe drought conditions modify the microbial
community structure, size and activity in amended and unamended soils. *Soil Biol Biochem*
**50**, 167–173 (2012).

38.  Rubin, R. L., van Groenigen, K. J. & Hungate, B. A. Plant growth promoting rhizobacteria are
more effective under drought: a meta-analysis. *Plant Soil* **416**, 309–323 (2017).

39.  Leizeaga, A., Hicks, L. C., Manoharan, L., Hawkes, C. V. & Rousk, J. Drought legacy affects
microbial community trait distributions related to moisture along a savannah grassland
precipitation gradient. *Journal of Ecology* 1–16 (2020) doi:10.1111/1365-2745.13550.

40.  Meisner, A., Jacquiod, S., Snoek, B. L., Ten Hooven, F. C. & van der Putten, W. H. Drought
legacy effects on the composition of soil fungal and prokaryote communities. *Front Microbiol*
**9**, 1–12 (2018).

41.  de Vries, F. *et al.* Soil bacterial networks are less stable under drought than fungal networks.
*Nat Commun* **9**, 3033 (2018).

42.  Müller, L. M. & Bahn, M. Drought legacies and ecosystem responses to subsequent drought.
*Glob Chang Biol* **28**, 5086–5103 (2022).

43.  Meisner, A., De Deyn, G. B., De Boer, W. & Van Der Putten, W. H. Soil biotic legacy effects of
extreme weather events influence plant invasiveness. *Proc Natl Acad Sci U S A* **110**, 9835–
9838 (2013).

44.  Essl, F. *et al.* Which Taxa Are Alien? Criteria, Applications, and Uncertainties. *Bioscience* **68**,
496–509 (2018).

45.  Essl, F. *et al.* A Conceptual Framework for Range-Expanding Species that Track Human-
Induced Environmental Change. *Bioscience* **69**, 908–919 (2019).

46.  Peng, S., Kinlock, N. L., Gurevitch, J. & Peng, S. Correlation of native and exotic species
richness: a global meta-analysis finds no invasion paradox across scales. *Ecology* **100**, 1–10
(2019).

47.  Bever, J. D., Westover, K. M. & Antonovics, J. Incorporating the Soil Community into Plant
Population Dynamics: The Utility of the Feedback Approach. *J Ecol* **85**, 561 (1997).

48.  Van der Putten, W. H. *et al.* Plant-soil feedbacks: The past, the present and future challenges.
*Journal of Ecology* **101**, 265–276 (2013).

49.  Harrison, S., Spasojevic, M. J. & Li, D. Climate and plant community diversity in space and
time. *Proceedings of the National Academy of Sciences* 201921724 (2020)
doi:10.1073/pnas.1921724117.

50.  Schwalm, C. R. *et al.* Global patterns of drought recovery. *Nature* **548**, 202–205 (2017).

51.  Prugh, L. R. *et al.* Ecological winners and losers of extreme drought in California. *Nat Clim
Chang* **8**, 819–824 (2018).

52.  Griffin-Nolan, R. J. *et al.* Shifts in plant functional composition following long-term drought in
grasslands. *Journal of Ecology* **107**, 2133–2148 (2019).

53. Ploughe, L. W. *et al.* Community Response to Extreme Drought (CRED): a framework for
drought-induced shifts in plant–plant interactions. *New Phytologist* **222**, 52–69 (2019).

54. Fuchslueger, L. *et al.* Drought history affects grassland plant and microbial carbon turnover
during and after a subsequent drought event. *Journal of Ecology* **104**, 1453–1465 (2016).

55. Manea, A., Sloane, D. R. & Leishman, M. R. Reductions in native grass biomass associated
with drought facilitates the invasion of an exotic grass into a model grassland system.
*Oecologia* **181**, 175–183 (2016).

56. Jiménez, M. A. *et al.* Extreme climatic events change the dynamics and invasibility of semi-
arid annual plant communities. *Ecol Lett* **14**, 1227–1235 (2011).

57. Schrama, M. & Bardgett, R. D. Grassland invasibility varies with drought effects on soil
functioning. *Journal of Ecology* **104**, 1250–1258 (2016).

58. Vannier, N., Bittebiere, A. K., Mony, C. & Vandenkoornhuyse, P. Root endophytic fungi impact
host plant biomass and respond to plant composition at varying spatio-temporal scales.
*Fungal Ecol* **44**, 100907 (2020).

59. Rodriguez, R. J., White, J. F., Arnold, A. E. & Redman, R. S. Fungal endophytes: Diversity and
functional roles: Tansley review. *New Phytologist* **182**, 314–330 (2009).

60. Dickie, I. A. *et al.* The emerging science of linked plant–fungal invasions. *New Phytologist* **215**,
1314–1332 (2017).

61. Inderjit & van der Putten, W. H. Impacts of soil microbial communities on exotic plant
invasions. *Trends Ecol Evol* **25**, 512–519 (2010).

62. Pringle, A. *et al.* Mycorrhizal symbioses and plant invasions. *Annu Rev Ecol Evol Syst* **40**, 699–
715 (2009).

63. Aschehoug, E. T., Metlen, K. L., Callaway, R. M. & Newcombe, G. Fungal endophytes directly
increase the competitive effects of an invasive forb. *Ecology* **93**, 3–8 (2012).

64. Perez, L. I. *et al.* Can seed-borne endophytes promote grass invasion by reducing host
dependence on mycorrhizas? *Fungal Ecol* **52**, (2021).

65. Gonzalez Mateu, M., Baldwin, A. H., Maul, J. E. & Yarwood, S. A. Dark septate endophyte
improves salt tolerance of native and invasive lineages of Phragmites australis. *ISME Journal*
**14**, 1943–1954 (2020).

66. Geisen, S. *et al.* Seed and root endophytic fungi in a range expanding and a related plant
species. *Front Microbiol* **8**, 1–11 (2017).

67. Mattoo, A. J. & Nonzom, S. Endophytic fungi: understanding complex cross-talks. *Symbiosis*
**83**, 237–264 (2021).

68. Hane, J. K., Anderson, J. P., Williams, A. H., Sperschneider, J. & Singh, K. B. Genome
Sequencing and Comparative Genomics of the Broad Host-Range Pathogen Rhizoctonia solani
AG8. *PLoS Genet* **10**, 11–12 (2014).

69. Anees, M., Edel-Hermann, V. & Steinberg, C. Build up of patches caused by Rhizoctonia solani.
*Soil Biol Biochem* **42**, 1661–1672 (2010).

70. Cubeta, M. & Vilgalys, R. Population Biology of the Rhizoctonia solani Complex.
*Phytopathology* **87**, 480–484 (1997).

71. Staddon, P. L., Gregersen, R. & Jakobsen, I. The response of two Glomus mycorrhizal fungi
and a fine endophyte to elevated atmospheric CO2, soil warming and drought. *Glob Chang*
*Biol* **10**, 1909–1921 (2004).

72. Lagueux, D. *et al.* Experimental drought re-ordered assemblages of root-associated fungi
across North American grasslands. *Journal of Ecology* **109**, 776–792 (2021).

73. Buchenau, N., van Kleunen, M. & Wilschut, R. A. Direct and legacy-mediated drought effects
on plant performance are species-specific and depend on soil community composition. *Oikos*
1–14 (2022) doi:10.1111/oik.08959.

74. Lozano, Y. M., Aguilar-Trigueros, C. A., Roy, J. & Rillig, M. C. Drought induces shifts in soil
fungal communities that can be linked to root traits across 24 plant species. *New Phytologist*
**232**, 1917–1929 (2021).

75. Adams, P. B. The potential of mycoparasites for biological control of plant diseases. *Annual*
*review of phytopathology. Vol. 28* 59–72 (1990) doi:10.1146/annurev.py.28.090190.000423.

76. Kardol, P., Souza, L. & Classen, A. T. Resource availability mediates the importance of priority
effects in plant community assembly and ecosystem function. *Oikos* **122**, 84–94 (2013).

77. Franklin, J., Serra-Diaz, J. M., Syphard, A. D. & Regan, H. M. Global change and terrestrial
plant community dynamics. *Proc Natl Acad Sci U S A* **113**, 3725–3734 (2016).

78. Komatsu, K. J. *et al.* Global change effects on plant communities are magnified by time and
the number of global change factors imposed. *Proc Natl Acad Sci U S A* **116**, 17867–17873
(2019).

79. Kaisermann, A., de Vries, F. T., Griffiths, R. I. & Bardgett, R. D. Legacy effects of drought on
plant–soil feedbacks and plant–plant interactions. *New Phytologist* **215**, 1413–1424 (2017).

80. Zhang, F. *et al.* When does extreme drought elicit extreme ecological responses? *Journal of*
*Ecology* **107**, 2553–2563 (2019).

81. Rillig, M. C. *et al.* The role of multiple global change factors in driving soil functions and
microbial biodiversity. *Science (1979)* **366**, 886–890 (2019).

82. Meisner, A., De Deyn, G. B., De Boer, W. & Van Der Putten, W. H. Soil biotic legacy effects of
extreme weather events influence plant invasiveness. *Proc Natl Acad Sci U S A* **110**, 9835–
9838 (2013).

83. Olsen, S. R. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*.
(US Department of Agriculture, 1954).

84. Ihrmark, K. *et al.* New primers to amplify the fungal ITS2 region - evaluation by 454-
sequencing of artificial and natural communities. *FEMS Microbiol Ecol* **82**, 666–677 (2012).

85. Gweon, H. S. *et al.* PIPITS: An automated pipeline for analyses of fungal internal transcribed
spacer sequences from the Illumina sequencing platform. *Methods Ecol Evol* **6**, 973–980
(2015).

86.    Bengtsson-Palme, J. *et al.* Improved software detection and extraction of ITS1 and ITS2 from
ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental
sequencing data. *Methods Ecol Evol* **4**, 914–919 (2013).

87.    Kõljalg, U. *et al.* The taxon hypothesis paradigm—On the unambiguous detection and
communication of taxa. *Microorganisms* **8**, 1–24 (2020).

88.    Nilsson, R. H. *et al.* The UNITE database for molecular identification of fungi: Handling dark
taxa and parallel taxonomic classifications. *Nucleic Acids Res* **47**, D259–D264 (2019).

89.    Nguyen, N. H. *et al.* FUNGuild: An open annotation tool for parsing fungal community
datasets by ecological guild. *Fungal Ecol* **20**, 241–248 (2016).

90.    Hannula, S. E. *et al.* Shifts in rhizosphere fungal community during secondary succession
following abandonment from agriculture. *ISME Journal* **11**, 2294–2304 (2017).

91.    Silk, M. J., Harrison, X. A. & Hodgson, D. J. Perils and pitfalls of mixed-effects regression
models in biology. *PeerJ* **8**, 1–20 (2020).

92.    Shipley, B. Confirmatory path analysis in a generalized multilevel context. *Ecology* **90**, 363–
368 (2009).

93.    R Core Team. R: A language and environment for statistical computing. Preprint at (2016).

94.    Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *J Open Source Softw*
**6**, 3139 (2021).

95.    Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear
Mixed Effects Models . *J Stat Softw* **82**, (2017).

96.    Wang, Y., Naumann, U., Wright, S. T. & Warton, D. I. Mvabund- an R package for model-based
analysis of multivariate abundance data. *Methods Ecol Evol* **3**, 471–474 (2012).

97.    Pinheiro, J., Bates, D., Debroy, S. & Sarkar, D. _nlme: Linear and Nonlinear Mixed Effects
Models_. R package version 3.1-152. Preprint at (2021).

98.    Lefcheck, J. S. piecewiseSEM: Piecewise structural equation modelling in r for ecology,
evolution, and systematics. *Methods Ecol Evol* **7**, 573–579 (2016).

**Table 1:** Effects of soil legacy of native plant diversity and extreme drought on the performance (shoot and root biomass) of invading plants of three categories (non-resident natives, range expanders and alien plants). The bold values indicate statistical significance (p-value<0.05). The degree of freedoms is estimated using Kenward-Roger approximations for mixed-effect models. Variance components are used as random intercept in mixed-effects models. Marginal $R^2$ represents model fit with only fixed effects, whereas conditional $R^2$ represents model fit with both fixed and random effects. sd stands for standard deviation. The upward arrows indicate increase of a given response variable (shown only for significant variables and those with

| Treatments (fixed effects) | Shoot biomass | | | | Root biomass (log-transformed) | | | |
|---|---|---|---|---|---|---|---|---|
| | F-value$_{df}$ | p-value | Marginal $R^2$ | Conditional $R^2$ | F-value$_{df}$ | p-value | Marginal $R^2$ | Conditional $R^2$ |
| Plant species richness (S) | $2.480_{1,\ 523.29}$ | 0.115 | | | **$4.747_{1,\ 523.88}$** | **0.029** ↑ | | |
| Extreme drought (D) | **$7.104_{1,523.31}$** | **0.007** ↑ | | | $0.101_{1,523.77}$ | 0.749 | | |
| Plant invader type (I) | **$4.487_{2,\ 14.51}$** | **0.030** | | | $2.066_{2,\ 18.89}$ | 0.154 | | |
| S×D | $0.081_{1,523.38}$ | 0.775 | 0.22 | 0.45 | $1.252_{1,523.89}$ | 0.263 | 0.06 | 0.24 |
| S×I | $1.742_{2,523.38}$ | 0.176 | | | $1.107_{2,524.03}$ | 0.331 | | |
| D×I | $0.647_{2,523.32}$ | 0.523 | | | $0.543_{2,523.73}$ | 0.581 | | |
| S×D×I | $1.245_{2,523.36}$ | 0.288 | | | $0.434_{2,523.81}$ | 0.647 | | |
| | | | | | | | | |
| **Variance components** | **Variance (sd)** | | | | **Variance (sd)** | | | |
| Soil collection site | <0.001 (0.019) | | | | <0.001 (0.027) | | | |
| Plant species | 0.004 (0.069) | | | | 0.080 (0.283) | | | |

**Table 2:** Effects of plant species richness, extreme drought and their interactions on plant biomass, soil abiotics and the abundance of soil fungal groups during the conditioning phase. The bold values indicate statistical significance (p-value<0.05) and italics are marginally significant (p-value<0.1). The upward and downward arrows indicate increase and decrease of a given response variable, respectively (shown only for statistically significant variables). Soil collection site was used as a random intercept in mixed models for which variance is reported. All mixed models were based on Gaussian error terms unless indicated by # for a given response variable (all other models included negative binomial error terms to account for overdispersion in the data).

| Response variables | Plant species richness (PSR) | | Extreme drought (D) | | PSR×D | | Soil collection site |
|---|---|---|---|---|---|---|---|
| | $F\text{-value}_{df}$ | p-value | $F\text{-value}_{df}$ | p-value | $F\text{-value}_{df}$ | p-value | Variance |
| ***Plant biomass*** | | | | | | | |
| Shoot biomass | $0.154_{1,183}$ | 0.695 | **$28.523_{1,183}$** | **<0.001** ↓ | $0.713_{1,183}$ | 0.399 | 0.371 |
| Root biomass (log) | **$20.491_{1,183}$** | **<0.001** ↓ | **$13.832_{1,183}$** | **<0.001** ↓ | $1.432_{1,183}$ | 0.232 | 0 |
| Total biomass | **$12.390_{1,183}$** | **<0.001** ↓ | **$64.956_{1,183}$** | **<0.001** ↓ | $0.064_{1,183}$ | 0.799 | 0.673 |
| ***Soil abiotics*** | | | | | | | |
| Soil C | $2.459_{1,183}$ | 0.118 | $0.057_{1,183}$ | 0.810 | $0.024_{1,183}$ | 0.875 | 0.295 |
| Soil N | $1.993_{1,183}$ | 0.159 | $0.697_{1,183}$ | 0.404 | $0.656_{1,183}$ | 0.418 | <0.001 |
| Soil $PO_4$ | **$15.596_{1,183}$** | **<0.001** ↓ | $2.622_{1,183}$ | 0.107 | $2.863_{1,183}$ | 0.092 | 11.180 |
| ***Soil fungal abundance*** | | | | | | | |
| Total fungi | $3.736_{1,168.99}$ | 0.054 | $0.001_{1,168.40}$ | 0.967 | $0.054_{1,168.29}$ | 0.815 | 0 |
| Plant pathogenic fungi (log) | *$2.882_{1,168.12}$* | *0.091* ↓ | $0.005_{1,168.04}$ | 0.942 | $0.115_{1,168.03}$ | 0.734 | 0.270 |
| Total endophytic fungi (log) | **$5.848_{1,168.31}$** | **0.016** ↓ | **$6.112_{1,168.12}$** | **0.014** ↓ | $1.970_{1,168.09}$ | 0.162 | 0.047 |
| Endophytic fungal pathogens (log) | $0.001_{1,168.03}$ | 0.965 | $0.684_{1,168.01}$ | 0.409 | *$3.935_{1,168.01}$* | *0.048* ↑ | 1.619 |
| Arbuscular mycorrhizal fungi (log) | **$8.266_{1,168.11}$** | **<0.001** ↑ | $0.022_{1,168.04}$ | 0.882 | $0.410_{1,168.03}$ | 0.522 | 0.550 |
| Ectomycorrhizal fungi (log) | **$6.738_{1,168.22}$** | **0.010** ↑ | $0.262_{1,168.08}$ | 0.608 | $0.213_{1,168.06}$ | 0.644 | 0.115 |
| Ericoid mycorrhizal fungi# | 1.940 | 0.208 | 0.746 | 0.376 | 2.140 | 0.168 | 0.479 |
| Animal pathogenic fungi (log) | $1.628_{1,168.08}$ | 0.203 | *$4.281_{1,168.03}$* | *0.040* ↓ | $1.907_{1,168.02}$ | 0.169 | 0.258 |
| Epiphytic fungi# | 0.598 | 0.178 | 3.270 | 0.409 | *4.133* | *0.039* ↑ | <0.001 |
| Fungal parasites# | **6.889** | **0.002** ↓ | 0.889 | 0.073 | 2.292 | 0.124 | <0.001 |
| Saprotrophic fungi (log) | $0.075_{1,168.22}$ | 0.783 | $1.517_{1,168.08}$ | 0.219 | *$2.772_{1,168.06}$* | *0.097* ↑ | 0.050 |
| Saprotrophic endophytic fungi (log) | *$4.101_{1,168.05}$* | *0.044* ↓ | $0.019_{1,168.02}$ | 0.889 | $0.080_{1,168.01}$ | 0.777 | 0.494 |
| Wood saprotrophic fungi (log) | $0.498_{1,168.26}$ | 0.481 | $0.012_{1,168.10}$ | 0.912 | $0.001_{1,168.07}$ | 0.971 | 0.073 |

# negative binomial model

**Figure 1: (A)** Experimental schema showing the conditioning phase with native plant diversity crossed with extreme drought, and feedback phase where those soils were used to grow three groups of invading plants (non-resident natives, range-expanding and aliens) each containing four plant species and grown separately. **(B)** Examples of the extreme drought treatment effects at the end of the conditioning phase: the left image is a monoculture of *Holcus lanatus* with permanent wilting in extreme drought conditions (right side in the image) compared to the one in ambient water conditions, whereas the right image is a mixed plant community with 4 species exposed to ambient and extreme drought treatments (right side in the image). **(C)** List of the plant species used during the conditioning and feedback phase of the experiment.

**(A)**

Native plant diversity (1 to 4 species) and extreme drought

Conditioning phase

Feedback phase

Native colonizers
4 species

Range expanding colonizers
4 species

Alien colonizers
4 species

**(B)**

**(C)**

| Plants | Family | Plant origin | Soil legacy phase |
|---|---|---|---|
| *Holcus lanatus* | Poaceae | Native | Conditioning |
| *Barbarea vulgaris* | Brassicaceae | Native | Conditioning |
| *Platago lanceolata* | Plantaginaceae | Native | Conditioning |
| *Crepis biennis* | Asteraceae | Native | Conditioning |
| *Poa trivialis* | Poaceae | Native | Feedback |
| *Trifolium pratense* | Fabaceae | Native | Feedback |
| *Jacobaea vulgaris* | Asteraceae | Native | Feedback |
| *Leucanthemum vulgare* | Asteraceae | Native | Feedback |
| *Setaria verticillata* | Poaceae | Range-expander | Feedback |
| *Lathyrus latifolius* | Fabaceae | Range-expander | Feedback |
| *Tragopogon dubius* | Asteraceae | Range-expander | Feedback |
| *Centaurea stoebe* | Asteraceae | Range-expander | Feedback |
| *Ceratochloa sitchensis* | Poaceae | Alien | Feedback |
| *Lupinus polyphyllus* | Fabaceae | Alien | Feedback |
| *Solidago canadensis* | Asteraceae | Alien | Feedback |
| *Senecio inaequidens* | Asteraceae | Alien | Feedback |

**Figure 2:** Feedback phase shoot biomass of three invader type plants (each group contained 4 plant species) in soils conditioned by native plant species richness and extreme drought. Control soils were with regular addition of water. Means are shown as dark colored points together with standard errors. Dashed lines indicated regression lines, whereas raw data points are shown as light colored points.

**Figure 3:** Feedback phase root biomass (log-transformed to meet linear model assumptions) of three groups of invading plants (each group contained 4 plant species) in soils conditioned by native plant species richness and extreme drought. Control soils were with regular addition of water. Means are shown as dark colored points together with standard errors. Dashed lines indicated regression lines, whereas raw data points are shown as light colored points.

**(A) Non-resident native invaders**

Model fit: Fisher's C = 4.009, p-value = 0.135

| Predictors | | Responses | Standardized estimates | p-value |
|---|---|---|---|---|
| Plant species richness (S) | → | Root biomass | **-0.506** | **0.015** |
| Extreme drought (D) | → | Root biomass | **-0.569** | **<0.001** |
| S×D | → | Root biomass | 0.293 | 0.255 |
| Plant species richness (S) | → | Endophytic fungal pathogens | *-0.325* | *0.051* |
| Extreme drought (D) | → | Endophytic fungal pathogens | -0.125 | 0.349 |
| S×D | → | Endophytic fungal pathogens | *0.366* | *0.073* |
| Root biomass | → | Endophytic fungal pathogens | **-0.156** | **0.011** |
| Endophytic fungal pathogens | → | Root biomass | -0.125 | 0.119 |
| Plant species richness (S) | → | Shoot biomass (feedback phase) | -0.088 | 0.687 |
| Extreme drought (D) | → | Shoot biomass (feedback phase) | 0.265 | 0.134 |
| S×D | → | Shoot biomass (feedback phase) | 0.192 | 0.475 |
| Plant species richness (S) | → | Root biomass (feedback phase) | 0.266 | 0.261 |
| Extreme drought (D) | → | Root biomass (feedback phase) | 0.010 | 0.954 |
| S×D | → | Root biomass (feedback phase) | -0.243 | 0.400 |
| Root biomass | → | Shoot biomass (feedback phase) | -0.063 | 0.430 |
| Endophytic fungal pathogens | → | Shoot biomass (feedback phase) | **-0.210** | **0.014** |
| Root biomass | → | Root biomass (feedback phase) | 0.054 | 0.528 |
| Endophytic fungal pathogens | → | Root biomass (feedback phase) | -0.055 | 0.553 |

**(B) Alien invaders**

Model fit: Fisher's C = 11.616, p-value = 0.071

| Predictors | | Responses | Standardized estimates | p-value |
|---|---|---|---|---|
| Plant species richness (S) | → | Root biomass | **-0.480** | **0.021** |
| Extreme drought (D) | → | Root biomass | **-0.570** | **<0.001** |
| S×D | → | Root biomass | 0.264 | 0.301 |
| Plant species richness (S) | → | Fungal parasites | **-0.490** | **0.007** |
| Extreme drought (D) | → | Fungal parasites | -0.531 | 0.107 |
| S×D | → | Fungal parasites | 0.216 | 0.065 |
| Root biomass | → | Fungal parasites | 0.040 | 0.378 |
| Fungal parasites | → | Root biomass | 0.036 | 0.597 |
| Plant species richness (S) | → | Shoot biomass (feedback phase) | 0.120 | 0.611 |
| Extreme drought (D) | → | Shoot biomass (feedback phase) | *0.327* | *0.082* |
| S×D | → | Shoot biomass (feedback phase) | -0.142 | 0.620 |
| Plant species richness (S) | → | Root biomass (feedback phase) | 0.162 | 0.502 |
| Extreme drought (D) | → | Root biomass (feedback phase) | -0.008 | 0.965 |
| S×D | → | Root biomass (feedback phase) | -0.155 | 0.597 |
| Root biomass | → | Shoot biomass (feedback phase) | -0.068 | 0.423 |
| Fungal parasites | → | Shoot biomass (feedback phase) | **0.161** | **0.034** |
| Root biomass | → | Root biomass (feedback phase) | **0.178** | **0.049** |
| Fungal parasites | → | Root biomass (feedback phase) | 0.049 | 0.521 |

**Figure 4:** Results from path model analyses showing how soils conditioned by native plant diversity and extreme drought independently (lack of any interaction effect) affect the performance of (A) non-resident native invaders and (B) alien invasive plants *via* shifting root biomass (conditioning phase) and the abundance of soil fungal groups (conditioning phase). In the figure, thick lines (red: negative, gray: positive) represent statistically significant (p-values<0.05) relationships, whereas faded dashed lines represent statistically non-significant paths (p-value>0.05). The effect size (standardized path coefficients) and its statistical significance are tabulated below the figures. Bold numbers in the table are statistically significant, whereas italics numbers represent marginal significant effects (p-values less than 0.1 but greater than 0.05). Details of model structure are provided in the Material & Methods section, whereas model fit details are provided in the Result section. Path model for range-expanding plants with plant and fungal responses failed to have a good fit, and thus not shown here (Supplementary figure 4).