Borrelia burgdorferi PlzA is a cyclic-di-GMP dependent DNA- and RNA-binding protein

Nerina Jusufovic[1], Andrew C. Krusenstjerna[1], Christina R. Savage[1], Timothy C. Saylor[1],

Catherine A. Brissette[2], Wolfram R. Zückert[3], Paula J. Schlax[4], Md A. Motaleb[5], and Brian

Stevenson[1,6,*]

[1] Microbiology, Immunology and Molecular Genetics, University of Kentucky College of

Medicine, University of Kentucky, Lexington, Kentucky, 40526-0001, USA [2] Department of

Biomedical Sciences, University of North Dakota, School of Medicine and Health Sciences,

Grand Forks, ND 58203-9061, USA, [3] Department of Microbiology, Molecular Genetics and

Immunology, University of Kansas School of Medicine, Kansas City, KS 66160, USA, [4]

Department of Chemistry and Biochemistry, Bates College, Lewiston, ME, 04240-6030, USA,

[5] Department of Microbiology and Immunology, Brody School of Medicine, East Carolina

University, Greenville, NC 27834-435, USA, and [6] Department of Entomology, University of

Kentucky, Lexington, Kentucky, 40526-0001, USA.

* To whom correspondence should be addressed. Tel: +1 859 257 9358; Fax: 859 257 8994;

Email: brian.stevenson@uky.edu

ABSTRACT

The PilZ domain-containing protein, PlzA, is the only cyclic di-GMP binding protein encoded

by all Lyme disease spirochetes. PlzA has been implicated in the regulation of many borrelial

processes, but the effector mechanism of PlzA was not previously known. Here we report that

PlzA can bind DNA and RNA, and that nucleic acid binding requires c-di-GMP. In studies with

the promoter and 5' UTR of the glycerol catabolism operon, *glpFKD*, the affinity of PlzA for

nucleic acids increased as concentrations of c-di-GMP were increased. A mutant PlzA that is

incapable of binding c-di-GMP did not bind to any tested nucleic acids. PlzA is a dual-domain

protein consisting of a PilZ-like N-terminal domain linked to a canonical C-terminal PilZ

domain. Dissection of the domains demonstrated that the separated N-terminal domain bound

nucleic acids independently of c-di-GMP. The C-terminal domain, which includes the c-di-

GMP binding motifs, did not bind nucleic acids under any tested conditions. Our data are

supported by computational docking, which predicts that c-di-GMP binding at the C-terminal

domain stabilizes the overall protein structure and facilitates PlzA-DNA interactions via

residues in the N-terminal domain. Based on our data, we propose that levels of c-di-GMP

during the various stages of the enzootic life cycle dictate PlzA regulatory targets and

functions.

INTRODUCTION

35 Cyclic bis-(3'→5')-dimeric guanosine monophosphate (c-di-GMP), is a ubiquitous bacterial

36 cyclic di-nucleotide. In many bacterial species, it engages in the regulation of key processes

37 such as transcription, motility, biofilm formation, and virulence (Ryjenkov *et al.*, 2005; Hengge,

38 2009; Romling *et al.*, 2013; Jenal *et al.*, 2017; Valentini and Filloux, 2019). The Lyme disease

39 spirochete, *Borrelia burgdorferi* sensu lato, referred to as *Borrelia burgdorferi* henceforth, has

40 a c-di-GMP regulatory network consisting of a transmembrane sensor histidine kinase (Hk1),

41 a response regulator diguanylate cyclase (Rrp1), two phosphodiesterases (PdeA and PdeB),

42 and a single, chromosomally encoded PilZ domain-containing protein, PlzA (Galperin *et al.*,

43 2001; Freedman *et al.*, 2009; Rogers *et al.*, 2009; Sultan *et al.*, 2010; Caimano *et al.*, 2011;

44 Sultan *et al.*, 2011; Novak *et al.*, 2014).

45 The borrelial Hk1/Rrp1 two-component system synthesizes c-di-GMP; activation of Hk1

46 (BB_0420) occurs through an unknown signal (Caimano *et al.*, 2011; Bauer *et al.*, 2015). Rrp1

47 (BB_0419), the only identified diguanylate cyclase in *B. burgdorferi,* synthesizes c-di-GMP

48 from two GTP molecules and is essential for acquisition and survival in the tick vector (Rogers

49 *et al.*, 2009; Kostick *et al.*, 2011). Turnover of c-di-GMP is mediated by PdeA (BB_0363) and

50 PdeB (BB_0374), the only phosphodiesterases present in *B. burgdorferi*, which degrade c-di-

51 GMP to pGpG and GMP, respectively (Sultan *et al.*, 2010; Sultan *et al.*, 2011). *B. burgdorferi*

52 must delicately balance c-di-GMP signaling, as c-di-GMP is required for both maintenance in

53 the tick vector and successful transmission into the vertebrate host, but constitutive synthesis

54 is detrimental to spirochete survival during vertebrate infection (Sultan *et al.*, 2010; Sultan *et*

55 *al.*, 2011; Caimano *et al.*, 2015; Groshong *et al.*, 2021).

56 Synthesized c-di-GMP is bound by PlzA, the only universal c-di-GMP binding protein found in

57 all isolates of Lyme disease *Borreliae* (Kostick-Dunn *et al.*, 2018). Mutation of *plzA* has been

58 implicated in many *B. burgdorferi* phenotypes, including mis-regulation of alternative

59 carbohydrate utilization, decreased motility, and virulence defects (Pitzer *et al.*, 2011; He *et*

60 *al.*, 2014; Mallory *et al.*, 2016; Zhang *et al.*, 2018; Groshong *et al.*, 2021). Binding of c-di-GMP

induces conformational changes in PlzA structure, which has led to the hypothesis that c-di-

GMP binding acts as a switching mechanism providing the holo and apo-forms of PlzA with

distinct functions (Mallory *et al.*, 2016; Groshong *et al.*, 2021; Grassmann *et al.*, 2023).

Specifically, c-di-GMP bound PlzA is thought to regulate genes required for survival in the tick

host, while apo-PlzA is speculated to play a role in regulation during vertebrate infection

(Groshong *et al.*, 2021; Grassmann *et al.*, 2023).

A common mechanism of action for c-di-GMP effector proteins is to bind nucleic acids (Wilksch

et al., 2011; Tschowri *et al.*, 2014; Tan *et al.*, 2015; Wang *et al.*, 2016; Schäper *et al.*, 2017;

Hsieh *et al.*, 2018; Skotnicka *et al.*, 2020). It has been hypothesized that PlzA could bind DNA

and regulate genes through impacts on transcription, such as directly interacting with RNA

polymerase (RNAP), but this has yet to be experimentally tested (Groshong *et al.*, 2021;

Grassmann *et al.*, 2023). Conversely, others have hypothesized that PlzA cannot bind DNA,

as it does not contain an obvious, conventional DNA binding motif (Caimano *et al.*, 2015;

Zhang *et al.*, 2018). Indeed, Zhang *et al.* concluded that PlzA does not bind DNA, as their

electrophoretic mobility shift assays (EMSA) did not identify binding (Zhang *et al.*, 2018). The

results and methodologies including protein and c-di-GMP concentrations, were not shown in

that report. After the submission of a preprint of our manuscript, another group published that

PlzA has RNA chaperone activity that is independent of c-di-GMP, supporting the notion that

holo- and apo-forms of PlzA have distinct functions (Van Gundy *et al.*, 2023).

Given the multiple, conflicting hypotheses regarding PlzA function, as well as the potential for

additional mechanisms between liganded and unliganded PlzA, we sought to further

characterize PlzA nucleic acid binding function. We decided to examine a known regulatory

target of PlzA, *glpFKD*, the glycerol catabolism operon. The *glpFKD* operon promotes *B.*

burgdorferi survival in ticks (Pappas *et al.*, 2011; Corona and Schwartz, 2015; Helble *et al.*,

2021). After a tick has completed the digestion of its blood meal, it is hypothesized that the

midgut is devoid of glucose and many other nutrients. To survive in that environment, Lyme

disease spirochetes would need to utilize alternate carbon sources such as glycerol, which is

available in the tick host (Lee and Baust, 1987). The *glpFKD* operon is regulated through the

Hk1/Rrp1 pathway, and borrelial *plzA* mutants are defective in glycerol metabolism (Rogers

et al., 2009; He *et al.*, 2011; Caimano *et al.*, 2015; Zhang *et al.*, 2018). Further, PlzA was found

to both positively and negatively affect the *glp* operon, at higher and lower c-di-GMP levels,

respectively (Zhang *et al.*, 2018).

PlzA is classified as an xPilZ-PilZ domain protein given its dual-domain structure. The binding

of c-di-GMP to the interdomain linker and C-terminal PilZ domain results in a conformational

change; locking the C-terminal PilZ and N-terminal xPilZ domains into rigid conformation,

which we hypothesized could permit DNA binding (Mallory *et al.*, 2016; Groshong *et al.*, 2021;

Singh *et al.*, 2021). While no canonical DNA binding domain is evident in the PlzA amino acid

sequence, our lab has discovered and characterized multiple nucleic acid binding proteins

possessing novel binding motifs (Babb *et al.*, 2006; Riley *et al.*, 2009; Jutras *et al.*, 2012;

Jutras *et al.*, 2013a; Jutras *et al.*, 2013b; Zhang *et al.*, 2018; Singh *et al.*, 2021). Here, we

report that PlzA can bind DNA and RNA directly downstream of the promoter and into the 5'

untranslated region (UTR) of the *glpFKD* operon and that binding to this region is c-di-GMP

dependent.

RESULTS

PlzA binds *glpFKD* DNA and the interaction is c-di-GMP dependent

Previous studies demonstrated that c-di-GMP and PlzA modulate expression of the glycerol

metabolism operon (Rogers *et al.*, 2009; He *et al.*, 2011; Caimano *et al.*, 2015; Zhang *et al.*,

2018). Activation of *glpFKD* is critical for spirochete survival in between blood meals of the tick

vector, which requires catabolism of glycerol (He *et al.*, 2011; Pappas *et al.*, 2011; Zhang *et*

al., 2018). Given that requirement, we hypothesized that PlzA might bind 5' of the *glpFKD*

operon. To that end, we sought to determine if PlzA could bind DNA between *glpFKD* and the

upstream ORF BB_0239, which encodes a deoxyguanosine/deoxyadenosine kinase. The

transcriptional boundaries and promoter elements of the *glpFKD* operon were previously

mapped (Adams *et al.*, 2017; Grove *et al.*, 2017). A fluorescently labeled DNA fragment was

generated by PCR amplification from a plasmid containing the 410 bp intergenic region

between the upstream gene, and the start of *glpF* (BB_0240) and is denoted as *glpFKD*(-219)

(Figure 1).

To assess if PlzA could bind this region of DNA, EMSAs with increasing concentrations of

recombinant PlzA were performed (Figure 2A). Binding was observed at higher micromolar

concentrations, as evidenced by the disappearance of free DNA. The experiment was

repeated with added non-specific competitor poly-dI-dC to a final concentration of 2.5 ng/μL.

Poly-dI-dC did not appreciably compete away the shifted protein-DNA complexes (Figure 2B).

Given that binding of c-di-GMP has been hypothesized to act as a switching mechanism for

PlzA function, we next assessed the effects of c-di-GMP on DNA binding. EMSA reaction

mixtures were supplemented with exogenous c-di-GMP to a final concentration of 100 μM.

This concentration was used to saturate all PlzA molecules present in the reaction to ensure

complete binding throughout electrophoresis. Saturating concentrations have similarly been

used in other studies with c-di-GMP binding proteins that also bind DNA (Wilksch *et al.*, 2011;

Yang *et al.*, 2013; Tan *et al.*, 2015; Zamorano-Sánchez *et al.*, 2015; Schäper *et al.*, 2017;

Skotnicka *et al.*, 2020). The EMSA with *glpFKD*(-219), poly-dI-dC, and increasing PlzA was

repeated with the supplemented c-di-GMP (Figure 2C). The addition of c-di-GMP enhanced

PlzA binding affinity for *glpFKD*(-219), resulting in binding at lower concentrations of protein,

and the formation of distinct protein-DNA complexes as indicated by the arrows. Further, as

PlzA concentration increased, the mobility of the shifted band decreased.

To further assess the requirement of c-di-GMP for PlzA DNA binding activity, a double mutant

PlzA protein (PlzA$_{RD-RD}$) that cannot bind c-di-GMP was produced by site-directed

mutagenesis which replaced key arginine residues in the conserved c-di-GMP binding motif,

RxxxR, with aspartic acids (Ryjenkov *et al.*, 2006; Amikam and Galperin, 2006; Chou and

Galperin, 2016). These mutations resulted in the amino-acid substitutions R145D and R149D

within the c-di-GMP binding site of PlzA and have previously been shown to abolish c-di-GMP

binding (Freedman *et al.*, 2009; Mallory *et al.*, 2016; Singh *et al.*, 2021). The *glpFKD*(-219)

EMSAs were repeated with PlzA$_{RD-RD}$ (Figure 2D-F). Binding to *glpFKD*(-219) was not

observed under any tested conditions with PlzA$_{RD-RD}$.

To ensure binding was not due to recognition of vector derived sequences within the probe,

competition EMSAs were performed with unlabeled DNA competitors derived from empty pCR

2.1 TA clones, or those containing the *glpFKD*(-219) sequence (Figure 3A). The nonspecific

DNA from the empty pCR 2.1 vector did not compete away PlzA$_{WT}$-*glpFKD*(-219) complexes

at a 20-fold higher concentration (Figure 3B, lanes 3-5). Meanwhile, as little as 6-fold molar

excess of unlabeled *glpFKD*(-219) was able to compete with the probe, and a 10-fold molar

excess successfully competed away the PlzA$_{WT}$-*glpFKD*(-219) complexes (Figure 3B, lanes

6-8). These results demonstrate PlzA preferentially bound the *glpFKD*-derived sequences.

Noting that c-di-GMP enhanced DNA binding by PlzA yet free DNA decreased with purified

recombinant PlzA without adding exogenous c-di-GMP, we investigated the possibility that

PlzA was being purified with c-di-GMP already bound. Purified recombinant proteins were

prepared and assayed by liquid chromatography-tandem mass spectrometry. Extracted ion

chromatograms showed the presence of c-di-GMP in PlzA$_{WT}$ protein preparations but not in

PlzA$_{RD-RD}$ preparations (Figure 4). The concentration of c-di-GMP present in recombinant

PlzA$_{WT}$ was extrapolated from a c-di-GMP standard curve at 99.29 nM. A total of 31.7 µM of

protein had been analyzed, indicating that approximately 0.3% of the protein was bound with

c-di-GMP. Prior studies have reported dissociation constants of 1.25 to 6.25 µM of PlzA for c-

di-GMP, which are typical K_D values for PilZ receptors and c-di-GMP (Pitzer *et al.*, 2011;

Mallory *et al.*, 2016). The amount of co-purified c-di-GMP detected is lower than the K_D. This

is likely due to removal of some c-di-GMP during the protein purification process, as well as

E. coli maintaining low cellular concentrations of c-di-GMP (~40-80 nM) (Sarenko *et al.*, 2017;

Junkermeier and Hengge, 2023). Taken together, these data indicate that PlzA DNA binding

to this region of the *glpFKD* operon is specific and c-di-GMP dependent.

167   **c-di-GMP levels modulate PlzA DNA binding affinity**

168   It was previously identified with *glpFKD-gfp* promoter fusion studies that the minimal *glpFKD*

169   promoter sequence (-46 relative to the *glpFKD* transcription start site) with the full 195bp UTR

170   region was expressed at higher levels than the core promoter sequence alone (Grove *et al.*,

171   2017). Given these findings and our data that PlzA can bind DNA, we focused further analysis

172   of PlzA binding on a smaller sequence within that region that was found to be important for

173   transcription (Grove *et al.*, 2017; Savage *et al.*, 2018). A 42 bp DNA probe encompassing the

174   -7 to +35 sites, called *glpFKD*(-7/+35), was generated by annealing complementary

175   oligonucleotides with the forward oligonucleotide 5' conjugated to IRDye800 (Figure 5A). An

176   RNA probe conjugated to Alexa Fluor 488 was also made for subsequent RNA binding studies.

177   To determine how different c-di-GMP levels impact PlzA binding affinity, EMSAs with

178   increasing PlzA$_{WT}$ concentrations were performed at concentrations of added c-di-GMP

179   ranging from 0 to 100 µM in the reaction mixtures with the *glpFKD*(-7/+35) probe (Figure 5B).

180   Competitor poly-dI-dC was added at a final concentration of 2.5 ng/µL to reduce nonspecific

181   interactions. PlzA$_{WT}$ bound to *glpFKD*(-7/+35), and binding was enhanced by increasing the

182   concentration of c-di-GMP. In reactions with no added c-di-GMP, we observed that the amount

183   of free DNA decreased with increasing PlzA concentrations but did not detect a band

184   corresponding to the protein-DNA complex. The decrease in free DNA with no added c-di-

185   GMP is presumably due to the presence of co-purified c-di-GMP in recombinant PlzA. When

186   equimolar c-di-GMP and protein levels were tested, a discernable shift (formation of a new

187   band) was evident only at protein concentrations greater than 2.5 µM (purple arrow). As c-di-

188   GMP concentrations were further increased, shifts became discernable at lower protein

189   concentrations, with the strongest binding observed at c-di-GMP concentrations of 25, 50, and

190   100 µM. Further, as protein concentration was increased, we consistently observed

191   progressive super-shifting of complexes in a dose-dependent manner (Figure 5B, colored

192   arrows). At protein concentrations greater than or equal to 5 µM coupled with c-di-GMP

193   concentrations greater than 25 µM, complexes appeared to smear, suggesting the observed

super-shifting is due to complex dissociation. The stoichiometry of this complex(es) is under

investigation.

Next, we calculated the apparent binding affinity ($K_{D(app)}$) under these experimental conditions

for PlzA for *glpFKD*(-7/+35) at the various c-di-GMP concentrations (Figure 5C). The

calculations were performed under the assumption that a single molecule of PlzA binds to a

single molecule of DNA. As noted above, it is possible that the protein-nucleic acid ratio is not

1:1, which could affect the calculated values. The values described below are the $K_{D(app)}$ ±

SEM (standard error of the mean). Without any supplemented c-di-GMP, the extrapolated

$K_{D(app)}$ of PlzA$_{WT}$-*glpFKD*(-7/+35) was calculated as 7.86 (± 1.84 µM), however we again note

that some c-di-GMP is already present in the PlzA$_{WT}$ protein preparations (see above and

Figure 4). Binding affinity increased approximately 1.5-fold under equimolar c-di-GMP and

protein concentrations with a calculated $K_{D(app)}$ of 4.22 (± 0.43 µM). Under saturating c-di-GMP

conditions (exceeding the K_D of PlzA for c-di-GMP), binding affinity increased further.

Supplementing binding reactions with 10 µM c-di-GMP improved PlzA-DNA binding 2.5-fold

($K_{D(app)}$ of 2.81 (± 0.50 µM). An approximately 6- to 7-fold increase in binding affinity was

observed when the binding reaction was supplemented with either 25, 50, or 100 µM of c-di-

GMP resulting in calculated $K_{D(app)s}$ of 1.33 (± 0.12 µM), 1.25 (± 0.08 µM), and 1.09 µM (± 0.06

µM), respectively.

To determine if the calculated $K_{D(app)}$ values differed significantly, Welch's ANOVA followed by

a Dunnett's T3 multiple comparisons test was performed (Figure 5D). All calculated $K_{D(app)}$

values differed significantly ($p \leq 0.05$) from the 0 µM c-di-GMP $K_{D(app)}$ except for the equimolar

and 10 µM c-di-GMP conditions. Compared to the equimolar condition, the 25, 50, and 100

µM $K_{D(app)}$ values differed significantly ($p \leq 0.001$), while the 10 µM c-di-GMP $K_{D(app)}$ value did

not ($p > 0.05$). When comparing the 10, 25, 50, and 100 µM $K_{D(app)}$ values of PlzA$_{WT}$-*glpFKD*(-

7/+35) interaction, there were no significant differences ($p > 0.05$). The mutant PlzA$_{RD-RD}$ did

not bind *glpFKD*(-7/+35) under any tested conditions (Figure 5E). Overall, we conclude that

levels of c-di-GMP significantly impact PlzA-DNA binding.

PlzA also binds *glp*-derived single-stranded nucleic acids

Several borrelial proteins that were initially identified as having affinities for DNA have

subsequently been found to also bind RNA (Jutras *et al.*, 2013a; Savage *et al.*, 2018;

Krusenstjerna *et al.*, 2023). Given the role the 5' UTR of the *glpFKD* operon plays in its

regulation, we hypothesized PlzA could potentially also bind RNA. RNA probes used were

designed to correspond to only transcribed regions (Figure 5A). The *glpFKD*(-7/+35)

counterpart RNA probe is designated *glpFKD*(UTR). EMSAs were performed with increasing

concentrations of PlzA$_{WT}$ with 0 µM, 100 µM, or equimolar c-di-GMP levels and labeled

glpFKD(UTR) (Figure 6A). We did not test additional c-di-GMP concentrations that were tested

for impacts on DNA binding for the RNA, as no significant difference was found between 100

µM c-di-GMP and the other tested c-di-GMP concentrations. PlzA$_{WT}$ bound the RNA, which

was enhanced by addition of c-di-GMP (Figure 6A). The binding pattern to the *glpFKD*(UTR)

RNA was comparable to the *glpFKD*(-7/+35) DNA, with increased protein and c-di-GMP

concentrations leading to an increase in the size of the PlzA-RNA shifts.

EMSAs were repeated with the PlzA$_{RD-RD}$ mutant protein at 0 or 100 µM c-di-GMP (Figure 6B).

The PlzA$_{RD-RD}$ protein did not bind *glpFKD*(UTR) RNA at any tested protein concentration,

regardless of whether c-di-GMP was included.

Given that PlzA can bind to both the DNA and RNA, we sought to determine if it could also

bind the cognate ssDNA corresponding to the coding strand. This is relevant as ssDNA is

exposed at open transcriptional complexes. The forward, IRDye labeled oligonucleotide used

to generate the *glpFKD*(+7/-35) dsDNA probe was used as a single-stranded DNA probe

(ss*glpF*) in EMSAs. No evident complexes were formed when exogenous c-di-GMP was

excluded from EMSA reactions, but free ssDNA did decrease at the highest tested PlzA

concentrations of 3.5-7.5 µM (Figure 6C right panel). Adding 100 µM c-di-GMP resulted in the

formation of PlzA-ssDNA complexes but at higher concentrations of protein than required to

achieve complex formation with dsDNA and RNA (Figure 6C left panel).

247 We next quantified the binding affinity of PlzA for the *glp* derived RNA and ssDNA sequences

248 through densitometry of the EMSA shifts, and the apparent $K_{D(app)s}$ calculated as described

249 above (Figure 6D). As with dsDNA, the binding affinity of PlzA for RNA was weakest when no

250 exogenous c-di-GMP was supplemented with a calculated $K_{D(app)}$ of 7.64 (± 1.85 µM). Binding

251 improved approximately 2-fold in equimolar c-di-GMP conditions with a $K_{D(app)}$ of 4.14 (± 0.74

252 µM). The $K_{D(app)}$ was 0.95 (± 0.11 µM) when the c-di-GMP concentration was 100 µM, an 8-

253 fold increase in PlzA binding affinity for the *glpFKD*(UTR) RNA as compared to when no c-di-

254 GMP was added. There was a significant difference between the PlzA$_{WT}$-*glpFKD*(UTR) RNA

255 $K_{D(app)}$ value calculated at 100 µM c-di-GMP and both the equimolar and 0 µM c-di-GMP values

256 (p ≤0.01) (Figure 7A). No significant differences were detected between the $K_{D(app)}$ values

257 calculated at 0 µM and equimolar c-di-GMP levels (p >0.05).

258 Although PlzA bound the ss*glpF* probe, it had a substantially lower affinity for ssDNA than

259 dsDNA or RNA (Figure 6D). The $K_{D(app)}$ of the PlzA$_{WT}$-ss*glpF* interaction was 8.74 (± 1.86 µM)

260 with 0 µM c-di-GMP and 4.42 (± 0.90 µM) with 100 µM c-di-GMP. These values were found to

261 be statistically significant from one another (p <0.05) (Figure 7B). Only a 2-fold increase in

262 binding affinity of PlzA for the single-strand DNA was observed when the c-di-GMP

263 concentration was increased to 100 µM, whereas the same concentration of c-di-GMP

264 increased PlzA binding affinity 7-fold for dsDNA and 8-fold for RNA.

265 Lastly, we also compared the binding affinities of PlzA for all three tested nucleic acids (Figure

266 7C). The calculated $K_{D(app)}$ values of PlzA$_{WT}$ for each nucleic acid in the presence of 100 µM

267 c-di-GMP were compared with a Welch's ANOVA and Dunnett's T3 multiple comparisons test.

268 No significant difference (p >0.05) was detected between the PlzA $K_{D(app)}$ values for dsDNA

269 (1.09 ± 0.06 µM) and RNA (0.95 ± 0.11 µM), indicating that PlzA binds the *glpFKD* dsDNA

270 and RNA with similar affinity. Both the dsDNA and RNA $K_{D(app)s}$ were significantly different from

271 the calculated ssDNA $K_{D(app)}$ (4.42 ± 0.90 µM) (p ≤0.01), suggesting that PlzA prefers *glp*-

272 derived dsDNA and RNA over ssDNA. All the binding data presented throughout the

manuscript are summarized in Table 1. Collectively, our data indicate that PlzA binding activity

is c-di-GMP dependent for all tested *glpFKD* nucleic acids.

Dissection of PlzA reveals the nucleic acid binding domain

The first discovered c-di-GMP binding receptors were the PilZ domain-containing proteins

(Ryjenkov *et al.*, 2006; Amikam and Galperin, 2006; Galperin and Chou, 2020). The PilZ

domain consists of two conserved c-di-GMP binding motifs, RxxxR and [D/N]ZSXXG

(Ryjenkov *et al.*, 2006; Amikam and Galperin, 2006; Chou and Galperin, 2016; Cheang *et al.*,

2019). The recently solved crystal structure of *B. burgdorferi* PlzA revealed a unique dual-

domain topology consisting of an amino-terminal PilZ-like domain, called PilZN3, which is

connected to the carboxy-terminal PilZ domain via a linker domain that contains the RxxxR c-

di-GMP binding motif (Singh *et al.*, 2021). The first 141 amino acid residues make up the N-

terminus of PlzA (PlzA$_{NTD}$), while residues 142-261 make up the C-terminal (PlzA$_{CTD}$) PilZ

domain. Having determined that PlzA is a c-di-GMP-dependent DNA and RNA binding protein,

we next sought to determine which domain of PlzA binds nucleic acid. To generate

recombinant PlzA$_{NTD}$ and PlzA$_{CTD}$ domains, truncated *plzA* genes called *plzA*-NTD and *plzA*-

CTD were cloned to encode the residues described above, which correspond to each

respective domain (Figure 8A).

EMSAs were performed with labeled *glpFKD*(-7/+35) DNA probe and increasing

concentrations of PlzA$_{NTD}$ or PlzA$_{CTD,}$ with or without 100 µM c-di-GMP. PlzA$_{NTD}$ bound DNA

independently of c-di-GMP, as supplementation with c-di-GMP did not affect binding (Figure

8B). PlzA$_{CTD}$ contains the motifs required for c-di-GMP binding, but the PlzA$_{CTD}$ domain did not

bind to DNA with or without added c-di-GMP (Figure 8C). It was also determined that PlzA$_{NTD}$

could bind *glpFKD*(UTR) RNA with or without c-di-GMP, while PlzA$_{CTD}$ does not (Figure 8D

and E respectively).

We also performed the LC/MS-MS analysis on PlzA$_{NTD}$ and PlzA$_{CTD}$ recombinant proteins to

determine if any di-nucleotides were present (Figure 9A). Co-purified c-di-GMP was detected

in PlzA$_{CTD}$ but not PlzA$_{NTD}$ by LC/MS-MS. The concentration of c-di-GMP present in

recombinant PlzA$_{CTD}$ was determined as 704.29 nM. A total of 66.7 µM of protein was sent for

analysis indicating approximately 1.1 % of PlzA$_{CTD}$ was bound with c-di-GMP. This indicates

that PlzA$_{CTD}$ retained the ability to bind c-di-GMP but cannot bind nucleic acid.

To confirm that the truncated proteins were stable, circular dichroism (CD) was performed,

and the CD spectra were obtained for PlzA$_{WT}$, PlzA$_{RD-RD}$, PlzA$_{NTD}$, and PlzA$_{CTD}$ in the binding

buffer used for EMSAs (Figure 9B). The CD results show that the mutated proteins retained

stable structures matching that of PlzA$_{WT}$. Therefore, the lack of DNA binding by PlzA$_{RD-RD}$ and

PlzA$_{CTD}$ is not due to improper folding.

Computational docking of PlzA-nucleic acid interactions

To gain a better understanding of the interactions between PlzA and DNA or RNA, we

performed a computational docking analysis with the HDOCK server (Huang and Zou, 2014;

Yan *et al.*, 2017a; Yan *et al.*, 2017b; Yan *et al.*, 2020). The *glpFKD*(-7/+35) DNA and

glpFKD(UTR) RNA sequences were input as ligands, while the PlzA crystal structure (PDB

ID: 7mie) was input as the receptor molecule. The modeling results indicate that interactions

between PlzA and *glpFKD*(-7/+35) DNA are facilitated primarily through the N-terminal domain

(Figure 10a). Beta-strands six and seven of the N-terminus are extended and protrude

outward, intercalating a major groove of the DNA 5' to the protein docking site (Figure 10a top

panel). Polar and positively charged amino acids, including N110 and K111, are located at the

loop region where beta-strands 6 and 7 converge, with the potential to directly interact with

bases of the DNA (Rooman *et al.*, 2002; Sathyapriya and Vishveshwara, 2004). Interactions

with a minor groove of the DNA are predicted to occur through the loop region of beta-strands

2 and 3 and an unstructured linker region (Figure 10a bottom panel). Several positively

charged residues, including arginine (R32 and R38) and lysine (K81), as well as an aromatic

phenylalanine residue (F45), are predicted to directly interact with the minor groove and

backbone of the DNA. An additional lysine residue K87 located in the unstructured region,

protrudes and interacts directly with bases of another major groove of *glpFKD*(-7/+35). No

residues of the C-terminal domain are predicted to interact directly with the DNA; however,

four residues are interfacing the DNA (K161, I213, D258, and N261). Residue K161 hovers

over the DNA backbone potentially interacting nonspecifically via electrostatic interactions.

The computational docking data supports our EMSA data showing that PlzA$_{NTD}$ binds DNA

while PlzA$_{CTD}$ does not (Figures 8B and C).

The docking analysis was also performed with PlzA$_{WT}$ and *glpFKD*(UTR) RNA (Figure10B).

Again, the N-terminal domain of PlzA facilitates most of the interactions. Similar regions of the

protein that interacted with dsDNA are implicated in the interaction with RNA including the

residues of the loop region of beta-strands 6 and 7 (K111 and Y113), the loop region of beta-

strands 2 and 3 (R38, Y42, and F45) and the unstructured linker region (R32, D85, K87, F92,

and F95). Unlike the docking model with dsDNA, the PlzA-RNA docking model indicates more

involvement of the C-terminal domain in RNA binding. Several residues of alpha-helixes 5

(D155, R158, K159, and K161) and 6 (D258 and Y259) in the C-terminal domain of PlzA, are

predicted to interact with RNA. Although our EMSA results show that PlzA$_{CTD}$ does not bind

any tested nucleic acids, the docking results indicate it could help stabilize PlzA$_{NTD}$-nucleic

acid interactions or could be involved in target recognition. Several PlzA residues predicted to

be involved in nucleic-acid binding are currently being investigated by our lab.

Computational docking was attempted with PlzA and ss*glpF*, but the HDOCK server could not

complete the analysis. Given that PlzA bound dsDNA and RNA better than ssDNA, it is

interesting to note that the computational modeling predicts the RNA to fold into a double

stranded secondary structure. Combined with our EMSA data, this suggests that PlzA could

have a preference towards double stranded nucleic acids and/or secondary structures.

DISCUSSION

Cyclic-di-GMP has wide-ranging effects on many bacterial processes. The responses to c-di-

GMP are commonly mediated through c-di-GMP binding effector proteins. PlzA is the only

chromosomally and universally encoded c-di-GMP binding protein of the Lyme disease

spirochetes. It has been shown to be crucial for borrelial infection processes and in the

regulation of the metabolism of alternative carbon sources during colonization of vector ticks

(Pappas *et al.*, 2011; Pitzer *et al.*, 2011; He *et al.*, 2014; Mallory *et al.*, 2016; Zhang *et al.*,

2018; Groshong *et al.*, 2021). Despite its singularity and importance to *B. burgdorferi*, little was

previously known about the biochemical functions of PlzA outside of binding c-di-GMP. Here,

we provide evidence that PlzA is a novel c-di-GMP-dependent nucleic acid-binding protein.

While PlzA lacks an obvious canonical DNA binding motif, we hypothesized that PlzA might

be a nucleic acid binding protein given its role in the regulation of the *glpFKD* operon. Binding

was only observed by the wild-type PlzA protein but not the mutant, PlzA$_{RD-RD}$, which is

incapable of binding c-di-GMP. It has previously been shown that structural rearrangements

are induced upon the binding of c-di-GMP to PlzA (Mallory *et al.*, 2016; Groshong *et al.*, 2021).

Without bound c-di-GMP, PlzA has a structure that is flexible, potentially preventing the

interactions necessary for DNA binding. This hypothesis is supported by the fact that

crystallization of PlzA was only achieved with c-di-GMP bound (Singh *et al.*, 2021). We

surmise that binding of c-di-GMP to the C-terminal PilZ domain results in a conformational

change of PlzA into a more rigid structure permitting nucleic acid binding through residues of

the N-terminal domain.

We identified the N-terminal domain of PlzA as the nucleic acid binding domain and that

PlzA$_{NTD}$ bound DNA independently of c-di-GMP. In contrast, the C-terminal domain, PlzA$_{CTD}$,

did not bind any tested nucleic acids although it still bound c-di-GMP. While PlzA$_{NTD}$ can

interact with nucleic acids, these interactions are not as strong as PlzA$_{WT}$. Binding of nucleic

acid by PlzA$_{NTD}$ resulted in smeared shifts that lacked defined bands. This suggests instability

of the PlzA$_{NTD}$-nucleic acid complexes resulting in dissociation of the complexes during

electrophoresis. Noting that binding of c-di-GMP appears to stiffen interactions between the

N- and C-terminal domains, we hypothesize the C-terminal domain stabilizes the interactions

of PlzA$_{NTD}$ with nucleic acids when c-di-GMP bound. On the other hand, we posit that the

flexibility of apo-PlzA results in the C-terminal domain disrupting the ability of the N-terminal

domain to bind DNA. Although the N- and C- terminal PlzA domains are similar, structural

comparisons of $PlzA_{NTD}$ and $PlzA_{CTD}$ reveal distinct differences (Figure 8). The N-terminus of

$PlzA_{NTD}$ contains an alpha-helix that is absent in the N-terminus of $PlzA_{CTD}$. Additionally, an

alpha-helix is present at the beta-barrel pore of $PlzA_{NTD}$ but not in the $PlzA_{CTD}$. These

differences suggest distinct roles in PlzA-nucleic acid binding dynamics. We are currently

exploring the sequence motifs recognized by PlzA and the exact amino acid residues involved

in nucleic acid binding.

Previous studies found that PlzA is a monomer in solution regardless of c-di-GMP (Mallory *et*

al., 2016). While many DNA-binding proteins bind DNA as dimers, alternative binding models

exist (Kim and Little, 1992; Kohler *et al.*, 1999). It has been shown that some proteins are

capable of binding sequentially as monomers which then multimerize on the DNA via protein-

protein interactions, such as some members of the Leucine zipper and helix-loop-helix zipper

families, the LexA repressor of *E. coli*, and BpaB of *B. burgdorferi* (Kim and Little, 1992; Kohler

et al., 1999; Burns *et al.*, 2010). Our studies revealed that increasing c-di-GMP and $PlzA_{WT}$

concentrations resulted in progressively slower migration of PlzA-*glpFKD*(-7/+35) DNA

complexes. The decreases in the migration of the complexes that are observed with increasing

protein concentration may indicate a stoichiometry that is not 1:1, or the observed smearing

may be the result of dynamic equilibrium of the complexes in the gels during electrophoresis

(Fried and Crothers, 1981; Fried, 1989). While increasing c-di-GMP enhances PlzA binding,

there could be a threshold where too much c-di-GMP could be inhibitory and cause

dissociation of PlzA from nucleic acids. A dimerization or protein-protein interaction

residue/motif/domain remains to be identified in PlzA, and work is ongoing to address the

DNA-binding mode and stoichiometry of PlzA-nucleic acid complexes.

We also observed probe signal in the gel wells at high protein and c-di-GMP concentrations.

This could be due to protein aggregation. Rather than simply being a negative consequence

of environmental stress, recent studies indicate a role for protein aggregation in regulation.

Specifically, certain conditions can induce a protein to aggregate, and thus be sequestered,

altering gene expression to permit bacterial adaptation to a specific environment (Schramm *et*

al., 2019). Whether high levels of c-di-GMP promote such processes remains to be tested, but

c-di-GMP levels certainly alter PlzA binding affinity.

Our study sought to identify if PlzA can bind nucleic acids and what role c-di-GMP plays in

that function. The data is conclusive that PlzA indeed binds DNA, both double and single-

stranded, and RNA, with a preference for the dsDNA and RNA. PlzA also displayed higher

affinity for certain nucleic acid types over others, and PlzA-*glp* DNA binding was competed

away by specific competitor but not nonspecific competitor. Further, the nucleic acid binding

at the *glpFKD* locus by PlzA is dependent on c-di-GMP. A caveat of the data is that the

calculated dissociation constants for PlzA-nucleic acids are lower than typically observed for

protein-nucleic acid interactions. The higher dissociation constants could be attributed to our

method of choice, EMSA, as some potential dissociation and protein aggregation were

observed. Attempts were made with various alterations aimed to improve the stability of the

complexes during electrophoresis, such as altering the gel percentage, pH of the running

buffer, salt/ion concentrations of the reaction buffer, and addition of detergents to the reaction

(Hellman and Fried, 2007). Altering the pH of the running buffer to 8.0 decreased the

aggregation in the wells but smearing still occurred at higher protein and c-di-GMP

concentrations (data not shown). The experimental conditions presented and reported here

yielded the most consistent results.

PlzA bound to a sequence adjacent to the promoter and extending 35 bp into the 5' UTR of

glpF. This site has been shown to positively affect *glpFKD* transcript levels and is a site where

two other borrelial regulators, SpoVG and BadR, also bind (Grove *et al.*, 2017; Savage *et al.*,

2018; Zhang *et al.*, 2022; Saylor *et al.*, 2023). BadR, the <u>B</u>orrelia Host <u>Ad</u>aption <u>R</u>egulator,

has recently been identified as a repressor of *glpFKD*, while PlzA has previously been shown

to exert both positive and negative effects on *glpFKD*, at high and low levels of c-di-GMP

respectively (Zhang *et al.*, 2018; Zhang *et al.*, 2022). Although SpoVG can bind the DNA and

RNA of *glpFKD*, the transcriptional consequences of these interactions are unknown (Savage

et al., 2018; Saylor *et al.*, 2023). How PlzA binding affects *glpFKD* expression, and how these

proteins work in tandem to regulate glycerol catabolism, is currently being investigated. Multi-

layered regulation of glycerol catabolism is found in other bacteria indicating this is not a

phenomenon unique to *Borrelia* (Bong *et al.*, 2019).

During the preparation of this manuscript, Van Gundy et al. published a report identifying RNA

chaperone activity by PlzA (Van Gundy *et al.*, 2023). The authors concluded that this activity

is independent of c-di-GMP. This greatly deviates from our results that c-di-GMP is required

for nucleic acid binding. This was acknowledged by those authors who suggested that

differences in methodologies and conditions might have contributed to the differences in

observed results. Some of these differences include their use of non-borrelial derived RNAs,

a different PlzA site-mutant, equimolar concentrations of c-di-GMP, and filter binding assays

used to calculate dissociation constants. Indeed, the differences in methodology and

conditions between the two studies make it difficult to draw comparisons between calculated

apparent K_D values, as they are only constant under the given conditions tested (Jarmoskaite

et al., 2020). However, some facets of the studies can still be compared. Their study utilized

short, unstructured, non-borrelial derived RNAs, while the nucleic acids used in our study are

borrelial derived, although examining just one locus PlzA is known to affect. Indeed, our data

indicates that PlzA preferentially binds double-stranded/structured nucleic acids. No condition

we tested yielded low nanomolar range binding. Regardless, the methodology yielded

reproducible data that allowed for the identification of PlzA c-di-GMP-dependent nucleic acid-

binding.

Given our finding that c-di-GMP levels greatly affect the affinity of PlzA for nucleic acids, we

tested binding to both *glpFKD* DNA and RNA under equimolar conditions like those used by

Van Gundy et al. In our hands, equimolar concentrations of c-di-GMP yielded significantly

reduced nucleic acid binding affinities than those exceeding the calculated K_D of PlzA for c-di-

GMP. Considering the higher K_D of the PlzA-c-di-GMP interaction (1.25-6.25 µM), under

equimolar conditions reaching the K_D of PlzA-c-di-GMP depends on the protein concentration

used. The protein concentrations used in the other study likely never reached a concentration

that exceeded the K_D of PlzA for c-di-GMP to saturate half of the PlzA molecules, except at

the higher protein ranges tested (1-2 μM). This could explain the observed weaker RNA

binding and/or chaperone activity with holo-PlzA.

We chose to evaluate saturating c-di-GMP concentrations to evaluate the maximal nucleic

acid binding activity of holo-PlzA and mitigate potential variability in the calculated apparent

K_D values. It was also never confirmed if their PlzA protein preparations co-purified with c-di-

GMP, which could contribute to observed binding in their apo-PlzA conditions. We also did not

observe binding under any tested conditions with the PlzA$_{RD-RD}$ mutant, which cannot bind c-

di-GMP. Lastly, the goals for our studies differed greatly as they sought to study RNA

chaperone activity, while we set out to investigate PlzA nucleic acid binding at the *glpFKD*

operon. Our two studies, however, do corroborate that PlzA can bind RNA and suggest that

this function plays a role in the regulation of borrelial gene expression. We agree that further

investigation is required to unravel the molecular mechanisms of PlzA's interactions with

nucleic acids, and the roles that those interactions play in gene regulation in this important

pathogen.

It is known that the affinity and specificity of proteins for target nucleic acids are impacted by

co-effectors (Carey, 2022). We observed that adding exogenous c-di-GMP increased binding

affinity of PlzA for *glp*-derived DNA and RNA. Studies in other bacteria with multiple PilZ

domain proteins have suggested that the differential affinity of the various PilZ receptors for c-

di-GMP sequentially regulates the processes that respond to c-di-GMP (Pultz *et al.*, 2012;

Junkermeier and Hengge, 2023). Given that PlzA is the only PilZ domain protein in most *B.*

burgdorferi isolates, an alternative to this could be that cellular levels of c-di-GMP alter the

affinity and/or specificity of PlzA. We propose a model in which c-di-GMP availability alters the

functionality of PlzA, and thus, the targets regulated by PlzA could depend on the c-di-GMP

levels available at different stages of the enzootic life cycle (Figure 11).

During vertebrate infection, no c-di-GMP is produced, favoring apo-PlzA functions such as

RNA chaperone activity (Van Gundy *et al.*, 2023). When c-di-GMP is present, holo-PlzA can

also anneal RNAs and potentially interact with RNAP (Grassmann *et al.*, 2023; Van Gundy *et*

al., 2023). At lower levels of c-di-GMP, PlzA nucleic-acid binding affinity is low for certain target

genes such as *glpFKD*. As c-di-GMP levels increase, such as in a tick or during the blood

meal, binding affinity for target nucleic acids is increased and potential multimerization of PlzA

molecules on DNA could occur. Lastly, we hypothesize that an unknown signal might regulate

PlzA activity at high concentrations resulting in the release of holo-PlzA from target DNAs.

This hypothesis requires further investigation, as does overall regulation of c-di-GMP in *B.*

burgdorferi.

In conclusion, PlzA is a c-di-GMP dependent nucleic acid binding protein. While other c-di-

GMP binding receptors have been identified as DNA-binding proteins, PlzA is the first that has

been demonstrated to also bind RNA, thus expanding the functional diversity of c-di-GMP

binding proteins. Given that PlzA can bind to both DNA and RNA, it could have a role in

coordinating transcriptional complexes, and it supports the notion that PlzA can interact with

RNAP (Grassmann *et al.*, 2023). Our study examined PlzA nucleic-acid binding at a locus

known to be affected by PlzA, the *glpFKD* catabolism operon, but sites/ligands of higher affinity

could exist. It remains to be seen what additional targets in the *B. burgdorferi*

genome/transcriptome fall under the umbrella of the PlzA regulon. Our work identifying PlzA

as a novel nucleic acid-binding protein provides a basis for its functional mechanism. This will

further inform our understanding of how *B. burgdorferi* regulates gene expression in response

to environmental cues and help unravel the PlzA regulon.

MATERIAL AND METHODS

Routine manipulations and molecular cloning

Oligonucleotides used in this study are listed in Table 2 and were ordered from Integrated

DNA Technologies (IDT). *B. burgdorferi* genomic DNA was purified utilizing the E.Z.N.A Tissue

DNA kit (OMEGA). Standard and high-fidelity PCR were performed with 2X DreamTaq Green

PCR Master Mix (Thermo Scientific) or Q5 High-Fidelity 2X Master Mix (NEB), respectively.

Plasmids were isolated using QIAprep Spin Miniprep kits (Qiagen) according to the Miraprep

protocol (Pronobis *et al.*, 2016). All plasmid constructs were transformed into chemically

competent *E. coli* Top10 (Invitrogen) or DH5α (Invitrogen) strains for cloning and plasmid

maintenance. Positive clones were identified by colony PCR screening with the appropriate

primer sets, and DNA sequencing was performed by Eurofins Genomics LLC (Louisville, KY).

Plasmid construction

The wild-type (WT) *plzA* gene from *B. burgdorferi* strain B31, was cloned between the

NdeI/XhoI sites of pET28a(+) and thus fused in-frame with an amino-terminal 6xHis-tag.

Truncated *plzA* genes, named *plzA*-NTD and *plzA*-CTD encoding the N-terminal (residues 1-

141) and C-terminal (residues 142-261) domains of PlzA, respectively, were generated via

gene synthesis and cloned to produce 6xHis-tagged proteins by Genscript (Piscataway, NJ).

Briefly, the synthesized truncations were cloned into the pET28b(+) vector with *plzA*-NTD

cloned between the BamHI/NotI sites producing an amino-terminal 6xHis-tag fusion, while

plzA CTD was cloned between the NcoI/XhoI sites producing a carboxy-terminal 6xHis-tag

fusion.

Generation of the site-directed PlzA R145D-R149D double mutant

Site-directed mutagenesis by overlap extension PCR was used to generate a PlzA mutant that

cannot bind c-di-GMP. Two-point mutations were introduced into a pTrcHis TOPO (Invitrogen)

construct containing the *plzA* gene. The successful introduction of the mutations was

confirmed by DNA sequencing. The site-directed PlzA double mutant is denoted as PlzA$_{RD-RD}$.

The oligonucleotide primers used for site-directed mutagenesis are listed in Table 2.

Recombinant protein overexpression and purification

For recombinant protein expression, *E. coli* Rosetta 2(DE3)(pLysS) (Novagen) chemically

competent cells were used for PlzA$_{WT}$, PlzA$_{RD-RD}$, and N-terminal PlzA domain (PlzA$_{NTD}$), while

One Shot BL21 Star (DE3) cells were used for the C-terminal PlzA domain (PlzA$_{CTD}$).

Overnight bacterial cultures were diluted 1:100 into Super broth (tryptone 32 g, yeast extract

20 g, NaCl 5 g per liter) supplemented with the appropriate antibiotics (kanamycin-50 µg/ml,

carbenicillin-100 µg/ml, and/or chloramphenicol-30 µg/ml). The cultures were allowed to grow

to an OD$_{600}$ of 0.5-1.0 and protein expression was subsequently induced by the addition of

0.25 mM (PlzA$_{NTD}$ only) or 0.5 mM isopropyl-β-D-thiogalactopyranoside (IPTG). Induced

cultures were incubated at either 37°C for 3-4 h (PlzA$_{WT}$ and PlzA$_{CTD}$), 29°C for 4-6 h (PlzA$_{WT}$),

or overnight at room temperature (PlzA$_{RD-RD}$ and PlzA$_{NTD}$). The cells were harvested by

centrifugation at 5400 x *g* for 30 min at 4°C, and the cell pellets were frozen at -80°C until

protein purification.

For protein extraction, cell pellets were thawed on ice and resuspended in equilibration buffer

(20mM sodium phosphate, 300 mM NaCl, 10 mM imidazole, pH 8.0) and lysed via sonication.

Lysates were clarified by centrifugation at 23700 x *g* for 20 min at 4°C, and the supernatant

was retained. When further clarification was required due to lysate viscosity, the supernatants

were passed through a sterile 0.22 µM Millex-GS Syringe Filter Unit (MilliporeSigma). Protein

purification was performed via column affinity chromatography using either HisPur Ni-NTA

resin (Thermo Scientific) or the Magne-His purification system (Promega Corporation) for

PlzA$_{RD-RD}$ per manufacturer protocol. Eluted protein solutions were dialyzed overnight into

EMSA buffer (50 mM Tris-HCl [pH 7.5], 50 mM KCl, 1 mM dithiothreitol (DTT), 1 mM

ethylenediaminetetraacetic acid (EDTA) [pH 8.0], 1 mM phenylmethanesulfonyl fluoride

(PMSF), 10% glycerol (v/v), and 0.01% Tween-20). The dialyzed proteins were then

concentrated using Amicon Ultra 3K (PlzA$_{N}$ and PlzA$_{C}$) or 10K (PlzA$_{WT}$ and PlzA$_{RD-RD}$)

centrifugal filter units (MilliporeSigma). Aliquots of dialyzed proteins were assessed for purity

by SDS-PAGE and Coomassie brilliant blue staining. Final recombinant protein concentrations

were determined using Quick Start Bradford Protein Assays (Bio-Rad) with bovine serum

albumin as the reference protein for standard curves. Purified protein aliquots were stored at

-80°C.

Nucleic acid fragments design and generation

All probe sequences were derived from the genomic sequence of *B. burgdorferi* type strain

B31, and the corresponding primers are reported in Table 2 (Fraser *et al.*, 1997; Casjens *et*

al., 2000). Large and small DNA substrates were used in this study. Larger fluorescently

tagged fragments >60 bp were generated by PCR amplification of the target sequences from

B31 genomic DNA and subsequent cloning into the pCR 2.1 TA vector (Invitrogen) following

TOPO TA protocols. Subsequently, target-specific or M13 F and R IRDye 800-labeled primers

were used for PCR of the plasmid templates to produce labeled DNA fragments for binding

studies. All PCR-generated probes were treated with exonuclease I (NEB) according to the

manufacturer's protocol to remove single-stranded DNA, followed by ethanol precipitation.

The DNA was resuspended in molecular-grade water (Ambion) and quantified by Nanodrop

UV spectrometry.

Smaller probes <60 bp long were produced by annealing complementary unlabeled and IRDye

800-labeled oligonucleotides at equimolar concentrations at 95°C for 5 min and were gradually

cooled overnight at RT. All probes were aliquoted and stored at -20°C until further use.

Unlabeled and 5' IRDye 800 (LI-COR Biosciences) labeled oligonucleotides were synthesized

by IDT. Alexa Fluor 488 conjugated RNA probes corresponding to the transcribed regions of

the respective DNA probes were also synthesized by IDT.

Electrophoretic mobility shift assays (EMSA)

EMSA reactions were performed in EMSA buffer at RT with a final concentration of dsDNA

probe of 10 nM or RNA probe of either 10 nM or 50 nM. EMSA reaction mixtures were

supplemented with 3'5' c-di-GMP (Sigma-Aldrich) at various concentrations ranging from 0 to

100 µM. c-di-GMP was not added to the gel or running buffer. Experiments using equimolar

concentrations of c-di-GMP are relative to the final protein concentration in the reaction.

Proteins were co-incubated with c-di-GMP for 5 min before the addition of any nucleic acids.

Prior to the addition of labeled probe, the non-specific competitor poly-dI-dC (Roche) was

added to the EMSA reactions at a final concentration of 2.5 ng/µL, and the reaction mix was

allowed to incubate another 5 min (Larouche *et al.*, 1996). Labeled probes were added last,

and the reactions were subsequently incubated for an additional 10 min prior to loading onto

gels. Competition EMSAs were conducted similarly but with the unlabeled, specific competitor

added after poly-dI-dC and then allowed to incubate with the reaction prior to the addition of

the labeled probe. Regarding RNA EMSAs, when RNase contamination was evident in

recombinant protein preparations, RiboGuard (Lucigen) or SUPERase•In (Ambion) RNase

inhibitor was added to the EMSA mixtures before the addition of RNA to a final concentration

of 4 U/µL and 2 U/µL respectively to inhibit RNase activity. The final reaction volume was 10

µL prior to the addition of loading dye.

For visualization during electrophoresis, 2 µL of EMSA loading dye (0.8 mg/mL Orange G, 15

mg/mL Ficoll 400) was added to each reaction prior to gel loading. Novex TBE 6 or 10% gels

(Invitrogen) were pre-run in 0.5x TBE running buffer (pH ~8.4 +/- 0.5), at 100V for a minimum

of 30 min. Due to the complexes of the larger *glpFKD*(-219) probe and higher concentrations

of PlzA not exiting the well, the pH of the running buffer was adjusted to a pH of 8.0

immediately prior to electrophoresis which allowed the complexes to enter the gel matrix. The

pH was not adjusted for EMSAs using nucleic acid fragments <60 bp as complexes readily

exited the well at the standard pH of 0.5x TBE running buffer. The entire reaction mixture was

then loaded onto the pre-run gels and resolved at 100 V for 60-90 min at RT. EMSA images

were acquired with a ChemiDoc Imaging System (Bio-Rad).

Densitometry and statistics

Densitometry of EMSAs was performed using Image Lab 6.1 (Bio-Rad) software. Analyses

were performed on triplicate EMSAs. Lanes as well as free and shifted bands were added

manually, and intensity values were reported as band percentage. For dsDNA EMSAs, free

dsDNA band percentages were normalized to the probe only control band percentage. The

free DNA band percentage values were then used to determine the percent shifted of the

respective probe relative to the free probe. For RNA and ssDNA EMSAs, lanes and bands

were selected manually, and the shifted bands were analyzed to obtain the percent shifted

values. Probe signal stuck in the gel wells, attributed to potential protein aggregation, were

not considered shifts, and thus omitted from densitometric analysis.

To calculate the apparent dissociation constant (K_D), nonlinear regression analysis was

performed to determine the best-fit values for each experimental condition tested. Briefly,

protein concentrations were plotted against the shifted band percentage values obtained from

densitometry using the one-site specific binding setting in Prism GraphPad 9.5.1. Confidence

was set at 95% with the analysis considering each replicate value from triplicate EMSAs

performed for the analysis. Either a Welch's t-test or Welch ANOVA followed by a Dunnett's

T3 multiple comparisons test, was used to compare fits.

Docking analysis and protein modeling

Computational modeling of holo-PlzA (PDB ID: 7mie) in complex with the nucleic acids of

interest was performed using the HDOCK server: http://hdock.phys.hust.edu.cn/. Briefly, the

PDB and chain ID of PlzA was entered for the input receptor molecule to retrieve the crystal

structure of the protein for the analysis. For each analysis, the corresponding nucleic acid

sequence of interest was then pasted as the input ligand molecule, and the type of nucleic

acid selected from the drop-down menu. The input nucleic acid sequences are listed in Table

2. All analyses were performed with default parameters. The top model for each computation

is shown. Docking and confidence scores are provided in the corresponding figure legends.

The more negative the docking score, the more probable the binding. A confidence score

above 0.7 indicates the molecules would be highly likely to bind.

Protein structure prediction for PlzA$_{NTD}$, PlzA$_{CTD}$, and PlzA$_{RD-RD}$ was performed via AlphaFold.

All output PDB files generated from the various analyses were viewed with ChimeraX 1.4

(Goddard *et al.*, 2018; Jumper *et al.*, 2021; Pettersen *et al.*, 2021).

Cyclic di-nucleotide LC-MS analysis

Detection of cyclic di-nucleotides in recombinant protein purifications was conducted by liquid

chromatography-tandem mass spectrometry (LC-MS/MS) at the Michigan State University

Mass Spectrometry core. Briefly, recombinant proteins were purified and quantified as

mentioned previously and subsequently dialyzed into EMSA buffer without Tween-20. For LC-

MS/MS analysis, approximately 1 mg/mL of PlzA$_{WT}$, PlzA$_{RD-RD,}$ PlzA$_{NTD}$, and PlzA$_{CTD}$ were

aliquoted and stored at -80°C until shipment to the core facility. Nucleotides bound to purified

protein samples (1mg/mL) were analyzed by precipitating protein with 3 volumes of acetonitrile

followed by centrifugation to pellet precipitated protein. The supernatant was transferred to a

new tube and evaporated to dryness. The sample was reconstituted in mobile phase solvent

A (10 mM tributylamine + 15 mM acetic acid in water/methanol, 97:3 v/v). LC-MS analysis was

performed on a Waters Xevo G2-XS Quadrupole-Time-of-Flight (QTof) mass spectrometer

interfaced with a Waters Acquity UPLC system. 10 μL of sample was injected onto a Waters

Acquity BEH-C18 UPLC column (2.1x50 mm) and compounds were separated using the

following gradient: initial conditions were 100% mobile phase A, hold at 100% A until 1 min,

ramp to 99% B at 7 min (mobile phase B: methanol), hold at 99% B until 8 min, return to 100%

A at 8.01 min and hold until 10 min. The flow rate through the column was 0.3 ml/min and the

column temperature was held at 40°C. Compounds were ionized by electrospray ionization

operated in negative mode. Capillary voltage was 2.0 kV, cone voltage was 35, source

temperature was 100C and desolvation temperature was 350C. The cone gas and desolvation

gas flows were 50 L/hr and 600 L/hr respectively. A TOF MS scan method with targeted

enhancement of m/z 689 was used with 0.5 second scan time. Lockmass correction was

performed using leucine enkephalin as the reference compound.

Data were processed using Waters Masslynx software. Extracted ion chromatograms were

performed to look for the presence of c-di-GMP (m/z 689.09), c-di-AMP (m/z 657.09) and c-

GAMP (673.09). The intensity values were converted to relative abundance as determined

from the base peak values. The c-di-GMP concentration was determined from the peak area

of the target compound against a standard curve of known c-di-GMP concentrations. These

concentrations were calculated using the Targetlynx tool in the Waters Masslynx software.

The percentage of protein bound with c-di-GMP was calculated from the total protein and

calculated c-di-GMP concentrations in each sample.

Circular dichroism spectroscopy

The CD analysis was performed on PlzA$_{WT}$, PlzA$_{RD-RD}$, PlzA$_{NTD}$, and PlzA$_{CTD}$ in EMSA dialysis

buffer (detailed above) at the University of Kentucky Center for Molecular Medicine Protein

Core. Briefly, 400 µL of each protein solution at a concentration of 180-210 µg/mL was pipetted

into a 1 mm quartz cuvette and loaded into a Jasco J-810 spectrophotometer. The samples

were scanned at 0.1 nm increments and CD spectra collected for wavelengths 200-260 nM.

EMSA buffer alone or supplemented with 50 µM c-di-GMP were also analyzed and used as

controls. Data were visualized in GraphPad Prism 9.5.1.

AVAILABILITY

HDOCK (http://hdock.phys.hust.edu.cn/)

ACKNOWLEDGEMENT

We dedicate this manuscript to our friend and colleague Christina R. Savage, Ph.D., whose

initial studies on PlzA were invaluable to this work. We also thank Jessamyn Moore and

Tatiana Castro Padovani for their support of the studies undertaken. Circular Dichroism was

performed at the University of Kentucky Center for Molecular Medicine Protein Core, and we

thank Martin Chow for his assistance on these studies. The LC-MS/MS to detect di-nucleotides

was performed by the Mass Spectrometry and Metabolomics Core at Michigan State

University. A sincere thank you to Tony Schilmiller at MSU for all his help with the mass

spectrometry services. Figures 1, 3a, 5a, 8a and 11 were created via BioRender.

FUNDING

Funding for open access charge: US National Institutes of Health (grant R01 AI144126-3 to

B.S.).

CONFLICT OF INTEREST

None declared.

REFERENCES

Adams, P.P., Avile, C.F., Popitsch, N., Bilusic, I., Schroeder, R., Lybecker, M., and
Jewett, M.W. (2017) In vivo expression technology and 5′ end mapping of the *Borrelia*
burgdorferi transcriptome identify novel RNAs expressed during mammalian infection.
Nucleic Acids Res **45**: 775–792.

Amikam, D., and Galperin, M.Y. (2006) PilZ domain is part of the bacterial c-di-GMP
binding protein. *Bioinformatics* **22**: 3–6
https://academic.oup.com/bioinformatics/article/22/1/3/218448. Accessed February 18,
2022.

Babb, K., Bykowski, T., Riley, S.P., Miller, M.C., Demoll, E., and Stevenson, B. (2006)
Borrelia burgdorferi EbfC, a novel, chromosomally encoded protein, binds specific DNA
sequences adjacent to erp loci on the spirochete's resident cp32 prophages. *J Bacteriol*
188: 4331–4339 https://journals.asm.org/journal/jb. Accessed April 8, 2022.

Bauer, W.J., Luthra, A., Zhu, G., Radolf, J.D., Malkowski, M.G., and Caimano, M.J.
(2015) Structural characterization and modeling of the *Borrelia burgdorferi* hybrid
histidine kinase Hk1 periplasmic sensor: A system for sensing small molecules
associated with tick feeding. *J Struct Biol* **192**: 48–58.

Bong, H.J., Ko, E.M., Song, S.Y., Ko, I.J., and Oh, J. Il (2019) Tripartite regulation of
the *glpFKD* operon involved in glycerol catabolism by gylr, crp, and sigf in
Mycobacterium smegmatis. *J Bacteriol* **201**.

Burns, L.H., Adams, C.A., Riley, S.P., Jutras, B.L., Bowman, A., Chenail, A.M., *et al.*
(2010) BpaB, a novel protein encoded by the Lyme disease spirochete's cp32
prophages, binds to *erp* Operator 2 DNA. *Nucleic Acids Res* **38**: 5443–5455.

Caimano, M.J., Dunham-Ems, S., Allard, A.M., Cassera, M.B., Kenedy, M., and Radolf,
J.D. (2015) Cyclic di-GMP modulates gene expression in Lyme disease spirochetes at
the tick-mammal interface to promote spirochete survival during the blood meal and

tick-to-mammal transmission. *Infect Immun* **83**: 3043–3060 http://dx.doi.org/10.1128.
Accessed May 1, 2021.

Caimano, M.J., Kenedy, M.R., Kairu, T., Desrosiers, D.C., Harman, M., Dunham-Ems,
S., *et al.* (2011) The hybrid histidine kinase Hk1 is part of a two-component system that
is essential for survival of *Borrelia burgdorferi* in feeding *Ixodes scapularis* ticks. *Infect*
Immun **79**: 3117–3130 http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml. Accessed
February 13, 2022.

Carey, J. (2022) Affinity, Specificity, and Cooperativity of DNA Binding by Bacterial
Gene Regulatory Proteins. *Int J Mol Sci* **23**: 1–16.

Casjens, S., Palmer, N., Vugt, R. Van, Huang, W.M., Stevenson, B., Rosa, P., *et al.*
(2000) A bacterial genome in flux: The twelve linear and nine circular
extrachromosomal DNAs in an infectious isolate of the Lyme disease spirochete
Borrelia burgdorferi. *Mol Microbiol* **35**: 490–516.

Cheang, Q.W., Xin, L., Chea, R.Y.F., and Liang, Z.X. (2019) Emerging paradigms for
PilZ domain-mediated C-di-GMP signaling. *Biochem Soc Trans* **47**: 381–388.

Chou, S.H., and Galperin, M.Y. (2016) Diversity of cyclic di-GMP-binding proteins and
mechanisms. *J Bacteriol* **198**: 32–46 http://ncbi.nlm.nih.gov/Complete_Genomes/c-di-
GMP.html. Accessed February 13, 2022.

Corona, A., and Schwartz, I. (2015) *Borrelia burgdorferi* : Carbon Metabolism and the
Tick-Mammal Enzootic Cycle. *Microbiol Spectr* **3**.

Fraser, C.M., Casjens, S., Huang, W.M., Sutton, G.G., Clayton, R., Lathigra, R., *et al.*
(1997) Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature*
390: 580–586.

Freedman, J.C., Rogers, E.A., Kostick, J.L., Zhang, H., Iyer, R., Schwartz, I., and
Marconi, R.T. (2009) Identification and molecular characterization of a cyclic-di-GMP
effector protein, PlzA (BB0733): Additional evidence for the existence of a functional
cyclic-di-GMP regulatory network in the Lyme disease spirochete, *Borrelia burgdorferi*.
FEMS Immunol Med Microbiol **58**: 285–294
https://academic.oup.com/femspd/article/58/2/285/597665. Accessed April 26, 2021.

Fried, M., and Crothers, D.M. (1981) Equilibria and kinetics of lac repressor-operator
interactions by polyacrylamide gel electrophoresis. *Nucleic Acids Res* **9**: 1981.

Fried, M.G. (1989) Measurement of protein-DNA interaction parameters by
electrophoresis mobility shift assay. *Electrophoresis* **10**: 366–376.

Galperin, M.Y., and Chou, S.H. (2020) Structural conservation and diversity of PilZ-
related domains. *J Bacteriol* **202**.

Galperin, M.Y., Nikolskaya, A.N., and Koonin, E. V. (2001) Novel domains of the
prokaryotic two-component signal transduction systems. *FEMS Microbiol Lett* **203**: 11–
21 https://academic.oup.com/femsle/article/203/1/11/478433. Accessed March 17,
2022.

Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H.,
and Ferrin, T.E. (2018) UCSF ChimeraX: Meeting modern challenges in visualization
and analysis. *Protein Science* **27**: 14–25.

Grassmann, A.A., Tokarz, R., Golino, C., McLain, M.A., Groshong, A.M., Radolf, J.D.,
and Caimano, M.J. (2023) BosR and PlzA reciprocally regulate RpoS function to
sustain *Borrelia burgdorferi* in ticks and mammals. *Journal of Clinical Investigation*
http://www.jci.org/articles/view/166710.

Groshong, A.M., Grassmann, A.A., Luthra, A., Mclain, M.A., Provatas, A.A., Radolf,
J.D., and Caimano, M.J. (2021) PlzA is a bifunctional c-di-GMP biosensor that
promotes tick and mammalian host- adaptation of *Borrelia burgdorferi*. *PLoS Pathog*
17: 1–31.

Grove, A.P., Liveris, D., Iyer, R., Petzke, M., Rudman, J., Caimano, M.J., *et al.* (2017)
Two Distinct Mechanisms govern RpoS-Mediated Repression of Tick-Phase Genes
during Mammalian Host Adaptation by *Borrelia burgdorferi*, the Lyme Disease
Spirochete. *mBio* **8** https://journals.asm.org/doi/full/10.1128/mBio.01204-17. Accessed
June 18, 2022.

He, M., Ouyang, Z., Troxell, B., Xu, H., Moh, A., Piesman, J., *et al.* (2011) Cyclic di-gmp
is essential for the survival of the lyme disease spirochete in ticks. *PLoS Pathog* **7**.

He, M., Zhang, J.J., Ye, M., Lou, Y., and Yang, X.F. (2014) Cyclic Di-GMP receptor
PlzA controls virulence gene expression through RpoS in *Borrelia burgdorferi*. *Infect*
Immun **82**: 445–452 http://iai.asm.org/. Accessed April 26, 2021.

Helble, J.D., McCarthy, J.E., and Hu, L.T. (2021) Interactions between *Borrelia*
burgdorferi and its hosts across the enzootic cycle. *Parasite Immunol* **43**.

Hellman, L.M., and Fried, M.G. (2007) Electrophoretic mobility shift assay (EMSA) for
detecting protein-nucleic acid interactions. *Nat Protoc* **2**: 1849–1861.

Hengge, R. (2009) Principles of c-di-GMP signalling in bacteria. *Nat Rev Microbiol* **7**:
263–273.

Hsieh, M.L., Hinton, D.M., and Waters, C.M. (2018) VpsR and cyclic di-GMP together
drive transcription initiation to activate biofilm formation in *Vibrio cholerae*. *Nucleic*
Acids Res **46**: 8876–8887.

Huang, S.Y., and Zou, X. (2014) A knowledge-based scoring function for protein-RNA
interactions derived from a statistical mechanics-based iterative method. *Nucleic Acids*
Res **42**.

Jarmoskaite, I., Alsadhan, I., Vaidyanathan, P.P., and Herschlag, D. (2020) How to
measure and evaluate binding affinities. *Elife* **9**: 1–34.

Jenal, U., Reinders, A., and Lori, C. (2017) Cyclic di-GMP: Second messenger
extraordinaire. *Nat Rev Microbiol* **15**: 271–284
http://dx.doi.org/10.1038/nrmicro.2016.190.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., *et al.*
(2021) Highly accurate protein structure prediction with AlphaFold. *Nature* **596**: 583–
589 http://dx.doi.org/10.1038/s41586-021-03819-2.

Junkermeier, E.H., and Hengge, R. (2023) Local signaling enhances output specificity
of bacterial c-di-GMP signaling networks. *microLife* **4**.

Jutras, B.L., Bowman, A., Brissette, C.A., Adams, C.A., Verma, A., Chenail, A.M., and
Stevenson, B. (2012) EbfC (YbaB) is a new type of bacterial nucleoid-associated
protein and a global regulator of gene expression in the Lyme disease spirochete. *J*
Bacteriol **194**: 3395–3406.

Jutras, B.L., Chenail, A.M., Carroll, D.W., Miller, M.C., Zhu, H., Bowman, A., and
Stevenson, B. (2013a) Bpur, the Lyme disease spirochete's PUR domain protein:
Identification as a transcriptional modulator and characterization of nucleic acid
interactions. *Journal of Biological Chemistry* **288**: 26220–26234 http://www.jbc.org/.
Accessed October 5, 2020.

Jutras, B.L., Chenail, A.M., Rowland, C.L., Carroll, D., Miller, M.C., Bykowski, T., and
Stevenson, B. (2013b) Eubacterial SpoVG Homologs Constitute a New Family of Site-
Specific DNA-Binding Proteins. *PLoS One* **8**: 66683 www.plosone.org. Accessed April
26, 2021.

Kim, B., and Little, J.W. (1992) Dimerization of a specific DNA-binding protein on the
DNA. *Science (1979)* **255**: 203–206.

Kohler, J.J., Metallo, S.J., Schneider, T.L., and Schepartz, A. (1999) DNA specificity
enhanced by sequential binding of protein monomers. *Proc Natl Acad Sci U S A* **96**:
11735–11739.

Kostick, J.L., Szkotnicki, L.T., Rogers, E.A., Bocci, P., Raffaelli, N., and Marconi, R.T.
(2011) The diguanylate cyclase, Rrp1, regulates critical steps in the enzootic cycle of
the Lyme disease spirochetes. *Mol Microbiol* **81**: 219–231
https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2958.2011.07687.x. Accessed
April 15, 2022.

Kostick-Dunn, J.L., Izac, J.R., Freedman, J.C., Szkotnicki, L.T., Oliver, L.D., and
Marconi, R.T. (2018) The *Borrelia burgdorferi* c-di-GMP Binding Receptors, PlzA and
PlzB, Are Functionally Distinct. *Front Cell Infect Microbiol* **8**: 213 www.frontiersin.org.
Accessed April 26, 2021.

Krusenstjerna, A.C., Arnold, W.K., Saylor, T.C., Tucker, J.S., and Stevenson, B. (2023)
Borrelia burgdorferi DnaA and the nucleoid-associated protein EbfC coordinate
expression of the *dnaX-ebfC* operon. *J Bacteriol* **205**: e00396-22
https://doi.org/10.1128/jb.00396-22.

Larouche, K., Bergeron, M.-J., Leclerc, S., and Guerin, S. (1996) Optimization of
competitor poly(dI-dC).poly(dI-dC) levels is advised in DNA-protein interaction studies
involving enriched nuclear proteins. *Biotechniques* **20**: 439–444.

Lee, R.E., and Baust, J.G. (1987) Cold-Hardiness in the Antarctic Tick, *Ixodes uriae*.
Physiol Zool **60**: 499–506 https://about.jstor.org/terms.

Mallory, K.L., Miller, D.P., Oliver, L.D., Freedman, J.C., Kostick-Dunn, J.L., Carlyon,
J.A., *et al.* (2016) Cyclic-di-GMP binding induces structural rearrangements in the PlzA
and PlzC proteins of the Lyme disease and relapsing fever spirochetes: A possible
switch mechanism for c-di-GMP-mediated effector functions. *Pathog Dis* **74**: 1–8.

Novak, E.A., Sultan, S.Z., and Motaleb, M.A. (2014) The cyclic-di-GMP signaling
pathway in the Lyme disease spirochete, *Borrelia burgdorferi*. *Front Cell Infect Microbiol*
4 www.frontiersin.org. Accessed April 26, 2021.

Pappas, C.J., Iyer, R., Petzke, M.M., Caimano, M.J., Radolf, J.D., and Schwartz, I.
(2011) *Borrelia burgdorferi* requires glycerol for maximum fitness during the tick phase
of the enzootic cycle. *PLoS Pathog* **7**.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., *et al.*
(2021) UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Science* **30**: 70–82.

Pitzer, J.E., Syed, S.Z., Hayakawa, Y., Hobbs, G., Miller, M.R., and Motaleb, M.A.
(2011) Analysis of the *Borrelia burgdorferi* Cyclic-di-GMP-Binding Protein PlzA Reveals
a Role in Motility and Virulence. *Infect Immun* **79**: 1815–1825 http://iai.asm.org/.
Accessed April 26, 2021.

Pronobis, M.I., Deuitch, N., and Peifer, M. (2016) The Miraprep: A protocol that uses a
Miniprep kit and provides Maxiprep yields. *PLoS One* **11**: 1–12.

Pultz, I.S., Christen, M., Kulasekara, H.D., Kennard, A., Kulasekara, B., and Miller, S.I.
(2012) The response threshold of Salmonella PilZ domain proteins is determined by
their binding affinities for c-di-GMP. *Mol Microbiol* **86**: 1424–1440.

Riley, S.P., Bykowski, T., Cooley, A.E., Burns, L.H., Babb, K., Brissette, C.A., *et al.*
(2009) *Borrelia burgdorferi* EbfC defines a newly-identified, widespread family of
bacterial DNA-binding proteins. *Nucleic Acids Res* **37**: 1973–1983 http://www.rcsb.org/.
Accessed April 8, 2022.

Rogers, E.A., Terekhova, D., Zhang, H.-M.M., Hovis, K.M., Schwartz, I., and Marconi,
R.T. (2009) Rrp1, a cyclic-di-GMP-producing response regulator, is an important
regulator of *Borrelia burgdorferi* core cellular functions. *Mol Microbiol* **71**: 1551–1573
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2843504&tool=pmcentrez&re
ndertype=abstract. Accessed February 18, 2022.

Romling, U., Galperin, M.Y., and Gomelsky, M. (2013) Cyclic di-GMP: the First 25
Years of a Universal Bacterial Second Messenger. *Microbiology and Molecular Biology*
Reviews **77**: 1–52.

Rooman, M., Lié Vin, J., Buisine, E., and Wintjens, R. (2002) Cation-p/H-bond Stair
Motifs at Protein-DNA Interfaces. *J Mol Biol* **319**: 67–76 http://www.idealibrary.com.

Ryjenkov, D.A., Simm, R., Römling, U., and Gomelsky, M. (2006) The PilZ domain is a
receptor for the second messenger c-di-GMP: The PilZ domain protein YcgR controls
motility in enterobacteria. *Journal of Biological Chemistry* **281**: 30310–30314.

Ryjenkov, D.A., Tarutina, M., Moskvin, O. V., and Gomelsky, M. (2005) Cyclic
diguanylate is a ubiquitous signaling molecule in bacteria: Insights into biochemistry of
the GGDEF protein domain. *J Bacteriol* **187**: 1792–1798
https://journals.asm.org/doi/abs/10.1128/JB.187.5.1792-1798.2005. Accessed March
17, 2022.

Sarenko, O., Klauck, G., Wilke, F.M., Pfiffer, V., Richter, A.M., Herbst, S., *et al.* (2017)
More than enzymes that make or break cyclic Di-GMP—local signaling in the
interactome of GGDEF/EAL domain proteins of Escherichia coli. *mBio* **8**.

Sathyapriya, R., and Vishveshwara, S. (2004) Interaction of DNA with clusters of amino
acids in proteins. *Nucleic Acids Res* **32**: 4109–4118.

Savage, C.R., Jutras, B.L., Bestor, A., Tilly, K., Rosa, P.A., Tourand, Y., *et al.* (2018)
Borrelia burgdorferi SpoVG DNA- and RNA-binding protein modulates the physiology of
the Lyme disease spirochete. *J Bacteriol* **200** http://jb.asm.org/. Accessed April 26,
2021.

Saylor, T.C., Savage, C.R., Krusenstjerna, A.C., Jusufovic, N., Zückert, W.R., Brissette,
C.A., *et al.* (2023) Quantitative analyses of interactions between SpoVG and RNA/DNA.
Biochem Biophys Res Commun **654**: 40–46.

Schäper, S., Steinchen, W., Krol, E., Altegoer, F., Skotnicka, D., Søgaard-Andersen, L.,
et al. (2017) AraC-like transcriptional activator CuxR binds c-di-GMP by a PilZ-like
mechanism to regulate extracellular polysaccharide production. *Proc Natl Acad Sci U S*
A **114**: E4822–E4831.

Schramm, F.D., Schroeder, K., and Jonas, K. (2019) Protein aggregation in bacteria.
FEMS Microbiol Rev **44**: 54–72.

Singh, A., Izac, J.R., Schuler, E.J.A., Patel, D.T., Davies, C., and Marconi, R.T. (2021)
High-resolution crystal structure of the *Borreliella burgdorferi* PlzA protein in complex
with c-di-GMP: new insights into the interaction of c-di-GMP with the novel xPilZ
domain. *Pathog Dis* **79**: 1–9.

Skotnicka, D., Steinchen, W., Szadkowski, D., Cadby, I.T., Lovering, A.L., Bange, G.,
and Søgaard-Andersen, L. (2020) CdbA is a DNA-binding protein and c-di-GMP
receptor important for nucleoid organization and segregation in *Myxococcus xanthus*.
Nat Commun **11** https://doi.org/10.1038/s41467-020-15628-8. Accessed February 13,
2022.

Sultan, S.Z., Pitzer, J.E., Boquoi, T., Hobbs, G., Miller, M.R., and Motaleb, M.A. (2011)
Analysis of the HD-GYP domain cyclic dimeric gmp phosphodiesterase reveals a role in
motility and the enzootic life cycle of *Borrelia burgdorferi*. *Infect Immun* **79**: 3273–3283.

Sultan, S.Z., Pitzer, J.E., Miller, M.R., and Motaleb, M.A. (2010) Analysis of a *Borrelia*
burgdorferi phosphodiesterase demonstrates a role for cyclic-di-guanosine
monophosphate in motility and virulence. *Mol Microbiol* **77**: 128–142.

Tan, J.W.H., Wilksch, J.J., Hocking, D.M., Wang, N., Srikhanta, Y.N., Tauschek, M., *et*
al. (2015) Positive Autoregulation of *mrkHI* by the Cyclic Di-GMP-Dependent MrkH
Protein in the Biofilm Regulatory Circuit of *Klebsiella pneumoniae*. *J Bacteriol* **197**:
1659–1667.

Tschowri, N., Schumacher, M.A., Schlimpert, S., Chinnam, N.B., Findlay, K.C.,
Brennan, R.G., and Buttner, M.J. (2014) Tetrameric c-di-GMP mediates effective
transcription factor dimerization to control *Streptomyces* development. *Cell* **158**: 1136–
1147 http://dx.doi.org/10.1016/j.cell.2014.07.022.

Valentini, M., and Filloux, A. (2019) Multiple roles of c-di-GMP signaling in bacterial
pathogenesis. *Annu Rev Microbiol* **73**: 387–406.

Van Gundy, T., Patel, D., Bowler, B.E., Rothfuss, M.T., Hall, A.J., Davies, C., *et al.*
(2023) c-di-GMP regulates activity of the PlzA RNA chaperone from the Lyme disease
spirochete. *Mol Microbiol* **119**: 711–727.

Wang, F., He, Q., Su, K., Gao, F., Huang, Y., Lin, Z., *et al.* (2016) The PilZ domain of
MrkH represents a novel DNA binding motif. *Protein Cell* **7**: 766–772
/pmc/articles/PMC5055493/. Accessed February 13, 2022.

Wilksch, J.J., Yang, J., Clements, A., Gabbe, J.L., Short, K.R., Cao, H., *et al.* (2011)
MrkH, a Novel c-di-GMP-Dependent Transcriptional Activator, Controls *Klebsiella*
pneumoniae Biofilm Formation by Regulating Type 3 Fimbriae Expression. *PLoS*
Pathog **7**: e1002204
https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1002204.
Accessed February 14, 2022.

Yan, Y., Tao, H., He, J., and Huang, S.Y. (2020) The HDOCK server for integrated
protein–protein docking. *Nat Protoc* **15**: 1829–1852.

Yan, Y., Wen, Z., Wang, X., and Huang, S.Y. (2017a) Addressing recent docking
challenges: A hybrid strategy to integrate template-based and free protein-protein
docking. *Proteins: Structure, Function and Bioinformatics* **85**: 497–512.

Yan, Y., Zhang, D., Zhou, P., Li, B., and Huang, S.Y. (2017b) HDOCK: A web server for
protein-protein and protein-DNA/RNA docking based on a hybrid strategy. *Nucleic*
Acids Res **45**: W365–W373.

Yang, J., Wilksch, J.J., Tan, J.W.H., Hocking, D.M., Webb, C.T., Lithgow, T., *et al.*
(2013) Transcriptional activation of the *mrkA* promoter of the *Klebsiella pneumoniae*
type 3 fimbrial operon by the c-di-GMP-dependent MrkH protein. *PLoS One* **8**.

Zamorano-Sánchez, D., Fong, J.C.N., Kilic, S., Erill, I., and Yildiz, F.H. (2015)
Identification and characterization of VpsR and VpsT binding sites in *Vibrio cholerae*. *J*
Bacteriol **197**: 1221–1235 http://dx.doi.org/10.1128. Accessed May 2, 2021.

Zhang, J.J., Chen, T., Yang, Y., Du, J., Li, H., Troxell, B., *et al.* (2018) Positive and
Negative Regulation of Glycerol Utilization by the c-di-GMP Binding Protein PlzA in
Borrelia burgdorferi. *J Bacteriol* **200** https://doi.org/10.1128/JB. Accessed April 14,
2021.

Zhang, J.J., Ranghunandanan, S., Wang, Q., Lou, Y., and Yang, X.F. (2022)
Mechanism of repression of the glycerol utilization operon in *Borrelia burgdorferi*.
bioRxiv **PREPRINT:NOT PEER REVIEWED**

FIGURES LEGENDS

FIGURE 1 The *glpFKD* operon, 5' UTR, and designed probe schematic. The glycerol

catabolism gene operon (*glpFKD*) contains four genes: *glpF* (BB_0240, glycerol uptake

facilitator protein), *glpK* (BB_0241, glycerol kinase), BB_0242 (hypothetical protein), and *glpD*

(BB_0243, glycerol-3-phosphate dehydrogenase). The promoter and transcriptional

boundaries were previously mapped (Grove *et al.*, 2017). The -35 and -10 sites are bolded

and underlined. The transcriptional start site nucleotide is highlighted in blue and topped with

an arrow indicating the direction of transcription. The start of the *glpF* ORF is marked by the

bolded ATG, while the start of the upstream BB_0239 ORF is labeled, and its nucleotides are

italicized. The long arrows indicate the location of primers used to generate a large probe

incorporating the entire intergenic region which is denoted as *glpFKD*(-219) for initial DNA

binding studies.

FIGURE 2 DNA binding by PlzA is c-di-GMP dependent. (**a**) EMSA with increasing

concentrations of PlzA$_{WT}$ showing binding to *glpFKD*(-219) DNA. The mixture was incubated

for 10 min before electrophoresis. (**b**) The same EMSA mixture as in (**a**) but with a final

concentration of 2.5 ng/µL of the nonspecific competitor poly-dI-dC. Protein was incubated

with poly-dI-dC for 5 min prior to the addition of labeled *glpFKD*(-219) probe. (**c**) The same

EMSA mixture as in (**b**) but supplemented with 100 µM c-di-GMP. Protein was incubated with

c-di-GMP for 5 min prior to the addition of any nucleic acids. Arrows indicate PlzA- *glpFKD*(-

219) complexes. Lane 1 in (**a-c**) is the probe only control, containing 50 nM of probe. Protein

concentrations are as follows in each subsequent lane: Lane 2: 0.5 µM, Lane 3: 1 µM, Lane

4: 2.5 µM, Lane 5: 5 µM, Lane 6: 7.5 µM, and Lane 7: 10 µM. (**d**), (**e**), (**f**) EMSA experiments

as in (**a**), (**b**), and (**c**), respectively, but performed with the mutant PlzA$_{RD-RD}$ protein. All

conditions were performed identically as with PlzA$_{WT}$. Each panel is a representative gel from

separate EMSAs which were performed in triplicate (PlzA$_{WT}$) or duplicate (PlzA$_{RD-RD}$).

FIGURE 3 PlzA recognizes *glpFKD*(-219) and not pCR 2.1 sequences. (**a**) A schematic of the

unlabeled DNA competitors derived from the empty pCR 2.1 vector backbone or the vector

containing the *glpFKD*(-219) sequence. A labeled M13R and an unlabeled M13F primer were

used to amplify the target DNAs from the respective plasmids to generate competitors. These

competitors were then used as a control in competition EMSAs performed with labeled

glpFKD(-219) DNA probe. (**b**) Lane 1 and lane 2 are probe only and probe + PlzA$_{WT}$ controls

respectively. All lanes contained a c-di-GMP concentration of 100 µM, a final poly-dI-dC

concentration of 2.5 ng/µL, and a labeled *glpFKD*(-219) probe concentration of 50 nM. Lanes

2-8 contained a constant PlzA$_{WT}$ protein concentration of 1.5 µM. The unlabeled pCR 2.1

competitor was titrated into the reactions at increasing molar excess relative to labeled probe:

Lane 3-300 nM (6x), lane 4- 500 nM (10x), and lane 5- 1000 nM (20x). Unlabeled *glpFKD*(-

219) competitor was titrated into the reactions at increasing molar excess relative to labeled

probe: Lane 6- 300 nM (6x), lane 7- 500 nM (10x), and lane 8- 600 nM (12x). The free arrow

indicates PlzA-*glpFKD*(-219) complexes. The pCR 2.1 competitor could not compete away the

PlzA-*glpFKD*(-219) complex at any tested concentration. Unlabeled *glpFKD*(-219) competitor

was able to compete at all tested concentrations as low as 6x molar excess.

FIGURE 4 Detection of c-di-GMP in recombinant PlzA$_{WT}$ and PlzA$_{RD-RD}$ by LC-MS/MS.

Recombinant PlzA$_{WT}$ and PlzA$_{RD-RD}$ were sent off for mass spectrometry to detect c-di-GMP.

Extracted ion chromatograms of the LC-MS/MS analysis for detection of c-di-GMP are

displayed. Chromatograms depict the control c-di-GMP standard at 1 µM, recombinant

PlzA$_{WT}$, and recombinant PlzA$_{RD-RD}$. The LC-MS/MS analysis reveals PlzA$_{WT}$ co-purified with

c-di-GMP, but PlzA$_{RD-RD}$ did not. The concentration of c-di-GMP in recombinant PlzA$_{WT}$ was

determined from a standard curve of c-di-GMP. PlzA$_{WT}$ co-purified with 99.29 nM of c-di-GMP

and approximately 0.3% of the protein was bound with c-di-GMP. Black lines indicate

chromatograms unrelated to this figure that were removed.

FIGURE 5 Levels of c-di-GMP alter DNA binding affinity of PlzA$_{WT}$. (**a**) A diagram of the

generated DNA probe, designated as *glpFKD*(-7/+35), corresponding to a 42 bp sequence

adjacent to the -10 site and proceeding 35 bp into the 5' UTR. An RNA probe was made

corresponding to only nucleotides that are within the 5 'UTR which was designated as

glpFKD(UTR) RNA. Probes were conjugated to a fluorescent molecule for the detection of

binding in EMSAs. (**b**) EMSAs were performed with increasing protein concentrations of

PlzA$_{WT,}$ and 10 nM fluorescently labeled probe at varying c-di-GMP concentrations (0,

equimolar, 10, 25, 50, or 100 µM). For each EMSA shown, the indicated c-di-GMP

concentration was held constant in all lanes. For each EMSA shown, lanes 1 and 7 are probe-

only controls. Protein concentrations were as follows: Lane 2- 1 µM, Lane 3- 2.5 µM, Lane 4-

3.5 µM, Lane 5- 5 µM, Lane 6- 7.5 µM. A final concentration of 2.5 ng/µL of the nonspecific

competitor poly-dI-dC was added to each lane. Arrows indicate higher-order complexes

formed as protein concentration increases. A representative gel for each tested c-di-GMP

concentration is shown from triplicate EMSAs. (**c**) Triplicate EMSAs were quantitated by

densitometry to determine the K$_{d(app)}$ of the PlzA$_{WT}$-*glpFKD*(-7/35+) interaction by nonlinear

regression analysis using the one-site specific binding setting in GraphPad Prism. Errors bars

represent one standard deviation (SD). (**d**) Bar graph displaying the calculated K$_{d(app)}$ ± the

standard error of the mean (SEM). The K$_{d(app)}$ values from three EMSAs were analyzed by a

Welch ANOVA with Dunnett's T3 multiple comparisons test via GraphPad Prism. Adjusted P-

values are indicated as follows: ns= p >0.05, *= p ≤0.05, and ***= p ≤0.001. Comparisons

between 10, 25, 50, and 100 µM c-di-GMP are not indicated on the graph as they were not

significant. (**e**) A similar EMSA experiment was performed as in Figure 5B but with PlzA$_{RD-RD}$

and either 100 (lanes 1-6) or 0 (lanes 7-12) µM of c-di-GMP supplemented. No binding is

observed.

FIGURE 6 PlzA$_{WT}$ also binds single stranded RNA and DNA. (**a**) Gradient EMSAs were

performed with increasing protein concentrations of PlzA$_{WT}$ without c-di-GMP (left panel),

equimolar (middle panel), or 100 µM c-di-GMP (right panel) and 10 nM fluorescently labeled

glpFKD(UTR) RNA probe. Lane 1 is the probe-only control in each EMSA. Lane protein

concentrations were as follows: Lane 2- 1 µM, Lane 3- 2.5 µM, Lane 4- 3.5 µM, Lane 5- 5 µM,

Lane 6- 7.5 µM. (**b**) A similar EMSA experiment was performed with PlzA$_{RD-RD}$ with (lanes 1-

8) or without (lanes 9-15) 100 µM c-di-GMP and 50 nM of *glpFKD*(UTR) RNA probe. Lane 1

is a probe-only control. Protein concentrations were as follows: Lanes 2 and 9- 1 µM, Lanes

3 and 10- 1.25 µM, Lanes 4 and 11- 2 µM, Lanes 5 and 12- 2.5 µM, Lanes 6 and 13- 3.3 µM,

Lanes 7 and 14- 5 µM, Lanes 8 and 15- 10 µM. Riboguard was supplemented in the PlzA$_{RD-}$

$_{RD}$ EMSA mixtures to a final concentration of 4 U/µL. (**c**) Gradient EMSAs were performed with

increasing protein concentrations of PlzA$_{WT}$ with 100 µM c-di-GMP (left panel) or 0 µM c-di-

GMP (right panel) with 10 nM fluorescently labeled ss*glpF* probe. Lanes 1 and 7 are the probe-

only control in each EMSA. Lane protein concentrations were as follows: Lanes 2 and 8- 1

µM, Lanes 3 and 9- 2.5 µM, Lanes 4 and 10- 3.5 µM, Lanes 5 and 11- 5 µM, Lanes 6 and 12-

7.5 µM. Representative gels are shown from triplicate (PlzA$_{WT}$) or duplicate (PlzA$_{RD-RD}$) EMSA

experiments. (**d**) Triplicate EMSAs from (**a**) and (**c**) were quantitated by densitometry to

determine the $K_{d(app)}$ of PlzA$_{WT}$ for the *glpFKD* RNA or ssDNA interactions by nonlinear

regression analysis using the one-site specific binding setting in GraphPad Prism. Errors bars

represent one standard deviation (SD).

FIGURE 7 c-di-GMP significantly impacts PlzA-nucleic acid binding and PlzA prefers dsDNA

and RNA over ssDNA. (**a**) The $K_{d(app)}$ values from triplicate RNA EMSAs at each tested c-di-

GMP concentration in Figure 6A were analyzed by a Welch ANOVA with Dunnett's T3 multiple

comparisons test. (**b**) The $K_{d(app)}$ values from triplicate ssDNA EMSAs at the tested c-di-GMP

concentrations in Figure 6C were analyzed by a Welch's t-test. (**c**) The $K_{d(app)}$ values from

EMSAs with PlzA and each tested nucleic acid at 100 µM c-di-GMP were compared via a

Welch ANOVA with Dunnett's T3 multiple comparisons test. Bar graphs display the calculated

$K_{d(app)s}$ ± the standard error of the mean (SEM). Adjusted P-values are indicated as follows:

ns= p >0.05, *= p ≤0.05, and **= p ≤0.01.

FIGURE 8 The N-terminal domain of PlzA is the nucleic acid binding domain. (**a**) The cloning

strategy used to produce recombinant N (PlzA$_{NTD}$) and C (PlzA$_{CTD}$) terminal PlzA domains.

The *plzA* gene portions corresponding to the N-terminus, *plzA*-NTD, and the C-terminus, *plzA*-

CTD, were generated and cloned into an expression vector to produce recombinant PlzA$_{NTD}$

and PlzA$_{CTD}$ proteins. Gradient EMSAs were performed with increasing protein concentrations

of PlzA$_{NTD}$ (**b, d**) or PlzA$_{CTD}$ (**c, e**) with (lanes 1-6) or without (lanes 7-12) 100 µM c-di-GMP

and 10 nM fluorescently labeled *glpFKD*(-7/35+) dsDNA (**b, c**) or *glpFKD*(UTR) RNA (**d, e**)

probe. Lanes 1 and 7 are probe-only controls. Protein concentrations were as follows: Lanes

2 and 8- 0.5 µM, Lanes 3 and 9- 1 µM, Lanes 4 and 10- 2.5 µM, Lanes 5 and 11- 5 µM, Lanes

6 and 12- 7.5 µM. SUPERase•In RNase inhibitor was supplemented to a final concentration

of 2 U/µL in RNA EMSA mixtures. The asterisk indicates PlzA$_{NTD}$-dsDNA shifts.

Representative gels are shown from triplicate (PlzA$_{NTD}$) or duplicate (PlzA$_{CTD}$) EMSAs. N= 2

or 3 independent experiments.

FIGURE 9 Detection of c-di-GMP in recombinant PlzA$_{NTD}$ and PlzA$_{CTD}$ by LC-MS/MS and

circular dichroism of proteins. (**a**) Recombinant PlzA$_{NTD}$ and PlzA$_{CTD}$ were sent off for mass

spectrometry to detect the presence of c-di-GMP. Extracted ion chromatograms of the LC-

MS/MS analysis are displayed. Chromatograms depict the control c-di-GMP standard at 1 µM,

recombinant PlzA$_{CTD}$, and recombinant PlzA$_{NTD}$. The LC-MS/MS analysis revealed that

PlzA$_{CTD}$ co-purified with c-di-GMP, but PlzA$_{NTD}$ did not. The concentration of c-di-GMP in

recombinant PlzA$_{CTD}$ was determined from a standard curve of c-di-GMP. PlzA$_{CTD}$ co-purified

with 704.29 nM of c-di-GMP and approximately 1.1 % of the protein was bound with c-di-GMP.

The c-di-GMP standard chromatogram shown is the same as in Figure 4. (**b**) Circular

dichroism spectra of recombinant PlzA$_{WT}$, PlzA$_{RD-RD}$, PlzA$_{NTD}$, and PlzA$_{CTD}$ measured in EMSA

buffer (see methods). EMSA buffer alone or plus 50 µM c-di-GMP were also measured as

controls.

FIGURE 10 Computational docking analysis of PlzA with *glp* DNA and RNA. Computational

docking was performed using the HDOCK server with PlzA and either (**a**) the 42 bp *glpFKD*(-

7/+35) dsDNA or (**b**) the *glpFKD*(UTR) RNA sequence. In (**a**) only 26 of 42 bp of the *glpFKD*(-

7/+35) sequence is shown. The dashed lines represent where bases were left out of the figure

to zoom in on the protein-DNA interactions. The dsDNA model was linear, and removal of the

bases has no impact on the interpretation of the data. The top predicted model from each

computation is shown. The HDOCK docking and confidence scores for each model were -

210.60 and 0.77 for PlzA-*glp* DNA, and -280.71 and 0.93 for PlzA-*glp* RNA, respectively. The

protein is rotated 180° from the top to the bottom panels. Select amino acid residues that

interface the nucleic acids are labeled.

FIGURE 11 Model depicting the modulation of PlzA function by levels of c-di-GMP. Apo and

holo-PlzA are hypothesized to have distinct functions in the vertebrate and tick hosts

respectively. Recent studies identified PlzA RNA chaperone activity that is c-di-GMP

independent (Van Gundy *et al.*, 2023). In environments with no c-di-GMP, such as the

vertebrate host, apo-PlzA engages in RNA chaperone activity to modulate gene expression

required for survival in the vertebrate host. The phosphodiesterases PdeA and PdeB are likely

to be highly active upon transmission and into vertebrate infection, lowering c-di-GMP levels

and facilitating apo-PlzA functions (Sultan *et al.*, 2010; Pitzer *et al.*, 2011; Sultan *et al.*, 2011;

Novak *et al.*, 2014; Groshong *et al.*, 2021). Phosphodiesterase activity is likely to occur at all

stages to regulate c-di-GMP levels and indirectly, PlzA function. When Hk1 is activated (signal

unknown), Rrp1 is phosphorylated and produces c-di-GMP. Holo-PlzA can also engage in

RNA chaperone activity by annealing RNAs, and potentially interact with RNAP (Grassmann

et al., 2023; Van Gundy *et al.*, 2023). We propose that levels of c-di-GMP impact the specific

genes regulated by PlzA through altering nucleic acid binding affinity. At lower c-di-GMP

levels, few holo-PlzA monomers exist resulting in weak PlzA binding affinity for certain target

genes such as *glpFKD*. As levels of c-di-GMP increase, such as in the tick host or during a

blood meal, more holo-PlzA molecules bind target genes with higher affinity, and PlzA could

potentially multimerize on DNA. At high protein and c-di-GMP concentrations, we hypothesize

release of DNA by holo-PlzA via a yet unknown signal.

FIGURE 1 The *glpFKD* operon, 5' UTR, and designed probe schematic. The glycerol catabolism gene operon (*glpFKD*) contains four genes: *glpF* (BB_0240, glycerol uptake facilitator protein), *glpK* (BB_0241, glycerol kinase), BB_0242 (hypothetical protein), and *glpD* (BB_0243, glycerol-3-phosphate dehydrogenase). The promoter and transcriptional boundaries were previously mapped (Grove et al., 2017). The -35 and -10 sites are bolded and underlined. The transcriptional start site nucleotide is highlighted in blue and topped with an arrow indicating the direction of transcription. The start of the *glpF* ORF is marked by the bolded ATG, while the start of the upstream BB_0239 ORF is labeled, and its nucleotides are italicized. The long arrows indicate the location of primers used to generate a large probe incorporating the entire intergenic region which is denoted as *glpFKD*(-219) for initial DNA binding studies.

FIGURE 2 DNA binding by PlzA is c-di-GMP dependent. (**a**) EMSA with increasing concentrations of PlzA$_{WT}$ showing binding to *glpFKD*(-219) DNA. The mixture was incubated for 10 min before electrophoresis. (**b**) The same EMSA mixture as in (a) but with a final concentration of 2.5 ng/µL of the nonspecific competitor poly-dI-dC. Protein was incubated with poly-dI-dC for 5 min prior to the addition of labeled *glpFKD*(-219) probe. (**c**) The same EMSA mixture as in (**b**) but supplemented with 100 µM c-di-GMP. Protein was incubated with c-di-GMP for 5 min prior to the addition of any nucleic acids. Arrows indicate PlzA- *glpFKD*(-219) complexes. Lane 1 in (**a-c**) is the probe only control, containing 50 nM of probe. Protein concentrations are as follows in each subsequent lane: Lane 2: 0.5 µM, Lane 3: 1 µM, Lane 4: 2.5 µM, Lane 5: 5 µM, Lane 6: 7.5 µM, and Lane 7: 10 µM. (**d**), (**e**), (**f**) EMSA experiments as in (**a**), (**b**), and (**c**), respectively, but performed with the mutant PlzA$_{RD-RD}$ protein. All conditions were performed identically as with PlzA$_{WT}$. Each panel is a representative gel from separate EMSAs which were performed in triplicate (PlzA$_{WT}$) or duplicate (PlzA$_{RD-RD}$).

FIGURE 3 PlzA recognizes *glpFKD*(-219) and not pCR 2.1 sequences. (**a**) A schematic of the unlabeled DNA competitors derived from the empty pCR 2.1 vector backbone or the vector containing the *glpFKD*(-219) sequence. A labeled M13R and an unlabeled M13F primer were used to amplify the target DNAs from the respective plasmids to generate competitors. These competitors were then used as a control in competition EMSAs performed with labeled *glpFKD*(-219) DNA probe. (**b**) Lane 1 and lane 2 are probe only and probe + PlzA_WT controls respectively. All lanes contained a c-di-GMP concentration of 100 μM, a final poly-dI-dC concentration of 2.5 ng/μL, and a labeled *glpFKD*(-219) probe concentration of 50 nM. Lanes 2-8 contained a constant PlzA_WT protein concentration of 1.5 μM. The unlabeled pCR 2.1 competitor was titrated into the reactions at increasing molar excess relative to labeled probe: Lane 3-300 nM (6x), lane 4- 500 nM (10x), and lane 5- 1000 nM (20x). Unlabeled *glpFKD(-219)* competitor was titrated into the reactions at increasing molar excess relative to labeled probe: Lane 6-300 nM (6x), lane 7- 500 nM (10x), and lane 8- 600 nM (12x). The free arrow indicates PlzA-*glpFKD*(-219) complexes. The pCR 2.1 competitor could not compete away the PlzA-*glpFKD*(-219) complex at any tested concentration. Unlabeled *glpFKD*(-219) competitor was able to compete at all tested concentrations as low as 6x molar excess.

FIGURE 4 Detection of c-di-GMP in recombinant PlzA$_{WT}$ and PlzA$_{RD-RD}$ by LC-MS/MS. Recombinant PlzA$_{WT}$ and PlzA$_{RD-RD}$ were sent off for mass spectrometry to detect c-di-GMP. Extracted ion chromatograms of the LC-MS/MS analysis for detection of c-di-GMP are displayed. Chromatograms depict the control c-di-GMP standard at 1 μM, recombinant PlzA$_{WT}$, and recombinant PlzA$_{RD-RD}$. The LC-MS/MS analysis reveals PlzA$_{WT}$ co-purified with c-di-GMP, but PlzA$_{RD-RD}$ did not. The concentration of c-di-GMP in recombinant PlzA$_{WT}$ was determined from a standard curve of c-di-GMP. PlzA$_{WT}$ co-purified with 99.29 nM of c-di-GMP and approximately 0.3% of the protein was bound with c-di-GMP. Black lines indicate chromatograms unrelated to this figure that were removed.

FIGURE 5 Levels of c-di-GMP alter DNA binding affinity of PlzA_WT. (a) A diagram of the generated DNA probe, designated as *glpFKD(-7/+35)*, corresponding to a 42 bp sequence adjacent to the -10 site and proceeding 35 bp into the 5' UTR. An RNA probe was made corresponding to only nucleotides that are within the 5 'UTR which was designated as *glpFKD*(UTR) RNA. Probes were conjugated to a fluorescent molecule for the detection of binding in EMSAs. (b) EMSAs were performed with increasing protein concentrations of PlzA_WT, and 10 nM fluorescently labeled probe at varying c-di-GMP concentrations (0, equimolar, 10, 25, 50, or 100 µM). For each EMSA shown, the indicated c-di-GMP concentration was held constant in all lanes. For each EMSA shown, lanes 1 and 7 are probe-only controls. Protein concentrations were as follows: Lane 2- 1 µM, Lane 3- 2.5 µM, Lane 4- 3.5 µM, Lane 5- 5 µM, Lane 6- 7.5 µM. A final concentration of 2.5 ng/µL of the nonspecific competitor poly-dI-dC was added to each lane. Arrows indicate higher-order complexes formed as protein concentration increases. A representative gel for each tested c-di-GMP concentration is shown from triplicated EMSAs. (c) Triplicate EMSAs were quantitated by densitometry to determine the $K_{d(app)}$ of the PlzA_WT-*glpFKD(-7/35+)* interaction by nonlinear regression analysis using the one-site specific binding setting in GraphPad Prism. Errors bars represent one standard deviation (SD). (d) Bar graph displaying the calculated $K_{d(app)}$ ± the standard error of the mean (SEM). The Kd(app) values from three EMSAs were analyzed by a Welch ANOVA with Dunnett's T3 multiple comparisons test via GraphPad Prism. Adjusted P-values are indicated as follows: ns= $p > 0.05$, *= $p ≤ 0.05$, and ***= $p ≤ 0.001$. Comparisons between 10, 25, 50, and 100 µM c-di-GMP are not indicated on the graph as they were not significant. (e) A similar EMSA experiment was performed as in Figure 5B but with PlzA_RD-RD and either 100 (lanes 1-6) or 0 (lanes 7-12) µM of c-di-GMP supplemented. No binding is observed.

FIGURE 6 PlzA_WT also binds single stranded RNA and DNA. (**a**) Gradient EMSAs were performed with increasing protein concentrations of PlzA_WT without c-di-GMP (left panel), equimolar (middle panel), or 100 µM c-di-GMP (right panel) and 10 nM fluorescently labeled *glpFKD*(UTR) RNA probe. Lane 1 is the probe-only control in each EMSA. Lane protein concentrations were as follows: Lane 2- 1 µM, Lane 3- 2.5 µM, Lane 4- 3.5 µM, Lane 5- 5 µM, Lane 6- 7.5 µM. (**b**) A similar EMSA experiment was performed with PlzA_RD-RD with (lanes 1-8) or without (lanes 9-15) 100 µM c-di-GMP and 50 nM of *glpFKD*(UTR) RNA probe. Lane 1 is a probe-only control. Protein concentrations were as follows: Lanes 2 and 9- 1 µM, Lanes 3 and 10- 1.25 µM, Lanes 4 and 11- 2 µM, Lanes 5 and 12- 2.5 µM, Lanes 6 and 13- 3.3 µM, Lanes 7 and 14- 5 µM, Lanes 8 and 15- 10 µM. Riboguard was supplemented in the PlzA_RD-RD EMSA mixtures to a final concentration of 4 U/µL. (**c**) Gradient EMSAs were performed with increasing protein concentrations of PlzA_WT with 100 µM c-di-GMP (left panel) or 0 µM c-di-GMP (right panel) with 10 nM fluorescently labeled ss*glpF* probe. Lanes 1 and 7 are the probe-only control in each EMSA. Lane protein concentrations were as follows: Lanes 2 and 8- 1 µM, Lanes 3 and 9- 2.5 µM, Lanes 4 and 10- 3.5 µM, Lanes 5 and 11- 5 µM, Lanes 6 and 12- 7.5 µM. Representative gels are shown from triplicated (PlzA_WT) or duplicated (PlzA_RD-RD) EMSA experiments. (**d**) Triplicated EMSAs from (**a**) and (**c**) were quantitated by densitometry to determine the Kd(app) of PlzA_WT for the *glpFKD* RNA or ssDNA interactions by nonlinear regression analysis using the one-site specific binding setting in GraphPad Prism. Errors bars represent one standard deviation (SD).

FIGURE 7 c-di-GMP significantly impacts PlzA-nucleic acid binding and PlzA prefers dsDNA and RNA over ssDNA. (**a**) The Kd(app) values from triplicate RNA EMSAs at each tested c-di-GMP concentration in Figure 6A were analyzed by a Welch ANOVA with Dunnett's T3 multiple comparisons test. (**b**) The Kd(app) values from triplicate ssDNA EMSAs at the tested c-di-GMP concentrations in Figure 6C were analyzed by a Welch's t-test. (c) The Kd(app) values from EMSAs with PlzA and each tested nucleic acid at 100 µM c-di-GMP were compared via a Welch ANOVA with Dunnett's T3 multiple comparisons test. Bar graphs display the calculated Kd(app)s ± the standard error of the mean (SEM). Adjusted P-values are indicated as follows: ns= p >0.05, *= p ≤0.05, and **= p ≤0.01.

Table 1. Binding interaction summary of PlzA$_{WT}$ and the tested nucleic acids at various c-di-GMP concentrations.

glpFKD(-7/+35) dsDNA

c-di-GMP (μM)	K$_{d(app)}$ (μM), 95% C.I. [LL, UL]	B$_{max}$ (% Shifted), 95% C.I. [LL, UL]	Adj. R-squared
100	1.09, [0.97, 1.23]	105.20, [102.20, 108.40]	0.986
50	1.25, [1.08, 1.45]	105.50, [101.50, 109.90]	0.981
25	1.33, [1.10, 1.59]	101.60, [96.58, 107.10]	0.971
10	2.81, [1.98, 4.02]	113.00, [98.79, 132.30]	0.940
Equimolar	4.22, [3.40, 5.26]	115.40, [104.40, 129.10]	0.983
0	7.86, [4.85, 14.06]	94.25, [71.72, 138.80]	0.950

glpFKD(UTR) RNA

c-di-GMP (μM)	K$_{d(app)}$ (μM), 95% C.I. [LL, UL]	B$_{max}$ (% Shifted), 95% C.I. [LL, UL]	Adj. R-squared
100	0.95, [0.73, 1.22]	102.6, [96.76, 109.20]	0.930
Equimolar	4.14, [2.79, 6.30]	106.7, [89.50, 132.60]	0.944
0	7.64, [4.71, 13.66]	87.73, [66.88, 128.80]	0.950

glpF ssDNA

c-di-GMP (μM)	K$_{d(app)}$ (μM), 95% C.I. [LL, UL]	Bmax (% Shifted), 95% C.I. [LL, UL]	Adj. R-squared
100	4.42, [2.94, 6.84]	102.80, [85.32, 129.90]	0.920
0	8.74, [5.67, 14.70]	82.04, [63.53, 116.90]	0.950

FIGURE 8 The N-terminal domain of PlzA is the nucleic acid binding domain. (**a**) The cloning strategy used to produce recombinant N (PlzA$_{NTD}$) and C (PlzA$_{CTD}$) terminal PlzA domains. The plzA gene portions corresponding to the N-terminus, plzA-NTD, and the C-terminus, plzA-CTD, were generated and cloned into an expression vector to produce recombinant PlzA$_{NTD}$ and PlzA$_{CTD}$ proteins. Gradient EMSAs were performed with increasing protein concentrations of PlzA$_{NTD}$ (**b**, **d**) or PlzA$_{CTD}$ (**c**, **e**) with (lanes 1-6) or without (lanes 7-12) 100 µM c-di-GMP and 10 nM fluorescently labeled *glpFKD*(-7/35+) dsDNA (b, c) or *glpFKD*(UTR) RNA (**d**, **e**) probe. Lanes 1 and 7 are probe-only controls. Protein concentrations were as follows: Lanes 2 and 8- 0.5 µM, Lanes 3 and 9- 1 µM, Lanes 4 and 10- 2.5 µM, Lanes 5 and 11- 5 µM, Lanes 6 and 12- 7.5 µM. SUPERase•In RNase inhibitor was supplemented to a final concentration of 2 U/µL in RNA EMSA mixtures. Representative gels are shown from triplicated (PlzA$_{NTD}$) or duplicated (PlzA$_{CTD}$) EMSAs.

FIGURE 9 Detection of c-di-GMP in recombinant PlzA_{NTD} and PlzA_{CTD} by LC-MS/MS and circular dichroism of proteins. (**a**) Recombinant PlzA_{NTD} and PlzA_{CTD} were sent off for mass spectrometry to detect the presence of c-di-GMP. Extracted ion chromatograms of the LC-MS/MS analysis are displayed. Chromatograms depict the control c-di-GMP standard at 1 µM, recombinant PlzA_{CTD}, and recombinant PlzA_{NTD}. The LC-MS/MS analysis revealed that PlzA_{CTD} co-purified with c-di-GMP, but PlzA_{NTD} did not. The concentration of c-di-GMP in recombinant PlzA_{CTD} was determined from a standard curve of c-di-GMP. PlzA_{CTD} co-purified with 704.29 nM of c-di-GMP and approximately 1.1 % of the protein was bound with c-di-GMP. The c-di-GMP standard chromatogram shown is the same as in Figure 4. (**b**) Circular dichroism spectra of recombinant PlzA_{WT}, PlzA_{RD-RD}, PlzA_{NTD}, and PlzA_{CTD} measured in EMSA buffer (see methods). EMSA buffer alone or plus 50 µM c-di-GMP were also measured as controls.

FIGURE 10 Computational docking analysis of PlzA with *glp* DNA and RNA. Computational docking was performed using the HDOCK server with PlzA and either (a) the 42 bp *glpFKD*(-7/+35) dsDNA or (b) the *glpFKD*(UTR) RNA sequence. In (a) only 26 of 42 bp of the *glpFKD*(-7/+35) sequence is shown. The dashed lines represent where bases were left out of the figure to zoom in on the protein-DNA interactions. The dsDNA model was linear, and removal of the bases has no impact on the interpretation of the data. The top predicted model from each computation is shown. The HDOCK docking and confidence scores for each model were -210.60 and 0.77 for PlzA-*glp* DNA and -280.71 and 0.93 for PlzA-*glp* RNA, respectively. The protein is rotated 180° from the top to the bottom panels. Select amino acid residues that interface the nucleic acids are labeled.

FIGURE 11 Model depicting the modulation of PlzA function by levels of c-di-GMP. Apo and holo-PlzA are hypothesized to have distinct functions in the vertebrate and tick hosts respectively. Recent studies identified PlzA RNA chaperone activity that is c-di-GMP independent (Van Gundy et al., 2023). In environments with no c-di-GMP, such as the vertebrate host, apo-PlzA engages in RNA chaperone activity to modulate gene expression required for survival in the vertebrate host. The phosphodiesterases PdeA and PdeB are likely to be highly active upon transmission and into vertebrate infection, lowering c-di-GMP levels and facilitating apo-PlzA functions (Sultan et al., 2010; Pitzer et al., 2011; Sultan et al., 2011; Novak et al., 2014; Groshong et al., 2021). Phosphodiesterase activity is likely to occur at all stages to regulate c-di-GMP levels and indirectly, PlzA function. When Hk1 is activated (signal unknown), Rrp1 is phosphorylated and produces c-di-GMP. Holo-PlzA can also engage in RNA chaperone activity by annealing RNAs, and potentially interact with RNAP (Grassmann et al., 2023; Van Gundy et al., 2023). We propose that levels of c-di-GMP impact the specific genes regulated by PlzA through altering nucleic acid binding affinity. At lower c-di-GMP levels, few holo-PlzA monomers exist resulting in weak PlzA binding affinity for certain target genes such as *glpFKD*. As levels of c-di-GMP increase, such as in the tick host or during a blood meal, more holo-PlzA molecules bind target genes with higher affinity, and PlzA could potentially multimerize on DNA. At high protein and c-di-GMP concentrations, we hypothesize release of DNA by holo-PlzA via a yet unknown signal.

Table 2. Oligonucleotides used in this study.

Target	Name and Sequence (5' → 3')	Size (bp)
Cloning and Sequencing Primers		
plzA	*plzA* F: GGCAGC**CATATG**CTTTTATCTAGAAAAATAAGAGATTATG *plzA* R: GTGGTG**CTCGAG**TTAATTGAAATAATCATGGATCAAC	786
MCS pET28a(+):*plzA*	T7 F: TAATACGACTCACTATAGGG T7 R: TAGTTATTGCTCAGCGGTGG	1026
pCR 2.1 TA clones	M13 F: GTAAAACGACGGCCAG M13 R: CAGGAAACAGCTATGAC	N/A
plzA-NTD	*plzA* 1F: ATGCTTTTATCTAGAAAAATAAGAGATTATG plzA N R: TTACTGATTTTGCCCAAGCTTTAAATC	435
plzA-CTD	*plzA* C F: ATGGATTTAAAGCTTGGGCAA plzA 1R: ATTGAAATAATCATGGATCAACATAGTATAC	378
Site-directed Mutagenesis Primers		
plzA R145-149D	*plzA*-mut F: AATCAGGACATTCATGAGGACATTATTATCGATAAAGATTCTATTAG plzA-mut R: GTCCTCATGAATGTCCTGATTTTGCCCAAGCTTTAAATCAAGAAGCTTTCC	786
EMSA Probes, Competitors, and Primers		
glpFKD(-219)	M13 F: GTAAAACGACGGCCAG M13 R 5'IRD800CW or unlabeled: CAGGAAACAGCTATGAC	654
glpFKD(-7/+35)	*glpF* F IRD800: ATTAAATATAATTTTAATAAGGCTTTTATTAGAAAAATTAAT *glpF* R: ATTAATTTTTCTAATAAAAGCCTTATTAAAATTATATTTAAT	42
ss*glpF*	*glpF* F IRD800: ATTAAATATAATTTTAATAAGGCTTTTATTAGAAAAATTAAT	42
glpFKD(UTR) RNA	*glpF* RNA 5' Alexa488: AUAAUUUUAAUAAGGCUUUUAUUAGAAAAAUUAAU	35
pCR 2.1 competitor	M13 F: GTAAAACGACGGCCAG M13 R: CAGGAAACAGCTATGAC	210
[a] Bolded and underlined nucleotides indicate restriction enzyme sites.		