Inferring condition in wild mammals: body condition indices confer no benefit over measuring body mass across ecological contexts

Andrea E. Wishart[1,*], Adriana L. Guerrero-Chacón[1], Rebecca Smith[1], Deborah M. Hawkshaw[1], Andrew G. McAdam[2], Ben Dantzer[3,4], Stan Boutin[5], Jeffrey E. Lane[1]

 *Author for correspondence: andrea.wishart@usask.ca

[1] Department of Biology, University of Saskatchewan, Saskatoon, SK, S7N 5E2, Canada

[2] Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, Colorado, USA

[3] Department of Psychology, University of Michigan, Ann Arbor, Michigan 48109-1043, USA

[4] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan 48109-1043, USA

[5] Department of Biological Sciences, University of Alberta, Edmonton, Alberta T6G 2E9, Canada

The utility of body condition indices to approximate energetically significant tissues (fat and lean mass) varies with a species' natural history, and when they are measured during the annual cycle.

Author Contributions. AEW designed the study, collected data, performed data and statistical analysis, and drafted the manuscript. JEL participated in the design. All authors contributed to data collection, provided valuable discussion and contributions to the writing of the manuscript, and gave final approval for publication.

Abstract

Many studies assume that it is beneficial for individuals of a species to be heavier, or have a

higher body condition index (BCI), without accounting for the physiological relevance of

variation in the composition of different body tissues. We hypothesized that the relationship

between BCI and masses of physiologically important tissues (fat and lean) would be conditional

on annual patterns of energy acquisition and expenditure. We studied three species with

contrasting ecologies in their respective natural ranges: an obligate hibernator (Columbian

ground squirrel, *Urocitellus columbianus*), a facultative hibernator (black-tailed prairie dog,

Cynomys ludovicianus), and a food-caching non-hibernator (North American red squirrel,

Tamiasciurus hudsonicus). We measured fat and lean mass in adults of both sexes using

quantitative magnetic resonance (QMR). We measured body mass and two measures of skeletal

structure (zygomatic width and right hind foot length) to develop sex- and species-specific BCIs,

and tested the utility of BCI to predict body composition in each species. Body condition indices

were more consistently, and more strongly correlated, with lean mass than fat mass. The indices

were most positively correlated with fat when fat was expected to be very high (pre-hibernation

prairie dogs). In all cases, however, BCI was never better than body mass alone in predicting fat

or lean mass. While the accuracy of BCI in estimating fat varied across the natural histories and

annual energetic patterns of the species considered, measuring body mass alone was as effective,

or superior in capturing sufficient variation in fat and lean in most cases.

43
Key words: body composition, condition, energetics, fat, food-caching, hibernation, lean,

phenology

Introduction

Body mass is among the most frequently measured physical traits of organisms, because

it often accounts for much of the observed variation in other traits of interest of the organism

(e.g., fecundity; Myers and Master 1983) as well as ecological processes (e.g., trophic structures;

Woodward et al. 2005). However, the mechanisms underlying these associations often may

depend on the different tissue components comprising body mass, rather than mass itself. For

processes in which energy stores or reserves (see Lindström and Piersma 1993 for distinction)

are the currency of interest, the non-structural components of body mass that represent

metabolizable tissues (fat and lean mass; Krebs and Singleton 1993) are likely more relevant

than size or mass alone. For example, relatively high fat stores facilitate successful migration

(Bairlein 2002) and hibernation (Humphries et al. 2003), while a higher proportion of lean mass

can enhance athletic performance (e.g., takeoff velocity in domestic cats; Harris and Steudel

2002). Understanding relationships between mass and the components of body mass may thus

provide greater insight into the mechanisms linking body mass to performance.

A given body mass may be distributed across structural elements of different sizes. Body

condition indices (BCIs) attempt to control for this by considering total body mass relative to

structural size (e.g., a linear skeletal measurement such as body length, tarsus length, foot length,

or a combination thereof; Jakob et al. 1996; Green 2001; Schulte-Hostedde et al. 2001), often to

give some indication of an individual's total lipid stores as a proxy for 'condition'. Typically,

individuals with relatively higher body mass:skeletal size ratios are considered to be in 'better'

condition than individuals with lower ratios, as the former are assumed to have more energetic

stores than individuals of similar structural size, but with a lower metabolizable fraction

(Schulte-Hostedde et al. 2001). Accurate measures of 'condition' in this energetic sense are

widely applicable to scenarios such as livestock breeding programs (e.g., maximizing meat or

offspring production), conservation biology (e.g., measuring responses to habitat degradation;

Stevenson and Woods 2006; Wikelski and Cooke 2006), and in exploring evolutionary and

ecological patterns (e.g., condition-dependent dispersal; Bonte and De La Peña 2009 and

migration; Andersen et al. 2000). However, because fat and lean mass have very different

energetic storage values (39.6 kJ/g for dry fat, in contrast to 18.1 kJ/g, for dry lean mass;

McGilvery 1983), and different functions, metrics that collapse fat and lean into one

energetically related fraction can obscure mechanistic relationships between physiology and

performance outcomes. Because metabolic rate scales with body size to a lower power than does

fat mass, controlling for body size in energetic studies is a logical solution; however, these

relationships can vary substantially within and between populations (Glazier 2005). Applications

of BCI therefore require a closer consideration of what physiological component of 'condition'

BCIs are describing (e.g., lean versus fat mass; Schulte-Hostedde et al. 2001).

The dynamic nature of body condition across time (Krebs and Singleton 1993) demands

that its correlates and consequences be considered within the context of a species' natural history

(Molnár et al. 2009). The composition of a body in 'good condition' in an adaptive sense (i.e.,

that which is associated with increased survival and/or fitness, *sensu* Wilson and Nussey 2010),

should therefore be expected to vary over time depending on circannual energetics. Fat reserves

are an important source of metabolizable energy (Jenni and Jenni-Eiermann 1998) that fuel

organisms through energetically-expensive behaviours or during periods of energetic shortfall.

For example, individuals with higher fat fractions entering hibernation are more likely to survive

over winter and breed successfully the following spring (Boyer and Barnes 1999). Fatter is not

always better though: during the active season, carrying more body mass can decrease running

speed (Trombulak 1989) and alter circulating hormones (Taylor et al. 1982). Arboreal species

may trade off fat stores for locomotion (Dittus 2013), and for species that primarily store energy

off-body as external food caches, fat reserves may not confer the same advantages during

resource-scarce seasons as they may for species that store energy exclusively on-body.

Furthermore, a higher lean fraction may be associated with increased activity, as shown in

captive rats (*Rattus norvegicus*; Swallow et al. 2010). The body composition associated with

better performance is therefore likely to be highly dependent on natural history, as well as on the

various activities an animal undertakes throughout the year (Wells et al. 2019).

Body mass and structural size can be measured from live animals, giving BCIs an

advantage over chemical composition analyses that require lethal sampling (Reynolds and Kunz

2001), especially in studies where repeated measures within individuals are valuable. In many

species, BCIs correlate well with chemical quantification of fat mass in vertebrates including

birds (Chang and Wiebe 2016) and reptiles (Weatherhead and Brown 1996), and invertebrates

including arthropods (Jakob et al. 1996; Moya-Laraño et al. 2008; Kelly et al. 2014). However,

BCIs are not without drawbacks. Acquiring the structural size component(s) of the index can be

challenging. Krebs and Singleton (1993) warned of low repeatability attributable to measurement

bias across different observers (although this can be greatly reduced by taking replicate

measurements; Blackwell et al. 2006). While estimating lengths of long bones yields accurate

estimates when measured from museum specimens (e.g., Dobson 1992), long bones can be

challenging to measure on live, unanesthetized animals in field conditions (Green 2001). Even

under anesthesia, measurement error can persist (e.g., Martin et al. 2013). Because skeletal

morphology often reflects natural history, as selection favours certain shapes for certain lifestyles

(e.g., arboreal vs. fossorial, e.g. Shimer 1903), the body component of interest may not be

accurately reflected by BCI if the linear measurement(s) of size does not correlate well with

overall body size. Furthermore, although BCI can correlate more strongly with lean mass rather

than fat mass (Schulte-Hostedde et al. 2001), the efficacy of BCI in predicting the masses of

body components may be no more effective than prediction through body mass alone (e.g., body

mass predicts fat stores in bats just as well as BCI; McGuire et al. 2018, Wells et al. 2019).

Finally, since BCI estimates the non-skeletal component of body composition in a general sense,

its interpretation and therefore relevance may differ depending on the energetic aspects of the

natural history of the focal species (e.g., for a fat-storing hibernator versus a food-caching non-

hibernator). We thus expect the correlation between BCIs and body composition (i.e., fat and

lean mass) to vary both across species and timing in annual cycles, reflecting the dynamic nature

of energy budgets in seasonally dependent activities.

128   We studied wild populations of three different mammal species to evaluate concordance

between BCIs and the mass of fat and lean tissue, and determined whether these relationships

differ according to the species' primary energy storage mode (e.g., on-body fat storage vs. off-

body food caching). We hypothesized that the relationship between BCI and body composition

depends on both the extent to which the species relies on on-body energy stores (i.e., fat) for

overwinter survival, and with the expected energetic balance within a stage of the annual cycle

(i.e., season).

135   We selected three species in the family Sciuridae which differ in patterns of energy

storage and metabolic demands: North American red squirrels (*Tamiasciurus hudsonicus*,

hereafter, red squirrels), black-tailed prairie dogs (*Cynomys ludovicianus*, hereafter prairie dogs),

and Columbian ground squirrels (*Urocitellus columbianus*, hereafter ground squirrels). We also

compare ground squirrels before and after hibernation to characterize within-species shifts in

energy stores. Red squirrels harvest and cache conifer cones in a central larder ('midden'; Smith

1968) from late summer through autumn (Fletcher et al. 2010) and weigh ~230-250 g on average

as adults (Boutin and Larsen 1993). Red squirrels remain euthermic throughout winter without

using torpor (Brigham and Geiser 2012), but are able to maintain low levels of energy

expenditure through behavioural adjustments (Humphries et al. 2005) and the use of well

insulated nests (Studd et al. 2016). They are not known to gain significant amounts of fat prior to

winter, during which they rely on cached resources for energy (Wishart 2023).

Prairie dogs can weigh up to 1710 g, but show high within- and between-individual

variation in body mass (Kusch et al. 2021). Throughout most of their range, prairie dogs are

active throughout the winter, however they are capable of hibernation (Lehmer et al. 2006). In

southwestern Saskatchewan, where our study population is located at the northern edge of the

species distribution, they hibernate for ~4 months during winter (Gummer 2005). Prairie dogs of

both sexes increase overall body mass, and fat mass specifically, leading up to winter (Lehmer

and Van Horne 2001; Kusch et al. 2021).

Ground squirrel body mass varies substantially across their active season (~400 g at

emergence from hibernation in spring, to up to ~700 g prior to immergence in late summer;

(Boag and Murie 1981, Dobson et al. 1992). These obligate hibernators are notable for their short

active season (~4 months) and extended time spent in hibernation (~8 months) each year

(Dobson et al. 1992). Ground squirrels reach lower minimum body temperatures (0 °C) and have

longer torpor bout durations (~390 hours for males; Young 1990) than prairie dogs.

We 1) characterized morphology and body composition of these three species in context

of their seasonal energetic demands, 2) evaluated the correlation of BCI with body composition

variables (lean and fat mass), and 3) determined whether BCI confers any benefit relative to

body mass in predicting body composition variables (i.e., whether the correlation between BCI

fat or lean mass was higher than between body mass fat or lean mass. We expected hibernators

(ground squirrels and prairie dogs) to have the highest fat fraction during the pre-winter season,

given the importance of on-body energetic reserves to sustain hibernation, compared to non-

hibernating red squirrels which rely mostly on hoarded food, and compared to post-emergence

ground squirrels in spring who have metabolized fat stores overwinter (Fletcher et al. 2013).

Because fat stores are expected to represent the bulk of the pre-hibernation weight gain in the

lead up to winter, we also expected high concordance between BCI and fat mass in pre-winter

hibernators. Conversely, we expected the lowest concordance between BCI and fat mass to be in

red squirrels, as lean mass is likely to be more important to sustained caching activity (Fletcher

et al. 2015).

Materials and Methods

Study sites and population monitoring

We sampled free-ranging, non-breeding adults from populations within the northern

regions of their respective ranges in Canada: red squirrels in the southwest Yukon (61° N, 138°

W, ~ 850 m above sea level (a.s.l.)), prairie dogs in Grasslands National Park, Saskatchewan

(49° N, 107° W, ~770 m a.s.l.), and ground squirrels in Sheep River Provincial Park, Alberta

(50° N, 114° W, ~ 1500 m a.s.l.). For all populations, we collected data through live-trapping.

All populations had been monitored for at least one year prior to data collection for the present

study. For detailed descriptions of population and reproductive monitoring, see McAdam et al.

(2007) and Dantzer et al. (2020) for red squirrels, Kusch et al. (2020) for prairie dogs, and Lane

et al. (2019) for ground squirrels. Briefly, all individuals received permanent uniquely marked

ear tags (National Band and Tag Company, Newport, KY) upon first trapping. We included only

adults (individuals one year of age or older) to minimize effects of skeletal growth dynamics. For

ground squirrels and prairie dogs, we knew ages for all individuals captured at first emergence

following birth. For red squirrels and some older ground squirrels and prairie dogs, exact ages

were not known, but we could confidently remove young-of-the-year based on size and

breeding/nipple status on first trapping (red squirrels excluded if under 150 g, ground squirrels

excluded if under 400 g, prairie dogs excluded if under 800 g; for all species, females also

excluded if they had small pink nipples at first trapping). We excluded all pregnant (assessed by

abdominal palpations) and lactating (assessed by milk expression) females to remove variance

related to maternal investment in offspring. Sample sizes for each sex/species/season group are

summarized in Table 1.

Morphometric measurements

We measured body mass and size for all live-trapped individuals (Table 1). We weighed

each prairie dog to the nearest 5 g using a Pesola spring scale (Pesola AG, Baar, Switzerland),

and weighed red squirrels and ground squirrels to the nearest 1 g on an electronic balance (Ohaus

C5 Series 2000 g). We measured zygomatic arch width ('ZW') to the nearest millimeter using

calipers (analogue for red squirrels and prairie dogs; digital for ground squirrels). We measured

right hind foot ('RHF') length from heel to longest toe (excluding claw) to the nearest millimeter

using a ruler fit with a heelstop at 0 mm. We measured both ZW and RHF three times per

handling, and used the mean value for analyses.

We took these measurements during the same handling occurrence as body composition

analyses for all red squirrels, all prairie dogs except one, and most ground squirrels. For the

remaining prairie dog and ground squirrels, we used skeletal measurements taken on the nearest

date to when composition analyses were completed. The prairie dog was an adult and unlikely to

be growing, so we used measurements taken 81 days prior. The median interval between skeletal

measurements and composition scans for ground squirrels was 0 days (range: 0-120). We

excluded data from ground squirrels that were younger than 3 years of age if their skeletal

measurements were taken more than two weeks before or after the date of body composition and

mass measurements, as younger squirrels may still be growing structurally (Dobson 1992). A

single yearling (female) remained in the spring dataset, so we removed all yearling ground

squirrels.

Body composition

We measured pre-winter body composition of red squirrels between late-September and

mid-October in 2018 and 2019, prairie dogs in late October 2018 (due to variable torpor patterns

in prairie dogs, exact immergence date into hibernation was not known), and ground squirrels

between late-July and late-August 2019 within a week of immergence for most individuals

(median = 3.5 days pre-immergence; range = 0-15). We measured spring body composition of

ground squirrels for which we were confident were captured within a few days of emergence

from hibernation (median = 1 day post-emergence; range = 0-3) between mid-April and early

May 2019. Most individual ground squirrels were measured in one season only, but 21

individuals were measured in both seasons. As we do not expect ground squirrels to change

skeletal size substantially after they have reached maturity, we recorded morphometric data once

per individual.

We used a quantitative magnetic resonance (QMR) body composition analyzer

(EchoMRI-1600, Echo Medical Systems, Houston, TX) to measure absolute lean and fat mass

(g). This technology provides a relatively non-invasive method of estimating body composition

precisely, and accurately (Tinsley et al. 2004). The remaining components not captured as fat

and lean mass are free water and skeletal mass (McGuire and Guglielmo 2010). Quantities

measured using QMR correlate well with carcass-derived quantities for numerous species across

multiple taxa, most relevant here being rodents, including laboratory rats (*Rattus norvegicus*

domestica; Johnson et al. 2009) and house mice (*Mus musculus domesticus*; Jones et al. 2009).

The QMR approach allows for repeated measures of live animals, both awake and sedated

(Tinsley et al. 2004; McGuire and Guglielmo 2010; Zanghi et al. 2013a, b).

We housed our QMR system in a custom-designed trailer to enable transportation to each

study site. The trailer was climate-controlled to stabilize the temperature, as the magnet within

the QMR system is temperature-sensitive. We targeted an ambient temperature of 21 °C as per

manufacturer recommendations, although field conditions widened the range of stabilized

temperatures to $\pm$ 7 °C. We calibrated the system daily to a 943 g canola oil standard at the

stabilized temperature, which was held for at least five hours prior to scanning animals

(following Guglielmo et al. 2011). Our QMR system was outfitted with an additional antenna to

measure animals from 100 g up to 1600 g to accommodate the range of body masses of the three

species. Details of similar systems are described elsewhere (McGuire and Guglielmo 2010), and

we followed similar protocols here. We live-trapped squirrels in the field, and transported them

to the trailer. We placed each squirrel in a clear plexiglass holding tube with perforations to

allow ample airflow to the animal, then inserted the tube into the QMR chamber. We recorded

body composition through a minimum of two scans, reporting the average values for each

individual. In 2019, we administered a mild sedative via thigh intramuscular injection to red

squirrels prior to scanning and collecting morphometric data (100 µg/kg of dexmedetomidine,

reversed by 1 mg/kg atipamezole) to minimize stress and movement during scans. Sedation was

not necessary for the semi-fossorial prairie dogs or ground squirrels, who remained still, and

even fell asleep in the chamber.

Calculating body condition indices (BCIs)

We retained individuals in the dataset for which we had measurements for all of the

following: ZW, RHF, body mass, and body composition. Within each species, we calculated

coefficients of variation for all variables, and analyzed relationships among variables. While

principal components analysis has been used previously as a general measure of structural size to

derive BCI (Schulte-Hostedde et al. 2005), we determined that it was not appropriate for our

dataset because correlation coefficients between ZW and RHF were not always positive

(Supplementary Fig. S1). Instead, we selected the single skeletal measure that had the greatest

coefficient of variation (CV) to generate a residual index (Jakob et al. 1996). Red squirrels

showed a negative, but non-significant, relationship between RHF and ZW (Supplementary

Table S1) so we chose to use ZW to generate the BCI for this species ('ZW index'). The

relationships between RHF and ZW appeared to differ in direction between male and female

prairie dogs; however, neither correlation was significant, so we generated a ZW index for this

species. In ground squirrels, only data for females pre-winter showed a significant (positive)

relationship. Females in spring, and males in both seasons, showed no significant relationship

between RHF and ZW. We therefore used RHF to generate the single-metric BCI ('RHF index')

for ground squirrels.

To calculate each BCI, we regressed either RHF (log transformed and standardized to

mean of zero and unit variance) alone on body mass (log transformed and standardized as above;

generating the RHF index), or ZW (log transformed and standardized as above) alone on body

mass (log transformed and standardized as above; generating the ZW index). Residual plots are

shown in Supplementary Fig. S2. We calculated BCIs within season (pre-winter or spring) and

within sex for each species using separate regressions. Because some CV values within

species/seasons were similar for RHF and ZW, we also ran models using the BCI derived from

the alternate skeletal measurement (Supplementary Online Material). We indicate in the results

when results differed from the primary BCI model.

In addition to the residual-derived BCIs described above, we also calculated the scaled

mass index (SMI; Peig and Green 2009) because it accounts for scaling relationships of linear

measurements (e.g., hind foot length), tissue components, and overall body size. We then ran

parallel statistical analyses as described below, substituting SMI values for BCI values in all

models (see Supplementary Table S11). We note in the Results when the relationships differed

between SMI and BCI.

Statistical analyses

We performed all analyses in R (v.4.0.3, R Core Team (2020)). We compared

morphological measures between the sexes within each species (and within season for ground

squirrels) using a Welch's two-sample t-test. To assess the efficacy of the selected BCI for each

species in predicting body composition variables, we modeled, separately for each species (and

for ground squirrels, separately for each season), linear models for fat mass (g) and lean mass

(g), each predicted by the selected BCI interacting with sex. We also modeled fat and lean mass

predicted by body mass interacting with sex. We compared the fit of the BCI model and body

mass model for each component for each species using Akaike's information criterion adjusted

for small sample size (Burnham and Anderson 2002), confirming no missing data. We

considered the model with the lowest AIC to be the best fit for explaining the variation in the

response variable, among candidate models. Models that were different by >2 AICc were

considered to be different from one another. To compare the utility of BCI to predict fat and lean

mass across species, we fit a linear model for data from all three species (including both seasons

for ground squirrels) together. For fat and lean mass separately, we defined an interaction term

between species-specific BCI and species as the independent variable. We fit a similar model set

with body mass (scaled within species) instead of BCI.

Results

Morphological measurements

Males were significantly heavier than females in all three species: males were 3.3% heavier

in red squirrels, 9.6% heavier in prairie dogs, 27.9% heavier in pre-winter ground squirrels, and

26.6% heavier in spring ground squirrels (Table 1). Right hind foot length was different between

sexes only for ground squirrels in spring, with male RHF 1.6 mm longer than in females.

Zygomatic width was significantly larger in male prairie dogs and ground squirrels than females,

but similar between red squirrel males and females.

Body composition

In autumn, red squirrels showed the lowest percent body fat, with 2.4% in males and

2.9% in females (Table 1) while prairie dogs had the highest percent body fat, with 30.9% and

35.4% in males and females, respectively. Ground squirrels were the only species to have

statistically significant sex differences in fat (in both seasons). They showed their highest body

fat levels pre-winter before entering hibernation (25.0% for males, 20.8% for females, t = -7.48,

d.f. = 26.9, $p < 0.001$), and lowest in spring (12.4% fat for males, 7.7% fat for females, t = -7.97,

d.f. = 58.5, $p < 0.001$).

Relationship between BCI, body mass, and body composition

The relationships between BCI/body mass and fat/lean mass was positive in almost all

cases (Figures 1-4; the exception being for male prairie dog BCI and lean mass, Figure 2b;

Supplementary Tables S3-6). For red squirrels, the BCI and body mass models performed

similarly in predicting fat, but correlations with fat were weak (adjusted $R^2 < 0.22$) for both BCI

and mass (Supplementary Table S3, Figure 1). Red squirrel lean mass was better predicted by

both BCI and mass models than fat mass, with the body mass model providing a better fit. This

pattern held for the alternate RHF BCI models (Supplementary Table S7). For prairie dogs, both

ZW BCI and body mass predicted fat well, but the body mass model was a better fit for both fat

and lean mass (Supplementary Table S4, Figure 2). For pre-winter ground squirrels, the two

models for fat were indistinguishable based on AICc and had similar correlation strengths

(Supplementary Table S5, Figure 3). The body mass model was a better fit for lean mass. This

pattern held for spring, with BCI and body mass models being similar for fat, and the body mass

model providing a better fit for lean mass (Supplementary Table S6, Figure 4). Patterns did not

change when using ZW BCI instead (Supplementary Tables S7-8). Additionally, for ground

squirrels, the variance in fat explained by the BCI and body mass models was higher pre-winter

(adjusted $R^2 = 0.88$) compared to the spring (adjusted $R^2 = 0.69$). The only comparison in which

the pattern for scaled mass index (SMI) differed from the BCI model sets was for red squirrel fat:

this was also the only instance when the SMI model was a better fit than the body mass model

(adjusted $R^2 = 0.25$; Supplementary Table S11). Otherwise, the body mass model outperformed

the SMI model in all parallel analyses.

In the models including data from all species (and for ground squirrels, both seasons),

the strength of the correlations in both BCI and body mass model sets were strong (adjusted $R^2 >$

0.9 for all) and similar within each component. However, body mass models were consistently a

better fit than the BCI models (ΔAICc = 26.7 for fat, 18 for lean mass, Figure 5, Supplementary

Table S12).

Discussion

We demonstrate that the utility of body condition indices in predicting physiologically

relevant tissues, specifically fat, is conditional on the expected energetic state of a species given

ecological and natural history considerations. If BCI is a reliable indicator of 'condition' as it

relates to on-body energy stores, it should be strongly correlated with fat mass. However, when

individuals were expected to be in a leaner state (e.g., food-caching red squirrels, spring ground

squirrels), correlations between BCI and fat were low to moderate. When individuals were

expected to have higher fat stores (e.g., pre-hibernation prairie dogs and ground squirrels), BCI

models predicted fat well, following similar patterns in seabirds (Jacobs et al. 2012).

Furthermore, the relationship between BCI and fat varied among species (and season, for ground

squirrels). This contingency creates a problem for using BCI to infer size of masses of fat;

specifically, the accuracy of the tool differs along the gradient of variation it is being used to

infer. In nearly all cases, BCI was positively correlated with lean mass. Because fat and lean

mass have significantly different energetic values (McGilvery 1983, Jenni and Jenni-Eiermann

1998), interpretations of BCI as they relate to metabolizable energy should take into account

species-specific natural history and annual energetic patterns. For example, some red squirrels

that would have ranked as lower condition based on BCI had nearly twice as much fat as some

individuals that had higher BCI values. Furthermore, models fit with body mass were almost

always more highly correlated with fat and lean mass than models with BCI, both within and

across species.

The selection of these three species, and the time of year at which they were studied,

provides insight into how body composition manifests in BCI when individuals are in a peak

positive energy balance after accumulating surplus energy to sustain them through upcoming

energetic shortfalls, and when they are expected to have depleted much of that accumulated

energy. By investigating these relationships at the ends of the continuum of energetic states that

organisms may experience throughout the year, we demonstrate that interpretation of 'condition'

indices should consider seasonal variation in energetic demands. For example, red squirrels

primarily store energy as cached food, so were expected to carry little fat. In comparison, we

expected prairie dogs to have high fat stores to sustain them through inefficient hibernation

(Gummer 2005; Hawkshaw 2022), and indeed they had the highest percent body fat of all three

species. The intermediate ground squirrels were the only species to show sex differences in fat,

likely arising from asymmetry in reproductive phenologies between males and females (males

arouse from torpor earlier to undergo spermatogenesis, relying on stored energy, while females

arouse later and have access to more food in the environment to support the cost of lactation,

Broussard et al. 2005). These results illustrate that the relative importance of fat and lean mass is

likely to vary with seasonal activities interacting with the natural history of the organisms.

Almost every relationship between either predictor variable (BCI or body mass) and body

component (fat or lean mass) was positive, save for male prairie dog lean mass. Echoing Schulte-

Hostedde et al. (2001), BCIs are capturing variation in both components; however, we did not

find BCIs to discriminate between the independent variation in each component. We found that

in general, lean mass had less variation around the line of best fit than fat mass, reflective of

previous studies on small-non hibernating mammals that also found that BCIs tend to be more

effective in predicting lean dry mass and water as compared to fat mass (Schulte-Hostedde et al.

2001; Tidhar and Speakman 2007). Given the dynamic nature of body composition in

hibernating species, inferring fat levels from BCIs can be particularly difficult since many

studies that use residual-derived BCIs to assess fat assume that lean mass scales with body size,

and that fat mass varies with condition (McGuire et al. 2018).

In some contexts, like large scale comparisons across species, body mass or BCIs could

be a rough proxy for both fat and lean mass. However, for comparisons among individuals or

within individuals of the same species, or research questions that require the precise estimation

of energy derived from the changes in body mass/BCI, a more precise technique like QMR is

needed (e.g., to estimate energy drawn from either fat lean mass during hibernation, Mejías et al.

2022; or migration, McGuire et al., 2022; Kelsey and Bairlein, 2019).

We have shown that the utility of body mass and/or BCI to describe energy-relevant

components is conditional on natural history and annual energetic cycles. This demands not only

within-species validation if BCI is to be used, but also careful consideration of how these

relationships may change across seasons or energetic states. If BCIs are only predictive of fat in

the upper end of the range of variation in fat stores, the value of BCI in comparing across

individuals breaks down, especially across individuals with relatively lower fat masses. As

shown within Columbian ground squirrels across two seasons, BCI (and the associated

challenges and handling stress involved in obtaining linear measurements of skeletal size can

impart on animal subjects) is not uniformly informative, therefore bringing into question its

utility. When balancing animal welfare, time, and research budgets with desired outcome of

informative data, we advise researchers to consider whether more precise measurements of body

composition are needed, or if simply measuring body mass is sufficient to capture the variation

of interest.

Ultimately, this study strengthens the case for using body mass as a covariate to capture

general variation in fat and lean masses in most scenarios rather than the less informative and

more difficult to assess BCI. BCIs rarely conferred an advantage over mass in predicting fat and

lean mass across the range of species, seasons, and ecological contexts considered here. Further

research into mass and composition dynamics across seasons and in different energetic contexts

will help determine the extent to which such relationships hold outside seasons of expected

extremes of energy budgets.

Acknowledgements. We thank Agnes MacDonald for long-term access to her trap-line, and to

the Champagne and Aishihik First Nations for allowing us to conduct fieldwork related to red

squirrels within their traditional territory. Fieldwork related to prairie dogs was conducted in

Treaty 4 territory, the traditional territory of the Oceti Sakowin and Niitsitpiis-stahkoii, and the

homeland of the Métis Nation. Fieldwork related to ground squirrels was conducted in Treaty 7

territory, the traditional territories of the Blackfoot Confederacy (Siksika, Piikani, and Kainai

First Nations), the Tsuut'ina First Nation, the Stoney Nakoda (Chiniki, Bearspaw, and Wesley

First Nations) and homeland of the Métis Nation. We thank the many volunteers, graduate

students, and field assistants, including Jillian Kusch, Tessie Aujla, Jack Hendrix, Ashley Mills,

Emily Kelvin, Megan Miller, M. Alejandra Hurtado, and Gabriela Heyer for field data collection,

and Grasslands National Park staff for assistance with permitting and logistical support. AEW

thanks Dr. Kurtis Swekla and Dr. Todd Shury for professional veterinary services in developing

anesthesia protocols for red squirrels.

Declarations

Funding The authors are grateful for ongoing funding from the Natural Sciences and

Engineering Research Council (SB, JEL, AGM), Canadian Foundation for Innovation (AGM,

JEL), and the National Science Foundation (BD, AGM). Fieldwork for this study was further

supported by the Northern Scientific Training Program (AEW), Sigma Xi Grant-in-Aid-of-

Research (AEW, RS), American Society of Mammalogists Grant-in-Aid-of-Research (AEW,

ALGC, RS), the American Museum of Natural History Theodore Roosevelt Memorial Fund

(ALGC) and the Alberta Conservation Association (ALGC, RS).

Conflict of Interest We have no competing interests.

Ethics approval All procedures were approved by the University of Saskatchewan Animal

Research Ethics Board. Fieldwork was completed under permits issued by Yukon Territorial

Government (red squirrels); Grasslands National Park and the Saskatchewan Ministry of the

Environment (prairie dogs); and Alberta Parks (ground squirrels).

Consent to participate Not applicable.

Consent for publication Not applicable.

Availability of data Data are archived on FigShare at the following private link:

https://figshare.com/s/51b0edc3df7e82bb9609. Data will be made public with an associated DOI

upon acceptance.

Code availability Code for analysis is available on GitHub at https://github.com/aewishart/bci-

composition.

References

Andersen R, Gaillard J-M, Linnell JD, Duncan P (2000) Factors affecting maternal care in an

income breeder, the European roe deer. Journal of Animal Ecology 69:672–682

Archibald DW, Fletcher QE, Boutin S, et al (2013) Sex-specific hoarding behavior in North

American red squirrels (Tamiasciurus hudsonicus). Journal of Mammalogy 94:761–770.

https://doi.org/10.1644/12-MAMM-A-213.1

Bairlein F (2002) How to get fat: Nutritional mechanisms of seasonal fat accumulation in

migratory songbirds. Naturwissenschaften 89:1–10. https://doi.org/10.1007/s00114-001-

0279-6

Blackwell GL, Bassett SM, Dickman CR (2006) Measurement error associated with external

measurements commonly used in small-mammal studies. Journal of Mammalogy

87:216–223. https://doi.org/10.1644/05-MAMM-A-215R1.1

Boag DA, Murie JO (1981) Weight in relation to sex, age, and season in Columbian ground

squirrels (Sciuridae: Rodentia). Canadian Journal of Zoology 59:999–1004.

https://doi.org/10.1139/z81-139

Bonte D, De La Peña E (2009) Evolution of body condition-dependent dispersal in

metapopulations. Journal of Evolutionary Biology 22:1242–1251.

https://doi.org/10.1111/j.1420-9101.2009.01737.x

Boutin S, Larsen KW (1993) Does food availability affect growth and survival of males and

females differently in a promiscuous small mammal, *Tamiasciurus hudsonicus*? Journal

of Animal Ecology 62:364–370

Boyer BB, Barnes BM (1999) Molecular and metabolic aspects of mammalian hibernation.

BioScience 49:713–724. https://doi.org/10.2307/1313595

Brigham RM, Geiser F (2012) Do red squirrels (*Tamiasciurus hudsonicus*) use daily torpor

during winter? Écoscience 19:127–132. https://doi.org/10.2980/19-2-3464

Broussard DR, Dobson FS, Murie JO (2005) The effects of capital on an income breeder:

evidence from female Columbian ground squirrels. Canadian Journal of Zoology 83:546–

552. https://doi.org/10.1139/z05-044

Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical

Information-Theoretic Approach (2nd ed).

Chang AM, Wiebe KL (2016) Body condition in Snowy Owls wintering on the prairies is greater

in females and older individuals and may contribute to sex-biased mortality. Auk

133:738–746. https://doi.org/10.1642/AUK-16-60.1

Dantzer B, McAdam AG, Humphries MM, et al (2020) Decoupling the effects of food and

density on life-history plasticity of wild animals using field experiments: Insights from

the steward who sits in the shadow of its tail, the North American red squirrel. Journal of

Animal Ecology 89:2397–2414. https://doi.org/10.1111/1365-2656.13341

Dittus WPJ (2013) Arboreal adaptations of body fat in wild toque macaques (Macaca sinica) and

the evolution of adiposity in primates. American Journal of Physical Anthropology

152:333–344. https://doi.org/10.1002/ajpa.22351

Dobson FS (1992) Body mass, structural size, and life-history patterns of the Columbian ground

squirrel. The American Naturalist 140:109–125

Dobson FS, Badry MJ, Geddes C (1992) Seasonal activity and body mass of Columbian ground

squirrels. Canadian Journal of Zoology 70:1364–1368. https://doi.org/10.1139/z92-192

Fisher D, Haines JA, Boutin S, et al (2019) Indirect effects on fitness between individuals that

have never met via an extended phenotype. Ecology Letters 442335.

https://doi.org/10.1101/442335

Fletcher QE, Boutin S, Lane JE, et al (2010) The functional response of a hoarding seed predator

to mast seeding. Ecology 91:2673–2683. https://doi.org/10.1890/09-1816.1

Fletcher QE, Landry-Cuerrier M, Boutin S, et al (2013) Reproductive timing and reliance on

hoarded capital resources by lacatating red squrrels. Oecologia 173: 1203-1215.

https://doi.org/10.1007/s00442-013-2699-3

Fletcher QE, Speakman JR, Boutin S, et al (2015) Daily energy expenditure during lactation is

strongly selected in a free-living mammal. Functional Ecology 29:195–208.

https://doi.org/10.1111/1365-2435.12313

Geiser F, Ruf T (1995) Hibernation versus daily torpor in mammals and birds: Physiological

variables and classification of torpor patterns. Physiological Zoology 68:935–966.

https://doi.org/10.1086/physzool.68.6.30163788

Glazier DS (2005) Beyond the '3/4-power law': variation in the intra- and interspecific scaling of

metabolic rate in animals. Biological Reviews 80: 611-662.

https://doi.org/10.1017/S1464793105006834

Green AJ (2001) Mass/length residuals: measures of body condition or generators of spurious

results? Ecology 82:1473. https://doi.org/10.2307/2680003

Guglielmo CG, McGuire LP, Gerson AR, Seewagen CL (2011) Simple, rapid, and non-invasive

measurement of fat, lean, and total water masses of live birds using quantitative magnetic

resonance. Journal of Ornithology 152:. https://doi.org/10.1007/s10336-011-0724-z

Gummer DL (2005) Geographic variation in torpor patterns: the northernmost prairie dogs and

kangaroo rats. Doctoral thesis, Department of Biology, University of Saskatchewan.

Haines JA, Delaney DM, Wishart AE, et al (2022) Sex-specific effects of capital resources on

reproductive timing and success in red squirrels. Behavioural Ecology and Sociobiology

Harris MA, Steudel K (2002) The relationship between maximum jumping performance and

hind limb morphology/physiology in domestic cats (*Felis silvestris catus*). Journal of

Experimental Biology 205:3877–3889. https://doi.org/10.1242/jeb.205.24.3877

Hawkshaw DM (2022) Covariates of intraspecific variation in hibernation expression in the

northernmost population of black-tailed prairie dogs. University of Saskatchewan

Hoogland JL (1995) The Black-tailed Prairie Dog: Social Life of a Burrowing Mammal.

University of Chicago Press, Chicago

Hoogland JL (2003) Sexual dimorphism of prairie dogs. Journal of Mammalogy 84:1254–1266.

https://doi.org/10.1644/BME-008

Humphries MM, Thomas DW, Kramer DL (2003) The role of energy availability in mammalian

hibernation: A cost-benefit approach. Physiological and Biochemical Zoology 76:165–

179. https://doi.org/10.1086/367950

Humphries MM, Boutin S, Thomas DW, et al. (2005) Expenditure freeze: the metabolic

response of small mammals to cold environments. Ecology Letters 8:13-26-1333. doi:

10.1111/j.1461-0248.2005.00839.x

Jacobs, S.R., Elliott, K., Guigueno, M.F., et al. (2012). Determining seabird body condition using

nonlethal measures. Physiological and Biochemical Zoology, 85: 85-95.

https://doi.org/10.1086/663832

Jakob EM, Marshall SD, Uetz GW, Estimating GW (1996) Estimating fitness: a comparison of

body condition indices. Oikos 77:61–67

Jenni L, Jenni-Eiermann S (1998) Fuel Supply and Metabolic Constraints in Migrating Birds.

Journal of Avian Biology 29:521–528

Johnson MS, Smith DL, Nagy TR, et al (2009) Validation of quantitative magnetic resonance

(QMR) for determination of body composition in rats. International Journal of Body

Composition Research 7:99–107.

https://doi.org/10.1097/MCA.0000000000000178.Endothelial

Jones AS, Johnson MS, Nagy TR (2009) Validation of quantitative magnetic resonance for the

determination of body composition of mice. International Journal of Body Composition

Research 7:67–72

Karpovich SA, Tøien Ø, Buck CL, Barnes BM (2009) Energetics of arousal episodes in

hibernating arctic ground squirrels. Journal of Comparative Physiology B: Biochemical,

Systemic, and Environmental Physiology 179:691–700. https://doi.org/10.1007/s00360-

009-0350-8

Kelly CD, Tawes BR, Worthington AM (2014) Evaluating indices of body condition in two

cricket species. Ecology and Evolution 4:4476–4487. https://doi.org/10.1002/ece3.1257

Kelsey NA, Bairlein F (2019) Migratory body mass increase in Northern Wheatears (*Oenanthe*

oenanthe) is the accumulation of fat as proven by quantitative magnetic resonance.

Journal of Ornithology 160: 389–397. https://doi.org/10.1007/s10336-018-1621-5

Krebs CJ, Singleton GR (1993) Indices of condition for small mammals. Australian Journal of

Zoology 41:317–323. https://doi.org/10.1071/ZO9930317

Kusch JM, Conway SE, Kapchinske A, Lane JE (2021) Reproductive phenology and seasonal

mass dynamics of black-tailed prairie dogs (*Cynomys ludovicianus*) at their northern

range limit. Can J Zool 99:257–268. https://doi.org/10.1139/cjz-2020-0054

Kusch JM, Matzke CC, Lane JE (2020) Reproductive failure predicts intracolony dispersal of

female black-tailed prairie dogs (*Cynomys ludovicianus*) in a northern population.

Western North American Naturalist 80:157–164. https://doi.org/10.3398/064.080.0203

Lane JE, Czenze ZJ, Findlay-Robinson R, Bayne E (2019) Phenotypic plasticity and local

adaptation in a wild hibernator evaluated through reciprocal translocation. American

Naturalist 194:516–528. https://doi.org/10.1086/702313

Lane JE, Kruuk LEB, Charmantier A, et al (2012) Delayed phenology and reduced fitness

associated with climate change in a wild hibernator. Nature 489:554–557.

https://doi.org/10.1038/nature11335

Lehmer EM, Savage LT, Antolin MF, Biggins DE (2006) Extreme plasticity in thermoregulatory

behaviors of free-ranging black-tailed prairie dogs. Physiological and Biochemical

Zoology 79:454–467. https://doi.org/10.1086/502816

Lehmer EM, Van Horne B (2001) Seasonal changes in lipids, diet, and body composition of free-

ranging black-tailed prairie dogs (*Cynomys ludovicianus*). Canadian Journal of Zoology

79:955–965. https://doi.org/10.1139/cjz-79-6-955

Lindström Å, Piersoma T (1993) Mass changes in migrating birds: the evidence for fat and

protein storage re☐examined. Ibis 135:70–78. https://doi.org/10.1111/j.1474-

919X.1993.tb02811.x

Martin JGA, Festa-Bianchet M, Côté SD, Blumstein DT (2013) Detecting between-individual

differences in hind-foot length in populations of wild mammals. Canadian Journal of

Zoology 91: 118–123. https://doi:10.1139/cjz-2012-0210

McAdam AG, Boutin S, Sykes AK, Humphries MM (2007) Life histories of female red squirrels

and their contributions to population growth and lifetime fitness. Écoscience 14:362–369.

https://doi.org/10.2980/1195-6860(2007)14[362:LHOFRS]2.0.CO;2

McGilvery, R.W. 1983. Biochemistry, a functional approach. Saunders, Philadelphia.

McGuire LP, Guglielmo CG (2010) Quantitative magnetic resonance: a rapid, noninvasive body

composition analysis technique for live and salvaged bats. Journal of Mammalogy

91:1375–1380. https://doi.org/10.1644/10-MAMM-A-051.1.Key

McGuire LP, Kelly LA, Baloun DE, et al (2018) Common condition indices are no more

effective than body mass for estimating fat stores in insectivorous bats. Journal of

Mammalogy 99:1065–1071. https://doi.org/10.1093/jmammal/gyy103

McGuire LP, Fuller NW, Haase CG, et al (2022). Lean mass dynamics in hibernating bats and

implications for energy and water budgets. Physiological and Biochemical Zoology 95:

317–325. https://doi.org/10.1086/720160

Mejías C, Navedo JG, Sabat P, et al (2022). Body composition and energy savings by

hibernation: lessons from the South American marsupial *Dromiciops gliroides*.

Physiological and Biochemical Zoology 95: 239–250. https://doi.org/10.1086/719932

Meyers P, Master LL (1983) Reproduction by *Peromyscus manuculatus*: size and compromise.

Journal of Mamalogy 64: 1-18. https://doi.org/10.2307/1380746

Molnár PK, Klanjscek T, Derocher AE, et al (2009) A body composition model to estimate

mammalian energy stores and metabolic rates from body mass and body length, with

application to polar bears. Journal of Experimental Biology 212:2313–2323.

https://doi.org/10.1242/jeb.026146

Moya-Laraño J, Macías-Ordóñez R, Blanckenhorn WU, Fernández-Montraveta C (2008)

Analysing body condition: Mass, volume or density? Journal of Animal Ecology

77:1099–1108. https://doi.org/10.1111/j.1365-2656.2008.01433.x

R Core Team (2020) R: A language and environment for statistical computing

Reynolds DS, Kunz TH (2001) Standard methods for destructive body composition analysis. In:

Speakman JR (ed) Body composition analysis of animals: a handbook of non-destructive

methods. Cambridge University Press

Schulte-Hostedde AI, Millar JS, Hickling GJ (2001) Evaluating body condition in small

mammals. Canadian Journal of Zoology 79:1021–1029. https://doi.org/10.1139/cjz-79-6-

1021

Schulte-Hostedde AI, Zinner B, Millar JS, Hickling GJ (2005) Restitution of mass-size residuals:

Validating body condition indices. Ecology 86:155–163. https://doi.org/10.1890/04-0232

Shimer HW (1903) Adaptations to aquatic, arboreal, fossorial and cursorial habits in mammals.

III. Fossorial habitats. American Naturalist 37: 819-825. https://doi.org/10.1086/278368

Smith MC (1968) Red squirrel responses to spruce cone failure in Interior Alaska. Journal of

Wildlife Management 32:305–317. https://doi.org/10.2307/3798975

Stevenson RD, Woods WA (2006) Condition indices for conservation: New uses for evolving

tools. Integrative and Comparative Biology 46:1169–1190.

https://doi.org/10.1093/icb/icl052

Studd EK, Boutin S, McAdam AG, Humphries MM (2016) Nest attendance of lactating red

squirrels (*Tamiasciurus hudsonicus*): influences of biological and environmental

correlates. Journal of Mammalogy 97:806-814. https://doi.org/10.1093/jmammal/gyw010

Swallow JG, Wroblewska AK, Waters RP, et al (2010) Phenotypic and evolutionary plasticity of

body composition in rats selectively bred for high endurance capacity. Journal of Applied

Physiology 109:778–785. https://doi.org/10.1152/japplphysiol.01026.2009

Taylor CR, Heglund NC, Maloiy GM (1982) Energetics and mechanics of terrestrial locomotion.

I. Metabolic energy consumption as a function of speed and body size in birds and

mammals. Journal of Experimental Biology 97:1–21

Tidhar WL, Speakman JR (2007) An evaluation of four non-destructive methods for predicting

body composition in a small rodent. 5:137–145

Tinsley FC, Taicher GZ, Heiman ML (2004) Evaluation of a quantitative magnetic resonance

method for mouse whole body composition analysis. Obesity Research 12:150–160.

https://doi.org/10.1038/oby.2004.20

Trombulak SC (1989) Running speed and body mass in Belding's ground squirrels. Journal of

Mammalogy 70:194–197

Weatherhead PJ, Brown GP (1996) Measurement versus estimation of condition in snakes.

Canadian Journal of Zoology 74:1617–1621. https://doi.org/10.1139/z96-179

Wells CP, Wilson JA, Kelt DA, Vuren DH Van (2019) Body mass as an estimate of female body

condition in a hibernating small mammal. The Canadian Field-Naturalist 133:34–42.

https://doi.org/10.22621/cfn.v133i1.2073

Wikelski M, Cooke SJ (2006) Conservation physiology. Trends in Ecology and Evolution

21:38–46. https://doi.org/10.1016/j.tree.2005.10.018

Wilson AJ, Nussey DH (2010) What is individual quality? An evolutionary perspective. Trends

in Ecology and Evolution 25:207–214. https://doi.org/10.1016/j.tree.2009.10.002

Wishart AE (2023) ariation in resource acquisition in a food-caching mammal, the North

American red squirrel (*Tamiasciurus hudsonicus*). Doctoral thesis, Department of

Biology, University of Saskatchewan. https://harvest.usask.ca/handle/10388/15058

Woodward G, Ebenman B, Emmerson M, et al (2005) Body size in ecological networks. Trends

in Ecology and Evolution 20:402–409. https://doi.org/10.1016/j.tree.2005.04.005

Young PJ (1990) Hibernating patterns of free-ranging Columbian ground squirrels. Oecologia

83:504–511

Zanghi BM, Cupp CJ, Pan Y, et al (2013a) Noninvasive measurements of body composition and

body water via quantitative magnetic x-ray absorptiometry in awake and sedated dogs.

American Journal of Veterinary Research 74:721–732.

https://doi.org/10.2460/ajvr.74.5.733

Zanghi BM, Cupp CJ, Pan Y, et al (2013b) Noninvasive measurements of body composition and

body water via quantitative magnetic resonance, deuterium water, and dual-energy x-ray

absorptiometry in cats. American Journal of Veterinary Research 74:721–732.

https://doi.org/10.2460/ajvr.74.5.733

Tables & Figures

Figure 1. Body composition (fat mass [A,C] and lean mass [B,D] in grams) as a function of

zygomatic-derived body condition index (ZW index, A-B) and body mass (C-D) for male (blue

triangles) and female (orange circles) North American red squirrels pre-winter.

Figure 2. Body composition (fat mass [A,C] and lean mass [B,D] in grams) as a function of

zygomatic-derived body condition index (ZW index, A-B) and body mass (C-D) for male (blue

triangles) and female (orange circles) adult non-breeding black-tailed prairie dogs pre-winter.

Figure 3. Body composition (fat mass [A,C] and lean mass [B,D] in grams) as a function of right

hind foot-derived body condition index (RHF index, A-B) and body mass (C-D) for male (blue

triangles) and female (orange circles) adult non-breeding Columbian ground squirrels pre-winter.

Figure 4. Body composition (fat mass [A,C] and lean mass [B,D] in grams) as a function of right

hind foot-derived body condition index (RHF index, A-B) and body mass (C-D) for male (blue

triangles) and female (orange circles) adult non-breeding Columbian ground squirrels in spring.

Figure 5. Partial plots showing the relationship between (A, B) species-specific body condition

indices and (C, D) body mass on (A, C) fat and (B, D) lean mass in two linear models for adult

non-breeding North American red squirrels (pre-winter), black-tailed prairie dogs (pre-winter),

and Columbian ground squirrels (pre-winter and spring).

Figure 1

718   Figure 2

729   Figure 3

742   Figure 4

754   Figure 5

768 **Table 1.** Summary of morphometric data (body mass, right hind foot length, and zygomatic

769 width) and body composition (fat mass, lean mass) for adult non-breeding male and female

770 North American red squirrels (pre-winter), black-tailed prairie dogs (pre-winter), and Columbian

771 ground squirrels (pre-winter and spring). Values reported as mean ± standard error of the mean

772 (SEM). Sample sizes for each sex ($\male$ = males, $\female$ = females) are indicated in the species column.

Species	Body mass (g)	Right hind foot length (mm)	Zygomatic width (mm)	Fat (g)	Lean (g)
North American red squirrels					
Male (*n* = 31)	251.8 ± 2.7 *	45.8 ± 0.3	28.9 ± 0.2	6.1 ± 0.5 (2.4%)	217.8 ± 2.9 (86.5%)
Female (*n* = 40)	243.8 ± 2.8 *	45.6 ± 0.2	28.3 ± 0.1	7.2 ± 0.6 (2.9%)	214.0 ± 2.3 (87.8%)
Black-tailed prairie dogs					
Male (*n* = 11)	1596.4 ± 28.7*	56.9 ± 0.4	49.8 ± 0.4*	495.6 ± 20.8 (30.9%)	928.8 ± 14.4* (58.3%)
Female (*n* = 16)	1456.9 ± 25.8*	56.2 ± 0.2	47.9 ± 0.3*	517.5 ± 18.2 (35.4%)	727.5 ± 15.9* (49.9%)
Columbian ground squirrels					
Pre-winter Males (*n* =15)	715.4 ± 11.5 *	46.4 ± 0.5	35.0 ± 0.4 *	179.3± 6.5* (25.0%)	471.7 ± 8.0* (66.0%)
Female (*n* = 15)	559.1 ± 13.3 *	45.3 ± 0.4	33.3 ± 0.3 *	116.4 ± 5.3* (20.8%)	388.4 ± 9.4* (69.5%)
Spring Males (*n* =33)	478.3± 89.5 *	46.7 ± 0.4 *	35.4 ± 0.2 *	59.7 ± 3.2 * (12.4%)	380.5 ± 7.5 * (79.6%)
Female (*n* = 48)	377.7 ± 6.7 *	45.1 ± 0.2 *	33.3 ± 0.2 *	29.5 ± 2.1 * (7.7%)	310.1 ± 5.3 * (82.2%)

773 *denotes significant differences between males and females; Welch two-sample t-test with α = 0.05*