Interpretation of body condition index should be informed by natural history

Andrea E. Wishart[1,*], Adriana L. Guerrero-Chacón[1], Rebecca Smith[1], Deborah M. Hawkshaw[1],

Andrew G. McAdam[2], Ben Dantzer[3,4], Stan Boutin[5], Jeffrey E. Lane[1]

*Author for correspondence: andrea.wishart@usask.ca

[1] Department of Biology, University of Saskatchewan, Saskatoon, SK, S7N 5E2, Canada

[2] Department of Ecology and Evolutionary Biology, Boulder, Colorado, USA

[3] Department of Psychology, University of Michigan, Ann Arbor, Michigan 48109-1043, USA

[4] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,

Michigan 48109-1043, USA

[5] Department of Biological Sciences, University of Alberta, Edmonton, Alberta T6G 2E9,

Canada

Abstract

1. Estimates of body condition are regularly made in wildlife studies, particularly those

focused on individual and/or population performance; however, many studies assume that

it is always beneficial to be heavier or have a higher body condition index (BCI), without

accounting for the physiological significance of variation in the composition of tissues

that differ in their function such as fat and lean mass.

2. We hypothesized that the relationship between BCI and masses of physiologically

important tissues (fat and lean) would be conditional on the annual patterns of energy

acquisition and expenditure of individuals under study, and tested relationships in three

species with contrasting ecologies in their respective natural ranges: an obligate

hibernator (Columbian ground squirrel, *Urocitellus columbianus*), a facultative hibernator

(black-tailed prairie dog, *Cynomys ludovicianus*), and a food-caching non-hibernator

(North American red squirrel, *Tamiasciurus hudsonicus*).

3. We measured fat and lean mass in adult males and females of these three species using

quantitative magnetic resonance (QMR). We measured body mass, two measures of

skeletal structure (zygomatic width and right hind foot length) to develop sex- and

species-specific BCIs, and tested the utility of BCI to predict body composition in each

species.

4. Body condition indices were more consistently and more strongly correlated with lean

mass than fat mass. The indices were most positively correlated with fat when fat was

expected to be very high (pre-hibernation prairie dogs). However, in all cases, fat and

lean mass were better predicted by overall body mass rather than BCI.

5. These results support our hypothesis that the utility of BCI in estimating fat is conditional

on the natural history and annual energetic patterns of the species with regards to

expected energy balances at the time of sampling, but measuring body mass alone is

likely capturing sufficient variation in fat and lean masses in most cases.

41
Key words: energetics, fat, food-caching, hibernation, lean, phenology

Introduction

Body mass is among the most frequently measured physical traits of organisms, because

it often accounts for much of the observed variation in other traits of interest as well as

ecological processes (Woodward et al., 2005). However, the mechanisms underlying these

associations may depend on the different components comprising body mass, rather than mass

itself. For processes in which energy stores or reserves (see (Lindström & Piersoma, 1993) for

distinction) are the currency of interest, the non-structural components of body mass that

represent metabolizable tissues (fat and lean mass; (Krebs & Singleton, 1993a) are likely more

relevant than size alone. For example, relatively high fat stores facilitate successful migration

(Bairlein, 2002) and hibernation (Humphries et al., 2003), while a higher proportion of lean mass

can enhance athletic performance (e.g., takeoff velocity in domestic cats *(Harris & Steudel,*

2002). Understanding relationships with the components of body mass may provide greater

insight into the mechanisms linking body mass to performance.

A given body mass may be distributed across structural elements of different sizes. Body

condition indices (BCIs) attempt to control for this by considering total body mass relative to

structural size (e.g., a linear skeletal measurement such as body length, tarsus length, foot length,

or a combination thereof; (Green, 2001; Jakob et al., 1996; Schulte-Hostedde et al., 2001)) to

give some indication of an individual's energetic 'condition'. Typically, individuals with

relatively higher body mass:skeletal size ratios are considered to be in 'better' condition than

individuals with lower ratios, as the former are assumed to have more energetic stores than

individuals of similar structural size, but with a lower metabolizable fraction (Schulte-Hostedde

et al., 2001). Accurate measures of 'condition' in this energetic sense are widely applicable to

scenarios such as livestock breeding programs (e.g., maximizing meat or offspring production),

conservation biology (e.g., measuring responses to habitat degradation; (Stevenson & Woods,

2006; Wikelski & Cooke, 2006), and in exploring evolutionary and ecological patterns (e.g.,

condition-dependent dispersal; (Bonte & De La Peña, 2009) and migration (Andersen et al.,

2000)). To better understand mechanisms underlying these relationships requires a closer

consideration of what physiological component of 'condition' BCIs are describing (e.g., lean

versus fat mass; (Schulte-Hostedde et al., 2001).

The dynamic nature of body condition across time (Krebs and Singleton 1993) demands

that its correlates and consequences be considered within the context of a species' natural history

(Molnár et al., 2009). The composition of a body in 'good condition' in an adaptive sense (i.e.,

that which is associated with increased survival and/or fitness, *sensu* (Wilson & Nussey, 2010),

should therefore be expected to vary over time depending on circannual energetics. Fat reserves

are an important source of metabolizable energy (~39.6 kJ/g for dry fat, in contrast to 17.8 kJ/g

for dry lean mass, (Jenni & Jenni-Eiermann, 1998)) that fuel organisms through energetically-

expensive behaviours or during periods of energetic shortfall. For example, individuals with

higher fat fractions entering hibernation are more likely to survive over winter and breed

successfully the following spring (Boyer & Barnes, 1999). Fatter is not always better though:

during the active season, carrying more body mass can decrease running speed (Trombulak,

1989) and alter circulating hormones (Taylor et al., 1982). Arboreal species may trade off fat

stores for locomotion (Dittus, 2013), and for species that primarily store energy off-body as

external food caches, fat reserves may not confer the same advantages during resource-scarce

seasons as they may for species that store energy exclusively on-body. Furthermore, a higher

lean fraction may be associated with increased activity, as shown in captive rats (*Rattus*

norvegicus; (Swallow et al., 2010). The body composition associated with better performance is

therefore likely to be highly dependent on natural history, as well as on the various activities an

animal undertakes throughout the year (Wells et al., 2019).

Because body mass and structural size can be measured from live animals, BCIs are less

invasive than chemical composition analyses that require lethal sampling (Reynolds & Kunz,

2001). In many species, BCIs correlate well with chemical quantification of fat mass in

vertebrates including birds (Chang & Wiebe, 2016) and reptiles (Weatherhead & Brown, 1996),

and invertebrates including arthropods (Jakob et al., 1996; Kelly et al., 2014; Moya-Laraño et al.,

2008). However, BCIs are not without drawbacks. Acquiring the structural size component(s) of

the index can be challenging. Krebs and Singleton (Krebs & Singleton, 1993b) warned of low

repeatability attributable to measurement bias across different observers (although this can be

significantly reduced by taking replicate measurements; (Blackwell et al., 2006), and estimating

lengths of long bones yield the most accurate estimates when measured from museum specimens

(e.g., (Dobson, 1992)), but can be challenging to measure on live, unanesthetized animals in field

conditions (Green, 2001). Because skeletal morphology often reflects natural history, as selection

favours certain shapes for certain lifestyles (e.g., arboreal vs. fossorial), the body component of

interest may not be accurately reflected by BCI. For example, BCI can correlate more strongly

with lean mass rather than fat mass (Schulte-Hostedde et al., 2001). Furthermore, the efficacy of

BCI in predicting the masses of body components may be no more effective than prediction

through body mass alone (e.g., body mass predicts fat stores in bats just as well as BCI;

(McGuire et al., 2018). Finally, since BCI estimates the non-skeletal component of body

composition in a general sense, its interpretation and therefore significance may differ depending

on the energetic aspects of the natural history of the focal species (e.g., for a fat-storing

hibernator versus a food-caching non-hibernator). We thus expect the correlation between BCIs

and body composition (i.e., fat and lean mass) to with both species and timing in annual cycles,

reflecting the dynamic nature of energy budgets in seasonally dependent activities.

We studied natural populations of three different mammal species to evaluate

concordance between BCIs and the energetic components of interest to many biologists (fat and

lean mass), and determined whether these relationships differ according to species' primary

energy storage mode (e.g., on-body fat storage vs. off-body food caching). We hypothesized that

the relationship between BCI and body composition depends on both the extent to which the

species relies on on-body energy stores (i.e., fat) for overwinter survival, and with the expected

energetic balance within a stage of the annual cycle (i.e., season).

We selected three species in the family Sciuridae which differ in patterns of energy

storage and metabolic demands: North American red squirrels (*Tamiasciurus hudsonicus*,

hereafter, red squirrels), black-tailed prairie dogs (*Cynomys ludovicianus*, hereafter prairie dogs),

and Columbian ground squirrels (*Urocitellus columbianus*, hereafter ground squirrels). We also

compare ground squirrels before and after hibernation to characterize within-species shifts in

energy stores. Red squirrels are arboreal mammals that store cached conifer cones in a central

larder ('midden'; (Smith, 1968) from late summer through autumn (Fletcher et al., 2010). Red

squirrels are relatively sexually monomorphic, weighing ~230-250 g on average as adults

(Boutin & Larsen, 1993), but males typically maintain larger cache sizes than females (Archibald

et al., 2013; Fisher et al., 2019; Haines et al., 2022). Red squirrels remain euthermic throughout

winter without using torpor (Brigham & Geiser, 2012), and are not known to gain significant

amounts of fat prior to winter, during which they rely on cached resources for energy.

Prairie dogs are semi-fossorial and can weigh up to 1710 g, but show high within- and

between-individual variation in body mass (Kusch et al., 2021). They are sexually dimorphic in

body mass (Hoogland, 1995, 2003); however, the extent of this dimorphism varies seasonally

(Hoogland, 1995, 2003; Kusch et al., 2021). Throughout most of their range, prairie dogs are

active throughout the winter; however, in southwestern Saskatchewan, where our study

population is located (the northern edge of the species distribution), they are known to hibernate

for ~4 months during winter (Gummer, 2005; Lehmer et al., 2006). Prairie dogs of both sexes

increase overall body mass, and fat mass specifically, leading up to winter (Kusch et al., 2021;

Lehmer & Van Horne, 2001). In hibernators, the greatest energy savings are achieved during

steady state torpor, when body temperature approximates ambient temperature, while most

energy is spent during interbout arousals to euthermy (Geiser & Ruf, 1995; Karpovich et al.,

2009). Prairie dogs in this population use torpor during winter, but are not considered efficient

hibernators, because they have a high minimum body temperatures during hibernation (16.9 °C

for males; Hawkshaw, 2022) and do not maintain steady state torpor (mean torpor bout duration

~126 hours for males, ~90 hours for females (Hawkshaw, 2022).

Ground squirrels are also semi-fossorial. Body mass varies substantially across their

active season (~400 g at emergence from hibernation in spring, to up to ~700 g prior to

immergence in late summer; (Dobson et al., 1992). These obligate hibernators are notable for

their short active season (~4 months) and extended time spent metabolically depressed in

hibernation (~8 months) each year (Dobson et al., 1992). Ground squirrels reach lower minimum

body temperatures (0 °C) and have longer torpor bout durations (~390 hours for males; Young,

1990) than prairie dogs. Ground squirrels exhibit male-biased sexual dimorphism, and

individuals experience significant body mass changes across time (Boag & Murie, 1981). Forage

quality and availability is low upon emergence in the spring (Lane et al., 2012; Young, 1990).

Consequently, body fat stores that remain after hibernation are presumed important for

supporting reproduction (Broussard et al., 2005).

We 1) characterized morphology and body composition of these three species, 2)

evaluated the correlation of BCI with body composition variables (lean and fat mass), and 3)

determined whether the correlation between BCI and body composition variables was higher

than between body mass and body composition variables. We expected hibernators (ground

squirrels and prairie dogs) to have the highest fat fraction within the pre-winter season given the

importance of on-body energetic reserves to sustain hibernation, compared to non-hibernating

red squirrels which rely on hoarded food, and compared to post-emergence ground squirrels in

spring who have metabolized fat stores overwinter. Because fat stores are expected to represent

the bulk of the pre-hibernation weight gain in the lead up to winter, we also expected high

concordance between BCI and fat mass in pre-winter hibernators. Conversely, we expected the

lowest concordance between BCI and fat mass to be in red squirrels, as lean mass is likely to be

more important to sustained caching activity (Fletcher et al., 2015). Finally, we test the null

hypothesis that BCI and body mass perform equally well in predicting lean and fat mass.

Materials and Methods

Study sites and population monitoring

We sampled free-ranging non-breeding adults from populations within the northern

regions of their respective ranges in Canada: red squirrels in the southwest Yukon (61° N, 138°

W, ~ 850 m a.s.l.), prairie dogs in Grasslands National Park, Saskatchewan (49°N, 107°W, ~770

m a.s.l.), and ground squirrels in Sheep River Provincial Park, Alberta (50°N, 114°W, ~ 1500 m

a.s.l.). For all populations, we collected data through live-trapping. All populations were

monitored for at least one year prior to data collection for the present study. For detailed

descriptions of population and reproductive monitoring, see (Dantzer et al., 2020; McAdam et

al., 2007)) for red squirrels, (Kusch et al., 2020)) for prairie dogs, and (Lane et al., 2019) for

ground squirrels. Briefly, all individuals received permanent uniquely marked ear tags (National

Band and Tag Company, Newport, KY) upon first trapping. We included only adults (individuals

one year of age or older) to minimize effects of skeletal growth dynamics. For ground squirrels

and prairie dogs, we knew ages for all individuals captured at first emergence following birth.

For red squirrels and some older ground squirrels and prairie dogs, exact ages were not known,

but we could confidently remove young-of-the-year based on size and breeding/nipple status on

first trapping (red squirrels excluded if under 150 g with small pink nipples at first trapping,

ground squirrels excluded if under 400 g, prairie dogs excluded if under 800 g a with small pink

nipples at first trapping). We excluded all pregnant (assessed by abdominal palpations) and

lactating (assessed by milk expression) females to remove variance related to maternal

investment in offspring.

Morphometric measurements

We measured body mass and size for all live-trapped individuals (Table 1). We weighed

each prairie dog to the nearest 5 g using a Pesola spring scale (Pesola AG, Baar, Switzerland),

and weighed red squirrels and ground squirrels to the nearest 1 g on an electronic balance. We

measured zygomatic arch width ('ZW') to the nearest millimeter using calipers (analogue for red

squirrels and prairie dogs; digital for ground squirrels). We measured right hind foot ('RHF')

length from heel to longest toe (excluding claw) to the nearest millimeter using a ruler fit with a

heelstop at 0 mm. We measured both ZW and RHF three times per handling, and used the mean

value for analyses.

We took these measurements during the same handling occurrence as body composition

analyses for all red squirrels, all prairie dogs except one, and most ground squirrels. For the

remaining prairie dog and ground squirrels, we used skeletal measurements taken on the nearest

date to when composition analyses were completed. The prairie dog was an adult and unlikely to

be growing, so we used measurements taken 81 days prior. The median interval between skeletal

measurements and composition scans for ground squirrels was 0 days (range: 0-120). We

excluded data from ground squirrels that were younger than 3 years of age if their skeletal

measurements were taken more than two weeks before or after the date of body composition and

mass measurements, as younger squirrels may still be growing structurally (Dobson, 1992). A

single yearling (female) remained in the spring dataset, so we removed all yearling ground

squirrels.

Body composition

We measured pre-winter body composition of red squirrels between late-September and

mid-October in 2018 and 2019, prairie dogs in late October 2018 (due to variable torpor patterns

in prairie dogs, immergence date into hibernation was not known), and ground squirrels between

late-July and late-August 2019 within a week of immergence for most individuals (median = 3.5

days; range = 0-15). We measured spring body composition of ground squirrels for which we

were confident were captured within a few days of emergence from hibernation (median = 1 day;

range = 0-3) between mid-April and early May 2019. Most individual ground squirrels were

measured in one season only, but 21 individuals were measured in both seasons. As we do not

expect ground squirrels to change skeletal size substantially after they have reached maturity, we

recorded morphometric data once per individual.

We used a quantitative magnetic resonance (QMR) body composition analyzer

(EchoMRI-1600, Echo Medical Systems, Houston, TX) to measure absolute lean and fat mass

(g). This technology provides a relatively non-invasive method of estimating body composition

precisely, and accurately (Tinsley et al., 2004). The remaining components not captured as fat

and lean mass are free water and skeletal mass (McGuire & Guglielmo, 2010). Quantities

measured using QMR correlate well with carcass-derived quantities for numerous species across

multiple taxa, most relevant here being rodents, including laboratory rats *Rattus norvegicus*

domestica (Johnson et al., 2009) and house mice *Mus musculus domesticus* (Jones et al., 2009).

The QMR approach allows for repeated measures of live animals, both awake and sedated

(McGuire & Guglielmo, 2010; Tinsley et al., 2004; Zanghi et al., 2013b, 2013a).

We housed our QMR system in a custom-designed trailer to enable transportation to each

study site. The trailer was climate-controlled to stabilize the temperature, as the magnet within

the QMR system is temperature-sensitive. We targeted an ambient temperature of 21 °C as per

manufacturer recommendations, although field conditions widened the range of stabilized

temperature to ± 7 °C. We calibrated the system daily to a 943 g canola oil standard at the

stabilized temperature, which was held for at least five hours prior to scanning animals

(Guglielmo et al., 2011). Our QMR system was outfitted with an additional antenna to measure

animals from 100 g up to 1600 g to accommodate the range of body masses of the three species.

Details of similar systems are described elsewhere (McGuire & Guglielmo, 2010), and we

followed similar protocols here. We live-trapped squirrels in the field and transported them to the

trailer. We placed each squirrel in a clear plexiglass holding tube with perforations to allow

ample airflow to the animal, then inserted the tube into the QMR chamber. We recorded body

composition through a minimum of two scans, reporting the average values for each individual.

In 2019, we administered a mild sedative via intramuscular injection to red squirrels prior to

scanning and collecting morphometric data (100 µg/kg of dexmedetomidine, reversed by 1

mg/kg atipamezole) to minimize stress and movement during scans. Sedation was not necessary

for the semi-fossorial prairie dogs or ground squirrels, who remained still and even fell asleep in

the chamber.

Calculating body condition indices (BCIs)

We retained individuals in the dataset for which we had measurements for all of the

following: ZW, RHF, body mass, and body composition. Within each species, we calculated

coefficients of variation for all variables, and analyzed relationships among variables. While

principal components analysis has been used previously as a general measure of structural size to

derive BCI (Schulte-Hostedde et al., 2005), we determined that it was not appropriate for our

dataset because correlation coefficients between ZW and RHF were not always positive

(Supplementary Fig. S1). Instead, we selected the single skeletal measure that had the greatest

coefficient of variation (CV) to generate a residual index (Jakob et al., 1996). Red squirrels

showed a negative but non-significant relationship between RHF and ZW (Table 1) so we chose

to use ZW to generate the BCI for this species ('ZW index'). The relationships between RHF and

ZW appeared to differ in direction between male and female prairie dogs; however, neither

correlation was significant, so we generated a ZW index for this species. In ground squirrels,

only data for females pre-winter showed a significant (positive) relationship. Females in spring

and males in both seasons showed no significant relationship between RHF and ZW. We

therefore used RHF to generate the single-metric BCI ('RHF index') for ground squirrels.

To calculate each BCI, we regressed either RHF (log transformed and standardized to

mean zero, unit variance) alone on body mass (log transformed and standardized to mean zero,

unit variance; generating the RHF index), or ZW (log transformed and scaled) alone on body

mass (log transformed and standardized to mean zero, unit variance; generating the ZW index).

Residual plots are shown in Supplementary Fig. S2. We calculated BCIs within season (pre-

winter/spring) and within sex for each species using separate regressions. Because some CV

values within species/seasons were similar for RHF and ZW, we also ran models using the BCI

derived from the alternate skeletal measurement (Supplementary Online Material). We indicate

in the results when results differed from the primary BCI model.

Statistical analyses

We performed all analyses in R (v.4.0.3, (R Core Team, 2020). We compared

morphological measures between the sexes within each species using a two-sample t-test. To

assess the efficacy of the selected BCI for each species in predicting body composition variables,

we modeled, separately for each species (and for ground squirrels, separately for each season),

linear models for fat mass (g) and lean mass (g), each predicted by the selected BCI interacting

with sex. We also modeled fat and lean mass predicted by body mass interacting with sex. We

compared the fit of the BCI model and body mass model for each component for each species

using Akaike's information criterion adjusted for small sample size (Burnham & Anderson,

2002), confirming no missing data. To compare the utility of BCI to predict fat and lean mass

across species, we fit a linear model for data from all three species (including both seasons for

ground squirrels) together. For fat and lean mass separately, we defined an interaction term

between species-specific BCI and species as the independent variable. We fit a similar model set

with body mass (scaled within species) instead of BCI.

Results

Morphological measurements

Males were significantly heavier than females in all three species: males were 3.3% heavier

in red squirrels, 9.6% heavier in prairie dogs, 27.9% heavier in pre-winter ground squirrels, and

26.6% heavier in spring ground squirrels (Table 1). Right hind foot length was different between

sexes only for ground squirrels in spring, with male RHF 1.6 mm longer than in females.

Zygomatic width was significantly larger in male prairie dogs and ground squirrels than females,

but similar between red squirrel males and females.

Body composition

In autumn, red squirrels showed the lowest percent body fat, with 2.4% in males and

2.9% in females (Table 1) while prairie dogs had the highest percent body fat, with 30.9% and

35.4% in males and females, respectively. Ground squirrels were the only species to have

significant sex differences in fat (in both seasons). They showed their highest body fat levels pre-

winter before entering hibernation (25.0% for males, 20.8% for females), and lowest in spring

(12.4% fat for males, 7.7% fat for females).

Relationship between BCI, body mass, and body composition

The relationship between BCI/body mass and fat/lean mass was positive in almost all

cases (Figures 1-4; the exception being for male prairie dog lean mass, Figure 2b). Both BCI and

body mass were significant predictors of fat and lean mass in their respective models for all

species, while sex and the BCI $\times$ sex interaction largely were not significant (Tables 2-5). For red

squirrels, the BCI and body mass models performed similarly in predicting fat but both had weak

correlations (Table 2, Figure 1). Red squirrel lean mass was better predicted by both BCI and

mass models than fat mass, with the body mass model providing a better fit. Sex was not

significant in the main ZW BCI model, but was in the alternate RHF BCI model (Supplementary

Table S3). For prairie dogs, both ZW BCI and body mass predicted fat well, but the body mass

model was a better fit for both fat and lean mass (Table 3, Figure 2). The BCI $\times$ sex interaction

was not significant in the main ZW BCI model, but was in the alternate RHF BCI model

(Supplementary Table S4). For pre-winter ground squirrels, the two models for fat were

indistinguishable based on AICc and had similar correlation strengths (Table 4, Figure 3). The

body mass model was a better fit for lean mass. This pattern held for spring, with BCI and body

mass models being similar for fat, and the body mass model providing a better fit for lean mass

(Table 5, Figure 4). Patterns did not change using ZW BCI instead (Supplementary Tables S5-6).

In the models including data from all species (and for ground squirrels, both seasons),

the effects of BCI, species, and the BCI $\times$ species interaction were significant for fat (Table 6,

Figure 5A). In the model for lean mass, BCI and species, but not their interaction, were

significant (Figure 5B). Patterns for models fitted with body mass instead of BCI differed

slightly, with body mass being significant in the fat model only in interaction terms except for

spring ground squirrels. In the lean model, all interaction terms were significant except for body

mass $\times$ spring ground squirrels. The strength of the correlations in both BCI and body mass

model sets were strong (adjusted $R^2 > 0.9$ for all) and similar within each component.

Discussion

We demonstrate that the utility of condition indices in predicting energetic components,

specifically fat, is conditional on expected energetic state given ecological considerations. If BCI

is a reliable indicator of 'condition' as it relates to on-body energy stores, it should be strongly

correlated with fat mass. However, when individuals were expected to be in a leaner state (e.g.,

food-caching red squirrels, spring ground squirrels), correlations between BCI and fat were low

to moderate. When individuals were expected to have higher fat stores (e.g., pre-hibernation

prairie dogs and ground squirrels), BCI models predicted fat well. Furthermore, the relationship

between BCI and fat depended on species (and season, for ground squirrels). This contingency

suggests that the assumption that high BCI is necessarily indicative of high fat stores should be

tempered depending on the species and/or time of year. In nearly all cases, BCI was positively

correlated with lean mass. Because fat and lean mass have significantly different energetic values

(Jenni & Jenni-Eiermann, 1998), interpretations of BCI as they relate to metabolizable energy

should take into account species-specific natural history and annual energetic patterns. For

example, some red squirrels that would have ranked as lower condition based on BCI had nearly

twice as much fat as some individuals who had higher BCI values. Furthermore, models fit with

body mass were almost always more highly correlated with fat and lean mass than models with

BCI.

The selection of these three species, and the time of year at which they were studied,

provides insight into how body composition manifests in BCI when individuals are in a peak

positive energy balance after accumulating surplus energy to sustain them through upcoming

energetic shortfalls, and when they are expected to have depleted much of that accumulated

energy. By investigating these relationships at the ends of the continuum of energetic states that

organisms may experience throughout the year, we demonstrate that interpretation of 'condition'

indices should be sensitive to seasonal energetic demands. For example, red squirrels primarily

store energy as cached food so were expected to carry little fat. In comparison, we expected

prairie dogs to have high fat stores to sustain them through inefficient hibernation (Gummer,

2005; Hawkshaw, 2022), and indeed they had the highest percent body fat of all three species.

These results illustrate that the relative importance of fat and lean mass is likely to vary with

seasonal activities interacting with natural history.

Almost every relationship between either predictor variable (BCI or body mass) and body

component (fat or lean mass) was positive, save for male prairie dog lean mass. Echoing Schulte-

Hostedde et al. (2001), BCIs are capturing variation in both components. There are, however,

fine scale, but important, differences worth discussing. We found that in general, lean mass had

less variation around the line of best fit than fat mass, reflective of previous studies on small-non

hibernating mammals that also found that BCIs tend to be more effective in predicting lean dry

mass and water compared with predicting fat mass (Schulte-Hostedde et al., 2001; Tidhar &

Speakman, 2007). Given the dynamic nature of body composition in hibernating species,

estimating fat levels can be particularly difficult since many studies using residual-derived BCIs

to assess fat assume that lean mass scales with body size, while fat mass is assumed to vary with

condition (McGuire et al., 2018).

We have shown that the utility of body mass and/or BCI to describe energy-relevant

components is conditional on natural history and annual energetic cycles. Quantitative magnetic

resonance provides fine-scale measurements that may be important for specific energetics

questions, but general relationships between body mass and fat/lean quantities are positive.

Ultimately, this study strengthens the case for using body mass as a covariate to capture general

variation in soft tissue in most scenarios, as we have demonstrated that BCIs do not confer an

advantage in predicting fat and lean mass in these three species. Further research into mass and

composition dynamics across seasons and in different energetic contexts will help determine the

extent to which such relationships hold outside seasons of expected extremes of energy budgets.

Ethics. All procedures were approved by the University of Saskatchewan Animal Research Ethics Board. Fieldwork was completed under permits issued by Yukon Territorial Government (red squirrels); Grasslands National Park and the Saskatchewan Ministry of the Environment (prairie dogs); and Alberta Parks (ground squirrels).

Acknowledgements. We thank Agnes MacDonald for long-term access to her trap-line, and to the Champagne and Aishihik First Nations for allowing us to conduct fieldwork related to red squirrels within their traditional territory. Fieldwork related to prairie dogs was conducted in Treaty 4 territory, the traditional territory of the Oceti Sakowin and Niitsitpiis-stahkoii, and the homeland of the Métis Nation. Fieldwork related to ground squirrels was conducted in Treaty 7 territory, the traditional territories of the Blackfoot Confederacy (Siksika, Piikani, and Kainai First Nations), the Tsuut'ina First Nation, the Stoney Nakoda (Chiniki, Bearspaw, and Wesley First Nations) and homeland of the Métis Nation. We thank the many volunteers, graduate students, and field assistants, including Jillian Kusch, Tessie Aujla, Jack Hendrix, Ashley Mills, and Emily Kelvin for field data collection, and Grasslands National Park staff for assistance with permitting and logistical support. AEW thanks Dr. Kurtis Swekla and Dr. Todd Shury for professional veterinary services in developing anesthesia protocols for red squirrels. The authors are grateful for ongoing funding from the Natural Sciences and Engineering Research Council (SB, JEL, AGM), Canadian Foundation for Innovation (JEL), and the National Science Foundation (BD, AGM). Fieldwork for this study was further supported by the Northern Scientific Training Program (AEW), Sigma Xi Grant-in-Aid-of-Research (AEW, RS), American Society of Mammalogists Grant-in-Aid-of-Research (AEW, ALGC, RS), the American Museum

of Natural History Theodore Roosevelt Memorial Fund (ALGC) and the Alberta Conservation

Association (ALGC, RS).

Conflict of Interest. We have no competing interests.

Author Contributions. AEW designed the study, collected data, performed data and statistical

analysis, and drafted the manuscript. JEL participated in the design. All authors contributed to

data collection, provided valuable discussion and contributions to the writing of the manuscript,

and gave final approval for publication.

Data availability: Data are archived on FigShare and will be made public with an associated

DOI upon acceptance.

426
427

References

Andersen, R., Gaillard, J.-M., Linnell, J. D., & Duncan, P. (2000). Factors affecting maternal

care in an income breeder, the European roe deer. *Journal of Animal Ecology*, *69*, 672–

682.

Archibald, D. W., Fletcher, Q. E., Boutin, S., McAdam, A. G., Speakman, J. R., & Humphries,

M. M. (2013). Sex-specific hoarding behavior in North American red squirrels

(*Tamiasciurus hudsonicus*). *Journal of Mammalogy*, *94*(4), 761–770.

https://doi.org/10.1644/12-MAMM-A-213.1

Bairlein, F. (2002). How to get fat: Nutritional mechanisms of seasonal fat accumulation in

migratory songbirds. *Naturwissenschaften*, *89*(1), 1–10. https://doi.org/10.1007/s00114-

001-0279-6

Blackwell, G. L., Bassett, S. M., & Dickman, C. R. (2006). Measurement error associated with

external measurements commonly used in small-mammal studies. *Journal of*

Mammalogy, *87*(2), 216–223. https://doi.org/10.1644/05-MAMM-A-215R1.1

Boag, D. A., & Murie, J. O. (1981). Weight in relation to sex, age, and season in Columbian

ground squirrels (Sciuridae: Rodentia). *Canadian Journal of Zoology*, *59*(6), 999–1004.

https://doi.org/10.1139/z81-139

Bonte, D., & De La Peña, E. (2009). Evolution of body condition-dependent dispersal in

metapopulations. *Journal of Evolutionary Biology*, *22*(6), 1242–1251.

https://doi.org/10.1111/j.1420-9101.2009.01737.x

Boutin, S., & Larsen, K. W. (1993). Does food availability affect growth and survival of males

and females differently in a promiscuous small mammal, *Tamiasciurus hudsonicus*?

Journal of Animal Ecology, *62*(2), 364–370.

Boyer, B. B., & Barnes, B. M. (1999). Molecular and metabolic aspects of mammalian

hibernation. *BioScience*, *49*(9), 713–724. https://doi.org/10.2307/1313595

Brigham, R. M., & Geiser, F. (2012). Do red squirrels (*Tamiasciurus hudsonicus*) use daily

torpor during winter? *Écoscience*, *19*(2), 127–132. https://doi.org/10.2980/19-2-3464

Broussard, D. R., Dobson, F. S., & Murie, J. O. (2005). The effects of capital on an income

breeder: Evidence from female Columbian ground squirrels. *Canadian Journal of*

Zoology, *83*(4), 546–552. https://doi.org/10.1139/z05-044

Burnham, K. P., & Anderson, D. R. (2002). Model Selection and Multimodel Inference: A

460   Practical Information-Theoretic Approach (2nd ed). In *Ecological Modelling* (Vol. 172).

461   https://doi.org/10.1016/j.ecolmodel.2003.11.004

462 Chang, A. M., & Wiebe, K. L. (2016). Body condition in Snowy Owls wintering on the prairies

463   is greater in females and older individuals and may contribute to sex-biased mortality.

464   *Auk*, *133*(4), 738–746. https://doi.org/10.1642/AUK-16-60.1

465 Dantzer, B., McAdam, A. G., Humphries, M. M., Lane, J. E., & Boutin, S. (2020). Decoupling

466   the effects of food and density on life-history plasticity of wild animals using field

467   experiments: Insights from the steward who sits in the shadow of its tail, the North

468   American red squirrel. *Journal of Animal Ecology*, *89*(11), 2397–2414.

469   https://doi.org/10.1111/1365-2656.13341

470 Dittus, W. P. J. (2013). Arboreal adaptations of body fat in wild toque macaques (*Macaca sinica*)

471   and the evolution of adiposity in primates. *American Journal of Physical Anthropology*,

472   *152*(3), 333–344. https://doi.org/10.1002/ajpa.22351

473 Dobson, F. S. (1992). Body mass, structural size, and life-history patterns of the Columbian

474   ground squirrel. *The American Naturalist*, *140*(1), 109–125.

475 Dobson, F. S., Badry, M. J., & Geddes, C. (1992). Seasonal activity and body mass of

476   Columbian ground squirrels. *Canadian Journal of Zoology*, *70*(7), 1364–1368.

477   https://doi.org/10.1139/z92-192

478 Fisher, D., Haines, J. A., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., & McAdam, A. G.

479   (2019). Indirect effects on fitness between individuals that have never met via an

480   extended phenotype. *Ecology Letters*, 442335. https://doi.org/10.1101/442335

Fletcher, Q. E., Boutin, S., Lane, J. E., LaMontagne, J. M., McAdam, A. G., Krebs, C. J., & Humphries, M. M. (2010). The functional response of a hoarding seed predator to mast seeding. *Ecology*, *91*(9), 2673–2683. https://doi.org/10.1890/09-1816.1

Fletcher, Q. E., Speakman, J. R., Boutin, S., Lane, J. E., Mcadam, A. G., Gorrell, J. C., Coltman, D. W., & Humphries, M. M. (2015). Daily energy expenditure during lactation is strongly selected in a free-living mammal. *Functional Ecology*, *29*, 195–208. https://doi.org/10.1111/1365-2435.12313

Geiser, F., & Ruf, T. (1995). Hibernation versus daily torpor in mammals and birds: Physiological variables and classification of torpor patterns. *Physiological Zoology*, *68*(6), 935–966. https://doi.org/10.1086/physzool.68.6.30163788

Green, A. J. (2001). Mass/length residuals: measures of body condition or generators of spurious results? *Ecology*, *82*(5), 1473. https://doi.org/10.2307/2680003

Guglielmo, C. G., McGuire, L. P., Gerson, A. R., & Seewagen, C. L. (2011). Simple, rapid, and non-invasive measurement of fat, lean, and total water masses of live birds using quantitative magnetic resonance. *Journal of Ornithology*, *152*. https://doi.org/10.1007/s10336-011-0724-z

Gummer, D. L. (2005). Geographic Variation in Torpor Patterns: The Northernmost Prarie Dogs and Kangaroo Rats. *Ph.D. Thesis. Department of Biology, University of Saskatchewan.*

Haines, J. A., Delaney, D. M., Wishart, A. E., McAdam, A. G., Coltman, D. W., Lane, J. E., Dantzer, B., & Boutin, S. (2022). Sex-specific effects of capital resources on reproductive timing and success in red squirrels. *Behavioural Ecology and Sociobiology*.

Harris, M. A., & Steudel, K. (2002). The relationship between maximum jumping performance

and hind limb morphology/physiology in domestic cats (*Felis silvestris catus*). *Journal of*

Experimental Biology, *205*(24), 3877–3889. https://doi.org/10.1242/jeb.205.24.3877

Hawkshaw, D. M. (2022). Covariates of intraspecific variation in hibernation expression in the

northernmost population of black-tailed prairie dogs. *M.Sc. Thesis. Department of*

Biology, University of Saskatchewan.

Hoogland, J. L. (1995). *The Black-tailed Prairie Dog: Social Life of a Burrowing Mammal.*

University of Chicago Press.

Hoogland, J. L. (2003). Sexual dimorphism of prairie dogs. *Journal of Mammalogy*, *84*(4),

1254–1266. https://doi.org/10.1644/BME-008

Humphries, M. M., Thomas, D. W., & Kramer, D. L. (2003). The role of energy availability in

mammalian hibernation: A cost-benefit approach. *Physiological and Biochemical*

Zoology, *76*(2), 165–179. https://doi.org/10.1086/367950

Jakob, E. M., Marshall, S. D., Uetz, G. W., & Estimating, G. W. (1996). Estimating fitness: A

comparison of body condition indices. *Oikos*, *77*(1), 61–67.

Jenni, L., & Jenni-Eiermann, S. (1998). Fuel supply and metabolic constraints in migrating birds.

Journal of Avian Biology, *29*(4), 521–528.

Johnson, M. S., Smith, D. L., Nagy, T. R., Smith Jr, D. L., & Nagy, T. R. (2009). Validation of

quantitative magnetic resonance (QMR) for determination of body composition in rats.

International Journal of Body Composition Research, *7*(3), 99–107.

Jones, A. S., Johnson, M. S., & Nagy, T. R. (2009). Validation of quantitative magnetic

resonance for the determination of body composition of mice. *International Journal of*

Body Composition Research, *7*(2), 67–72.

Karpovich, S. A., Tøien, Ø., Buck, C. L., & Barnes, B. M. (2009). Energetics of arousal episodes

in hibernating arctic ground squirrels. *Journal of Comparative Physiology B:*

Biochemical, Systemic, and Environmental Physiology, *179*(6), 691–700.

https://doi.org/10.1007/s00360-009-0350-8

Kelly, C. D., Tawes, B. R., & Worthington, A. M. (2014). Evaluating indices of body condition

in two cricket species. *Ecology and Evolution*, *4*(23), 4476–4487.

https://doi.org/10.1002/ece3.1257

Krebs, C. J., & Singleton, G. R. (1993a). Indices of condition for small mammals. *Australian*

Journal of Zoology, *41*(4), 317–323. https://doi.org/10.1071/ZO9930317

Krebs, C. J., & Singleton, G. R. (1993b). Indices of condition for small mammals. *Australian*

Journal of Zoology, *41*(1989), 317–323. https://doi.org/10.1071/ZO9930317

Kusch, J. M., Conway, S. E., Kapchinske, A., & Lane, J. E. (2021). Reproductive phenology and

seasonal mass dynamics of black-tailed prairie dogs (*Cynomys ludovicianus*) at their

northern range limit. *Canadian Journal of Zoology*, *99*(4), 257–268.

https://doi.org/10.1139/cjz-2020-0054

Kusch, J. M., Matzke, C. C., & Lane, J. E. (2020). Reproductive failure predicts intracolony

dispersal of female black-tailed prairie dogs (*Cynomys ludovicianus*) in a Northern

Population. *Western North American Naturalist*, *80*(2), 157–164.

https://doi.org/10.3398/064.080.0203

Lane, J. E., Czenze, Z. J., Findlay-Robinson, R., & Bayne, E. (2019). Phenotypic plasticity and

local adaptation in a wild hibernator evaluated through reciprocal translocation. *American*

Naturalist, *194*(4), 516–528. https://doi.org/10.1086/702313

Lane, J. E., Kruuk, L. E. B., Charmantier, A., Murie, J. O., & Dobson, F. S. (2012). Delayed

phenology and reduced fitness associated with climate change in a wild hibernator.

Nature, *489*(7417), 554–557. https://doi.org/10.1038/nature11335

Lehmer, E. M., Savage, L. T., Antolin, M. F., & Biggins, D. E. (2006). Extreme plasticity in

thermoregulatory behaviors of free-ranging black-tailed prairie dogs. *Physiological and*

Biochemical Zoology, *79*(3), 454–467. https://doi.org/10.1086/502816

Lehmer, E. M., & Van Horne, B. (2001). Seasonal changes in lipids, diet, and body composition

of free-ranging black-tailed prairie dogs (*Cynomys ludovicianus*). *Canadian Journal of*

Zoology, *79*(6), 955–965. https://doi.org/10.1139/cjz-79-6-955

Lindström, Å., & Piersoma, T. (1993). Mass changes in migrating birds: The evidence for fat and

protein storage re□examined. *Ibis*, *135*(1), 70–78. https://doi.org/10.1111/j.1474-

919X.1993.tb02811.x

McAdam, A. G., Boutin, S., Sykes, A. K., & Humphries, M. M. (2007). Life histories of female

red squirrels and their contributions to population growth and lifetime fitness.

Écoscience, *14*(3), 362–369. https://doi.org/10.2980/1195-

6860(2007)14[362:LHOFRS]2.0.CO;2

McGuire, L. P., & Guglielmo, C. G. (2010). Quantitative magnetic resonance: A rapid ,

noninvasive body composition analysis technique for live and salvaged bats. *Journal of*

Mammalogy, *91*(6), 1375–1380. https://doi.org/10.1644/10-MAMM-A-051.1.Key

McGuire, L. P., Kelly, L. A., Baloun, D. E., Boyle, W. A., Cheng, T. L., Clerc, J., Fuller, N. W.,

Gerson, A. R., Jonasson, K. A., Rogers, E. J., Sommers, A. S., & Guglielmo, C. G.

(2018). Common condition indices are no more effective than body mass for estimating

569   fat stores in insectivorous bats. *Journal of Mammalogy*, *99*(5), 1065–1071.

570   https://doi.org/10.1093/jmammal/gyy103

571 Molnár, P. K., Klanjscek, T., Derocher, A. E., Obbard, M. E., & Lewis, M. A. (2009). A body

572   composition model to estimate mammalian energy stores and metabolic rates from body

573   mass and body length, with application to polar bears. *Journal of Experimental Biology*,

574   *212*(15), 2313–2323. https://doi.org/10.1242/jeb.026146

575 Moya-Laraño, J., Macías-Ordóñez, R., Blanckenhorn, W. U., & Fernández-Montraveta, C.

576   (2008). Analysing body condition: Mass, volume or density? *Journal of Animal Ecology*,

577   *77*(6), 1099–1108. https://doi.org/10.1111/j.1365-2656.2008.01433.x

578 R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation

579   for Statistical Computing.

580 Reynolds, D. S., & Kunz, T. H. (2001). Standard methods for destructive body composition

581   analysis. In J. R. Speakman (Ed.), *Body composition analysis of animals: A handbook of*

582   *non-destructive methods*. Cambridge University Press.

583 Schulte-Hostedde, A. I., Millar, J. S., & Hickling, G. J. (2001). Evaluating body condition in

584   small mammals. *Canadian Journal of Zoology*, *79*(6), 1021–1029.

585   https://doi.org/10.1139/cjz-79-6-1021

586 Schulte-Hostedde, A. I., Zinner, B., Millar, J. S., & Hickling, G. J. (2005). Restitution of mass-

587   size residuals: Validating body condition indices. *Ecology*, *86*(1), 155–163.

588   https://doi.org/10.1890/04-0232

589 Smith, M. C. (1968). Red squirrel responses to spruce cone failure in Interior Alaska. *Journal of*

590   *Wildlife Management*, *32*(2), 305–317. https://doi.org/10.2307/3798975

Stevenson, R. D., & Woods, W. A. (2006). Condition indices for conservation: New uses for

evolving tools. *Integrative and Comparative Biology*, *46*(6), 1169–1190.

https://doi.org/10.1093/icb/icl052

Swallow, J. G., Wroblewska, A. K., Waters, R. P., Renner, K. J., Britton, S. L., & Koch, L. G.

(2010). Phenotypic and evolutionary plasticity of body composition in rats selectively

bred for high endurance capacity. *Journal of Applied Physiology*, *109*(3), 778–785.

https://doi.org/10.1152/japplphysiol.01026.2009

Taylor, C. R., Heglund, N. C., & Maloiy, G. M. (1982). Energetics and mechanics of terrestrial

locomotion. I. Metabolic energy consumption as a function of speed and body size in

birds and mammals. *Journal of Experimental Biology*, *97*(1970), 1–21.

Tidhar, W. L., & Speakman, J. R. (2007). An evaluation of four non-destructive methods for

predicting body composition in a small rodent. *International Journal of Body*

Composition Research 5(4), 137–145.

Tinsley, F. C., Taicher, G. Z., & Heiman, M. L. (2004). Evaluation of a quantitative magnetic

resonance method for mouse whole body composition analysis. *Obesity Research*, *12*(1),

150–160. https://doi.org/10.1038/oby.2004.20

Trombulak, S. C. (1989). Running speed and body mass in Belding's ground squirrels. *Journal*

of Mammalogy, *70*(1), 194–197.

Weatherhead, P. J., & Brown, G. P. (1996). Measurement versus estimation of condition in

snakes. *Canadian Journal of Zoology*, *74*(9), 1617–1621. https://doi.org/10.1139/z96-179

Wells, C. P., Wilson, J. A., Kelt, D. A., & Vuren, D. H. Van. (2019). Body mass as an estimate

of female body condition in a hibernating small mammal. *The Canadian Field-Naturalist*,

133(1), 34–42. https://doi.org/10.22621/cfn.v133i1.2073

Wikelski, M., & Cooke, S. J. (2006). Conservation physiology. *Trends in Ecology and Evolution*,

21(1), 38–46. https://doi.org/10.1016/j.tree.2005.10.018

Wilson, A. J., & Nussey, D. H. (2010). What is individual quality? An evolutionary perspective.

Trends in Ecology and Evolution, *25*(4), 207–214.

https://doi.org/10.1016/j.tree.2009.10.002

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J. M., Olesen, J. M., Valido, A., &

Warren, P. H. (2005). Body size in ecological networks. *Trends in Ecology and*

Evolution, *20*(7), 402–409. https://doi.org/10.1016/j.tree.2005.04.005

Young, P. J. (1990). Hibernating patterns of free-ranging Columbian ground squirrels.

Oecologia, *83*, 504–511.

Zanghi, B. M., Cupp, C. J., Pan, Y., Tissot-favre, D. G., Milgram, N. W., Nagy, T. R., &

Dobson, H. (2013a). Noninvasive measurements of body composition and body water via

quantitative magnetic resonance, deuterium water, and dual-energy x-ray absorptiometry

in cats. *American Journal of Veterinary Research*, *74*(5), 721–732.

https://doi.org/10.2460/ajvr.74.5.733

Zanghi, B. M., Cupp, C. J., Pan, Y., Tissot-favre, D. G., Milgram, N. W., Nagy, T. R., &

Dobson, H. (2013b). Noninvasive measurements of body composition and body water via

quantitative magnetic x-ray absorptiometry in awake and sedated dogs. *American Journal*

of Veterinary Research, *74*(5), 721–732. https://doi.org/10.2460/ajvr.74.5.733

Tables & Figures

Figure 1. Body composition (fat [A,C] and lean [B,D] in grams) as a function of zygomatic-

derived body condition index (ZW index, A-B) and body mass (C-D) for male (blue triangles)

and female (orange circles) North American red squirrels pre-winter.

Figure 2. Body composition (fat [A,C] and lean [B,D] in grams) as a function of zygomatic-

derived body condition index (ZW index, A-B) and body mass (C-D) for male (blue triangles)

and female (orange circles) adult non-breeding black-tailed prairie dogs pre-winter.

Figure 3. Body composition (fat [A,C] and lean [B,D] in grams) as a function of right hind foot-

derived body condition index (RHF index, A-B) and body mass (C-D) for male (blue triangles)

and female (orange circles) adult non-breeding Columbian ground squirrels pre-winter.

Figure 4. Body composition (fat [A,C] and lean [B,D] in grams) as a function of right hind foot-

derived body condition index (RHF index, A-B) and body mass (C-D) for male (blue triangles)

and female (orange circles) adult non-breeding Columbian ground squirrels in spring.

Figure 5. Partial plots showing the relationship between (A, B) species-specific body condition indices

and (C, D) body mass on (A, C) fat and (B, D) lean mass in two linear models for adult non-breeding

North American red squirrels (pre-winter), black-tailed prairie dogs (pre-winter), and Columbian ground

squirrels (pre-winter and spring).

670 **Table 1.** Summary of morphometric data (body mass, right hind foot length, and zygomatic width) and

671 body composition (fat mass, lean mass) for adult non-breeding male and female North American red

672 squirrels (pre-winter), black-tailed prairie dogs (pre-winter), and Columbian ground squirrels (pre-winter

673 and spring). Values reported as mean ± SEM. Sample sizes for each sex ($\male$ = males, $\female$ = females) are

674 indicated in the species column.

Species	Body mass (g)	Right hind foot length (mm)	Zygomatic width (mm)	Fat (g)	Lean (g)
North American red squirrels					
Male (*n* = 31)	251.8 ± 2.7 *	45.8 ± 0.3	28.9 ± 0.2	6.1 ± 0.5 (2.4%)	217.8 ± 2.9 (86.5%)
Female (*n* = 40)	243.8 ± 2.8 *	45.6 ± 0.2	28.3 ± 0.1	7.2 ± 0.6 (2.9%)	214.0 ± 2.3 (87.8%)
Black-tailed prairie dogs					
Male (*n* = 11)	1596.4 ± 28.7*	56.9 ± 0.4	49.8 ± 0.4*	495.6 ± 20.8 (30.9%)	928.8 ± 14.4* (58.3%)
Female (*n* = 16)	1456.9 ± 25.8*	56.2 ± 0.2	47.9 ± 0.3*	517.5 ± 18.2 (35.4%)	727.5 ± 15.9* (49.9%)
Columbian ground squirrels					
Pre-winter Males (*n* =15)	715.4 ± 11.5 *	46.4 ± 0.5	35.0 ± 0.4 *	179.3± 6.5* (25.0%)	471.7 ± 8.0* (66.0%)
Female (*n* = 15)	559.1 ± 13.3 *	45.3 ± 0.4	33.3 ± 0.3 *	116.4 ± 5.3* (20.8%)	388.4 ± 9.4* (69.5%)
Spring Males (*n* =33)	478.3± 89.5 *	46.7 ± 0.4 *	35.4 ± 0.2 *	59.7 ± 3.2 * (12.4%)	380.5 ± 7.5 * (79.6%)
Female (*n* = 48)	377.7 ± 6.7 *	45.1 ± 0.2 *	33.3 ± 0.2 *	29.5 ± 2.1 * (7.7%)	310.1 ± 5.3 * (82.2%)

675   **denotes significant differences between males and females; Welch two-sample t-test with α = 0.05*
676

677 **Table 2.** Model coefficients for linear models testing whether a body condition index (BCI) derived from

678 zygomatic width (ZW index) explains observed variation in fat and lean mass in North American red

679 squirrels pre-winter. Reference group for sex: female.

Component	Model	Independent terms	Estimate	*t*	*p*	AICc	Adjusted R^2
Fat	~ BCI*sex	Intercept	7.22 ± 0.44	16.54	< 0.001 *	390.7	0.215
		BCI	1.87 ± 0.44	4.26	< 0.001 *		
		Sex	-1.25 ± 0.68	-1.85	0.069		

			Estimate	t	p	AICc	Adjusted R²
		BCI*sex	-1.03 ± 0.73	-1.42	0.161		
	~mass*sex	Intercept	-21.67 ± 6.36	-3.41	< 0.002 *	389.4	0.223
		Body mass (g)	0.12 ± 0.03	4.56	< 0.001 *		
		Sex	19.26 ± 10.72	1.80	0.077		
		Mass*sex	-0.09 ± 0.04	-1.977	0.051		
Lean	~ BCI*sex	Intercept	214.51 ± 1.33	160.89	< 0.001 *	562.6	0.644
		BCI	12.28 ± 1.34	9.12	< 0.001 *		
		Sex	3.46 ± 2.07	1.67	0.099		
		BCI*sex	0.85 ± 2.22	0.38	0.702		
	~mass*sex	Intercept	24.13 ± 15.53	1.55	0.125	527.0	0.776
		Body mass (g)	0.78 ± 0.06	12.30	< 0.001 *		
		Sex	-28.60 ± 26.20	-1.09	0.279		
		Mass*sex	0.10 ± 0.11	1.00	0.323		

*denotes significant p-value at α = 0.05

Table 3. Model coefficients for linear models testing whether a body condition index (BCI) derived from zygomatic width (ZW index) explains observed variation in fat and lean mass in black-tailed prairie dogs pre-winter. Reference group for sex: female.

Component	Model	Independent terms	Estimate	t	p	AICc	Adjusted R²
Fat	~ BCI*sex	Intercept	517.47 ± 10.48	49.36	< 0.001 *	286.9	0.649
		BCI	69.42 ± 12.04	5.77	< 0.001 *		
		Sex	-21.91 ± 16.43	-1.33	0.195		
		BCI*sex	20.57 ± 25.45	0.808	0.427		
	~mass*sex	Intercept	-397.6 ± 128.5	-3.41	0.005 *	277.4	0.753
		Body mass (g)	0.628 ± 0.09	7.14	< 0.001 *		

		Sex	-109.1 ± 227.0	-0.48	0.635		
		Mass*sex	-0.00 ± 0.15	-0.00	0.999		
Lean	~ BCI*sex	Intercept	727.53 ± 12.38	58.79	< 0.001 *	295.9	0.817
		BCI	47.44 ± 14.21	3.34	0.003 *		
		Sex	201.26 ± 19.39	10.38	< 0.001 *		
		BCI*sex	-56.48 ± 30.04	-1.88	0.073		
	~mass*sex	Intercept	-56.00 ± 140.98	-0.40	0.695	282.4	0.889
		Body mass (g)	0.54 ± 0.10	5.57	< 0.001 *		
		Sex	675 ± 249.01	2.71	0.013 *		
		Mass*sex	-0.34 ± 0.16	-2.14	0.043		

687 *denotes significant p-value at α = 0.05*
688

704 **Table 4.** Model coefficients for linear models testing whether a body condition index (BCI) derived from

705 right hind foot length (RHF index) explains observed variation in fat and lean mass in Columbian ground

706 squirrels pre-winter. Reference group for sex: female.

Component	Model	Independent terms	Estimate	t	p	AICc	Adjusted R^2
Fat	~ BCI*sex	Intercept	116.42 ± 4.53	25.71	< 0.001 *	265.2	0.80
		BCI	15.68 ± 5.16	3.04	0.005 *		
		Sex	62.85 ±6.40	9.81	< 0.001 *		
		BCI*sex	1.78 ± 7.09	0.25	0.804		
	~mass*sex	Intercept	-49.70 ± 52.39	-0.95	0.352	266.7	0.80
		Body mass (g)	0.30 ± 0.09	3.18	0.004 *		
		Sex	-10.43 ± 93.38	-0.11	0.912		
		Mass*sex	0.04 ± 0.14	0.26	0.794		
Lean	~ BCI*sex	Intercept	388.37 ± 5.73	67.81	< 0.001 *	279.3	0.83
		BCI	34.54 ± 6.52	5.30	< 0.001 *		
		Sex	83.35 ± 8.1	10.29	< 0.001 *		
		BCI*sex	-14.23 ± 8.96	-1.89	0.124		
	~mass*sex	Intercept	12.61 ± 48.80	0.26	0.789	262.5	0.90
		Body mass (g)	0.67 ± 0.09	7.73	< 0.001 *		
		Sex	78.36 ± 87.00	0.90	0.376		
		Mass*sex	-0.14 ± 0.13	-1.053	0.302		

707 *denotes significant p-value at α = 0.05*
708

714   **Table 5.** Model coefficients for linear models testing whether a body condition index (BCI) derived from

715   right hind foot length (RHF index) explains observed variation in fat and lean mass in Columbian ground

716   squirrels in spring. Reference group for sex: female.

Component	Model	Independent terms	Estimate	t	p	AICc	Adjusted R^2
Fat	~ BCI*sex	Intercept	29.50 ± 2.03	14.56	< 0.001 *	664.6	0.59
		BCI	8.34 ± 2.23	3.74	< 0.001 *		
		Sex	30.16 ± 3.18	9.50	< 0.001 *		
		BCI*sex	0.50 ± 3.36	0.148	0.883		
	~mass*sex	Intercept	-35.48 ± 16.84	-2.11	0.038 *	664.9	0.59
		Body mass (g)	0.17 ± 0.04	3.87	< 0.001 *		
		Sex	23.41 ± 27.65	0.85	0.400		
		Mass*sex	-0.02 ± 0.06	-0.35	0.729		
Lean	~ BCI*sex	Intercept	310.09 ± 2.55	121.85	< 0.001 *	701.5	0.89
		BCI	34.54 ± 6.52	12.41	< 0.001 *		
		Sex	70.46 ± 3.99	17.67	< 0.001 *		
		BCI*sex	5.95 ± 4.22	1.41	0.163		
	~mass*sex	Intercept	22.57 ± 14.30	1.58	0.119	638.3	0.95
		Body mass (g)	0.76 ± 0.04	20.25	< 0.001 *		
		Sex	-2.22 ± 23.47	-0.10	0.925		
		Mass*sex	-0.001 ± 0.05	-0.15	0.880		

717   *denotes significant p-value at α = 0.05
718

 Table 6. Model coefficients for a set of linear models testing the interactive and singular effects of body

 condition index (BCI) or body mass (scaled within species) and species on fat and lean mass in red

 squirrels, prairie dogs, and ground squirrels (pre-winter and spring). Reference species: red squirrels.

Predictor	Component	Independent terms	Estimate	t	p	AICc	Adjusted R^2
BCI	Fat (g)	Intercept	6.70 ± 2.70	2.485	0.014 *	1981.6	0.98
		BCI	1.57 ± 2.84	0.53	0.596		
		Species (pre-winter prairie dogs)	501.84 ± 5.30	-94.78	< 0.001 *		
		Species (pre-winter ground squirrels)	-141.14 ± 5.10	27.70	< 0.001 *		
		Species (spring ground squirrels)	35.09 ± 3.77	9.31	< 0.001 *		
		BCI*species (pre-winter prairie dogs)	72.52 ± 6.63	10.94	< 0.001 *		
		BCI*species (pre-winter ground squirrels)	15.11 ± 5.55	2.73	0.007 *		
		BCI*species (spring ground squirrels)	7.05 ± 4.00	1.77	0.079		
	Lean (g)	Intercept	215.94 ± 5.74	37.70	< 0.001 *	2305.3	0.93
		BCI	12.56 ± 6.02	2.09	0.039 *		
		Species (pre-winter prairie dogs)	593.58 ± 11.24	52.80	< 0.001 *		
		Species (pre-winter ground squirrels)	214.10 ± 10.82	19.79	< 0.001 *		
		Species (spring ground squirrels)	122.85 ± 8.00	15.36	< 0.001 *		
		BCI*species (pre-winter prairie dogs)	22.24 ± 14.07	1.58	0.115		
		BCI*species (pre-winter ground squirrels)	14.43 ± 11.79	1.23	0.222		
		BCI*species (spring ground squirrels)	24.84 ± 8.48	2.93	0.004 *		
Body mass (scaled)	Fat (g)	Intercept	6.69 ± 2.54	2.64	< 0.001 *	1954.9	0.98
		Body mass (scaled)	1.40 ± 2.62	0.54	< 0.001 *		
		Species (pre-winter prairie dogs)	501.85 ± 54.98	100.84	< 0.001 *		
		Species (pre-winter ground squirrels)	141.15 ± 4.79	29.47	< 0.001 *		

	Species (spring ground squirrels)	35.10 ± 3.54	9.91	< 0.001 *		
	Mass *species (prairie dogs)	61.61 ± 65.16	11.93	< 0.001 *		
	Mass*species (pre-winter ground squirrels)	13.74 ± 4.95	2.77	0.006 *		
	Mass *species (spring ground squirrels)	6.92 ± 3.62	1.91	0.057		
Lean (g)	Intercept	215.83 ± 5.49	39.29	< 0.001 *	2287.3	0.94
	Body mass (scaled)	13.42 ± 5.67	2.37	0.019		
	Species (pre-winter prairie dogs)	593.69 ± 10.78	55.06	< 0.001 *		
	Species (pre-winter ground squirrels)	214.21 ± 10.38	20.65	< 0.001 *		
	Species (spring ground squirrels)	122.96 ± 7.67	16.03	< 0.001 *		
	Mass *species (pre-winter prairie dogs)	27.89 ± 11.18	2.47	0.014 *		
	Mass *species (pre-winter ground squirrels)	15.67 ± 10.73	1.46	0.146		
	Mass *species (spring ground squirrels)	23.89 ± 7.85	3.05	0.003 *		

725   *denotes significant p-value at α = 0.05*