Metabolic impact of trans 10, cis 12-conjugated linoleic acid in pai transgenic mice

Yu Rao [1,2,3,#], Shi-Li Li [4,#], Mei-Juan Li [5,#], Bao-Zhu Wang [1,#], Yang-Yang Wang [1], Lu-Wen Liang [1], Shuai Yu [1], Zong-Ping Liu [1], Sheng Cui [3], Ke-Mian Gou [1,2,3,*]

[1] *Institute of Comparative Medicine,* [2] *Jiangsu Co-innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses,* [3] *Institute of Reproduction and Metabolism, College of Veterinary Medicine, Yangzhou University, Yangzhou 225009, China*;

[4] *Department of Internal Medicine, the University of Texas Southwestern Medical Center, Dallas, Texas 75390, USA*;

[5] *Institute of Animal Husbandry and Veterinary Science, Guizhou Academy of Agricultural Sciences, Guiyang 550005, China.*

[#] YR, SLL, MJL, and BZW contributed equally to this work.

* Corresponding author: Ke-Mian Gou, College of Veterinary Medicine, Yangzhou University, Yangzhou 225009, China. **Email**: gou@yzu.edu.cn

Declarations of interest: none

Abbreviations: CLA, conjugated linoleic acid; LA, linoleic acid; t10c12, *trans* 10, *cis* 12; pai, *Propionibacterium acnes* isomerase; wt, wild-type; FGF21, fibroblast growth factor 21; UCP, uncoupling protein; AMPK, AMP-activated protein kinase; BAT, brown adipose tissue; WAT, white adipose tissue; FAs, fatty acids

BAT: mass ↑, thermogenesis ↑

Hypothalamus: aberrant gene expression

Heat release ↑
Activities ↓

Blood: triglycerides ↓ FGF 21 ↑

Liver: weight ↑ no steatosis

WAT: mass ↓

PAI transgenic mice
(producing trans 10, cis 12-CLA)

Graphical Abstract

1. A transgenic mouse itself can produce trans 10, cis 12-conjugated linoleic acid.

2. Mice showed more heat release and few activities.

3. Conjugated linoleic acid induces excess FGF21 and reduced triglycerides in the blood.

4. Conjugated linoleic acid promotes the mass and thermogenesis of brown adipose tissues.

5. Conjugated linoleic acid results in aberrant gene expression in the hypothalamus.

Highlight

1   **Abstract**

Introduction: Trans 10, *cis* 12-conjugated linoleic acid (t10c12-CLA) from ruminant-

derived foodstuffs can induce body fat loss in mammals after oral administration, while its

mechanism on fat reduction has yet to be clarified fully until now.

Methods: In the current study, a transgenic mouse that produces t10c12-CLA has been

generated via the insertion of the Propionibacterium acnes isomerase (Pai) expression cassette

into the Rosa26 locus, and its male offspring were used to decipher an irreversible long-term

impact of t10c12-CLA on health and its mechanism of action.

Results: Compared to their wild-type C57BL/6J littermates, comprehensive phenotype

profiling of biallelic pai/pai mice indicated that white fat was decreased while brown fat was

increased reversely; meanwhile, more heat was released and the central activities were reduced.

Besides decreased plasma triglycerides in both pai genotypes and increased serum FGF21 in

pai/wt mice, RNA and protein analysis revealed that the thermogenesis capacity of brown

adipose tissues was increased via pAMPK and UCP2 over-expression.

Conclusion: The results indicate that the t10c12-CLA-induced fat loss might be caused

by the excess FGF21 and the increased mass and extra thermogenesis of brown adipose tissue

in transgenic mice.

Keywords: conjugated linoleic acid; adipose tissue; obesity; thermogenesis; FGF21;

hypothalamus

1. Introduction

Trans 10, *cis* 12-conjugated linoleic acid (t10c12-CLA), one of the isomers of linoleic acid (18:2n-6, LA), is naturally present in ruminant-derived foodstuffs. It is estimated that a person can take about 9~20 mg of t10c12-CLA from foodstuffs daily [1]. Since the effect of t10c12-CLA on body fat loss in mice [2] and milk fat depression in cows [3] was reported, numerous studies tried to explore its mechanism on lipid metabolism. Some studies revealed that t10c12-CLA induces fat reduction by reducing food intake, modulating lipid metabolism, increasing energy expenditure, and/or browning WAT [4-6]. Still, some of these reports showed contradictory results sometimes. That is, the mechanism of action of t10c12-CLA on fat reduction has yet to be clarified fully until now.

Meanwhile, other studies also indicated that t10c12-CLA influenced the metabolism or physiological roles of some critical fatty acids (FAs), such as palmitic, stearic, oleic, linoleic, arachidonic, docosahexaenoic acids and/or their derived leukotrienes signals/hormonal molecules via affecting their contents in various tissues [7-10] based on the level of dietary fat and the degree of host obesity [9, 11]. Furthermore, these studies in mice fed with a t10c12-CLA diet revealed approximately 0.1-1.5% of t10c12-CLA in the brain and eye [12], milk [8], adipose tissues [7], liver [7, 9], heart, spleen, kidney, muscle, and serum [9], suggesting that t10c12-CLA might have a systemic impact on the body. Actually, besides anti-obesity, a few studies also showed the connections between t10c12-CLA and hepatic steatosis [5, 13], pro-inflammatory effects [13], and even intestinal microbiota in mice [14], suggesting the potential risk of use of t10c12-CLA.

For naturally t10c12-CLA present at very low concentrations, most t10c12-CLA studies or applications used a chemically CLA mixture prepared by industrial isomerisation. CLA

mixture usually contains approximately equal amounts of t10c12-CLA and *cis* 9, *trans* 11-CLA

(c9t11-CLA), typically 80–95% of total CLA, and other minor isomers, such as *trans*, *trans*

isomers [6]. To reveal the whole and exact t10c12-CLA's impact that might be offset or masked

by c9t11-CLA [6] or trans fatty acids [15] in the CLA mixtures, we seek the possibility of

producing t10c12-CLA by an animal itself so we might investigate it's long-term impact and

mechanism of action in a novel way.

Based on the reports that *Propionibacterium acnes* isomerase (PAI) can successfully

convert LA into t10c12-CLA in recombinant lactic acid bacteria [7, 9] and transgenic plants

[16], We produced t10c12-CLA efficiently in murine 3T3 cells [17]. Then we established a

pai-transgenic mouse previously [18]; unfortunately, the pai-expressing cassette was randomly

integrated into the Myh11 gene in this mouse, resulting in the death of homozygous mice after

birth due to gastric inflation and bladder without urinating ability [19].

In the present study, we produced a novel transgenic founder in which the pai-expressing

cassette was knocked into the Rosa26 locus by the CRISPR-Cas9 technique. The results

indicated that the pai/pai mice reduced white fat and lowered triglycerides (TGs) in the blood

but had no changes in hormones involved in energy intake, stress response, inflammation, and

fatty liver, suggesting long-term exposure to t10c12-CLA is safe, at least in male mammals.

Furthermore, we first revealed that increased FGF21 in the blood and more heat released by

brown fat resulted in white fat reduction in pai mice. The results shed light on the mechanism

of action of t10c12-CLA. Moreover, aberrant expression of the hypothalamic genes suggests

that t10c12-CLA could have other potential impacts on health. Thus, the use of t10c12-CLA

in anti-obesity practice or clinic trials should be closely monitored to ensure proper and safe

application.

2. Materials and Methods

2.1. Chemicals

Unless otherwise stated, all chemicals were purchased from Sigma (USA) or Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). Restriction endonucleases were purchased from New England Biolabs Inc (USA).

2.2. Vector construction

The pRosa-pai vector was constructed as follows. In brief, based on the nucleotide sequence of the Pai gene of *Propionibacterium acnes* (GenBank accession number AX062088)*,* the codon-optimized Pai gene was synthesised, which added the 6 x His affinity tag sequence immediately upstream of the terminal codon. It has been confirmed that gene transfection with this vector in murine 3T3 cells led to t10c12-CLA production [17]. On this basis, the Pai expression vector was firstly generated after insertion into the *Eco*R I site of the pCAGGS vector (Gift from Dr Timothy J. Ley, The Washington University, St. Louis), in which the Pai gene was under the control of the cytomegalovirus enhancer and the chicken beta-actin promoter (CAG promoter). After two molecular cloning steps of enzymic modification, the CAG-pai expression cassette was inserted into the Pac I and Asc I sites of the pROSA26-PA backbone vector (Gift from Dr Frank Costantini, Addgene plasmid#21271; http://n2t.net/addgene:21271) to generate the pRosa-pai vector.

The pRosa-Cas9 vector was constructed as follows. Briefly, the CAG promoter sequence of Cas9 in pDG330 plasmid (Gift from Dr Paul Thomas; Addgene plasmid # 100898; http://n2t.net/addgene:100898) was replaced by the EF-1α promoter from the plasmid pEF.myc.ER-E2-Crimson (Gift from Dr Benjamin Glick; Addgene plasmid # 38770;

http://n2t.net/addgene:38770) followed by insertion of the sgRNAs sequences (5-

actccagtctttctagaaga-3, from reference [20]) into Bbs I sites to generate the pRosa-Cas9 vector

which included the Cas9 gene driven by EF-1α promoter and dual gRNA expression cassettes.

2.3. Pronuclear microinjection

Purified pRosa-Cas9 plasmid DNA at a final concentration of 10 ng/μl and the fragments

of pRosa-pai plasmid after digestion by Pvu I + Sal I enzymes at a final concentration of

50~100 ng/μl were mixed well and used for pronuclear co-microinjection immediately.

2.4. Mice

Mice were obtained from the Laboratory Animal Centre, Yangzhou University, China.

All animal experiments were performed following the Committee for Experimental Animals

of Yangzhou University. Mice were maintained in a light-controlled room (12L:12D, lights on

at 0700 h) at a temperature of 22-23°C and were fed ad libitum with a standard diet containing

10% kcal% fat.

Embryos from superovulated D6B2F1 females mated with C57BL/6J male mice were

used to produce the transgenic mice. The transgenic founders were backcrossed to C57BL/6J

mice more than eight generations before analysis in the current study. The biallelic (pai/pai),

monoallelic (pai/wt), and wild-type (wt) male offsprings from the pai/wt matings were used in

this study.

2.5. Diets

New diets containing 10% kcal% fat were prepared per month under sterile conditions

according to formula No. D12450H of OpenSource DIETS™ (Research Diets Inc., NJ, USA).

The formula D12450H and the sources of the food-grade ingredients for the diets are listed in

Supplementary Table 1. Fatty acid compositions in diets were spot-checked by gas

chromatography to examine the contents of linoleic acid and t10c12-CLA. One kilogram of

diet contained approximately 20 g of LA, and no t10c12-CLA was detected in diet samples.

2.6. Transgene characterisation

Genomic DNA from the tail tissues was used for nucleic acid analysis. The presence of

the pai transgene was assayed to amplify the 514-bp pai fragments (All primer sequences and

annealing temperatures were listed in Supplementary Table 2). The same insertion site was

assayed to amplify the 605-bp fragments that spanned the insertional Rosa26 area. The PCR

amplification was performed as follows: 30 cycles at 94°C for 30 sec, 57°C for 30 sec, and

72°C for 40 sec.

Southern blot was performed according to the standard procedure. In brief, 10 μg of

genomic DNA was first digested with the EcoR I enzyme for 12-16 h at 37°C and subjected to

1% agarose gel electrophoresis, then transferred onto the Hybond-N$^+$ membrane (Amersham,

UK). The membrane was subsequently hybridised with the DIG (Roche, Germany) labelled

probes according to the manufacturer's protocol, using alkaline phosphatase labelled anti-Dig

antibody (Roche) and CDP-STAR solution to develop the photos.

2.7. Western blot

Western blot was performed according to the standard procedures. Briefly, total protein

extracts from 20 mg tissues were quantified, and 20 μg protein extracts were separated with

10-15% SDS-PAGE gel and transferred to the PVDF membranes (Millipore ISEQ00010,

Merck). Membranes were blocked and hybridised with the respective primary antibody alone

or mixed with the anti-beta-actin antibody (Proteintech Group, Inc, China, 20536-1-AP)

followed by HPR-labelled goat anti-rabbit IgG (Santa Cruz Biotechnology, sc-2004). The

image was developed using the SuperSignal™ West Femto substrate (ThermoFisher, USA). After subtracting the local background, the bands were quantified using Image software (Media Cybernetics, USA). The primary antibodies included rabbit antibodies to His-tag (Proteintech Group, 10001-0-AP), CPT1B (Proteintech, 22170-1-AP), UCP2 (Proteintech, 11081-1-AP), FGF21 (Abcam Inc, USA, ab171941), pAMPK (Thr183 + Thr172, Abcam, AB133448), AMPK (Abcam, ab207442), FASN (Abcam, AB128870), UCP1 (Abcam, ab234430), GRP78 BiP (Abcam, ab108613), PPAR-Γ (Wanleibio Ltd, China, WL01800), PGC1A (Wanleibio, WL02123), and Phospho-HSL (Ser563, Cell Signaling Technology, #4139).

2.8. RNA analysis

Total RNA was extracted from tissues using the RNAiso Plus kit and treated with DNase I (TaKaRa, China). The purified RNA was used for first-strand cDNA synthesis, and reverse transcription was performed using an M-MLV reverse transcriptase with oligo-dT primers according to the manufacturer's instructions (Promega, USA). For RT-PCR, the resulting cDNA was used for PCR amplification with the Pai-specific primers (Supplementary Table 2) that produced a 164-bp fragment. A 78-bp fragment of the Gapdh gene was amplified under the same conditions and used as an endogenous control. PCR amplification was performed as follows: 30 cycles of 94°C for 30 sec, 60°C for 30 sec, and 72°C for 30 sec.

All real-time PCRs were carried out in 96-well plates using ChamQ SYBR qPCR Master Mix kit in the ABI Prism 7500 Sequence Detection System (Applied Biosystems, USA) at 95°C, 2 min, one cycle; 95°C, 5 sec, 60°C, 32 sec, 40 cycles. Three replicates were included for each sample. The relative transcriptional level of a target gene was normalised to the one of endogenous gene expression of Gapdh, ApoB, or 36B4 using the method of $2^{-\Delta\Delta Ct}$. All genes' full names and these primer sequences are listed in Supplementary Table 3.

2.9. RNA sequencing

The RNA-seq was used to identify whole transcriptome differences in the BAT from wt (n=2) and pai/pai (n=3) mice at 12 weeks. Total RNAs were analysed by BGI Genomics company (China) using the DNBSEQ platform. Clean reads were aligned to *Mus musculus* genome GCF_000001635.26_GRCm38.p6, and the expression level was normalised as FPKM with gene annotation file. Differential expressional genes and functional enrichment for KEGG had plotted an online platform for data analysis and visualisation (http://www.bioinformatics.com.cn).

2.10. Gas chromatography

According to the modified method described by Jenkins [21], the procedure of two-step trans-esterification that utilised sodium methoxide followed by shorter methanolic HCl was used to methylate each organ/tissue homogenised by grinding in liquid nitrogen. Fatty acid methyl esters were separated on an HP-88 fused-silica capillary column (60 m X 0.25 mm i.d., 0.2-µm film thickness, J & W 112-88A7, Agilent Technologies, USA) and quantified using a fully automated 7890 Network GC System with a flame-ionization detector (Agilent). The program setting details followed Jenkins' method [21]. C19:0 was used as the internal standard, and the peaks were identified by comparison with FA standards (Sigma, 47885U and O5632). The area percentage for all resolved peaks was analysed using GC ChemStation Software (Agilent).

2.11. Hepatic parameters measurements

Hepatic tissues (0.1 g) were homogenised and resuspended in PBS solution. The concentrations of total cholesterol (TC) and triglycerides (TGs) were respectively measured using the corresponding detecting kits (Meimian Industrial Co. Ltd., China).

2.12. Blood parameters measurements

Unless otherwise stated, all blood samples were collected from the 11~15-week-old mice fed ad libitum. Fresh blood from the tail veins was used to measure the circulating glucose using a hand-held glucose monitor (Accu-Chek® Performa Blood Glucose Meter, Roche). Heparin-treated blood from the tail vein was respectively used to measure the plasma TGs or high-density lipoprotein (HDL) using a hand-held cholesterol monitor (On-Call® CCM-111 Blood Cholesterol System, Aikang Biotech Co. Ltd., China). Blood samples from the submandibular vein of conscious mice were collected at the first 4~5 hrs of the light phase and used to measure the serum corticosterone using the Cort ELISA kit (Ruixin Biotech Co. Ltd., China). Serums from the orbital sinus vein of anaesthetic mice were used to measure the circulating TC, free fatty acids (FFA), prostaglandin E2 (PGE2) using the respective ELISA kits (Meimian), as well as insulin, ghrelin, leptin, fibroblast growth factor 21 (FGF21), interleukin-6 (IL-6), adrenaline glucagon tumour necrosis factor-alpha (TNFα), and C reactive protein (CRP) using the respective ELISA kits (Ruixin).

2.13. Intraperitoneally glucose tolerance test

Mice fasted overnight were intraperitoneally injected with D-glucose (2 g/kg body weight). Blood glucose was measured in a drop of blood collected from the tail vein before, 15, 30, 60, 90, and 120 min post glucose injection.

2.14. Histological analysis

Histological analysis was performed following the standard methods. Briefly, the tissues were fixed by immersion in 4% formaldehyde in PBS and then dehydrated before embedding. The 5-μm tissue sections (Leica CM1950) embedded with paraffin were dewaxed in xylene and rehydration. Histological details were examined using hematoxylin-eosin staining kits

(Beijing Solarbio Science & Technology Co., Ltd., China). For red oil staining, 10-μm tissue

slices embedded by Tissue-Tek® O.C.T. Compound (Sakura, USA) were treated with a 60%

isopropyl alcohol solution twice, and histological details were examined after staining with oil

red O (Wuhan Servicebio Technology Co., Ltd., China). Stained slides were examined with an

Olympus microscope equipped with a digital camera.

Each cellular cross-sectional area was estimated following the method described by Chen

and Farese [22]. In detail, at least six slices from each mouse sample were randomly chosen

for image analysis. Firstly, three photos were randomly taken from non-overlapping

microscope fields of every slice under the 20 x objective. The cross-sectional area of each cell

on an image was calculated one by one using ImageJ software (Version 2.0.0), and the means

of the top 100 cell areas per picture represented the value of a microscope sampling field.

Finally, the means of at least 18 values per mouse sample (3 fields x 6 slices) were looked upon

as this sample's average cellular cross-sectional area for statistical analysis.

2.15. Metabolic cage measurements

Indirect calorimetry, heat production, and movement of 11-week-old mice were

measured using the Automated Home Cage Phenotyping TSE PhenoMaster V4.5.3 system

(TSE Systems Inc., Germany). The climate chamber was set to 22°C with a 12-h:12-h light-

dark cycle (lights on at 0700 h). Mice were individually housed in plexiglass cages and fed ad

libitum. VO_2, VCO_2, and food intake were measured every 39 min. After a 48 hrs acclimation

period, the data were collected for the following 72 hrs, and the calculated lean mass was

adjusted for all measurements.

2.16. Statistical analysis

All values are presented as mean $\pm$ SD. The statistical significance was assessed by

unpaired student's t-test or one-way ANOVA (Brown-Forsythe and Welch) among wt, pai/wt,

and pai/pai mice. A value of $p < 0.05$ was considered statistically significant.

Figure 1. Transgene analysis. (A) Strategy for inserting the pai-expressing cassette into the Rosa26 locus by the CRISPR/Cas9 technique. The upper panel illustrates the recombined homological vector in which the codon-optimised Pai gene is driven by a CAG promoter and ended by a rabbit beta-globin polyA signal (rβ-pA). Rosa L and Rosa R represent the left and right sequences of the insertional site within the first intron of Rosa26, respectively. Linear DNA fragments digested with Pvu I and Sal I enzymes are used for co-microinjection. The down panel shows the CRISPR/Cas9 plasmid containing EF-1α-driven Cas9 and dual gRNA target sequences for co-microinjection. The numbers in brackets indicate the sequence length. (B) PCR and (C) Southern blot analysis of the genomic DNA samples from transgenic (tg), wild-type (wt) mice, and wt DNA mixed with a pRosa-Pai plasmid (P) shows that the Pai gene is inserted into

the Rosa26 locus in the tg sample. PCR products are 514-bp fragments spanning the Pai gene and 605-bp fragments spanning the insertional site of the Rosa26 sequence. (D) RT-PCR analysis of RNAs shows that the Pai gene is expressed in all tg tissues, including brown (BAT) and white (WAT) adipose tissues, not in all wt samples. The amplified product of Pai is 164-bp fragments, and the Gapdh is 78-bp fragments as an internal control. (E) Western blot analysis reveals that PAI antigen existed in the tg heart, liver, spleen, and kidney, not in wt samples. (F) The lipid profiles of partial gas chromatography traces show that the t10c12-CLA is produced in the kidney of the tg mouse compared to its wt littermate. (G) Real-time PCR analysis showed that the relative mRNA expression level of the Pai gene is quite different in each sample from biallelic pai/pai or monoallelic pai/wt mice (n=3), and no Pai mRNA is detected in samples of their wt littermates. Each value represents the mean ± SD after normalisation to the mRNA levels of Gapdh.

3. Results

3.1. Transgene analysis

The strategy to insert the pai-expressing cassette into the Rosa26 locus is shown in Figure 1A. PCR and Southern blot (Fig. 1 B-C) and DNA sequencing analysis demonstrated that the correct transgene was inserted into the Rosa26 locus in a founder mouse among 96 mice derived from injected embryos. RT-PCR and western blot analyses (Fig. 1 D-E) indicated that the pai gene was expressed in all tested tissues. Gas chromatography verified that the functional PAI enzyme produced the t10c12-CLA successfully (Fig. 1 F). Real-time PCR analysis (Fig. 1 G) revealed that the mRNA levels of Pai were quite different between biallelic pai/pai and monoallelic pai/wt mice; thus, two pai genotypes were analysed separately in the following tests.

3.2. Fatty acids analysis

T10c12-CLA and FAs compositions in the hearts, livers, kidneys, tibialis anterior muscle, BAT, or WAT were analysed by gas chromatography. Compared to wt littermates, the contents (μg/g of tissue weight) of t10c12-CLA had increased in the kidneys ($p = 0.004$) and heart ($p = 0.055$) from pai/wt mice and WAT from pai/wt or pai/pai mice ($p < 0.05$; Table 1), not in the other examined tissues, suggesting that the t10c12-CLA is produced in trace amounts or degraded quickly. Simultaneously, the quantities of LA substrates had decreased in the livers from both pai genotypes and increased in the pai/wt kidneys ($p < 0.05$; Table 1).

The contents of other FAs, such as myristic (14:0), palmitic (16:0), palmitoleic (16:1n-7), stearic (18:0), oleic (18:1n-9), cis-vaccenic (18:1n-7), or arachidonic (20:4n-6) acids, were changed by varying degrees in pai heart, kidney, liver or WAT tissues (Supplementary Table

Table 1. Contents of t10c12-CLA and linoleic acid in wt and pai mice

	T10c12-CLA (µg/g)			Linoleic acid (mg/g)		
	wt (n)	pai/wt (n)	pai/pai (n)	wt (n)	pai/wt (n)	pai/pai (n)
Heart	7.12 ± 0.95 (11)	8.85 ± 1.20 (8)	8.47 ± 2.35 (8)	2.06 ± 0.31 (11)	2.03 ± 0.31 (8)	2.11 ± 0.19 (8)
Liver	11.93 ± 8.27 (12)	16.65 ± 12.53 (11)	9.93 ± 1.31 (10)	1.29 ± 0.20[a] (12)	1.09 ± 0.25[b] (11)	1.07 ± 0.23[b] (10)
Kidney	6.77 ± 1.16[a] (12)	11.33 ± 4.18[b] (13)	6.87 ± 1.17[a] (8)	1.13 ± 0.16[a] (10)	1.38 ± 0.26[b] (13)	1.20 ± 0.19[a] (8)
Muscle	ND (9)	ND (8)	ND (8)	0.87 ± 0.41 (9)	0.84 ± 0.31 (8)	0.61 ± 0.17 (8)
BAT	25.87 ± 16.37 (8)	25.63 ± 10.00 (8)	28.80 ± 7.52 (7)	1.88 ± 0.31 (8)	1.70 ± 0.42 (8)	1.69 ± 0.31 (7)
WAT	ND (5)	4.10 ± 1.05 (3)	4.89 ± 0.83 (3)	0.50 ± 0.34[ab] (5)	0.33 ± 0.01[a] (3)	0.44 ± 0.01[b] (3)

Note: Tissues of mice at the age of 11 weeks are used for gas chromatography, and 19:0 is an internal standard. ND, Not detected. Each value represents the mean ± SD, and the numbers in brackets indicate the size of the tested samples. For all fatty acid contents, see Supplementary Table 4. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

4). In contrast, no evident changes in FA compositions were observed in pai muscle and BAT

tissue, suggesting the content changes of some FAs are genotype- or tissue-specific.

Simultaneously, the amounts of total FAs were decreased in the pai/pai livers (9.19 ± 1.55 in

wt versus 8.19 ± 1.78 mg/g in pai/pai; $p < 0.05$) and contrarily increased in both pai kidneys

(9.59 ± 0.99 in wt versus 12.97 ± 2.21 in pai/wt versus 10.66 ± 1.06 in pai/pai; $p < 0.05$;

Supplementary Table 4), suggesting fat loss in livers but fat accumulation in kidneys.

3.3. Weaning and growth analysis

Considering the potential impact of maternal t10c12-CLA from the placenta and/or milk,

pups delivered by wt or pai/wt mothers were measured separately. In the group of wt mothers

(Fig. 2A), the pai/wt pups showed no bodyweight difference from the wt littermates during the

post-weaning development. In the group of pai/wt mothers (Fig. 2B), the pai/wt or pai/pai pups

also showed no bodyweight difference from the wt littermates during the post-weaning

development. However, the pai/wt genotypes were overweight ($p < 0.05$) than pai/pai mice

from 10 weeks onwards.

Interestingly, within the group of pai/wt mothers, the weaning weights of three genotypes

were significantly different (pai/pai < pai/wt < wt, respectively; $p < 0.05$. Fig. 2 B-C),

suggesting the endogenous t10c12-CLA produced by foetus/pups themselves would result in

the reduced bodyweight during embryonic/weaning development. On the other hand, the

weaning weights of wt or pai/wt pups from pai/wt mothers were respectively overweight than

that from wt mother ($p < 0.05$; Fig. 2C), suggesting that the maternal t10c12-CLA might result

in the increased bodyweight of the embryos/pups.

3.4. Energy metabolism and activity studies

Figure 2. Growth curves of different genotype pups from wt (A) or pai/wt (B) mother and comparison of the body weight (C) of pups at the age of 3 weeks. Each group contained 15-20 (A, B) or 30-85 (C) pups. Bars represent the mean ± SD. # and different letters indicate $p < 0.05$ among three genotypes; * indicates $p < 0.05$ between pai/pai and pai/wt genotypes.

Figure 3. Pai mice have abnormal energy metabolism and activities.

Comparison of body weight gain during 72 h period (A), food intake (B), water intake

(C), total distance travelled during 72 h period (D), O_2 consumption (E), CO_2

production (F), heat release and its time course (G-H), respiratory exchange ratio (REF) and RER time course (I-J), locomotor activities in the centre of the cage and its time course (K-L), central ambulating and stereotypic activities (M-N), locomotor activities in the margin or corners of the cage and its time course (O-P), marginal ambulating and stereotypic activities (Q-R), distance travelled during a 39-min sampling duration (S), and speed (T) among wt (n=5) and pai/pai (n=5) mice at the age of 11 weeks. Lights turn on and off at 7:00 am and 7:00 pm, respectively. The data are normalised to lean body weight. Bars represent the mean $\pm$ SD. * indicate $p < 0.05$ within the same phase.

Energy metabolism and activity parameters were determined under the consideration of

body weight in the 11-week-old pai/pai mice and wt littermates over 72 hrs in metabolic cages

(Fig. 3). Compared to the wt littermates, pai/pai mice showed no different ($p > 0.05$) changes

in food intake, respiratory exchange ratio, marginal activities, and speed in both light and dark

phases; as well as body weight gain and total distance travelled during the 72 hrs period.

However, pai/pai mice had fewer central activities significantly ($p < 0.05$) in both light

(56%) and dark (26%) phases, but they consumed 32% and 48% more water, 4.1% and 9.9%

more O_2, and produced 4.3% and 12% more CO_2, and 4.2% and 10.4% more heat during the

light ($p > 0.05$) and dark ($p < 0.05$) periods, respectively (Fig. 3). That was, pai/pai mice

reduced their activities, but consumed more oxygen (energy) to release more heat for

maintenance.

3.5. Metabolic complications

To match energy demand and expenditure, the circulating hormonal signals were

examined under the non-fasted state. The concentrations of serum insulin, leptin, ghrelin, and

circulating glucose (Table 2), as well as the length of the small intestine (Supplementary Table

5), showed no differences ($p > 0.05$) among the three genotypes. It suggests that the t10c12-

CLA does not play a role in the energy intake, consistent with the observation in the metabolic

cage.

Glucose tolerance tests showed no difference in the circulating glucose concentrations or

the area under the curve among the three genotypes (Fig. 4 A). However, the blood glucose

levels before glucose injection were remarkably ($p < 0.05$; Fig. 4 A) increased in pai/wt mice

and maintained typically when fasted 24-h duration (Fig. 4 B). It suggests that the t10c12-CLA

might ameliorate the glucose sensitivity in pai/wt mice at the beginning of starvation, which

Table 2. Hormones/factors in blood in wt and pai mice

	wt (n)	pai/wt (n)	pai/pai (n)
Serum total cholesterol (μM)	304 ± 32 (12)	303 ± 44 (9)	301 ± 50 (8)
Plasma triglycerides (mg/dL)	126 ± 27ᵃ (33)	107 ± 30ᵇ (30)	110 ± 28ᵇ (30)
Plasma high-density lipoprotein (mg/dL)	68.5 ± 17.0 (36)	71.3 ± 19.1 (30)	66.7 ± 17.9 (31)
Serum free fatty acids (μM)	407 ± 125 (14)	394 ± 90 (19)	465 ± 110 (10)
Blood glucose (mM)	7.98 ± 1.44 (28)	8.45 ± 0.92 (22)	7.95 ± 1.32 (26)
Serum insulin (μg/L)	0.41 ± 0.06 (12)	0.41 ± 0.08 (8)	0.44 ± 0.12 (9)
Serum leptin (μg/L)	1.05 ± 0.48 (12)	1.15 ± 0.47 (13)	1.21 ± 0.55 (11)
Serum ghrelin (ng/L)	184 ± 27 (7)	216 ± 56 (8)	194 ± 19 (9)
Serum fibroblast growth factor 21 (ng/L)	441 ± 38ᵃ (15)	480 ± 43ᵇ (12)	423 ± 27ᵃ (15)
Serum prostaglandin E2 (ng/L)	342 ± 42 (15)	327 ± 31 (9)	337 ± 42 (14)
Serum glucagon (mU/L)	75.1 ± 53.5 (14)	78.8 ± 51.4 (14)	103 ± 62 (13)
Serum cortosterone (μg/L)	9.03 ± 2.56 (16)	9.44 ± 2.44 (8)	9.44 ± 2.49 (16)
Serum adrenaline (μg/L)	4.56 ± 4.34 (11)	3.51 ± 3.06 (18)	6.30 ± 5.53 (5)
Serum interleukin-6 (ng/L)	17.3 ± 16.1 (8)	22.3 ± 23.5 (15)	23.7 ± 12.2 (9)
Serum tumour necrosis factor α (ng/L)	157 ± 139 (20)	249 ± 168 (12)	184 ± 181 (11)
Serum C reactive protein (μg/L)	20.1 ± 14.1 (9)	17.9 ± 15.8 (7)	34.0 ± 41.4 (11)

Note: Blood samples were collected from non-fasted mice at the ages of 11~15 weeks. Each value represents the mean ± SD, and the numbers in brackets indicate the size of the tested samples. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

Figure 4. Blood glucose tolerance test and dynamic glucose levels in wt and pai mice. (A) Comparison of glucose tolerance and the area under curve among wt and pai mice. (C) Dynamic blood glucose changes in fasting mice starved from Zeitgeber time 0 to 24. Zeitgeber times 0 and 12 are defined as lights-on and -off times, respectively. Bars represent the mean $\pm$ SD. # indicates $p < 0.05$ between pai/wt with wt or pai/pai mice.

may be associated with the elevated FGF21 in pai/wt mice (p < 0.05; Table 2) for FGF21 plays

an active role in insulin-sensitising and energy expenditure [23].

In addition, there were no differences in the circulating TC, FFAs, and HDL among the

three genotypes, but the serum levels of TGs were reversely decreased in pai/wt (p = 0.011) or

pai/pai mice (p = 0.021; Table 2) compared to wt mice.

Furthermore, the serum PGE2, glucagon, corticosterone, adrenaline, TNFα, IL-6, and

CRP in pai/wt or pai/pai mice were respectively kept at normal levels compared with wt mice

(p > 0.05; Table 2), suggesting no pathological symptoms such as cellular toxicity of expressing

the bacterial protein, response to illness-related stimuli, and inflammation in pai mice. It also

indicates that the phenotype profiling of pai mice is just a result of the t10c12-CLAs' impact.

3.6. WAT reduction

Mice at 11 weeks were used to determine the WAT features by dissection, magnetic

resonance (MR) imaging, histological, and RNA analyses. The results exhibited genotype-

specific and described as follows. Compared to wt mice, in pai/wt mice, no evident changes in

organs/tissues, adipocyte volume or cross-sectional areas per adipocyte (Fig. 5 B-D) were

observed. On the contrary, in pai/pai mice, organs such as livers, spleens, and kidneys were

significantly enlarged (p < 0.05; Supplementary Table 5). Simultaneously, the white fat was

lost (Fig. 5 A-B), consistent with the reduced adipocyte volume or cross-sectional areas per

adipocyte (Fig. 5 C-D), as well as the lowered RNA levels of leptin, Cebpβ (Full name seen in

Table 3) that plays a role in the early stages of adipogenesis, macrophage markers F4/80, CD68,

and CD11c in WAT (Table 3). The results suggest that the t10c12-CLA can result in fat loss

in pai/pai mice.

Figure 5. Magnetic resonance imaging and histological analysis of white adipose tissues. (A) Coronal sections were obtained on the whole-body midsection for five wt or pai/pai mice at 11 weeks in each group during magnetic resonance imaging (bar = 1 cm). The wt mice show adipose tissue in the subcutaneous and intra-abdominal regions as areas of increased signal intensity on images. The pai/pai mice show less fat. (B) The WAT weight and (C) Imaging by hematoxylin-eosin staining (bar = 10 µm), as well as (D) analysis of cross-sectional area per adipocyte, show that the volumes of white adipocytes are reduced in pai/pai mice. Bars represent the mean ± SD. Different letters indicate $p < 0.05$.

Table 3. Relative expression of mRNAs in the white (WAT) and brown (BAT) adipose tissues

Genes	WAT		BAT	
	wt (n)	pai/pai (n)	wt (n)	pai/pai (n)
Fatty acid synthase (*Fasn*)	1.01 ± 0.32 (22)	4.70 ± 2.25*** (28)	0.99 ± 0.30 (15)	1.61 ± 0.58** (16)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	1.04 ± 0.28 (17)	1.01 ± 0.46 (12)	1.01 ± 0.18 (13)	1.41 ± 0.26*** (15)
Adipose triglyceride lipase (*Atgl*)	1.08 ± 0.45 (7)	0.69 ± 0.07 (5)	1.07 ± 0.45 (7)	0.76 ± 0.44 (8)
Hormone-sensitive lipase (H*sl*)	1.04 ± 0.30 (7)	0.82 ± 0.06 (5)	1.05 ± 0.35 (8)	0.58 ± 0.23** (8)
Monoacylglycerol lipase (*Mgl*)	1.06 ± 0.41 (8)	1.05 ± 0.19 (6)	1.04 ± 0.30 (8)	0.54 ± 0.24** (8)
Perilipin 1a	1.03 ± 0.28 (7)	0.95 ± 0.16 (6)	1.15 ± 0.62 (8)	0.61 ± 0.15* (7)
Comparative gene identification 58 (*Cgi58*)	1.03 ± 0.26 (8)	1.16 ± 0.49 (7)		
Lipoprotein lipase (*Lpl*)	1.01 ± 0.12 (12)	0.59 ± 0.24*** (12)	1.00 ± 0.26 (15)	0.69 ± 0.27** (14)
Carnitine palmitoyltransferase 1a (*Cpt1a*)	1.08 ± 0.45 (7)	1.60 ± 1.13 (8)	1.03 ± 0.26 (8)	0.68 ± 0.28* (7)
Carnitine palmitoyltransferase 1b (*Cpt1b*)			1.04 ± 0.33 (7)	0.42 ± 0.21** (8)
Long-chain 3-hydroxyacyl-CoA dehydrogenase (*Lchad*)	1.05 ± 0.34 (8)	1.16 ± 0.47 (7)		
Acetyl-CoA acyltransferase 2 (*Acaa2*)	1.03 ± 0.28 (8)	1.66 ± 0.59* (7)		
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.04 ± 0.32 (17)	1.69 ± 0.65** (16)	1.01 ± 0.13 (13)	0.99 ± 0.48 (14)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.05 ± 0.38 (8)	1.18 ± 0.39 (8)	1.01 ± 0.20 (6)	1.04 ± 0.37 (7)
Uncoupling protein 1 (*Ucp1*)	0.97 ± 0.34 (8)	0.59 ± 0.36* (10)	1.04 ± 0.29 (18)	0.65 ± 0.29*** (22)
Uncoupling protein 2 (*Ucp2*)	1.04 ± 0.38 (12)	0.31 ± 0.22*** (13)	1.01 ± 0.14 (9)	0.64 ± 0.15*** (8)
Ppar-gamma coactivator 1 alpha (*Pgc1α*)	1.04 ± 0.31 (8)	1.94 ± 0.42** (5)	1.04 ± 0.31 (6)	1.63 ± 0.53* (7)
PR domain containing 16 (*Prdm16*)	1.02 ± 0.24 (6)	0.65 ± 0.21* (7)	1.04 ± 0.28 (7)	1.20 ± 0.55 (8)
Sterol regulatory element binding protein-1a (*Srebp1a*)	1.05 ± 0.33 (13)	0.98 ± 0.48 (8)	1.03 ± 0.25 (5)	0.61 ± 0.15* (5)
Sterol regulatory element binding protein (*Srebp2*)	1.03 ± 0.42 (12)	0.99 ± 0.40 (13)	1.01 ± 0.19 (5)	1.46 ± 0.81 (4)

Table 3. Relative expression of mRNAs in the white (WAT) and brown (BAT) adipose tissues

Genes	WAT		BAT	
	wt (n)	pai/pai (n)	wt (n)	pai/pai (n)
Insulin receptor	1.03 ± 0.25 (7)	1.15 ± 0.38 (8)		
Insulin receptor substrate-1 (*Irs1*)	1.00 ± 0.31 (23)	0.50 ± 0.21*** (17)		
Insulin receptor substrate-2 (*Irs2*)	1.12 ± 0.54 (8)	1.09 ± 0.50 (8)		
Phosphoinositide 3-kinase (*Pi3k*)	1.06 ± 0.36 (6)	5.11 ± 3.08* (6)		
Insulin-like growth factor-1 (*Igf1*)	1.05 ± 0.34 (8)	0.85 ± 0.32 (7)		
IGF binding protein 1 (*Igfbp1*)	1.19 ± 0.66 (6)	1.08 ± 0.42 (6)		
Glucose-6-phosphate dehydrogenase (*G6pd*)	0.97 ± 0.23 (13)	0.99 ± 0.68 (10)	1.01 ± 0.13 (5)	1.48 ± 1.02 (5)
Glucose transporter type 4 (*Glut4*)	1.01 ± 0.15 (18)	0.45 ± 0.26*** (18)	1.01 ± 0.18 (4)	3.15 ± 0.76** (4)
Carbohydrate response element binding protein (*Chrebp*)	0.98 ± 0.25 (12)	1.84 ± 0.78** (11)	1.02 ± 0.20 (5)	0.88 ± 0.33 (5)
Fibroblast growth factor 21 (*Fgf21*)	1.04 ± 0.32 (7)	1.10 ± 0.33 (7)		
Glucose-6-phosphatase (*G6p*)	0.96 ± 0.38 (16)	0.39 ± 0.17*** (24)		
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.08 ± 0.39 (8)	1.04 ± 0.67 (7)		
AMP-activated protein kinase (*Ampk*)	1.02 ± 0.35[a] (15)	1.41 ± 0.57* (15)	1.01 ± 0.17 (8)	2.24 ± 1.46* (8)
Leptin	1.01 ± 0.16 (6)	0.34 ± 0.12*** (7)		
Adiponecin	1.03 ± 0.28 (8)	0.87 ± 0.28 (11)		
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.02 ± 0.23 (8)	0.94 ± 0.39 (8)		
F4/80	1.04 ± 0.30 (28)	0.50 ± 0.27*** (30)		
CD68	1.00 ± 0.34 (31)	0.48 ± 0.31*** (36)		
CD11c	1.04 ± 0.32 (17)	0.34 ± 0.27*** (20)		
Acyl-CoA-binding protein (*Acbp*)	1.02 ± 0.20 (9)	0.67 ± 0.20** (8)	1.00 ± 0.08 (5)	1.02 ± 0.39 (5)

Table 3. Relative expression of mRNAs in the white (WAT) and brown (BAT) adipose tissues

Genes	WAT		BAT	
	wt (n)	pai/pai (n)	wt (n)	pai/pai (n)
CCAAT/enhancer-binding-protein beta (*Cebpβ*)	1.05 ± 0.37 (6)	0.58 ± 0.34* (8)		
Pai	0.99 ± 0.50 (14)	1320 ± 622*** (19)	1.08 ± 0.45 (8)	3108 ± 1504*** (9)

Numbers in brackets of the parameters indicate the size of tested samples. Each value represents the mean ± SD. * indicates $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$ within the same tissue, respectively.

In pai/pai white adipocytes, significant ($p < 0.05$) changes in RNA levels appeared in

50% (20/40) tested genes, including seven up-regulated and 13 down-regulated (Table 3). The

energy regulator Ampk was up-regulated, suggesting the inhibition of lipid synthesis, which is

consistent with the down-regulated lipoprotein lipase that supports the uptake of FAs from very

low-density lipoproteins in the bloodstream, regular expression of Srebp1a and Srebp2

involved in the sterol pathway and transcriptional factor Ppar-γ, while the fatty acid synthase

over-expressed. Genes involved in lipolyses, such as Atgl, hormone-sensitive lipase, Mgl,

Perilipin 1a, and Ggi58 kept normal RNA levels. Normal RNA levels of genes involved in

beta-oxidation, including Cpt1a, Lchad, and Mcad and the reduced RNA levels of the acyl-

CoA-binding protein that act as an intracellular carrier of acyl-CoA esters were detected in

pai/pai WAT, suggesting the weakening, not enhancing the process of beta-oxidation in

mitochondria. On the contrary, the RNA levels of Acaa2 and Lcad were increased, suggesting

the up-regulation of beta-oxidation of very long-chain FAs in peroxisomes.

In pai/pai adipocytes, although Pgc1α, which activates mitochondrial biogenesis genes

through interaction with Ppar-γ during energy stress, were up-regulated. Simultaneously,

Prdm16, Ucp1, and Ucp2 were down-regulated, suggesting the declined WAT browning.

Consistent with the normal levels of the circulating insulin or glucocorticoids, their

receptor genes Insr and Nr3c1, also expressed commonly on the pai/pai adipocyte surface;

interestingly, the Irs1 was down-regulated while its downstream Pi3K gene was up-regulated.

The RNA levels of other genes involved in insulin/IGF-1 signalling, such as Irs2, Igf1, and

Igfbp1 had no changes compared to the wt samples. Meanwhile, reduced expression of Chrebp

and glucose transporter Glut4, as well as up-regulated glucose-6-phosphatase involved in

gluconeogenesis, suggesting the energy mobilisation for glucose homeostasis. These results

indicate that the t10c12-CLA has induced the broadly positive or negative feedback regulation of lipid and glucose metabolism in WAT.

3.7. BAT thermoregulation

Parameters of BAT were measured to investigate the molecular mechanism of more heat production in pai mice. MR imaging and dissection analysis revealed that the weight (% bodyweight) of interscapular BAT was increased from 0.39 ± 0.07 in wt to 0.50 ± 0.18 in pai/wt ($p = 0.059$) and 0.54 ± 0.18 in pai/pai mice ($p < 0.05$; Fig. 6 A-C). HE staining showed that there were more small-sized and irregular lipid drops within the pai/wt or pai/pai adipocytes, quite different from the wt adipocytes containing relatively uniform and medium-sized lipid drops (Fig. 6 D); simultaneously, the size of cross-sectional area per brown adipocyte was enlarged in both pai genotypes ($p < 0.05$; Fig. 6 E). It suggests that t10c12-CLA-induced BAT mass increased in pai mice.

RNA sequencing was performed to provide a snapshot of the transcriptional dynamics between the wt and pai/pai BAT. The heatmap (Fig. 6 F) and a volcano plot (Fig. 6 G) revealed that the global transcriptional profile of pai/pai BAT differed from that of wt BAT. KEGG pathway enrichment analysis showed that the up-regulated pathways in pai/pai BAT included fatty acid biosynthesis, pyruvate metabolism, carbohydrate digestion and absorption, Insulin signalling pathway, steroid biosynthesis, biosynthesis of amino acids, citrate cycle, and propanoate metabolism (Fig. 6 H). In contrast, the down-regulated pathway included NK cell-mediated cytotoxicity, C-type lectin receptor signalling pathway, cell adhesion molecules, phagosome, tuberculosis, Fc epsilon RI signalling pathway, adherens junction, antigen processing and presentation, and Fc gamma R-mediated phagocytosis signalling pathways (Fig. 6 I).

Figure 6. Aspects of brown adipose tissues. Axial sections of the interscapular BAT show more prominent areas of grey signal intensity on magnetic resonance images (Brown arrows) in five pai/pai mice than that in five wt mice at the age of 11 weeks (A). The BAT mass increases in pai/pai mice (B, C). Imaging by hematoxylin-eosin staining show more small-sized and irregular lipid drops in the pai/wt or pai/pai adipocytes and relatively uniform and medium-sized lipid drops in the wt adipocytes (D, bar = 10 μm) and analysis of cross-sectional area per cell show that the volumes of brown adipocytes are increased in pai mice (E). RNA-Seq analysis shows transcriptional changes of differentially expressed genes in BAT from two wt and three pai/pai mice using heatmap (F) and the up- or down-regulated genes in pai/pai BAT using volcano plots (G). Maps of KEGG enrichment analysis show the up- (H) and down-regulated (I) metabolic pathways in pai/pai BAT. Western blot images (J) and relative intensities (K) analyses of some critical proteins in the BAT from wt and pai/pai mice. ACTIN, beta-actin; (p)AMPK, (phosphorylated) AMP-activated protein kinase; CPT1B, carnitine palmitoyltransferase 1b; FASN, fatty acid synthase; FGF21, fibroblast growth factor 21; PGC1α, pparγ coactivator 1α; pHSL, phosphorylated hormone-sensitive lipase; PPARγ, peroxisome proliferator-activated receptor-γ; UCP, uncoupling protein. Bars represent the mean ± SD. Different letters in superscript or * indicate $p < 0.05$.

Based on the prediction of transcriptome analysis and the conventional physiological

factors that can impact BAT's activity, 24 genes were chosen for real-time PCR analysis and

fourteen (56%) of them altered their transcriptional patterns in pai/pai brown adipocytes.

Among them, five up-regulated genes were Ampk, Fasn, Glut4, Ppar-r, and Pgc1α and nine

down-regulated genes were Srebp1a, Lpl, Hsl, Mgl, Perilipin1a, Cpt1a, Cpt1b, Ucp1, and Ucp2

(Table 3). In addition, western blot analysis revealed that the protein level of AMPK was not

altered. In contrast, the pAMPK level was increased significantly in pai/pai BAT compared

with the wt samples (Fig. 6 J-K), suggesting the AMPK activation to rewire metabolism to

decrease anabolic processes and increase catabolism. Coordinately, the elevation of CPT1B

and UCP2 proteins suggests the increased lipid beta-oxidation and BAT thermogenesis in

pai/pai mice. However, no difference in protein levels of UCP1, PGC1, FASN, PPAR-γ, pHSL,

and FGF21 in the samples of BAT from the pai/pai and wt mice (Fig. 6 J-K). The inconsistency

between some RNA and protein levels suggests the possibility of feedback regulation in pai

mice.

3.8. Hepatic features

Compared to wt mice, the hepatic analysis also showed varying degree differences

between pai/wt and pai/pai livers. Besides liver hypertrophy (p < 0.05; Fig. 7 A) and the amount

decrease of total FAs (p < 0.05; Supplementary Table 4) in pai/pai livers, the hepatic TC or

TGs levels had not changed (p > 0.05; Fig. 7 B-C), and no steatosis was observed by

histological staining of liver slices from pai/wt and pai/pai mice, while the estimation of cross-

sectional area per cell showed hepatocytes were swollen (Fig. 7 D-G), consistent with liver

hypertrophy in both pai genotypes. In addition, the mRNA levels of 58% (28/48) critical genes

were significantly changed in pai/pai livers, including 11 down-regulated and 17 up-regulated

Figure 7. Lipid and histological analysis of livers. Comparison of weight (A), total cholesterol (B), and triglycerides (C) levels in wt and pai livers. Imaging of hematoxylin-eosin staining shows the abnormal morphology and oedema of pai hepatocytes. Analysis of cross-sectional area per cell indicates that the cellular volumes gradually enlarge in pai/wt and pai/pai hepatocytes, respectively (D-E, Bar = 10 μm). Oil red staining and analysis of the cross-sectional area of lipids per field show no lipid accumulation in either pai liver (G~H). Bars represent the mean ± SD, and different letters indicate p < 0.05.

Table 4. Relative expression of mRNAs in the livers

Genes	wt (n)	pai/pai (n)
Fatty acid and TGs synthesis		
Fatty acid synthase (*Fasn*)	1.03 ± 0.26 (14)	1.72 ± 0.56*** (13)
Stearoyl-CoA desaturase 1 (*Scd1*)	1.06 ± 0.41 (7)	1.19 ± 0.50 (5)
Fatty acid desaturase 1 (*Fads1*)	1.01 ± 0.21 (3)	1.08 ± 0.18 (3)
Fatty acid desaturase 2 (*Fads2*)	1.04 ± 0.33 (3)	0.92 ± 0.26 (3)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	1.04 ± 0.35 (28)	0.65 ± 0.31*** (23)
Glycerol-3-phosphate acyltransferase 1 (*Gpat1*)	1.05 ± 0.35 (7)	1.04 ± 0.37 (8)
1-acylglycerol-3-phosphate O-acyltransferase 2 (*Agpat2*)	1.02 ± 0.18 (15)	1.11 ± 0.25 (14)
Diacylglycerol acyltransferase-1 (*Dgat1*)	1.04 ± 0.28 (17)	1.37 ± 0.39* (15)
Diacylglycerol acyltransferase-2 (*Dgat2*)	1.03 ± 0.26 (9)	1.05 ± 0.32 (9)
CD36	1.07 ± 0.38 (8)	0.58 ± 0.32* (8)
Lipolysis and fatty acid oxidation		
Hepatic triglyceride lipase (*Htgl*)	1.06 ± 0.37 (9)	1.25 ± 0.28 (9)
Adipose triglyceride lipase (*Atgl*)	1.04 ± 0.33 (9)	1.03 ± 0.36 (9)
Hormone-sensitive lipase (*Hsl*)	1.07 ± 0.38 (9)	1.08 ± 0.26 (9)
Comparative gene identification 58 (*Cgi58*)	1.03 ± 0.27 (9)	1.40 ± 0.38* (9)
Lipoprotein lipase (*Lpl*)	0.99 ± 0.16 (11)	1.37 ± 0.15*** (6)
Carnitine palmitoyltransferase Ia (*Cpt1a*)	1.05 ± 0.37 (9)	1.01 ± 0.35 (9)
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.01 ± 0.18 (14)	1.11 ± 0.49 (14)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.06 ± 0.38 (15)	0.95 ± 0.36 (13)
Uncoupling protein 1 (*Ucp1*)	1.00 ± 0.31[a] (20)	0.59 ± 0.36** (13)
Uncoupling protein 2 (*Ucp2*)	1.04 ± 0.28 (16)	0.82 ± 0.28* (14)
Sterol pathway		
Liver X receptor α (*Lxrα*)	1.01 ± 0.16 (9)	1.15 ± 0.40 (9)
Liver X receptor β (*Lxrβ*)	1.02 ± 0.23 (9)	1.22 ± 0.47 (9)
Sterol Regulatory Element Binding Protein-1a (*Srebp1a*)	1.07 ± 0.39 (7)	1.54 ± 0.28* (7)
Sterol Regulatory Element Binding Protein-1c (*Srebp1c*)	1.12 ± 0.69 (4)	1.29 ± 0.44 (5)
Sterol Regulatory Element Binding Protein-2 (*Srebp2*)	1.00 ± 0.30 (16)	1.40 ± 0.71* (18)
HMG-CoA reductase (*Hmgcr*)	1.03 ± 0.27 (6)	0.65 ± 0.28* (7)
Insulin and IGF-1 signaling		
Insulin receptor (*Insr*)	0.99 ± 0.24 (16)	1.40 ± 0.34*** (17)
Insulin receptor substrate-1 (*Irs1*)	1.03 ± 0.28 (29)	1.19 ± 0.68 (25)
Insulin receptor substrate-2 (*Irs2*)	1.02 ± 0.34 (15)	1.55 ± 0.52** (16)

Table 4. Relative expression of mRNAs in the livers

Genes	wt (n)	pai/pai (n)
Insulin-like growth factor-1 (*Igf1*)	1.01 ± 0.23 (21)	1.40 ± 0.55** (22)
IGF binding protein 1 (*Igfbp1*)	1.01 ± 0.23 (15)	0.53 ± 0.26*** (18)
Forkhead box protein a2 (*Foxa2*)	1.03 ± 0.31 (27)	1.75 ± 0.77*** (30)
Igf1 receptor (*Igf1r*)	1.07 ± 0.37 (5)	1.63 ± 0.28* (4)
NADPH-producing		
Malic enzyme	1.02 ± 0.23 (8)	1.89 ± 0.93* (7)
Glucose-6-phosphate dehydrogenase (*G6pd*)	0.98 ± 0.19 (14)	2.34 ± 0.63*** (17)
6-Phosphogluconate dehydrogenase (*Pgd*)	1.24 ± 0.93 (8)	1.29 ± 0.80 (9)
Glucose metabolism and gluconeogenesis		
Glucose transporter type 4 (*Glut4*)	0.99 ± 0.23[a] (20)	0.59 ± 0.23*** (15)
Glucokinase	1.02 ± 0.23 (12)	0.66 ± 0.28** (13)
Carbohydrate response element binding protein (*Chrebp*)	1.00 ± 0.28 (13)	1.98 ± 0.60*** (11)
Fibroblast growth factor 21 (*Fgf21*)	1.32 ± 1.07 (9)	1.34 ± 0.91 (8)
Glucose-6-phosphatase (*G6p*)	1.02 ± 0.19 (14)	2.34 ± 0.63*** (25)
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.06 ± 0.35 (13)	0.69 ± 0.31** (14)
Others		
AMP-activated protein kinase (*Ampk*)	1.06 ± 0.37 (9)	1.07 ± 0.35 (9)
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.03 ± 0.31 (3)	1.26 ± 0.36 (3)
F4/80	1.02 ± 0.20 (14)	1.45 ± 0.53* (13)
CD68	1.05 ± 0.34 (10)	0.76 ± 0.22* (10)
CD11c	1.05 ± 0.31 (19)	1.93 ± 0.89** (17)
Forkhead box protein c2 (*Foxc2*)	1.01 ± 0.17 (8)	1.59 ± 1.02 (10)
Pai	1.17 ± 0.69 (7)	10830 ± 8060* (7)

Numbers in brackets of the parameters indicate the size of tested samples. Each value represents the mean ± SD. * indicates p < 0.05; ** indicates p < 0.01; *** indicates p < 0.001, respectively.

genes (p < 0.05; Table 4), suggesting the t10c12-CLA might influence the liver metabolism in

mice.

Although the hepatic TC and TGs levels were normal in pai/pai mice, their hepatic

Srebp1a and Srebp2 were up-regulated, and the HMG-CoA reductase (Hmgcr) was down-

regulated. Some genes involved in FAs/TGs synthesis had also altered their transcription

levels. For instance, the RNA levels of Fasn, Dgat1 which convert diacylglycerol into TGs,

and Lpl which support FAs uptake, were increased; and the RNA levels of Ppar-γ and CD36,

which transport long-chain FAs into the cells, were decreased. Simultaneously, a series of

crucial enzymes involved in lipolysis and beta-oxidation kept standard transcription except for

the up-regulated Cgi58. In addition, the Ucp 1 and 2 engaged in thermogenesis were down-

regulated; the malic enzyme and G6pd, which both produce NADPH, were significantly up-

regulated; the inflammatory factors F4/80 and CD11c were up-regulated, but CD68 was down-

regulated (Table 4).

For genes involved in insulin/IGF1 signalling, except for reduced Igfbp1 and commonly

expressed Irs1, the other five essential genes, including Insr, Irs2, Igf1, Igf1 receptor, and

pioneering transcription factor Foxa2 were elevated together. For genes involved in glucose

metabolism and gluconeogenesis, the RNA levels of Glut4, glucokinase, and Pepck were

reduced, whereas the RNA levels of G6p and Chrebp were elevated. These transcriptional

changes suggest that t10c12-CLA easily influences hepatic glucose metabolism and insulin

sensitivity.

3.9. Hypothalamic gene analysis

To determine whether gene expression has been affected by t10c12-CLA in the

hypothalamus, which contains highly conserved neural circuitry controlling energy

Table 5. Relative expression of mRNAs in the hypothalamus

Genes	wt (n)	pai/wt (n)	pai/pai (n)
Insulin receptor (Insr)	1.01 ± 0.14^a (8)	1.49 ± 0.64^{ab} (4)	1.40 ± 0.34^b (8)
Leptin receptor (Lepr)	1.02 ± 0.21^a (9)	0.97 ± 0.39^a (7)	1.60 ± 0.60^b (8)
Ghrelin receptor (Ghsr)	1.03 ± 0.25^a (8)	1.21 ± 0.43^a (7)	1.80 ± 0.41^b (6)
AMP-activated protein kinase (Ampk)	1.02 ± 0.20^a (8)	1.41 ± 0.47^b (9)	1.61 ± 0.51^b (8)
Signal transducer and activator of transcription 3 (Stat3)	1.03 ± 0.27 (6)	1.11 ± 0.16 (7)	0.82 ± 0.45 (6)
Protein kinase B (Akt/PKB)	1.01 ± 0.14^a (9)	1.84 ± 0.70^{ab} (4)	1.63 ± 0.79^b (10)
DEP domain containing MTOR-interacting protein (Deptor)	1.02 ± 0.24 (9)	1.49 ± 0.46 (4)	0.96 ± 0.28 (10)
Agouti-related peptide (Agrp)	1.03 ± 0.24^a (14)	1.00 ± 0.33^a (13)	1.88 ± 0.81^b (11)
Neuropeptide Y (Npy)	1.02 ± 0.21^a (14)	1.31 ± 0.35^b (15)	1.74 ± 0.77^b (11)
NPY postsynaptic receptor Y1 (Npy1r)	1.02 ± 0.22^a (13)	1.44 ± 0.40^b (8)	1.05 ± 0.35^a (15)
Proopiomelanocortin (Pomc)	1.02 ± 0.21^a (10)	1.31 ± 0.46^b (14)	1.20 ± 0.31^{ab} (8)
Cocaine- and amphetamine-regulated transcript (Cart)	1.00 ± 0.10^a (7)	1.01 ± 0.07^a (4)	0.80 ± 0.18^b (8)
Melanocortic 4 receptor (Mc4r)	1.02 ± 0.24 (9)	1.16 ± 0.18 (4)	0.91 ± 0.20 (11)
glucose-regulated protein 78 (Grp78)	1.00 ± 0.03^a (9)	1.31 ± 0.27^b (7)	1.39 ± 0.31^b (8)
Orexins	1.12 ± 0.40 (6)	0.96 ± 0.43 (7)	0.69 ± 0.63 (5)
Gonadtrophin releasing hormone (Gnrh)	1.03 ± 0.23^a (12)	1.33 ± 0.24^b (11)	1.44 ± 0.45^b (13)
Arginine vasopressin (Avp)	1.02 ± 0.22^a (8)	1.53 ± 0.34^b (8)	2.15 ± 1.21^b (8)
Corticotropin releasing hormone (Crh)	1.02 ± 0.22 (8)	1.05 ± 0.29 (10)	1.10 ± 0.37 (9)
Oxytocin	1.00 ± 0.10^a (5)	1.47 ± 0.36^b (7)	1.49 ± 0.40^b (6)
Nuclear receptor subfamily 3 group C member 1 (Nr3c1)	1.03 ± 0.19^a (7)	1.01 ± 0.26^a (7)	1.93 ± 0.56^b (6)
H19	1.02 ± 0.23 (9)	0.84 ± 0.23 (7)	0.82 ± 0.27 (7)
Ribonuclease type III (Dicer)	1.01 ± 0.15 (8)	0.86 ± 0.20 (7)	0.88 ± 0.21 (6)
DNA (cytosine-5)-methyltransferase 1 (Dnmt1)	1.00 ± 0.09^a (6)		1.37 ± 0.32^b (6)
Pai	1.05 ± 0.34^a (8)	258 ± 137^b (4)	356 ± 203^b (6)

Numbers in brackets of the parameters indicate the size of tested samples. Each value represents the mean ± SD. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

metabolism, fluid and electrolyte balance, thermoregulation, responses to stressors, wake-sleep

cycles, and reproduction. The current study randomly chose 23 critical hypothalamic genes and

measured the transcriptional levels (Table 5).

Compared to wt samples, in the pai/pai hypothalamus, the RNA level changes were

observed in 14 (61%) genes. Only Cart, a marker of appetite-suppressive POMC neuron cells,

was down-regulated among them. The remaining 13 genes were up-regulated, including

appetite-relative genes such as the Insulin receptor, ghrelin receptor, agouti-related peptide,

neuropeptide Y, leptin receptor, and the intracellularly signal molecule Akt/PKB, suggesting

the sensitivity in regulating energy balance. In addition, the up-regulated gonadotrophin-

releasing hormone, oxytocin, arginine vasopressin, DNA methyltransferase-1, and Nr3c1,

suggesting that t10c12-CLA might affect the various neural circuitry in the hypothalamus.

Furthermore, the RNA levels of Ampk and molecular chaperone Grp78 which releases the

endoplasmic reticulum pressure and transmits thermogenesis signals to the BAT tissues were

increased in the pai/pai hypothalamus. However, western blot analysis revealed that the protein

levels of AMPK, pAMPK and GRP78 were not altered in the pai/pai hypothalamus compared

with the wt samples (Fig. 8 A-B).

In the pai/wt hypothalamus, the RNA levels of eight (36%) genes, including Ampk, Npy,

Npy postsynaptic receptor Y1, Pomc, Grp78, GnRH, oxytocin, and arginine vasopressin, were

increased, and no tested genes were down-regulated in pai/wt hypothalamus. Between pai/wt

and pai/pai samples, the RNA levels of six (27%) genes, including Lepr, Ghsr, Agrp, Npy

postsynaptic receptor Y1, Cart, and Nr3c1 exhibited differences ($p < 0.05$), suggesting the

genotype-specific transcription in the hypothalamus.

Figure 8. Western blot images (A) and relative intensities (B) analyses of some critical proteins in the hypothalamus from wt and pai/pai mice. ACTIN, beta-actin; GRP78, glucose-regulated protein 78; (p)AMPK, (phosphorylated) AMP-activated protein kinase. Bars represent the mean ± SD.

4. Discussion

Unlike the temporary fat loss in the body [24] or milk fat depression which the fat de novo synthesis would reverse after the termination of the t10c12-CLA supplementation in mice [8], in pai mice, the chronic t10c12-CLA resulted in an irreversible fat reduction. It may cause an adaptive metabolic response throughout the entire developmental stage. Metabolic compensation or adaptation to t10c12-CLA is intrinsically complicated owing to its diverse metabolic functions in multiple target organs. Thus, the interpretation of these data could be more explicit in pai mice.

Although the fat reduction in milk did not impact the body weight of piglets from the energy supply perspective [25], previous studies in mice indicated that the dietary t10c12-CLA could reduce milk fat and retard pup growth [8, 26]. However, we found that the wt pups fostered by the pai mother were reversely overweight at three weeks, suggesting that the natural t10c12-CLA from the mothers might play an active role during offspring development and more work needs to be done in future.

When a mouse was fed a low-fat diet containing t10c12-producing lactic acid bacteria or 0.4~1% t10c12-CLA, its fat loss happened quickly, as early as seven days after starting the t10c12-CLA diet [13, 24, 27] and the body fat could reduce from approximately 2.8 g to less than 0.4 g in C57BL/6J males [28]. Simultaneously, the contents of t10c12-CLA were 0.1% in the livers of Balb/c males [7], approximately 0.2% in various tissues of ICR females [9], 0.45% in the milk of C57BL/6J females [8], 0.013 μmol/g in the brain or 0.089 μmol/g t10c12-CLA in eye lipids of C57BL/6N females [12]. Similarly, in the current study, the comparable values of t10c12-CLA were also detected in transgenic hearts (8.85 μg/g, 0.09%), livers (16.65 μg/g,

0.19%), kidneys (11.33 μg/g, 0.09%), BAT (28.8 μg/g, 0.23%), or WAT (4.89 μg/g, 0.13%;

Supplemental Table 4) in pai mice.

Studies on whether the dietary t10c12-CLA plays a role in food intake or appetite showed

contradictory results in mice [4, 6, 27, 29]. In pai mice, we had not observed any changes in

food intake within metabolic cages or crucial circulating leptin, insulin, glucose, and ghrelin.

The results suggest that the anti-obesity property of t10c12-CLA is independent of food intake

or appetite. On the other hand, the hypothalamic RNA increases in receptors for insulin, leptin,

and ghrelin, and vital appetite-relative genes Npy, Argp, and an RNA decrease in appetite-

suppressive Cart imply a tendency to promote food intake, which the feedback regulation

mechanism might cause after more heat release, for the hormonal receptors, neuropeptides and

energy sensor AMPK is in a complex regulation and interplay of energy homeostasis in the

hypothalamic cells.

Studies of oral t10c12-CLA administration suggested that the fat loss mainly occurred in

fat depots through modulating adipocyte metabolism, such as increased fatty acid oxidation

and browning WAT [2, 13, 30], or increasing the number of beige adipocytes in mice [28]. The

pai/pai mice here reduced the WAT mass by approximately 20% at 11 weeks. However, RNA

analysis indicated reduced pai/pai WAT browning and gluconeogenesis. Interestingly, the

beta-oxidation of very long-chain FAs in peroxisomes was increased. That was, the pai/pai

mice had adapted to long-term t10c12-induced fat reduction and could decrease the browning

of WAT and mobilise the energy from the very long-chain FAs to maintain energy

homeostasis.

Studies on whether the dietary t10c12-CLA modulates hepatic lipid metabolism and even

induces fatty liver also showed contradictory results in mice [5, 13, 24, 31, 32]. It was supposed

that the t10c12-induced hepatic steatosis might be due to the uptake and accumulation of lipids

mobilised from the adipose tissue [5]. In the current study, although the essential genes Fasn,

Dgat1, and Lpl involved in TGs uptake and synthesis were up-regulated in hepatocytes,

analysis of hepatic TGs and TC levels and slices indicated no steatosis in pai livers.

Furthermore, the blood TGs concentration was also lowered in pai mice, similar to the reduced

plasma TGs in mice treated with a high dose of t10c12-CLA (0.6% w/w) [33]. A study in

lactating mice suggested that dietary t10c12-CLA might not induce fat accumulation in the

liver when plenty of energy was consumed for lactation [8]. In pai mice, more energy

expenditure via increased thermogenesis also hinted that no surplus energy could be

accumulated in the liver. Our results suggest that the t10c12-CLA has yet to break local lipid

homeostasis in the livers thoroughly.

A study in female 129Sv/J retired breeders indicated that dietary CLA mixture could

induce body fat reduction via increasing energy expenditure [4]. Unfortunately, this work has

yet to clarify whether the energy expenditure is via exercise strengthening or fat burning. Here,

increased heat release accompanied by reduced activities in pai/pai mice suggests that body fat

reduction results from increased thermogenesis. Additionally, the levels of circulating PGE2,

adrenaline, corticosterone, TNF-α, Il-6, and CRP in pai mice also suggest that the increased

heat production is not due to the response to bacterial PAI enzyme-induced stress and/or

inflammation in pai mice.

BAT thermogenesis can account for up to 50% of the body's energetic maintenance

demand. An increase [9] or decrease in BAT mass [33] were respectively observed in t10c12-

CLA treated mice, but in pai/pai mice, the increased BAT mass might correlate with more heat

release; additionally, the over-expression of pAMPK that phosphorylates key targets to rewire

a variety of metabolic processes and UCP2 proteins in the pai/pai brown adipocytes also

suggest increased thermogenesis via BAT activation. Based on the phenotypic characteristics

of pai mice, the authors suspected that the excess FGF21 in the blood and the central regulatory

mechanism might be involved in the t10c12-CLA-induced BAT thermogenesis.

FGF21 functions as a master sensitiser of specific hormonal signalling involved in

enhancing insulin sensitivity and lowering serum glucose through direct actions on adipose

tissues, possibly through the FGF21/adiponectin/ceramide axis [34], increasing energy

expenditure through immediate efforts on the central nervous system to stimulate BAT

thermogenesis, decreasing hepatic oxidative stress [35], lowering plasma TGs by accelerating

lipoprotein catabolism in WAT and BAT [36], and regulating macronutrient preference [23].

In the current study, excess FGF21 in the blood and local FGF21 expression in the BAT tissues

suggest that the circulating FGF21 plays an active role in BAT thermogenesis. Additionally,

the observable characteristics in pai/wt mice, such as an increase in FAs content in the kidney,

body weight increase during adulthood, blood glucose sensitivity, serum TGs decrease, and

genotype-specific gene expression in various tissues would be associated with the excess

FGF21. We have yet to pay more attention to unique features in pai/wt mice here, and further

work needs to be done in the future.

AMPK in the ventromedial nucleus of the hypothalamus can decrease ceramide-induced

endoplasmic reticulum (ER) stress which stimulates the thermogenic program in BAT through

the sympathetic nervous system [37]. In addition, GRP78 over-expression in the ventromedial

nucleus can ameliorate hypothalamic ER stress, leading to weight loss, reduced hepatic

steatosis, improved insulin and leptin sensitivities, increased BAT thermogenesis and

stimulation WAT browning [38, 39]. In the pai hypothalamus, the up-regulated transcriptional

levels of Ampk and Grp78 suggest that the hypothalamic neurons might promote heat

production via ameliorating ER stress in pai/pai mice. Unfortunately, we had not observed

evident changes in the hypothalamic AMPK, pAMPK and GRP78 proteins.

Focused on increased thermogenesis, more work needs to clarify if the t10c12-CLA

might affect ceramide in various ways: 1) the sphingomyelinase pathway to break down

sphingomyelin in the plasma membrane to release ceramide; 2) de novo synthesis of ceramide

from the condensation of palmitic acid (16:0) and serine to form 3-keto-dihydrosphingosine in

the ER; or 3) the salvage pathway to break down sphingolipids to sphingosine in the lysosomes

and to form ceramide by re-acylation in the ventromedial nucleus.

In addition, the expression modification of other hypothalamic genes, such as GCs

receptor Nr3c1, required negative regulation of the hypothalamic–pituitary–adrenal axis at a

basal condition and under stress [40]; vasopressin involved in fluid and electrolyte balance;

GnRH and oxytocin involved in reproduction; even imprinted H19 and Dnmt1 involved in the

epigenetic modification, were also observed in the pai hypothalamus. These aberrant

expressions of hypothalamic genes suggest that the t10c12-CLA might play active roles in the

myriad neural circuits in the hypothalamus and a more complex mechanism of action of

t10c12-CLA needs to be investigated further.

Acknowledgement

We thank Dr Li-Li Song and graduate students Yun-Cheng Zhuang, Yu-Huang Chen, Chao Wang, Qi Wang, Mei Liu, and Jia-Qi Lu in our lab for their generous assistance in partial tests and Ms Yu-Yang Wang in our university for technique help in gas chromatography. We also thank Professor Hong-Jun Wang at the Medical University of South Carolina, USA, for the helpful discussion during the writing process and Ms Wen-Jing Gou's language help in preparing this manuscript.

Funding sources

This work was supported by the National Key Research and Development Program of China (2018YFC1003800) and a Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD). The Jiangsu Funding Program for Excellent Postdoctoral Talent supported Rao Y.

CRediT Authorship Contributions

Yu Rao: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Software; Roles/Writing - original draft. Shi-Li Li: Conceptualization; Formal analysis. Mei-Juan Li, Bao-Zhu Wang, Yang-Yang Wang, Lu-Wen Liang: Data curation; Investigation; Methodology. Shuai Yu, Zong-Ping Liu: Data curation; Software; Validation; Visualization. Sheng Cui: Conceptualization; Funding acquisition; Project administration. Ke-Mian Gou: Conceptualization; Formal analysis; Funding acquisition; Project administration; Supervision; Writing - review & editing.

All authors proofread, made comments, and approved the report.

Competing Interest Statement

563  The authors declare no competing interests.

564  **Data access**

565  The data supporting the findings of this study are available from the corresponding author

566 upon reasonable request.

567  **Supplementary data**

568  This manuscript contains supplementary data.

References

[1]   Ritzenthaler KL, McGuire MK, McGuire MA, Shultz TD, Koepp AE, Luedecke LO, et al. Consumption of conjugated linoleic acid (CLA) from CLA-enriched cheese does not alter milk fat or immunity in lactating women. J Nutr 2005;135:422-30. http://doi.org/10.1093/jn/135.3.422.

[2]   Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. Effect of conjugated linoleic acid on body composition in mice. Lipids 1997;32:853-8. http://doi.org/10.1007/s11745-997-0109-x.

[3]   Baumgard LH, Corl BA, Dwyer DA, Saebo A, Bauman DE. Identification of the conjugated linoleic acid isomer that inhibits milk fat synthesis. Am J Physiol Regul Integr Comp Physiol 2000;278:R179-84. http://doi.org/10.1152/ajpregu.2000.278.1.R179.

[4]   Park Y, Park Y. Conjugated nonadecadienoic acid is more potent than conjugated linoleic acid on body fat reduction. J Nutr Biochem 2010;21:764-73. http://doi.org/10.1016/j.jnutbio.2009.05.003.

[5]   Vyas D, Kadegowda AK, Erdman RA. Dietary conjugated linoleic Acid and hepatic steatosis: species-specific effects on liver and adipose lipid metabolism and gene expression. J Nutr Metab 2012;2012:932928. http://doi.org/10.1155/2012/932928.

[6]   Kim JH, Kim Y, Kim YJ, Park Y. Conjugated Linoleic Acid: Potential Health Benefits as a Functional Food Ingredient. Annu Rev Food Sci Technol 2016;7:221-44. http://doi.org/10.1146/annurev-food-041715-033028.

[7]   Rosberg-Cody E, Stanton C, O'Mahony L, Wall R, Shanahan F, Quigley EM, et al. Recombinant lactobacilli expressing linoleic acid isomerase can modulate the fatty acid composition of host adipose tissue in mice. Microbiol-Sgm 2011;157:609-15. http://doi.org/10.1099/mic.0.043406-0.

[8]   Harvatine KJ, Robblee MM, Thorn SR, Boisclair YR, Bauman DE. Trans-10, cis-12 CLA dose-dependently inhibits milk fat synthesis without disruption of lactation in C57BL/6J mice. J Nutr 2014;144:1928-34. http://doi.org/10.3945/jn.114.198911.

[9]   Li SL, Ma SY, Xu BR, Fan ZY, Li MJ, Cao WG, et al. Effects of trans-10, cis-12-conjugated linoleic acid on mice are influenced by the dietary fat content and the degree of murine obesity. Eur J Lipid Sci Tech 2015;117:1908-18. http://doi.org/10.1002/ejlt.201400568.

[10]   Adkins Y, Belda BJ, Pedersen TL, Fedor DM, Mackey BE, Newman JW, et al. Dietary Docosahexaenoic Acid and trans-10, cis-12-Conjugated Linoleic Acid Differentially Alter Oxylipin Profiles in Mouse Periuterine Adipose Tissue. Lipids 2017;52:399-413. http://doi.org/10.1007/s11745-017-4252-3.

[11]   So MH, Tse IM, Li ET. Dietary fat concentration influences the effects of trans-10, cis-12 conjugated linoleic acid on temporal patterns of energy intake and hypothalamic expression

of appetite-controlling genes in mice. J Nutr 2009;139:145-51. http://doi.org/10.3945/jn.108.093849.

[12] Vemuri M, Adkins Y, Mackey BE, Kelley DS. Docosahexaenoic Acid and Eicosapentaenoic Acid Did not Alter trans-10,cis-12 Conjugated Linoleic Acid Incorporation into Mice Brain and Eye Lipids. Lipids 2017;52:763-9. http://doi.org/10.1007/s11745-017-4282-x.

[13] Shen W, Baldwin J, Collins B, Hixson L, Lee KT, Herberg T, et al. Low level of trans-10, cis-12 conjugated linoleic acid decreases adiposity and increases browning independent of inflammatory signaling in overweight Sv129 mice. J Nutr Biochem 2015;26:616-25. http://doi.org/10.1016/j.jnutbio.2014.12.016.

[14] Marques TM, Wall R, O'Sullivan O, Fitzgerald GF, Shanahan F, Quigley EM, et al. Dietary trans-10, cis-12-conjugated linoleic acid alters fatty acid metabolism and microbiota composition in mice. Br J Nutr 2015;113:728-38. http://doi.org/10.1017/S0007114514004206.

[15] Oteng AB, Kersten S. Mechanisms of action of trans fatty acids. Adv Nutr 2020;11:697-708. http://doi.org/10.1093/advances/nmz125.

[16] Kohno-Murase J, Iwabuchi M, Endo-Kasahara S, Sugita K, Ebinuma H, Imamura J. Production of trans-10, cis-12 conjugated linoleic acid in rice. Transgenic Res 2006;15:95-100. http://doi.org/10.1007/s11248-005-3736-8.

[17] Wang C, Wang YY, Rao Y, Li MJ, Liu ZP, Li SL, et al. Heterologous expression of Propionibacterium acnes isomerase in mouse (Mus musculus) cells and production of t10c12-conjugated linoleic acid. Chinese Journal of Agricultural Biotechnology 2021;29:2304-11. http://doi.org/10.3969/j.issn.1674-7968.2021.12.004.

[18] Li SL. Analysis of the effect of trans 10, cis 12-conjugated linoleic acid on mice growth and metabolism. Ph.D. Dissertation. Beijing: China Agricultural University; 2015.

[19] Li M, Li S, Rao Y, Cui S, Gou K. Loss of smooth muscle myosin heavy chain results in the bladder and stomach developing lesion during foetal development in mice. J Genet 2018;97:469-75.

[20] Chu VT, Weber T, Graf R, Sommermann T, Petsch K, Sack U, et al. Efficient generation of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. BMC Biotechnol 2016;16:4. http://doi.org/10.1186/s12896-016-0234-4.

[21] Jenkins TC. Technical note: common analytical errors yielding inaccurate results during analysis of fatty acids in feed and digesta samples. J Dairy Sci 2010;93:1170-4. http://doi.org/10.3168/jds.2009-2509.

[22] Chen HC, Farese RV, Jr. Determination of adipocyte size by computer image analysis. J Lipid Res 2002;43:986-9.

[23] Flippo KH, Potthoff MJ. Metabolic Messengers: FGF21. Nat Metab 2021;3:309-17. http://doi.org/10.1038/s42255-021-00354-2.

[24] Cordoba-Chacon J, Sugasini D, Yalagala PCR, Tummala A, White ZC, Nagao T, et al. Tissue-dependent effects of cis-9,trans-11- and trans-10,cis-12-CLA isomers on glucose and lipid metabolism in adult male mice. J Nutr Biochem 2019;67:90-100. http://doi.org/10.1016/j.jnutbio.2019.01.020.

[25] Sandri EC, Harvatine KJ, Oliveira DE. Trans-10, cis-12 conjugated linoleic acid reduces milk fat content and lipogenic gene expression in the mammary gland of sows without altering litter performance. Br J Nutr 2020;123:610-8. http://doi.org/10.1017/S0007114519003325.

[26] Robblee MM, Boisclair YR, Bauman DE, Harvatine KJ. Dietary Fat Does Not Overcome trans-10, cis-12 Conjugated Linoleic Acid Inhibition of Milk Fat Synthesis in Lactating mice. Lipids 2020;55:201-12. http://doi.org/10.1002/lipd.12228.

[27] Shelton VJ, Shelton AG, Azain MJ, Hargrave-Barnes KM. Incorporation of conjugated linoleic acid into brain lipids is not necessary for conjugated linoleic acid-induced reductions in feed intake or body fat in mice. Nutr Res 2012;32:827-36. http://doi.org/10.1016/j.nutres.2012.10.003.

[28] Yeganeh A, Zahradka P, Taylor CG. Trans-10,cis-12 conjugated linoleic acid (t10-c12 CLA) treatment and caloric restriction differentially affect adipocyte cell turnover in obese and lean mice. Journal of Nutritional Biochemistry 2017;49:123-32. http://doi.org/10.1016/j.jnutbio.2017.08.003.

[29] Park Y, Albright KJ, Storkson JM, Liu W, Pariza MW. Conjugated linoleic acid (CLA) prevents body fat accumulation and weight gain in an animal model. J Food Sci 2007;72:S612-7. http://doi.org/10.1111/j.1750-3841.2007.00477.x.

[30] den Hartigh LJ, Wang SR, Goodspeed L, Wietecha T, Houston B, Omer M, et al. Metabolically distinct weight loss by 10,12 CLA and caloric restriction highlight the importance of subcutaneous white adipose tissue for glucose homeostasis in mice. Plos One 2017;12. http://doi.org/ARTN e0172912

10.1371/journal.pone.0172912.

[31] Ashwell MS, Ceddia RP, House RL, Cassady JP, Eisen EJ, Eling TE, et al. Trans-10, cis-12-conjugated linoleic acid alters hepatic gene expression in a polygenic obese line of mice displaying hepatic lipidosis. J Nutr Biochem 2010;21:848-55. http://doi.org/10.1016/j.jnutbio.2009.06.013.

[32] Kostogrys RB, Franczyk-Zarow M, Maslak E, Gajda M, Mateuszuk L, Chlopicki S. Effects of margarine supplemented with t10c12 and C9T11 CLA on atherosclerosis and steatosis in apoE/LDLR -/- mice. J Nutr Health Aging 2012;16:482-90. http://doi.org/10.1007/s12603-011-0354-4.

[33] Shen W, Chuang CC, Martinez K, Reid T, Brown JM, Xi L, et al. Conjugated linoleic acid reduces adiposity and increases markers of browning and inflammation in white adipose tissue of mice. J Lipid Res 2013;54:909-22. http://doi.org/10.1194/jlr.M030924.

[34]   Holland WL, Adams AC, Brozinick JT, Bui HH, Miyauchi Y, Kusminski CM, et al. An FGF21-adiponectin-ceramide axis controls energy expenditure and insulin action in mice. Cell Metab 2013;17:790-7. http://doi.org/10.1016/j.cmet.2013.03.019.

[35]   Ye D, Wang Y, Li H, Jia W, Man K, Lo CM, et al. Fibroblast growth factor 21 protects against acetaminophen-induced hepatotoxicity by potentiating peroxisome proliferator-activated receptor coactivator protein-1alpha-mediated antioxidant capacity in mice. Hepatology 2014;60:977-89. http://doi.org/10.1002/hep.27060.

[36]   Schlein C, Talukdar S, Heine M, Fischer AW, Krott LM, Nilsson SK, et al. FGF21 Lowers Plasma Triglycerides by Accelerating Lipoprotein Catabolism in White and Brown Adipose Tissues. Cell Metab 2016;23:441-53. http://doi.org/10.1016/j.cmet.2016.01.006.

[37]   Martinez-Sanchez N, Seoane-Collazo P, Contreras C, Varela L, Villarroya J, Rial-Pensado E, et al. Hypothalamic AMPK-ER Stress-JNK1 Axis Mediates the Central Actions of Thyroid Hormones on Energy Balance. Cell Metab 2017;26:212-29 e12. http://doi.org/10.1016/j.cmet.2017.06.014.

[38]   Contreras C, Gonzalez-Garcia I, Seoane-Collazo P, Martinez-Sanchez N, Linares-Pose L, Rial-Pensado E, et al. Reduction of Hypothalamic Endoplasmic Reticulum Stress Activates Browning of White Fat and Ameliorates Obesity. Diabetes 2017;66:87-99. http://doi.org/10.2337/db15-1547.

[39]   Contreras C, Fondevila MF, Lopez M. Hypothalamic GRP78, a new target against obesity? Adipocyte 2018;7:63-6. http://doi.org/10.1080/21623945.2017.1405878.

[40]   Laryea G, Schutz G, Muglia LJ. Disrupting hypothalamic glucocorticoid receptors causes HPA axis hyperactivity and excess adiposity. Mol Endocrinol 2013;27:1655-65. http://doi.org/10.1210/me.2013-1187.