Alpha-synuclein pre-formed fibrils injected into prefrontal cortex primarily spread to cortical and subcortical structures and lead to isolated behavioral symptoms

Matthew A. Weber, PhD[a], Gemma Kerr[a], Ramasamy Thangavel, PhD[a], Mackenzie M. Conlon[b], Hisham A. Abdelmotilib, MD, PhD[a], Oday Halhouli, MD[a], Qiang Zhang, MD[a], Joel C. Geerling MD, PhD[a], Nandakumar S. Narayanan, MD, PhD[a], Georgina M. Aldridge, MD, PhD[#,a]

Affiliations:

[a] Department of Neurology, Carver College of Medicine, University of Iowa, Iowa City.

[b] Medical Scientist Training Program, Carver College of Medicine, University of Iowa, Iowa City.

Running Title: Cortical/Subcortical α-Syn Related Behavior

Funding: This work was supported by NIH R01s MH116043, NS120987 to NSN and NIH K08 NS109287 and the Iowa Neuroscience Institute to GMA. This work was also supported by NIH P20NS123151 to NSN and GMA.

Conflict of Interest:
There are no conflicts of interest.

Abstract: 172
Main Text: 4653
Figures: 5
Supplemental Figures: 2
Tables: 0

\# Corresponding Author
Georgina M. Aldridge
georgina-aldridge@uiowa.edu
Phone: 319-384-1635
169 Newton Road
Pappajohn Biomedical Discovery Building - 5334
University of Iowa, Iowa City, 52242

Abstract

Parkinson's disease dementia (PDD) and dementia with Lewy bodies (DLB) are characterized by

diffuse spread of alpha-synuclein (α-syn) throughout the brain. Patients with PDD and DLB have a

neuropsychological pattern of deficits that include executive dysfunction, such as abnormalities in

planning, timing, working memory, and behavioral flexibility. The prefrontal cortex (PFC) plays a major

role in normal executive function and often develops α-syn aggregates in DLB and PDD. To investigate

the consequences of α-syn pathology in the cortex, we injected human α-syn pre-formed fibrils into the

PFC of wildtype mice. We report that PFC PFFs: 1) induced α-syn aggregation in multiple cortical and

subcortical regions with sparse aggregation in midbrain and brainstem nuclei; 2) did not affect interval

timing or spatial learning acquisition but did mildly alter behavioral flexibility as measured by intraday

reversal learning; 3) increased open field exploration; and 4) did not affect susceptibility to an

inflammatory challenge. This model of cortical-dominant pathology aids in our understanding of how

local α-syn aggregation might impact some symptoms in PDD and DLB.

Key Words: Prefrontal cortex, alpha-Synuclein, Parkinson's disease, Behavioral Flexibility, Exploratory

behavior

INTRODUCTION

Patients with brainstem alpha-synuclein (α-syn) pathology have motor characteristics of Parkinson's disease (PD), including rigidity, bradykinesia, and tremor [1]. However, patients with PD dementia (PDD) and dementia with Lewy bodies (DLB), collectively termed Lewy Body dementias (LBD), have diffuse spread of α-syn throughout the brain, including cortical regions [2,3]. These patients often suffer from severe symptoms, including executive and working memory deficits, fluctuations in arousal and attention, hallucinations and psychosis, and increased risk for delirium [3–6]. It is not known whether the spread of α-syn to the cortex causes the symptoms of dementia, such as impaired cognition, or whether cortical α-syn simply reflects a more general process of spread and severity that does not directly affect cortical and cognitive function. To date, there are very few effective therapies approved to manage the debilitating cognitive symptoms of dementia [7,8]; thus, it is critical that we understand the etiological factors of the diverse symptoms seen in LBD.

To investigate the impact of cortical-predominant pathology, we induced α-syn aggregation using human pre-formed fibrils (PFFs) injected in the prefrontal cortex (PFC) of wild-type mice, which can induce aggregation of native α-syn that resembles the pathology described in PDD and DLB [9–12]. Prior literature has extensively studied both mouse and human α-syn injected in the rodent striatum, the predominant model currently used primarily due to its unique ability to induce cell loss in the substantia nigra [11,13]. A few studies have used varying injection locations, including duodenal, pedunculopontine nucleus, nucleus accumbens, olfactory bulb, and others [14–20]. These studies have offered mixed results regarding behavioral deficits and often restrict analyses to 6–7 months post-injection (mpi). Furthermore, only a few studies have investigated PFC PFF injections [21–23]. A prior report from our group found no behavioral effects of PFC PFFs [23]. These animals had a combined viral overexpression/PFF expression strategy, aged animals to 15 months (6 mpi), and examined interval timing, novel object recognition, and open field behavior [23]. In this study, we investigated the effect of PFC PFFs on cortical, subcortical,

midbrain, and brainstem α-syn pathology in mice aged up to 2 years (21 mpi) with a broader range of

behavioral assays and challenges.

We report four results: 1) PFFs injected in the PFC caused a consistent pattern of

moderate-to-severe pathological α-syn aggregation in cortical and subcortical regions with sparse

aggregation in midbrain and brainstem nuclei; 2) cognitive behaviors were mildly affected in mice

injected with PFFs in the PFC, with no change in interval timing or spatial working memory acquisition,

but a small, consistent alteration in behavioral flexibility as measured by intraday reversal learning; 3)

mice with cortical PFFs showed greater exploration of the center of an open field, without a change in

total distance traveled; 4) mice with cortical PFFs did not have enhanced susceptibility to a delirium-like

trigger. Together, these data suggest that aggregation of α-syn in cortical regions impacts specific

behaviors even in the absence of midbrain/brainstem inclusion. These results provide insight into cortical

synucleinopathy, which is relevant for LBD.

MATERIALS AND METHODS

Mice: All experimental procedures were performed in accordance with relevant guidelines by the

University of Iowa Institutional Animal Care and Use Committee (IACUC). Wild-type male C57BL/6

mice were received from Jackson Labs (Bar Harbor, ME) at approximately 8–10 weeks of age and

acclimated to the animal holding facility for 1–2 weeks. All mice were communally housed on a 12-hour

light/dark cycle with *ad lib* access to standard laboratory chow and water, except as noted.

Monomer and fibril preparation: Monomeric alpha-synuclein and human pre-formed fibrils

(PFFs) were a contribution from Dr. Andrew West and prepared according to previously used protocols

[23,24]. On the day of surgery, aliquoted monomers and PFFs were thawed and kept on ice immediately

prior to surgical procedures. Potentially contaminated surfaces were then thoroughly cleaned with 10%

bleach.

Surgical Procedures: Mice were anesthetized under 4.0% isoflurane and surgical levels of

isoflurane (1.5–3.0%) were sustained for the duration of the surgery (SomnoSuite, Kent Scientific,

Torrington, CT, USA). Analgesic (carprofen) and local anesthetic (bupivacaine) were administered prior

to the start of surgery. An incision was made along midline and bilateral burr holes drilled above the

prefrontal cortex (PFC), and a Hamilton syringe was lowered to the target coordinates (AP +1.8, ML

±0.5, DV -1.5). Mice were assigned to bilateral microinjections of either 2 µL monomers or 2 µL PFFs.

Monomeric controls or PFFs were infused over 20 minutes (0.1 µl/min; Legato 130 Syringe Pump, kd

Scientific, Holliston, MA, USA), with a 5-minute wait period before removing the needle. The incision

was then sutured, and mice were moved to a clean cage with *ad lib* access to food and water. Behavioral

testing began approximately 12 months later (Fig 1).

Interval Timing Switch Task: All mice were trained to perform an interval timing switch task

described in detail previously [25–28]. Interval timing behavior was assessed on two occasions: 1) 12

months post-injection (mpi); 2) 20 mpi. Briefly, food-restricted mice are trained in standard operant

chambers (MedAssociates, St. Albans, VT) equipped with two nosepoke response ports (left and right)

flanking a reward hopper and another nosepoke response port on the back wall. Either the left or right

nosepoke was designated for short trials and the contralateral nosepoke for long trials (counterbalanced

across mice and experimental groups). Trials are initiated by a response at the back nosepoke, which

generated identical cues for both the left and right nosepokes. In short trials, mice receive a reward for the

first response after 6 seconds. In long trials, mice receive a reward for the first response after 18 seconds.

Because cues are identical for both short and long trials, the optimal strategy is to start responding at the

short nosepoke and switch to the other nosepoke to receive a reward if enough time passed without

reward.

Rapid reversal Barnes maze: A modified Barnes maze was used to evaluate spatial learning

acquisition and behavioral flexibility using a modified Barnes maze adapted from established protocols

[29,30], with the following notable exceptions: 1) the spatial target was changed every day to a new

random location and every 4 trials to an opposite location in order to assess rapid acquisition of a spatial

target (first 4 trials) and intraday reversal learning (last 4 trials); 2) to minimize the use of olfactory cues,

every hole was equipped with its own 3D printed escape box "shuttle" such that no shuttle was ever

re-used during a session, and the maze itself was also rotated. A plastic grate made of identical material

prevented entry to incorrect shuttles; 3) following a correct response (entry), the shuttle was connected by

tunnel with the home cage as a potent reward, during a 3-minute inter-trial interval; 4) a 40-hole circular

Barnes maze apparatus was used in which only 10 holes were available. Total distance traveled was

captured with overhead cameras and analyzed using ANY-Maze 7.14d (Stoelting, Wood Dale, IL).

Open field: Open field activity was used to assess exploration and activity levels on two

occasions (12- and 16 mpi), using methods described previously [31,32]. On each occasion, mice were

habituated to behavior rooms for at least 30 minutes prior to testing. Mice were placed in the center of an

empty open field arena (40 cm × 40 cm × 30 cm) and allowed to explore uninterrupted for 10 minutes.

Mice were returned to their home cage immediately, and the open field arenas were thoroughly cleaned

with 70% ethanol and allowed to air dry between trials. Rescue disinfectant (0.5% hydrogen peroxide)

was used between cohorts. Total distance traveled and thigmotaxis (distance traveled outside a 25.5 ×

25.5 cm central square divided by total distance traveled) were analyzed using ANY-Maze 7.14d.

Home cage activity: Home cage activity was assessed in parallel with nest building and food

burying (described below) at 15 mpi. Overhead cameras captured morning and nighttime activity using an

infrared camera. ANY-Maze 7.14d software was then used to quantify periods of activity and rest over a

one-hour period during the morning (~6–7 am) and night (~2–3 am). Midday activity could not be

quantified due to unexpected interference with lighting through cage bars. Periods of immobility were

defined as periods 40-seconds or longer during which ANY-Maze did not detect movement. Percent

immobile was then defined as the number of seconds the mouse spent immobile compared with total

seconds tracked. Bout length was based on prior published data using EEG suggesting that periods of

immobility greater than 40-seconds are likely to be sleep bouts [33].

Nesting: Nest building was used to approximate general health and welfare, using slightly

modified methods described previously [34]. Nesting behavior for each mouse was assessed on two

occasions (15- and 16 mpi). At 15 mpi, three different materials (shredded paper, paper twist, and pressed

cotton square) were provided on separate days. At 16 mpi, only the paper nest building material was

provided. Photographs were taken of each nest and scored by blinded observers on a scale of 1–5

according to previous literature [34,35].

Food burying: Food burying was assessed in parallel with nest building at 16 mpi. Pre-weighed

food was provided in a glass jar equipped with a PLA 3D printed cone-shaped hopper to prevent mice

from nesting in the jar. The jar was weighed the following day to calculate food buried or eaten.

Lipopolysaccharide injection challenge: Systemic infection increases risk of developing

delirium and dementia in humans [36–38] and peripheral injections of lipopolysaccharide (LPS) are

widely used as a model of systemic infection and resultant behavioral changes in rodents [39–41]. At 16

mpi, 0.5 mg/kg LPS (Sigma #L2880, St. Louis, MO) diluted in 0.9% sterile saline was injected

intraperitoneally (IP) in five mice with PFC monomers and five mice with PFC PFFs. This dose of LPS

was chosen as previous studies demonstrated accelerated decline in rodent models of tauopathy and prion

disease [42,43]. The remaining three mice with PFC monomers and three mice with PFC PFFs were

injected with equivalent volumes of saline. Behavior in the open field, food burying assay, and nest

building was then assessed. At 21 mpi, a lower dose (0.25 mg/kg LPS) was injected IP to all mice to

examine acute effects of the challenge in the open field and rapid reversal Barnes maze 48 hours prior to

perfusion (Fig. S2).

Histology: Mice were anesthetized with ketamine (100 mg/kg IP) and xylazine (10 mg/kg IP) and

transcardially perfused with phosphate-buffered saline (PBS) and 4% paraformaldehyde (PFA). Brains

were fixed in 5% PFA overnight and immersed in 30% sucrose for approximately 48 hours. Brains were

then serially sliced into 40-μm coronal sections using a cryostat (Leica Biosystems, Deer Park, IL). Brain

sections were blocked for one hour in 2% normal goat serum (NGS) in PBST (0.3% Triton X-100 in 1x

PBS) and then incubated for approximately 20 hours at 4°C in either rabbit anti-α-synuclein (α-syn)

phosphorylated serine 129 (P-S129; Abcam #EP1536Y, Cambridge, United Kingdom) diluted to 1:1000

in blocking solution or rabbit anti-tyrosine hydroxylase (TH; Millipore #AB152, Burlington, MA) diluted

to 1:1000 in blocking solution. Sections were then washed in PBS three times and incubated overnight at

4°C in goat anti-rabbit IgG (H+L) Alexa Fluor 647 secondary antibody (Invitrogen #A11036, Waltham,

MA) diluted to 1:1000 in blocking. Sections were washed three more times. To stain for nissl, sections

were washed in PBST for 10 minutes and two additional 5-minute washes in PBS. Sections were

incubated in NeuroTrace™ 435/455 (Thermo Fisher Scientific Inc. #N21479, Waltham, MA) diluted

1:300 in PBS for 20 minutes at room temperature and then washed in PBST for 10 minutes and twice in

PBS for 5 minutes each. After one more wash at room temperature for 2 hours, sections were mounted

with ProLong Diamond Antifade Mountant (Invitrogen #P36961) on Superfrost microscope slides (Fisher

Scientific).

All sections were imaged using VS-ASW-S6 imaging software (Olympus, Center Valley, PA)

and an Olympus Slide Scanner (Olympus VS120) with a 20x objective. To characterize pathological

spread of α-syn, we semi-quantitively graded the degree of P-S129 inclusions in multiple regions of

interest (ROI; see Fig. 2 for complete list) referenced from Franklin & Paxinos, 2008. The α-syn

pathology was graded using the rubric images from previously published literature [45]. Briefly,

pathology was assigned on a scale of 0–4 separately for neurites and somas, with the following general

guidelines, in addition to published rubric, to assist and calibrate blinded grading; 0: no soma or neurite

P-S129 positive staining; 1: sparse/light soma or neurite staining (1–3 somas or neurites per 750 µm

square area within ROI); 2: mild burden (4 or more somas or neurites per 750 µm, with definitive areas

showing no P-S129 staining); 3: dense burden (>10 soma or neurites per 250 µm); 4: heavy burden (>20

somas or neurites per 250 µm). For every brain, one out of every six 40 µm sections were graded for

α-syn inclusion. Results from brain regions that spanned multiple sections were averaged across sections

and hemispheres, even if some sections contained no pathology.

Statistics: All data was analyzed using custom written MATLAB routines and Graphpad Prism.

All tests with ordinal outcomes were analyzed using appropriate non-parametric statistics. Pathology

scores (ordinal) were analyzed using Friedman Rank Sum test. Tyrosine hydroxylase values (quantitative

intensity) were analyzed using a repeated-measures ANOVA. Behavioral tasks were analyzed using

separate unpaired t-tests or repeated measures ANOVA. Separate nest building scores (ordinal) were

analyzed using separate Mann-Whitney U tests. Behavioral tasks at 16 mpi – 0.5 mg/kg LPS challenge –

were analyzed using two- or three-way ANOVA or Friedman Rank Sum test (nest building only)

followed by separate Mann-Whitney U tests corrected for multiple comparisons.

RESULTS

Pathology

Injections of PFFs in the PFC led to α-syn aggregation in the injected cortical area as well as several downstream regions. We found clear and dense pathological spread of α-syn in orbitofrontal, secondary motor M2, perirhinal, entorhinal, piriform, nucleus accumbens, amygdala, and thalamus (Fig. 2C-D). Subregions of the striatum were differentially affected by PFFs injected in the PFC (Fig. 2F). A Friedman test was conducted and there was a statistically significant difference in α-syn aggregation between subregions of the striatum ($X^2(3) = 22.2$, p < 0.0001). Post-hoc analyses using Dunn's multiple comparisons test revealed significant differences between dorsomedial and dorsolateral striatum ($p = 0.04$), dorsomedial and ventrolateral striatum ($p < 0.0001$), and ventromedial and ventrolateral ($p = 0.04$). Unlike other subcortical regions, subregions of the hippocampus were considerably less dense, and similar sparse-to-absent pathological α-syn aggregation was observed in regions of the midbrain and brainstem.

We also quantified tyrosine hydroxylase positive (TH+) fluorescence levels in the PFC, striatum, ventral tegmental area, and substantia nigra (Fig. 2G), similar to prior methods established in the lab [28]. As expected, there was a statistically significant difference in TH+ fluorescence levels between regions (Fig. 3E; two-way repeated-measures ANOVA; main effect of region ($F_{(1.105, 15.47)} = 122.9$; $p < 0.0001$), but we found no significant change in TH+ fluorescent levels in PFF-injected mice compared with controls in any of the four regions (main effect of treatment ($F_{(1, 14)} = 0.2078$; $p = 0.66$). Further, we did not find any differences in markers for astrocyte (GFAP) or microglia (IBA-1) activation (Fig. S2D).

Cognitive behaviors: Interval timing and behavioral flexibility

Interval timing requires mice to estimate an interval of several seconds and requires working memory for temporal rules as well as attention to the passage of time. Our prior work showed no deficits in interval timing using a fixed-interval timing task at 15 months of age (6 months post injection (mpi) of

PFFs) [23]. Here, we sought to investigate interval timing performance in mice at 12+ mpi, using an

interval timing switch task that enables us to track mean temporal estimates as well as the variability of

temporal estimates. We found that at 12 mpi, mice injected with PFFs showed no difference in mean

switch times (shifting early or late as an estimate of timing accuracy; Fig. 3A; unpaired t-test; $t(13) =$

0.66; $p = 0.52$) and no difference in switch time coefficient of variability (estimate of timing precision;

Fig. 3B; unpaired t-test; $t(13) = 1.05$; $p = 0.31$). One PFF injected mouse was excluded because no trials

initiated or rewards obtained, likely due to a lack of motivation to perform the task. We repeated interval

timing switch task training at 20 mpi. Two monomer-treated mice were sacrificed at 19 mpi due to

age-related health concerns per the University of Iowa Office of Animal Resources guidelines, so this

round of interval timing training included six monomer and eight PFF mice. Similar to interval timing at

12 mpi, we found no difference in mean switch times (Fig. 3C; unpaired t-test; $t(12) = 0.29$; $p = 0.78$) or

the coefficient of variability of switch times (Fig. 3D; unpaired t-test; $t(12) = 1.20$; $p = 0.25$). We found

no difference in the proportion of rewarded trials at either timepoint (Fig. S1A & B). Together, these

results suggest that PFC PFFs do not affect interval timing behavior even at 20 mpi.

After interval timing training, two mice treated with PFFs were sacrificed due to age-related

health concerns. Therefore, six monomer and six PFF mice remained. To test learning and behavioral

flexibility, all remaining mice were tested using a rapid reversal Barnes maze at 21 mpi. PFF mice

showed an altered pattern of behavior flexibility, with PFF mice learning quickly during the initial trials,

but then travelling a greater distance to reach the target compared with control mice during the reversal

phase (Fig. 3E; two-way repeated-measures ANOVA; main effect of trial ($F_{(2.145, 19.31)} = 6.71$; $p = 0.005$)

and trial $\times$ treatment interaction ($F_{(7, 63)} = 2.17$; $p = 0.049$); post-hoc analyses revealed no significant

differences when corrected for multiple-comparisons). A similar pattern of behavior during the reversal

phase, although not statistically different, was observed during a single day rapid reversal Barnes protocol

at 17 mpi (Fig. S1C).

Open field

We assessed exploration and activity levels in the open field. At 12 mpi, mice injected with PFFs

in the PFC did not differ from monomer controls in distance traveled (meters) over the 10-minute test

(Fig. 4A; unpaired t-tests; $t(13) = 0.35$; $p = 0.73$), but did display reduced thigmotaxis, showing more

exploration of the center of the arena (Fig. 4B; unpaired t-tests; $t(13) = 2.49$; $p = 0.03$). These results may

suggest greater exploration or reduced anxiety-like behavior.

Home cage activity and nest building (15 mpi)

To assess general health and welfare as well as naturalistic behaviors, we used assays to examine

home cage activity and nest building. At 15 mpi, mice injected with PFFs in the PFC did not differ on

amount of home cage activity during either the light or dark cycle (Fig. 4C; mixed-effects two-way

ANOVA; main effect of treatment $F_{(1, 14)} = 0.04$; $p = 0.84$), but there was a significant main effect of time

of day ($F_{(1, 12)} = 50.45$; $p < 0.0001$). At the same time-point, there were no differences in nest building

score across three different types of material (Fig. 4D; Mann-Whitney *U*; paper $p = 0.61$; twist $p = 0.42$;

squares $p = 0.18$). Altogether, these data suggest that PFC PFFs do not reduce naturalistic innate

behaviors or overall well-being.

Peripheral immune challenge (16 mpi)

We tested if mice previously injected with PFF were more susceptible to peripheral immune

activation by lipopolysaccharide (LPS) injections. At 16 mpi, we injected five mice with PFC PFFs and

five mice with PFC monomers with 0.5 mg/kg LPS. The remaining three mice in each group received

equivalent volumes of saline. We then tested behavior in the open field, food burying, and nest building

assays. Although LPS caused a significant decrease in distance travelled (Fig. 5A; two-way ANOVA; $F_{(1,}$

$_{12)} = 44.82$; $p = < 0.0001$) and food burying (Fig. 5C; three-way ANOVA; $F_{(1, 12)} = 25.64$; $p = 0.0003$), we

found no evidence of an enhanced response in PFF injected mice (Fig. 5A; distance travelled $F_{(1, 12)} =$

1.33; $p = 0.27$; Fig. 5B; two-way ANOVA; thigmotaxis $F_{(1, 11)} = 0.52$; $p = 0.49$; Fig. 5C; food burying $F_{(1,}$

$_{12)} = 1.32$; $p = 0.27$). Unlike 12 mpi, there was no significant effect of PFFs on thigmotaxis ($F_{(1, 11)} = 2.46$;

$p = 0.15$), possibly due to a lack of statistical power, as the pattern remained similar even in the presence

of LPS. Similarly, although LPS impaired nest building ($X^2(4) = 12.18$, $p = 0.02$), this response was not

enhanced in PFC PFF mice (Fig. 5D; 1-day post LPS injection; Mann-Whitney U; $p = 0.16$). These

results suggest that PFC PFF mice are not more susceptible to peripheral immune challenge. This is

further supported by a final lower dose challenge of 0.25 mg/kg LPS given at 21 mpi. PFC PFF mice

were not more susceptible to LPS challenge in the open field or rapid reversal Barnes maze (Fig. S2).

DISCUSSION

We tested the effects of prefrontal cortex (PFC) human pre-formed fibrils (PFFs) in aged mice on multiple behavioral assays and then evaluated spread of α-synuclein (α-syn) aggregation in cortical, subcortical, and brainstem regions. First, we found that PFC PFF injection triggered widespread cortical and subcortical α-syn aggregation, without significant spread to brainstem regions, allowing evaluation of the behavioral impact of α-syn limited to these regions. Second, and in line with our prior work, we found no reliable deficits in interval timing, but we did find mild alterations in reversal learning. Third, we also found changes in exploration without changes in distance travelled in the open field assay. Finally, we found no effects of inflammatory challenge. Taken together, these data contribute to understanding potential consequences of cortical α-syn, which is important for efforts investigating the role of this pathological finding in synucleinopathies such as Parkinson's disease dementia (PDD) and dementia with Lewy bodies (DLB).

The rodent PFC is a highly connected brain region, sending projections to multiple cortical, subcortical, midbrain, and brainstem nuclei [46,47]. Our prior work suggested that human PFFs injected into PFC cause a consistent but spatially limited pattern of α-syn pathology including PFC, temporal cortex, and striatum. The consequences of α-syn outside of the brainstem, as occurs in PDD and DLB, are still relatively unknown. Using the model of cortically-injected PFFs helps tease out the consequences of cortical/subcortical α-syn pathology in the absence of dramatic loss of neurons in brainstem regions. Using semi-quantitative analysis of 25 brain regions, we demonstrate dense pathological spread of α-syn in multiple cortical regions such as the orbitofrontal, secondary motor M2, perirhinal, entorhinal, and piriform. A consistent pattern of α-syn was also observed in subcortical regions, including the striatum, nucleus accumbens, amygdala, and thalamus. We note that there is differential aggregation of α-syn in subregions of the striatum, matching the connectivity of our prefrontal injection site and in line with prior work from Gabbott and colleagues (2005). We targeted the dorsal prelimbic and anterior cingulate subregions of the PFC, which contain a greater number of neurons that project to the dorsal striatum compared to the ventral prelimbic and infralimbic subregions of the PFC that contain more ventral striatal

projecting neurons [48]. We also note that, unlike other subcortical regions, the subregions of the

hippocampus were relatively spared from α-syn aggregation. This finding was again in line with prior

literature suggesting that cortical input to the hippocampus passes through the perirhinal and entorhinal

cortices first [49], two areas that show dense α-syn deposition here. While there was clear α-syn in

cortical and subcortical regions, we found little or no α-syn pathology in midbrain and brainstem nuclei,

including substantia nigra, locus coeruleus, and dorsal raphe. Together, these findings extend our prior

work to show that the aggregation induced in cortical and subcortical regions by human PFFs injected

into PFC are not cleared over the course of aging, as has been reported in some brain regions [50].

Notably, prior work from our group has shown that human PFFs injected in the PFC does not

affect interval timing or open field behavior 6 mpi in mice aged 15 months [23]. By contrast, prior work

has shown that people with PD display reliable deficits in interval timing [51–54] and that the PFC is

crucial for interval timing behavior [51,55,56]. The data presented here, and our prior report demonstrate

that PFC PFFs are not sufficient to affect interval timing. There are several interesting hypotheses that

could explain these results; 1) midbrain dopaminergic cell loss, which is absent in this model, may be the

main driver for interval timing deficits seen in PD and mouse models [28,51]; 2) human PFFs may

predominate in a PFC cell type that is unnecessary for interval timing; 3) cortical cells may function even

in the setting of α-syn aggregation, either through cellular or circuit-level compensation.

While α-syn aggregates were widespread and dense in cortical and subcortical regions, mice that

received PFC PFF injections displayed only mild behavioral deficits. PFF mice did not display any

difference in measures of interval timing, including variability of or mean switch times. This result in is

line with prior work from our group [23] and suggests that PFC PFFs and the resulting α-syn aggregation

is not sufficient to induce an interval timing deficit, even up to 20 mpi. We also found that PFF mice

displayed reduced thigmotaxis at 12 mpi, with the same trend observed at 16 mpi, which may suggest a

difference in explorative and anxiolytic behaviors in the open field as a result of PFC PFF injections.

Further, we also found that PFF mice showed rapid spatial acquisition but required a longer search route

to locate the target exit hole during the reversal phase of a daily spatial training Barnes maze task at 21

mpi, suggesting altered behavioral flexibility.

366   This study has several limitations. First, our control animals received monomer injections, which

has been shown previously to occasionally lead to mild and transient aggregation [57]. We did note some

mild p-α-syn positive puncta, but morphology was distinct and punctate, and may have been caused by

aging alone. Fibrillar and "lewy-neurite-like" structures were only seen in PFF injected mice, and this was

confirmed by confocal microscopy. Second, we did not investigate cell-type preference for human

PFF-induced α-syn pathology which could impact the response of the PFC and may be different from the

response to alternate models, such as brain homogenate-derived α-syn fibrils; for example, recent

publications have suggested human PFFs also induce significant glial inclusions [58,59]. Third, we did

not include a group of young mice to determine whether our lack of behavioral change was primarily due

to age-related decline in cognitive function or interval timing, which has been observed previously

[60,61], although the interval timing behavior presented here is similar to previously published work from

our group using healthy, young C57BL/6 mice [26]. Fourth, our measures of interval timing and

behavioral flexibility do not fully interrogate cognitive dysfunctions related to PDD and LBD and it is

possible that other cognitive behaviors such as attention or impulse control were negatively affected by

PFC PFF injections. Fifth, behavioral experiments at the later timepoints are underpowered to detect

subtle differences due to age-related loss. Despite these limitations, due to careful pathological

confirmation of each mouse in this two-year study, we believe these results inform future research and are

important for understanding this model and considerations for testing other models to induce aggregation.

Finally, this study was limited to male mice. Recent evidence suggest female mice may show reduced

pathology compared with male mice [62]. Human males are more likely to be diagnosed with PD and

DLB [63–65]. Understanding potential differences in pathology due to sex is an essential goal of future

research and fully powered studies of both sexes are needed.

388   In summary, our results describe widespread cortical and subcortical α-syn aggregation from PFC

389 PFFs with mild behavioral alterations that were not modulated by an inflammatory challenge. The lack of

390   brainstem involvement is notable and could provide insight into the brain networks that drive behavioral

391   deficits in PD, PDD, and DLB.

ACKNOWLEDGMENTS

This work was supported by NIH R01s MH116043, NS120987 to NSN and NIH K08 NS109287

and the Iowa Neuroscience Institute to GMA. This work was also supported by NIH P20NS123151 to

NSN and GMA. We thank Dr. Andrew West for generously providing the pre-formed fibrils and Travis

Larson for assisting with behavioral experiments.

DATA AVAILABILITY

All raw data are available at https://aldridge.lab.uiowa.edu/open-science.

CODE AVAILABILITY

All code is available at https://aldridge.lab.uiowa.edu/open-science.

AUTHOR CONTRIBUTIONS

MAW, HAA, QZ, NSN, and GMA designed the experiments. MAW, MMC, HAA, GK, OH, and

GMA performed all experiments. GK, RT, JG, and GMA performed histological analyses. MAW and

GMA performed all statistical analyses. MAW and GMA wrote the manuscript, and all authors reviewed

and revised the manuscript.

COMPETING INTERESTS

The authors have no conflict of interest to report.

REFERENCES

1. Braak, H., Tredici, K. D., Rüb, U., de Vos, R. A. I., Jansen Steur, E. N. H., & Braak, E. (2003). Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiology of Aging*, *24*(2), 197–211. https://doi.org/10.1016/S0197-4580(02)00065-9

2. Irwin, D. J., Lee, V. M.-Y., & Trojanowski, J. Q. (2013). Parkinson's disease dementia: Convergence of α-synuclein, tau and amyloid-β pathologies. *Nature Reviews Neuroscience*, *14*(9), 626–636. https://doi.org/10.1038/nrn3549

3. McKeith, I. G., Boeve, B. F., Dickson, D. W., Halliday, G., Taylor, J.-P., Weintraub, D., Aarsland, D., Galvin, J., Attems, J., Ballard, C. G., Bayston, A., Beach, T. G., Blanc, F., Bohnen, N., Bonanni, L., Bras, J., Brundin, P., Burn, D., Chen-Plotkin, A., … Kosaka, K. (2017). Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium. *Neurology*, *89*(1), 88–100. https://doi.org/10.1212/WNL.0000000000004058

4. Aldridge, G. M., Birnschein, A., Denburg, N. L., & Narayanan, N. S. (2018). Parkinson's Disease Dementia and Dementia with Lewy Bodies Have Similar Neuropsychological Profiles. *Frontiers in Neurology*, *9*, 123. https://doi.org/10.3389/fneur.2018.00123

5. FitzGerald, J. M., Perera, G., Chang-Tave, A., Price, A., Rajkumar, A. P., Bhattarai, M., O'Brien, J. T., Ballard, C., Aarsland, D., Stewart, R., & Mueller, C. (2019). The Incidence of Recorded Delirium Episodes Before and After Dementia Diagnosis: Differences Between Dementia With Lewy Bodies and Alzheimer's Disease. *Journal of the American Medical Directors Association*, *20*(5), 604–609. https://doi.org/10.1016/j.jamda.2018.09.021

6. Lawson, R. A., McDonald, C., & Burn, D. J. (2019). Defining delirium in idiopathic Parkinson's disease: A systematic review. *Parkinsonism & Related Disorders*, *64*, 29–39. https://doi.org/10.1016/j.parkreldis.2018.09.025

7. Halhouli, O., Zhang, Q., & Aldridge, G. M. (2022). Caring for patients with cognitive dysfunction, fluctuations and dementia caused by Parkinson's disease. In *Progress in Brain Research* (Vol. 269, pp. 407–434). Elsevier. https://doi.org/10.1016/bs.pbr.2022.01.018

8. Zhang, Q., Aldridge, G. M., Narayanan, N. S., Anderson, S. W., & Uc, E. Y. (2020). Approach to

Cognitive Impairment in Parkinson's Disease. *Neurotherapeutics*, *17*(4), 1495–1510.

https://doi.org/10.1007/s13311-020-00963-x

9. Kim, S., Kwon, S.-H., Kam, T.-I., Panicker, N., Karuppagounder, S. S., Lee, S., Lee, J. H., Kim, W. R.,

Kook, M., Foss, C. A., Shen, C., Lee, H., Kulkarni, S., Pasricha, P. J., Lee, G., Pomper, M. G.,

Dawson, V. L., Dawson, T. M., & Ko, H. S. (2019). Transneuronal Propagation of Pathologic

α-Synuclein from the Gut to the Brain Models Parkinson's Disease. *Neuron*, *103*(4), 627-641.e7.

https://doi.org/10.1016/j.neuron.2019.05.035

10. Luk, K. C., Song, C., O'Brien, P., Stieber, A., Branch, J. R., Brunden, K. R., Trojanowski, J. Q., &

Lee, V. M.-Y. (2009). Exogenous α-synuclein fibrils seed the formation of Lewy body-like

intracellular inclusions in cultured cells. *Proceedings of the National Academy of Sciences*,

106(47), 20051–20056. https://doi.org/10.1073/pnas.0908005106

11. Luk, K. C., Kehm, V., Carroll, J., Zhang, B., O'Brien, P., Trojanowski, J. Q., & Lee, V. M.-Y. (2012).

Pathological α-Synuclein Transmission Initiates Parkinson-like Neurodegeneration in

Nontransgenic Mice. *Science*, *338*(6109), 949–953. https://doi.org/10.1126/science.1227157

12. Volpicelli-Daley, L. A., Luk, K. C., Patel, T. P., Tanik, S. A., Riddle, D. M., Stieber, A., Meaney, D.

F., Trojanowski, J. Q., & Lee, V. M.-Y. (2011). Exogenous α-Synuclein Fibrils Induce Lewy

Body Pathology Leading to Synaptic Dysfunction and Neuron Death. *Neuron*, *72*(1), 57–71.

https://doi.org/10.1016/j.neuron.2011.08.033

13. Milanese, C., Cerri, S., Ulusoy, A., Gornati, S. V., Plat, A., Gabriels, S., Blandini, F., Di Monte, D.

A., Hoeijmakers, J. H., & Mastroberardino, P. G. (2018). Activation of the DNA damage

response in vivo in synucleinopathy models of Parkinson's disease. *Cell Death & Disease*, *9*(8),

818. https://doi.org/10.1038/s41419-018-0848-7

14. Ayers, J. I., Riffe, C. J., Sorrentino, Z. A., Diamond, J., Fagerli, E., Brooks, M., Galaleldeen, A., Hart,

P. J., & Giasson, B. I. (2018). Localized Induction of Wild-Type and Mutant Alpha-Synuclein

Aggregation Reveals Propagation along Neuroanatomical Tracts. *Journal of Virology*, *92*(18), e00586-18. https://doi.org/10.1128/JVI.00586-18

15. Challis, C., Hori, A., Sampson, T. R., Yoo, B. B., Challis, R. C., Hamilton, A. M., Mazmanian, S. K., Volpicelli-Daley, L. A., & Gradinaru, V. (2020). Gut-seeded α-synuclein fibrils promote gut dysfunction and brain pathology specifically in aged mice. *Nature Neuroscience*, *23*(3), 327–336. https://doi.org/10.1038/s41593-020-0589-7

16. Henrich, M. T., Geibl, F. F., Lakshminarasimhan, H., Stegmann, A., Giasson, B. I., Mao, X., Dawson, V. L., Dawson, T. M., Oertel, W. H., & Surmeier, D. J. (2020). Determinants of seeding and spreading of α-synuclein pathology in the brain. *Science Advances*, *6*(46), eabc2487. https://doi.org/10.1126/sciadv.abc2487

17. Rahayel, S., Mišić, B., Zheng, Y.-Q., Liu, Z.-Q., Abdelgawad, A., Abbasi, N., Caputo, A., Zhang, B., Lo, A., Kehm, V., Kozak, M., Yoo, H. S., Dagher, A., & Luk, K. C. (2022). Differentially targeted seeding reveals unique pathological alpha-synuclein propagation patterns. *Brain*, *145*(5), 1743–1756. https://doi.org/10.1093/brain/awab440

18. Rey, N. L., Steiner, J. A., Maroof, N., Luk, K. C., Madaj, Z., Trojanowski, J. Q., Lee, V. M.-Y., & Brundin, P. (2016). Widespread transneuronal propagation of α-synucleinopathy triggered in olfactory bulb mimics prodromal Parkinson's disease. *Journal of Experimental Medicine*, *213*(9), 1759–1778. https://doi.org/10.1084/jem.20160368

19. Rey, N. L., George, S., Steiner, J. A., Madaj, Z., Luk, K. C., Trojanowski, J. Q., Lee, V. M.-Y., & Brundin, P. (2018). Spread of aggregates after olfactory bulb injection of α-synuclein fibrils is associated with early neuronal loss and is reduced long term. *Acta Neuropathologica*, *135*(1), 65–83. https://doi.org/10.1007/s00401-017-1792-9

20. Uemura, N., Ueda, J., Yoshihara, T., Ikuno, M., Uemura, M. T., Yamakado, H., Asano, M., Trojanowski, J. Q., & Takahashi, R. (2021). α Synuclein Spread from Olfactory Bulb Causes Hyposmia, Anxiety, and Memory Loss in BAC SNCA Mice. *Movement Disorders*, *36*(9), 2036–2047. https://doi.org/10.1002/mds.28512

21. Espa, E., Clemensson, E. K. H., Luk, K. C., Heuer, A., Björklund, T., & Cenci, M. A. (2019). Seeding

of protein aggregation causes cognitive impairment in rat model of cortical synucleinopathy.

Movement Disorders, *34*(11), 1699–1710. https://doi.org/10.1002/mds.27810

22. Peeraer, E., Bottelbergs, A., Van Kolen, K., Stancu, I.-C., Vasconcelos, B., Mahieu, M.,

Duytschaever, H., Ver Donck, L., Torremans, A., Sluydts, E., Van Acker, N., Kemp, J. A.,

Mercken, M., Brunden, K. R., Trojanowski, J. Q., Dewachter, I., Lee, V. M. Y., & Moechars, D.

(2015). Intracerebral injection of preformed synthetic tau fibrils initiates widespread tauopathy

and neuronal loss in the brains of tau transgenic mice. *Neurobiology of Disease*, *73*, 83–95.

https://doi.org/10.1016/j.nbd.2014.08.032

23. Zhang, Q., Abdelmotilib, H., Larson, T., Keomanivong, C., Conlon, M., Aldridge, G. M., &

Narayanan, N. S. (2021). Cortical alpha-synuclein preformed fibrils do not affect interval timing

in mice. *Neuroscience Letters*, *765*, 136273. https://doi.org/10.1016/j.neulet.2021.136273

24. Abdelmotilib, H., Maltbie, T., Delic, V., Liu, Z., Hu, X., Fraser, K. B., Moehle, M. S., Stoyka, L.,

Anabtawi, N., Krendelchtchikova, V., Volpicelli-Daley, L. A., & West, A. (2017). α-Synuclein

fibril-induced inclusion spread in rats and mice correlates with dopaminergic Neurodegeneration.

Neurobiology of Disease, *105*, 84–98. https://doi.org/10.1016/j.nbd.2017.05.014

25. Balci, F., Papachristos, E. B., Gallistel, C. R., Brunner, D., Gibson, J., & Shumyatsky, G. P. (2008).

Interval timing in genetically modified mice: A simple paradigm. *Genes, Brain and Behavior*,

7(3), 373–384. https://doi.org/10.1111/j.1601-183X.2007.00348.x

26. Bruce, R. A., Weber, M. A., Volkman, R. A., Oya, M., Emmons, E. B., Kim, Y., & Narayanan, N. S.

(2021). Experience-related enhancements in striatal temporal encoding. *The European Journal of*

Neuroscience, *54*(3), 5063–5074. https://doi.org/10.1111/ejn.15344

27. Larson, T., Khandelwal, V., Weber, M. A., Leidinger, M. R., Meyerholz, D. K., Narayanan, N. S., &

Zhang, Q. (2022). Mice expressing P301S mutant human tau have deficits in interval timing.

Behavioural Brain Research, *432*, 113967. https://doi.org/10.1016/j.bbr.2022.113967

28. Weber, M. A., Sivakumar, K., Tabakovic, E. E., Oya, M., Aldridge, G. M., Zhang, Q., Simmering, J.

E., & Narayanan, N. S. (2023). Glycolysis-enhancing α1-adrenergic antagonists modify cognitive

symptoms related to Parkinson's disease. *Npj Parkinson's Disease*, *9*(1), 32.

https://doi.org/10.1038/s41531-023-00477-1

29. Gawel, K., Gibula, E., Marszalek-Grabska, M., Filarowska, J., & Kotlinska, J. H. (2019). Assessment

of spatial learning and memory in the Barnes maze task in rodents—Methodological

consideration. *Naunyn-Schmiedeberg's Archives of Pharmacology*, *392*(1), 1–18.

https://doi.org/10.1007/s00210-018-1589-y

30. O'Leary, T. P., & Brown, R. E. (2013). Optimization of apparatus design and behavioral measures for

the assessment of visuo-spatial learning and memory of mice on the Barnes maze. *Learning &*

Memory, *20*(2), 85–96. https://doi.org/10.1101/lm.028076.112

31. Kraeuter, A.-K., Guest, P. C., & Sarnyai, Z. (2019). The Open Field Test for Measuring Locomotor

Activity and Anxiety-Like Behavior. In P. C. Guest (Ed.), *Pre-Clinical Models* (Vol. 1916, pp.

99–103). Springer New York. https://doi.org/10.1007/978-1-4939-8994-2_9

32. Tatem, K. S., Quinn, J. L., Phadke, A., Yu, Q., Gordish-Dressman, H., & Nagaraju, K. (2014).

Behavioral and Locomotor Measurements Using an Open Field Activity Monitoring System for

Skeletal Muscle Diseases. *Journal of Visualized Experiments*, *91*, 51785.

https://doi.org/10.3791/51785

33. Fisher, S. P., Godinho, S. I. H., Pothecary, C. A., Hankins, M. W., Foster, R. G., & Peirson, S. N.

(2012). Rapid Assessment of Sleep-Wake Behavior in Mice. *Journal of Biological Rhythms*,

27(1), 48–58. https://doi.org/10.1177/0748730411431550

34. Neely, C. L. C., Pedemonte, K. A., Boggs, K. N., & Flinn, J. M. (2019). Nest Building Behavior as an

Early Indicator of Behavioral Deficits in Mice. *Journal of Visualized Experiments*, *152*, 60139.

https://doi.org/10.3791/60139

35. Gaskill, B. N., Karas, A. Z., Garner, J. P., & Pritchett-Corning, K. R. (2013). Nest Building as an

Indicator of Health and Welfare in Laboratory Mice. *Journal of Visualized Experiments*, *82*,

51012. https://doi.org/10.3791/51012

36. Matsuo, N., Morita, T., Matsuda, Y., Okamoto, K., Matsumoto, Y., Kaneishi, K., Odagiri, T., Sakurai,

H., Katayama, H., Mori, I., Yamada, H., Watanabe, H., Yokoyama, T., Yamaguchi, T., Nishi, T.,

Shirado, A., Hiramoto, S., Watanabe, T., Kohara, H., … Iwase, S. (2017). Predictors of Delirium

in Corticosteroid-Treated Patients with Advanced Cancer: An Exploratory, Multicenter,

Prospective, Observational Study. *Journal of Palliative Medicine*, *20*(4), 352–359.

https://doi.org/10.1089/jpm.2016.0323

37. Quispel⬜Aggenbach, D. W. P., Schep⬜de Ruiter, E. P. R., Bergen, W., Bolling, J. R., Zuidema, S.

U., & Luijendijk, H. J. (2021). Prevalence and risk factors of delirium in psychogeriatric

outpatients. *International Journal of Geriatric Psychiatry*, *36*(1), 190–196.

https://doi.org/10.1002/gps.5413

38. Spiller, J. A., & Keen, J. C. (2006). Hypoactive delirium: Assessing the extent of the problem for

inpatient specialist palliative care. *Palliative Medicine*, *20*(1), 17–23.

https://doi.org/10.1191/0269216306pm1097oa

39. Culley, D. J., Snayd, M., Baxter, M. G., Xie, Z., Lee, I. H., Rudolph, J., Inouye, S. K., Marcantonio,

E. R., & Crosby, G. (2014). Systemic inflammation impairs attention and cognitive flexibility but

not associative learning in aged rats: Possible implications for delirium. *Frontiers in Aging*

Neuroscience, *6*, 107. https://doi.org/10.3389/fnagi.2014.00107

40. Field, R. H., Gossen, A., & Cunningham, C. (2012). Prior Pathology in the Basal Forebrain

Cholinergic System Predisposes to Inflammation-Induced Working Memory Deficits:

Reconciling Inflammatory and Cholinergic Hypotheses of Delirium. *Journal of Neuroscience*,

32(18), 6288–6294. https://doi.org/10.1523/JNEUROSCI.4673-11.2012

41. Sparkman, N. L., Buchanan, J. B., Heyen, J. R. R., Chen, J., Beverly, J. L., & Johnson, R. W. (2006).

Interleukin-6 Facilitates Lipopolysaccharide-Induced Disruption in Working Memory and

Expression of Other Proinflammatory Cytokines in Hippocampal Neuronal Cell Layers. *Journal*

of Neuroscience, *26*(42), 10709–10716. https://doi.org/10.1523/JNEUROSCI.3376-06.2006

42. Cunningham, C., Campion, S., Lunnon, K., Murray, C. L., Woods, J. F. C., Deacon, R. M. J.,

Rawlins, J. N. P., & Perry, V. H. (2009). Systemic Inflammation Induces Acute Behavioral and

Cognitive Changes and Accelerates Neurodegenerative Disease. *Biological Psychiatry*, *65*(4),

304–312. https://doi.org/10.1016/j.biopsych.2008.07.024

43. Torvell, M., Hampton, D. W., Connick, P., MacLullich, A. M. J., Cunningham, C., & Chandran, S.

(2019). A single systemic inflammatory insult causes acute motor deficits and accelerates disease

progression in a mouse model of human tauopathy. *Alzheimer's & Dementia: Translational*

Research & Clinical Interventions, *5*(1), 579–591. https://doi.org/10.1016/j.trci.2019.09.001

44. Franklin, K. B. J., & Paxinos, G. (2008). *The mouse brain in stereotaxic coordinates* (Compact 3. ed).

Elsevier Academic Press.

45. Rey, N. L., Bousset, L., George, S., Madaj, Z., Meyerdirk, L., Schulz, E., Steiner, J. A., Melki, R., &

Brundin, P. (2019). α-Synuclein conformational strains spread, seed and target neuronal cells

differentially after injection into the olfactory bulb. *Acta Neuropathologica Communications*,

7(1), 221. https://doi.org/10.1186/s40478-019-0859-3

46. Anastasiades, P. G., & Carter, A. G. (2021). Circuit organization of the rodent medial prefrontal

cortex. *Trends in Neurosciences*, *44*(7), 550–563. https://doi.org/10.1016/j.tins.2021.03.006

47. Souza, R., Bueno, D., Lima, L. B., Muchon, M. J., Gonçalves, L., Donato, J., Shammah-Lagnado, S.

J., & Metzger, M. (2022). Top-down projections of the prefrontal cortex to the ventral tegmental

area, laterodorsal tegmental nucleus, and median raphe nucleus. *Brain Structure and Function*,

227(7), 2465–2487. https://doi.org/10.1007/s00429-022-02538-2

48. Gabbott, P. L. A., Warner, T. A., Jays, P. R. L., Salway, P., & Busby, S. J. (2005). Prefrontal cortex in

the rat: Projections to subcortical autonomic, motor, and limbic centers. *The Journal of*

Comparative Neurology, *492*(2), 145–177. https://doi.org/10.1002/cne.20738

49. Burwell, R. D., & Amaral, D. G. (1998). Cortical afferents of the perirhinal, postrhinal, and entorhinal

cortices of the rat. *The Journal of Comparative Neurology*, *398*(2), 179–205.

https://doi.org/10.1002/(SICI)1096-9861(19980824)398:2<179::AID-CNE3>3.0.CO;2-Y

50. Sun, F., Salinas, A. G., Filser, S., Blumenstock, S., Medina☐Luque, J., Herms, J., & Sgobio, C.

(2022). Impact of α☐synuclein spreading on the nigrostriatal dopaminergic pathway depends on

the onset of the pathology. *Brain Pathology*, *32*(2). https://doi.org/10.1111/bpa.13036

51. Kim, Y.-C., Han, S.-W., Alberico, S. L., Ruggiero, R. N., De Corte, B., Chen, K.-H., & Narayanan,

N. S. (2017). Optogenetic Stimulation of Frontal D1 Neurons Compensates for Impaired

Temporal Control of Action in Dopamine-Depleted Mice. *Current Biology*, *27*(1), 39–47.

https://doi.org/10.1016/j.cub.2016.11.029

52. Malapani, C., Rakitin, B., Levy, R., Meck, W. H., Deweer, B., Dubois, B., & Gibbon, J. (1998).

Coupled Temporal Memories in Parkinson's Disease: A Dopamine-Related Dysfunction. *Journal*

of Cognitive Neuroscience, *10*(3), 316–331. https://doi.org/10.1162/089892998562762

53. Pastor, M. A., Artieda, J., Jahanshahi, M., & Obeso, J. A. (1992). TIME ESTIMATION AND

REPRODUCTION IS ABNORMAL IN PARKINSON'S DISEASE. *Brain*, *115*(1), 211–225.

https://doi.org/10.1093/brain/115.1.211

54. Singh, A., Cole, R. C., Espinoza, A. I., Evans, A., Cao, S., Cavanagh, J. F., & Narayanan, N. S.

(2021). Timing variability and midfrontal ~4 Hz rhythms correlate with cognition in Parkinson's

disease. *Npj Parkinson's Disease*, *7*(1), 14. https://doi.org/10.1038/s41531-021-00158-x

55. Emmons, E. B., De Corte, B. J., Kim, Y., Parker, K. L., Matell, M. S., & Narayanan, N. S. (2017).

Rodent Medial Frontal Control of Temporal Processing in the Dorsomedial Striatum. *The Journal*

of Neuroscience, *37*(36), 8718–8733. https://doi.org/10.1523/JNEUROSCI.1376-17.2017

56. Narayanan, N. S., Land, B. B., Solder, J. E., Deisseroth, K., & DiLeone, R. J. (2012). Prefrontal D1

dopamine signaling is required for temporal control. *Proceedings of the National Academy of*

Sciences, *109*(50), 20726–20731. https://doi.org/10.1073/pnas.1211258109

57. Manfredsson, F. P., Luk, K. C., Benskey, M. J., Gezer, A., Garcia, J., Kuhn, N. C., Sandoval, I. M.,

Patterson, J. R., O'Mara, A., Yonkers, R., & Kordower, J. H. (2018). Induction of

alpha-synuclein pathology in the enteric nervous system of the rat and non-human primate results

in gastrointestinal dysmotility and transient CNS pathology. *Neurobiology of Disease*, *112*, 106–

118. https://doi.org/10.1016/j.nbd.2018.01.008

58. Uemura, N., Uemura, M. T., Lo, A., Bassil, F., Zhang, B., Luk, K. C., Lee, V. M.-Y., Takahashi, R.,

& Trojanowski, J. Q. (2019). Slow Progressive Accumulation of Oligodendroglial

Alpha-Synuclein (α-Syn) Pathology in Synthetic α-Syn Fibril-Induced Mouse Models of

Synucleinopathy. *Journal of Neuropathology & Experimental Neurology*, *78*(10), 877–890.

https://doi.org/10.1093/jnen/nlz070

59. Uemura, N., Marotta, N., Ara, J., Meymand, E., Zhang, B., Kameda, H., Koike, M., Luk, K.,

Trojanowski, J., & Lee, V. (2023). *Distinct biological activity of Lewy body α-Synuclein strain in*

mice [Preprint]. In Review. https://doi.org/10.21203/rs.3.rs-2579805/v1

60. Beas, B. S., McQuail, J. A., Bañuelos, C., Setlow, B., & Bizon, J. L. (2017). Prefrontal cortical

GABAergic signaling and impaired behavioral flexibility in aged F344 rats. *Neuroscience*, *345*,

274–286. https://doi.org/10.1016/j.neuroscience.2016.02.014

61. Gür, E., Duyan, Y. A., Arkan, S., Karson, A., & Balcı, F. (2020). Interval timing deficits and their

neurobiological correlates in aging mice. *Neurobiology of Aging*, *90*, 33–42.

https://doi.org/10.1016/j.neurobiolaging.2020.02.021

62. Gezer, A. O. (2020). Sex-specific Effects of Developmental Dieldrin Exposure on Synucleinopathy

[Ph.D., Michigan State University]. In *ProQuest Dissertations and Theses* (2404294669).

Dissertations & Theses @ CIC Institutions; ProQuest Dissertations & Theses Global.

http://login.proxy.lib.uiowa.edu/login?url=https://www.proquest.com/dissertations-theses/sex-spe

cific-effects-developmental-dieldrin/docview/2404294669/se-2?accountid=14663

63. Savica, R., Grossardt, B. R., Bower, J. H., Boeve, B. F., Ahlskog, J. E., & Rocca, W. A. (2013).

Incidence of Dementia With Lewy Bodies and Parkinson Disease Dementia. *JAMA Neurology*,

70(11), 1396. https://doi.org/10.1001/jamaneurol.2013.3579

64. Van Den Eeden, S. K. (2003). Incidence of Parkinson's Disease: Variation by Age, Gender, and

Race/Ethnicity. *American Journal of Epidemiology*, *157*(>11), 1015–1022.

https://doi.org/10.1093/aje/kwg068

65. Wooten, G. F. (2004). Are men at greater risk for Parkinson's disease than women? *Journal of*

Neurology, Neurosurgery & Psychiatry, *75*(4), 637–639.

https://doi.org/10.1136/jnnp.2003.020982

FIGURES

Figure 1. Experimental timeline.

Figure 2. Evaluation of pathology in monomer control and pre-formed fibril (PFF) injected mice.

A) Top: representative high-resolution images across the entire rostral-caudal aspect of the brain in two

mice treated with PFFs (one saline and one LPS treated at 16 mpi, sacrificed at 21 mpi). Bottom:

representative images of two monomer control mice (one saline and one LPS treated at 16 mpi, sacrificed

at 21 mpi). Semi-quantitative scores of phospho α-syn (p-α-syn) positive **C)** somas and **D)** neurites of 25

brain regions, with each column representing an individual mouse, averaged over the entire rostral-caudal

extent for each brain region examined. Neurites for monomer mice are not displayed as there were zero

p-α-syn positive neurites in any region of any mouse. **E)** Confocal images showing examples of fibrillar

pathology in multiple brain regions from PFF injected mice (panels 1–4). Monomer mice showed rare

p-α-syn positive puncta, without clear fibrillar morphology (panel 5). No fibrillar or neurite pathology

was found in monomer mice using confocal microscopy. **F)** Averaged semi-quantitative score of

pathological α-syn within subregions of the striatum restricted to 0–1.7 mm bregma. Horizontal bars

indicate group medians. * $p < 0.05$, **** $p < 0.0001$ **G)** Tyrosine hydroxylase intensity (arbitrary units)

from 4 different brain regions PFF treated mice: substantia nigra (SN), ventral tegmental area (VTA),

striatum, and prefrontal cortex (PFC). Tyrosine hydroxylase intensity expressed as mean ± SEM, and

each dot represents a single mouse.

Figure 3. Interval timing and reversal learning in monomer control and PFC PFF mice. A) Diagram

for the interval timing switch task outlining optimal performance during long trials. **B)** Mean switch time

and **C)** switch time coefficient of variance during the interval timing task at 12 mpi. Data from 8

monomer (blue) and 7 PFF (red) mice. **D)** Mean switch time and **E)** switch time coefficient of variance

during the interval timing task. Data from 6 monomer (blue) and 8 PFF (red) mice. Each dot represents a

single mouse. **F)** Diagram for the rapid reversal Barnes maze. **G)** Average total distance travelled

averaged across three-days. There was a main effect of trial and trial × treatment interaction. * $p < 0.05$,

but post-hoc testing at specific trials did not survive multiple comparisons. Data from 6 monomer (blue)

and 6 PFF (red) mice. Data are expressed as mean ± SEM.

Figure 4. Open field, home cage activity, and nest building in monomer control and PFC PFF mice at 12–15 mpi. A) Distance travelled and **B)** thigmotaxis during the open field test. The right panel is an example trace from one monomer and one PFF mouse during the open field assay. **C)** Percent immobility time between ~2–3 am (night) and ~6–7 am (morning). **D)** Nest building score for three different materials (shredded paper, cotton twist, and precut cotton squares) with representative images from each material shown in the right panel. Data from 8 monomer (blue) and 7-8 PFF (red) mice. All data are expressed as mean ± SEM, and each dot represents a single mouse. * p < 0.05

Figure 5. Open field, food burying, and nest building during 0.5 mg/kg LPS challenge. A) Distance

travelled and **B)** thigmotaxis during the open field after either saline or lipopolysaccharide (LPS)

injection. There was a main effect of LPS on distance traveled, but this effect was not different for

monomer and PFF mice. One monomer mouse treated with LPS was excluded from thigmotaxis analysis

as no movement was recorded during the assay. **C)** Amount of food buried (in grams) during the food

burying assay and **D)** nest building score (shredded paper only) for 2 days before and 3 days after either

saline or LPS injection. All data are expressed as mean ± SEM, and each dot represents a single mouse. *

$p < 0.05$; *** $p < 0.001$; **** $p < 0.0001$

SUPPLEMENTAL MATERIALS

Figure S1. Additional measures of interval timing performance and behavioral flexibility.

Proportion of rewarded trials during the interval timing switch task at **A)** 12 mpi and **B)** 20 mpi. Unpaired

t-tests revealed no differences in the proportion correct at either timepoint (12 mpi: $t(13) = 2.03$; $p = 0.06$;

20 mpi: $t(12) = 1.2$; $p = 0.25$) **C)** Average total distance travelled during a single day rapid reversal

Barnes maze protocol. Compared to the multiple day Barnes maze protocol at 21 mpi, PFC PFF mice

show a similar, but non-significant, altered pattern of behavioral flexibility with a greater distance

travelled to reach the target during the reversal phase (two-way repeated-measures ANOVA; trial: $F_{(2.182, 30.55)} = 4.232$; $p = 0.02$; treatment: $F_{(1, 14)} = 0.2317$; $p = 0.64$) . All data are expressed as mean ± SEM, and

each dot represents a single mouse.

0.25 mg/kg LPS: 21 mpi

Figure S2. Lower dose lipopolysaccharide (LPS) challenge (0.25 mg/kg) at 21 mpi. A) Distance travelled and **B)** thigmotaxis during the open field test. PFC PFF injected mice did not differ from controls in distance travelled (meters) either 6- or 24-hours post LPS injection (two-way repeated-measures ANOVA; timepoint: $F_{(1, 9)} = 5.115$; $p = 0.05$; treatment: $F_{(1, 9)} = 1.272$; $p = 0.29$). However, PFC PFF mice did display a non-significant trend for reduced thigmotaxis (two-way repeated-measures ANOVA; timepoint: $F_{(1, 10)} = 0.003$; $p = 0.96$; treatment: $F_{(1, 9)} = 4.096$; $p = 0.07$), regardless of timepoint post injection ($F_{(1, 10)} = 0.003$; $p = 0.96$). **C)** Average total distance travelled averaged across two-days post LPS of the rapid reversal Barnes maze (two-way repeated-measures ANOVA; trial: $F_{(2.626, 21.01)} = 2.667$; $p = 0.08$; treatment: $F_{(1, 8)} = 0.034$; $p = 0.86$). **D)** Fluorescent intensity (arbitrary units) of GFAP and IBA1 in whole striatum. All data are expressed as mean ± SEM, and each dot represents a single mouse.