1 **Hybridization Mediated Range Expansion and Climate Change Resilience in Two Keystone Tree**
2 **Species of Boreal Forests**
3

Piyal Karunarathne[1,2,3]*, Qiujie Zhou[1,2], Martin Lascoux[1,2], and Pascal Milesi[1,2]†
1. Plant Ecology and Evolution, Department of Ecology and Genetics, Uppsala University,
Norbyvägen 18D, 75236 Uppsala, Sweden
2. Science for Life Laboratory (SciLifeLab), Uppsala, Sweden
3. Institute of Population Genetics, Heinrich-Heine University, Düsseldorf, Universitäts Straße 1,
40225 Düsseldorf, Germany
* piyal.karunarathne@hhu.de
† pascal.milesi@scilifelab.uu.se

12
13

14 **Abstract**
15

Current global climate change is expected to affect biodiversity negatively at all scales leading to mass
biodiversity loss. Many studies have shown that the distribution of allele frequencies across a species'
range is often influenced by specific genetic loci associated with local environmental variables. This
association reflects local adaptation and allele changes at those loci could thereby contribute to the
evolutionary response to climate change. However, predicting how species will adapt to climate change
from this type of data alone remains challenging. In the present study, we combined exome capture
sequences and environmental niche reconstruction, to test multiple methods for assessing local
adaptation and climate resilience in two widely distributed conifers, Norway spruce and Siberian
spruce. Both species are keystone species of the boreal forest and share a vast hybrid zone. We show
that local adaptation in conifers can be detected through allele frequency variation, population-level
ecological preferences, and historical niche movement. Moreover, we integrated genetic and ecological
information into genetic offset predictive models to show that hybridization plays a central role in
expanding the niche breadth of the two conifer species and may help both species to cope better with
future changing climates. This joint genetic and ecological analysis also identified genetically isolated
populations that are at risk under current climate change.

31
32

33 **Keywords:** *Eco-evolution, Climate change, Landscape genomics, hybridization, forest trees*

Background

Human activities have led to the destruction of the habitat of an increasing number of species and climate change has only exacerbated this situation (Bellard et al., 2012; Grimm et al., 2013; Walther et al., 2002). Given the rapid pace of global warming, ecological communities face an urgent need for rapid adaptation in their structure and composition to cope with these ongoing challenges. Interestingly, recent studies have revealed that communities tend to respond more strongly to regional than global changes (Altieri & Gedan, 2015; Araújo & Rahbek, 2006; Atkins & Travis, 2010; Thuiller et al., 2005; Williams et al., 2021; Wuethrich, 2000). Climate change impacts vary across ecosystems, leading to diverse ecological dynamics and unpredictable changes within local environments (Carbeck et al., 2022; Walther et al., 2002). In order to mitigate the risk of demographic collapse and potential extinction, organisms must either track their climatic niche, be phenotypically plastic, or adapt to the changing environment from existing genetic variation or through de novo mutations. Adaptations of populations to local conditions result from a subtle balance between the intensity of natural selection and migration between the populations. While the former pushes the populations towards their local optima the latter tends to homogenize the gene pools between the populations (for a review see Bonnet et al., 2022; Lenormand, 2002). Still, some level of gene flow, although not an absolute requirement, contributes to promoting local and global genetic diversity and preserving the adaptive potential of populations (e.g., Frankham, 2005; Markert et al., 2010; Pauls et al., 2013). Global climate change poses significant challenges as it can affect simultaneously many populations, placing species and their populations at greater vulnerability in limiting their ability to cope with environmental changes and potentially leading to extinction (Leroy et al., 2020; Pauls et al., 2013; Rellstab et al., 2016).

Plants are probably even more affected by global change than animals due to their inherent disadvantage in tracing suitable habitats, especially species with long generation times (Dauphin et al., 2021; Leroy et al., 2020). Therefore, the survival of a long-lived plant species depends critically upon the potential of the species to expand into new regions through past adaptations to such environments as well as the potential for populations at the edge of its distribution range to face new environmental conditions (Pfennig et al., 2016; Taylor et al., 2015). Therefore, to comprehensively understand the interplay between environmental shifts and their genetic consequences, it is imperative to employ eco-evolutionary models (Bush et al., 2016; Thuiller et al., 2013). For instance, Cotto et al., (2017) pioneered an innovative approach by integrating ecological niche modeling with individual-based genetics modeling to explore how populations in the Alpine range respond to evolving environmental conditions. Similarly, genetic offset methodologies are designed to predict the disparity between a population's current status and the necessary adaptations to be able to survive under future climatic scenarios (e.g., Rellstab et al., 2021).

However, these approaches encounter certain limitations when applied on a continental scale, particularly with regard to accounting for intricate demographic histories and ongoing gene flow dynamics. In our present study, we have taken a multifaceted approach by seamlessly merging ecological niche modeling, landscape genomics, and genotype-environment association analyses. Our aim is to unravel the pivotal role of recurrent hybridization in facilitating the adaptation of two cornerstone species within the boreal forest ecosystem to the challenges posed by global environmental changes.

In Western Europe, recurrent cycles of population contraction and expansion following glacial cycles shaped both genetic diversity (Milesi et al., 2023; Petit et al., 2003) and local adaptation patterns along environmental gradients that also correspond to post-glacial recolonization routes (L. Li et al., 2022; Savolainen et al., 2013; Thörn et al., 2021). Genetic admixture following range expansion from refugia can serve as a major source of genetic diversity in forest trees (J. Chen et al., 2019; Lexer et al., 2010; Müller et al., 2019; Tsuda et al., 2017; Zhou et al., 2023). It can promote increased climate resilience in wild populations (e.g., Inoue & Berg, 2017; Meester et al., 2018) and aid wider phenotypic plasticity against rapid climate changes (see Catullo et al., 2019)). Admixture caused by recurrent colonization by superior ecotypes can promote better resistance to environmental stress (e.g., (Hamilton & Miller, 2016; Ortego et al., 2015) notably through an effective population size increase, persistence at the range edge and niche expansion (e.g., Gillies et al., 2016; Teukeng et al., 2022; Zalapa et al., 2010). Hybridization and admixture are thus effective mechanisms to increase overall genetic diversity and can further promote the transfer of adaptive alleles from one genetic background to another, enhancing local adaption and allowing populations to respond quickly to selective pressures (Rendón-Anaya et al., 2021; Rieseberg & Willis, 2007; Stelkens et al., 2014; Taylor et al., 2015). Here we hypothesize that the recurrent hybridization and admixture events between two closely related coniferous species through cycles of historic range shifts due to ice cycles may help both species to better cope with future climatic changes.

Norway spruce (*Picea abies* (L.) H. Karst) and Siberian spruce (*Picea obovata* Ledeb.) are two closely related conifer species that diverged several million years ago (J. Chen et al., 2019; Tsuda et al., 2016; Zhou et al., 2023). Today, the two species have clear western (Western Europe) and eastern (Siberian Russia) distributions, albeit with a known hybrid zone located between west of the Ural Mountains and expanding up to Northern Fennoscandia (Tollefsrud et al., 2015; Tsuda et al., 2016; Zhou et al., 2023: See map Fig. 1). Currently, the distribution of *P. abies* is subdivided into seven main genetic clusters (J. Chen et al., 2019; L. Li et al., 2022). They correspond to three ancestral clusters (Alpine, Carpathian, Fennoscandian) as well as to admixture between these three main clusters (J. Chen et al., 2019; L. Li et al., 2022). Siberian spruce, on the other hand, expanded westwards from Northeast Asia to Western Russia. Following glacial cycles, the two species came into contact on several occasions, leading to

introgression (e.g., J. Chen et al., 2019; Tollefsrud et al., 2015; Tsuda et al., 2016; Zhou et al., 2023). Recently, Nota et al., (2022), using ancient DNA, and Li et al., (2022), using contemporary nuclear markers, confirmed conclusions drawn from pollen records (e.g. Giesecke, 2005) that recolonization of the Scandinavian peninsula by Norway spruce after the last glacial maximum (LGM) occurred through two different entry points, a southern one, from Eastern location across the Baltic Sea and a northern one through northern Finland with a contribution from *P. obovata*. Li et al. (2022) demonstrated that the contact zone between the two main genetic clusters aligns with the boundary between Scandinavia's primary climatic regions, revealing the influence of natural selection in maintaining this zone and shaping spruce genetic diversity. While pollen records are valuable for inferring species distribution and population movement during the Holocene (e.g., Trondman et al. 2014), they offer limited insights, particularly at the species level, for events predating that period, as observed in the case of Norway and Siberian spruce (J. Chen et al., 2019; Tsuda et al., 2016; Zhou et al., 2023).

We here used a combination of genome scans, genotype-environment association studies, and ecological niche modeling (for both past and future scenarios) to demonstrate that recurrent hybridization significantly contributes to the expansion of both species' niche breadth, potentially enabling them to better cope with changing climates. Given the close genetic affinity, the extensive hybrid zone, historic recolonization events, and the climatic gradient occupied by Norway and Siberian spruce, they provide a unique system for studying the role of hybridization in shaping climate resilience through genetic variation and range expansion. Our study first provides evidence of local adaptation and identifies key climatic variables influencing species distributions. We then use environmental niche models with these climatic variables and species occurrences to define the ecological niche preferences of both species and their historical movements. Finally, we integrate genetic and ecological information into genetic offset predictive models to assess the resilience of each population to future climatic conditions.

Results

Populations exhibiting a high level of admixture are spread across three climatic groups

We used the same sampling of 55 populations covering the whole range of *P. abies* and the western part of the distribution range of *P. obovata* as in Zhou et al. (2023). Using a combination of population genetics analyses, the authors showed that there are two main genetic entities: one western cluster, corresponding to Norway spruce (*P. abies*), centered on the Alps and the Carpathians, and extending towards Scandinavia and the Russian Plains, and one eastern cluster stemming from Siberia, east of the Urals Mountains and corresponding to what has generally been recognized as Siberian spruce (*P. obovata*). The populations previously recognized as of hybrid nature located west of the Ural Mountains and east of the Baltic region formed sub-clusters within the two main clusters (Fig. S1A also see Zhou et al. 2023 Fig. S3). Following this initial clustering, in order to quantify the degree of admixture between *P. abies* and *P. obovata*, we derived a *hybrid index* for each individual using ADMIXTURE

(K=2, see materials and methods and Fig. 1A & S1B) and averaged it per population. The distribution

of the *hybrid index* is highly consistent with the Bayesian clustering. To investigate whether specific

DNA polymorphisms were associated with given climatic zones, we used redundancy analysis (RDA)

incorporating both genetic and climatic data (see *Genetic diversity distribution among populations* in

materials and methods). We defined five major environmental clusters (i.e., genetically informed

climatic groups) among all the populations studied, with populations in the hybrid zone belonging to

three different clusters (Fig. 1B). Interestingly, this divided western populations (i.e., *P. abies*) into

three groups: a southern group consisting of Alpine, Carpathian and Central European populations, a

northern group made of Fennoscandian populations and, a third group with central European and Baltic

populations, while eastern populations (i.e., *P. obovata*) were separated into a northern and a southern

group along the southern border of the west Siberian taiga (Fig. 1B).

169
170

171   **Figure 1: Climatic clusters and hybrid index (*hi*) between *P. abies* and *P. obovata;* A.** Spatial

172   projection of the hybrid index computed from ancestry coefficient using K=2 [For the analyzed alleles,

173   *hi* = 1 = *P. abies* = red, *hi* = 0 = *P.obovata* = blue]. **B.** Genetically informed climatic clusters of the

174   studied populations; clusters were identified with k-means clustering of environmental and adaptive

175   index variables (see methods) [Cent: central, FenoSc: Fennoscandian, Nr: northern, Sth: southern].

Spatial heterogeneity and environmental conditioning of genetic diversity

We investigated changes in total nucleotide diversity (π) across the whole distribution range of the

species. Globally, π ranging from 0.00518 to 0.00695, the highest values were found in the North-

Eastern range for *P. obovata* and Southern Scandinavia for *P. abies*. Overall, *P. obovata* populations

exhibit higher π values than *P. abies*, likely reflecting more stable historical distribution and more

continuous distribution through space (Fig. 2A). In *P. obovata*, a clear shift in π matches the location

of the Ural Mountain, which likely limits the gene flow. Pockets of high diversity are found in the

Southernmost range of *P. obovata*, west of the Urals, possibly reflecting refugia during the ice age.

While the overall genetic diversity within *P. abies* also ranges between 0.0054 and 0.0068, the northern-

and southern-most populations had the lowest nucleotide diversity. Even though southern locations in

Italy and the Alpine mountains region have been refugia for Norway spruce during the ice ages, the

current relative isolation of the populations because of the topography, probably explains the relatively

lower genetic diversity. On the other hand, the northernmost populations of *P. abies* in Sweden harbor
the lowest genetic diversity. These populations are also the youngest and the low nucleotide diversity
may reflect the colonization forefront. Nevertheless, the highest π values are found in Southern
Scandinavia, an area that was also recolonized after the LGM but from different populations. Finally,
populations located in contact zones (e.g., in Scandinavia between the Northern and the Southern range)
or located within the large hybrid zone between *P. abies* and *P. obovata* exhibit higher π values than
neighboring populations, exhibiting the effect of introgression and/or admixture on genetic diversity.
As expected, hybrid populations also show intermediate heterozygosity between *P. abies* and *P.*
obovata (linear model, $F = 6.55$, $df = 34$, $p = 0.015$, Fig. 2B). More importantly, hybrid populations
tend to have more homogeneous allele frequencies from one SNP to another than in *P. abies* (Welch *t*-
test, $t = -2.85$, $df = 11.172$, $p = 0.02$) or *P. obovata* ($t = -5.63$, $df = 7.93$, $p < 0.001$, Fig. 2C): Note that
standard deviation in allele frequencies vary quantitively with the hybrid index, meaning that they are
both minimum for a hybrid index closed to 0.5 (Fig. S4 A&B).

202
203 **Figure 2. A.** The distribution of nucleotide diversity (π) throughout the complete distribution range of
204 the two species. **B.**, and **C.** depict population-wise heterozygosity and average and standard deviation
205 in allele frequencies across SNPs for *P. obovata*, *P. abies,* or hybrid populations.
206

Local adaptation

We further investigated whether the observed pattern of genetic differentiation could be due to local
adaptations using *pcAdapt*, a genome scan method to detect outlier SNPs exhibiting a higher degree of
differentiation than the genetic background while controlling for population structure. We detected
1650 SNPs, clustered into specific areas of the genome (Fig. S3 A&C) suggesting that natural selection
has shaped the current genetic diversity, besides the population movements and demographic history.
In order to understand the role of environmental variation in shaping genetic diversity, we conducted a
redundancy analysis (RDA) with bioclimatic factors as predictors (Fig. S3B). After an ad-hoc selection
to remove high levels of correlation between climatic variables, we retained three climatic variables:

Bio8, the mean temperature of the wettest quarter, *Bio9*, the temperature of the driest quarter, and *Bio18*,

the precipitation of the warmest quarter. While *Bio8* mainly varies along latitudes, *Bio9* and *Bio18*

mainly reflect continentality and vary longitudinally. Together the climatic variables explained 5.4% of

the total genetic variation in the climatic partial RDA (pRDA) where co-variables were introduced to

control for population structure (Fig. S3B). Still, we detected 1536 outlier loci significantly associated

with at least one of the bioclimatic variables; out of which 298 were recognized as top outliers, based

on the overlap of a simple RDA model and the climatic model (see Capblancq & Forester, 2021 for

details). These top outliers clustered together in various regions of the genome (Fig. S3A). Outlier SNPs

showed substantial congruence with the enrichment observed in the *pcAdapt* output (82 % overlap, Fig.

S3D) indicating that a significant part of the allele frequency variation observed among populations is

driven by environmental heterogeneity. From the pRDA, we derived an *adaptive index* (see materials

and methods) that measures the match between the genotypes and their environment to capture the

spatial distribution of population-level adaptation to current climatic conditions (Fig. 3). The adaptive

index and the hybrid index were highly correlated (Spearman's rho = 0.90, S = 2156, $p < 0.001$ and rho

= 0.40, S= 13339, $p = 0.003$, for adaptive index computed along RDA1 and 2 respectively). While the

populations at both extremes of the sampling range show the highest degree of adaptation (strongly

positive or negative values), populations located within the hybrid zone exhibit intermediate values

indicating a lower sensitivity to environmental constraints (Fig. 3A & B).

We subsequently tested whether the observed pattern is primarily influenced by candidate SNPs specific

to the environmental variables included in the pRDA or if it demonstrates a broader, overarching trend.

It is worth noting that the phenomenon of adaptation across environmental gradients typically involves

a polygenic basis, wherein alterations in allele frequencies across multiple genetic loci contribute to the

observed effects. Consequently, our hypothesis anticipates that populations demonstrating heightened

adaptation will exhibit distinct allele frequencies when contrasted with the wider global dataset. For

each population, we calculated the ratio of major alleles specific to that population. For individual SNPs,

we initially identified the minor allele as the one with the lowest frequency across the entire dataset (f_{all}

< 0.5). Subsequently, we determined, for each population individually, the percentage of sites at which

the minor allele's frequency exceeded 0.5. In other words, while the allele might be minor in the broader

global context (frequency < 0.5), it assumed a major role within the focal population (i.e. $f_{all} < 0.5$ but

$f_{pop_i} > 0.5$). First, the hybrid populations showed a significantly lower proportion of major alleles than

that in *P. abies* or *P. obovata* populations (linear model, $r^2 = 0.65$, $df = 33$, $p < 0.001$, Fig. 3C and Fig.

S4C). Second, a strong positive correlation exists between the absolute adaptive index of a population

and the proportion of major alleles in that population (Spearman's *rho* = 0.67, S= 2532, $p < 0.001$, also

see Fig. 3E); the same relationship but weaker is observed when computing the adaptive index along

RDA2 (Spearman's *rho* = 0.33, S= 2502, $p = 0.05$). The reduced adaptiveness observed in hybrid

populations is not solely attributed to markers associated with the environmental variables considered

in the pRDA analysis. Instead, it is indicative of widespread shifts in allele frequencies throughout their

genome. Additionally, *P. obovata* populations, characterized by a greater number of population-specific major alleles, tend to exhibit higher genetic diversity. Therefore, it is probable that the observed pattern is a result of selection pressures rather than genetic drift, especially in populations that may be somewhat isolated.

Figure 3. The spatial distribution of population fitness in Norway and Siberian spruces – the adaptive index calculated from RDA1 & B – RDA2. The negative and positive adaptive index scores depict the adaptedness across continuous spectrum of extreme climatic conditions (Capblancq et al. 2021). **C.** Distribution of the adaptive index between *P. abies, P. obovata,* and their hybrids. **D.** Distribution of the proportion of population-specific major alleles between *P. abies, P. obovata* and their hybrids **E.** Adaptive index as a function of the proportion of population-specific major alleles gradient represents the hybrid index for each group as in **C**.

Environmental preference, historical niche movement, and genetic refugia

Using the ordination of climatic ranges of each species from the occurrence data, we defined the niche envelop of the two species to contextualize the niche optima of the two species. The range of environmental preferences varied substantially between the two species albeit with visible overlap (CEG: non-parametric environmental gradient Fig. 4A). Interestingly, the niches were in our environment niches two species, suggesting that marginal populations of both species are adapted to similar environmental conditions (Fig. 4B). Further, when hybrid populations are considered in the analysis, the overlap becomes wider, suggesting niche expansion owing to hybridization (Fig. 4C).

Figure 4. Environmental preference of Norway spruce and Siberian spruce; A. Collective environmental gradient (following Karunarathne et al. 2018): the sum of transformed environmental variables (i.e., Bio8, Bio9, and Bio18) for all populations plotted against Kernel density; B&C. Niche optima of the two species (following Bronnimann et al. 2016).

To shed light on spatial variation in nucleotide diversity, we used niche modeling to infer species' suitable habitat during LGM (~25,000 years ago) and mid-Holocene (~4,000-10,000 years ago Fig. 5). It confirmed that the southern range of *P. abies* (e.g., Alpine, Carpathian, and western European forests) harbored suitable habitat and must have served as refugia during the last glacial maximum. With the warming climate, the species migrated back to higher latitudes, eventually establishing its present distribution. Niche modeling for the period before the Last glacial maximum, specifically for the interglacial period (i.e., ~140,000 years ago), shows that optimal habitats for *P. abies* existed in the northernmost regions of the current distribution (e.g., northern Sweden and Norway) as well as in the Alpine region. The distribution of Norway spruce thus followed the movement of its niche during cycles of glaciation which drastically affected population connectivity. In contrast, the alteration in suitable habitats for *P. obovata* was considerably more modest across glacial cycles. This species maintained a broader distribution compared to *P. abies*, with an almost uninterrupted range. Notably, the suitable habitat ranges of both species frequently intersected over the course of the glacial cycle, with a notable convergence observed in the western Russian forests during LGM. This overlap could have facilitated multiple instances of hybridization and genetic exchange between the two species. This phenomenon mirrors the current distribution scenario where a modest spatial overlap of the suitable ecological niche is evident in North European Russia (*sensu* TDWG classification) (Fig. 5).

Figure 5. Distribution of suitable habitat/niche for *P. abies* and *P. obovata* from the last interglacial period to future climatic conditions, showing the niche movement from the past to present, determined by maxent modeling of the niche optima.

Genetic offset and resilience to climate change

Lastly, we explored the capacity of present-day populations of both species to respond to climate change. To achieve this, we quantified population adaptability to future climates by assessing the discrepancy in allele frequencies between current and projected values based on anticipated climatic conditions, referred to as the "genetic offset." Risk of non-adaptedness (RONA: Restalb et al. 2016) measures the allele frequency change necessary for future climate based on a linear regression of current values. We calculated RONA for three bioclimatic variables using climate projections for the years 2080 and 2100 (i.e., Bio8, Bio9, Bio18). Overall, the RONA assessment indicates a relatively low risk of non-adaptedness to all three bioclimatic variables in regions characterized by high genetic diversity (Fig. 6). This underscores that genetic diversity serves as a pivotal factor in mitigating the risk of non-adaptedness, thereby providing populations with a sheltered wellspring of adaptive potential. We observed high risk of non-adaptedness scores in populations from central Europe and at the Southernmost range of *P. abies* (0.18 – 0.22) indicating that populations in these regions may be at higher risk of extinction under future climatic conditions. However, this trend was not seen in populations in the Alpine region which has served as a refugium during the glacial period and also most likely in warmer conditions during interglacial periods (see Fig. 5).

Even more intriguingly, in the case of both species, the risk of non-adaptedness reaches its nadir within the hybrid zones (Fig. 6). This observation further bolsters the proposition that admixture amplifies genetic diversity, thereby fostering the broadening of ecological niche range, a phenomenon conducive to species adaptation to future climatic scenarios. Despite the fact that the Siberian spruce populations harbor the broadest genetic diversity overall (with a π range of 0.0059-0.0069), they paradoxically exhibit a greater susceptibility to non-adaptedness compared to the hybrid populations. Finally, we integrated insights from the genotype-environment-association study with those of the RONA analysis; about two-thirds of the SNPs detected as outliers in the pRDA are among the SNPs with the highest RONA values. This convergence signifies that the SNPs responsible for local adaptation to current climatic conditions are potentially detrimental in the context of long-term response to climate change.

Figure 6. Top row: The map of RONA (Risk of non-adaptedness) for Norway spruce and Siberian spruce. RONA was assessed according to Rellstab et al. 2016, using the coefficient of variation from the linear regression of present environmental variables and allele frequencies. Middle row: Predicted linear response of top 10 alleles to each climate variable. Bottom row: Range of RONA for each bioclimatic variable among the two species and the hybrid populations.

Discussion

Extensive evidence now highlights the detrimental influence of global climate change on biodiversity at multiple scales (see Altieri & Gedan, 2015; Walther et al., 2002; Williams et al., 2021). Population-level genetic variation equips species with the capacity to navigate shifting climates, safeguarding them from extinction (Meester et al., 2018). This is often observed through species' adaptations to their local environment and reflects their potential for future climate change resilience (Sang et al., 2022). Nevertheless, identifying these environment-linked alleles necessitates extensive whole-genome sequencing of multiple populations and/or long-running common garden experiments. Nonetheless, this approach proves impractical for species characterized by extended generation times, frequently overlooking essential evolutionary parameters such as demographic history and the extent of gene flow. In the present study, using exome capture sequences, we investigate the role of species hybridization in shaping the current ecological niches of Norway and Siberian spruce. Our findings reveal that recurrent admixture has heightened the climate change resilience and adaptive capacity of both species. Additionally, this approach has enabled us to pinpoint populations vulnerable to prevailing climate change conditions.

Niche modeling suggests recurrent shifts from Southern to Northern distribution ranges

Our analysis of niche optima movement of Norway spruce during and post-LGM agrees with fossil records and ancient DNA analyses (Brewer et al., 2017; Nota et al., 2022; Parducci et al., 2012;

Tollefsrud et al., 2015). Tollefsrud et al. (2015) traced the optimal ecological niches of Norway spruce and Siberian spruce during the most recent glaciation cycle. They also delineated the distribution ranges of these species during the LGM and the mid-Holocene. The study's findings showcased the gradual migration of both species to and from their southern refugia. This is of course expected since many species in temperate regions followed the same path during the last glaciation cycle (Feliner, 2011; Schönswetter et al., 2005; Sommer & Nadachowski, 2006). However, inferring *P. abies* and *P. obovata* ecological niches further back in time (~140K years ago), a period where macrofossils fail to provide species-specific information, unveiled a substantial overlap in the distribution ranges of both species in their northern habitats. This convergence could have paved the way for multiple instances of admixture. This is highly consistent with Chen et al. (2019) and Zhou et al. (2023) estimates of a massive admixture event ~103 - 187 kya between *P. abies* and *P. obovata* populations using genome-wide SNPs. Our niche modeling approach further reveals that the Southern European Mountain ranges, including the Alps and the Carpathians, likely acted as refugia for *P. abies* during both glacial and interglacial periods. Meanwhile, the species' primary distribution underwent a transition from southern to northern latitudes in tandem with the shifts in glaciation cycles. The recurrent utilization of the same discrete and well-defined refugia across ice-cycles may, at least partially, explain the difference observed between Norway spruce and two other boreal tree species - *Pinus sylvestris* (Scots pine) and *Betula pendula* (Silver birch) – both of which display notably less population structure (Bruxaux et al., 2023; Milesi et al., 2023; Tsuda et al., 2017). The latter may have instead survived the glacial periods in a large metapopulation south of the glaciated regions (Hewitt, 2000; Palmé et al., 2003).

Genetic diversity does not reflect niche-centrality but captures distribution range variations

Niche centrality suggests that a species' performance is best at its optimal niche (also niche centroid) conditions (see Lira-Noriega & Manthey, 2014). Although the niche center of a species may or may not be the geographical center of the species, it hypothesizes that the genetic diversity in the niche center is higher than in the peripheral populations (Manel & Holderegger, 2013; Pironon et al., 2015). Our results, however, show little congruence with niche centrality expectations. For *P. abies*, we recognized southern Scandinavia as the niche centroid (Fig. 4: see the populations in the center of eigen ellipses) and populations in this region show the highest nucleotide diversity (Fig. 2A). It is, however, imperative to mention that the currently observed high genetic diversity of *P. abies* in the southern Scandinavian forests is also a result of recent introductions of genetic material from the whole range of the species (Chen et al., 2019 and reference therein). For *P. obovata*, our sampling does not represent the complete current distribution of the species, especially east of the Yenisei River. For the western range of the species, the highest genetic diversity is found along the Yenisei River and the niche centroid points towards the northern Ural Mountains. While acquiring a more continuous sampling would enhance the informativeness of our conclusions, various other factors could also account for the absence of support for niche centrality. First, both Norway and Siberian spruces, as boreal species with intricate

demographic histories, experienced shifts in their suitable habitats and ecological niches across multiple

ice-age cycles. For *P. abies*, populations were even extirpated during the LGM from what represents

today the niche centroid. Additionally, the long generation time and long-range pollen dispersal of

coniferous species could also contribute to the lack of fit to the Niche centrality expectations in these

species, as is observed in *P. sylvestris* across its whole distribution range (Bruxaux et al. 2023). Our

findings align with recent research that observed minimal variation in estimates of genetic diversity

among populations from various dominant forest tree species in Western Europe (James et al., 2023;

Milesi et al., 2023).

Recurrent hybridization enlarged the ecological niche breadth

Our niche modeling strategy and retrospective projections offer additional ecological validation for the

influence of hybridization on shaping the genetic diversity of both species. Our results are in agreement

with genetic data, demographic modeling (Zhou et al. 2023), and paleoecological records (Semerikov

et al., 2013 and references therein), which suggest Siberia's state during the last glacial cycle, and

possibly more ancient glacial cycles, as a dry desert interspersed with pockets of forested areas. In

agreement with these data, our niche modeling results (Fig. 5) indicate that the suitable habitat of *P.*

obovata remained notably larger than that of *P. abies* during the late Pleistocene and Holocene periods.

Furthermore, it can be seen in Fig. 5 that the suitable habitat for both species overlapped in many

instances during the Pleistocene and Holocene, in particular in their northern range, representing

opportunities for genetic introgression. Contrary to limited gene flow, Zhou et al. (2023) demonstrated

that secondary contacts among distinct genetic entities of the two species amplified local genetic

diversity. This phenomenon is evident in populations situated within present contact zones or within

the expansive hybrid zone between *P. abies* and *P. obovata*, which displays high π values compared to

neighboring populations (Fig. 2). More importantly, Zhou et al. (2023) highlighted that the interactions

between *P. abies* and *P. obovata* led to the emergence of novel genetic clusters with admixed

characteristics. These clusters have endured through to the present day, persisting within cryptic refugia

during glacial periods. Further, they proposed that during the expansion phases, these entities helped to

bridge *P. abies* and *P. obovata* genetic backgrounds, serving as stepping stones (see also Tsuda et al.,

2016).

The role of hybridization in allowing niche expansion has been well documented in many cases of

invasive species (e.g., Pfennig et al., 2016 and reference therein), particularly in the presence of long-

distance gene-flow (e.g., Berthouly-Salazar et al., 2013) that allows for the maintenance of genetic

diversity in peripheral populations and for the introgression of favorable alleles from other populations.

Hybridization can also directly contribute to favorable traits (e.g., Darwin's finches' beak shape, Grant

& Grant, 2019) and foster their evolution through an increase in standing genetic variation, while

adaptation from *de novo* mutations can be slow (Rieseberg, 1997; Taylor et al., 2015); especially in

species with long generation time like spruces.

In Norway and Siberian spruce, primary phenotypic traits related to growth phenology or reproduction

exhibit a continuous spectrum of variation between the two species. Hybrid individuals, in this context,

manifest intermediate phenotypes (Lagercrantz & Ryman, 1990; Orlova et al., 2020; Popov, 2010).

Categorized by their phenotypes, the hybrid populations can be distinguished into two forms, each

exhibiting "abies" or "obovata" characteristics (Nakvasina et al., 2019). These distributions align with

the delineation of the hybrid zone as outlined by Zhou et al. (2023) and precisely coincide with the

distribution of the hybrid index generated in our study (see Fig. 1A). Provenance tests (i.e., breeding

experiments equivalent to common garden experiment) also revealed that the genotypes of hybrid

nature had higher survival rates in the hybrid zone (Nakvasina et al., 2019). Consistent with these

findings, our study demonstrates that the hybrid populations do not necessarily fall within the ecological

niches of either species. Instead, they carve out a niche of their own that partially intersects with the

ecological preferences of both species (see Fig. 4). Therefore, it is likely that hybridization is

responsible for expanding the ecological niche breadth of both species.

Integrating niche modeling, population genetics, and genetic-offset analysis

Given the current pace of global climate change, the central question is, will long-lived organisms such

as spruces have enough time to adapt to the new environmental conditions before facing an inevitable

demographic collapse? First and foremost, our niche modeling to future scenarios suggests that suitable

habitats would be available for both species in their current northern distribution range and our genetic

offset analysis indicates that these populations should be poised to counter climate change (as indicated

by a low risk of non-adaptedness, see Fig. 5 and 6). However, it's worth noting that a considerable

portion of the current range of *P. abies*, as well as a substantial segment of the western distribution

range of *P. obovata*, might not be suitable for their continued survival.

The populations situated at the opposite ends of the adaptive index, as indicated by the genetic offset

analysis, correspondingly represent those populations most susceptible to future climate conditions

(Fig. 3A & 6). This observation aligns with the expectation that populations displaying a pronounced

degree of adaptedness can be regarded as "specialists" tailored to their specific environments. However,

according to our modeling of the future distribution ranges, regions with populations showing an

elevated risk of non-adaptedness are projected to accommodate suitable habitats for species persistence

(Fig. 5). At first glance, these two results may seem contradictory. The essence of the RONA analysis

lies in gauging the average change in allele frequency of SNPs associated with key climatic variables.

This change is required to maintain an equivalent level of adaptedness under future climatic scenarios.

This suggests that the alleles that are responsible for local adaptation to the current climate might

potentially be deleterious under future environmental conditions, and consequently, they could be at

risk of being lost. However, RONA, like many other genetic off-set prediction models, suffers from

well-identified limitations, including its failure to account for demography history, population structure,

or gene flow (Gain et al., 2023; Rellstab et al., 2021).

Both *P. abies* and *P. obovata* exhibit extensive effective population sizes alongside far-reaching pollen

dispersal (Zhou et al., 2023). This characteristic could potentially aid in the dissemination of adaptive

alleles, particularly considering the distinct ecological preferences of the two species (see Fig. 4). Our

study has further revealed that hybrid populations occupy their own ecological niche while having a

unique genetic makeup. Moreover, these hybrid populations could serve as potential sources for future

recolonizations, in addition to acting as a reservoir of genetic diversity. For instance, hybrid populations

display the highest survival rate within the hybrid zone (Nakvasina et al., 2019). Both species hold

significant economic importance, and human-assisted transfer of trees from different provenances could

play a role in bolstering adaptation to future conditions. For instance, *P. abies* trees of Southern origin

planted in Southern Scandinavia for breeding purposes (J. Chen et al., 2019) exhibited higher growth

rates in Southern Sweden compared to trees of local origin (Milesi et al., 2019). Intriguingly, Southern

Sweden presently showcases the highest genetic diversity in *P. abies* (Fig. 2A). Consequently, these

trees can potentially introgress with local counterparts leading to the introduction of favorable alleles

for future climatic conditions. Notably, the plants' response to global change can be rapid, even in long-

lived species, like oak (*Quercus petraea*), which despite its ~50-year generation time, displayed a

genome-wide response to the Little Ice Age (~1650 AD, Saleh et al., 2022).

Finally, while neither of the studied spruce species might face a large risk of global extinction under

future climatic conditions, it is worth noting that certain areas are under particular threat. Among

populations with the highest risk of non-adaptedness, two are located south of the Italian Alps and one,

Indigo, in the Northern range of *P. obovata* western domain. As highlighted by Zhou et al. (2023), these

populations are geographically isolated from the rest, exhibit low nucleotide diversity, and have

undergone more pronounced genetic drift. These characteristics often point to populations that might

struggle to contend with the challenges posed by global climate change.

Conclusion and Perspective

When species are unable to track their ecological niche, standing genetic variation is the main

determinant for species' ability to cope with climate change. This is even more true for species with

long-generation time where the probability of adaptation through *de novo* mutation is low. Cycles of

range contractions and expansions can enhance genetic diversity in a feedback loop (Barrett & Schluter,

2008; Sexton et al., 2009) albeit with expected diversity loss during contraction events (Pauls et al.,

2013). In Western Europe, species-wise genetic diversity of seven forest tree species globally increased

across multiple ice-cycles (Milesi et al., 2023). Among them, at least five (*Betula pendula*, *Quercus*

petraea, *Fagus sylvatica*, *Populus nigra,* and *P. abies*) are known to hybridize with closely related

species and are all long-lived species with long-range gene flow. Hybridization could thus be a main

factor in coping with climate change in temperate and boreal forest tree species, as our study suggests
in the *P.abies* / *P. obovata* species complex. There is a burst of new methodology development aiming
at integrating niche modeling and genetic offset prediction in order to better apprehend species response
to global change (e.g., Barratt et al., 2023; Y. Chen et al., 2022). If this clearly represents a step forward,
our study also evidenced the need for a more integrative framework to also consider species
demography history, population structure, and gene flow.
508

Materials and methods
Sampling, genetic material, and SNP calling

The populations come from a large-scale study uncovering the demography history and spatial
population structure of the closely related Norway spruce (*Picea abies*) and Siberian spruce (*Picea*
obovata) species (Zhou et al., 2023). DNA was extracted from the needles of 546 individual trees
coming from 55 populations representing five predetermined geographic groups (including introgress
individuals and putative hybrid populations: following (J. Chen et al., 2019; L. Li et al., 2022;
Nakvasina et al., 2019); see map Fig. 1) and genomic DNA was extracted for NextGen sequencing
using Qiagen® plant DNAeasy kit. Paired-end Illumina libraries were constructed and sequenced by
RAPiD Genomic, USA, using 40,018 exome capture probes targeting 26,219 genes (Vidalis et al.,
2018). Raw reads can be found on SRA (https://www.ncbi.nlm.nih.gov/sra) under PRJNA511374 and
PRJNA1007582.

Clean sequence reads were aligned to the *P. abies* reference genome (Nystedt et al., 2013) with BWA
(H. Li & Durbin, 2009), and individual genotypes were identified using GATK HaplotypeCaller (Van
der Auwera & O'Connor, 2020) followed by SNP calling across all samples with GenotypeGVCFs and
hard-filtering of SNPs (QD < 2.0, MQ < 40.0, SOR > 3.0, QUAL < 20.0, MQRankSum < -12.5,
ReadPosRankSum < -8.0, and FS > 60.0). For population genetics analyses, we retained only putatively
neutral SNPs (i.e. synonymous mutations, mutations located in the introns or into the intergenic space)
and pruned the dataset for singleton and SNPs in high linkage disequilibrium (plink v1.9, www.cog-
genomics.org/plink/1.9/, pairwise $r^2 > 0.2$, window size 1 kb), resulting in a 390,903 SNPs data set. For
all other analyses, we kept all SNPs with minor allele frequency higher than 0.05 including both neutral
and non-neutral alleles (450,000 SNPs).
531

Climatic data selection

Environmental variables for current conditions were downloaded from the WorldClim 2.0 database at
30 arc seconds resolution (Fick & Hijmans, 2017; Hijmans et al., 2005). They are annual averages of
both temperature and precipitation-related variables. Climatic records at population locations were
extracted with the TERRA R package (Hijmans et al., 2022). In order to avoid multi-collinearity and
model overfitting, we performed pairwise Pearson-correlation between all the bioclimatic variables,
sequentially removing the variable with the highest correlation coefficient, *r*, until it reached $r_{max} = 0.6$.

Additionally, to ensure that multi-collinearity is minimal in the composite variables of RDA, we
assessed the variance inflation factor (VIF) and retained the environmental variables for which the VIF
was lower than five, as recommended by Capblancq & Forester (2021). We used this final set of
environmental variables (predictor variables hereafter) for all downstream analyses.

Hybrid index
We calculated a *hybrid index* using the ADMIXTURE software (v.1.3.0 Alexander et al. 2009)
assuming $K = 2$ ancestral populations. The hybrid index was defined as being the proportion of *P. abies*
ancestry component and ranges from 0 "pure" *P. obovata* to 1 "pure" *P. abies*. The population-wise
hybrid index was obtained by averaging individual *hybrid index.* Populations with a *hybrid index* higher
than 0.2 and below 0.8 were classified as hybrid populations in qualitative analysis. Extrapolation of
the *hybrid index* to the whole distribution of the two species was done using custom-R scripts (available
in the [GitHub](#) repository) with nearest neighbor interpolation and was mapped to visualize the spatial
distribution of genetic diversity of the two species.

We calculated global nucleotide diversity (π: Nei and Li, 1979) to assess population-level genetic
diversity with the POPGENOME R package (Pfeifer et al., 2014). Extrapolation of the π values to the
whole distribution of the two species was done using the same approach as for the *hybrid index.*
We then investigated the relationship between the *hybrid index* and various metrics of genetic diversity
and variation in allele frequencies. In order to minimize the skewed allele frequencies due to the varying
number of samples in each population, we (i) randomly sampled ten individuals per population for
populations with more than ten individuals and (ii) grouped close-by populations that were also
genetically similar before downsampling to ten individuals. The resulting dataset consisted of thirty-six
populations.

Genetically informed climatic group definition
In order to find genetically informed "climatic groups" of both species, we combined two sets of data:
i) adaptively enriched genetic space scores from pRDA – the RDA scores represent local adaptation in
populations derived from allele frequencies, ii) bioclimatic data – this dataset represents the
environmental variability of populations after removing of multi-collinearity. The former was used as
a proxy for population-level adaptations to local climatic conditions and the latter to explain the spatial
variation of local conditions. A *k-means* clustering was performed for the total within the sum of squares
to determine the optimal number of genetically informed climatic clines. This was performed separately
for tentative hybrid and non-hybrid populations to avoid cluster overlaps that could arise from specific
adaptations in hybrid populations. In non-hybrid populations, we identified five distinct climatic groups
while there were only three within putatively hybrid populations. After obtaining the optimal number
of clusters as five, a second k-means clustering was performed with K=5 for all populations (i.e.,
including hybrid and non-hybrid populations) to determine the climatic group membership of each

population. This step was performed specifically to group the putative hybrid populations with their

closest non-hybrid populations to identify their closest climatic groups.

Identification of alleles putatively involved in local adaptation

To study the genetic architecture of local adaptation in Norway spruce and Siberian spruce, we ran

genome scans to detect outlier-loci (i.e., loci showing an excess of genetic differentiation in comparison

to the rest of the genetic background) using the principal component-based outlier detection method,

"pcadapt", implemented in the R package "pcadapt" v.4.3.2 (Luu et al., 2017; Privé et al., 2020).

Briefly, a Z-score matrix, obtained from regressing all SNPs to PCs, is used to produce one multivariate

distance for each variant. Assuming that SNP variation along principal component axes largely reflects

demographic processes, this method allows controlling for population structure when testing for local

adaptation. In our analysis, to control for false positive risk discovery associated with multiple testing,

we used FDR q-value < 0.1 to obtain the significant SNPs putatively involved in local adaptation.

Redundancy Analysis (RDA) is an asymmetric canonical analysis that combines ordination and

regression, where multiple regression on the response variable (i.e., allele frequencies) and explanatory

variables (climatic variables) is followed by ordination of fitted values of the regression (Van Den

Wollenberg 1977). We used Redundancy Analysis (RDA) to find the main climatic variables that

explain most of the variance in allele frequencies between the two species and to identify candidate loci

associated with environmental variation.

In the special case of partial RDA (pRDA), additional explanatory variables (covariates) are used to

control for various confounding effects (see Capblancq et al., 2018; Capblancq & Forester, 2021). This

aspect of RDA is extremely useful in association analyses to control for the confounding effect of

factors that are not the focus of the study, such as population structure and geographic distance. Thus,

to isolate different drivers of genetic variation, in addition to a full model (including all predictor

variables), we performed three independent pRDA models controlling for geographic distance,

population structure, and environmental conditions. We adopted the method described in Capblancq &

Forester (2021) where the Mahalanobis distance of each locus to the center of the RDA is calculated

using two axes (K=2) of partial RDA (i.e., the number of axes that explain more than 60% variation;

following Capblancq et al. 2021), followed by obtaining the top outlier SNPs from this distance matrix.

For geographic distance, the cartesian distance of populations was used as a proxy; to control for

population structure, we used the first three axes of a PCA performed on the set of putatively neutral

loci.

Using the output from the pRDA, we computed an *Adaptive index (Aij)* as implemented by Capblancq

& Forester (2021) from Steane et al. (2014) using the RDA 1 and 2 independently. For the top outliers,

RDA1 and RDA2 explained 88% and 10% of the variation respectively. For each location, j, the

adaptive index is given by the sum over each climatic variable, *i*, of the products between its score along

a given RDA axis and its value at location *j*.

$$A_{ij} = \sum_{i=1}^{n} a_i b_{ij}$$

A_{ij} ranges from negative to positive values and essentially measures the match between the genotypes

and their environment, 0 meaning no match.

Niche expansion, species distribution modeling, and historical niche optima movement analysis

The climatic niche of the two species was defined as the ordination space of all the populations with

PCA eigenvalues. Therein, a PCA was performed on the selected climatic variables (i.e., Bio8, Bio9,

Bio18) for all the populations of *P. abies* and *P. obovata* separately. The optimal niche was determined

as the centroid of the eigenvalues and the niche breath was defined as the 95% ellipsoid of the PCA

scorings. The analysis was performed with and without hybrid populations to visualize the niche breadth

overlap. As a complementary analysis, we also used a non-parametric method, collective environmental

gradient (CEG; Karunarathne et al. 2018) to assess the total environmental aptitude of each species and

to visualize their overlap. CEG is a single-value decomposition approach to determine the composite

two-dimensional environmental/niche breach. We performed CEG according to Karunarathne et al.,

(2018) for the selected bioclimatic variables (i.e., Bio8, Bio9, Bio18).

Species distribution models for the current, mid-Holocene (ca. 6,000 ya), Last Glacial Maximum (LGM,

ca. 22,000 ya), Last Inter-glacial period (ca. 120,000 - 140,000 ya), and future climatic scenarios (2021-

2080 and 2081-2100) were constructed using maximum entropy (Maxent) modeling (Phillips et al.,

2006). The optimal niche for separate species was determined with selected climatic variables (i.e.,

Bio8, Bio9, Bio18) using the *maxent* function of the *dismo* R package (Hijmans et al., 2016). Maxent

uses presence-only data to estimate the probability distribution of occurrence based on environmental

constraints (Phillips et al., 2006). In order to test the models and to facilitate a more accurate prediction

of species distribution, we downloaded species occurrence data for *P. abies* and *P. obovata* from GBIF

(www.GBIF.org) and filtered the data to include only confirmed and natural occurrences of the species.

occurrence data were then subsampled to retain only one occurrence per grid cell of 1 arc minute. The

final dataset consisted of 619 occurrences (*P. abies*: 288; *P. obovata*: 331), hybrid populations have

been excluded. All bioclimatic variables were downloaded from WorlClim 1.4. database at 30 arc-

second resolution. The climatic variables for the Mid-Holocene, LGM, and Last inter-glacial were

downloaded from the model CCSM4 and the future climate variables were downloaded for ACCESS-

CM2 model. The projection of past and future niche distribution of the two species was done using the

predict function of the *dismo* R package. All the models were evaluated using the area under the

operating curve (AUC) and only the models with an AUC of at least 0.85 were considered.

Assessment of genetic offset and resilience to future climatic conditions

The genetic offset of an entity (e.g., population) is the allele frequency difference between the current and required frequency in the future to survive in future climatic scenarios. We calculated the genetic offset of our populations with risk of nonadaptedness (RONA) as in Rellstab et al. (2016). RONA is the average theoretical change in allele frequency at climate-associated loci required to match future climatic scenarios. This is established by regressing the selected loci allele frequencies with the selected environmental variables and using the regression coefficients to predict the expected allele frequencies of the respective loci to future conditions. The difference between the two allele frequencies is the risk of nonadaptedness. Here, we first performed linear regression on all loci with the selected environmental variables (i.e., Bio8, Bio9, Bio18) and kept only the loci that showed significant association (p-value < 0.05) with the three environmental variables. Then we obtained the intersection between these alleles and the top outliers from the RDA analysis for the RONA analysis. To avoid linkage between selected loci, we filtered loci within 10,000 base pairs in the same scaffold; the locus with the highest R^2 from the regression was retained. In total, ~25,000 candidate SNPs for environmental associations were kept for the analysis. Finally, we used nearest neighbor interpolation (as in Rukundo & Cao 2012: custom scripts available in the GitHub repository) to predict the RONA values for the entire distribution of the two species for spatial visualization (Fig. 6).

Acknowledgments

We would like to thank Elena Nakvasina, Vladimir Semerikov, Lars Opgenoorth, Katrin Heer, and Giovanni Giuseppe Vendramin for their valuable efforts in providing us with plant materials. We also thank the Swedish National Infrastructure for Computing (SNIC) for resource allocation for high-power computing under the project numbers SNIC 2021/6-313 and SNIC 2021/5-540. The Nilsson-Ehle grant by the Royal Physiographic Society of Lund awarded to Piyal Karunarathne (42191) was immensely helpful for acquiring genome sequences to fill data gaps.

Data availability

All the raw reads studied in this study can be found on the NCBI SRA database (https://www.ncbi.nlm.nih.gov/sra) under PRJNA511374 and PRJNA1007582. The scripts used for all the analyses, simulations and respective input data are available in the GitHub repository.

Conflict of interest statement

The authors declare no conflict of interest.

References

Altieri, A. H., & Gedan, K. B. (2015). Climate change and dead zones. *Global Change Biology*, *21*(4), 1395–1406. https://doi.org/https://doi.org/10.1111/gcb.12754

Araújo, M. B., & Rahbek, C. (2006). How Does Climate Change Affect Biodiversity? *Science*, *313*(5792), 1396–1397. https://doi.org/10.1126/science.1131758

Atkins, K. E., & Travis, J. M. J. (2010). Local adaptation and the evolution of species' ranges under climate change. *Journal of Theoretical Biology*, *266*(3), 449–457. https://doi.org/https://doi.org/10.1016/j.jtbi.2010.07.014

Barratt, C. D., Onstein, R. E., Pinsky, M. L., Steinfartz, S., Kühl, H. S., Forester, B. R., & Razgour, O.
(2023). Life on the edge: a new toolbox for population-level climate change vulnerability
assessments. *BioRxiv*, 2023.06.23.543988. https://doi.org/10.1101/2023.06.23.543988
Barrett, R. D. H., & Schluter, D. (2008). Adaptation from standing genetic variation. *Trends in
Ecology & Evolution*, *23*(1), 38–44.
Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate
change on the future of biodiversity. *Ecology Letters*, *15*(4), 365–377.
https://doi.org/https://doi.org/10.1111/j.1461-0248.2011.01736.x
Berthouly-Salazar, C., Hui, C., Blackburn, T. M., Gaboriaud, C., Van Rensburg, B. J., Van Vuuren,
B. J., & Le Roux, J. J. (2013). Long-distance dispersal maximizes evolutionary potential during
rapid geographic range expansion. *Molecular Ecology*, *22*(23), 5793–5804.
Bonnet, T., Morrissey, M. B., de Villemereuil, P., Alberts, S. C., Arcese, P., Bailey, L. D., Boutin, S.,
Brekke, P., Brent, L. J. N., Camenisch, G., Charmantier, A., Clutton-Brock, T. H., Cockburn, A.,
Coltman, D. W., Courtiol, A., Davidian, E., Evans, S. R., Ewen, J. G., Festa-Bianchet, M., …
Kruuk, L. E. B. (2022). Genetic variance in fitness indicates rapid contemporary adaptive
evolution in wild animals. *Science*, *376*(6596), 1012–1016.
https://doi.org/10.1126/science.abk0853
Brewer, S., Giesecke, T., Davis, B. A. S., Finsinger, W., Wolters, S., Binney, H., de Beaulieu, J.-L.,
Fyfe, R., Gil-Romera, G., & Kühl, N. (2017). Late-glacial and Holocene European pollen data.
Journal of Maps, *13*(2), 921–928.
Bruxaux, J., Zhao, W., Hall, D., Curtu, A. L., Androsiuk, P., Drouzas, A. D., Gailing, O., Konrad, H.,
Sullivan, A. R., Semerikov, V., & Wang, X.-R. (2023). Low genetic differentiation across an
entire continent: the Scots pine paradox. *BioRxiv*, 2023.06.09.543371.
https://doi.org/10.1101/2023.06.09.543371
Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., McEvey, S., & Ferrier, S.
(2016). Incorporating evolutionary adaptation in species distribution modelling reduces
projected vulnerability to climate change. *Ecology Letters*, *19*(12), 1468–1478.
https://doi.org/10.1111/ele.12696
Capblancq, T., & Forester, B. R. (2021). Redundancy analysis: A Swiss Army Knife for landscape
genomics. *Methods in Ecology and Evolution*, *12*(12), 2298–2309.
Capblancq, T., Luu, K., Blum, M. G. B., & Bazin, E. (2018). Evaluation of redundancy analysis to
identify signatures of local adaptation. *Molecular Ecology Resources*, *18*(6), 1223–1233.
Carbeck, K., Wang, T., Reid, J. M., & Arcese, P. (2022). Adaptation to climate change through
seasonal migration revealed by climatic versus demographic niche models. *Global Change
Biology*, *28*(14), 4260–4275. https://doi.org/https://doi.org/10.1111/gcb.16185
Catullo, R. A., Llewelyn, J., Phillips, B. L., & Moritz, C. C. (2019). The Potential for Rapid Evolution
under Anthropogenic Climate Change. *Current Biology*, *29*(19), R996–R1007.
https://doi.org/https://doi.org/10.1016/j.cub.2019.08.028
Chen, J., Li, L., Milesi, P., Jansson, G., Berlin, M., Karlsson, B., Aleksic, J., Vendramin, G. G., &
Lascoux, M. (2019). Genomic data provide new insights on the demographic history and the
extent of recent material transfers in Norway spruce. *Evolutionary Applications*, *12*(8), 1539–
1551. https://doi.org/10.1111/eva.12801
Chen, Y., Jiang, Z., Fan, P., Ericson, P. G. P., Song, G., Luo, X., Lei, F., & Qu, Y. (2022). The
combination of genomic offset and niche modelling provides insights into climate change-driven
vulnerability. *Nature Communications*, *13*(1), 4821. https://doi.org/10.1038/s41467-022-32546-
z
Cotto, O., Wessely, J., Georges, D., Klonner, G., Schmid, M., Dullinger, S., Thuiller, W., &
Guillaume, F. (2017). A dynamic eco-evolutionary model predicts slow response of alpine
plants to climate warming. *Nature Communications*, *8*(1), 15399.
https://doi.org/10.1038/ncomms15399
Dauphin, B., Rellstab, C., Schmid, M., Zoller, S., Karger, D. N., Brodbeck, S., Guillaume, F., &
Gugerli, F. (2021). Genomic vulnerability to rapid climate warming in a tree species with a long
generation time. *Global Change Biology*, *27*(6), 1181–1195.
https://doi.org/https://doi.org/10.1111/gcb.15469

Feliner, G. N. (2011). Southern European glacial refugia: A tale of tales. *TAXON*, *60*(2), 365–372. https://doi.org/https://doi.org/10.1002/tax.602007

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. https://doi.org/https://doi.org/10.1002/joc.5086

Frankham, R. (2005). Stress and adaptation in conservation genetics. *Journal of Evolutionary Biology*, *18*(4), 750–755.

Gain, C., Rhoné, B., Cubry, P., Salazar, I., Forbes, F., Vigouroux, Y., Jay, F., & François, O. (2023). A Quantitative Theory for Genomic Offset Statistics. *Molecular Biology and Evolution*, *40*(6), msad140. https://doi.org/10.1093/molbev/msad140

Giesecke, T. (2005). Moving front or population expansion: How did Picea abies (L.) Karst. become frequent in central Sweden? *Quaternary Science Reviews*, *24*(23–24), 2495–2509. https://doi.org/10.1016/j.quascirev.2005.03.002

Gillies, S., Clements, D. R., & Grenz, J. (2016). Knotweed (Fallopia spp.) Invasion of North America Utilizes Hybridization, Epigenetics, Seed Dispersal (Unexpectedly), and an Arsenal of Physiological Tactics. *Invasive Plant Science and Management*, *9*(1), 71–80. https://doi.org/DOI: 10.1614/IPSM-D-15-00039.1

Grant, P. R., & Grant, B. R. (2019). Hybridization increases population variation during adaptive radiation. *Proceedings of the National Academy of Sciences*, *116*(46), 23216–23224. https://doi.org/10.1073/pnas.1913534116

Grimm, N. B., Chapin III, F. S., Bierwagen, B., Gonzalez, P., Groffman, P. M., Luo, Y., Melton, F., Nadelhoffer, K., Pairis, A., Raymond, P. A., Schimel, J., & Williamson, C. E. (2013). The impacts of climate change on ecosystem structure and function. *Frontiers in Ecology and the Environment*, *11*(9), 474–482. https://doi.org/https://doi.org/10.1890/120282

Hamilton, J. A., & Miller, J. M. (2016). Adaptive introgression as a resource for management and genetic conservation in a changing climate. *Conservation Biology*, *30*(1), 33–41.

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), 907–913. https://doi.org/10.1038/35016000

Hijmans, R. J., Bivand, R., Forner, K., Ooms, J., Pebesma, E., & Sumner, M. D. (2022). Package 'terra.' *Maintainer: Vienna, Austria*.

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*(15), 1965–1978. https://doi.org/10.1002/joc.1276

Hijmans, R. J., Phillips, S., Leathwick, J., & Elith, V. (2016). *dismo: Distribution Modeling. R package version 1.0-15*. https://cran.r-project.org/package=dismo

Inoue, K., & Berg, D. J. (2017). Predicting the effects of climate change on population connectivity and genetic diversity of an imperiled freshwater mussel, Cumberlandia monodonta (Bivalvia: Margaritiferidae), in riverine systems. *Global Change Biology*, *23*(1), 94–107. https://doi.org/https://doi.org/10.1111/gcb.13369

James, J., Kastally, C., Budde, K., Gonzalez-Martinez, S., Milesi, P., Pyhäjärvi, T., GenTree-Consortium, & Lascoux, M. (2023). Between but not within species variation in the distribution of fitness effects. *Molecular Biology and Evolution*.

Karunarathne, P., Schedler, M., Martínez, E. J., Honfi, A. I., Novichkova, A., & Hojsgaard, D. (2018). Intraspecific ecological niche divergence and reproductive shifts foster cytotype displacement and provide ecological opportunity to polyploids. *Annals of Botany*, *121*(6), 1183–1196. https://doi.org/10.1093/aob/mcy004

Lagercrantz, U., & Ryman, N. (1990). Genetic structure of Norway spruce (Picea abies): concordance of morphological and allozymic variation. *Evolution*, *44*(1), 38–53.

Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, *17*(4), 183–189. https://doi.org/https://doi.org/10.1016/S0169-5347(02)02497-7

Leroy, T., Louvet, J.-M., Lalanne, C., Le Provost, G., Labadie, K., Aury, J.-M., Delzon, S., Plomion, C., & Kremer, A. (2020). Adaptive introgression as a driver of local adaptation to climate in European white oaks. *New Phytologist*, *226*(4), 1171–1182. https://doi.org/https://doi.org/10.1111/nph.16095

Lexer, C., Joseph, J. A., van Loo, M., Barbará, T., Heinze, B., Bartha, D., Castiglione, S., Fay, M. F.,
& Buerkle, C. A. (2010). Genomic admixture analysis in European Populus spp. reveals
unexpected patterns of reproductive isolation and mating. *Genetics*, *186*(2), 699–712.
Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics, *25*(14), 1754–1760.
Li, L., Milesi, P., Tiret, M., Chen, J., Sendrowski, J., Baison, J., Chen, Z., Zhou, L., Karlsson, B.,
Berlin, M., Westin, J., Garcia-Gil, M. R., Wu, H. X., & Lascoux, M. (2022). Teasing apart the
joint effect of demography and natural selection in the birth of a contact zone. *New Phytologist*,
236(5), 1976–1987. https://doi.org/https://doi.org/10.1111/nph.18480
Lira-Noriega, A., & Manthey, J. D. (2014). Relationship of genetic diversity and niche centrality: a
survey and analysis. *Evolution*, *68*(4), 1082–1093. https://doi.org/10.1111/evo.12343
Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt: an R package to perform genome scans for
selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1), 67–77.
Manel, S., & Holderegger, R. (2013). Ten years of landscape genetics. *Trends in Ecology and*
Evolution, *28*(10), 614–621. https://doi.org/10.1016/j.tree.2013.05.012
Markert, J. A., Champlin, D. M., Gutjahr-Gobell, R., Grear, J. S., Kuhn, A., McGreevy, T. J., Roth,
A., Bagley, M. J., & Nacci, D. E. (2010). Population genetic diversity and fitness in multiple
environments. *BMC Evolutionary Biology*, *10*(1), 1–13.
Meester, L. De, Stoks, R., & Brans, K. I. (2018). Genetic adaptation as a biological buffer against
climate change: Potential and limitations. *Integrative Zoology*, *13*(4), 372–391.
https://doi.org/https://doi.org/10.1111/1749-4877.12298
Milesi, P., Berlin, M., Chen, J., Orsucci, M., Li, L., Jansson, G., Karlsson, B., & Lascoux, M. (2019).
Assessing the potential for assisted gene flow using past introduction of Norway spruce in
southern Sweden: Local adaptation and genetic basis of quantitative traits in trees. *Evolutionary*
Applications, *12*(10), 1946–1959. https://doi.org/https://doi.org/10.1111/eva.12855
Milesi, P., Kastally, C., Dauphin, B., Cervantes, S., Bagnoli, F., Budde, K. B., Cavers, S., Ojeda, D. I.,
Fady, B., Faivre-Rampant, P., González-Martínez, S. C., Grivet, D., Gugerli, F., Jorge, V.,
Lesur-Kupin, I., Olsson, S., Opgenoorth, L., Pinosio, S., Plomion, C., … Pyhäjärvi, T. (2023).
Synchronous effective population size changes and genetic stability of forest trees through
glacial cycles. *BioRxiv*, 2023.01.05.522822. https://doi.org/10.1101/2023.01.05.522822
Müller, M., Lopez, P. A., Papageorgiou, A. C., Tsiripidis, I., & Gailing, O. (2019). Indications of
genetic admixture in the transition zone between Fagus sylvatica L. and Fagus sylvatica ssp.
orientalis Greut. & Burd. *Diversity*, *11*(6), 90.
Nakvasina, E., Demina, N., Prozherina, N., & Demidova, N. (2019). Assessment of phenotypic
plasticity of spruce species (L.) Karst. and (Ledeb.) on provenances tests in European North of
Russia. *Central European Forestry Journal*, *65*(2), 121–128.
Nota, K., Klaminder, J., Milesi, P., Bindler, R., Nobile, A., van Steijn, T., Bertilsson, S., Svensson, B.,
Hirota, S. K., & Matsuo, A. (2022). Norway spruce postglacial recolonization of Fennoscandia.
Nature Communications, *13*(1), 1333.
Nystedt, B., Street, N. R., Wetterbom, A., Zuccolo, A., Lin, Y. C., Scofield, D. G., Vezzi, F.,
Delhomme, N., Giacomello, S., Alexeyenko, A., Vicedomini, R., Sahlin, K., Sherwood, E.,
Elfstrand, M., Gramzow, L., Holmberg, K., Hällman, J., Keech, O., Klasson, L., … Jansson, S.
(2013). The Norway spruce genome sequence and conifer genome evolution. *Nature*, *497*(7451),
579–584. https://doi.org/10.1038/nature12211
Orlova, L., Gussarova, G., Glazkova, E., Egorov, A., Potokin, A., & Ivanov, S. (2020). Systematics
and distribution of spruce species in the North-West of Russia. *Dendrobiology*, *84*, 12–29.
https://doi.org/10.12657/denbio.084.002
Ortego, J., Gugger, P. F., & Sork, V. L. (2015). Climatically stable landscapes predict patterns of
genetic structure and admixture in the Californian canyon live oak. *Journal of Biogeography*,
42(2), 328–338. https://doi.org/https://doi.org/10.1111/jbi.12419
Palmé, A. E., Su, Q., Rautenberg, A., Manni, F., & Lascoux, M. (2003). Postglacial recolonization
and cpDNA variation of silver birch, Betula pendula. *Molecular Ecology*, *12*(1), 201–212.
https://doi.org/https://doi.org/10.1046/j.1365-294X.2003.01724.x

Parducci, L., Jørgensen, T., Tollefsrud, M. M., Elverland, E., Alm, T., Fontana, S. L., Bennett, K. D.,
Haile, J., Matetovici, I., & Suyama, Y. (2012). Glacial survival of boreal trees in northern
Scandinavia. *Science*, *335*(6072), 1083–1086.
Pauls, S. U., Nowak, C., Bálint, M., & Pfenninger, M. (2013). The impact of global climate change on
genetic diversity within populations and species. *Molecular Ecology*, *22*(4), 925–946.
Petit, R. J., Aguinagalde, I., de Beaulieu, J.-L., Bittkau, C., Brewer, S., Cheddadi, R., Ennos, R.,
Fineschi, S., Grivet, D., Lascoux, M., Mohanty, A., Müller-Starck, G., Demesure-Musch, B.,
Palmé, A., Martín, J. P., Rendell, S., & Vendramin, G. G. (2003). Glacial Refugia: Hotspots But
Not Melting Pots of Genetic Diversity. *Science*, *300*(5625), 1563–1565.
https://doi.org/10.1126/science.1083264
Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S. E., & Lercher, M. J. (2014). PopGenome: an efficient
Swiss army knife for population genomic analyses in R. *Molecular Biology and Evolution*,
31(7), 1929–1936.
Pfennig, K. S., Kelly, A. L., & Pierce, A. A. (2016). Hybridization as a facilitator of species range
expansion. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1839), 20161329.
Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species
geographic distributions. *Ecological Modeling*, *190*(2006), 231–259.
https://doi.org/10.1016/j.ecolmodel.2005.03.026
Pironon, S., Villellas, J., Morris, W. F., Doak, D. F., & Garc??a, M. B. (2015). Do geographic,
climatic or historical ranges differentiate the performance of central versus peripheral
populations? *Global Ecology and Biogeography*, *24*(6), 611–620.
https://doi.org/10.1111/geb.12263
Popov, P. P. (2010). Form structure and geographic differentiation of spruce populations in
Northwestern Russia. *Russian Journal of Ecology*, *41*, 378–385.
Privé, F., Luu, K., Vilhjálmsson, B. J., & Blum, M. G. B. (2020). Performing highly efficient genome
scans for local adaptation with R package pcadapt version 4. *Molecular Biology and Evolution*,
37(7), 2153–2154.
Rellstab, C., Dauphin, B., & Exposito-Alonso, M. (2021). Prospects and limitations of genomic offset
in conservation management. *Evolutionary Applications*, *14*(5), 1202–1212.
https://doi.org/https://doi.org/10.1111/eva.13205
Rellstab, C., Zoller, S., Walthert, L., Lesur, I., Pluess, A. R., Graf, R., Bodénès, C., Sperisen, C.,
Kremer, A., & Gugerli, F. (2016). Signatures of local adaptation in candidate genes of oaks
(Quercus spp.) with respect to present and future climatic conditions. *Molecular Ecology*,
25(23), 5907–5924.
Rendón-Anaya, M., Wilson, J., Sveinsson, S., Fedorkov, A., Cottrell, J., Bailey, M. E. S., Ruņģis, D.,
Lexer, C., Jansson, S., Robinson, K. M., Street, N. R., & Ingvarsson, P. K. (2021). Adaptive
Introgression Facilitates Adaptation to High Latitudes in European Aspen (Populus tremula L.).
Molecular Biology and Evolution, *38*(11), 5034–5050. https://doi.org/10.1093/molbev/msab229
Rieseberg, L. H. (1997). Hybrid origins of plant species. *Annual Review of Ecology and Systematics*,
359–389.
Rieseberg, L. H., & Willis, J. H. (2007). Plant Speciation. *Science*, *317*(5840), 910–914.
https://doi.org/10.1126/science.1137729
Saleh, D., Chen, J., Leplé, J.-C., Leroy, T., Truffaut, L., Dencausse, B., Lalanne, C., Labadie, K.,
Lesur, I., Bert, D., Lagane, F., Morneau, F., Aury, J.-M., Plomion, C., Lascoux, M., & Kremer,
A. (2022). Genome-wide evolutionary response of European oaks during the Anthropocene.
Evolution Letters, *6*(1), 4–20. https://doi.org/10.1002/evl3.269
Sang, Y., Long, Z., Dan, X., Feng, J., Shi, T., Jia, C., Zhang, X., Lai, Q., Yang, G., & Zhang, H.
(2022). Genomic insights into local adaptation and future climate-induced vulnerability of a
keystone forest tree in East Asia. *Nature Communications*, *13*(1), 6541.
Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature*
Reviews Genetics, *14*(11), 807–820. https://doi.org/10.1038/nrg3522
Schönswetter, P., Stehlik, I., Holderegger, R., & Tribsch, A. (2005). Molecular evidence for glacial
refugia of mountain plants in the European Alps. *Molecular Ecology*, *14*(11), 3547–3555.
Semerikov, V. L., Semerikova, S. A., Polezhaeva, M. A., Kosintsev, P. A., & Lascoux, M. (2013).
Southern montane populations did not contribute to the recolonization of West Siberian Plain by

Siberian larch (*Larix sibirica*): a range-wide analysis of cytoplasmic markers. *Molecular Ecology*, *22*(19), 4958–4971. https://doi.org/10.1111/mec.12433

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of species range limits. *Annu. Rev. Ecol. Evol. Syst.*, *40*, 415–436.

Sommer, R. S., & Nadachowski, A. (2006). Glacial refugia of mammals in Europe: evidence from fossil records. *Mammal Review*, *36*(4), 251–265.

Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., & Greig, D. (2014). Hybridization facilitates evolutionary rescue. *Evolutionary Applications*, *7*(10), 1209–1217. https://doi.org/https://doi.org/10.1111/eva.12214

Taylor, S. A., Larson, E. L., & Harrison, R. G. (2015). Hybrid zones: windows on climate change. *Trends in Ecology & Evolution*, *30*(7), 398–406. https://doi.org/https://doi.org/10.1016/j.tree.2015.04.010

Teukeng, F. F. D., Blin, M., Bech, N., Gomez, M. R., Zein-Eddine, R., Simo, A. M. K., Allienne, J.-F., Tchuem-Tchuenté, L. A., & Boissier, J. (2022). Hybridization increases genetic diversity in Schistosoma haematobium populations infecting humans in Cameroon. *Infectious Diseases of Poverty*, *11*(1), 37. https://doi.org/10.1186/s40249-022-00958-0

Thörn, F., Rödin-Mörch, P., Cortazar-Chinarro, M., Richter-Boix, A., Laurila, A., & Höglund, J. (2021). The effects of drift and selection on latitudinal genetic variation in Scandinavian common toads (Bufo bufo) following postglacial recolonisation. *Heredity*, *126*(4), 656–667. https://doi.org/10.1038/s41437-020-00400-x

Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(23), 8245–8250. https://doi.org/10.1073/pnas.0409902102

Thuiller, W., Münkemüller, T., Lavergne, S., Mouillot, D., Mouquet, N., Schiffers, K., & Gravel, D. (2013). A road map for integrating eco-evolutionary processes into biodiversity models. *Ecology Letters*, *16*(s1), 94–105. https://doi.org/10.1111/ele.12104

Tollefsrud, M. M., Latałowa, M., van der Knaap, W. O., Brochmann, C., & Sperisen, C. (2015). Late Quaternary history of North Eurasian Norway spruce (*Picea abies*) and Siberian spruce (Picea obovata) inferred from macrofossils, pollen and cytoplasmic DNA variation. *Journal of Biogeography*, *42*(8), 1431–1442. https://doi.org/https://doi.org/10.1111/jbi.12484

Tsuda, Y., Chen, J., Stocks, M., Källman, T., Sønstebø, J. H., Parducci, L., Semerikov, V., Sperisen, C., Politov, D., Ronkainen, T., Väliranta, M., Vendramin, G. G., Tollefsrud, M. M., & Lascoux, M. (2016). The extent and meaning of hybridization and introgression between Siberian spruce (Picea obovata) and Norway spruce (Picea abies): cryptic refugia as stepping stones to the west? *Molecular Ecology*, *25*(12), 2773–2789. https://doi.org/https://doi.org/10.1111/mec.13654

Tsuda, Y., Semerikov, V., Sebastiani, F., Vendramin, G. G., & Lascoux, M. (2017). Multispecies genetic structure and hybridization in the Betula genus across Eurasia. *Molecular Ecology*, *26*(2), 589–605. https://doi.org/https://doi.org/10.1111/mec.13885

Van der Auwera, G. A., & O'Connor, B. D. (2020). *Genomics in the cloud: using Docker, GATK, and WDL in Terra*. O'Reilly Media.

Vidalis, A., Scofield, D. G., Neves, L. G., Bernhardsson, C., García-Gil, M. R., & Ingvarsson, P. K. (2018). Design and evaluation of a large sequence-capture probe set and associated SNPs for diploid and haploid samples of Norway spruce (Picea abies). *BioRxiv*, 291716. https://doi.org/10.1101/291716

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M., Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, *416*(6879), 389–395.

Williams, J. W., Ordonez, A., & Svenning, J.-C. (2021). A unifying framework for studying and managing climate-driven rates of ecological change. *Nature Ecology & Evolution*, *5*(1), 17–26. https://doi.org/10.1038/s41559-020-01344-5

Wuethrich, B. (2000). How Climate Change Alters Rhythms of the Wild. *Science*, *287*(5454), 793–795. https://doi.org/10.1126/science.287.5454.793

Zalapa, J. E., Brunet, J., & Guries, R. P. (2010). The extent of hybridization and its impact on the genetic diversity and population structure of an invasive tree, Ulmus pumila (Ulmaceae).

963   *Evolutionary Applications*, *3*(2), 157–168. https://doi.org/https://doi.org/10.1111/j.1752-
964   4571.2009.00106.x
965   Zhou, Q., Karunarathne, P., Andersson-Li, L., Chen, C., Opgenoorth, L., Heer, K., Vendramin, B. G.,
966   Nakvasina, E., Lascoux, M., & Milesi, P. (2023). Recurrent hybridization and gene flow shaped
967   Norway and Siberian spruce evolutionary history over multiple glacial cycles. *BioRxiv*.
968