Article summary line: The present work describes the first protocol for *Plasmodium ovale*

wallikeri microsatellite genotyping and its application to relapse characterization.

Running title: Genetic profiling of *P. ovale wallikeri* relapses

Keywords: *Plasmodium ovale wallikeri*; microsatellites; relapses; WGS; sWGA

Title: Genetic profiling of *Plasmodium ovale wallikeri* relapses with microsatellite markers

and whole-genome sequencing

Authors: Valentin Joste[1,2], Emma Colard-Itté[2], Émilie Guillochon[1], Frédéric Ariey[3], Romain

Coppée[4], Jérôme Clain[1,2] and Sandrine Houzé[1,2] on behalf of the French National Reference

Center for Imported Malaria Study Group

Author affiliations:

1. Université Paris Cité, IRD, MERIT, F-75006 Paris, France

2. Centre National de Référence du Paludisme, AP-HP, Hôpital Bichat – Claude-Bernard, F-

75018 Paris, France

3. INSERM U1016, Institut Cochin, Laboratoire de parasitologie-mycologie, Hôpital Cochin,

AP-HP, Université de Paris, Paris, France

4. Université Paris Cité & Sorbonne Paris Nord, Inserm, IAME, F-75018 Paris, France

Corresponding author:

Valentin Joste, valentinjoste@gmail.com,

phone: +33617435543

Abstract

Plasmodium ovale wallikeri and *P. ovale curtisi* were recently classified as two distinct

species. Like *P. vivax*, both *P. ovale curtisi* and *P. ovale wallikeri* have the ability to cause

relapse in humans. *P. ovale* spp. are still understudied and the development of genomic tools

to study parasite genetic diversity parameters or medically-important traits such as relapses

are needed. Here, we describe a protocol to type a set of eight putatively neutral *P. ovale*

wallikeri microsatellite markers. Expected heterozygosity ranged from 0.764 to 0.940 in 69 *P.*

ovale wallikeri samples from Sub-Saharan Africa and polyclonality was detected in 7 isolates.

We genotyped 15 pairs of isolates from primary (D_0) and relapsing infections ($D_{relapse}$) that

occurred in metropolitan France from 2013 to 2021. For most pairs, the D_0 and the $D_{relapse}$

isolates were highly genetically related, an observation that was confirmed by whole-genome

sequencing for the four pairs we further studied.

Main article

Introduction

Malaria is one of the deadliest infectious diseases, especially in Sub-Saharan Africa with

approximatively 619,000 deaths in 2021 (1). While *Plasmodium falciparum* is the best

characterized species in Africa with a well-described epidemiology (1), *P. ovale* spp. are also

endemic in Africa (2) but their exact prevalence are still imperfectly established and have

been probably largely underestimated (3). The increased use of qPCR makes the diagnosis of

non-*falciparum* or *Plasmodium* co-infections easier (4) and will help in the future to better

define the *P. ovale* spp. prevalence in endemic area.

P. ovale wallikeri and *P. ovale curtisi* were recently classified as two different species based

on distinct genetic patterns (5). These species are sympatric and display different clinical

characteristics that are well-described in imported malaria patients, such as deeper

thrombocytopenia and shorter latency period for *P. ovale wallikeri* compared to *P. ovale*

curtisi (6,7). Like *P. vivax*, both *P. ovale curtisi* and *P. ovale wallikeri* have the ability to

cause relapse (6,8), defined as recurring asexual parasitaemia originating from liver dormant

forms subsequent to a primary infection (9). Despite *P. ovale* spp. relapses are rarer than in *P.*

vivax infections, primaquine has demonstrated its ability to reduce their prevalence in

endemic settings (10), but not in non-endemic regions (11). *P. ovale* spp. are still

understudied and the development of genomic tools to study the multiplicities of infection

(MOI), population genetic parameters, and relapses are of utmost interest. We recently

developed a method to selectively amplify *P. ovale* spp. genomes over the human genome

(12), permitting whole-genome sequencing (WGS) of these parasites by next-generation

sequencing (NGS) from clinical blood samples. This is a promising approach but WGS

remains expensive and requires bioinformatics skills not available everywhere. Therefore, a

set of easy to type and highly polymorphic markers would complement the genetic tools to

study *P. ovale* spp. population genetics.

Microsatellite loci are useful molecular markers to study the population genetics and

epidemiology of malaria because they exhibit high level of polymorphism, are generally

neutral, and are frequently found in eukaryote genomes. While approximatively 30,000

microsatellites are found in their genomes (13,14), microsatellite typing protocols are missing

for both *P. ovale* spp. species, contrarily to other human-infecting *Plasmodium* species (15–

18).

In this study, we describe protocols to type *P. ovale wallikeri* microsatellite markers. Based

on the reference genome, we developed and validated eight markers located on eight different

chromosomes and genotyped 69 *P. ovale wallikeri* samples. Using these eight microsatellite

markers, we also genotyped 15 pairs of isolates from primary (D_0) and relapsing infections

($D_{relapse}$) that occurred in metropolitan France in the context of imported malaria from 2013 to

2021. For most of the pairs, the D_0 and the $D_{relapse}$ isolates were highly genetically related, an

observation that was confirmed by whole-genome sequencing for the four pairs we further

studied.

Methods

<u>Sample collection for microsatellite validation</u>

P. ovale wallikeri isolates from 2013 to 2021 were retrospectively selected from the French

National Malaria Reference Center (FNMRC) DNA bank. Genomic DNA was extracted from

200 µL of whole blood using MagNA Pure automaton (Roche diagnostics, United States of

America) and eluted in 100 µL. *P. ovale* spp. infections were confirmed using the

Plasmodium typage kit (Bio-Evolution, France). In-house qPCR-High Resolution Melting

(HRM) was used to differentiate *P. ovale wallikeri* from *P. ovale curtisi* (19).

<u>Relapse definition</u>

A *P. ovale wallikeri* relapse infection ($D_{relapse}$) in the context of imported malaria was defined

as asexual parasitaemia recurring 28 days or more after a primary *P. ovale wallikeri* infection

(D_0) treated in France and without any reported trip to a malaria-endemic country between the

two infections, thus excluding reinfections. All isolates' pairs collected between 2013 and

2021 that fit those requirements were included in the study. Epidemiological, clinical and

biological data were collected through an online patient form completed by FNMRC

correspondents.

<u>Ethical statement</u>

No specific consent from patients was required since clinical and biological data were

collected from the FNMRC database in accordance with the common public health mission of

all National Reference Centers in France, in coordination with the Santé Publique France

organization for malaria surveillance and care. The study was considered as non-

interventional research according to article L1221–1.1 of the public health code in France and

only requires the non-opposition of the patient (per article L1211–2 of the public health code).

<u>Microsatellite development and analysis</u>

We used the PowCR01 reference chromosomes (GenBank assembly accession:

GCA_900090025.2) and the software Tandem Repeat Finder (20) to identify genome-wide

microsatellites. The criteria for microsatellite selection were restricted to perfect or imperfect

microsatellite with unit motifs of 2-10 base pairs (bp) and a minimum threshold of 10 repeats.

Once the markers were selected *in silico*, several ones were experimentally tested and further

selected based on (i) efficacy of the PCR; (ii) absence of nonspecific bands on agarose gel;

(iii) location on different chromosomes; and (iv) easy-to-interpretate fragment analysis

profiles.

Forward primers were marked in 5' with 6-FAM or HEX fluorophores (Eurogentec®) and

when useful, reverse primers were pig-tailed to reduce non-templated addition of a nucleotide

to the 3' end of the amplicon (21). We used touchdown-PCR to rise sensibility and specificity

of the amplification (22). The PCR mix and cycle are detailed in the Supplemental methods.

All PCR products were run on 2% agarose gel electrophoresis. They were diluted before

fragment analysis, if necessary, in molecular biology grade water two to ten times according

to the intensity on agarose gel and run with either GeneScan^{TM} 350 ROX^{TM} or GeneScan^{TM}

500 ROX^{TM} (ThermoFisher) on an ABI 3130 genetic analyzer (ThermoFisher Scientific).

Fragment analysis profiles were interpreted with the Microsatellite Analysis software on the

Thermo Fisher Connect^{TM} website. When multiple peaks were detected, minor peaks were

scored only if their height was at least one-third of the one of the dominant peak.

Amplification of *P. falciparum*, *P. vivax*, *P. malariae* and *P. ovale curtisi* was tested for each

marker and compared to *P. ovale wallikeri*.

<u>Measurement of genetic diversity</u>

We measured allele frequencies at each marker using all the isolates ($D_{relapse}$ excluded) that

were successfully genotyped. We calculated the expected heterozygosity (H_E) for each marker

using the following formula: $H_E = \left[\frac{n}{n-1}\right]\left[1 - \sum_{i=1}^{n}p^2\right]$ where p is the frequency of the i^{th}

allele and *n* the number of different alleles (23). We counted the number of alleles (A) at each

marker in each country for which more than ten isolates were successfully genotyped. The

probability of having two isolates sharing the same allele(s) at all markers was calculated by

combining individual probabilities of each marker following this equation: $\pi p_i = P_1 \times P_2 \times$

$(\ldots) \times P_i$ where $P_i = \sum p_i^2$ with the assumption of independent infections (17). The

discriminatory power of the microsatellite haplotype was also evaluated with the Simpson

index of diversity (*D*): $D = 1 - \frac{1}{N(N-1)}\sum_{j=1}^{S}nj(nj-1)$ where *N* is the total number of

isolates within the population, *s* is the total number of different haplotypes described and *nj* is

the number of isolates belonging to the j^{th} haplotype.

Polyclonal isolates were defined as isolates with at least two scored peaks for at least one

microsatellite marker. Multiplicity of infection (MOI) was calculated for each isolate and

averaged across markers.

<u>Genetic relationship analysis</u>

We built a matrix of the allele count (only the major peak was considered for polyclonal

isolates) using the *adegenet* R package (24). Principal Component Analyses (PCA) were

made with *ade4* and plotted with the *factoextra* R package (25). Phylogenetic tree was built

by the neighbor-joining method with the *poppr* R package (26) from a matrix of Bruvo's

distance (27) and plotted with the interactive tree of life (iTOL) software (28).

<u>Comparison with microsatellite markers of primary (D_0) and relapse ($D_{relapse}$) infections</u>

We compared the microsatellite haplotypes obtained for the primary and the relapse

infections. We classified the two isolates within each pair according to three categories: (i)

homologous isolates: all of the alleles detected in the primary infections are detected in the

relapse infections; (ii) heterologous isolates: no shared allele(s) between the primary and

relapse infections for at least one microsatellite marker; (iii) related isolates: when the

primary infection is polyclonal, the D_0 and the $D_{relapse}$ isolates shared alleles at each marker

but some extra-allele(s) is detected in the primary and not in the relapse isolate.

<u>Selective whole-genome amplification and variant calling analysis</u>

To confirm the results obtained for relapses with microsatellite typing, genome-wide single

nucleotide polymorphisms (SNPs) obtained from WGS were compared for four D_0-$D_{relapse}$

isolate pairs. Prior to WGS, *P. ovale wallikeri* DNA was enriched following a sWGA strategy

and specific primers (12). The protocol is detailed in the Supplemental methods.

After WGS, raw reads were mapped to the *P. ovale wallikeri* reference genome

(GCA_900090025.2) and duplicate reads were removed using Picard MarkDuplicates (v.

2.26.10). Genome-wide SNPs were identified using BCFtools *mpileup* version 1.13 following

previously published quality criteria (12). We only kept positions with an allelic depth ≥ 5 for

the reference and/or the alternative allele. We defined as homozygous a position with $\geq 70\%$

of reads for the reference (homozygous wild-type) or the alternative allele (homozygous

mutant) (29). Otherwise, positions were defined as heterozygous. For genome-wide SNP

comparisons, only homozygous SNPs located within the fourteen chromosomes of the nuclear

genome were analyzed. Homozygous SNPs located on the contigs and heterozygous SNPs

were filtered out.

Genome-wide SNP comparisons were done for each D_0 and $D_{relapse}$ pair and the proportion of

identical genotypes between positions included for both isolates were calculated. We used

several WGS data from previous published work (12) or previous unpublished sequencing as

controls. The first control was an isolate sequenced twice (either with a sWGA or

leukodepleted approach) and named "Matched control pair". The two other controls were

built with one isolate from a relapse pair and one isolate previously sequenced from the same

country with no common history and named "Unmatched control pairs". Finally, to increase

the number of comparisons, SNPs from D_0 and $D_{relapse}$ isolates from different pairs were

compared and called "Unmatched infection pairs".

Data analysis

All statistical analyses and graphs were performed on R software v.4.0.3 (30). Quantitative

results were expressed as median [Interquartile Interval] or mean +/- Standard Deviation

(SD). Spearman's rank test was used to assess correlations. Proportions were compared using

the χ2 or Fisher's exact tests. We used the Mann-Whitney U test or the Wilcoxon signed-rank

test to compare medians.

Results

Sample collection

Isolates were selected to cover several countries in Sub-Saharan Africa and large range of

parasite density to evaluate the PCR success rate, especially for low parasite density

infections. Sixty-nine isolates covering 16 African countries and collected between January

2013 and December 2021 were included. Most of the isolates originated from West or Central

Africa and all were *P. ovale wallikeri* mono-infections (Figure 1). Among the 69 included

isolates, 52 were an isolate from an infection with no relapse history, 16 were an isolate from

a D_0 infection and 1 from a $D_{relapse}$. Isolates from $D_{relapse}$ with identical haplotype than isolate

from D_0 were not included for microsatellite marker diversity evaluation. Three isolates were

from patients who traveled through multiple African countries (not shown on the map). Ivory

Coast (n= 15 isolates), Cameroon (n= 14 isolates) and Guinea (n= 12 isolates) were the most

represented countries. Median parasite density was 3,836 [555-9,474] parasites/µL (p/µL).

Patients with relapse mainly lived in metropolitan France and experienced a primary malaria

infection after a short travel to Africa (Table S1). Median time between the D_0 and the $D_{relapse}$

infections was 108 [60-145] days with three relapses that occurred less than 50 days after the

primary infection. Delay between D_0 and $D_{relapse}$ was not different whether the primary

infection occurred during high (from July to October) or low transmission malaria season (60

days [46-107] *vs* 121 days [86-142]; $p=0.19$, Mann-Whitney *U*-test) nor in chloroquine or

non-chloroquine treated-patients (61 days [50-155] *vs* 128 days [66-142]; $p=0.39$, Mann-

Whitney *U*-test).

<u>Microsatellite marker development and validation</u>

We developed one-round PCR protocols for eight microsatellite markers, each located on a

different chromosome of *P. ovale wallikeri* (Table 1 and Figure S1) and genotyped 69 *P.*

ovale wallikeri isolates. Three markers (named 5.2, 11.1 and 14.2) displayed a sensitivity of

100% (no PCR failure; Table 2). The remaining five markers displayed a sensitivity ranging

from 94 to 99%. Failed amplifications were associated with a lower parasite density compared

to successful amplifications: 6.6% of PCR failures (n=9/136) for parasite densities ≤ 550 p/µL

vs 1% (n=4/416) for parasite densities > 550 p/µL ($p=0.001$, Fisher's exact test). Sixty-one

samples had a complete haplotype with the eight microsatellites that were successfully

genotyped and the remaining had seven (n=5 isolates), six (n=1 isolates) or five (n=2 isolates)

microsatellites successfully genotyped. Nonspecific amplification was observed for some

markers when tested against *P. falciparum* (marker 2.1) and against *P. malariae* and *P. vivax*

(markers 2.1, 8.3 and 14.2). It was significantly decreased when annealing temperature was

increased or when DNA templates contained also *P. ovale wallikeri* (Figure S2). As mono-

specific *P. ovale wallikeri* infections were analyzed in this study, this caveat had no impact on

our results.

<u>Genetic diversity</u>

From the 69 isolates investigated, we counted from 9 to 17 different alleles per microsatellite

(Table 2) and 65 eight-marker haplotypes were unique (Table S2). The expected

heterozygosity H_E was calculated for each marker and varied from 0.764 to 0.940. For each

microsatellite, allele frequencies were calculated in the total cohort (Figure 2 and Table S3)

and in Cameroon, Guinea and Ivory Coast (Table S3). The combined likelihood that two

random samples had the same eight-markers haplotype was 2.1×10^{-6} (Table S4). The Simpson

index of diversity was equivalent (*D*=0.9987) using 8 or only 6 markers (2.1 and 10.1 markers

excluded, the two loci with the lower H_E). *D* value remained high with the four markers with

the highest H_E (*D*=0.9978; combination of the 8.3, 11.1, 12.1 and 14.2 markers), highlighting

their combined high discriminatory power.

The diversity (based on the mean number of alleles, A, and on the mean expected

heterozygosity, H_E) was found to be high in Cameroon and Guinea (Table S5). Interestingly,

seven of the 69 isolates displayed two peaks for at least one marker and were considered as

polyclonal (Table S2).

<u>Population structure analysis</u>

We next studied population structure between West (n=44) and Central African (n=21)

isolates, and between isolates from Cameroon (n=14), Guinea (n=12) and Ivory Coast (n=15),

respectively. In both analyses, PCA based on microsatellite allele sizes was not able to

separate isolates according to their geographic origin (Figure S3). Similarly, the phylogenetic

tree reconstructed using eight-marker haplotypes was not able to separate the isolates from

West and Central Africa or from Cameroon, Guinea and Ivory Coast (Figure S4 and S5).

<u>Application of the microsatellite markers for relapse typing</u>

Among the 17 D_0-$D_{relapse}$ isolate pairs, two were not analyzable because of PCR failures at

more than four markers, likely due to very low parasite density (Table S6). Eleven pairs had

homologous parasites (Table S6). Three pairs (pairs 12, 14 and 17) had at least one marker

with multiple peaks for the D_0 but a single shared peak for the $D_{relapse}$ and were then

considered as related but not strictly homologous. For two of those three pairs (pairs 12 and

14), the $D_{relapse}$ was monoclonal. For the last pair (pair 17), both D_0 and $D_{relapse}$ displayed

multiple peaks for respectively 4 and 1 microsatellite markers. Finally, the pair 15 had

heterologous parasites, with no shared allele for the marker 5.2 (allele sizes were 253 and 250

for D_0 and $D_{relapse}$ respectively, corresponding to one trinucleotide TAC motif).

<u>Genome-wide SNP analysis</u>

To complete the microsatellite results, we performed WGS for four homologous D_0-$D_{relapse}$

isolate pairs (Table S7) and counted the number of SNPs that differed within each pair. For

the matched control pair (same isolate sequenced twice), 99.93% of the SNPs were identical,

compared to 31.7% for the unmatched control pairs and 30.7% for the unmatched infection

pairs (Figure 3 and Table S8). For the four D_0-$D_{relapse}$ pairs, (pairs 1, 2, 4 and 5; see Table S8)

an average of 99.64% SNPs were identical (Figure 3) with very few different SNPs within

pairs (Figure 4). No statistical difference in the count of identical SNPs was observed between

the matched control pair and the relapses (p=0.93, χ2 test). In contrast, the number of identical

SNPs statistically differed between the unmatched control pairs or the unmatched infection

pairs and the relapses (p<0.001, χ2 test).

Discussion

Despite several epidemiological evidences of *P. ovale* spp. relapses during the last seventy

years (6,8,11,31–33), biological realness of *P. ovale* spp. hypnozoites is still debated (9,34).

Indeed, proofs of hypnozoite existence are scarce, but Soulard *et al*. (35) described late-

developing schizontes in humanized mice infected with *P. ovale* spp. sporozoites, consistent

with a hypnozoite origin of those maturing forms (35).

Easy-to-use molecular tools are missing to study the genetic relatedness of *P. ovale* spp.

isolates. We developed eight microsatellite markers that displayed very high H_E and high

combined probability value (Table 2), suggesting a great discriminative potential for

epidemiological studies. Interestingly, the combination of just four markers (8.3, 11.1, 12.1

and 14.2) remains highly informative. These could be used as a minimal set to genetically

profile *P. ovale wallikeri* isolates for time and resource saving. We observed high diversity

with 65 unique haplotypes and only 3 non-unique haplotypes each shared by two isolates

from West Africa that did not have reported common history (Figure S4 and Table S2). More

studies, in particular at a country level, are needed to better investigate *P. ovale wallikeri*

population structure in West and Central Africa.

With these eight newly developed microsatellite markers, we genotyped 17 pairs of *P. ovale*

wallikeri isolates that fitted with an epidemiological relapse definition. As our study is the

first one that genetically analyzed *P. ovale wallikeri* relapses, we used *P. vivax* relapses as a

reference. Our relapse definition is purely based on the epidemiology and history of cases as

we did not have complete parasitemia follow-up for each patient. As the World Health

Organization recommends a 28 days parasitaemia follow-up for *P. vivax* (36), we did not

include in this study pairs of isolates with time to recurring parasitemia below this threshold.

Of note, three pairs had short delay between the two infections (28, 33 and 39 days). Even

with precise molecular tools, these relapse infections could not be formally distinguished

from a recrudescent infection caused by incomplete treatment clearance of the blood-stage

primary infection. We decided to classify those three pairs as homologous relapse and not

recrudescence because : (i) in *P. vivax*, recrudescences seem to be much less frequent than

relapses (37,38) and (ii) to the best we know, *P. ovale* spp. treatment failure is rarely reported

in the literature (no failure reported in published clinical trials (39–43), except one persistent

P. ovale wallikeri on Day-28 in a *P. falciparum* co-infection in Gabon (44)).

Among the 15 analyzable pairs, 11 were scored as genetically homologous. For four out of

these 11 genetically homologous pairs also tested by WGS, the very high genetic relatedness

of D_0 and $D_{relapse}$ isolates was confirmed. Contrary to most of *P. vivax* relapses studied in

Thailand where heterologous relapses are the norm (45) (exception made of the relapses

observed in young infants that were mainly homologous (46)), we report high percentage of

homologous relapses. This has to be interpreted in the light of two key features: (i) we studied

imported malaria cases. Patients mainly live in metropolitan France with short travel to Africa

limiting the probabilities, compared to patients living in endemic area, of a previous *P. ovale*

wallikeri infection or multiple infections (two ways to get heterologous relapses (47,48)); and

(ii) according to the microsatellite results, the polyclonal infection (second way to get

heterologous relapses (45)) rate in *P. ovale wallikeri* isolates from imported malaria appears

to be much lower compared to the one in *P. vivax* (10% for *P. ovale wallikeri* in our work *vs* a

mean proportion of polyclonal infections of 43% for *P. vivax* in endemic zone (49)).

The genetic relatedness within the four last pairs (12, 14, 15 and 17; Table S7) was more

complex. For pairs 12, 14 and 17, D_0 isolates were polyclonal while $D_{relapse}$ carried no new

alleles, confirming their high genetic relatedness. Interestingly, $D_{relapse}$ from pairs 12 and 14

were monoclonal contrary to relapse from pair 17 that was polyclonal. This may result from

different ways of hypnozoite reactivation, stochastic for pairs 12 and 14 (*i.e.* one hypnozoite

reactivation) and external stimulus for pair 17 (*i.e.* at least two hypnozoites reactivation)(47).

Finally, the two isolates from pair 15 were both monoclonal and differed at a single marker by

just one tandem repeat. Although initially classified as heterologous, the new allele in $D_{relapse}$

could be simply derived by mutation of the D_0 allele early during the course of the relapse

infection. Slippage during the PCR was excluded since PCR was repeated.

In conclusion, using a novel set of eight microsatellite markers we genotyped for the first time

P. ovale wallikeri isolates from primary and relapse infections in the context of imported

malaria. As expected, most paired primary and relapse isolates appeared to be highly

genetically related. WGS should help in the future to get a more complete genetic comparison

of D_0-$D_{relapse}$ isolate pairs and also to look for putative genetic markers associated with *P.*

ovale wallikeri dormancy.

Data availability

The genome Illumina sequencing reads from the *P. ovale wallikeri* samples were deposited in

the European Nucleotide Archive under the accession number PRJEB51041.

About the author

Valentin Joste is a medical microbiologist working in pathology laboratory medicine at the

Bichat hospital and at the French National Malaria Reference Center. His research interests

include *P. ovale* spp. diagnosis and epidemiology.

Acknowledgments

We thank the hospital correspondents who sent us *P. ovale wallikeri* samples and filled in the

online patient forms.

Additional members of the French National Reference Center for Imported Malaria Study

Group who contributed data: Sylvain Clauser (Boulogne), Ludovic de Gentile (Angers), Marc

Pihet (Angers), Jean-Philippe Lemoine (Angers), Vincent Dubee (Angers), Anthony Marteau

(Avicenne), Arezki Izri (Avicenne), Cécile Ficko (Bégin), Jean-Etienne Pilo (Bégin),

Sébastien Larréché (Bégin), Anne-Pauline Bellanger (Besançon), Adela Enache Angoulvant

(Bicêtre), Nadia Guennouni (Bicêtre), Didier Raffenot (Chambéry), Dr Olivier Rogeaux

(Chambéry), Céline Nourrisson (Clermont-Ferrand), Philippe Poirier (Clermont-Ferrand),

Naima Dahane (Cochin hospital, Paris), Cécile Garnaud (Grenoble), Marie Gladys Robert

(Grenoble), Cécile Garnaud (Grenoble), Céline Dard (Grenoble), Hervé Pelloux (Grenoble),

Dominique Maubon (Grenoble), Eric Dannaoui (Hôpital Européen Georges Pompidou

hospital, Paris), Françoise Botterel-Chartier (Henri-Mondor hospital), Dieudonné Bemba

(Jean-Verdier hospital), Isabelle Poilane (Jean-Verdier hospital), Muriel Silva (Le Havre),

Céline Malassigne (Le Havre), Sophie Deleplancque (Lille), Boualem Sendid (Lille), Marie-

Laure Darde (Limoges), Marie-Fleur Durieux (Limoges), Anne Debourgogne (Nancy), Fakhri

Jeddi (Nantes), Rose-Anne Lavergne (Nantes), Marie-Elisabeth Bougnoux (Necker hospital,

Paris), Emilie Sitterle (Necker hospital, Paris), Bernadette Buret (Niort), Ali Cherif Touil

(Niort), Didier Poisson (Orléans), Jérôme Guinard (Orléans), Aurélie Guigon (Orléans), Marc

Thellier (Pitié-Salpétrière hospital, Paris), Estelle Perraud-Cateau (Poitiers), Gwénaël le Moal

(Poitiers), Antoine Huguenin (Reims), Sorya Belaz (Rennes), Odile Fenneteau (Robert Debré

hospital, Paris), Jean-Yves Siriez (Robert Debré Hospital, Paris), Lauren Pull (Robert Debré

hospital, Paris), Loïc Favennec (Rouen), Gilles Gargala (Rouen), Alicia Moreno-Sabater

(Saint-Antoine hospital, Paris), Cécile Tournus (Saint-Denis), Samia Hamane (Saint-Louis

hospital, Paris), Stéphane Bretagne (Saint-Louis hospital, Paris), Julie Brunet (Strasbourg),

Ahmed Abou-Bacar (Strasbourg), Ghania Belkacem Belkadi (Tenon hospital), Pamela

Chauvin (Toulouse), Antoine Berry (Toulouse), Nathalie Desuremain (Trousseau hospital,

Paris), Gisèle Dewulf (Valenciennes), Edith Mazars (Valenciennes), Odile Eloy (Versailles).

Bibliography

1.   World malaria report 2022 [Internet]. [cited 2022 Dec 11]. Available from:
https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2022
2.   Hawadak J, Dongang Nana RR, Singh V. Global trend of Plasmodium malariae and
Plasmodium ovale spp. malaria infections in the last two decades (2000–2020): a systematic
review and meta-analysis. Parasites & Vectors. 2021 Jun 3;14(1):297.
3.   Yman V, Wandell G, Mutemi DD, Miglar A, Asghar M, Hammar U, et al. Persistent
transmission of Plasmodium malariae and Plasmodium ovale species in an area of declining
Plasmodium falciparum transmission in eastern Tanzania. PLOS Neglected Tropical
Diseases. 2019 May 28;13(5):e0007414.
4.   Groger M, Lutete GT, Mombo-Ngoma G, Ntamabyaliro NY, Mesia GK, Mujobu
TBM, et al. Effectiveness of pyronaridine-artesunate against Plasmodium malariae,
Plasmodium ovale spp, and mixed-Plasmodium infections: a post-hoc analysis of the
CANTAM-Pyramax trial. The Lancet Microbe. 2022 Aug 1;3(8):e598–605.
5.   Sutherland CJ, Tanomsing N, Nolder D, Oguike M, Jennison C, Pukrittayakamee S, et
al. Two nonrecombining sympatric forms of the human malaria parasite Plasmodium ovale
occur globally. J Infect Dis. 2010 May 15;201(10):1544–50.
6.   Joste V, Bailly J, Hubert V, Pauc C, Gendrot M, Guillochon E, et al. Plasmodium
ovale wallikeri and P. ovale curtisi Infections and Diagnostic Approaches to Imported
Malaria, France, 2013-2018. Emerg Infect Dis. 2021 Feb;27(2).
7.   Rojo-Marcos G, Rubio-Muñoz JM, Angheben A, Jaureguiberry S, García-Bujalance
S, Tomasoni LR, et al. Prospective comparative multi-centre study on imported Plasmodium
ovale wallikeri and Plasmodium ovale curtisi infections. Malar J. 2018 Oct 30;17(1):399.
8.   Groger M, Veletzky L, Lalremruata A, Cattaneo C, Mischlinger J, Manego Zoleko R,
et al. Prospective Clinical and Molecular Evaluation of Potential Plasmodium ovale curtisi
and wallikeri Relapses in a High-transmission Setting. Clin Infect Dis. 2019 Nov
27;69(12):2119–26.
9.   Markus MB. Do hypnozoites cause relapse in malaria? Trends Parasitol. 2015
Jun;31(6):239–45.
10.   Robinson LJ, Wampfler R, Betuela I, Karl S, White MT, Li Wai Suen CSN, et al.
Strategies for understanding and reducing the Plasmodium vivax and Plasmodium ovale
hypnozoite reservoir in Papua New Guinean children: a randomised placebo-controlled trial
and mathematical model. PLoS Med. 2015 Oct;12(10):e1001891.
11.   Wångdahl A, Sondén K, Wyss K, Stenström C, Björklund D, Zhang J, et al. Relapse
of Plasmodium vivax and Plasmodium ovale Malaria With and Without Primaquine
Treatment in a Nonendemic Area. Clinical Infectious Diseases. 2022 Apr 1;74(7):1199–207.
12.   Joste V, Guillochon E, Clain J, Coppée R, Houzé S. Development and Optimization of

a Selective Whole-Genome Amplification To Study Plasmodium ovale Spp. Microbiol
Spectr. 2022 Sep 13;e0072622.
13. Rutledge GG, Böhme U, Sanders M, Reid AJ, Cotton JA, Maiga-Ascofare O, et al.
Plasmodium malariae and P . ovale genomes provide insights into malaria parasite evolution.
Nature. 2017 Feb;542(7639):101–4.
14. Mathema VB, Nakeesathit S, White NJ, Dondorp AM, Imwong M. Genome-wide
microsatellite characteristics of five human Plasmodium species, focusing on Plasmodium
malariae and P. ovale curtisi. Parasite. 2020;27:34.
15. Imwong M, Nair S, Pukrittayakamee S, Sudimack D, Williams JT, Mayxay M, et al.
Contrasting genetic structure in Plasmodium vivax populations from Asia and South America.
Int J Parasitol. 2007 Jul;37(8–9):1013–22.
16. Greenhouse B, Myrick A, Dokomajilar C, Woo JM, Carlson EJ, Rosenthal PJ, et al.
VALIDATION OF MICROSATELLITE MARKERS FOR USE IN GENOTYPING
POLYCLONAL PLASMODIUM FALCIPARUM INFECTIONS. The American journal of
tropical medicine and hygiene. 2006 Nov;75(5):836.
17. Mathema VB, Nakeesathit S, Pagornrat W, Smithuis F, White NJ, Dondorp AM, et al.
Polymorphic markers for identification of parasite population in Plasmodium malariae.
Malaria Journal. 2020 Jan 28;19(1):48.
18. Divis PCS, Singh B, Anderios F, Hisam S, Matusop A, Kocken CH, et al. Admixture
in Humans of Two Divergent Plasmodium knowlesi Populations Associated with Different
Macaque Host Species. PLOS Pathogens. 2015 May 28;11(5):e1004888.
19. Joste V, Kamaliddin C, Kendjo E, Hubert V, Argy N, Houzé S. Distinction of
Plasmodium ovale wallikeri and Plasmodium ovale curtisi using quantitative Polymerase
Chain Reaction with High Resolution Melting revelation. Scientific Reports. 2018 Jan
10;8(1):1–8.
20. Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic
Acids Res. 1999 Jan 15;27(2):573–80.
21. Brownstein MJ, Carpten JD, Smith JR. Modulation of non-templated nucleotide
addition by Taq DNA polymerase: primer modifications that facilitate genotyping.
Biotechniques. 1996 Jun;20(6):1004–6, 1008–10.
22. Korbie DJ, Mattick JS. Touchdown PCR for increased specificity and sensitivity in
PCR amplification. Nat Protoc. 2008 Sep;3(9):1452–6.
23. Sutton PL, Luo Z, Divis PCS, Friedrich VK, Conway DJ, Singh B, et al.
Characterizing the genetic diversity of the monkey malaria parasite Plasmodium cynomolgi.
Infect Genet Evol. 2016 Jun;40:243–52.
24. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics. 2008 Jun 1;24(11):1403–5.
25. Thioulouse J, Chessel D, Dole´dec S, Olivier JM. ADE-4: a multivariate analysis and
graphical display software. Statistics and Computing. 1997 Mar 1;7(1):75–83.
26. Kamvar ZN, Tabima JF, Grünwald NJ. Poppr: an R package for genetic analysis of
populations with clonal, partially clonal, and/or sexual reproduction. PeerJ. 2014 Mar
4;2:e281.
27. Bruvo R, Michiels NK, D'souza TG, Schulenburg H. A simple method for the
calculation of microsatellite genotype distances irrespective of ploidy level. Molecular
Ecology. 2004;13(7):2101–6.
28. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic Acids Research. 2021 Jul 2;49(W1):W293–6.
29. Moser KA, Drábek EF, Dwivedi A, Stucke EM, Crabtree J, Dara A, et al. Strains used
in whole organism Plasmodium falciparum vaccine trials differ in genome structure,
sequence, and immunogenic potential. Genome Medicine. 2020 Jan 8;12(1):6.

30.   Wickham, H. and Henry, L. (2018). tidyr: Easily Tidy Data with "spread()" and
"gather()" Functions. R package version 0.8.1. https://CRAN.R-project.org/package=tidyr.
31.   Garnham PCC, Bray RS, Cooper W, Lainson R, Awad FI, Williamson J. The pre-
erythrocytic stage of Plasmodium ovale. Trans R Soc Trop Med Hyg. 1955 Mar 1;49(2):158–
67.
32.   Austin SC, Stolley PD, Lasky T. The History of Malariotherapy for Neurosyphilis:
Modern Parallels. JAMA. 1992 Jul 22;268(4):516–9.
33.   Collins WE, Jeffery GM. A retrospective examination of sporozoite-induced and
trophozoite-induced infections with Plasmodium ovale: development of parasitologic and
clinical immunity during primary infection. Am J Trop Med Hyg. 2002 May;66(5):492–502.
34.   Richter J, Franken G, Mehlhorn H, Labisch A, Häussinger D. What is the evidence for
the existence of Plasmodium ovale hypnozoites? Parasitol Res. 2010 Nov;107(6):1285–90.
35.   Soulard V, Bosson-Vanga H, Lorthiois A, Roucher C, Franetich JF, Zanghi G, et al.
Plasmodium falciparum full life cycle and Plasmodium ovale liver stages in humanized mice.
Nature Communications. 2015 Jul 24;6:7690.
36.   Methods for surveillance of antimalarial drug efficacy [Internet]. [cited 2022 Dec 21].
Available from: https://www.who.int/publications-detail-redirect/9789241597531
37.   Taylor AR, Watson JA, Chu CS, Puaprasert K, Duanguppama J, Day NPJ, et al.
Resolving the cause of recurrent Plasmodium vivax malaria probabilistically. Nat Commun.
2019 Dec 6;10(1):5595.
38.   Popovici J, Pierce-Friedrich L, Kim S, Bin S, Run V, Lek D, et al. Recrudescence,
Reinfection, or Relapse? A More Rigorous Framework to Assess Chloroquine Efficacy for
Plasmodium vivax Malaria. J Infect Dis. 2019 Jan 7;219(2):315–22.
39.   Visser BJ, Wieten RW, Kroon D, Nagel IM, Bélard S, van Vugt M, et al. Efficacy and
safety of artemisinin combination therapy (ACT) for non-falciparum malaria: a systematic
review. Malaria Journal. 2014 Nov 26;13(1):463.
40.   Mombo-Ngoma G, Kleine C, Basra A, Würbel H, Diop DA, Capan M, et al.
Prospective evaluation of artemether-lumefantrine for the treatment of non-falciparum and
mixed-species malaria in Gabon. Malar J. 2012 Jul 10;11:120.
41.   Lwin KM, Phyo AP, Tarning J, Hanpithakpong W, Ashley EA, Lee SJ, et al.
Randomized, double-blind, placebo-controlled trial of monthly versus bimonthly
dihydroartemisinin-piperaquine chemoprevention in adults at high risk of malaria. Antimicrob
Agents Chemother. 2012 Mar;56(3):1571–7.
42.   Siswantoro H, Russell B, Ratcliff A, Prasetyorini B, Chalfein F, Marfurt J, et al. In
Vivo and In Vitro Efficacy of Chloroquine against Plasmodium malariae and P. ovale in
Papua, Indonesia. Antimicrobial Agents and Chemotherapy. 2011 Jan;55(1):197–202.
43.   Groger M, Fischer HS, Veletzky L, Lalremruata A, Ramharter M. A systematic review
of the clinical presentation, treatment and relapse characteristics of human Plasmodium ovale
malaria. Malar J. 2017 11;16(1):112.
44.   Groger M, Veletzky L, Lalremruata A, Cattaneo C, Mischlinger J, Zoleko-Manego R,
et al. Prospective Clinical Trial Assessing Species-Specific Efficacy of Artemether-
Lumefantrine for the Treatment of Plasmodium malariae, Plasmodium ovale, and Mixed
Plasmodium Malaria in Gabon. Antimicrob Agents Chemother. 2018 Mar;62(3):e01758-17.
45.   Imwong M, Snounou G, Pukrittayakamee S, Tanomsing N, Kim JR, Nandy A, et al.
Relapses of Plasmodium vivax infection usually result from activation of heterologous
hypnozoites. J Infect Dis. 2007 Apr 1;195(7):927–33.
46.   Imwong M, Boel ME, Pagornrat W, Pimanpanarak M, McGready R, Day NPJ, et al.
The first Plasmodium vivax relapses of life are usually genetically homologous. J Infect Dis.
2012 Feb 15;205(4):680–3.
47.   Popovici J, Friedrich LR, Kim S, Bin S, Run V, Lek D, et al. Genomic Analyses

Reveal the Common Occurrence and Complexity of Plasmodium vivax Relapses in
Cambodia. mBio. 2018 Jan 23;9(1):e01888-17.
48.   Bright AT, Manary MJ, Tewhey R, Arango EM, Wang T, Schork NJ, et al. A High
Resolution Case Study of a Patient with Recurrent Plasmodium vivax Infections Shows That
Relapses Were Caused by Meiotic Siblings. PLoS Negl Trop Dis. 2014 Jun 5;8(6):e2882.
49.   Lopez L, Koepfli C. Systematic review of Plasmodium falciparum and Plasmodium
vivax polyclonal infections: Impact of prevalence, study population characteristics, and
laboratory procedures. PLOS ONE. 2021 Jun 11;16(6):e0249382.
525
526

527 **Tables**

528 **Table 1 – List of the microsatellite markers for *P. ovale wallikeri* genotyping.**

Name	Chr	Primer sequence (5'-3')$^\$$	Position : end-start*	Repeat sequence*
2.1	2	F: 6FAM-TTTTAAAGTTGCCGCATTGTC	103,121-103,270	$(GTT)_{12}(GCT)_9$
		R: gtttcttATACCAAAAGCGCTGTCTCC	intergenic	
5.2	5	F: HEX-ACACCCGAAGTGGTATCAGC	664,493-664,733	$(TAC)_{10}$
		R: ACGGATGCGCTATAATTTGG	POWCR01_050018800	
8.3	8	F: HEX-CCCCTTGCAAACGTGTAAAT	855,305-855,486	$(CTAC)_{16}$
		R: GTGTTTGTGCACATGGAAGG	intergenic	
10.1	10	F: HEX-GATTCATCGGTGCTTTCTCC	432,312-432,527	$(CGT)_{12}$
		R: GGAAGAAACGGAAGAAGTGG	POWCR01_100015100	
11.1	11	F: HEX-AAGGGGTGGGGGAGTTTAC	539,226-539,442	$(GTT)_{16}$
		R: gtttcttCAGGCACGTACTAGCATGGA	intergenic	
12.1	12	F: 6FAM-ACATGCAACGCAACTTTCAG	1,540,305-1,540,570	$(CATA)_{14}$
		R: GTGCGCCTCCACTTTTTACT	intergenic	
13.3	13	F: 6FAM-CGAATGAAGTAGCACAGTTAAGC	268,664-268,882	$(TTC)_{10}$
		R: GCTCAATATCGCTCTTTTCCA	intergenic	
14.2	14	F: 6FAM-GGCATTTTTCCATTTCGTCA	1,141,610-1,141,858	$(ATAC)_{11}(AT)_5$
		R: AAAATTGATAAGGCGCGAGA	intergenic	

529 $^\$$F: forward, R: reverse, gtttctt: pig-tailed primer, Chr: chromosome.

530 *on the PowCR01 reference genome (GenBank assembly accession: GCA_900090025.2)

533   **Table 2 – Characteristics of the microsatellite markers tested on 69** *P. ovale wallikeri*

534   **isolates.**

Microsatellite	PCR length range (bp)	PCR success rate	Allelic diversity, A	Expected heterozygosity (H_E)	MOI$^\$$ mean (max*)
2.1	132-159	99%	10	0.764	1.04 (2)
5.2	233-263	100%	9	0.849	1.01 (2)
8.3	127-608	99%	15	0.902	1.04 (2)
10.1	203-239	97%	10	0.786	1.01 (2)
11.1	185-279	100%	17	0.932	1.03 (2)
12.1	227-356	94%	13	0.906	1.06 (2)
13.3	183-267	94%	14	0.834	1.00 (1)
14.2	227-344	100%	14	0.940	1.01 (2)

535    $^\$$MOI: multiplicity of infection.

536   *max: maximum number of peaks detected in one sample.

537   bp stands for base pair.

Figure 1 – Geographic origins of the *P. ovale wallikeri* isolates included in the study.
Countries with included isolates are colored in blue. Number of isolates per country is indicated in salmon circle.

Figure 2 - Number of alleles and their frequencies for the eight microsatellite markers.

Figure 3 - Proportion of identical genome-wide SNPs within infection pairs.
Matched control pair and D_0-$D_{relapse}$ pairs are in light brown. Unmatched control pairs and unmatched pairs are in gray. Each dot represents one pair. NS stands for non-significant (χ_2 tests).

Figure 4 – Genetic relatedness within matched or unmatched infection pairs.
The distribution of the different alleles within each D_0-$D_{relapse}$ pair is shown throughout the fourteen chromosomes of the nuclear genome. Each cross represents a 10 kb window containing at least one differing SNP, represented according to its genomic position (x axis, nucleotide position in base pair; y axis, chromosome). Beige triangle shows the end of each chromosome.