Risk assessment during nest defense against three simulated predators by female

northern house wrens (*Troglodytes aedon*)

Ross C. Eggleston[1,‡], Josephina H. Fornara[1,2,‡], Kyle M. Davis[1,3], Jess Dong[1,4], and Dustin G.

Reichard[1,*]

[1]Ohio Wesleyan University, Department of Zoology, 61 S Sandusky St, Delaware, OH 40315

[2]Current address: Indiana University, Department of Biology, 1001 E Third St, Bloomington,

Indiana 47405

[3]Current address: Michigan State University, Department of Integrative Biology, 288 Farm Ln,

East Lansing, MI 48824

[4]Current address: Ohio State University, School of Environment and Natural Resources, 2021

Coffey Rd, Columbus, OH 43210

*Corresponding author email: dgreicha@owu.edu

‡Equal author contribution

Conflict of Interest Statement

None of the authors declare any conflicts of interest.

Data Availability Statement

The data collected and analyzed to support these findings are freely available in the

Mendeley Data Repository at DOI:10.17632/zjb5ynz5n4.2.

Funding Statement

This study was funded by the Ohio Wesleyan University Summer Science Research

Program.

Ethics Approval Statement

All procedures were reviewed and approved by the Ohio Wesleyan University

Institutional Animal Care and Use Committee (Protocol Numbers: Spring_2020-21-

01_Reichard and 05-2021-06), U.S. Federal Bird Banding Lab (Permit Number: 24035), and

State of Ohio Department of Natural Resources (Permit Numbers: 23-009, 23-010).

Abstract

Offspring predation is one of the greatest obstacles to an organism's reproductive success, but parents vary in the strength of their response to potential predators. One explanation for this variable investment is that defending current offspring has the potential to lower future reproductive success if the predator is also capable of injuring or killing the parent. Northern house wrens (*Troglodytes aedon*) are cavity-nesting songbirds that defend against multiple species of nest predators including small mammals, birds of prey, and snakes. Here, we used three different predator decoys: two nest predators - an eastern chipmunk (*Tamias striatus*) and an eastern ratsnake (*Pantherophis alleghaniensis*) - as well as a predator of both offspring and adults - a juvenile Cooper's hawk (*Accipiter cooperi*) - to elicit nest defense and test whether females use risk assessment to modulate their antipredator behavior. We found that antipredator behaviors were not significantly different between the two nest predators, which posed a high risk to the nestlings, but lower risk to the parents as neither species frequently captures adult wrens outside the nest box. However, female wrens never dove at or attacked the Cooper's hawk while they frequently attacked both the snake and chipmunk decoys. Neighboring house wrens from adjacent territories were also less likely to respond to the hawk, but more heterospecifics mobbed the hawk than the snake decoy. Collectively, these results show that risk assessment and the strength of the antipredator response varies substantially both within and among species. Female house wrens exhibit plasticity in their nest defense behavior, and they respond to different types of predators in a way that could maximize lifetime fitness while risking the loss of their current offspring.

Introduction

Parental investment can be defined as any behaviors that increase an offspring's chance of survival while simultaneously decreasing the parent's ability to invest in other offspring both currently and in the future (Trivers 1974; Montgomerie and Weatherhead 1988; Clutton-Brock 1991). This tradeoff between current and future reproduction creates a parent-offspring conflict (*sensu* Trivers 1974) because the optimal parenting strategy, one that maximizes the parent's lifetime fitness, often requires submaximal investment in the current offspring, which can decrease their survival and fitness. The amount of parental investment varies widely both within and among species (Fontaine and Martin 2006; Dulac et al. 2014), and this variation can be further explained by differences in life history strategies (Ricklefs 1977; Montgomerie and Weatherhead 1988; Clutton-Brock 1991; Ghalambor and Martin 2001). For example, both adult mortality rates and offspring number and age should correlate positively with parental investment because species with shorter lifespans and larger current investments will have fewer reproductive opportunities in the future (Caro 2005).

Optimally modulating parental investment is particularly critical in the context of antipredator behavior, as parents that vigorously defend their young risk completely eliminating their future reproduction if they are mortally injured or killed. On the other hand, parents that limit their antipredator behavior are more likely to lose their current offspring as predation is one of the most common causes of reproductive failure across species (Ricklefs 1977; Caro 2005; Martin and Briskie 2009; Ibáñez-Álamo et al. 2015). The most successful parents should be those that exhibit plastic antipredator behavior that maximizes their lifetime reproductive success, and selection should favor individuals with accurate risk assessment during encounters with predators near their offspring (Montgomerie and Weatherhead 1988).

An abundance of evidence supports the hypothesis that animals from across the

taxonomic spectrum use risk assessment to adjust their antipredator behavior relative to the

perceived threat level (reviewed in Caro 2005; Lima 2009). Species with altricial young, such

as passerine songbirds, are particularly vulnerable to predation at the nest because their

offspring have limited capabilities for self-defense or escape without parental assistance (e.g.,

Suzuki 2011). Consequently, parents respond to different predation pressures with both

proactive and reactive behavior. When the perceived risk of predation is high, parents can

proactively reduce egg or clutch sizes (Fontaine and Martin 2006; Morosinotto et al. 2010;

Travers et al. 2010) or limit visits to the nest, including offspring provisioning (Fontaine and

Martin 2006; Ghalambor et al. 2013; Dudeck et al. 2018), which minimizes their potential

energetic losses and draws less attention to the nest site. Alternatively, parents may simply

avoid nesting in locations with high predator abundance or even forgo reproduction entirely

(Spaans et al. 1998; Morosinotto et al. 2010; Szymkowiak and Thomson 2019).

When a predator appears at the nest site, parents can react in ways that might divert the

predator's attention toward the parents rather than the offspring such as alarm calling or visual

distraction displays (e.g. broken-wing display; de Framond et al. 2022). Alarm calling also has

the potential to initiate a community-level response by attracting nearby conspecifics and

heterospecifics to collectively harass (i.e., "mob") the predator, which may increase the

likelihood of expelling the predator from the area (Carlson and Griesser 2022). Although alarm

calls and visual displays increase risk to the parents, they are often less costly than more

aggressive defense behaviors such as diving at or hitting the predator in an attempt to

physically drive it away from the offspring. Studies from multiple species have shown that

parents exhibit plasticity in their antipredator responses and adjust their behavior according to

factors such as the distance between the predator and the offspring (Colombelli-Négrel et al.

2010; Fernández and Carro 2022) and the relative risks posed to both adults and offspring by

different types of predators (Gottfried 1979; Winkler 1992; Mahr et al. 2015; Duré Ruiz et al.

2018; Teunissen et al. 2020). Similarly, the willingness of nearby conspecifics and

heterospecifics to respond to alarm calls with mobbing behavior varies according to the

intensity of the calls (Templeton et al. 2005, Fernández et al. 2023) and the risks posed by the

predator (Forsman and Mönkkönen 2001, Scott and Robinson 2023). In this study, we tested

for evidence of risk assessment by female northern house wrens (*Troglodytes aedon*) during

simulated interactions with predators at their nest that posed different threats to adults and

nestlings.

Previous research in both northern and southern house wrens (Troglodytes musculus) has

found evidence for risk assessment and plasticity in parental behavior in response to predation

pressure. For example, wren parents produced more alarm calls when an owl model was placed

closer to their nest (Fernández and Carro 2022), consistent with an ability to perceive levels of

risk. After interacting with a simulated nest predator, wren parents have also been shown to

reduce their offspring provisioning (Freed 1981; Ghalambor et al. 2013) and take longer to re-

enter the nest box when compared to control observations (Corral et al. 2013; Fernandez and

Llambías 2013; Fernández et al. 2015; Duré Ruiz et al. 2018). When presented with different

simulated species of raptors of varying threat levels, male and female southern house wrens

took longer to resume normal parental behavior after encountering a predator of both adults

and nestlings than a predator of nestlings only (Duré Ruiz et al. 2018). This result suggests that

risk assessment may shape parental behavior in adult wrens by shifting investment away from

offspring and towards self-preservation when perceived threat levels are high.

Here, we expand on these observations by testing whether wren parents assess risks

posed by both avian and non-avian predators and adjust the strength of their nest defense

relative to risks associated with each type of predator (see below). Over two breeding seasons,

we placed three simulated predators of varying risks on top of house wren nest boxes during

the early nestling stage. One predator, a rubber eastern ratsnake decoy (*Pantherophis*

alleghaniensis), was presented in both years, while the second predator differed between years

and consisted of a rubber eastern chipmunk decoy (*Tamias striatus*) in year one and a

taxidermied juvenile Cooper's Hawk (*Accipiter cooperii*) in year two. We assumed that the

ratsnake and chipmunk decoys posed a high risk to nestlings, but a low-to-moderate risk to

adult wrens, which are unlikely to be captured by either predator outside of the nest box (Reidy

et al., 2009; Johnson, 2020). In contrast, the hawk posed a high risk to adults, but a low-to-

moderate risk to nestlings, which are still vulnerable to raptors and other avian predators

despite the protection offered by an enclosed nest box (Milsap et al. 2013; Johnson 2020). If

the hypothesis that house wrens use risk assessment was supported, we predicted that female

wrens would respond equally aggressively to the ratsnake and chipmunk decoys, which are

similar in threat level to both nestlings and adults. Female responses to the hawk were

predicted to be less aggressive overall due to the higher threat that the hawk posed to adults

and lower threat posed to nestlings. Finally, we examined the community-level response to

predators of varying risk by quantifying the number of conspecific and heterospecific

individuals that responded to each predator treatment.

Methods

Study System and Sites

Northern house wrens (*Troglodytes aedon*) are socially monogamous, cavity-nesting

songbirds that readily use artificial nest boxes to raise clutches of up to 8 offspring (Johnson

2020). The guild of house wren nestling predators is diverse and includes mammals ranging in

size from mice (e.g., *Peromyscus* spp.) to bears (*Ursus spp.*), passerines (e.g., *Cyanocitta* spp.),

woodpeckers (e.g., *Melanerpes* spp.), raptors, and snakes (Johnson 2020). At our study sites

the most common house wren nest predators are small mammals and snakes, but multiple

raptor species also frequent our sites and pose risks to adults and offspring (DGR, personal

observation).

We monitored house wren breeding success at 180 nest boxes divided between six

locations in Delaware and Union Counties, Ohio, USA (Table S1, Supplementary Materials)

during the 2020 and 2021 breeding seasons. The nest boxes varied slightly in dimensions due

to landowner preferences, but the average internal dimensions were 110 x 115 x 195 mm and

the entrance hole size was 38 mm. Each nest box was mounted on a conduit pipe

approximately 1.75 m above the ground, and the pipe was covered in axle grease to limit

access by predators. The majority of the boxes were located in edge habitat near the transition

from grass or field to forest, but a small subset was located in suburban backyards (*N*=7).

Banding of Adults and Nestlings

We captured male house wrens using mist nets and a brief playback of conspecific song.

Each male was banded with a USFWS aluminum leg band and a unique combination of 2-3

additional colored bands for identification from a distance. We also collected morphological

measurements and a small blood sample for an unrelated study. Banding only males allowed us

to differentiate between the sexes during our behavioral trials (see below), and because most

females were never captured, it also minimized the likelihood that the capture process

disrupted the female's antipredator response. Any females that were captured accidentally

(Experiment 1, $N = 6$ of 38 females; Experiment 2, $N = 10$ of 38 females) were rapidly banded

and immediately released within a few minutes of capture to limit any potential effects of

handling on subsequent behavior. All birds were captured and banded at least 24 hours before

any behavioral trials, and most females that were captured were handled more than one week

before their first trial (Experiment 1, $N = 4/6$; Experiment 2, $N = 5/10$). We assumed that the

small number of females captured, the limited handling time, and the delay after any female

was captured before conducting behavioral trials minimized any effects of unintentional

handling on female responses.

We also monitored the number of eggs laid, hatching success, and offspring development

for every nest box. Nest checks occurred every 3-4 days, and any checks on the day of a

behavioral trial were conducted after the trial concluded to minimize any effect on behavior.

We counted the number of nestlings at the end of each behavioral trial (see below), and banded

nestlings when they were 8 days old. Each nestling received a single USFWS aluminum band,

and we collected morphological measurements and a small blood sample. After banding, we

passively monitored each nest box until all visible and acoustic activity ceased. Unless the

contents of the nest suggested otherwise (i.e., presence of dead nestlings and/or their parts), we

assumed that empty nests successfully fledged.

Experiment 1: Chipmunk v. Snake Decoy

In 2020, we conducted behavioral trials between June 9 and July 24 at 47 nest boxes

containing nestlings between the ages of 3-7 days old. A smaller age range was impossible due

to the distances between our sites and other field-related challenges (e.g., weather). Two life-

like rubber predator decoys, an eastern chipmunk (*Tamias striatus*; Safari Ltd.) and an eastern

ratsnake (*Pantherophis alleghaniensis*; Flybanboo), were used to elicit nest defense behavior.

We chose these predators because both species are common at our study sites and had been

previously observed either depredating nests (ratsnake) or being actively attacked by house

wrens near nests (both predators; personal observations). This comparison allowed us to

empirically test the prediction that wrens should respond equally to predators of similar risk

despite the fact that the two predators are morphologically and ecologically distinct.

To initiate a trial, we waited for each female to enter or land on the nest box before

flushing them away during setup. One predator was placed on top of the nest box, and then on

the following day the other predator was presented in a counterbalanced order to control for

order effects. Although these stimuli are terrestrial predators, both are capable of climbing

nearby vegetation and approaching the nest box from above, making their appearance on top of

the box biologically relevant.

All trials occurred between 7:00-13:00, but we were unable to conduct trials exactly 24

hours apart due to the distant locations of our field sites and unpredictability in how rapidly

females would return to their boxes. Trials were recorded with a video camera mounted on a

tripod placed roughly 2-3 meters away from the nest box to later confirm behaviors and

differentiate between the male and female wren in cases of uncertainty. We positioned the

camera in tall grass or other vegetation to minimize its disruption to the focal birds, and

observers maintained a distance of roughly 15 m away from the nest box. Each decoy was

attached to a fishing line, and at the completion of the trial we used a fishing rod to rapidly reel

the decoy away from the box. This approach eliminated the need for the observer to approach

the box and remove the decoy. We allowed the video camera to continue to record for an

additional 20 minutes to measure the amount of time it took for the female to re-enter the nest

box after removal of the decoy (N=36). Post-trial data for some of the females were lost due to

technical issues or user error.

We quantified the nest defense behavior of female house wrens for a total of seven

minutes after the decoy was placed on the nest box. We counted the number of hits, defined as

any contact the female made with the decoy, and the number of flyovers, defined as flights by

the female over the decoy. We also measured the amount of time females spent within five

meters of the nest box, the amount of time that females spent physically on or in the nest box,

and the female's closest approach (m) to the decoy. The five-meter threshold was chosen

because all boxes had potential perches within five meters of the nest box despite variation in

vegetative cover among our sites. Any female that hit the decoy was given a closest approach

value of zero meters. We also noted whether or not the female produced alarm calls, and

whether the female's mate was present during the trial. Neighboring conspecifics and

heterospecifics sometimes responded to the predator decoys by approaching, alarm calling, and

in rare cases flying over and hitting the decoy. During each trial, we noted the number of

individuals of each species that approached within 5 meters of the decoy. To avoid counting

incidental birds that were not responding to the predator, we only counted individuals that

either oriented towards the decoy or produced alarm calls.

Behavioral data from both predator trials for an individual female were excluded from

our final analysis if either of the decoys were knocked off the box during either trial (N=3 of 47

total trials; 6.4%). However, if the chipmunk decoy was knocked over but remained on the

box, the trials were not excluded (N=5 of 47 trials; 10.6%). We also excluded trials if the

female did not return to the box within seven minutes of initiating the trial (*N*=6 of 47 trials,

12.8%). In response to this limitation, we adjusted our protocol early in the season so that we

removed the decoy and reset the trial if the female did not return within 4 minutes after the trial

was initiated. Once the female returned to the box, we flushed her away, placed the predator

decoy, and started the seven-minute observation again. In summary, nine trials (of 47 trials;

19.1%) were excluded and our final sample size was 38 paired behavioral trials.

Experiment 2: Hawk v. Snake Decoy

In 2021 between June 6 and July 30, we replicated this experiment in the same house

wren population, but we replaced the chipmunk decoy with a taxidermied, juvenile Cooper's

hawk (*Accipiter cooperii*), which is commonly observed at our study sites and elicits alarm

calls from passerines and small mammals (DGR, personal observation). Although Cooper's

hawks primarily forage on small or medium-sized adult songbirds and mammals, they have

been frequently observed capturing nestlings from both cavity nests in bird boxes and open-cup

nests in the forest canopy (Milsap et al. 2013, Rosenfield et al. 2020). Thus, the comparison of

the response to the Cooper's hawk versus the ratsnake allowed us to test the prediction that

wrens should respond less aggressively to the higher risk predator of adults (hawk). The

taxidermied hawk was mounted with wire onto a wooden plank, and two large binder clips

were attached to the bottom of the plank with jewelry wire. This design allowed the decoy to

be quickly attached and removed from the roof of the nest box while eliminating the possibility

that it could be knocked off the box.

We conducted behavioral trials at 41 nest boxes with nestlings aged 2-7 days. A different

observer (JHF) scored the trials in 2021, but she was trained by the observer (RCE) from 2020

to ensure consistency between years. All of our sampling methods were replicated from

experiment 1, but we were unable to continue to use the fishing rod and line to rapidly remove

the decoys. As a result, one of the observers quickly ran to the box and removed the decoy at

the end of each seven-minute trial. The video camera continued to record for an additional 20

minutes (N=38), but the approach of the observer may have affected the return to normal

behavior. The snake decoy was knocked off the nest box during three trials (of 41 total; 7.3%),

and we excluded both the snake and hawk trials from those subjects in our final analysis for a

total sample of 38 paired behavioral trials.

Note on Experimental Design

Our decision to split this study into two separate experiments rather than one experiment

including all three predator stimuli was driven by uncertainty stemming from the global Covid-

19 pandemic. This event shutdown numerous research labs in 2020 and disruptions continued

into 2021. The potential for an unpredictable and abbreviated 2020 field season led us to err on

the side of a less complicated protocol to maximize our sample size. We used the STRANGE

framework to identify and address potential issues in sampling methodology in order to help

prevent misinterpretation of results due to sampling biases (see Supp. Materials; Webster and

Rutz 2020).

An additional limitation of this design was that we did not include a non-predatory

control in either experiment (e.g., a heterospecific passerine that would pose no threat to a

house wren nest). Thus, the behaviors we observed could be typical of a generalized nest

defense response rather than a specifically "antipredator" response. We think this interpretation

is unlikely for a few reasons. First, in a subsequent experiment, we compared the response of

female wrens in our population to a novel object (car dice) placed on their nest box versus the

ratsnake decoy. Female wrens responded significantly more aggressively to the snake and were

significantly less likely to produce alarm calls when the novel object was present (DGR,

unpublished data). This result suggests that female wrens in our population do not react to

predatory and non-predatory stimuli with similar behaviors. Second, similar nest defense

experiments in southern house wrens, a closely-related species, have shown that both male and

female wrens respond less aggressively to non-predatory control birds at their nest (Fasanella

and Fernández 2009, Duré Ruiz et al. 2017). Finally, a study in a different population of

northern house wrens found that parents decreased their provisioning more in response to a

simulated nest predator than a non-predatory control bird (Ghalambor et al. 2013). Although

this study did not measure nest defense, it is consistent with wrens behaving differently around

predators relative to non-predators near the nest.

Statistical analysis

For each experiment, we generated either linear (LMM) or generalized linear (GLMM)

mixed-effects models in R version 4.4.2 (R Core Team 2024; packages: "lme4," "car,"

"lmerTest") to test whether female behaviors including flyovers, hits, closest approach to the

decoy, time spent within five meters of the decoy, the production of alarm calls, and latency to

re-enter the nest box after removal of the predator differed between predator types.

Measurements of time spent on or inside the box were excluded due to extremely low rates of

occurrence in both experiments (Table S2). Each model included female behavior as the

dependent variable with predator type, treatment order, and their interaction as fixed effects.

Day of the year and male presence (yes or no) were included as covariates and female ID was

included as a random effect to control for seasonal shifts in behavior, the presence of the

female's mate, and repeated sampling of the same individuals. We did not include location as a

random effect in our models because all of our nest boxes are located in similar edge habitats

and the sample size for each individual site is small (N<10), which substantially reduced our

statistical power to interpret any potential differences. We applied an alpha value of 0.05 for all

analyses.

The number of hits and flyovers as well as the closest approach to the decoy were not

normally distributed owing to a large amount of zeros. Because transformations could not

resolve this issue, we used GLMMs that assumed a Poisson distribution to analyze those

behaviors. Closest approach scores were multiplied by 100 before the analysis to generate

integers, which are required for the Poisson distribution. Time within 5m of the decoy was

analyzed with LMMs that assumed a Gaussian distribution. Although the model residuals

marginally failed a Shapiro-Wilk test of normality (package: "stats"), we visually inspected the

residuals to confirm that they approximated a normal distribution. An abundance of evidence

has indicated that LMMs are robust to violations of this assumption, except in cases where the

sample sizes are prohibitively small or the distributions are bimodal, neither of which occurred

here (Schielzeth et al. 2020; Knief and Forstmeier 2021; and further citations within). Time to

return to the nest was also analyzed with LMMs that assumed a Gaussian distribution, but we

square root transformed the data to achieve normality of the residuals based on Shapiro-Wilk

tests and visual inspection of the residuals. The same transformation of the time within 5 m

data made the residuals less normal, which is why we did not transform the data for that

behavior. Because alarm calls were scored as a categorical variable (yes or no), we used

GLMMs that assumed a binomial distribution.

To further evaluate model assumptions across our analyses, we tested for

homoscedasticity using Levene's test (package: "car"). Homoscedasticity was met for all

models in experiment 1 but none of the models in experiment 2. These violations were

seemingly due to the uniformly weak responses elicited by the hawk decoy in experiment 2.

Although these violations likely had minimal effects on the validity of the models (Schielzeth

et al. 2020; Knief and Forstmeier 2021), we ran a separate series of nonparametric Wilcoxon

tests (package: "stats") that lack the assumptions of normality and homoscedasticity to confirm

the results of the mixed-effects models. For each behavior, we split the data based on treatment

order and analyzed responses to each predator for the first trial separately from responses to the

predators for the second trial. This approach allowed us to isolate the fixed effect of interest

(predator type) from the fixed effect of order and the random effect of female identity. The

qualitative results of these nonparametric tests were nearly identical to the mixed-effects

models (Tables S5-6).

We also used nonparametric Wilcoxon tests to determine whether the number of

neighboring conspecific and heterospecific individuals and the number of heterospecific

species that responded to the trials differed between predator types in both years. This

nonparametric test was chosen due to the lack of normality and low variance in these datasets

caused by a large proportion of zero or near zero values. We considered applying a GLMM

with a binomial distribution, but decided against this approach because it did not allow us to

analyze both the total number of heterospecifics and the number of heterospecific species that

responded because both reduced data sets (i.e., presence or absence of heterospecifics) would

have been identical.

Ethical Note

All procedures were reviewed and approved by the Ohio Wesleyan University

Institutional Animal Care and Use Committee (Protocol Numbers: Spring_2020-21-

01_Reichard and 05-2021-06), U.S. Federal Bird Banding Lab (Permit Number: 24035), and

State of Ohio Department of Natural Resources (Permit Numbers: 23-009, 23-010). To

minimize stress in the free-living birds used in this study, we attempted to capture and mark

only the male of each pair, which was the minimum required to facilitate individual

identification during our behavioral trials. All birds were immediately removed from mist nets

after capture, processed rapidly to limit handling time, and then safely released at the capture

site. No adverse effects were observed from these widely established techniques. Although our

behavioral trials involved the observation of a simulated predator at the nest box, all subjects

returned to normal parental care after the completion of the trial and no nests were abandoned

as a result of these experiments.

Results

Experiment 1: Chipmunk v. Snake Decoy

Female wrens exhibited no significant differences in the number of hits (χ^2_1=1.76,

P=0.18), number of flyovers ($\chi^2_{1=2.34}$, *P*=0.13), time spent within 5 meters of the nest box

($F_{1,39.1}$=2.68, *P*=0.11), or in their closest approach (χ^2_1=2.46, *P*=0.12) to the snake and

chipmunk decoys (Figure 1A-D). There was also no difference in the number of females (36 of

38 total for both predators; 94.7%) that produced alarm calls in response to each predator

(χ^2_1<0.001, *P*>0.98). After the predator decoys were removed, we found no significant

difference between predator types in the amount of time females took to re-enter the nest box

($F_{1,32.6}$=0.68, P=0.42; Figure 2A).

Day of the year had a significant effect on the number of hits (χ^2_1=8.42, P=0.003) with

females hitting the decoy less often later in the breeding season. Similarly, females took longer

to return to the nest box later in the season ($F_{1,27}$=8.87, P=0.006). There was no detectable

effect of date on any other behaviors ($P \geq 0.46$ in all cases; Table S3). The presence of the

female's mate had a significant effect on the time spent within 5 m of the decoy ($F_{1,34}$=4.20,

P=0.048) and how quickly the female returned to the nest (χ^2_1=7.86, P=0.009). When their

mate was present, females spent more time within 5 m of the decoy and they returned to the

nest faster. There was no detectable effect of male presence on any other behaviors ($P \geq 0.13$ in

all cases; Table S3). For all six behaviors, we found no evidence for significant effects of

treatment order ($P \geq 0.055$ in all cases; Table S3) or significant interactions between treatment

order and predator type ($P \geq 0.13$ in all cases; Table S1). In one exception, we noted a marginal

effect of treatment order on the closest approach to the decoy with females tending to approach

more closely during the first trial (χ^2_1=3.67, P=0.055).

Figure 1 — Responses of female house wrens to a snake or chipmunk decoy on top of their nest

box during a seven-minute trial. There were no significant differences in the number of (A) hits

(P=0.18), (B) flyovers (P=0.12), (C) how closely females approached each model (P=0.12), or

(D) how much time females spent within 5 m of each model (P=0.11). The thick center line

represents the median, and the box encloses the interquartile range. Whiskers show the range

of data within 1.5 times the interquartile range, and dots are data points exceeding that range.

415
416
417

Figure 2 — No detectable differences in how long it took female house wrens to re-enter the

nest box after encountering a snake decoy versus a chipmunk decoy (P=0.42) or a hawk decoy

(P=0.41). Box plots as in Fig. 1.

423   The number of conspecific ($W=724.5$, $P=0.98$) and heterospecific individuals ($W=693.5$,

424    $P=0.74$) and the number of heterospecific species ($W=700.5$, $P=0.80$) that responded to the

425   predator decoys did not differ detectably between the two predator types.

429 Table 1. Mean ± standard deviation (range) for the number of conspecific and heterospecific

430 individuals and heterospecific species that responded to the predator stimuli in Experiment 1.

Observation	Snake	Chipmunk
Conspecific Individuals	0.53 ± 1.0 (0-5)	0.47 ± 0.8 (0-3)
Heterospecific Individuals	1.08 ± 2.7 (0-15)	0.76 ± 1.4 (0-6)
Heterospecific Species	0.61 ± 1.0 (0-5)	0.58 ± 0.9 (0-4)

Experiment 2: Hawk v. Snake Decoy

Similar to Experiment 1, female wrens frequently performed hits and flyovers while

defending against the snake decoy, with 25 out of 41 females (61.0%) hitting the snake model

at least once and 38 out of 41 females (92.7%) flying over at least once. Notably, zero of the

female wrens performed hits or flyovers while defending against the hawk decoy. Our GLMM

approach could not assess statistical significance with this data structure (i.e. no variation in

response to the hawk), but our nonparametric analysis confirmed that females hit and flew over

the snake significantly more often than the hawk ($P<0.001$ in both cases, Table S6; Figure 3A-

B). Females approached significantly closer to the snake ($\chi^2_1=21.8$, $P < 0.001$; Figure 3C), but

the amount of time spent within 5 meters was not significantly different from the response to

the hawk ($F_{1,39.6}=1.28$, $P=0.27$; Figure 3D). There was also no difference in the number of

females (34 of 38 total for both predators; 89.5%) that produced alarm calls in response to each

predator ($\chi^2_1=0.10$, $P=0.75$). After the removal of the predator decoys, female wrens did not

differ detectably in the time to re-enter the nest box after the predator was removed

($F_{1,37.5}=0.70$, $P=0.41$; Figure 2B). Finally, we found no evidence for an effect of date ($P\geq0.09$

in all cases; Table S4), the presence of the female's mate (*P*>0.27 in all cases; Table S4),

treatment order (*P*>0.75 in all cases; Table S4) or a predator type by treatment order interaction

(*P*>0.50 in all cases; Table S4) for any behaviors.

Figure 3 — Responses of female house wrens to a snake or hawk decoy on top of their nest box

during a seven-minute trial. Female house wrens performed significantly more (A) hits

(P<0.001) and (B) flyovers (P<0.001) against the snake decoy than against the hawk decoy.

Female wrens also (C) made significantly closer approaches to (P<0.001), but (D) they did not

significantly differ in the time spent within 5 meters of the decoys (P=0.27). Box plots as in

Fig. 1. * *P<0.001.*

Significantly more heterospecific individuals responded to the hawk decoy than to the

snake decoy (*W*=531.5, *P*=0.04), but there was no detectable difference in the number of

heterospecific species that responded to each predator (*W*=599.5, *P*=0.18). In contrast,

significantly more conspecific individuals mobbed the snake decoy than mobbed the hawk

(W=908, *P*=0.01).

Table 2. Mean ± standard deviation (range) for the number of conspecific and heterospecific

individuals and heterospecific species that responded to the predator stimuli in Experiment 2.

Bold text denotes *P*<0.05.

Observation	Snake	Hawk
Conspecific Individuals*	**0.55 ± 0.8** **(0-3)**	**0.13 ± 0.3** **(0-1)**
Heterospecific Individuals*	**0.97 ± 1.1** **(0-4)**	**2.05 ± 2.2** **(0-9)**
Heterospecific Species	0.84 ± 0.9 (0-3)	1.29 ± 1.4 (0-6)

Discussion

Defending offspring from predators can be dangerous, and parents that accurately assess

the risks posed by different predators and scale their responses appropriately should have

higher lifetime fitness (Montgomerie and Weatherhead 1988; Clutton-Brock 1991; Caro 2005).

We found evidence for risk assessment and plastic antipredator behavior in nesting house

wrens. Female wrens responded with similar levels of aggressive nest defense behavior when

encountering a simulated snake or chipmunk at their nest box, both of which posed a higher

risk to offspring than adults. In contrast, the wrens responded significantly less aggressively to

a simulated hawk that posed a higher risk to adults than nestlings. This plasticity in nest

defense behavior is potentially adaptive because parents that avoid high risk predators like the

hawk are more likely to live longer and have higher lifetime fitness despite the potential costs

to the current clutch of offspring (Trivers 1974; Montgomerie and Weatherhead 1988).

Collectively, these results are consistent with the predictions of parental investment theory

(Montgomerie and Weatherhead 1988; Caro 2005), and similar parental risk assessment in

different contexts has been previously observed in house wrens and other species (Lima 2009;

Mahr et al. 2015; Duré Ruiz et al. 2018).

We observed substantial individual variation among females in how strongly they

responded to each of the simulated predators. For example, the most aggressive females hit the

snake and chipmunk decoys over 50 times in a seven-minute period while the least aggressive

females never dove at or hit the decoys. Some of this variation could be explained by parental

investment theory, which predicts a positive relationship between the intensity of nest defense

and various life history traits such as clutch size, nestling age, and adult age and longevity

(Montgomerie and Weatherhead 1988, Caro 2005). However, testing these predictions was not

a main focus of our study; rather, we sought to control for differences in parental investment

across the nestling stage. As a result, we only captured a fraction of the possible variation in

clutch size (80% of pairs defended 4-6 offspring) and nestling age (85% of offspring were 3-6

days post hatch), which limits our ability to draw conclusions about the relationship between

aggression and current reproductive investment. In Experiment 1, we observed that females hit

the predator decoy less and took longer to return to the nest when trials occurred later in the

breeding season. This result is consistent with a redirection of investment away from offspring

towards the individual, which could be adaptive in a species with multiple breeding attempts.

House wrens, however, are short-lived with an average lifespan of less than two years (Johnson

2020), so the benefit, if any, of this shift remains unclear. Alternatively, reduced effort later in

the season could result from a seasonal decline in female condition due the cumulative effects

of multiple breeding attempts (Laiolo et al. 2004, Mitchell et al. 2012). More research is

necessary to evaluate how these energetic tradeoffs might shape the different components of

antipredator behavior, and in turn, affect both adult and offspring survival in house wrens and

other species (Lind and Cresswell 2005).

Two important exceptions to the persistent variation among individuals in nest defense

behavior occurred with (1) the production of alarm calls when any predator decoy was present,

and (2) the behavioral responses to the simulated hawk. In both experiments, females almost

universally produced alarm calls despite varying levels of risk posed by the different predators

to the adults and their nestlings. Alarm calls are a lower risk behavior than the other nest

defense behaviors sampled here, which might explain their consistent production. When

produced near the nest, alarm calls can serve many adaptive functions including distracting the

predator (Fasanella and Fernández 2009), inducing neighboring birds to mob the predator

(Templeton et al. 2005), and alerting the young inside the nest that a predator is nearby (Suzuki

2011). Previous research in another house wren population found that parents produced alarm

calls even in the presence of a non-predatory heterospecific bird at their nest (Fasanella and

Fernández 2009), suggesting that alarm calls might function as a general response to any

disturbance at the nest box. However, wren parents also produced more alarm calls in response

to a potential predator rather than a non-predatory control bird (Fasanella and Fernández 2009)

and more alarm calls were produced as a simulated snake moved closer to the nest (Fernández

and Carro 2022). Although we did not measure whether the quantity of alarm calls differed

between predator types in our study, it seems clear that alarm calls are a consistent and

important component of the antipredator response, and the intensity of these calls might signal

information about the predator type and level of risk posed to the nestlings (Fasanella and

Fernández 2009, Fernández and Carro 2022, Fernández et al. 2023).

When responding to the hawk decoy, females varied in how closely they approached and

how much time they spent within five meters of the hawk, but none of the 41 females ever hit

the hawk or even flew over it. This consistency among individuals indicates that when the

predatory threat surpasses a certain level it is possible to see uniformity in a typically plastic

behavior. Furthermore, the total absence of physical aggression towards the hawk supports the

hypothesis that selection favors risk assessment, likely by acting against high-risk behaviors,

such as closely approaching or physically attacking a dangerous predator (Desrocher et al.

2002; Mahr et al. 2015). However, our observations of the community-level response to the

hawk and other simulated predators show that risk assessment is not uniform across songbird

species.

Neighboring conspecifics and heterospecific songbirds frequently responded to our

simulated predators by approaching and alarm calling, but very few approached closely enough

to attack or fly directly over the decoys, including the snake and chipmunk (RCE, DGR, JHF,

personal observations). This observation is consistent with mobbing behavior, and the reliance

on alarm calls rather than overt aggression likely reflects the reduced threat posed to these

neighbors by a predator that was not near their own offspring (Colombelli-Négrel et al. 2010;

Kryštofková et al. 2011). Similar to the antipredator behavior of the wren parents, the number

of conspecifics and heterospecifics that responded did not differ significantly between the

snake and chipmunk. In contrast, more conspecifics responded to the snake while more

heterospecifics responded to the hawk in our second experiment. This result mirrors the

cautious behavior of house wrens when responding to a hawk at their nest box, which likely

explains the lack of mobbing behavior from conspecific neighbors. In contrast, larger

heterospecific songbirds including tree swallows (*Tachycineta bicolor*), northern cardinals

(*Cardinalis cardinalis*), and blue jays (*Cyanocitta cristata*) appeared in higher numbers when

the hawk was present, and tree swallows frequently dove at the hawk, which was in stark

contrast with all other species (RCE, DGR, JHF, personal observations).

The heterospecific response to the hawk was surprising given that Cooper's hawks

primarily forage for larger songbirds (Roth and Lima 2003; Rosenfield et al. 2020), which

elevated the risk of predation for those that approached. Although some heterospecifics were

likely defending nearby nests from the hawk (e.g., tree swallows from nearby boxes with active

nests), others may have been gathering information through eavesdropping behavior, which

can be used to inform future decisions related to predation threats (Magrath et al. 2015). House

wrens, as an insectivorous species that must be highly vigilant in search of prey, may be better

able to detect predators than less vigilant species or those that spend more time in denser

substrate, thereby providing more accurate spatial information for eavesdropping

heterospecifics (Magrath et al. 2015; Johnson, 2020). Future experiments could examine the

ecological relationship between defensive cooperation, social learning, and information flow

through social networks of different avian communities.

Reactive nest defense behavior can be subdivided into (1) behaviors that distract or

redirect the predator towards the signaler and (2) behaviors that involve acts of aggression

towards the predator (Caro 2005; Lima 2009). Smaller organisms like house wrens may rely on

distraction rather than attacking predators above a certain size because they lack the body size

or armaments necessary to successfully drive the predator away (Montgomerie and

Weatherhead 1988; Winkler 1992). If the likelihood of physically deterring the predator is low,

then any amount of risk to the parent from attacking the predator could justify a distraction-

based antipredator response or even no response at all. However, it is unlikely that size

thresholds universally govern antipredator behavior given that certain species of limited size

and armaments frequently attack larger predators, including humans (Levey et al. 2009,

Lawson et al. 2021). In addition to size differences, birds of prey are substantially more mobile

in the air than terrestrial predators, which further elevates the risks associated with physical

aggression and could explain distraction-based responses regardless of an avian predator's size

(Mahr et al. 2015). More broadly, risk-taking thresholds and antipredator behavior can vary

according to other factors such as predator abundance, resource availability, or competition as

adult birds employ strategies that balance the reproduction-survival tradeoff (Díaz et al., 2013;

Mikula et al., 2018). Future work should continue to focus on elucidating the proximate and

ultimate causes of persistent intra- and interspecific variation in parental antipredator behavior

and risk assessment.

Data Availability Statement

The data collected and analyzed to support these findings are freely available in the

Mendeley Data Repository at DOI:10.17632/zjb5ynz5n4.2.

Acknowledgements

We thank Ohio Wesleyan University, the Big Walnut School System, and the Davis,

Koban, and Fink families for access to facilities and field sites. We thank Holly Keating for

analyzing videos, Tamaki Yuri for preparing the taxidermied Cooper's Hawk, Lily Bonar for

field assistance, Chandler Carr for measuring our nest boxes, and Eric Gangloff for advice on

statistics. This project was funded by the Ohio Wesleyan University Summer Science Research

Program.

Literature cited

Caro T. 2005. Antipredator Defenses in Birds and Mammals. University of Chicago Press.

Carlson NV, Griesser G. 2022. Mobbing in animals: A thorough review and proposed future

directions. Advances in the Study of Behavior. 54:1-41.

doi.org/10.1016/bs.asb.2022.01.003.

Clutton-Brock TH. 1991. The Evolution of Parental Care. Princeton University Press.

Colombelli-Négrel D, Robertson J, Kleindorfer S, Sulloway F. 2010. Extended parental care of

fledglings: parent birds adjust anti-predator response according to predator type and

distance. Behaviour. 147(7):853–870. doi:10.1163/000579510X495771.

Corral MG, Llambías PE, Fernández GJ. 2013. Effect of conspecific alarm calls in the parental

behaviour of nesting southern house wrens. Acta Ethol. 16(1):47–51.

doi:10.1007/s10211-012-0129-x.

Desrochers A, Bélisle M, Bourque J. 2002. Do mobbing calls affect the perception of risk by

forest birds?. Anim Behav. 64:709-714. doi:10.1006/anbe.2002.4013.

Díaz M, Møller AP, Flensted-Jensen E, Grim T, Ibáñez-Álamo JD, Jokimäki J, Markó G,

Tryjanowski, P. 2013. The Geography of Fear: A Latitudinal Gradient in Anti-Predator

Escape Distances of Birds across Europe. PLOS ONE. 8(5):e64634.

https://doi.org/10.1371/journal.pone.0064634

Dudeck BP, Clinchy M, Allen MC, Zanette LY. 2018. Fear affects parental care, which

predicts juvenile survival and exacerbates the total cost of fear on demography. Ecology.

99(1):127–135. doi:10.1002/ecy.2050.

Dulac C, O'Connell LA, Wu Z. 2014. Neural control of maternal and paternal behaviors.

Science. 345(6198):765–770. doi:10.1126/science.1253291.

Duré Ruiz NM, Fasanella M, Fernández GJ. 2018. Breeding southern house wrens exhibit a

threat-sensitive response when exposed to different predator models. J Ethol. 36(1):43–

53. doi:10.1007/s10164-017-0528-6.

Fasanella M, Fernández, GJ. 2009. Alarm calls of the Southern House Wren *Troglodytes*

musculus: variation with nesting stage and predator model. Journal of Ornithology

150:853–863.

Fernández GJ, Carro ME. 2022. Alarm calls of southern house wrens, *Troglodytes aedon*

bonariae, convey information about the level of risk. Ethology. 128(4):293–302.

doi:10.1111/eth.13255.

Fernández GJ, Corral MG, Llambías PE. 2015. Sexual differences in risk-taking behaviour of

the Southern house wren during the nestling rearing stage. Acta Ethol. 18(1):11–18.

doi:10.1007/s10211-013-0177-x.

Fernández GJ, Dutour M, Carro ME. 2023. Information transfer during mobbing: call rate is

more important than the number of callers in a southern temperate passerine. Behav Ecol

Sociobiol. 77:78. doi.org/10.1007/s00265-023-03357-z.

Fernández GJ, Llambías PE. 2013. Parental risk-taking behaviour and nest defence during the

nestling rearing stage in Northern House Wrens *Troglodytes aedon*. Acta Ornith.

48(1):55–63. doi:10.3161/000164513X670016.

Fontaine JJ, Martin TE. 2006. Parent birds assess nest predation risk and adjust their

reproductive strategies. Ecology Letters. 9(4):428–434. doi:10.1111/j.1461-

0248.2006.00892.x.

Forsman JT, Mönkkönen M. 2001. Responses by breeding birds to heterospecific song and

mobbing call playbacks under varying predation risk. Anim Behav. 62(6): 1067-1073.

doi.org/10.1006/anbe.2001.1856

de Framond L, Brumm H, Thompson WI, Drabing SM, Francis CD. 2022. The broken-wing

display across birds and the conditions for its evolution. Proc R Soc B: Biol Sci.

289(1971):20220058. doi:10.1098/rspb.2022.0058.

Freed LA. 1981. Breeding ecology of House Wrens: new views of avian life history

phenomena [PhD Thesis]. [Ames, Iowa]: University of Iowa.

Ghalambor CK, Martin TE. 2001. Fecundity-survival trade-offs and parental risk-taking in

birds. Science. 292(5516):494–497. doi:10.1126/science.1059379.

Ghalambor CK, Peluc SI, Martin TE. 2013. Plasticity of parental care under the risk of

predation: how much should parents reduce care? Biology Letters. 9(4):20130154.

doi:10.1098/rsbl.2013.0154.

Gottfried BM. 1979. Anti-predator aggression in birds nesting in old field habitats: an

experimental analysis. The Condor. 81(3):251–257. doi:10.2307/1367626.

Ibáñez-Álamo JD, Magrath RD, Oteyza JC, Chalfoun AD, Haff TM, Schmidt KA, Thomson

RL, Martin TE. 2015. Nest predation research: recent findings and future perspectives. J

Ornithol. 156(1):247–262. doi:10.1007/s10336-015-1207-4.

Johnson LS. 2020. House Wren (*Troglodytes aedon*), version 1.0. In: Poole AF, editor. Birds

of the World. Cornell Lab of Ornithology, Ithaca, NY, USA.

Knief U, Forstmeier W. 2021. Violating the normality assumption may be the lesser of two

evils. Behav Res Methods 53, 2576–2590.

Kryštofková M, Haas M, Exnerová A. 2011. Nest defense in Blackbirds *Turdus merula*: effect

of predator distance and parental sex. Acta Ornith. 46(1):53–61.

doi:10.3161/000164511X589938.

Laiolo P, Tella JL, Carrete M, Serrano D, López G. 2004. Distress calls may honestly signal

bird quality to predators. Proc. R. Soc. Lond. B. 271S513–S515.

http://doi.org/10.1098/rsbl.2004.0239

Lawson SL, Enos JK, Antonson ND, Gill SA, Hauber ME. 2021. Do hosts of avian brood

parasites discriminate parasitic vs. predatory threats? A meta-analysis. Adv Study Behav.

53:63-95.

Levey DJ, Londoño GA, Ungvari-Martin J, Hiersoux MR, Jankowski JE, Poulsen JR, Stracey

CM, Robinson SK. 2009. Urban mockingbirds quickly learn to identify individual

humans. PNAS. 106(22):8959–8962. doi:10.1073/pnas.0811422106.

Lind J, Cresswell W, Determining the fitness consequences of antipredation behavior. 2005.

Behav Ecol. 16(5):945–956. doi:10.1093/beheco/ari075

Lima SL. 2009. Predators and the breeding bird: behavioral and reproductive flexibility under

the risk of predation. Biol Rev. 84(3):485–513. doi:10.1111/j.1469-185X.2009.00085.x.

Magrath RD, Haff TM, Fallow PM, Radford AN. 2015. Eavesdropping on heterospecific alarm

calls: From mechanisms to consequences. Biol Rev. 90(2):560–586.

https://doi.org/10.1111/brv.12122

Mahr K, Riegler G, Hoi H. 2015. Parental risk management in relation to offspring defence:

bad news for kids. Proc R Soc B: Biol Sci. 282(1798):20141670.

doi:10.1098/rspb.2014.1670.

Martin TE, Briskie JV. 2009. Predation on dependent offspring: a review of the consequences

for mean expression and phenotypic plasticity in avian life history traits. Ann N Y Acad

Sci. 1168:201–217. doi:10.1111/j.1749-6632.2009.04577.x.

Mikula P, Díaz M, Albrecht T, Jokimäki J, Kaisanlahti-Jokimäki ML, Kroitero G, Møller AP,

Tryjanowski P, Yosef R, Hromada M. 2018. Adjusting risk-taking to the annual cycle of

long-distance migratory birds. Sci Reports. 8(1): 13989. https://doi.org/10.1038/s41598-

018-32252-1

Milsap, BA, Breen, TF, Phillips, LM. 2013. Ecology of the Cooper's Hawk in North Florida.

North American Fauna 78: 1–58.

Mitchell GW, Wheelwright NT, Guglielmo CG, Norris DR. 2012. Short- and long-term costs

of reproduction in a migratory songbird. Ibis. 154(2):325-337.

Montgomerie RD, Weatherhead PJ. 1988. Risks and rewards of nest defence by parent birds. Q

Rev Biol. 63(2):167–187. doi:10.1086/415838.

Morosinotto C, Thomson RL, Korpimäki E. 2010. Habitat selection as an antipredator

behaviour in a multi-predator landscape: all enemies are not equal. J Anim Ecol.

79(2):327–333. doi:10.1111/j.1365-2656.2009.01638.x.

R Core Team. 2024. R: A language and environment for statistical computing. https://www.R-

project.org.

Reidy JL, Stake MM, Thompson FR. 2009. Nocturnal predation of females on nests: An

important source of mortality for golden-cheeked warblers? Wilson J Ornith. 121(2):416–

421. https://doi.org/10.1676/08-076.1

Ricklefs RE. 1977. On the evolution of reproductive strategies in birds: reproductive effort. Am Nat. 111(979):453–478.

Rosenfield RN, Madden KK, Bielefeldt J, Curtis OE. 2020. Cooper's Hawk (*Accipiter cooperii*), version 1.0. Birds of the World. doi:10.2173/bow.coohaw.01.

Roth TC II, Lima SL. 2003. Hunting behavior and diet of cooper's hawks: an urban view of the small-bird-in-winter paradigm. The Condor. 105(3):474–483. doi:10.1093/condor/105.3.474.

Schielzeth, H, Dingemanse, NJ, Nakagawa, S, et al. 2020. Robustness of linear mixed-effects models to violations of distributional assumptions. Methods Ecol Evol 11: 1141– 1152.

Scott M, Robinson WD. 2023. Mobbing behavior of songbirds in response to calls of an ambush-predator, the Northern Pygmy-Owl. Front. Ecol. Evol. 11:1092323. doi: 10.3389/fevo.2023.1092323.

Spaans B, Blijleven HJ, Popov I, Rykhlikova M, Ebbinge B. 1998. Dark-bellied Brent Geese *Branta bernicla bernicla* forego breeding when Arctic Foxes Alopex lagopus are present during nest initiation. Ardea. 86:11–20.

Suzuki, TN. 2011. Parental alarm calls warn nestlings about different predatory threats. Current Biology. 21:R15-R16.

Szymkowiak J, Thomson RL. 2019. Nest predator avoidance during habitat selection of a songbird varies with mast peaks and troughs. Behav Ecol Sociobiol. 73(7):91. doi:10.1007/s00265-019-2702-z.

Templeton CN, Greene E, Davis K. 2005. Allometry of alarm calls: black-capped chickadees

728   encode information about predator size. Science 308: 1934-1937.

729 Teunissen N, Kingma SA, Peters A. 2020. Predator defense is shaped by risk, brood value and

730   social group benefits in a cooperative breeder. Behav Ecol. 31(3):761–771.

731   doi:10.1093/beheco/araa012.

732 Travers M, Clinchy M, Zanette L, Boonstra R, Williams TD. 2010. Indirect predator effects on

733   clutch size and the cost of egg production. Ecology Letters. 13(8):980–988.

734   doi:10.1111/j.1461-0248.2010.01488.x.

735 Trivers RL. 1974. Parent-Offspring Conflict. American Zoologist. 14(1):249–264.

736   doi:10.1093/icb/14.1.249.

737 Webster MM, Rutz C. 2020. How STRANGE are your study animals? Nature 582, 337–340.

738 Winkler DW. 1992. Causes and Consequences of Variation in Parental Defense Behavior by

739   Tree Swallows. The Condor. 94(2):502–520. doi:10.2307/1369222.

Supplementary Materials

Statement on STRANGE framework

Studies of animal behavior can be misinterpreted due to sampling biases, and the

STRANGE framework provides a method for identifying these potential issues (Webster and

Rutz 2020). All subjects in this study were free-living songbirds, and we included all

individuals that nested on our study sites regardless of age or nesting location, making it

unlikely that social background biased our results. As detailed above, a small number of

individuals were excluded, namely those that failed to respond within 7 minutes during

Experiment 1 before our protocol was adjusted to be more inclusive or those that responded so

forcefully that they knocked the predator decoy from the box. Thus, we may have failed to

sample a subset of both low and high responders. Although this approach may have biased our

sample towards intermediate responders, we think this potential bias is unlikely to have skewed

our results for a few reasons. First, our sample size is large for a field study and it includes

individuals that responded both aggressively and weakly to the predator decoys, so it is

unlikely that we failed to sample the extremes of behavioral variation in this population.

Second, we conducted repeated-measures sampling, which required subjects to be excluded

from all treatments rather than only those trials where we were unable to collect data. This

approach limits the likelihood of bias as each individual effectively serves as its own control.

With respect to seasonal changes in behavior, we sampled evenly across the entire breeding

season and included measurement date as a covariate in our models to test for this effect.

Because each subject experienced two sampling events it is possible that habituation affected

responses to the second treatment. We limited this effect by presenting trials on separate days

in a counterbalanced order and included treatment order in our models to test for an effect of

habituation.

It is also possible that carryover effects caused by sampling the same individuals between

years could bias the behavioral responses in Experiment 2. We think that this outcome is

unlikely given the length of time between the two experiments, but if the same females were

sampled in both years carryover effects remain possible. It is difficult to determine how many

females were sampled in both experiments because only 13 of 47 females (27.7%) were

banded in experiment one, including six females banded in previous breeding seasons. Of those

banded females, only one (7.7.% return rate) was definitely sampled in experiment two. It is

likely that some of the unbanded females also returned to be sampled again, and some

inferences can be drawn from males, which were banded at much higher rates. In experiment

one, 43 of 47 males (91.5%) were banded and eight males returned to be sampled again in

experiment two (18.5% return rate). Based on these data, it seems likely that 10-20% of

females were sampled in both experiments.

798   *Table S1 - Locations of field sites and number of nest boxes present.*

Site	Location	Number of Boxes
Big Walnut Elementary School, Sunbury, Ohio	40°14'02.6"N 82°51'53.1"W	15
Big Walnut High School, Sunbury, Ohio	40°14'00.9"N 82°51'37.3"W	41
Davis Family Property, Sunbury, Ohio	40°15'50.2"N 82°49'52.8"W	36
Fink Family Property, Marysville, Ohio	40°16'42.4"N 83°15'43.3"W	21
Kraus Research Preserve, Delaware, Ohio	40°11'45.8"N 83°03'47.1"W	42
Ohio Wesleyan University and nearby homes, Delaware, Ohio	40°17'47.7"N 83°03'53.8"W	25

800   *Table S2 - Summary of the number of trials that include an observation of time spent in or on*
801   *the next box. Experiment 1 tested responses to a simulated snake and chipmunk. Experiment 2*
802   *tested responses to a simulated snake and hawk.*

Experiment	Time spent in the nest box	Time spent on the nest box
1	4/76 (5.3%)	12/76 (15.8%)[1]
2	4/76 (5.3%)	10/76 (13.2%)[2]

803   [1] 9 of 12 observations were for 10 seconds or less (75%)
804   [2] 7 of 10 observations were for 10 seconds or less (70%)

815 *Table S3 – Analysis of model fixed effects and covariates for Experiment 1. *P<0.05*

	Day of the Year		Male Presence		Order		Interaction with Order	
Behavior	X^2 or F	P	X^2 or F	P	X^2 or F	P	X^2 or F	P
Hits	8.42	0.004*	0.38	0.54	2.11	0.15	1.87	0.17
Flyovers	0.02	0.88	2.09	0.15	2.11	0.15	2.34	0.13
Time within 5 m	0.33	0.57	4.20	0.048*	0.004	0.95	2.30	0.14
Closest Approach	0.55	0.46	2.25	0.13	3.67	0.055	1.91	0.17
Latency to Return	8.87	0.006*	7.86	0.009*	1.59	0.23	0.67	0.42
Alarm Calls	0.18	0.67	1.01	0.32	0.10	0.76	<0.001	0.99

816
817
818 *Table S4 – Analysis of model fixed effects and covariates for Experiment 2.*

	Day of the Year		Male Presence		Order		Interaction with Order	
Behavior	*X^2 or F*	*P*	*X^2 or F*	*P*	*X^2 or F*	*P*	*X^2 or F*	*P*
Time within 5 m	*2.16*	*0.15*	*0.54*	*0.47*	*0.02*	*0.89*	*0.47*	*0.50*
Closest Approach	*2.83*	*0.09*	*1.22*	*0.27*	*0.006*	*0.94*	*0.22*	*0.64*
Latency to Return	*1.06*	*0.31*	*0.10*	*0.75*	*<0.001*	*0.98*	*0.001*	*0.97*
Alarm Calls	*0.13*	*0.71*	*0.07*	*0.80*	*0.10*	*0.75*	*0.11*	*074*

819
820
821

824 *Table S5 - Results of nonparametric analysis of Experiment 1 behavior.*

	First Trial		Second Trial	
Behavior	W	P	W	P
Hits	200	0.41	118	0.09
Flyovers	220.5	0.15	126.5	0.17
Time within 5 m	204	0.35	168	0.91
Closest Approach	147	0.39	210.5	0.23
Latency to Return	116	0.82	109	0.98

827 *Table S6 - Results of nonparametric analysis of Experiment 2 behavior. *P<0.05.*

	First Trial		Second Trial	
Behavior	W	P	W	P
Hits	280	<0.001*	288	<0.001*
Flyovers	350	<0.001*	351	<0.001*
Time within 5 m	223	0.21	170	0.78
Closest Approach	4	<0.001*	8.5	<0.001*
Latency to Return	92	0.048*	111.5	0.19