Risk assessment during nest defense against three simulated predators by female

northern house wrens (*Troglodytes aedon aedon*)

Ross C. Eggleston[1,‡], Josephina H. Fornara[1,2,‡], Kyle M. Davis[1,3], Jess Dong[1,4], and Dustin G.

Reichard[1,*]

[1]Ohio Wesleyan University, Department of Zoology, 61 S Sandusky St, Delaware, OH 40315

[2]Current address: Indiana University, Department of Biology, 1001 E Third St, Bloomington,

Indiana 47405

[3]Current address: Michigan State University, Department of Integrative Biology, 288 Farm Ln,

East Lansing, MI 48824

[4]Current address: Ohio State University, School of Environment and Natural Resources, 2021

Coffey Rd, Columbus, OH 43210

*Corresponding author email: dgreicha@owu.edu

[‡]Equal author contribution

Abstract

Offspring predation is one of the greatest obstacles to an organism's reproductive success, but parents vary in the strength of their response to potential predators. One explanation for this variable investment is that defending current offspring has the potential to lower future reproductive success if the predator is also capable of injuring or killing the parent. House wrens (*Troglodytes aedon aedon*) are cavity-nesting songbirds that defend against multiple species of nest predators including small mammals, birds of prey, and snakes. Here, we used three different predator decoys: two nest predators - an eastern chipmunk (*Tamias striatus*) and an eastern ratsnake (*Pantherophis alleghaniensis*) - as well as a predator of both offspring and adults - a juvenile Cooper's hawk (*Accipiter cooperi*) - to elicit nest defense and test whether females use risk assessment to modulate their antipredator behavior. We found that antipredator behaviors were not significantly different between the two nest predators, which posed a high risk to the nestlings, but lower risk to the parents as neither species frequently captures adult wrens. Females also did not differ significantly in their responses to the snake between years, suggesting that the population-level response might be consistent through time despite substantial individual variation among females. However, female wrens never dove at or attacked the Cooper's hawk while they frequently attacked both the snake and chipmunk decoys. Neighboring house wrens from adjacent territories were also less likely to respond to the hawk, but more heterospecifics mobbed the hawk than the snake decoy. Collectively, these results show that risk assessment and the strength of the antipredator response varies substantially both within and among species. Female house wrens exhibit plasticity in their nest defense behavior, and they respond to different types of predators in a way that could maximize lifetime fitness while risking the loss of their current offspring.

Introduction

Parental investment can be defined as any behaviors that increase an offspring's chance of survival while simultaneously decreasing the parent's ability to invest in other offspring both currently and in the future (Trivers 1974; Montgomerie and Weatherhead 1988; Clutton-Brock 1991). This tradeoff between current and future reproduction creates a parent-offspring conflict (*sensu* Trivers 1974) because the optimal parenting strategy, one that maximizes the parent's lifetime fitness, often requires submaximal investment in the current offspring, which can decrease their survival and fitness. The amount of parental investment varies widely both within and among species (Fontaine and Martin 2006; Dulac et al. 2014), and this variation can be further explained by differences in life history strategies (Ricklefs 1977; Montgomerie and Weatherhead 1988; Clutton-Brock 1991; Ghalambor and Martin 2001). For example, both adult mortality rates and offspring number and age should correlate positively with parental investment because species with shorter lifespans and larger current investments will have fewer reproductive opportunities in the future (Caro 2005).

Optimally modulating parental investment is particularly critical in the context of antipredator behavior, as parents that vigorously defend their young risk completely eliminating their future reproduction if they are mortally injured or killed. On the other hand, parents that limit their antipredator behavior are more likely to lose their current offspring as predation is one of the most common causes of reproductive failure across species (Ricklefs 1977; Caro 2005; Martin and Briskie 2009; Ibáñez-Álamo et al. 2015). The most successful parents should be those that exhibit plastic antipredator behavior that maximizes their lifetime reproductive success, and selection should favor individuals with accurate risk assessment during encounters with predators near their offspring (Montgomerie and Weatherhead 1988).

An abundance of evidence supports the hypothesis that animals from across the

taxonomic spectrum use risk assessment to adjust their antipredator behavior relative to the

perceived threat level (reviewed in Caro 2005; Lima 2009). Species with altricial young, such

as passerine songbirds, are particularly vulnerable to predation at the nest because their

offspring have limited capabilities for self-defense or escape without parental assistance (e.g.,

Suzuki 2011). Consequently, parents respond to different predation pressures with both

proactive and reactive behavior. When the perceived risk of predation is high, parents can

proactively reduce egg or clutch sizes (Fontaine and Martin 2006; Morosinotto et al. 2010;

Travers et al. 2010) or limit visits to the nest, including offspring provisioning (Fontaine and

Martin 2006; Ghalambor et al. 2013; Dudeck et al. 2018), which minimizes their potential

energetic losses and draws less attention to the nest site. Alternatively, parents may simply

avoid nesting in locations with high predator abundance or even forgo reproduction entirely

(Spaans et al. 1998; Morosinotto et al. 2010; Szymkowiak and Thomson 2019).

When a predator appears at the nest site, parents can react in ways that might divert the

predator's attention toward the parents rather than the offspring such as alarm calling or visual

distraction displays (e.g. broken-wing display; de Framond et al. 2022). Alarm calling also has

the potential to initiate a community-level response by attracting nearby conspecifics and

heterospecifics to collectively harass (i.e., "mob") the predator, which may increase the

likelihood of expelling the predator from the area (Carlson and Griesser 2022). Although alarm

calls and visual displays increase risk to the parents, they are often less costly than more

aggressive defense behaviors such as diving at or hitting the predator in an attempt to

physically drive it away from the offspring. Studies from multiple species have shown that

parents exhibit plasticity in their antipredator responses and adjust their behavior according to

factors such as the distance between the predator and the offspring (Colombelli-Négrel et al.

2010; Fernández and Carro 2022) and the relative risks posed to both adults and offspring by

different types of predators (Gottfried 1979; Winkler 1992; Mahr et al. 2015; Duré Ruiz et al.

2018; Teunissen et al. 2020). Similarly, the willingness of nearby conspecifics and

heterospecifics to respond to alarm calls with mobbing behavior varies according to the

intensity of the calls (Templeton et al. 2005, Fernández et al. 2023) and the risks posed by the

predator (Forsman and Mönkkönen 2001, Scott and Robinson 2023). In this study, we tested

for evidence of risk assessment by female northern house wrens (*Troglodytes aedon aedon*)

during simulated interactions with predators at their nest that posed different threats to adults

and nestlings.

Previous research in both the northern and southern house wren subspecies has found

evidence for risk assessment and plasticity in parental behavior in response to predation

pressure. For example, wren parents produced more alarm calls when a snake model was

placed closer to their nest (Fernández and Carro 2022), consistent with an ability to perceive

levels of risk. After interacting with a simulated nest predator, wren parents have also been

shown to reduce their offspring provisioning (Freed 1981; Ghalambor et al. 2013) and take

longer to re-enter the nest box when compared to control observations (Corral et al. 2013;

Fernandez and Llambías 2013; Fernández et al. 2015; Duré Ruiz et al. 2018). When presented

with different simulated species of raptors of varying threat levels, male and female southern

house wrens (*Troglodytes aedon musculus*) took longer to resume normal parental behavior

after encountering a predator of both adults and nestlings than a predator of nestlings only

(Duré Ruiz et al. 2018). This result is consistent with house wrens assessing the threat posed by

a heterospecific bird at their nest and responding in a potentially adaptive way by behaving

more cautiously in the presence of a predator of adult wrens.

Here, we expand on these observations by testing whether wren parents assess risks

posed by both avian and non-avian predators and adjust the strength of their nest defense

relative to risks associated with each type of predator (see below). Over two breeding seasons,

we placed three simulated predators of varying risks on top of house wren nest boxes during

the early nestling stage. One predator, a rubber eastern ratsnake decoy (*Pantherophis*

alleghaniensis), was presented in both years, while the second predator differed between years

and consisted of a rubber eastern chipmunk decoy (*Tamias striatus*) in year one and a

taxidermied juvenile Cooper's Hawk (*Accipiter cooperii*) in year two. We assumed that the

ratsnake and chipmunk decoys posed a high risk to nestlings, but a low-to-moderate risk to

adult wrens, while the hawk posed a high risk to adults, but a low-to-moderate risk to nestlings,

which are still vulnerable to raptors and other avian predators despite the protection offered by

an enclosed nest box (Milsap et al. 2013; Johnson 2020). If the hypothesis that house wrens use

risk assessment is supported, we predicted that female wrens would respond equally

aggressively to the ratsnake and chipmunk decoys, which are similar in threat level to both

nestlings and adults. Female responses to the hawk were predicted to be less aggressive overall

due to the higher threat that the hawk posed to adults and lower threat posed to nestlings.

Parental investment theory also predicts a positive relationship between the intensity of nest

defense and the size of the reproductive investment (Montgomerie and Weatherhead 1988,

Caro 2005). We tested this prediction by comparing nest defense behavior to clutch size and

nestling age, which varied slightly across our predator treatments. Finally, we examined the

community-level response to predators of varying risk by quantifying the number of

conspecific and heterospecific individuals that responded to each predator treatment.

Methods

Study System and Sites

Northern house wrens (*Troglodytes aedon aedon*) are socially monogamous, cavity-

nesting songbirds that readily use artificial nest boxes to raise clutches of up to 8 offspring

(Johnson 2020). The guild of house wren nestling predators is diverse and includes mammals

ranging in size from mice (e.g., *Peromyscus* spp.) to bears (*Ursus spp.*), songbirds (e.g.,

Cyanocitta spp.), woodpeckers (e.g., *Melanerpes* spp.), raptors, and snakes (Johnson 2020). At

our study sites the most common house wren nest predators are small mammals and snakes,

but multiple raptor species also frequent our sites and pose risks to adults and offspring (DGR,

personal observation).

We monitored house wren breeding success at 180 nest boxes divided between six

locations in Delaware and Union Counties, Ohio, USA (Table S1, Supplementary Materials)

during the 2020 and 2021 breeding seasons. The nest boxes varied slightly in dimensions due

to landowner preferences, but the average internal dimensions were 110 x 115 x 195 mm and

the entrance hole size was 38 mm. Each nest box was mounted on a conduit pipe

approximately 1.75 m above the ground, and the pipe was covered in axle grease to limit

access by predators. The majority of the boxes were located in edge habitat near the transition

from grass or field to forest, but a small subset was located in suburban backyards (*N*=7).

Banding of Adults and Nestlings

We captured male house wrens using mist nets and a brief playback of conspecific song.

Each male was banded with a USFWS aluminum leg band and a unique combination of 2-3

additional colored bands for identification from a distance. We also collected morphological

measurements and a small blood sample for an unrelated study. Banding only males allowed us

to differentiate between the sexes during our behavioral trials (see below), and because most

females were never captured, it also minimized the likelihood that the capture process

disrupted the female's antipredator response. Any females that were captured accidentally

(Experiment 1, $N = 6$ of 38 females; Experiment 2, $N = 10$ of 38 females) were rapidly banded

and immediately released within a few minutes of capture to limit any effects of handling on

subsequent behavior. All subjects were captured and banded at least 24 hours before any

behavioral trials, and most females were captured more than one week before their first trial

(Experiment 1, $N = 4/6$; Experiment 2, $N = 5/10$).

We also monitored the number of eggs laid, hatching success, and offspring development

for every nest box. We counted the number of nestlings at the end of each behavioral trial (see

below), and banded nestlings when they were 8 days old. Each nestling received a single

USFWS aluminum band, and we collected morphological measurements and a small blood

sample. After banding, we passively monitored each nest box until all visible and acoustic

activity ceased. Unless the contents of the nest suggested otherwise (i.e., presence of dead

nestlings and/or their parts), we assumed that empty nests successfully fledged.

Experiment 1: Chipmunk v. Snake Decoy

In 2020, we conducted behavioral trials between June 9 and July 24 at 47 nest boxes

containing nestlings between the ages of 3-7 days old. A smaller age range was impossible due

to the distances between our sites and other field-related challenges (e.g., weather). Two life-

like rubber predator decoys, an eastern chipmunk (*Tamias striatus*; Safari Ltd.) and an eastern

ratsnake (*Pantherophis alleghaniensis*; Flybanboo), were used to elicit nest defense behavior.

Both of these species are commonly observed at our study sites. We waited for each female to

enter or land on the nest box before flushing them away during setup. One predator was placed

on top of the nest box, and then on the following day the other predator was presented in a

counterbalanced order to control for order effects. Although these stimuli are terrestrial

predators, both are capable of climbing nearby vegetation and approaching the nest box from

above, making their appearance on top of the box biologically relevant.

All trials occurred between 7:00-13:00, but we were unable to conduct trials exactly 24

hours apart due to the distant locations of our field sites and unpredictability in how rapidly

females would return to their boxes. Trials were recorded with a video camera mounted on a

tripod placed roughly 2-3 meters away from the nest box to later confirm behaviors and

differentiate between the male and female wren in cases of uncertainty. We positioned the

camera in tall grass or other vegetation to minimize its disruption to the focal birds, and

observers maintained a distance of roughly 15 m away from the nest box. Each decoy was

attached to a fishing line, and at the completion of the trial we used a fishing rod to rapidly reel

the decoy away from the box. This approach eliminated the need for the observer to approach

the box and remove the decoy. We allowed the video camera to continue to record for an

additional 20 minutes to measure the amount of time it took for the female to re-enter the nest

box after removal of the decoy (*N*=36). Post-trial data for some of the females were lost due to

technical issues or user error.

We quantified the nest defense behavior of female house wrens for a total of seven

minutes after the decoy was placed on the nest box. We counted the number of hits, defined as

any contact the female made with the decoy, and the number of flyovers, defined as flights by

the female over the decoy. We also measured the amount of time females spent within five

meters of the nest box, the amount of time that females spent physically on or in the nest box,

and the female's closest approach (m) to the decoy. The five-meter threshold was chosen

because all boxes had potential perches within five meters of the nest box despite variation in

vegetative cover among our sites. Any female that hit the decoy was given a closest approach

value of zero meters. We also noted whether or not the female produced alarm calls, and

whether the female's mate was present during the trial. Neighboring conspecifics and

heterospecifics sometimes responded to the predator decoys by approaching, alarm calling, and

in rare cases flying over and hitting the decoy. During each trial, we noted the number of

individuals of each species that approached within 5 meters of the decoy. To avoid counting

incidental birds that were not responding to the predator, we only counted individuals that

either oriented towards the decoy or produced alarm calls.

Behavioral data from both predator trials for an individual female were excluded from

our final analysis if either of the decoys were knocked off the box during either trial ($N=3$ of 47

total trials; 6.4%). However, if the chipmunk decoy was knocked over but remained on the

box, the trials were not excluded ($N=5$ of 47 trials; 10.6%). We also excluded trials if the

female did not return to the box within seven minutes of initiating the trial ($N=6$ of 47 trials,

12.8%). In response to this limitation, we adjusted our protocol early in the season so that we

removed the decoy and reset the trial if the female did not return within 4 minutes after the trial

was initiated. Once the female returned to the box, we flushed her away, placed the predator

decoy, and started the seven-minute observation again. In summary, nine trials (of 47 trials;

19.1%) were excluded and our final sample size was 38 paired behavioral trials.

Experiment 2: Hawk v. Snake Decoy

In 2021 between June 6 and July 30, we replicated this experiment in the same house

wren population, but we replaced the chipmunk decoy with a taxidermied, juvenile Cooper's

hawk (*Accipiter cooperii*), which is commonly observed at our study sites. Although Cooper's

hawks primarily forage on small or medium-sized adult songbirds and mammals, they have

been frequently observed capturing nestlings from both cavity nests in bird boxes and open-cup

nests in the forest canopy (Milsap et al. 2013, Rosenfield et al. 2020). The hawk was mounted

with wire onto a wooden plank, and two large binder clips were attached to the bottom of the

plank with jewelry wire. This design allowed the decoy to be quickly attached and removed

from the roof of the nest box while eliminating the possibility that it could be knocked off the

box.

We conducted behavioral trials at 41 nest boxes with nestlings aged 2-7 days. A different

observer (JHF) scored the trials in 2021, but she was trained by the observer (RCE) from 2020

to ensure consistency between years. All of our sampling methods were replicated from

experiment 1, but we were unable to continue to use the fishing rod and line to rapidly remove

the decoys. As a result, one of the observers quickly ran to the box and removed the decoy at

the end of each seven-minute trial. The video camera continued to record for an additional 20

minutes (*N*=38), but the approach of the observer may have affected the return to normal

behavior. The snake decoy was knocked off the nest box during three trials (of 41 total; 7.3%),

and we excluded both the snake and hawk trials from those subjects in our final analysis for a

total sample of 38 paired behavioral trials.

It is difficult to determine how many females were sampled in both experiments because

only 13 of 47 females (27.7%) were banded in experiment one, including six females banded in

previous breeding seasons. Of those banded females, only one (7.7.% return rate) was

definitely sampled in experiment two. It is likely that some of the unbanded females also

returned to be sampled again, and some inferences can be drawn from males, which were

banded at much higher rates. In experiment one, 43 of 47 males (91.5%) were banded and eight

males returned to be sampled again in experiment two (18.5% return rate). Based on these data,

it seems likely that 10-20% of females were sampled in both experiments.

Note on Experimental Design

Our decision to split this study into two separate experiments rather than one experiment

including all three predator stimuli was driven by uncertainty stemming from the global Covid-

19 pandemic. This event shutdown numerous research labs in 2020 and disruptions continued

into 2021. The potential for an unpredictable and abbreviated 2020 field season led us to err on

the side of a less complicated protocol to maximize our sample size. We chose to focus on two

nest predators (snake, chipmunk) in 2020, and then based on the results of that experiment, we

transitioned to one nest predator (snake) and one predator of nests and adults (hawk) in 2021.

One additional limitation of this design is that we lack a non-predatory control, such as a

heterospecific passerine not known to depredate or destroy other species' nests. Thus, the

responses we observed could be typical of general nest defense rather than specifically

antipredator responses. We think this interpretation is unlikely to be supported because similar

nest defense experiments in southern house wrens, a closely-related subspecies, have shown

that both male and female wrens respond less aggressively to non-predatory control birds at

their nest (Fasanella and Fernández 2009, Duré Ruiz et al. 2017). Furthermore, a study in a

different population of northern house wrens found that parents decreased their provisioning

more in response to a simulated nest predator than a non-predatory control bird (Ghalambor et

al. 2013). Although this study did not measure nest defense, it is consistent with wrens

behaving differently around predators relative to non-predators near the nest.

Statistical analysis

For each experiment, we generated linear mixed models in R version 4.1.3 (R Core Team

2022; packages: "lme4," "car") to test whether female behaviors including flyovers, hits,

closest approach to the decoy, time spent within five meters of the decoy, and latency to re-

enter the nest box after removal of the predator differed between predator types. Measurements

of time spent on or inside the box were excluded due to extremely low rates of occurrence in

both experiments (Table S2). Each model included female behavior as the dependent variable

with predator type, treatment order, and their interaction as fixed effects. Day of the year was

included as a covariate and female ID was included as a random effect to control for seasonal

shifts in behavior and repeated sampling of the same individuals. Alarm calls were recorded as

a categorical variable (presence/absence), and we compared the proportion of females that

produced calls between predator types for each experiment using a two-proportion z-test. We

also compared the proportion of males that responded to each predator type in each experiment

using a two-proportion z-test.

We used Mann-Whitney U Tests (package: "stats") to determine whether the number of

neighboring conspecific and heterospecific individuals and the number of heterospecific

species that responded to the trials differed between predator types in both years. This

nonparametric test was chosen due to the lack of normality and low variance in these datasets

caused by a large proportion of zero or near zero values.

We also used linear mixed models to conduct inter-annual comparisons of female

behavior for the common predator present during both years, the eastern ratsnake. Due to the

change in retrieval protocol of the predator models between the two years (see above), we were

unable to compare latency to re-enter the box between years. Data points for the inter-annual

analysis were excluded if the subject was also excluded from the analysis for either individual

experiment (see above). In each model, female behavior was the dependent variable and year

was the fixed effect. Day of the year was included as a covariate and treatment order was a

random effect.

We used linear regressions (package: "stats") to test for relationships between female

antipredator behaviors and both clutch size and nestling age. To expand our sample size, we

also pooled female responses to the snake decoy across years and ran linear regressions

between behavior and both clutch size and nestling age. In addition, we used generalized linear

mixed models to test whether female antipredator behaviors predicted whether nests fledged or

failed. Nest outcome was included as the dependent variable with female behavior as a fixed

effect, day of the year as a covariate, and female ID as a random effect. These models assumed

a binomial distribution to account for nest outcome being scored as a categorical variable

(fledge/fail).

To evaluate model assumptions across our analyses, we tested for normality of model

residuals using Shapiro-Wilk tests and for homoscedasticity using Levene's test. As is often

the case with behavioral data, few of our models' residuals achieved normality and most were

slightly skewed towards one end of the distribution. Homoscedasticity was met for all models

in experiment 1 and the inter-annual comparisons, but only for the female return time data in

experiment 2. This violation was seemingly due to the uniformly weak responses elicited by

the hawk decoy in experiment 2. An abundance of evidence has indicated that linear mixed

models are robust to violations of these assumptions, except in cases where the sample sizes

are prohibitively small or the distributions are bimodal, neither of which occurs in our data

(Schielzeth et al. 2020; Knief and Forstmeier 2021; and further citations within).

Ethical Note

All procedures were reviewed and approved by the Ohio Wesleyan University

Institutional Animal Care and Use Committee (Protocol Numbers: Spring_2020-21-

01_Reichard and 05-2021-06), U.S. Federal Bird Banding Lab (Permit Number: 24035), and

State of Ohio Department of Natural Resources (Permit Numbers: 23-009, 23-010). To

minimize stress in the free-living birds used in this study, we attempted to capture and mark

only the male of each pair, which was the minimum required to facilitate individual

identification during our behavioral trials. All birds were immediately removed from mist nets

after capture, processed rapidly to limit handling time, and then safely released at the capture

site. No adverse effects were observed from these widely established techniques. Although our

behavioral trials involved the observation of a simulated predator at the nest box, all subjects

returned to normal parental care after the completion of the trial and no nests were abandoned

as a result of these experiments.

Statement on STRANGE framework

Studies of animal behavior can be misinterpreted due to sampling biases, and the

STRANGE framework provides a method for identifying these potential issues (Webster and

Rutz 2020). All subjects in this study were free-living songbirds, and we included all

individuals that nested on our study sites regardless of age or nesting location, making it

unlikely that social background biased our results. As detailed above, a small number of

individuals were excluded, namely those that failed to respond within 7 minutes during

Experiment 1 before our protocol was adjusted to be more inclusive or those that responded so

forcefully that they knocked the predator decoy from the box. Thus, we may have failed to

sample a subset of both low and high responders. Although this approach may have biased our

sample towards intermediate responders, we think this potential bias is unlikely to have skewed

our results for a few reasons. First, our sample size is large for a field study and it includes

individuals that responded both aggressively and weakly to the predator decoys, so it is

unlikely that we failed to sample the extremes of behavioral variation in this population.

Second, we conducted repeated-measures sampling, which required subjects to be excluded

from all treatments rather than only those trials where we were unable to collect data. This

approach limits the likelihood of bias as each individual effectively serves as its own control.

With respect to seasonal changes in behavior, we sampled evenly across the entire breeding

season and included measurement date as a covariate in our models to test for this effect.

Because each subject experienced two sampling events it is possible that habituation affected

responses to the second treatment. We limited this effect by presenting trials on separate days

in a counterbalanced order and included treatment order in our models to test for an effect of

habituation.

Results

Experiment 1: Chipmunk v. Snake Decoy

Female wrens exhibited no significant differences in the number of hits (χ^2_1=0.02,

P=0.88), number of flyovers (χ^2_1<0.001, *P*>0.99), time spent within 5 meters of the nest box

(χ^2_1=0.45, *P*=0.50), or in their closest approach (χ^2_1=1.45, *P*=0.23) to the snake and chipmunk

decoys (Figure 1A-D). There was also no difference in the number of females (36 of 38 total

for both predators; 94.7%) that produced alarm calls in response to each predator (χ^2_1<0.001,

P>0.99). After the predator decoys were removed, we found no significant difference between

predator types in the amount of time females took to re-enter the nest box (χ^2_1=0.02, *P*=0.88;

Figure 2A). Males were present for 87% of snake trials (*N*=33 of 38 total) and 92% of

chipmunk trials (*N*=35 of 38), but this difference was not statistically significant (χ^2_1=0.14,

P=0.71).

Day of the year had a significant effect on the closest approach to the predator (χ^2_1=3.98,

P=0.046) with females approaching more closely later in the breeding season. We also noted

trends towards an effect of date on the time spent within 5 m of the predator (χ^2_1=3.73,

P=0.054) and the time to return to the nest box (χ^2_1=3.13, *P*=0.08). Later in the breeding

season, females tended to spend more time within 5 m of the predator and took more time to

return to the nest box after the predator was removed. There was no detectable effect of date on

hits or flyovers (P>0.27 in both cases; Table S3). For all five behaviors, we found no evidence

for significant effects of treatment order (P>0.12 in all cases; Table S3) or significant

interactions between treatment order and predator type (P>0.07 in all cases; Table S1). The

only exception was a trend towards an interaction between predator types and order for the

time spent within 5 m (χ^2_1=3.38, P=0.07). Female wrens tended to spend less time within 5m

of the chipmunk decoy when it was presented first, but the opposite was true for the snake

decoy (less time when presented first).

Figure 1 — Responses of female house wrens to a snake or chipmunk decoy on top of their nest

box during a seven-minute trial. There were no significant differences in the number of (A) hits

(P=0.88), (B) flyovers (P>0.99), (C) how closely females approached each model (P=0.23), or

(D) how much time females spent within 5 m of each model (P=0.50). The thick center line

represents the median, and the box encloses the interquartile range. Whiskers show the range

of data within 1.5 times the interquartile range, and dots are data points exceeding that range.

399
400

Figure 2 — Female house wrens modulate how quickly they resume normal behavior after

encountering different nest predator decoys. (A) There was no detectable difference in how

long it took female wrens to resume normal behavior after encountering a snake decoy versus

a chipmunk decoy (P=0.88). (B) Females took significantly longer to resume normal behavior

after encountering a hawk decoy than after encountering a snake decoy (P=0.03). Box plots as

*in Fig. 1. *P<0.05.*

The number of conspecific (W=724.5, P=0.98) and heterospecific individuals (W=693.5,

P=0.74) and the number of heterospecific species (W=700.5, P=0.80) that responded to the

predator decoys did not differ detectably between the two predator types.

Table 1. Mean ± standard deviation (range) for the number of conspecific and heterospecific

individuals and heterospecific species that responded to the predator stimuli in Experiment 1.

Observation	Snake	Chipmunk
Conspecific Individuals	0.53 ± 1.0 (0-5)	0.47 ± 0.8 (0-3)
Heterospecific Individuals	1.08 ± 2.7 (0-15)	0.76 ± 1.4 (0-6)
Heterospecific Species	0.61 ± 1.0 (0-5)	0.58 ± 0.9 (0-4)

We found no relationship between clutch size and the number of hits, number of flyovers,

time spent within 5 meters of the nest box, closest approach, or latency for female wrens to

return to the nest box after the trial (P>0.10 in all cases; Table S4). Similarly, there were no

significant relationships between any of the female behaviors and nestling age (P>0.40 in all

cases; Table S5). We also found that none of the nest defense behaviors were predictive of

whether nests fledged (N=21) or failed (N=4; P>0.22 for all behaviors; Table S6). The sample

size for this comparison is reduced (N=25) because multiple nests remained active after the end

of our data collection and their end point is unknown.

Experiment 2: Hawk v. Snake Decoy

Similar to Experiment 1, female wrens frequently performed hits and flyovers while

defending against the snake decoy, with 25 out of 41 females (61.0%) hitting the snake model

at least once and 38 out of 41 females (92.7%) flying over at least once. Notably, none of the

female wrens performed hits or flyovers while defending against the hawk decoy (Figure 3).

Statistically, females performed significantly more hits (χ^2_1=14.88, $P < 0.001$) and flyovers

(χ^2_1=42.75, $P < 0.001$) in response to the snake decoy (Figure 3A-B). Females approached

significantly closer to the snake (χ^2_1=116.9, $P < 0.001$; Figure 3C), but the amount of time

spent within 5 meters was not significantly different (χ^2_1=3.04, P=0.08) despite a marginal

trend towards more time in close proximity to the snake (Figure 3D).

We found no evidence for an effect of date (P>0.09 in all cases; Table S7), treatment

order (P>0.14 in all cases; Table S6) or a predator type by treatment order interaction (P>0.22

in all cases; Table S7) on any behaviors with two exceptions. There was a trend towards an

order effect for time spent within 5 meters of the predator decoy (χ^2_1=3.70, P=0.06; Table S7).

Female wrens tended to spend more time within 5 meters of the decoy during the second trial

than during the first trial regardless of the order in which predators were presented. In addition,

there was a marginal effect of date on the number of flyovers (χ^2_1=2.88, P=0.09; Table S7)

such that females tended to produce more flyovers later in the breeding season.

An equal number of females (34 of 38 total; 89.5%) produced alarm calls in response to

each of the two predator decoys (χ^2_1<0.001, P>0.99). After the removal of the predator decoys,

female wrens took significantly more time to re-enter the nest box after the hawk trials than

after the snake trials (χ^2_1=4.90, P=0.03; Figure 2B). There was no detectable effect of treatment

order (χ^2_1<0.01, P=0.93) or an interaction between order and predator type (χ^2_1<0.001, P=0.98)

for this behavior. Males were present for 87% of snake trials (*N*=33 of 38 total) and 95% of

hawk trials (*N*=36 of 38), which was not statistically different (χ^2_1=0.63, *P*=0.43).

Figure 3 — Responses of female house wrens to a snake or hawk decoy on top of their nest box

during a seven-minute trial. Female house wrens performed significantly more (A) hits

(P<0.001) and (B) flyovers (P<0.001) against the snake decoy than against the hawk decoy.

Female wrens also (C) made significantly closer approaches to (P<0.001) and (D) tended to

spend more time within 5 meters of the snake decoy than the hawk decoy (P=0.08). Box plots

as in Fig. 1. ✱ *P<0.001.* ‡ *P<0.10.*

Significantly more heterospecific individuals responded to the hawk decoy than to the

snake decoy (*W*=531.5, *P*=0.04), but there was no detectable difference in the number of

heterospecific species that responded to each predator (W=599.5, P=0.18). In contrast,

significantly more conspecific individuals mobbed the snake decoy than mobbed the hawk

(W=908, P=0.01).

Table 2. Mean ± standard deviation (range) for the number of conspecific and heterospecific

individuals and heterospecific species that responded to the predator stimuli in Experiment 2.

Bold text denotes P<0.05.

Observation	Snake	Hawk
Conspecific Individuals*	**0.55 ± 0.8 (0-3)**	**0.13 ± 0.3 (0-1)**
Heterospecific Individuals*	**0.97 ± 1.1 (0-4)**	**2.05 ± 2.2 (0-9)**
Heterospecific Species	0.84 ± 0.9 (0-3)	1.29 ± 1.4 (0-6)

As in 2020, we found no evidence of a relationship between clutch size (P>0.14 in all

cases; Table S4) or nestling age (P>0.63 in all cases; Table S5) and any female nest defense

behaviors. We were unable to compare nest defense behavior between nests that successfully

fledged and those that failed due to a limited sample size of failed nests (N=1).

Interannual Comparison — Snake 2020 Data v. Snake 2021 Data

In addition to comparing how female house wrens responded to two different nest

predators during a single breeding season, we compared how the wrens responded to the same

predator—the snake—during two different breeding seasons. There was no significant

difference between the number of hits (χ^2_1=1.34, P=0.25), the number of flyovers (χ^2_1=0.87,

$P=0.35$), or the time that female wrens spent within 5 meters of the snake decoy ($\chi^2_1=0.23$,

$P=0.63$) between years (Figure 4). We found a significant difference ($\chi^2_1=3.91$, $P=0.048$)

between the female wrens' closest approach to the snake decoy in 2020 ($x̄=0.72$ meters)

versus in 2021 ($x̄=0.18$ meters; Figure 4). However, after removing one low-responding

outlier from the 2020 data set (closest approach = 7 m), the difference was no longer

significant ($\chi^2_1=2.84$, $P=0.09$; Figure 4). A slightly higher proportion of females (36 of 38;

94.7%) produced alarm calls in 2020 than in 2021 (34 of 38; 89.5%), but these proportions

were not statistically different ($\chi^2_1=0.18$, $P=0.67$).

Because the majority of the female behaviors were not statistically different between the

two years, we combined both years of snake response data to increase our sample size and test

for a relationship between both clutch size and nestling age and nest defense behavior.

However, the results remained unchanged as we found no evidence for any significant

relationships between female behavior and clutch size ($P>0.33$ for all behaviors; Table S8) or

nestling age ($P>0.09$ for all behaviors; Table S9). The only potential exception was a slight

trend towards females spending less time within 5 m of the snake decoy as nestling age

increased ($P=0.09$; Table S9).

Figure 4 — Interannual comparison of female house wren nest defense behaviors in response to a snake decoy across two breeding seasons. There were no significant differences in (A) the number of hits (P=0.25) or (B) the number of flyovers (P=0.87). (C) Female wrens made significantly closer approaches in 2021 than in 2020 (P<0.05), but when one extreme outlier was removed from the 2020 data set, the difference was no longer significant (P=0.09). (D) There was no significant difference between the amount of time spent within 5 meters of the snake model between years (P=0.67). Box plots as in Fig. 1. † P<0.05 before outliers were removed from 2020 data.

Discussion

Defending offspring from predators is a risky behavior that varies across species and contexts (Dugatkin and Godin 1992; Lima 2009). Predator guilds are often diverse, and all

predators do not pose equal threats to both adults and their young. As a result, parents that

accurately assess the risks posed by different predators and scale their responses appropriately

should have higher lifetime fitness (Montgomerie and Weatherhead 1988; Clutton-Brock 1991;

Caro 2005). We found evidence for risk assessment and plastic antipredator behavior in nesting

house wrens. Female wrens responded with similar levels of aggressive nest defense behavior

when encountering a simulated snake or chipmunk at their nest box, both of which posed a

higher risk to offspring than adults. In contrast, the wrens responded significantly less

aggressively to a simulated hawk that posed a higher risk to adults than nestlings, and females

also took longer to return to their nest box after encountering the hawk than the snake. This

plasticity in nest defense behavior is potentially adaptive because parents that avoid high risk

predators like the hawk are more likely to live longer and have higher lifetime fitness despite

the potential costs to the current clutch of offspring (Trivers 1974; Montgomerie and

Weatherhead 1988). Collectively, these results are consistent with the predictions of parental

investment theory (Montgomerie and Weatherhead 1988; Caro 2005), and similar parental risk

assessment in different contexts has been previously observed in house wrens and other species

(Lima 2009; Mahr et al. 2015; Duré Ruiz et al. 2018).

We also found that female wrens responded similarly to a simulated snake at the nest box

between the two breeding seasons, which indicates that the mean antipredator response was

stable at the population level despite limited sampling of the same females between years. This

population-level consistency is in stark contrast to the large amount of individual variation

among females in how strongly they responded to each of the simulated predators. For

example, the most aggressive females hit the snake and chipmunk decoys over 50 times in a

seven-minute period while the least aggressive females never dove at or hit the decoys.

Parental investment is predicted to be higher when clutch sizes are larger and offspring

are older. However, we found no evidence of a relationship between offspring number or age

and nest defense behavior, and both of these predictions have received mixed support more

broadly (Montgomerie and Weatherhead 1988; Caro 2005). One limitation of our study is that

we may have lacked a sufficient range and samples of clutch sizes and nestling ages to reveal

any relationships with antipredator behavior. For example, 80% of the wren pairs in our study

were defending clutches of 4-6 offspring, and 87% had nestlings between 3-6 days post hatch.

A proper test of these relationships could involve experimental manipulations of clutch size or

sampling nest defense at two distant points in the nestling phase such as soon after hatch and

again near fledging. Additionally, there are other variables such as parental age and experience

that could explain some of the individual variation in nest defense behavior (Caro 2005).

Two important exceptions to the persistent variation among individuals in nest defense

behavior occurred with the production of alarm calls across the predator stimuli and also in the

behavioral responses to the simulated hawk. Females almost universally produced alarm calls

in response to all three of our predator stimuli despite varying levels of risk. Alarm calls are a

lower risk behavior than the other nest defense behaviors sampled here, which might explain

their consistent production. When produced near the nest, alarm calls can serve many adaptive

functions including distracting the predator (Fasanella and Fernández 2009), inducing

neighboring birds to mob the predator (Templeton et al. 2005), and alerting the young inside

the nest that a predator is nearby (Suzuki 2011). Previous research in another house wren

population found that parents produced alarm calls even in the presence of a non-predatory

heterospecific bird at their nest (Fasanella and Fernández 2009), suggesting that alarm calls

might function as a general response to any disturbance at the nest box. However, wren parents

also produced more alarm calls in response to a potential predator rather than a non-predatory

control bird (Fasanella and Fernández 2009) and more alarm calls were produced as a

simulated snake moved closer to the nest (Fernández and Carro 2022). Although we did not

measure whether the quantity of alarm calls differed between predator types in our study, it

seems clear that alarm calls are a consistent and important component of the antipredator

response, and the intensity of these calls might signal information about the predator type and

level of risk posed to the nestlings (Fasanella and Fernández 2009, Fernández and Carro 2022,

Fernández et al. 2023).

When responding to the hawk decoy, females varied in how closely they approached and

how much time they spent within five meters of the hawk, but none of the 41 females ever hit

the hawk or even flew over it. This consistency among individuals indicates that when the

predatory threat surpasses a certain level it is possible to see uniformity in a typically plastic

behavior. Furthermore, the total absence of overt aggression towards the hawk supports the

hypothesis that selection favors risk assessment as well as the prediction that selection can act

strongly against behaviors that carry higher risks such as attacking a predator capable of

injuring or killing an adult. However, our observations of the community-level response to the

hawk and other simulated predators show that risk assessment is not uniform across songbird

species.

Neighboring conspecifics and heterospecific songbirds frequently responded to our

simulated predators by approaching and alarm calling, but very few attacked or flew directly

over the decoys, including the snake and chipmunk. This observation is consistent with

mobbing behavior, and the reliance on alarm calls rather than overt aggression likely reflects

the reduced threat posed to these neighbors by a predator that was not in close proximity to

their own offspring (Colombelli-Négrel et al. 2010; Kryštofková et al. 2011). Similar to the

antipredator behavior of the wren parents, the number of conspecifics and heterospecifics that

responded did not differ significantly between the snake and chipmunk. In contrast, more

conspecifics responded to the snake while more heterospecifics responded to the hawk in our

second experiment. This result mirrors the broader avoidance of the hawk by house wrens,

particularly when the hawk is not posing a direct threat to their own offspring. It also highlights

that certain heterospecific species actually respond in greater numbers to the hawk than the

snake. Specifically, larger songbirds including tree swallows (*Tachycineta bicolor*), northern

cardinals (*Cardinalis cardinalis*), and blue jays (*Cyanocitta cristata*) appeared in higher

numbers when the hawk was present, and tree swallows frequently dove at the hawk, which

was in stark contrast with all other species. The willingness of these heterospecifics to risk

mobbing the hawk might be explained by the observation that Cooper's hawks tend to prefer

larger songbirds as prey (Roth and Lima 2003; Rosenfield et al. 2020), so the hawk posed a

greater risk to both adults and offspring in these species. Although the intensity of aggression

from tree swallows may seem surprising given that their nests are concealed and protected by

cavities (Montgomerie and Weatherhead 1988), observations of Cooper's Hawks have revealed

that they regularly check and depredate nest boxes while foraging (Milsap et al. 2013).

Reactive nest defense behavior can be subdivided into (1) behaviors that distract or

redirect the predator towards the signaler and (2) behaviors that involve acts of aggression

towards the predator (Caro 2005; Lima 2009). Smaller organisms like house wrens may rely on

distraction rather than attacking predators above a certain size because they lack the body size

or armaments necessary to successfully drive the predator away (Montgomerie and

Weatherhead 1988; Winkler 1992). If the likelihood of physically deterring the predator is low,

then any amount of risk to the parent from attacking the predator could justify a distraction-

based antipredator response or even no response at all. For example, when responding to

human researchers that are actively removing nestlings from the nest and posing a major threat,

house wrens in our population have never been observed diving at or hitting the researchers,

only approaching and alarm calling. Thus, it is possible that house wrens and other organisms

employ a size-based threshold process in which physical attacks are limited to predators below

the size threshold and distraction methods are used regardless. Multiple species employ a

similar threshold-based approach when judging predator proximity by increasing the intensity

of their antipredator behavior as predators get closer to their nest or offspring (Colombelli-

Négrel et al. 2010; Kryštofková et al. 2011; Mahr et al. 2015). The size-threshold hypothesis

proposed above could be tested by comparing parental responses to predators of equivalent risk

but varying sizes. However, it is unlikely that size thresholds universally govern antipredator

behavior given that certain species of limited size and armaments like northern mockingbirds

(*Mimus polyglottos*) are notorious for attacking humans and other large predators that approach

their nests (Levey et al. 2009).

Although risk assessment and plasticity in nest defense behavior appears to be common

among organisms with altricial offspring, including house wrens, understanding the proximate

and ultimate factors that drive the individual variation within and among species continues to

be challenging (Montgomerie and Weatherhead 1988; Caro 2005; Butler et al. 2009). To

adequately test the determinants of nest defense, it is both important and extremely difficult in

field-based systems to isolate and vary a small number of potentially important variables while

controlling for all others (Regelmann and Curio 1983; Winkler 1992). We tested for

connections between nest defense and fitness, but we found no evidence for a relationship

between nest defense behavior and clutch size or nestling age or any differences in behavior

between nests that fledged and those that failed. However, each of those comparisons was

limited by small sample sizes, insufficient controls for potentially important variables such as

female age, and the fact that our nests benefitted from extra protection of artificial predator

guards.

Despite these limitations, we did observe evidence for risk assessment in how female

wrens responded to different types of predators at the nest box, and their behavior matched our

predictions of the relative threats posed by each predator type. Furthermore, we observed

differences in the community-level response as heterospecific songbirds responded in greater

numbers to the hawk than conspecifics from neighboring territories. These results indicate that

risk assessment is not consistent among species, and species like house wrens that respond

minimally to certain predators can gain some protection from the antipredator behavior of

nearby heterospecifics. Future work should continue to focus on elucidating the causes and

consequences of persistent intra- and interspecific variation in parental antipredator behavior.

Data Availability Statement

The data collected and analyzed to support these findings are freely available in the

Mendeley Data Repository at DOI:10.17632/zjb5ynz5n4.2.

Acknowledgements

We thank Ohio Wesleyan University, the Big Walnut School System, and the Davis,

Koban, and Fink families for access to facilities and field sites. We thank Holly Keating for

analyzing videos, Tamaki Yuri for preparing the taxidermied Cooper's Hawk, Lily Bonar for

field assistance, Chandler Carr for measuring our nest boxes, and Eric Gangloff for advice on

statistics. This project was funded by the Ohio Wesleyan University Summer Science Research

Program.

Literature cited

Butler LK, Bisson I-A, Hayden TJ, Wikelski M, Romero LM. 2009. Adrenocortical responses

to offspring-directed threats in two open-nesting birds. Gen Comp Endocrinol.

162(3):313–318. doi:10.1016/j.ygcen.2009.04.008.

Caro T. 2005. Antipredator Defenses in Birds and Mammals. University of Chicago Press.

Carlson NV, Griesser G. 2022. Mobbing in animals: A thorough review and proposed future

directions. Advances in the Study of Behavior. 54:1-41. doi.org/10.1016/bs.asb.2022.01.003.

Clutton-Brock TH. 1991. The Evolution of Parental Care. Princeton University Press.

Colombelli-Négrel D, Robertson J, Kleindorfer S, Sulloway F. 2010. Extended parental care of

fledglings: parent birds adjust anti-predator response according to predator type and

distance. Behaviour. 147(7):853–870. doi:10.1163/000579510X495771.

Corral MG, Llambías PE, Fernández GJ. 2013. Effect of conspecific alarm calls in the parental

behaviour of nesting southern house wrens. Acta Ethol. 16(1):47–51.

doi:10.1007/s10211-012-0129-x.

Dudeck BP, Clinchy M, Allen MC, Zanette LY. 2018. Fear affects parental care, which

predicts juvenile survival and exacerbates the total cost of fear on demography. Ecology.

99(1):127–135. doi:10.1002/ecy.2050.

Dugatkin LA, Godin J-GJ. 1992. Prey approaching predators: a cost-benefit perspective.

Annales Zoologici Fennici. 29(4):233–252.

Dulac C, O'Connell LA, Wu Z. 2014. Neural control of maternal and paternal behaviors.

Science. 345(6198):765–770. doi:10.1126/science.1253291.

Duré Ruiz NM, Fasanella M, Fernández GJ. 2018. Breeding southern house wrens exhibit a

threat-sensitive response when exposed to different predator models. J Ethol. 36(1):43–

53. doi:10.1007/s10164-017-0528-6.

Fasanella M, Fernández, GJ. 2009. Alarm calls of the Southern House Wren *Troglodytes*

musculus: variation with nesting stage and predator model. Journal of Ornithology

150:853–863.

Fernández GJ, Carro ME. 2022. Alarm calls of southern house wrens, *Troglodytes aedon*

bonariae, convey information about the level of risk. Ethology. 128(4):293–302.

doi:10.1111/eth.13255.

Fernández GJ, Corral MG, Llambías PE. 2015. Sexual differences in risk-taking behaviour of

the Southern house wren during the nestling rearing stage. Acta Ethol. 18(1):11–18.

doi:10.1007/s10211-013-0177-x.

Fernández GJ, Dutour M, Carro ME. 2023. Information transfer during mobbing: call rate is

more important than the number of callers in a southern temperate passerine. Behav Ecol

Sociobiol. 77: 78. doi.org/10.1007/s00265-023-03357-z.

Fernández GJ, Llambías PE. 2013. Parental Risk-Taking Behaviour and Nest Defence During

the Nestling Rearing Stage in Northern House Wrens *Troglodytes aedon*. Acta Ornith.

48(1):55–63. doi:10.3161/000164513X670016.

Fontaine JJ, Martin TE. 2006. Parent birds assess nest predation risk and adjust their

reproductive strategies. Ecology Letters. 9(4):428–434. doi:10.1111/j.1461-

0248.2006.00892.x.

Forsman JT, Mönkkönen M. 2001. Responses by breeding birds to heterospecific song and

mobbing call playbacks under varying predation risk. Anim Behav. 62(6): 1067-1073.

doi.org/10.1006/anbe.2001.1856

de Framond L, Brumm H, Thompson WI, Drabing SM, Francis CD. 2022. The broken-wing

display across birds and the conditions for its evolution. Proc R Soc B: Biol Sci.

289(1971):20220058. doi:10.1098/rspb.2022.0058.

Freed LA. 1981. Breeding ecology of House Wrens: new views of avian life history

phenomena [PhD Thesis]. [Ames, Iowa]: University of Iowa.

Ghalambor CK, Martin TE. 2001. Fecundity-Survival Trade-Offs and Parental Risk-Taking in

Birds. Science. 292(5516):494–497. doi:10.1126/science.1059379.

Ghalambor CK, Peluc SI, Martin TE. 2013. Plasticity of parental care under the risk of

predation: how much should parents reduce care? Biology Letters. 9(4):20130154.

doi:10.1098/rsbl.2013.0154.

Gottfried BM. 1979. Anti-Predator Aggression in Birds Nesting in Old Field Habitats: An

Experimental Analysis. The Condor. 81(3):251–257. doi:10.2307/1367626.

Ibáñez-Álamo JD, Magrath RD, Oteyza JC, Chalfoun AD, Haff TM, Schmidt KA, Thomson

RL, Martin TE. 2015. Nest predation research: recent findings and future perspectives. J

Ornithol. 156(1):247–262. doi:10.1007/s10336-015-1207-4.

Johnson LS. 2020. House Wren (*Troglodytes aedon*), version 1.0. In: Poole AF, editor. Birds

of the World. Cornell Lab of Ornithology, Ithaca, NY, USA.

Knief U, Forstmeier W. 2021. Violating the normality assumption may be the lesser of two

evils. Behav Res Methods 53, 2576–2590.

Kryštofková M, Haas M, Exnerová A. 2011. Nest defense in Blackbirds *Turdus merula*: effect

of predator distance and parental sex. Acta Ornith. 46(1):53–61.

doi:10.3161/000164511X589938.

Levey DJ, Londoño GA, Ungvari-Martin J, Hiersoux MR, Jankowski JE, Poulsen JR, Stracey

CM, Robinson SK. 2009. Urban mockingbirds quickly learn to identify individual

humans. PNAS. 106(22):8959–8962. doi:10.1073/pnas.0811422106.

Lima SL. 2009. Predators and the breeding bird: behavioral and reproductive flexibility under

the risk of predation. Biol Rev. 84(3):485–513. doi:10.1111/j.1469-185X.2009.00085.x.

Mahr K, Riegler G, Hoi H. 2015. Parental risk management in relation to offspring defence:

bad news for kids. Proc R Soc B: Biol Sci. 282(1798):20141670.

doi:10.1098/rspb.2014.1670.

Martin TE, Briskie JV. 2009. Predation on dependent offspring: a review of the consequences

for mean expression and phenotypic plasticity in avian life history traits. Ann N Y Acad

Sci. 1168:201–217. doi:10.1111/j.1749-6632.2009.04577.x.

Milsap, BA, Breen, TF, Phillips, LM. 2013. Ecology of the Cooper's Hawk in North Florida.

North American Fauna 78: 1–58.

Montgomerie RD, Weatherhead PJ. 1988. Risks and Rewards of Nest Defence by Parent Birds.

Q Rev Biol. 63(2):167–187. doi:10.1086/415838.

Morosinotto C, Thomson RL, Korpimäki E. 2010. Habitat selection as an antipredator

behaviour in a multi-predator landscape: all enemies are not equal. J Anim Ecol.

79(2):327–333. doi:10.1111/j.1365-2656.2009.01638.x.

R Core Team. 2022. R: A language and environment for statistical computing. https://www.R-

project.org.

Regelmann K, Curio E. 1983. Determinants of Brood Defence in the Great Tit *Parus major* L.

Behav Ecol Sociobiol. 13(2):131–145.

Ricklefs RE. 1977. On the Evolution of Reproductive Strategies in Birds: Reproductive Effort.
Am Nat. 111(979):453–478.

Rosenfield RN, Madden KK, Bielefeldt J, Curtis OE. 2020. Cooper's Hawk (*Accipiter cooperii*), version 1.0. Birds of the World. doi:10.2173/bow.coohaw.01.

Roth TC II, Lima SL. 2003. Hunting Behavior and Diet of Cooper's Hawks: An Urban View of the Small-Bird-In-Winter Paradigm. The Condor. 105(3):474–483. doi:10.1093/condor/105.3.474.

Schielzeth, H, Dingemanse, NJ, Nakagawa, S, et al. 2020. Robustness of linear mixed-effects models to violations of distributional assumptions. Methods Ecol Evol 11: 1141– 1152.

Scott M, Robinson WD. 2023. Mobbing behavior of songbirds in response to calls of an ambush-predator, the Northern Pygmy-Owl. Front. Ecol. Evol. 11:1092323. doi: 10.3389/fevo.2023.1092323.

Spaans B, Blijleven HJ, Popov I, Rykhlikova M, Ebbinge B. 1998. Dark-bellied Brent Geese *Branta bernicla bernicla* forego breeding when Arctic Foxes Alopex lagopus are present during nest initiation. Ardea. 86:11–20.

Suzuki, TN. 2011. Parental alarm calls warn nestlings about different predatory threats. Current Biology. 21:R15-R16.

Szymkowiak J, Thomson RL. 2019. Nest predator avoidance during habitat selection of a songbird varies with mast peaks and troughs. Behav Ecol Sociobiol. 73(7):91. doi:10.1007/s00265-019-2702-z.

Templeton CN, Greene E, Davis K. 2005. Allometry of alarm calls: black-capped chickadees encode information about predator size. Science 308: 1934-1937.

762   Teunissen N, Kingma SA, Peters A. 2020. Predator defense is shaped by risk, brood value and

763   social group benefits in a cooperative breeder. Behav Ecol. 31(3):761–771.

764   doi:10.1093/beheco/araa012.

765   Travers M, Clinchy M, Zanette L, Boonstra R, Williams TD. 2010. Indirect predator effects on

766   clutch size and the cost of egg production. Ecology Letters. 13(8):980–988.

767   doi:10.1111/j.1461-0248.2010.01488.x.

768   Trivers RL. 1974. Parent-Offspring Conflict. American Zoologist. 14(1):249–264.

769   doi:10.1093/icb/14.1.249.

770   Webster MM, Rutz C. 2020. How STRANGE are your study animals? Nature 582, 337–340.

771   Winkler DW. 1992. Causes and Consequences of Variation in Parental Defense Behavior by

772   Tree Swallows. The Condor. 94(2):502–520. doi:10.2307/1369222.

     Supplementary Materials

786   *Table S1 - Locations of field sites and number of nest boxes present.*

Site	Location	Number of Boxes
Big Walnut Elementary School, Sunbury, Ohio	40°14'02.6"N 82°51'53.1"W	15
Big Walnut High School, Sunbury, Ohio	40°14'00.9"N 82°51'37.3"W	41
Davis Family Property, Sunbury, Ohio	40°15'50.2"N 82°49'52.8"W	36
Fink Family Property, Marysville, Ohio	40°16'42.4"N 83°15'43.3"W	21
Kraus Research Preserve, Delaware, Ohio	40°11'45.8"N 83°03'47.1"W	42
Ohio Wesleyan University and nearby homes, Delaware, Ohio	40°17'47.7"N 83°03'53.8"W	25

788   *Table S2 - Summary of the number of trials that include an observation of time spent in or on*
789   *the next box. Experiment 1 tested responses to a simulated snake and chipmunk. Experiment 2*
790   *tested responses to a simulated snake and hawk.*

Experiment	Time spent in the nest box	Time spent on the next box
1	4/76 (5.3%)	12/76 (15.8%)[1]
2	4/76 (5.3%)	10/76 (13.2%)[2]

791   [1] 9 of 12 observations were for 10 seconds or less (75%)
792   [2] 7 of 10 observations were for 10 seconds or less (70%)

801　*Table S3 – Analysis of predator order effects and interactions between predator type and order*
802　*for Experiment 1. *P<0.05*
803

	Day of the Year Effect		Order Effect		Interaction with Order	
Behavior	χ^2	P	χ^2	P	χ^2	P
Hits	0.99	0.32	<0.01	0.93	2.41	0.12
Flyovers	1.21	0.27	<0.01	0.93	1.29	0.26
Time within 5 m	3.73	0.054	<0.01	0.92	3.39	0.07
Closest approach	3.98	0.046*	1.19	0.28	1.84	0.18
Latency to return	3.13	0.08	2.41	0.12	0.10	0.76

805　*Table S4 – There were no detectable relationships between clutch size and nest defense*
806　*behaviors by female house wrens in either experiment.*

	2020 (Snake vs. Chipmunk)		2021 (Snake vs. Hawk)	
Behavior	t	P	t	P
Hits	0.09	0.93	-0.58	0.56
Flyovers	-0.13	0.90	0.86	0.39
Time within 5 m	-1.69	0.10	1.48	0.14
Closest approach	1.52	0.13	0.08	0.93
Latency to return	-0.58	0.56	-0.15	0.88

814 *Table S5 – There were no detectable relationships between nestling age and nest defense*
815 *behaviors by female house wrens in either experiment.*

Behavior	2020 (Snake vs. Chipmunk)		2021 (Snake vs. Hawk)	
	t	P	t	P
Hits	0.84	0.40	-0.27	0.79
Flyovers	-0.02	0.98	-0.15	0.88
Time within 5 m	0.10	0.93	-0.27	0.79
Closest approach	0.72	0.48	-0.10	0.93
Latency to return	0.04	0.97	-0.15	0.88

817 *Table S6 – Female nest defense behavior in Experiment 1 did not predict nest success.*

Behavior	z	P
Hits	-0.80	0.42
Flyovers	0.90	0.37
Time within 5 m	1.22	0.22
Closest approach	0.18	0.86
Latency to return	-0.36	0.72

831 *Table S7 – Analysis of predator order effects and interactions between predator type and order*
832 *for Experiment 2.*
833

Behavior	Day of the Year Effect		Order Effect		Interaction with Order	
	χ^2	*P*	χ^2	*P*	χ^2	*P*
Hits	0.86	0.35	1.18	0.28	1.36	0.24
Flyovers	2.88	0.09	1.91	0.17	1.51	0.22
Time within 5 m	2.73	0.10	3.70	0.055	0.48	0.49
Closest approach	2.01	0.16	2.23	0.14	1.14	0.29
Latency to return	1.42	0.23	<0.01	0.93	<0.01	0.98

835 *Table S8 – Focusing only on female responses to the snake decoy and pooling samples from*
836 *both experiments, we again found no detectable relationships between clutch size and the*
837 *strength of the nest defense response in house wrens.*

Behavior	*t*	*P*
Hits	-0.40	0.69
Flyovers	0.99	0.33
Time within 5 m	0.26	0.80
Closest approach	-0.31	0.76
Latency to return	-0.29	0.77

849   *Table S9 – Focusing only on female responses to the snake decoy and pooling samples from*
850   *both experiments, we again found no detectable relationships between nestling age and the*
851   *strength of the nest defense response in house wrens. However, there was a trend towards*
852   *females spending less time within 5 m of the snake decoy as nestlings aged.* [‡]*P<0.10.*

Behavior	t	P
Hits	0.01	0.99
Flyovers	-0.77	0.45
Time within 5 m	-1.70	0.09[‡]
Closest approach	1.20	0.23
Latency to return	0.18	0.86