Title: Differential modulation of cortical codes for sounds of varying intensity by distinct inhibitory neurons.

Abbreviated title: Inhibitory modulation of cortical codes

Authors: Melanie Tobin[1], Janaki Sheth[1], Katherine C. Wood[1], Erin K. Michel[1], and Maria N. Geffen[1,2,3]

Affiliations:
1. Department of Otorhinolaryngology, University of Pennsylvania, Philadelphia, PA 19104, United States
2. Department of Neuroscience, University of Pennsylvania, Philadelphia, PA 19104, United States
3. Department of Neurology, University of Pennsylvania, Philadelphia, PA 19104, United States

Corresponding author: Maria N. Geffen, mgeffen@pennmedicine.upenn.edu

CONFLICT OF INTEREST

The authors declare no competing interests.

ACKNOWLEDGEMENTS

The authors thank Dr. Anna Schapiro and members of the Geffen laboratory for helpful discussions and advice. This work was supported by NIDCD R01DC015527, R01DC014479, NINDS R01NS113241 to MNG and NIDCD K99DC019504 to KCW.

ABSTRACT

Cortical circuits contain multiple types of inhibitory neurons which shape how information is processed within neuronal networks. Here, we asked whether somatostatin-expressing (SST) and vasoactive intestinal peptide-expressing (VIP) inhibitory neurons may have distinct effects on population neuronal responses to noise bursts of varying intensities. We optogenetically stimulated SST or VIP neurons while simultaneously measuring the calcium responses of populations of hundreds of neurons in the auditory cortex of male and female awake, head-fixed mice to sounds. Upon SST neuronal activation, noise bursts representations became more discrete for different intensity levels, relying on cell identity rather than strength. In contrast, upon VIP neuronal activation, noise bursts of different intensity level activated overlapping neuronal populations, albeit at different response strengths. At the single-cell level, SST and VIP neuronal activation differentially activated the response-level curves of monotonic and nonmonotonic neurons. SST neuronal activation effects were consistent with a shift of the neuronal population responses toward a more localist code with different cells responding to sounds of different intensity. By contrast, VIP neuronal activation shifted responses towards a more distributed code, in which sounds of different intensity level are encoded in the relative response of similar populations of cells. These results delineate how distinct inhibitory neurons in the auditory cortex dynamically control cortical population codes. Different inhibitory neuronal populations may be recruited under different behavioral demands, depending on whether categorical or invariant representations are advantageous for the task.

SIGNIFICANCE

Information about sounds is represented in the auditory cortex by neuronal population activity that has a characteristic sparse structure. Cortical neuronal populations comprise multiple types of excitatory and inhibitory neurons. Here, we find that activating different types of inhibitory neurons differentially controls population neuronal representations, with one type of inhibitory neurons increasing the differences in the identity of the cells recruited to represent the different sounds, and another inhibitory neuron type changing the relative activity level of overlapping neuronal populations. Such transformations may be beneficial for different types of auditory behaviors, suggesting that these different types of inhibitory neurons may be recruited under different behavioral constraints in optimizing neuronal representations of sounds.

INTRODUCTION

Sensory cortical neuronal networks are comprised of multiple subtypes of neurons, including excitatory and inhibitory neurons. Inhibitory neurons can be further divided into multiple sub-classes, including somatostatin-expressing (SST) and vasoactive intestinal peptide-expressing (VIP) neurons, which mutually inhibit each other (Campagnola et al., 2022). The activity of these neurons modulates stimulus representations in auditory cortex (AC). Specifically, activating SST neurons reduces and decorrelates cortical activity (Chen et al., 2015), sharpens frequency tuning of AC neurons (Phillips and Hasenstaub, 2016) and contributes to surround suppression (Lakunina et al., 2020) and adaptation to stimulus context (Natan et al., 2015, 2017). By contrast, VIP neurons disinhibit excitatory neurons (Millman et al., 2020), largely via their projections onto SST neurons (Pfeffer et al., 2013; Pi et al., 2013), without affecting frequency tuning (Bigelow et al., 2019) and can enable high-excitability states in the cortex (Jackson et al., 2016). Both SST and VIP neurons can be modulated by noradrenergic and cholinergic inputs (Kawaguchi and Shindou, 1998; Fanselow et al., 2008; Chen et al., 2015) for SST neurons, and multiple neuromodulators for VIP neurons (Fu et al., 2014; Zhang et al., 2014; Chen et al., 2015) and therefore may serve differential modulatory functions in the context of different behavioral demands.

Sound pressure level representation supports sound detection, hearing in noise, source localization and distance to target calculation (Litovsky and Clifton, 1992). Most neurons in AC respond selectively to sounds at different sound pressure levels, either in a monotonic or a non-monotonic fashion. Monotonic neurons increase their firing rate with sound intensity, differing in their threshold and slope of the response functions. Non-monotonic neurons exhibit preference for specific sound pressure level ranges, differing in their preferred sound pressure level (Zhang et al., 2013). Previous work found that a mix of monotonic and non-monotonic neurons in the auditory cortex is important for sound encoding (Sun et al., 2017). Because the excitatory neurons in the cortex form tightly connected circuits with inhibitory neurons, inhibitory neuronal activity can shift the sound level response functions of excitatory neurons across monotonic and non-monotonic neurons. These changes can in turn affect the representation of sound pressure level by cortical populations. Whereas multiple studies have examined the effects of SST and VIP neuronal modulation on sound responses in individual neurons in AC (Natan et al., 2015; Seybold et al., 2015; Phillips and Hasenstaub, 2016; Natan et al., 2017; Bigelow et al., 2019; Millman et al., 2020; Seay et al., 2020), their effect on population representation of sound pressure level remains to be fully understood.

Here, we studied whether and how cortical inhibitory neurons control the representation of sound pressure levels in populations of neurons, by presenting periodic noise bursts at different sound pressure levels to awake head-fixed mice and imaging Calcium responses in populations of hundreds of neurons while simultaneously activating SST or VIP neurons optogenetically (Figure 1B, Figure 2). First, we tested whether and how activation of SST or VIP neurons differentially modulated sound pressure level responses at the level of individual neuronal response functions in AC. Next, we tested how the representation of sound pressure level changed in the neuronal space with and without SST and VIP neuronal activation. We tested for the effects of SST and VIP neuronal activation on response sparseness and separation angle of population response vectors. Finally, we tested whether and how changes in the response-level curves of monotonic and nonmonotonic individual neurons upon SST or VIP neuronal activation mediated the changes in representation at the population for monotonic and nonmonotonic neurons. Our results suggest that SST and VIP neuronal activation differentially affect both monotonic and non-monotonic neuronal sound pressure level response functions, thereby shifting the neuronal population codes between localist and distributed representations.

METHODS

Animals

We performed experiments in fourteen adult mice (7 males and 7 females), which were crosses between Cdh23 mice (B6.CAST-Cdh23[Ahl+/]Kjn, JAX: 002756) and Sst-Cre mice (Sst[tm2.1(cre)Zjh]/J, JAX: 013044; n=5 in experimental group) or Vip-IRES-Cre mice (Vip[tm1(cre)Zjh]/J, JAX: 010908; n=4 in experimental group, n=5 in control group) (Table 1). Mice had access to food and water ad libitum and were exposed to light/dark on a reversed 12h cycle at 28°C. Experiments were performed during the animals' dark cycle. Mice were housed individually after the cranial window implant. All experimental procedures were in accordance with NIH guidelines and approved by the Institutional Animal Care and Use Committee at the University of Pennsylvania.

Surgery procedures

Mice were implanted with cranial windows over Auditory Cortex following a published procedure (Wood et al., 2022). Briefly, mice were anesthetized with 1.5-3% isoflurane and the left side of the skull was exposed and perforated by a 3mm biopsy punch over the left Auditory Cortex. We injected in that region 3x750nL of an adeno-associated virus (AAV) mix of AAV1.Syn.GCaMP (6m: Addgene 100841 or 7f: Addgene 104488; dilution 1:10 ~ $1x10^{13}$ GC/mL) and AAV1.Syn.Flex.Chrimson.tdTomato (UNC Vector Core; dilution 1:2 ~ $2x10^{12}$ GC/mL). In the control mice, we injected a mix of AAV1.Syn.jGCaMP7f (Addgene 104488; dilution 1:10 ~ $1x10^{13}$ GC/mL) and AAV1.Syn.Flex.tdTomato (Addgene 28306; dilution 1:100-1:20 ~ $2x10^{11}$ – $1x10^{12}$ GC/mL) in VIP-Cre mice. We then sealed the craniotomy with a glass round window, attached a head plate to the mouse and let it recover for 3-4 weeks. After habituating the mouse to being head fixed for 3 days, we mapped the sound-responsive areas of the brain and located Auditory Cortex using wide field imaging, then performed two-photon imaging in Auditory Cortex (Figure 2C).

Two-photon imaging

We imaged calcium activity in neurons in layer 2/3 of Auditory Cortex of awake, head-fixed mice (VIP-Cre mice: 3321 neurons over 16 recordings, SST-Cre mice: 2284 neurons over 13 recordings) using the two-photon microscope (Ultima *in vivo* multiphoton microscope, Bruker) with a laser at 940nm (Chameleon Ti-Sapphire). The fluorescence from the tissue went through a Primary Dichroic long pass (620 LP), through an IR Blocker (625 SP), through an Emission Dichroic Long pass (565 LP) which separated the light in two beams. The shorter wavelengths went through an additional bandpass filter (525/70) before being captured by a PMT ("green channel"); the longer wavelengths went through a bandpass filter (595/50) before being captured by a PMT ("red channel"). This set up was used to minimize the contamination of the green channel by the optogenetic stimulus at 635nm. There was nevertheless some bleedthrough during the optogenetic stimulus which was small enough not to saturate the green channel, and thus the activity of neurons could be recorded continuously without interruption during optogenetic stimulation. We checked there was no saturation for the highest laser power used by plotting the average grayscale profile over the dimension of the image perpendicular to the scanning, and verifying it was well below saturation. During a 5-ms laser pulse, while the whole field of view is illuminated by the laser, it appears on the image only for the lines that were being scanned during the laser stimulus. These bands were identified using the average grayscale value of each image line, and the contaminated pixels inside these bands were removed before processing the recordings with Suite2p. We imaged a surface of 512x512 pixels^2 at 30Hz. If we recorded from a mouse several times, we changed the location or depth within layer 2/3 of Auditory Cortex in order to not image the same neurons twice.

Optogenetic laser: Power calibration

We first calibrated the laser power by measuring the curve of command voltage versus output power for the laser (Optoengine LLC, MRL-III-635-300mW). The laser's peak frequency was 635nm. Prior to every recording, we calibrated the laser power at the tissue level as follows: we used an empty cannula to reduce the power of the laser by a factor 10-15 and positioned the optical fiber on the objective so it would shine a spot of 1mm diameter centered on the focal point of the objective. Thus, the calibrated power at the imaging plane was for the medium laser power: 0.3 ± 0.09 mW/mm^2 (mean $\pm$ std, n=29; range: 0.14-0.47 mW/mm^2) and for the high laser power: 3.4 ± 1.0 mW/mm^2 (mean $\pm$ std, n=29; range: 1.6-5.3 mW/mm^2).

Identification of interneurons being stimulated

We started each recording by taking a 2600 frame video both in the green and the red channels (thus imaging GCaMP and tdTomato). As tdTomato is not dependent on the cell activity, any modulation in the signal in the red channel is due to bleedthrough from the GCaMP. We plotted for all cells the raw signal from the red channel versus the signal from the green channel and did a linear fit to extract the bleedthrough coefficient. We then subtracted the bleedthrough in the red signal and calculated the average fluorescence of the processed red signal for every cell. We then z-scored the signal of the red channel to the background fluorescence and selected the cells with a fluorescence higher than 2 σ (standard deviation of the background) as the targeted interneurons. The percentage of cells labeled as VIP or SOM interneurons with this criterion was consistent with the percentage of VIP or SOM expected within cortex (Rudy et al., 2011).

Stimulus presentation

We presented combinations of sound and optogenetic stimuli. The auditory stimulus consisted in 1-s long click trains of 25-ms pulses of broadband white noise (range 3−80 kHz) at 10Hz, at 7 sound pressure levels within 0-90 dB SPL (0; 30; 50; 60; 70; 80; 90 dB SPL). The optogenetic stimulus consisted in a 1-s long pulse train of 635nm laser with 5-ms pulses at 20 Hz, at 3 amplitudes with no, medium or high laser power (see power at tissue level in section Optogenetic laser: Power calibration). The two stimuli were presented simultaneously, with the optogenetic stimulus preceding the sound stimulus by 20 ms (Blackwell et al., 2020) for maximal optogenetic effect, the inter-stimulus interval was 5 s. All 21 combinations of sound and optogenetic stimuli were presented randomly and with 10 repeats per combination.

Analysis of single-cell activity: Optimal time window versus fixed time window

For each trial of a stimulus, the response was defined as the mean $\Delta F/F_{std}$ over the baseline one-second window fluorescence $F_{baseline}$ preceding the stimulus: $\Delta F/F_{std} = (F - \text{mean}(F_{baseline}))/\text{std}(F_{baseline})$, with F the fluorescence of the cell. *Optimal window:* In order to compute the best responses across different neurons, which may respond with different time courses of calcium signals, we defined the optimal window of neuronal response of each neuron for each stimulus combination as the one-second window for which the average response most reliably differs from the baseline activity. The optimal time window was selected as the 1-s averaging window which maximized the sensitivity index (d') between [0-1s] and [4-5s]. The optimal time window was only used in Figure 2, panels A,D,E,F,G,K,L,N. *Fixed window:* In order to compare how neuronal responses changed with laser stimulation, keeping all parameters similar besides that one, we defined the fixed window of neuronal response for each recording (one window for all stimulus combinations) as the one-second window with the largest number of responsive neurons. The fixed time window was selected as the 1-s averaging window which maximized the number of cells with a significant response to at least one of the stimuli pairs for each recording compared to the pre-stimulus fluorescence (paired t-test, p<0.01 with multiple comparison correction).

179 **Sparseness of a neuron's response and activity sparseness of a population of neurons**

180 To quantify how many stimuli a neuron responds to, we calculated the sparseness of each neuron adapted
181 from (Vinje and Gallant, 2000):

$$S = \frac{1}{1-\frac{1}{n}}\left(1 - \frac{\left(\Sigma\frac{r_i}{n}\right)^2}{\Sigma\left(\frac{r_i^2}{n}\right)}\right),$$

where r_i is the average response of a neuron to the ith sound pressure level calculated over its optimal time
window minus the neuron's lowest response to any combination of sound pressure level and laser activation,
and n is the number of sound pressure levels. We subtracted the lowest response of each neuron to the
response at a given sound amplitude and laser power to adapt this measure to fluorescence data. Indeed,
this sparseness measure can only be computed with positive responses and is usually computed with the
firing rate of each neuron (Rolls and Tovee, 1995; Vinje and Gallant, 2000). A sparseness value of 0%
indicates that a neuron's responses to all sound pressure levels are equal, and a sparseness value of 100%
indicates that a neuron only responds to one sound pressure level.

To quantify how many neurons in a population are active in response to a given stimulus, we calculated the
activity sparseness of each population (Willmore and Tolhurst, 2001) at a given sound level pressure and
laser power (Figure 2E and L) as the ratio of neurons that had an increase in response above threshold from
silence at the same laser power. The activity sparseness from silence and no laser was computed by
subtracting each neuron's response over its optimal window to its response at 0dB and no laser power, and
computing the ratio of neurons with an increase in response above threshold (Figure 2G and N). The
threshold was set as the standard deviation of the population's response to no sound and no laser power and
the response was calculated over each neuron's optimal time window. An activity sparseness value of 0%
indicates that all neurons in a population are active, and a value of 100% indicates that none of the neurons
are active.

Separation angle and Vector length

To quantify whether mean population vectors were collinear in the neuronal space, we calculated the
separation angle between mean population vectors adapted from (Vinje and Gallant, 2000). For each
recording, we computed the mean population vectors over the fixed window time at each laser power from
0dB to each non-zero sound pressure level (Figure 3A). We then computed the angle between each pair of
mean population vectors at a given laser power (Figure 3B), and represented the mean ± s.e.m (Figure 3D-
E and G-F) or the mean difference in separation angle from a given laser power to no laser power (Figure
3F and I) across recordings for each sound pair. To quantify whether mean population responses were close
in the neuronal space, we computed the vector length between mean population vectors (Figure 3C). For
each recording, we computed the mean population vectors over the fixed window time at each laser power
between all pairs of sound pressure levels (Figure 3A). We then computed the Euclidian norm of each mean
population vector at a given laser power (Figure 3C), and represented the mean ± s.e.m (Figure 3J-K and
M-N) or the mean difference in vector length from a given laser power to no laser power (Figure 3L and O)
across recordings for each sound pair.

Fitting of response-level curves

Mean response curves and the standard error of the mean (s.e.m) for every neuron were determined by
averaging over the fixed-time window its responses to all 10 trials of each sound pressure level. Thus, for
a   given   cell   we   constructed   three   response   curves,   one   for   every   light   condition.

To characterize responses as monotonic or nonmonotonic, we first normalized the response curves such that abs(max(response)) ≤ 1 and computed the monotonicity index (MI). This metric refers to the relative responses at higher stimulus levels (Watkins and Barbour, 2011) and was calculated from the mean curve as

$$MI = \frac{|response_{max_{level}}| - |response_{spontaneous}|}{|max(response) - min(response)|} \quad (1)$$

where $response_{max_{level}}$ is the response to 90 dB which is the highest level of sound presented and $response_{spontaneous}$ is the spontaneous response measured at 0 dB. A response curve was classified as nonmonotonic if its MI was less than 0.3 and monotonic if it was greater than 0.7. We refrain from a hard cutoff at 0.5 since preliminary analysis of the response curves indicated that due to stochasticity both monotonic and nonmonotonic curves may have MI values between 0.3 and 0.7. Furthermore, note that a given cell could change its monotonicity in the presence of optogenetic stimulation.

After determining the monotonicity of the neuronal response, we fitted the monotonic and nonmonotonic curves with a 4-parameter sigmoid function and a 4-parameter Gaussian function, respectively. The sigmoid function is given by the equation,

$$y = y_0 + \frac{y_{range}}{1 + e^{\frac{x_0 - soundlevel}{\Delta x}}} \quad (2)$$

while the Gaussian function can be written as,

$$y = y_0 + y_{range} * e^{\frac{-(soundlevel - x_{mean})^2}{2*\sigma^2}} \quad (3)$$

y refers to the response curve, y_0 is the offset response, y_{range} is its range in amplitude, x_0 is the x value of the sigmoid midpoint and Δx denotes the width of the sigmoid. In the Gaussian, the parameter y_0 is the offset response. The amplitude, mean and standard deviation of the Gaussian are denoted by y_{range}, x_{mean} and σ, respectively, and have their regular interpretations. During the fitting procedure, we minimize (1 – McFadden pseudo R-squared) using the Powell optimizer (scipy.minimize.optimize in Python). The formula for this error value is,

$$1 - R^2 = \frac{ln\hat{L}(M_{full})}{ln\hat{L}(M_{intercept})} \quad (4)$$

Assuming $\hat{L}(M_{full})$ is gaussian with the experimentally computed response average as its mean value and the response s.e.m as its standard deviation, we can rewrite the formula as,

$$1 - R^2 = \frac{\sum \frac{(mean(response) - response)^2}{sem^2}}{\sum \frac{(y - response)^2}{sem^2}} \quad (5)$$

We chose the McFadden R-squared since it allows us to account for different values of s.e.m at the different intensities. The regular R-squared equation constrains the s.e.m values to be equal at all intensities. A cell's response curve is considered well fit by its respective function if the Mcfadden R^2 is greater than 0.8. Due to the nonlinear nature of our optimization we chose 16 random starting points for the optimizer and cells fitted using 2 or more of the starting conditions were characterized using the fitting curve which had the

highest R^2 value. For neurons whose mean response curve MI lay between 0.3 and 0.7 we follow a similar
procedure but fit the curve with both the sigmoid and Gaussian functions to find the better fitting function.
Furthermore, we constrain the mean of all our Gaussian fits to lie between 10 and 80 dB. Response curves
with means less than 10 dB or greater than 80 dB were recharacterized using the sigmoid function since
only one sound pressure level data point (0 dB or 90 dB) is insufficient to adequately distinguish if the cell
is monotonic or nonmonotonic. Roughly ~5% of the total response curves (combined across all three light
conditions) were refitted in this manner. Lastly, we tested if the fitting curve was overfitted to the empirical
sound intensities by calculating a new variable – interpolated error. The interpolated sigmoid/Gaussian
curve is constructed by interpolating the fitted curve at the intermediate sound pressure levels – (15, 40, 55,
65, 75, 85) dB. The interpolated error is the regular R-squared value evaluated using the equation,

262
$$R^2 = 1 - \frac{\sum\left(y_{interpolatedlevels} - interpolated\,fitted\,curve\right)^2}{\sum\left(interpolated\,fitted\,curve - mean(interpolated\,fitted\,curve)\right)^2} . (6)$$

$y_{interpolatedlevels}$ refers to the Gaussian/sigmoid equations (2, 3) computed at the sound pressure levels 0,
15, 30, 40, 50, 55, 60, 65, 70, 75, 80, 85, and 90 dB. When computing statistics on different parameters we
remove neurons with interpolated error less than 0.25 for both the Sigmoid and Gaussian fits. This threshold
allows us to select at least 90% of the fitted curves.

Decoding sound pressure level using an SVM Decoder

We linearly decoded the 7 different sound pressure levels at each opto-stimulated condition using an SVM
decoder with a linear kernel. Specifically, we decoded each individual pressure level versus the remaining
six. Input to the SVM consisted of the fixed time-window responses of all neurons in the population.

To individually decode each of the 7 amplitudes, for every given experimental dataset we projected the
average responses of all neurons derived using a fixed window onto a lower-dimensional space using PCA.
The lower-dimensional space had (n) dimensions such that these dimensions accounted for 70% of the
variance in the dataset.

Next, because at a given opto-stimulated condition we had 10 trials per sound pressure level, the input data
to the SVM decoder was unbalanced as 10:60. We balanced the dataset by oversampling the sound pressure
level of interest, i.e., if we were decoding the 0dB stimulus from the rest, we oversampled to construct 60
trials for the 0dB stimulus. Specifically, we constructed these 60 trials by fitting the 10 experimentally
obtained 0dB trials using a Gaussian kernel and sampling the corresponding distribution. The resulting
oversampled dataset comprising 120 trials total was input into the two-class SVM decoder which was
trained using 10-fold cross-validation. Because we were not tuning the hyperparameters of the SVM, we
did not have separate validation and test sets, rather we computed the model's accuracy directly using the
validation set which was chosen randomly for each of the 10 iterations. Figure panels 2F and 2M illustrate
the decoding accuracies for all 3 conditions of opto-stimulation of SST and VIP interneurons.

Statistics

All responses are plotted as mean ± s.e.m (standard error of the mean) *with the number of measurements*
above the figure. We tested significance with a Generalized Linear Mixed Effects (GLME) Model with the
matlab function fitglme, using laser power, sound pressure level and the interaction term between laser
power and sound pressure level as fixed-effect terms and cell identity or session number as grouping
variables. All results from the statistical analyses are reported in Table 2.

RESULTS

SST and VIP neuronal activation modulate the response of sound-increasing neurons

To investigate whether and how distinct classes of inhibitory neurons in AC, SST and VIP neurons, affect sound representation at population level, we imaged Calcium activity of neurons in AC of awake, head-fixed mice presented with sounds while activating SST or VIP neurons using sub-millisecond optogenetic manipulation with Chrimson (Figure 1A) (Klapoetke et al., 2014). We monitored calcium activity by measuring fluorescence of GCaMP expressed in hundreds of neurons at a time (VIP-Cre mice: 3321 neurons over 16 recordings, SST-Cre mice: 2284 neurons over 13 recordings) and identified the cells expressing the opsin through co-expression of tdTomato (Figure 1B and 1C). This approach allowed us to quantify the transformations of sound representations within a large population of cortical neurons driven by SST or VIP neuronal activation.

FIGURE 1: EXPERIMENTAL SET UP AND OPTOGENETIC STIMULATION OF INTERNEURON POPULATIONS. A. Two-photon imaging and laser stimulation through the round window of a mouse injected with viruses encoding Syn.jGCaMP7f and Flex.ChrimsonR.tdTomato in the left Auditory Cortex. Flex.ChrimsonR.tdTomato is injected in AC of SST-Cre and VIP-Cre mice, and is activated by a 635-nm laser. The mouse lines used were SST-Cre x Cdh23+/+ and VIP-Cre x Cdh23+/+. A speaker delivers a broadband noise stimulus at sound pressure levels within 0-90 dB to the right ear. B. Cell tissue with two-photon imaging in the green channel (left) and cell identification (right) using Suite2p software, with yellow lines delineating cell borders, and red lines indicating the neurons expressing ChrimsonR.tdTomato. C. Left: Outline of the spread of the viral injection in a representative brain. Signal in the green channel (center

panels) and red channel (right panels) of an SST-Cre mouse (top panels) and an VIP-Cre mouse (bottom panels). Cells identified as VIP or SST interneurons are indicated with an arrow (see Methods). D. (Top) Diagrams for optogenetic manipulation in the circuit and experimental set-up. (Bottom left) Response of a SST neuron to no sound and sound stimulation (50 dB) when activating SST neurons with different laser powers. (Bottom right) Average fluorescence over a 1-s fixed time window as a function of sound pressure level for the example cell in the left panel. E. (Left) Response of a sound-increasing neuron to no sound and sound stimulation (70 dB) when activating SST neurons with different laser powers. (Right) Average fluorescence over a 1-s fixed time window as a function of sound pressure level for the example cell in the left panel. F. (Top) Diagrams for optogenetic manipulation in the circuit and experimental set-up. (Bottom left) Response of a VIP neuron to no sound and sound stimulation (70 dB) when activating VIP neurons with different laser powers. (Bottom right) Average fluorescence over a 1-s fixed time window as a function of sound pressure level for the example cell in the left panel. G. (Left) Response of a sound-increasing neuron to no sound and sound stimulation (70 dB) when activating VIP neurons with different laser powers. (Right) Average fluorescence over a 1-s fixed time window as a function of sound pressure level for the example neuron in the left panel. H. (Top) Experimental set-up for the control experiment – laser effect in the absence of an opsin. (Bottom) Average fluorescence over a 1-s fixed window as a function of sound pressure level for the whole population of VIP neurons recorded, tagged with Flex.tdTomato, when the laser illuminates AC. I. Average fluorescence over a 1-s fixed window as a function of sound pressure level for the whole population of neurons recorded (VIP neurons excluded) when the laser illuminates AC. For all panels, black, pink and red colors correspond to no laser power (0 mW/mm2), medium laser power (~0.3 mW/mm2) and high laser power (~3.5 mW/mm2), respectively (see Methods for power calibration). The gray and red bars below the example traces in panels D-G indicate the presence of the sound and laser stimulus, respectively.

We first confirmed that the optogenetic manipulations produced expected responses. SST neurons directly inhibit excitatory cells and other cells within the neuronal population, and so we expected that optogenetic manipulation would increase SST neuronal activity, but decrease the responses of other cells. By contrast, VIP neurons mostly inhibit other inhibitory neurons (Campagnola et al., 2022), and therefore we expected the VIP neuronal activation would increase both VIP neuronal activity and provide a release of inhibition to other cells in the network. In SST-Cre mice, a representative SST neuron increased activity at all sound pressure levels with laser power (example neuron, ***p_{laser}=1.8e-13, GLME, Figure 1D). The change in the response of a representative non-SST neuron to SST neuronal activation was sound level-dependent, with a decrease at most sound pressure levels for the medium laser power. Sound responses were abolished with strong SST neuronal activation at the high laser power (p_{laser}=0.74, ***$p_{laser:sound}$=1.0e-14, GLME, Figure 1E). In VIP-Cre mice, the response of a representative VIP neuron increased at all sound pressure levels with laser power and the increase was sound-level dependent (example neuron, ***p_{laser}=1.8e-7, *$p_{laser:sound}$=1.1e-2, GLME, Figure 1F). The activity of a representative non-VIP neuron increased at most sound pressure levels during activation of VIP neurons, with a larger increase at the high than medium laser power (*p_{laser}=1.5e-2, GLME, Figure 1G). As a control, we injected mice with Flex.tdTomato instead of the opsin in VIP-Cre x Cdh23 mice (n=5), and verified that laser stimulation of AC in the absence of the opsin did not lead to significant changes in the neuronal responses of VIP neurons (p_{laser}=0.20, GLME, Figure 1H) and non-VIP neurons (p_{laser}=0.64, GLME, Figure 1I). These representative effects of SST or VIP neuronal activation are consistent with previous reports, suggesting that the activation method worked as expected(Natan et al., 2015; Seybold et al., 2015; Phillips and Hasenstaub, 2016; Bigelow et al., 2019).

Differential distribution of population activity and sparseness with SST and VIP neuronal activation

We next tested whether and how SST and VIP neuronal activation differentially affects the sound pressure level response functions of neurons in AC. We characterized the response of each cell over its optimal time window for each sound and laser combination (Figure 2 panels A and H; see Methods) and the response of each population to any stimulus combination over its fixed time window (Figure 2 panels B-C and I-J, see Methods).

FIGURE 2: POPULATION RESPONSE TO SST AND VIP ACTIVATION AND SPARSENESS A. Rasters of the average fluorescence versus sound pressure level for all neurons imaged in the SST-Cre mice, calculated over the optimal time window for each cell and each sound and laser condition (see Methods). Rasters from left to right correspond to SST activation with no laser power, medium laser power and high laser power. The thick blue line at the bottom of each raster indicates the SST interneurons. Cells are ordered given their response at 90dB and no laser power. B. Average fluorescence over a 1-s fixed window as a function of sound pressure level for the whole population of SST neurons recorded (132 neurons), when the SST population is activated. C. Average fluorescence over a 1-s fixed window as a function of sound pressure level for the whole population of non-SST neurons recorded (2152 neurons), when the SST population is activated. D. Cumulative distribution function of sparseness for the population of non-SST neurons, when the SST population is activated. E. Average activity sparseness as a function of sound pressure level for each population of neurons (SST neurons excluded), when the SST population is activated. F. Decoding accuracy of SVM decoder at each laser power, decoding individual sound pressure levels using non-SST neuronal responses within 1-s fixed window, when the laser illuminates AC and stimulates SST neurons. G. Average activity sparseness measured from silence and no laser as a function of sound pressure level for each population of neurons (SST neurons excluded), when the SST population is activated. There was no significant change in activity sparseness measured from silence and no laser upon SST activation (p_{laser}=0.11, GLME). The point at 0dB for no

laser power was by design 100% and thus omitted from the plot and the statistical test. H. Rasters of the average fluorescence versus the sound
pressure level for all neurons imaged in the VIP-Cre mice, calculated over the optimal time window for each cell and each sound and laser
condition (see Methods). Rasters from left to right correspond to VIP activation with no laser power, medium laser power and high laser power.
The thick green line at the bottom of each raster indicates the VIP interneurons. Cells are ordered given their response at 90dB and no laser
power. I. Average fluorescence over a 1-s fixed window as a function of sound pressure level for the whole population of VIP neurons recorded
(226 neurons), when the VIP population is activated. J. Average fluorescence over a 1-s fixed window as a function of sound pressure level
for the whole population of non-VIP neurons recorded (3095 neurons), when the VIP population is activated. K. Cumulative distribution
function of sparseness for the population of non-VIP neurons, when the VIP population is activated. L. Average activity sparseness as a function
of sound pressure level for each population of neurons (VIP neurons excluded) when the VIP population is activated. M. Decoding accuracy
of SVM decoder at each laser power, decoding individual sound pressure levels using non-VIP neuronal responses within 1-s fixed window,
when the laser illuminates AC and stimulates VIP neurons. N. Average activity sparseness measured from silence and no laser as a function of
sound pressure level for each population of neurons (VIP neurons excluded), when the VIP population is activated. There was a significant
decrease in activity sparseness measured from silence and no laser upon VIP activation (***p_{laser}=4.9e-4, GLME). The point at 0dB for no laser
power was by calculation 100% and thus omitted from the plot and the statistical test. For all panels, black, pink and red colors correspond to
no laser power (0 mW/mm2), medium laser power (~0.3 mW/mm2) and high laser power (~3.5 mW/mm2), respectively (see Methods for power
calibration).

Because SST neurons directly inhibit excitatory cells, we hypothesized that SST neuronal activation would
lead to fewer neurons responding at increasing sound pressure levels. During the SST neuronal activation,
the average response of SST neurons increased at all sound pressure levels (n=132 SST neurons,
***p_{laser}<1e-100, GLME, Figure 2A-B), whereas non-SST neurons exhibited a mix of decreased and
increased responses (Figure 2A). The overall effect of SST neuronal activation on the population of non-
SST neurons had a significant interaction between sound and laser amplitude, but no significant sound-
independent laser effect (n=2152 neurons, p_{laser}=0.20, ***$p_{laser:sound}$=2.2e-10, GLME, Figure 2C). The
average response-level curve for non-SST neurons shifted downwards for the medium laser power, and at
a high laser power, the modulation of the population's response by sound was lost with an average response
to silence equal to the average response to sounds, consistent with expectations for a sparser. more localist
population code (Polley et al., 2004).

To further assess the representation of sound pressure level in the neuronal population, we studied two
characteristics of sparse distributed representations: (1) each neuron responds only to a few stimuli (high
sparseness) and (2) only a few neurons respond to each stimulus (high activity sparseness). To measure
how many stimuli a neuron responds to, we computed the sparseness of each non-SST neuron (see
Methods). The sparseness of non-SST neurons increased for 70% of the neurons, with a significant increase
from 31% to 33% upon SST neuronal activation (median, n=2152 neurons, ***p_{laser}=7.1e-4, GLME, Figure
2D), indicating that neurons responded to fewer stimuli. To measure how many neurons responded to each
stimulus, we computed the activity sparseness for each population of non-SST neurons, which is the ratio
of neurons that are *not* active in response to a given stimulus, compared to silence at a given laser power.
The activity sparseness increased significantly upon SST neuronal activation (n=13 populations,
**p_{laser}=1.6e-3, GLME, Figure 2E), indicating that at each successive sound pressure level, there were fewer
non-SST neurons that were active. However, whereas decoding accuracy differed across different sound
pressure levels, and was slightly lower with SST neuronal inactivation, there was no interaction between
the laser and sound amplitudes (n=13 populations, p_{sound}=0.0049, p_{laser} =1.11e-4, $p_{laser:sound}$=0.056, GLME,
Figure 2F). This suggests that despite the neuronal population responses becoming relatively sparser across
sound pressure levels, the relative decoding accuracy was preserved across SPLs. Combined these results
point to a more localist representation for sound pressure level with SST neuronal activation.

Because VIP neuron activity provides a release of SST neuron inhibition on excitatory cells, we
hypothesized that VIP neuronal activation would lead to more neurons being active in response to each
sound level and having stronger responses. When VIP neurons were activated, the average response of VIP

neurons increased at all sound pressure levels (n=226 VIP neurons, ***p_{laser}<1e-100, GLME, Figure 2H-I). Non-VIP neurons similarly exhibited an increase in response, both in silence and to sounds at different sound pressure levels (Figure 2H). The average response-level curve for non-VIP neurons shifted upwards at all sound pressure levels as VIP neurons were activated (n=3095 neurons, ***p_{laser}<1e-100, GLME, Figure 2J), reflective of a more distributed stimulus representation. At all laser powers, the modulation of the population's response by sound was maintained: the population average to sounds was higher than to silence.

Neuronal sparseness decreased for all neurons with VIP neuronal activation, with a significant decrease from 32% to 22% upon VIP neuronal activation (median, n=3095 neurons, ***p_{laser}<1e-100, GLME, Figure 2K), indicating that each neuron responded more equally to the different sound pressure levels. The activity sparseness measured from silence at a given laser power did not change upon VIP neuronal activation (n=16 populations, p_{laser}=0.44, GLME, Figure 2L), indicating that the same number of neurons showed an increase in response at each sound pressure level from silence at a given laser power. Consistent with the overall increase in responses with VIP neuronal activation in silence, the activity sparseness measured from silence at no laser power significantly decreased (Figure 2N). Importantly, this change in population activity did not affect the decoding performance (n=13 populations, p_{sound}=3.99e-4, p_{laser}=0.49, $p_{laser:sound}$=0.91, GLME, Figure 2M). Combined, these results suggest that activating VIP neurons transforms population responses to a more distributed representation, while preserving the decoding accuracy.

Overall, SST neuronal activation led to weaker and sparser responses in the population, with neurons responding to fewer stimuli and each stimulus eliciting a response in fewer neurons, shifting the population responses toward a more localist stimulus representation. By contrast VIP neuronal activation leads to a global increase in the neuronal population's response, along with each neuron responding to more stimuli, leading to a more distributed stimulus representation.

Sound pressure level is represented more discretely or continuously in the neuronal population with SST or VIP neuronal activation, respectively.

There are various ways that a neuronal network can implement a representation of a sensory feature. For example, a distributed code may rely on the magnitude of response of the population of neurons or on the relative response of each cell. To investigate this, we examined next the properties of the representation of sound pressure level upon SST and VIP neuronal activation in the neuronal space. We computed the mean population vector from 0dB at a given laser amplitude to each nonzero sound pressure level at that laser amplitude (Figure 3A), and computed the separation angle between pairs of mean population vectors at the same laser power (Figure 3B), as well as the length of mean population vectors between pairs of sounds (Figure 3C).

FIGURE 3: SEPARATION ANGLE AND VECTOR LENGTH ARE DIFFERENTIALLY CONTROLLED BY SST AND VIP ACTIVATION. A. Schematic representation of the population's response to sound and laser stimulation, for single trials (small dots) and the average response over all trials (large dots). Sound pressure level is represented by the shading of the dot, laser power by the outline of the dot. We simplify the representation of the population's response to two dimensions, keeping only the color code for the laser power. B. Low-dimensional schematic of the separation angle between mean population vectors to each sound pressure level at a given laser power, starting from 0dB at each laser power. C. Low-dimensional schematic of the vector length of mean population vectors between sound pressure levels at a given laser power. D. Confusion matrix of the separation angle between pairs of sound stimuli for no (left), medium (middle) and high (right) laser power of SST activation. E. Separation angle between pairs of mean population vectors to different sound pressure levels as a function of the difference in sound pressure level for no, medium and high laser powers of SST activation (circles). The dotted lines are the result from the GLME fit at the different laser powers of SST activation. F. Confusion matrix of the difference in separation angle between pairs of sound stimuli from medium (left) or high (right) laser power to no laser power of SST activation. G. Confusion matrix of the separation angle between pairs of sound stimuli for no (left), medium (middle) and high (right) laser power of VIP activation. H. Separation angle between pairs of mean population vectors to different sound pressure levels as a function of the difference in sound level for no, medium and high laser powers of VIP activation (circles). The dotted lines are the result from the GLME fit at the different laser powers of VIP activation. I. Confusion matrix of the difference in separation angle between pairs of sound stimuli from medium (left) or high (right) laser power to no laser power of VIP activation. J. Confusion matrix of the vector length between pairs of sound stimuli for no (left), medium (middle) and high (right) laser power of SST activation. K. Length of the mean population vector between pairs of sound pressure levels as a function of the difference in sound pressure level for no, medium and high laser powers of SST activation (circles). The dotted lines are the result from the GLME fit at the different laser powers of SST activation. L. Confusion matrix of the difference in vector length between pairs of sound stimuli from medium (left) or high (right) laser power to no laser power of SST activation. M. Confusion matrix of the vector length between pairs of sound stimuli for no (left), medium (middle) and high (right) laser power of VIP activation. N. Length of the mean population vector between pairs of sound pressure levels as a function of the difference in sound pressure level for no, medium and high laser powers of VIP activation (circles). The dotted lines are the result from the GLME fit at the different laser powers of VIP activation. O. Confusion matrix of the difference in vector length between pairs of sound stimuli from

The separation angle (Vinje and Gallant, 2000) computes the angle between the mean population vectors to two different sound pressure levels taking 0dB at each laser power as the origin (Figure 3B). A smaller angle indicates the population vectors are more collinear, meaning similar neurons respond to the different stimuli, although perhaps with differing magnitude. A larger angle indicates that there is less overlap between the populations of neurons responding to each stimulus.

When there is no interneuron activation, the separation angle increased with the difference in sound pressure level (Figure 3E and H, black circles), indicating that there is less overlap in the groups of neurons responding to sounds with a large difference in sound pressure level than a small difference in sound pressure level. Upon SST neuronal activation, the curve flattened around 60°, meaning that there was an increase in separation angle for small differences in sound pressure level, and a decrease in separation angle for large differences in sound pressure level (dotted lines correspond to GLME estimates, ***p_{laser}=1.6e-17, ***$p_{\Delta sound}$=2.3e-31, ***$p_{laser:\Delta sound}$=3.6e-13, GLME, Figure 3D-E). On average, the difference in separation angle between sound pairs from medium or high laser to no laser was positive, at $3.7° \pm 1.7°$ for the high laser power (mean ± s.e.m, 15 angles, Figure 3F). This indicates that the population vectors to the different tested sound pressure levels were more equally distributed in the neuronal space. There is, however, still an overlap between groups of neurons responding to different sound pressure levels, as a fully orthogonal coding of sound pressure level would lead to a 90° separation angle. In contrast, upon VIP neuronal activation, the separation angle decreased equally for all differences in sound pressure level (dotted lines correspond to GLME estimates, **p_{laser}=6.1e-3, ***$p_{\Delta sound}$=6.9e-13, $p_{laser:\Delta sound}$=0.47, GLME, Figure 3G-H). On average, the difference in separation angle between sound pairs from medium or high laser to no laser was negative, at $-4.6° \pm 0.7°$ degrees at the high laser power (mean ± s.e.m, 15 angles, Figure 3I). This indicates that the population vectors to the different tested sound pressure levels were more collinear, with more overlap between groups of neurons responding to different sound pressure levels with VIP neuronal activation.

The vector length computes the Euclidian norm of the mean population vector between two sound pressure levels at a given laser power (Figure 3C). A small length indicates that the responses to two different stimuli are close in the neuronal space, either due to small magnitudes of response, to small differences in separation angle or both, whereas a large length indicates that there is a large difference in magnitude, in separation angle or both. Therefore, we tested whether SST and VIP neuronal activation differentially affected the vector length.

Upon SST neuronal activation, the vector length decreased for all sound pressure level differences to about 2 a.u. at high laser power, along with a decrease in the slope of the GLME estimate by 81% at high laser power (dotted lines correspond to GLME estimates, ***p_{laser}=2.9e-6, ***$p_{\Delta sound}$=1.6e-8, ***$p_{laser:\Delta sound}$=3.8e-4, GLME, Figure 3J-K). The average change in length from medium or high laser power to no laser was negative, at -1.00 ± 0.10 a.u. for the high laser power (mean ± s.e.m, 21 lengths, Figure 3L). Upon VIP neuronal activation, the vector length increased for all sound pressure level differences, and the slope of the GLME estimate also increased by 72% at high laser power (dotted lines correspond to GLME estimates, *p_{laser}=1.2e-2, ***$p_{\Delta sound}$=6.1e-5, ***$p_{laser:\Delta sound}$=1.5e-6, GLME, Figure 3M-N). The average change in length from medium or high laser power to no laser was positive, at 1.27 ± 0.12 a.u. for the high laser power (mean ± s.e.m, 21 lengths, Figure 3O).

Combined with the differences in the separation angle, these results point to emergence of two types of modulations of population codes: Upon SST neuronal activation, the encoding of sound pressure level resembles a localist pattern coding where the magnitude of response is less relevant than the identity of responding cells: the strength of response is reduced and similar for all sound pressure levels, but the population vector angles are more spread out in neuronal space, indicating that there is less overlap between groups of neurons responding to different sound pressure levels. Upon VIP neuronal activation, the encoding of sound pressure level resembles a rate code, which is a type of distributed representation in which the varying strength of the whole population encodes a continuously varying parameter of the stimulus. Therefore, VIP neuronal activation promotes the strength of response more than the identity of responding neurons: there is more overlap between the groups of neurons responding to different sound pressure levels, but the strength of response is increased.

FIGURE 4: SINGLE-CELL FITS OF RESPONSE-LEVEL CURVES FOR MONOTONIC CELLS. A. Example neuron with a monotonic response-level curve (solid black line) with a sigmoid fit estimated at the probed sound amplitudes (blue dashed line with circles) and the sigmoid function with the same parameters (dotted black line). The parameters for the sigmoid fit are: Offset amplitude $y_0 = 0.1$; Range $y_{range} = 2.4$; Midpoint $x_0 = 60$ dB; Width $\Delta x = 5$ dB. B. Example of changes to the response-level curve of a sound-increasing monotonic neuron upon SST activation. Response-level curves (solid lines) are fit by sigmoid functions (dotted lines) at no (black lines) and high (red lines) laser powers of SST activation. The parameters for the sigmoid fit are, for no SST activation: $y_0 = -0.3$; $y_{range} = 13.4$; $x_0 = 93$ dB; $\Delta x = 11$ dB; and for SST activation: $y_0 = -0.2$; $y_{range} = 8.6$; $x_0 = 96$ dB; $\Delta x = 8$ dB. C. Example of changes to the response-level curve of a sound-increasing monotonic neuron upon VIP activation. Response-level curves (solid lines) are fit by sigmoid functions (dotted lines) at no (black lines) and high (red lines) laser powers of VIP activation. The parameters for the sigmoid fit are, for no VIP activation: $y_0 = -0.1$; $y_{range} = 5.6$; $x_0 = 66$ dB; $\Delta x = 5$ dB; and for VIP activation: $y_0 = -0.4$; $y_{range} = 9.9$; $x_0 = 60$ dB; $\Delta x = 5$ dB. D. Schematic showing the offset amplitude y_0 parameter of the sigmoid fit (left panel) and cumulative distribution function of y_0 for monotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. E. Schematic showing the amplitude range y_{range} parameter of the sigmoid fit (left panel) and cumulative distribution function of y_{range} for monotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. F. Schematic showing the midpoint x_0 parameter of the sigmoid fit (left panel) and cumulative distribution function of x_0 for monotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. G. Schematic showing the width Δx parameter of the sigmoid fit (left panel) and cumulative distribution function of Δx for monotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. H. Schematic of the mean significant changes to the response-level curve of a monotonic sound-increasing cell (gray line) upon SST (blue line) and VIP (green line) activation. For all panels, black, pink and red colors correspond to no laser power (0 mW/mm2), medium laser power (~0.3 mW/mm2) and high laser power (~3.5 mW/mm2), respectively (see Methods for power calibration). For all panels, the statistical test GLME was performed on the distributions at the three different levels of interneuron activation, with n.s. corresponding to non-significant, * corresponds to p<0.05, ** corresponds to p<0.01, *** corresponds to p<0.001. There are n=109, n=103 and n=64 sound-increasing monotonic cells fit at no, medium and high laser powers of SST activation, respectively. There are n=267, n=239 and n=269 sound-increasing monotonic cells fit at no, medium and high laser powers of VIP activation, respectively.

FIGURE 5: SINGLE-CELL FITS OF RESPONSE-LEVEL CURVES FOR NONMONOTONIC CELLS A. Example neuron with a nonmonotonic response-level curve (solid black line) with a Gaussian fit estimated at the probed sound pressure levels (blue dashed line with circles) and the Gaussian function with the same parameters (dotted black line). The parameters for the Gaussian fit are: Offset amplitude $y_0 = 0.15$; Range $y_{range} = 3.8$; Mean $x_{mean} = 49$ dB; Standard Deviation $\sigma = 9$ dB. B. Example of changes to the response-level curve of a sound-increasing nonmonotonic neuron upon SST activation. Response-level curves (solid lines) are fit by Gaussian functions (dotted lines) at no (black lines) and high (red lines) laser powers of SST activation. The parameters for the Gaussian fit are, for no SST activation: $y_0 = 0.2$; $y_{range} = 3.8$; $x_{mean} = 49$ dB; $\sigma = 9$ dB; and for SST activation: $y_0 = 0.1$; $y_{range} = 0.8$; $x_{mean} = 47$ dB; $\sigma = 11$ dB. C. Example of changes to the response-level curve of a sound-increasing nonmonotonic neuron upon VIP activation. Response-level curves (solid lines) are fit by Gaussian functions (dotted lines) at no (black lines) and high (red lines) laser powers of VIP activation. The parameters for the Gaussian fit are, for no VIP activation: $y_0 = 0.0$; $y_{range} = 0.2$; $x_{mean} = 49$ dB; $\sigma = 4$ dB; and for VIP activation: $y_0 = 0.0$; $y_{range} = 0.4$; $x_{mean} = 66$ dB; $\sigma = 7$ dB. D. Schematic showing the offset amplitude y_0 parameter of the Gaussian fit (left panel) and cumulative distribution function of y_0 for nonmonotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. E. Schematic showing the amplitude range y_{range} parameter of the Gaussian fit (left panel) and cumulative distribution function of y_{range} for nonmonotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. F. Schematic showing the mean x_{mean} parameter of the Gaussian fit (left panel) and cumulative distribution function of x_{mean} for nonmonotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. G. Schematic showing the standard deviation σ parameter of the Gaussian fit (left panel) and cumulative distribution function of σ for nonmonotonic sound-increasing neurons at different laser powers of SST (middle panel) and VIP (right panel) activation. H. Schematic of the mean significant changes to the response-level curve of a nonmonotonic sound-increasing cell (gray line) upon SST (blue line) and VIP (green line) activation. For all panels, black, pink and red colors correspond to no laser power (0 mW/mm2), medium laser power (~0.3 mW/mm2) and high laser power (~3.5 mW/mm2), respectively (see Methods for power calibration). For all panels, the statistical test GLME was performed on the distributions at the three different levels of interneuron activation, with n.s. corresponding to non-

647 significant, * corresponds to p<0.05, ** corresponds to p<0.01, *** corresponds to p<0.001. There are n=224, n=175 and n=130 sound-
648 increasing nonmonotonic cells fit at no, medium and high laser powers of SST activation, respectively. There are n=243, n=278 and n=310
649 sound-increasing nonmonotonic cells fit at no, medium and high laser powers of VIP activation, respectively.

Response-level curves of sound-modulated cells exhibit a narrower response upon SST neuronal activation, and a broader response upon VIP neuronal activation.

We next tested how the shifts in stimulus representation mediated by SST and VIP neurons at the scale of the neuronal population are implemented at the single-cell level by analyzing the changes with SST or VIP neuronal activation in response-level curves of single neurons responding positively to sound. Some AC neurons exhibit increased responses with increased sound pressure levels (monotonic response-level curve) while others are tuned with a peak response to a specific sound pressure level (nonmonotonic response-level curve) (Schreiner et al., 1992; Phillips et al., 1995; Wu et al., 2006). We classified cells depending on their Monotonicity Index (MI, see Methods) and fit response functions of monotonically responding cells with a Sigmoid function (Figure 4 see Methods) and those of non-monotonically responding cells with a Gaussian function (Figure 5, see Methods). We then tested how the parameters of the fits change with interneuron activation.

We first characterized the responses of sound-modulated cells that exhibited a monotonic response-level curve by fitting this curve for individual cells at different levels of laser power (Figure 4A-C). Out of the four sigmoidal fit parameters (Figure 4D-G, middle panels), only the midpoint of the sigmoid fit exhibited a significant change upon SST neuronal activation from 58 dB at no laser power to 73 dB at high laser power (median values, n=109; 103 and 64 neurons for no, medium and high laser power, respectively; offset: p_{laser}=0.41; range: p_{laser}=0.22; midpoint: **p_{laser}=8.6e-3; width: p_{laser}=0.99; GLME, Figure 4D-G, middle panels). With VIP activation, only the range of the sigmoid fit showed a significant increase (n=267, 239 and 269 neurons for no, medium, and high laser power, respectively; offset: p_{laser}=0.081; range: **p_{laser}=1.9e-3; midpoint: p_{laser}=0.38; width: p_{laser}=0.57; GLME, Figure 4D-G, right panels). Thus, SST neuronal activation leads to monotonic response-level functions that are shifted rightwards towards higher sound pressure levels, leading to responses to a narrower range of sounds at higher sound pressure levels. VIP activation expanded the neuronal response-level curves upwards, leading to responses to a broader range of sound pressure levels (Figure 4H).

We then characterized the responses of sound-modulated cells that exhibited a nonmonotonic response-level curve by fitting this curve for individual cells at different levels of laser power (Figure 5A-C). Whereas the mean and standard deviation of the Gaussian fit remained unchanged (mean: p_{laser}=0.73; standard deviation: p_{laser}=0.53; GLME, Figure 5F-G, middle panels), the offset and the range of the Gaussian fit decreased with SST neuronal activation (n=224, 175 and 130 neurons for no, medium and high laser power, respectively; offset: **p_{laser}=1.7e-3; range: ***p_{laser}=9.4e-8; GLME, Figure 5D-E, middle panels). The offset of response did not change with VIP neuronal activation (offset: p_{laser}=0.48, GLME, Figure 5D, right panel), whereas the range increased significantly as well as the Gaussian mean and standard deviation (n=243, 278 and 310 neurons for no, medium and high laser power, respectively; range: ***p_{laser}=2.9e-8; mean: ***p_{laser}=8.6e-4; standard deviation: ***p_{laser}=8.1e-5; GLME, Figure 5E-G, right panels). Thus, SST neuronal activation led to nonmonotonic response-level functions that were shifted downwards with a decreased range of responses, leading to responses above noise level to a narrower range of sound pressure levels. VIP neuronal activation shifted the neuronal response-level curves rightwards with an increased range of response, leading to increased peak responses at higher sound pressure levels (Figure 5H).

690 These results demonstrated that SST neuronal activation promoted a more localist representation of sound
691 pressure level at the population scale by eliciting responses above noise level over a narrower range of
692 sound pressure levels, and increasing separation between the sound pressure levels covered by monotonic
693 and nonmonotonic neurons. By contrast, VIP neuronal activation promoted a more distributed
694 representation of sound pressure level by broadening response-level curves of single neurons and increasing
695 the overlap between the sound pressure levels covered by monotonic and nonmonotonic neurons.

696 Could the changes to the fitted parameters of sound-modulated cells (Figure 4 and Figure 5) explain the
697 differential effect of SST and VIP neuronal activation on the separation angle and length between mean
698 population vectors to different sound pressure levels (Figure 3)? To answer this question, we constructed a
699 qualitative model including a monotonic and a nonmonotonic cell, with response-level fit parameters for
700 no and high laser power taken as the mean parameters from the data in Figure 4 and Figure 5 for no and
701 high laser power (Figure 6A-B). With this simple two-cell population, we could qualitatively reproduce the
702 increase in separation angle upon SST neuronal activation and the decrease in separation angle upon VIP
703 activation over the range of sound pressure levels we tested (Figure 6D-F), and similarly the decrease in
704 vector length upon SST neuronal activation and the increase in vector length upon VIP (Figure 6G-I). Thus,
705 the changes to the fit parameters observed in Figures 4-5 can explain the change in representation of sound
706 pressure levels observed in Figure 3.

A Response-level curve of the monotonic neuron

B Response-level curve of the nonmonotonic neuron

C interneuron activation:
none
SST
VIP

D No activation   SST activation   VIP activation
Angle

E

F SST activation   VIP activation
Difference in angle

G No activation   SST activation   VIP activation
Length

H

I SST activation   VIP activation
Difference in length

J Two-cell Model Monotonic cell

K Two-cell Model Nonmonotonic cell

FIGURE 6: TWO-CELL MODEL A. Response-level curve of the monotonic cell with parameters taken as the mean parameters from Figure 4 at no and high laser power. The parameters for the sigmoid curve with no interneuron activation (black) are: Offset amplitude $y_0 = -0.12$; Range $y_{range} = 0.88$; Midpoint $x_0 = 55$ dB; Width $Dx = 11$ dB. Upon SST activation (blue), all parameters remain constant except for: Midpoint $x_0^{SST} = 68$ dB; Upon VIP activation (green), all parameters remain constant except for: Range $y_{range}^{VIP} = 1.43$. B. Response-level curve of the nonmonotonic cell with parameters taken as the mean parameters from Figure 5 at no and high laser power. The parameters for the Gaussian curve with no interneuron activation (black) are: Offset amplitude $y_0 = -0.04$; Range $y_{range} = 0.54$; Mean $x_{mean} = 50$ dB; Standard Deviation $s = 13$ dB. Upon SST activation (blue), all parameters remain constant except for: Offset amplitude $y_0^{SST} = -0.07$; Range $y_{range}^{SST} = 0.27$; Upon VIP activation (green), the offset amplitude remains constant, and: Range $y_{range}^{VIP} = 0.86$; Mean $x_{mean}^{VIP} = 54$ dB; Standard Deviation $s^{VIP} = 17$ dB. C. Trajectory of the population's response from 0dB to 90dB in the neural space, with the response of the monotonic cell on the x-axis and the response of the nonmonotonic cell on the y-axis. The response of both cells at 0dB has been subtracted from the curves, thus the dots at the (0,0) coordinate are the response to 0dB, and the end of the curves on the right indicate the response to 90dB. The trajectories are computed from 0 dB to 90dB with 1dB increments, and circles on a line represent 10dB increments from 0dB to 90dB. D. Confusion matrix of the separation angle between population responses to each sound and laser power from silence at a given laser power, for no (left), SST (middle) and VIP (right) activation. Sound pressure level is in 1dB increments, and the gray box indicates the sound levels sampled in the experiments (Figure 3D, and G) E. Schematic in the neural space (see panel C) of the angle between 50 dB and 70dB when there is no (black), SST (blue) or VIP (green) activation, starting from the population's response to silence for each case of (or lack of) interneuron activation. The angle is greatest when SST neurons are activated, and smallest when VIP neurons are activated. F. Confusion matrix of the difference in separation angle from SST (left) or VIP (right) activation to no interneuron activation, with the angles calculated as in (D). Sound level is in 1dB increments, and the gray box indicates the sound pressure levels sampled in the experiments (Figure 3F and I). The mean angle difference for SST activation is, over 1-90dB: $+ 3.6°$ and over 30-90dB: $+ 3.7°$; for VIP activation, over 1-90dB: $- 4.0°$, and over 30-90dB: $- 4.1°$. G. Confusion matrix of the length of the population vector between each sound pressure level at a given laser power, for no (left), SST (middle) and VIP (right) activation. Sound pressure level is in 1dB increments, and the gray box indicates the sound pressure levels sampled in the experiments (Figure 3J and M). H. Schematic in the neural space (see panel C) of the vector length between 50 dB and 70dB when there is no (black), SST (blue) or VIP (green) activation. The vector length is greatest when VIP neurons are activated, and smallest when SST neurons are activated. I. Confusion matrix of the difference in vector length from SST (left) or VIP (right) activation to no interneuron activation, with the lengths calculated as in (G). Sound pressure level is in 1dB increments, and the gray box indicates the sound pressure levels sampled in the experiments (Figure 3I and L). The mean length difference for SST activation is, over 1-90dB: $- 0.12$ a.u. and over 30-90dB: $- 0.07$ a.u.; for VIP activation, over 1-90dB, $+ 0.27$ a.u. and over 30-90dB: $+ 0.21$ a.u.J. Response of the monotonic cell with laser activation versus no laser activation. SST activation shows a divisive regime, a subtractive regime, a combination of divisive and subtractive or multiplicative and subtractive regimes depending on the range of responses sampled. VIP activation shows a multiplicative regime or an additive and multiplicative regime depending on the range of responses sampled. K. Response of the nonmonotonic cell with laser activation versus no laser activation. SST activation shows a combination of divisive and subtractive regime. VIP activation shows a multiplicative regime, and additive regime or an additive and multiplicative regime depending on the range of responses sampled. For all panels, black indicates no interneuron activation, blue indicates SST activation and green indicates VIP activation.

Overall, when neither SST or VIP neurons are activated, the neuronal population encoded different sound pressure levels using two strategies: the identity of the responsive cells (different cells respond to different sound pressure levels, discrete encoding of sound pressure level) and the strength of the neuronal response (continuous encoding of sound pressure level). When SST neurons are activated, the neuronal population shifts towards a more localist representation of sound pressure level. Specifically, the encoding of sound pressure level relies more on the identity of the responsive cells and less on the magnitude of response: there is less overlap between populations of cells responding to different sound pressure levels, but the strength of response is similar for all sound pressure levels. This can be explained with the narrower bandwidths of response, albeit of reduced magnitude, of both monotonic and nonmonotonic sound-increasing cells with SST neuronal activation. By contrast, when VIP neurons are activated, the neuronal population shifts towards a more distributed representation of sound pressure level. Specifically, the encoding of sound pressure level by the magnitude of the neuronal response is enhanced, while the representation by different cell groups declines: the neuronal responses are of higher magnitude and over a higher range, but there is more overlap between neurons responding to different sound pressure levels. This can be explained with the larger and broader responses of monotonic and nonmonotonic sound-increasing neuronal responses with VIP neuronal activation.

DISCUSSION

Within the brain, neurons form intricate networks, which represent sensory information. A sensory stimulus, such as a specific sound or a visual image, elicits activity in a subset of neurons in a network. A neuronal network can use a multitude of codes to represent information. A stimulus can be encoded discretely with a localist, pattern-separated representation, in which a specific group of neurons represents a specific stimulus, and different stimuli elicit activity in different groups of neurons (Figure 7A). Such localist representations have the advantage of discreteness: they can separate stimuli in different categories. Alternatively, in a distributed representation, stimulus-evoked activity can be distributed across the network, such that the relative activity of neurons within a group represent different stimuli (Figure 7A-B). An example of a distributed representation is a rate code, in which the firing rate of the active neurons represent a continuously varying stimulus feature, such as intensity or sound location (Belliveau et al., 2014). Distributed representations have the advantage of invariance: a small change in stimulus parameter will elicit a small variation in the neuronal response. Neuronal population responses have been measured at various positions along the localist to distributed representation spectrum across many features and areas (such as memory: (Wixted et al., 2014), sound (Hromádka et al., 2008), sound localization (Lesica et al., 2010; Belliveau et al., 2014), vision (Christensen and Pillow, 2022)) and can change dynamically along the spectrum (Kato et al., 2015; Honey et al., 2017; Kuchibhotla et al., 2017). A defining feature of the auditory cortex is sparse coding (DeWeese et al., 2003), which can lead to both distributed and localist representations. Along the auditory pathway, the coexistence of neurons with monotonic and nonmonotonic response-level curves indicates that sound pressure level is represented by both localist and distributed codes (Schreiner et al., 1992; Wu et al., 2006; Tan et al., 2007; Watkins and Barbour, 2011). More generally, stimulus representation within neuronal networks is mixed between local and distributed codes, in so-called sparse distributed representations, with both the level of activity and identity of activated neurons encoding the stimulus (Figure 7A) (Rolls and Tovee, 1995; Vinje and Gallant, 2000; Hromádka et al., 2008; Wixted et al., 2014). Based on the environmental and behavioral demands, it may be beneficial for neuronal representations to shift dynamically towards a more localist or a more distributed representation of a stimulus feature.

Our results suggest that distinct inhibitory neurons in the auditory cortex affect population neuronal response code by differentially shifting the responses toward a distributed or a localist representations. Previous work found that SST neuronal activation decreases the activity of the neuronal population to sounds (Phillips and Hasenstaub, 2016; Natan et al., 2017), and leads to a rightward shift of monotonic response-level curves (Wilson et al., 2012), while VIP neuronal activation, through a disinhibitory circuit, increases the activity of the population (Pfeffer et al., 2013; Pi et al., 2013; Zhang et al., 2014). We found that activation of SST neurons, led to a sparser, more localist representation (Figure 2D-E), where sound pressure level is encoded in discrete steps by distinct groups of neurons (Figure 3E) with a similar low strength (Figure 3K). By contrast, activation of VIP neurons, led to a more distributed representation (Figure 2K-L), with more overlap between the cell populations responding to different sound pressure levels (Figure 3H): sound pressure level is encoded continuously by varying the strength of response of a large group of neurons (Figure 3N). These shifts in representation are implemented at the single-neuron level through changes to the response-level function of monotonic (Figure 4) and nonmonotonic neurons (Figure 5). SST neuronal activation shifts the response-level curves of sound-modulated neurons by further separating the sound pressure levels that monotonic and nonmonotonic neurons represent (Figure 4H and 5H). With VIP neuronal activation, the changes to the response-level curves of sound-modulated neurons allow for stronger responses and more overlap in bandwidth between monotonic and nonmonotonic neurons (Figure 4H and 5H).

As distinct inhibitory populations can be recruited during different behaviors, their ability to transform the neuronal code can be advantageous to distinct behaviors and computations. Localist versus distributed representations, modulated by the relative strength of global (SST) versus local (PV or VIP) inhibition respectively, may provide support for neuronal computations such as discreteness versus invariance (Kuchibhotla and Bathellier, 2018), segmentation versus concatenation (Haga and Fukai, 2021), integration of bottom-up versus top-down information (Honey et al., 2017; Hertäg and Sprekeler, 2019). The two types of interneurons receive neuromodulatory inputs and may dynamically change the network's state towards one or the other type of representation for a given task: SST neuronal activation may help with tasks requiring focused attention such as discriminating different stimuli (Lee and Middlebrooks, 2011) or detecting in noise (Lakunina et al., 2022), by sharpening tuning, decreasing the activity for non-relevant stimuli, and enhancing the information-per-spike (Phillips and Hasenstaub, 2016). In contrast, VIP neuronal activation may help with tasks requiring receptive-attention such as active sensing (Gentet et al., 2012) or detecting small stimuli by amplifying weak signals (Millman et al., 2020), increasing detectability (Cone et al., 2019) without increasing the stimulus-response mutual information (Bigelow et al., 2019).

FIGURE 7: Predictions for testing for differential control of localist and distributed representations of sounds by inhibitory
neurons. A. A schematic of localist versus distributed neuronal codes. B. Localist versus distributed representations can be implemented in
many ways. An example of localist code is the pattern code without redundancy (known as the "Grandmother cell", (Bowers, 2009)). An
example of distributed code is a code relying only on the relative response of each cell and not the magnitude of response of the population
vector. C. Neuronal circuit manipulation: Optogenetic stimulation of SST and VIP neurons in Auditory Cortex (simplified connectivity
circuit). D. Noise bursts at different sound pressure levels (SPL) were presented to an awake mouse. E. For each sound pressure level, some
neurons responded (filled circles) while many cells didn't respond (empty circles). F. The response to different sound pressure levels can be
described in the neuronal space by the separation angle between the mean population vectors and the length separating them. G. Changes to
the representation of sound pressure level at the population level are implemented through changes to the response-level curve of each
neuron. Gray: baseline; blue: response transformation with SST activation; green: response transformation with VIP activation.

Our results expand on the results of previous studies that measured more general effects of
interneuron modulation. The changes to the response-level curves (Figures 4 and 5) we measured upon SST
or VIP neuronal activation can explain how the frequency-response functions of neurons in AC change with
interneuron modulation. The excitatory and inhibitory inputs to pyramidal cells of AC are frequency-tuned
(Isaacson and Scanziani, 2011; Li et al., 2013; Kato et al., 2017), and intracortical inhibition further shapes
the tuning (Wu et al., 2008). From the response-level curves, we can plot the response with interneuron
activation versus without and thus predict within which range of amplitudes we can expect multiplicative
or additive effects to the frequency tuning curve. Previous studies have shown that SST neuronal activation
leads to either subtractive, divisive or a combination of both subtractive and divisive effects on the
frequency tuning curve (Wilson et al., 2012; Seybold et al., 2015; Phillips and Hasenstaub, 2016). Our data
are consistent with these results: for monotonic neurons, SST neuronal activation can lead to divisive and/or
subtractive effects at a low range of sound amplitudes and for nonmonotonic neurons, SST neuronal
activation leads to both divisive and subtractive effects (Figure 6J-K). Previous studies have also shown
that VIP leads to an additive shift in the frequency tuning curve (Pi et al., 2013; Bigelow et al., 2019) and
similarly SST inactivation leads to a multiplicative or additive shift mostly (Phillips and Hasenstaub, 2016).
Our data can explain these results as well, with multiplicative effects for monotonic neurons and a range of
additive and multiplicative effects for nonmonotonic neurons (Figure 6J-K). One component that can
contribute to a change in representation is a change in the noisiness of the responses. Indeed, changes in the
signal-to-noise ratio (SNR) could drive populations to appear as more localist or distributed in their coding.
The change in SNR may be one of the components explaining how the change in representation is
implemented, however it is not the sole factor as assessed by the decoding accuracy (Figure 2F and M).

Nonmonotonic neurons in AC either can have their nonmonotonicity inherited from the
nonmonotonic excitatory input into those cells while the monotonic inhibitory input, which shows a peak
in delay at the cell's best pressure level, sharpens the nonmonotonicity (Wu et al., 2006) or can be
constructed *de novo* with an imbalance between excitatory and inhibitory inputs (Wu et al., 2006; Tan et
al., 2007). This means that for some of the nonmonotonic cells we recorded from, the input into the cell
does not covary with the sound pressure level, and so for these cells, we are not assessing the input-output
function of the cell through the response-level curve, rather how inhibition further shapes the already
intensity-tuned input. From our experiments, we observe that SST neuronal activation decreases the range
of responses of nonmonotonic neurons but does not change the sound pressure level of peak response nor
the width of responses (Figure 5). This may indicate that SST neuronal activation does not change the
timing of the inhibitory input into the nonmonotonic cells, but rather the overall strength of inhibition across
all sound pressure levels. In contrast, VIP neuronal activation leads to a shift of the sound pressure level of
peak response towards higher levels, along with a broadening of the response and an increase in the range
of response (Figure 5). This could simply be explained if VIP neuronal activation changes the timing of the
inhibition, with the delay between excitatory and inhibitory inputs peaking at a higher sound pressure level.

A limitation of our study is that we measured the response only from a subset of neurons from layer 2/3 while broadly stimulating many SST or VIP neurons across different cortical layers. However, neurons across the cortical column may perform additional computations in other layers, which would be important to record in future experiments. Furthermore, whereas we combined imaging sessions across the auditory cortex, future studies should also examine differences in function of inhibitory neurons across the different primary and non-primary auditory areas.

A related caveat in interpreting our results is that optogenetic activation of inhibitory neurons may differ across samples, and can potentially drive higher activity level of excitatory or inhibitory neurons than physiological levels, saturating the responses. To mitigate this limitation, we included multiple activation levels in each of the imaging sessions by modulating the strength of the laser between high and low levels. This allowed within sample comparison of activity patterns, rather than comparisons of absolute changes across multiple imaging sessions. Furthermore, by comparing the activity levels of imaged neurons between low and high laser intensities, we ensured that the modulation of activity was not saturating due to the laser.

Another potential caveat in interpreting the data is the while we were able to identify a subset of neurons as SST or VIP positive, the analysis combined multiple types of neurons as non-SST or non-VIP neurons, and for some neurons the tdTomato expression might not have been strong enough for detection as SST or VIP positive. A targeted experimental approach which would allow either in vivo or post-hoc identification of imaged neuronal types would allow to further distinguish response parameters among different types of inhibitory and excitatory neurons (Kerlin et al., 2010; Khan et al., 2018). Furthermore, future studies inactivating these inhibitory neurons would further complement and extend our findings (Phillips and Hasenstaub, 2016).

In our experiments, we tested the results of SST or VIP neuronal activation on the neuronal representation of sound pressure level, and an important next step will be to investigate whether the changes in neuronal representation of sound pressure level correlate with behavioral effects. One approach would be to image the activity of SST neurons while a mouse is engaged in a task that may require SST neuronal activation, or similarly VIP neurons. SST neurons may be involved in tasks leading to a sharpening in the neuronal tuning properties, or to a filtering out of irrelevant stimuli through the overall decrease in firing rate, such as discrimination tasks and signal-in-noise tasks (Otazu et al., 2009; Lee and Middlebrooks, 2011; Kuchibhotla et al., 2017; Christensen et al., 2019; Lakunina et al., 2022). VIP neurons may be involved in tasks requiring amplification of weak signals, such as a detection at threshold task or active sensing (Fritz et al., 2003; Gentet et al., 2012; Bennett et al., 2013; Kato et al., 2015; Cone et al., 2019; Millman et al., 2020). A second approach would be to measure how SST or VIP neuronal activity changes the performance of a mouse engaged in a task. In a detection task, we may expect that detection thresholds increase with SST neuronal activation and decrease with VIP neuronal activation, as seen in the visual cortex (Cone et al., 2019). In a discrimination task or detection of sounds in background task, we may expect SST neuronal activation to increase the performance, while VIP neuronal activation may decrease or not affect performance: SST neuronal inactivation decreases performance in the detection of sounds in background noise (Lakunina et al., 2022), and at the neuronal level, VIP neuronal activation decreases "encoding efficiency" (Bigelow et al., 2019) while SST neuronal activation may increase it (Phillips and Hasenstaub, 2016).

It should be noted that the dichotomy between the functional roles of SST and VIP neurons might not be so clear cut: VIP and SST neurons may cooperate to simultaneously amplify relevant stimuli and filter out irrelevant stimuli, respectively, or VIP neurons may be more active for weak stimuli and SST neurons for loud stimuli (Zhang et al., 2014; Mesik et al., 2015; Karnani et al., 2016; Kuchibhotla et al.,

2017; Dipoppa et al., 2018a; Millman et al., 2020). For example, activation of the cingulate cortex can elicit
spiking activity in PV, SST and VIP neurons, with SSTs contributing to surround suppression, and VIPs to
facilitation of the center of the receptive field (Zhang et al., 2014); cholinergive modulation depolarizes
multiple types of inhibitory neurons, including SSTs and VIPs (Kuchibhotla et al., 2017); and locomotion
increases activity in VIPs primarily for small stimuli, but in SSTs for large stimuli (Dipoppa et al., 2018b).
VIPs inhibit SSTs, creating attentional spotlights through targeted disinhibition (Karnani et al., 2016).
Therefore, examination of the function of SST and VIP neurons in complex behaviors needs to take the
circuit mechanisms into account. Interestingly, the connectivity between cortical neurons is largely
conserved across layers, primary and non-primary cortices, sensory and non-sensory areas (Douglas and
Martin, 2004; Markram et al., 2004; Yu et al., 2019; Campagnola et al., 2022), with SST and VIP neurons
mutually inhibiting each other as a common motif: perhaps the change in representation we observe with
varying sound level pressure upon SST or VIP neurons extends to other stimulus features, sensory and non-
sensory alike.

REFERENCES

Belliveau LAC, Lyamzin DR, Lesica NA (2014) The Neural Representation of Interaural Time Differences in Gerbils Is Transformed from Midbrain to Cortex. Journal of Neuroscience 34:16796–16808.

Bennett C, Arroyo S, Hestrin S (2013) Subthreshold Mechanisms Underlying State-Dependent Modulation of Visual Responses. Neuron 80:350–357.

Bigelow J, Morrill RJ, Dekloe J, Hasenstaub AR (2019) Movement and VIP Interneuron Activation Differentially Modulate Encoding in Mouse Auditory Cortex. eNeuro 6:ENEURO.0164-19.2019.

Blackwell JM, Lesicko AM, Rao W, De Biasi M, Geffen MN (2020) Auditory cortex shapes sound responses in the inferior colliculus Shinn-Cunningham BG, Carr CE, Woolley S, Read H, eds. eLife 9:e51890.

Bowers JS (2009) On the biological plausibility of grandmother cells: Implications for neural network theories in psychology and neuroscience. Psychological Review 116:220–251.

Campagnola L et al. (2022) Local connectivity and synaptic dynamics in mouse and human neocortex. Science 375 Available at: https://www.science.org/doi/10.1126/science.abj5861.

Chen N, Sugihara H, Sur M (2015) An acetylcholine-activated microcircuit drives temporal dynamics of cortical activity. Nat Neurosci 18:892–902.

Christensen AJ, Pillow JW (2022) Reduced neural activity but improved coding in rodent higher-order visual cortex during locomotion. Nat Commun 13:1676.

Christensen RK, Lindén H, Nakamura M, Barkat TR (2019) White Noise Background Improves Tone Discrimination by Suppressing Cortical Tuning Curves. Cell Reports.

Cone JJ, Scantlen MD, Histed MH, Maunsell JHR (2019) Different inhibitory interneuron cell classes make distinct contributions to visual contrast perception. eNeuro.

DeWeese MR, Wehr M, Zador AM (2003) Binary spiking in auditory cortex. J Neurosci 23:7940–7949.

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD (2018a) Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron.

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD (2018b) Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron.

Douglas RJ, Martin KAC (2004) Neuronal Circuits of the Neocortex. Annual Review of Neuroscience 27:419–451.

Fanselow EE, Richardson KA, Connors BW (2008) Selective, state-dependent activation of somatostatin-expressing inhibitory interneurons in mouse neocortex. J Neurophysiol 100:2640–2652.

Fritz J, Shamma S, Elhilali M, Klein D (2003) Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. Nat Neurosci 6:1216–1223.

Fu Y, Tucciarone JM, Espinosa JS, Sheng N, Darcy DP, Nicoll RA, Huang ZJ, Stryker MP (2014) A cortical circuit for gain control by behavioral state. Cell 156:1139–1152.

Gentet LJ, Kremer Y, Taniguchi H, Huang ZJ, Staiger JF, Petersen CCH (2012) Unique functional properties of somatostatin-expressing GABAergic neurons in mouse barrel cortex. Nat Neurosci 15:607–612.

Haga T, Fukai T (2021) Multiscale representations of community structures in attractor neural networks. PLOS Computational Biology 17:e1009296.

Hertäg L, Sprekeler H (2019) Amplifying the redistribution of somato-dendritic inhibition by the interplay of three interneuron types Cuntz H, ed. PLoS Comput Biol 15:e1006999.

Honey CJ, Newman EL, Schapiro AC (2017) Switching between internal and external modes: A multiscale learning principle. Network Neuroscience 1:339–356.

Hromádka T, DeWeese MR, Zador AM (2008) Sparse Representation of Sounds in the Unanesthetized Auditory Cortex. PLOS Biology 6:e16.

Isaacson JS, Scanziani M (2011) How inhibition shapes cortical activity. Neuron 72:231–243.

Jackson J, Ayzenshtat I, Karnani MM, Yuste R (2016) VIP+ interneurons control neocortical activity across brain states. Journal of Neurophysiology 115:3008–3017.

Karnani MM, Jackson J, Ayzenshtat I, Sichani XH, Manoocheri K, Kim S, Yuste R (2016) Opening holes in the blanket of inhibition: Localized lateral disinhibition by vip interneurons. Journal of Neuroscience 36:3471–3480.

Kato HK, Asinof SK, Isaacson JS (2017) Network-Level Control of Frequency Tuning in Auditory Cortex. Neuron.

Kato HK, Gillet SN, Isaacson JS (2015) Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. Neuron.

Kawaguchi Y, Shindou T (1998) Noradrenergic excitation and inhibition of GABAergic cell types in rat frontal cortex. J Neurosci 18:6963–6976.

Kerlin AM, Andermann ML, Berezovskii VK, Reid RC (2010) Broadly tuned response properties of diverse inhibitory neuron subtypes in mouse visual cortex. Neuron 67:858–871.

Khan AG, Poort J, Chadwick A, Blot A, Sahani M, Mrsic-Flogel TD, Hofer SB (2018) Distinct learning-induced changes in stimulus selectivity and interactions of GABAergic interneuron classes in visual cortex. Nature neuroscience 21:851–859.

Klapoetke NC et al. (2014) Independent optical excitation of distinct neural populations. Nat Methods 11:338–346.

Kuchibhotla K, Bathellier B (2018) Neural encoding of sensory and behavioral complexity in the auditory cortex. Current Opinion in Neurobiology 52:65–71.

Kuchibhotla KV, Gill JV, Lindsay GW, Papadoyannis ES, Field RE, Sten TAH, Miller KD, Froemke RC (2017) Parallel processing by cortical inhibition enables context-dependent behavior. Nat Neurosci 20:62–71.

Lakunina AA, Menashe N, Jaramillo S (2022) Contributions of Distinct Auditory Cortical Inhibitory Neuron Types to the Detection of Sounds in Background Noise. eneuro 9:ENEURO.0264-21.2021.

Lakunina AA, Nardoci MB, Ahmadian Y, Jaramillo S (2020) Somatostatin-expressing interneurons in the auditory cortex mediate sustained suppression by spectral surround. Journal of Neuroscience.

Lee C-C, Middlebrooks JC (2011) Auditory cortex spatial sensitivity sharpens during task performance. Nat
Neurosci 14:108–114.

Lesica NA, Lingner A, Grothe B (2010) Population Coding of Interaural Time Differences in Gerbils and Barn Owls.
Journal of Neuroscience 30:11696–11702.

Li L, Li Y, Zhou M, Tao HW, Zhang LI (2013) Intracortical multiplication of thalamocortical signals in mouse auditory
cortex. Nat Neurosci 16:1179–1181.

Litovsky RY, Clifton RK (1992) Use of sound-pressure level in auditory distance discrimination by 6-month-old
infants and adults. J Acoust Soc Am 92:794–802.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical
inhibitory system. Nat Rev Neurosci 5:793–807.

Mesik L, Ma WP, Li LY, Ibrahim LA, Huang ZJ, Zhang L, Tao HW (2015) Functional response properties of VIP-
expressing inhibitory neurons in mouse visual and auditory cortex. Frontiers in Neural Circuits 9.

Millman DJ, Ocker GK, Caldejon S, Kato I, Larkin JD, Lee EK, Luviano J, Nayan C, Nguyen TV, North K, Seid S, White
C, Lecoq J, Reid C, Buice MA, de Vries SEJ (2020) VIP interneurons in mouse primary visual cortex
selectively enhance responses to weak but specific stimuli. eLife 9:1–22.

Natan RG, Briguglio JJ, Mwilambwe-Tshilobo L, Jones SI, Aizenberg M, Goldberg EM, Geffen MN (2015)
Complementary control of sensory adaptation by two types of cortical interneurons King AJ, ed. eLife
4:e09868.

Natan RG, Rao W, Geffen MN (2017) Cortical Interneurons Differentially Shape Frequency Tuning following
Adaptation. Cell Reports 21:878–890.

Otazu GH, Tai L-H, Yang Y, Zador AM (2009) Engaging in an auditory task suppresses responses in auditory cortex.
Nat Neurosci 12:646–654.

Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of inhibition in visual cortex: The logic of
connections between molecularly distinct interneurons. Nature Neuroscience.

Phillips DP, Semple MN, Kitzes LM (1995) Factors shaping the tone level sensitivity of single neurons in posterior
field of cat auditory cortex. Journal of Neurophysiology 73:674–686.

Phillips EA, Hasenstaub AR (2016) Asymmetric effects of activating and inactivating cortical interneurons Uchida
N, ed. eLife 5:e18383.

Pi HJ, Hangya B, Kvitsiani D, Sanders JI, Huang ZJ, Kepecs A (2013) Cortical interneurons that specialize in
disinhibitory control. Nature 503:521–524.

Polley DB, Heiser MA, Blake DT, Schreiner CE, Merzenich MM (2004) Associative learning shapes the neural code
for stimulus magnitude in primary auditory cortex. Proceedings of the National Academy of Sciences
101:16351–16356.

Rolls ET, Tovee MJ (1995) Sparseness of the neuronal representation of stimuli in the primate temporal visual
cortex. Journal of Neurophysiology 73:713–726.

Rudy B, Fishell G, Lee S, Hjerling-Leffler J (2011) Three groups of interneurons account for nearly 100% of
neocortical GABAergic neurons. Developmental Neurobiology 71:45–61.

Schreiner CE, Mendelson JR, Sutter ML (1992) Experimental Brain Research Functional topography of cat primary
auditory cortex: representation of tone intensity.

Seay MJ, Natan RG, Geffen MN, Buonomano DV (2020) Differential short-term plasticity of PV and SST neurons
accounts for adaptation and facilitation of cortical neurons to auditory tones. Journal of Neuroscience
40:9224–9235.

Seybold BA, Phillips EAK, Schreiner CE, Hasenstaub AR (2015) Inhibitory Actions Unified by Network Integration.
Neuron 87:1181–1192.

Sun W, Marongelli EN, Watkins PV, Barbour DL (2017) Decoding sound level in the marmoset primary auditory
cortex. J Neurophysiol 118:2024–2033.

Tan AYY, Atencio CA, Polley DB, Merzenich MM, Schreiner CE (2007) Unbalanced synaptic inhibition can create
intensity-tuned auditory cortex neurons. Neuroscience 146:449–462.

Vinje WE, Gallant JL (2000) Sparse Coding and Decorrelation in Primary Visual Cortex During Natural Vision.
Science 287:1273–1276.

Watkins PV, Barbour DL (2011) Rate-level responses in awake marmoset auditory cortex. Hearing Research
275:30–42.

Willmore B, Tolhurst DJ (2001) Characterizing the sparseness of neural codes. Network: Computation in Neural
Systems 12:255–270.

Wilson NR, Runyan CA, Wang FL, Sur M (2012) Division and subtraction by distinct cortical inhibitory networks in
vivo. Nature 488:343–348.

Wixted JT, Squire LR, Jang Y, Papesh MH, Goldinger SD, Kuhn JR, Smith KA, Treiman DM, Steinmetz PN (2014)
Sparse and distributed coding of episodic memory in neurons of the human hippocampus. Proceedings of
the National Academy of Sciences 111:9621–9626.

Wood KC, Angeloni CF, Oxman K, Clopath C, Geffen MN (2022) Neuronal activity in sensory cortex predicts the
specificity of learning in mice. Nat Commun 13:1167.

Wu GK, Arbuckle R, Liu B, Tao HW, Zhang LI (2008) Lateral Sharpening of Cortical Frequency Tuning by
Approximately Balanced Inhibition. Neuron 58:132–143.

Wu GK, Li P, Tao HW, Zhang LI (2006) Nonmonotonic Synaptic Excitation and Imbalanced Inhibition Underlying
Cortical Intensity Tuning. Neuron 52:705–715.

Yu J, Hu H, Agmon A, Svoboda K (2019) Recruitment of GABAergic Interneurons in the Barrel Cortex during Active
Tactile Behavior. Neuron.

Zhang JW, Lau C, Cheng JS, Xing KK, Zhou IY, Cheung MM, Wu EX (2013) Functional magnetic resonance imaging of
sound pressure level encoding in the rat central auditory system. NeuroImage 65:119–126.

Zhang S, Xu M, Kamigaki T, Do JPH, Chang WC, Jenvay S, Miyamichi K, Luo L, Dan Y (2014) Long-range and local
circuits for top-down modulation of visual cortex processing. Science.

1062

1063

1064 **Table 1. Mouse strains and numbers**

Experiment	Figures	Strain	Number of mice	Number of recordings
GCaMP7f + ChrimsonR	Figs 1-5	CDH23 x VIP-Cre	2	7
		CDH23 x SST-Cre	5	13
GCaMP6m + ChrimsonR	Figs 1-5	CDH23 x VIP-Cre	2	9
		CDH23 x SST-Cre	0	0
Control: GCaMP7f + Flex.tdTomato – VIP cells	Fig 1H	CDH23 x VIP-Cre	4	7
Control: GCaMP7f + Flex.tdTomato – Non-VIP cells	Fig 1I	CDH23 x VIP-Cre	2	4

1065

Table 2. Statistics Table. We used a Generalized Linear Mixed-Effects (GLME) model and Wilcoxon
signed-rank tests to compute the statistics for the data.
For Figure 1, Figure 2B,C,E,G,I,J,L,N; the data ('`table`') had four columns: cell, sound level, laser
power, output. The formula used was (Matlab): `glme=fitglme(table,'output ~ sound + laser`
`+ sound*laser + (1|cell)');`
For Figure 3D,K, Figure 4 and Figure 5, the data ('`table`') had three columns: cell, laser power, output.
The formula used was (Matlab): `glme=fitglme(table,'output ~ laser + (1|cell)');`
For Figure 3E,H,K,N, the data ('`table`') had four columns: cell, sound level difference, laser power,
output. The formula used was (Matlab): `glme=fitglme(table,'output ~ sounddiff + laser +`
`sounddiff*laser + (1|cell)');`
For Figure 2F,M, we compared each sound amplitude across different light conditions using Wilcoxon
tests.
1078

Comparison	Figure	N	Test	Test Statistic	p-value	Effect size
FIGURE 1						
SST neuron with SST activation	Fig 1D	10 repeats	GLME	t_{laser}=7.89 t_{sound}=0.34 $t_{laser:sound}$=-0.55 DF = 206	***p_{laser}=1.8e-13** p_{sound}=0.74 $p_{laser:sound}$=0.58	η_{laser}^2=0.58 η_{sound}^2=3.8e-3 $\eta_{laser:sound}^2$=1.5e-2

Sound-increasing neuron with SST activation	Fig 1E	10 repeats	GLME	t_{laser}=0.33 t_{sound}=12.37 $t_{laser:sound}$=-8.34 DF = 206	p_{laser}=0.74 ***p_{sound}=1.2e-26 ***$p_{laser:sound}$=1.0e-14	η_{laser}^2=2.3e-3 η_{sound}^2=0.84 $\eta_{laser:sound}^2$=0.78
VIP neuron with VIP activation	Fig 1F	10 repeats	GLME	t_{laser}=5.40 t_{sound}=0.93 $t_{laser:sound}$=-2.56 DF = 206	***p_{laser}=1.8e-7 p_{sound}=0.35 *$p_{laser:sound}$=1.1e-2	η_{laser}^2=0.39 η_{sound}^2=2.8e-2 $\eta_{laser:sound}^2$=0.25
Sound-increasing neuron with VIP activation	Fig 1G	10 repeats	GLME	t_{laser}=2.45 t_{sound}=3.06 $t_{laser:sound}$=1.11 DF = 206	*p_{laser}=1.5e-2 **p_{sound}=2.5e-3 $p_{laser:sound}$=0.27	η_{laser}^2=0.12 η_{sound}^2=0.24 $\eta_{laser:sound}^2$=5.8e-2
Control: VIP neurons with laser activation	Fig 1H	54 cells	GLME	t_{laser}=1.27 t_{sound}=2.99 $t_{laser:sound}$=-0.11 DF = 11336	p_{laser}=0.20 **p_{sound}=2.8e-3 $p_{laser:sound}$=0.91	η_{laser}^2=6.5e-4 η_{sound}^2=5.5e-3 $\eta_{laser:sound}^2$=1.1e-5
Control: All neurons (VIP excluded) with laser activation	Fig 1I	492 cells	GLME	t_{laser}=0.46 t_{sound}=12.23 $t_{laser:sound}$=3.27 DF = 103316	p_{laser}=0.64 ***p_{sound}=2.1e-34 **$p_{laser:sound}$=1.1e-3	η_{laser}^2=8.5e-6 η_{sound}^2=9.0e-3 $\eta_{laser:sound}^2$=9.8e-4
FIGURE 2						
SST neurons with SST activation	Fig 2B	132 cells	GLME	t_{laser}=36.91 t_{sound}=1.32 $t_{laser:sound}$=0.16 DF = 27716	***p_{laser}=3.1e-291 p_{sound}=0.19 $p_{laser:sound}$=0.88	η_{laser}^2=0.14 η_{sound}^2=3.2e-4 $\eta_{laser:sound}^2$=6.7e-6
All non-SST neurons with SST activation	Fig 2C	2152 cells	GLME	t_{laser}=-1.27 t_{sound}=10.75	p_{laser}=0.20 ***p_{sound}=5.9e-27	η_{laser}^2=1.5e-5 η_{sound}^2=1.6e-3

				$t_{laser:sound}=-6.35$ DF = 451916	***$p_{laser:sound}=2.2e-10$	$\eta_{laser:sound}^2=8.5e-4$
Sparseness with SST activation	Fig 2D	2152 cells	GLME	$t_{laser}=3.39$ DF = 6454	***$p_{laser}=7.1e-4$	$\eta_{laser}^2=1.7e-3$
Activity sparseness with SST activation	Fig 2E	13 populations	GLME	$t_{laser}=3.20$ $t_{sound}=0.69$ $t_{laser:sound}=-0.99$ DF = 230	**$p_{laser}=1.6e-3$ $p_{sound}=0.49$ $p_{laser:sound}=0.33$	$\eta_{laser}^2=0.26$ $\eta_{sound}^2=9.8e-3$ $\eta_{laser:sound}^2=4.8e-2$
SST: Decoding accuracy of a linear SVM decoder with laser activation	Fig 2F	13 populations	GLME	$t_{laser}=-3.92$ $t_{sound}=-2.84$ $t_{laser:sound}=1.92$ DF = 269	***$p_{laser}=1.11e-4$ **$p_{sound}=0.0049$ $p_{laser:sound}=0.056$	$\eta_{laser}^2=0.19$ $\eta_{sound}^2=0.16$ $\eta_{laser:sound}^2=0.11$
Activity sparseness from 0dB and no laser power with SST activation	Fig 2F	13 populations	GLME	$t_{laser}=1.60$ $t_{sound}=1.17$ $t_{laser:sound}=-1.54$ DF = 230	$p_{laser}=0.11$ $p_{sound}=0.24$ $p_{laser:sound}=0.12$	$\eta_{laser}^2=5.5e-2$ $\eta_{sound}^2=1.9e-2$ $\eta_{laser:sound}^2=7.7e-2$
VIP neurons with VIP activation	Fig 2I	226 cells	GLME	$t_{laser}=41.50$ $t_{sound}=2.71$ $t_{laser:sound}=-1.96$ DF = 47456	***$p_{laser}=0$ **$p_{sound}=6.7e-3$ *$p_{laser:sound}=4.95e-2$	$\eta_{laser}^2=0.12$ $\eta_{sound}^2=9.3e-4$ $\eta_{laser:sound}^2=7.3e-4$
All non-VIP neurons with VIP activation	Fig 2J	3095 cells	GLME	$t_{laser}=34.18$ $t_{sound}=11.32$ $t_{laser:sound}=8.41$ DF = 649946	***$p_{laser}=7.8e-256$ ***$p_{sound}=1.1e-29$ ***$p_{laser:sound}=4.1e-17$	$\eta_{laser}^2=6.0e-3$ $\eta_{sound}^2=1.0e-3$ $\eta_{laser:sound}^2=8.4e-4$
Sparseness with VIP activation	Fig 2K	3095 cells	GLME	$t_{laser}=-22.95$	***$p_{laser}=2.2e-113$	$\eta_{laser}^2=4.8e-2$

				DF = 9283		
Activity sparseness with VIP activation	Fig 2L	16 populations	GLME	$t_{laser}=0.78$ $t_{sound}=-0.01$ $t_{laser:sound}=-0.49$ DF = 284	$p_{laser}=0.44$ $p_{sound}=0.99$ $p_{laser:sound}=0.62$	$\eta_{laser}^2=1.3e\text{-}2$ $\eta_{sound}^2=1.8e\text{-}6$ $\eta_{laser:sound}^2=8.2e\text{-}3$
VIP: Decoding accuracy of a linear SVM decoder with laser activation	Fig 2M	13 populations	GLME	$t_{laser}=-0.69$ $t_{sound}=-3.58$ $t_{laser:sound}=-0.11$ DF = 332	$p_{laser}=0.49$ ***$p_{sound}=\mathbf{3.99e\text{-}4}$ $p_{laser:sound}=0.91$	$\eta_{laser}^2=4.3e\text{-}3$ $\eta_{sound}^2=0.15$ $\eta_{laser:sound}^2=2.7e\text{-}4$
Activity sparseness from 0dB and no laser power with VIP activation	Fig 2N	16 populations	GLME	$t_{laser}=-3.52$ $t_{sound}=-0.49$ $t_{laser:sound}=0.0059$ DF = 284	***$p_{laser}=\mathbf{4.9e\text{-}4}$ $p_{sound}=0.62$ $p_{laser:sound}=0.995$	$\eta_{laser}^2=0.15$ $\eta_{sound}^2=2.1e\text{-}3$ $\eta_{laser:sound}^2=7.6e\text{-}7$
FIGURE 3						
Separation angle from 0dB at each laser power – SST activation	Fig 3E	15 angles, 13 recordings	GLME	$t_{laser}=8.80$ $t_{\Delta sound}=12.37$ $t_{laser:\Delta sound}=-7.44$ DF = 581	***$p_{laser}=\mathbf{1.6e\text{-}17}$ ***$p_{\Delta sound}=\mathbf{2.3e\text{-}31}$ ***$p_{laser:\Delta sound}=\mathbf{3.6e\text{-}13}$	$\eta_{laser}^2=0.30$ $\eta_{\Delta sound}^2=0.58$ $\eta_{laser:\Delta sound}^2=0.43$
Separation angle from 0dB at each laser power – VIP activation	Fig 3H	15 angles, 16 recordings	GLME	$t_{laser}=-2.75$ $t_{\Delta sound}=7.32$ $t_{laser:\Delta sound}=0.73$ DF = 716	**$p_{laser}=\mathbf{6.1e\text{-}3}$ ***$p_{\Delta sound}=\mathbf{6.9e\text{-}13}$ $p_{laser:\Delta sound}=0.47$	$\eta_{laser}^2=3.0e\text{-}2$ $\eta_{\Delta sound}^2=0.26$ $\eta_{laser:\Delta sound}^2=5.2e\text{-}3$
Vector length at each laser power – SST activation	Fig 3K	21 lengths, 13 recordings	GLME	$t_{laser}=-4.71$ $t_{\Delta sound}=5.71$ $t_{laser:\Delta sound}=-3.57$ DF = 815	***$p_{laser}=\mathbf{2.9e\text{-}6}$ ***$p_{\Delta sound}=\mathbf{1.6e\text{-}8}$ ***$p_{laser:\Delta sound}=\mathbf{3.8e\text{-}4}$	$\eta_{laser}^2=6.1e\text{-}2$ $\eta_{\Delta sound}^2=0.16$ $\eta_{laser:\Delta sound}^2=9.1e\text{-}2$

Vector length at each laser power – VIP activation	Fig 3N	21 lengths, 16 recordings	GLME	t_{laser}=2.50 $t_{\Delta sound}$=4.03 $t_{laser:\Delta sound}$=4.84 DF = 1004	*p_{laser}=**1.2e-2** ***$p_{\Delta sound}$=**6.1e-5** ***$p_{laser:\Delta sound}$=**1.5e-6**	η_{laser}^2=9.7e-3 $\eta_{\Delta sound}^2$=4.8e-2 $\eta_{laser:\Delta sound}^2$=9.0e-2
FIGURE 4						
Sigmoid fit – offset – with SST activation	Fig 4D	None: 109 Med: 103 High: 64	GLME	t_{laser}=0.83 DF = 274	p_{laser}=0.41	η_{laser}^2=2.4e-3
Sigmoid fit – offset – with VIP activation	Fig 4D	None: 267 Med: 239 High: 269	GLME	t_{laser}=1.74 DF = 773	p_{laser}=8.1e-2	η_{laser}^2=3.6e-3
Sigmoid fit – range – with SST activation	Fig 4E	None: 109 Med: 103 High: 64	GLME	t_{laser}=-1.24 DF = 274	p_{laser}=0.22	η_{laser}^2=3.1e-3
Sigmoid fit – range – with VIP activation	Fig 4E	None: 267 Med: 239 High: 269	GLME	t_{laser}=3.11 DF = 773	**p_{laser}=1.9e-3**	η_{laser}^2=9.4e-3
Sigmoid fit – midpoint – with SST activation	Fig 4F	None: 109 Med: 103 High: 64	GLME	t_{laser}=2.65 DF = 274	**p_{laser}=8.6e-3**	η_{laser}^2=2.1e-2
Sigmoid fit – midpoint – with VIP activation	Fig 4F	None: 267 Med: 239 High: 269	GLME	t_{laser}=-0.88 DF = 773	p_{laser}=0.38	η_{laser}^2=5.9e-4
Sigmoid fit – width – with SST activation	Fig 4G	None: 109 Med: 103 High: 64	GLME	t_{laser}=-0.019 DF = 274	p_{laser}=0.99	η_{laser}^2=8.8e-7
Sigmoid fit – width – with VIP activation	Fig 4G	None: 267 Med: 239 High: 269	GLME	t_{laser}=0.56 DF = 773	p_{laser}=0.57	η_{laser}^2=2.2e-4
FIGURE 5						
Gaussian fit – offset – with SST activation	Fig 5D	None: 224 Med: 175	GLME	t_{laser}=-3.16	**p_{laser}=1.7e-3**	η_{laser}^2=1.8e-2

		High: 130		DF = 527		
Gaussian fit – offset – with VIP activation	Fig 5D	None: 243 Med: 278 High: 310	GLME	t_{laser}=0.71 DF = 829	p_{laser}=0.48	η_{laser}^2=5.3e-4
Gaussian fit – range – with SST activation	Fig 5E	None: 224 Med: 175 High: 130	GLME	t_{laser}=-5.41 DF = 527	***p_{laser}=9.4e-8	η_{laser}^2=3.8e-2
Gaussian fit – range – with VIP activation	Fig 5E	None: 243 Med: 278 High: 310	GLME	t_{laser}=5.60 DF = 829	***p_{laser}=2.9e-8	η_{laser}^2=1.7e-2
Gaussian fit – mean – with SST activation	Fig 5F	None: 224 Med: 175 High: 130	GLME	t_{laser}=0.35 DF = 527	p_{laser}=0.73	η_{laser}^2=1.9e-4
Gaussian fit – mean – with VIP activation	Fig 5F	None: 243 Med: 278 High: 310	GLME	t_{laser}=3.34 DF = 829	***p_{laser}=8.6e-4	η_{laser}^2=7.4e-3
Gaussian fit – standard deviation – with SST activation	Fig 5G	None: 224 Med: 175 High: 130	GLME	t_{laser}=-0.63 DF = 527	p_{laser}=0.53	η_{laser}^2=7.3e-4
Gaussian fit – standard deviation – with VIP activation	Fig 5G	None: 243 Med: 278 High: 310	GLME	t_{laser}=3.96 DF = 829	***p_{laser}=8.1e-5	η_{laser}^2=1.7e-2

1079

1080