Insights into the mechanism of mycelium transformation of *Streptomyces*

***toxytricini* into pellet**

Punit Kumar[1], Khushboo[2], Deepanshi Rajput[3], Kashyap Kumar Dubey[3]*

[1]Department of Morphology and Physiology, Karaganda Medical University,

Karaganda, Kazakhstan 100008

[2]Department of Biotechnology, Central University of Haryana, Mahendergarh, India

123031

[3]Biomanufacturing and Process Development Laboratory, School of Biotechnology,

Jawaharlal Nehru University, New Delhi, India 110067

*Corresponding author

E-mail: kashyapdubey@gmail.com (KKD)

Telephone No.: +91-9996122280

Abstract

Formation of the mycelial pellet in submerged cultivation of *Streptomycetes* is unwanted in industrial fermentation processes as it imposes mass transfer limitations, changes in the rheology of a medium, and affects the production of secondary metabolites. Though detailed information is not available about the factors involved in regulating mycelial morphology but it is studied that culture conditions and genetic information of strain play a key role. Moreover, the proteomic study has revealed the involvement of low molecular weight proteins such as; DivIVA, FilP, ParA, Scy, and SsgA proteins in apical growth and branching of hyphae which results in the establishment of the mycelial network. The present study proposes the mechanism of pellet formation of *Streptomyces toxytricini* (NRRL B-5426) with the help of microscopic and proteomic analysis. The microscopic analysis revealed that growing hyphae followed a certain organized path of growth and branching, which was further converted into the pellet, and proteomic analysis revealed the production of low molecular weight proteins, which possibly participate in the regulation of pellet morphology.

Keywords: *Streptomyces*; pellet morphology; apical tip extension; mass transfer limitations; hyphae branching; lipstatin; microscopic analysis of pellet

1. Introduction

During submerged fermentation conditions, filamentous microorganisms exhibit morphologies between diffused mycelia and pellets as per the culture conditions and type of microbial strain. It is evident that mycelial morphologies are associated with the production of the desired natural product. Moreover, the pellet morphologies affect the rheology of a medium, heterogeneous mass transfer, and downstream processing (Rioseras et al., 2014; Wang et al., 2017). Therefore, regulation of pellet morphology is necessary for industrial processes.

Studies conducted at the molecular level suggested the involvement of many genes and proteins that control the mycelium transformation or morphological differentiation of Streptomyces (Vassallo et al., 2020; Wu et al., 2020). The growth of *Streptomyces* takes place from spore into vegetative hyphae which is under the regulation of AMP receptor protein Crp (Derouaux et al., 2004). Further growth of the hyphae takes place by tip extension and branching, where polarity protein DivIVA plays a key role (Flardh et al., 2012). It was suggested that during the growth of hyphae, the development of cross-walls also takes place, which protects hyphae from fission and forms a multinucleated compartment (Jakimowicz and van Wezel, 2012). Moreover, many protein complexes of *Streptomyces* like Scy, Tat secretion system, SsgA, and CslA are also found to be associated with apical tip growth of *Streptomyces* (Hempel et al., 2012; Holmes et al., 2013; Noens et al., 2007; Willemse et al., 2012; Xu, 2008). It was reported that SsgA is involved in the identification of a location for the development of septum and germination site (Noens et al., 2007). Interestingly, Scy, ParA, and FilP proteins are assumed to interact with DivIVA to control the apical growth of mycelia (Ditkowski et al., 2013; Holmes et al., 2013). One more protein HyaS has been found conserved in

streptomycetes and found to be associated with the regulation of pellet morphology by maintaining hyphal contacts (Koebsch et al., 2009).

Moreover, it was also suggested that some extracellular materials (proteins, sugars, and DNA) work as adhesives and are involved in the formation of pellet-like structures and provide protection. Extracellular DNA, hyaluronic acid, teichoic acids, and CslA have been found to be associated with the pellet morphology of *Streptomyces* (Kim and Kim, 2004; Ultee et al., 2020). It was found that cellulose synthase-like protein CslA was also identified near the hyphal tip which was involved in growth and structural transformation. It was also studied that it interacts with DivIVA and regulate the formation of biopolymer with glycosidic bonds near hyphal tips. The physical phenomenon, such as oxygen transfer and shear rate, also influence the morphology of *Streptomyces* (Ribeiro et al., 2021). Directive and quick-acting approaches, like the addition of microparticles and macroparticles, have also been used to regulate mycelial morphology (Böl et al., 2020; Yue et al., 2021).

Streptomyces genus is a group of Gram's positive bacteria which are one of the most explored microorganisms at the commercial scale for the synthesis of natural products. During submerged fermentation, these microorganisms represent pellet morphology (Hobbs et al., 1989). *Streptomyces toxytricini* is recognized for the production of lipase-inhibitory natural product lipstatin; which is used in an FDA-approved anti-obesity drug (Kumar and Dubey, 2015,16). Like other species of the genus *Streptomyces toxytricini* also exhibits pellet morphology in submerged fermentation (Kumar and Dubey, 2017).

In the previous report, authors have reported that culture conditions (pH of culture medium, medium composition, agitation rate, and inoculum size) influence the pellet size of *S. toxytricini* (Kumar and Dubey, 2017), but the mechanism of mycelial

transformation into pellet has not been elaborated. Though pellet size was reduced in

with change in culture conditions but biomass formation was also reduced. Thus in

order to reduce the pellet size or to maintain dispersed mycelial morphology without

affecting the biomass, mechanism of pellet formation must be studied. In the present

study authors made an attempt to microscopic analysis of hyphae growth and branching

which further transformed into the pellet. Further, partial analysis of proteins (produced

by bacteria in pellet and culture medium) was performed to understand the role of

proteins in pellet formation.

2. Materials and methods

2.1 Microorganism and growth conditions

S. toxytricini NRRL B-5426 strain was obtained from the Agricultural Research Service

(NRRL), Department of Agriculture, United States. The bacterium was grown in a

shake flask as per the conditions mentioned in (Kumar and Dubey, 2017). Incubation

temperature was maintained at $27.5^{o}C$ till the appearance of visible colonies. Well-

grown colonies of *S. toxytricini* were pink in colour, elevated, circular in shape, and

having a specific odour.

2.2 Analysis of pellet morphology

For pellet morphology analysis, an inoculum of *S. toxytricini* was prepared by

transferring a loopful of bacteria (*S. toxytricini*) from a Petri plate into a shake flask,

modified from (Kumar and Dubey, 2017).

After incubation, pellets were settled down and centrifuged at 5000 rpm for 10 min to

remove the culture medium. Aggregated pellets were washed thrice by 0.1M sodium

phosphate buffer of pH 7.2 ± 0.1 and stored for further processing. The Gram's staining

and methylene blues staining of pellets were performed to analyze morphology and

shape. Visualization was done in a compound microscope. For scanning electron

microscopy (SEM) of pellets, primary fixation of pellets was performed in 2.5%

glutaraldehyde and 2% paraformaldehyde in 0.1M sodium phosphate buffer at pH

7.2 ± 0.1. Further dehydration, fixation and coating, and SEM analysis were done at

Advanced Instrumentation Research Facility, Jawaharlal Nehru University, New Delhi

(India). For biomass analysis, pellets were kept for drying at 50 $^{\circ}$C in a hot air oven till

the appearance of constant weight.

2.3 Proteomic analysis of samples

Intracellular and extracellular proteins produced by *S. toxytricini* during growth were

extracted from the growth medium and pellets separately. For protein extraction from

culture broth, the acetone precipitation method was used and from protein isolation

from pellets, pellets were processed in lysis buffer at pH 7.0 [7,21]. Isolated proteins

were analyzed by 10% SDS PAGE.

3 Results and Discussion

3.1 Growth of *S. toxytricini*

The growth of *S. toxytricini* in submerged fermentation showed that if inoculum was

transferred from Petri plate to shake flask, it produced diffused mycelia and pellets of

smaller size while the transfer of inoculum from shake flask to shake flask, produced

visible pellets in the culture broth. Thus it can be assumed that in Petri plate bacterial

mycelium was not programmed for pellet formation, but in the culture broth, *S.*

toxytricini produces some chemical compound (intracellular or extracellular) that directs

the formation of pellets at the submerged level. Microscopic examination of *S.*

toxytricini, which was grown using inoculum from a Petri plate revealed that mycelial

aggregation took place, but the majority of these aggregations were clumps, loose

mycelia, and smaller pellets (upto 20μm). While shake flask to shake flask inoculation

produced pellets of larger size (30μm to 2mm) as major morphological form and very

less loose mycelia. Thus, it may be assumed that the transformation of mycelia into

pellets is influenced by culture conditions, and when *S. toxytricini* is grown in

submerged conditions, it produced some bio-chemicals or proteins which controlled

pellet formation. Including this heterogeneity of mycelial forms (loose, clump and

pellet) is maintained in broth medium which is according to previously reported study

(Kumar and Dubey, 2017). Similarly, some studies have reported that variation in

culture conditions directly affected pellet morphology of bacterial strain but such

mycelial morphology is unwanted for industrial processes (Flardh et al., 2012; Rioseras

et al., 2014; van Veluw et al., 2012).

3.2 Morphology of pellets

For better insight into pellet structure, the morphology of pellets was analyzed by

scanning electron microscope (Fig 1). In this study, it was observed that pellet

formation takes place by interwoven hyphae (Fig 1A) where hyphae are joined with

each other and form a compact network (Kim and Kim, 2004; Koebsch et al., 2009).

The surface of pellet has loose mycelia which are sites of growth and these mycelia

undergo fragmentation during agitation to form new pellet (Fig 1B) (Koebsch et al.,

2009). Further magnification (2μm scale) revealed that hyphae are attached to each

other and form tight junctions (Fig 1C). Microscopic analysis revealed that growing

hyphae bears bud-like structures behind the apical tip and these buds like structures

form branches which were developed at regular intervals (Fig 1D) (Hempel et al.,

2012). Thus it may be assumed that branching behind apical tip and tight junctions

assist in compact pellet formation. As reported earlier such types of association protect

hyphae from fission and produce multinucleoid structure (Jakimowicz and van Wezel,

2012). The microscopic analysis strongly supported the observations of previous

researchers such as; generation of multiple polarity centers (Holmes et al., 2013),

growth and development of pellet by tip extension, branching and cross-wall formation

(Hempel et al., 2012; Flardh et al., 2012).

3.3 Proteomics analysis

Researchers have reported many proteins (DivIVA, Scy, FilP, SsgA, ParA, Tat, CslA,

Afsk, CRP, HyaS, TrpM, and Mat complex etc) involved with apical growth, formation

of septa, branching in growing hyphae (Ditkowski et al., 2013; Hempel et al., 2012;

Holmes et al., 2013; Noens et al., 2007; van Dissel et al., 2018; Vassallo et al., 2020;

Willemse et al., 2012). To further elucidate the involvement of different proteins behind

pellet formation of *S. toxytricini*, secreted proteins in culture medium and intracellular

proteins were analyzed by electrophoresis. In this study, the partial analysis of proteins

(intracellular and extracellular) was conducted and It was observed that large numbers

of proteins of different molecular weight were produced at intracellular level and

extracellular level ranging from 95Kda to less than 20kDa (Fig 2). It is assumed that

many of these proteins are involved in pellet formation. It was observed in Fig 1 D, that

growing hyphae forms branch behind apical tip and studies have reported the expression

of DivIVA (21.5kDA) at growing tip which is suggested to interact with Scy, ParA and

FilP to regulate pellet formation (Ditkowski et al., 2013; Flardh et al., 2013; Holmes et

al., 2013). It has been suggested that Scy regulate number of polarity centers by

associating with DivIVA for new tip construction during branching (Holmes et al.,

2013). Including this, Scy was found to sequester DivIVA and initiate the formation of

new polarity centers (Holmes et al., 2013). The protein Scy is also assumed to associate

with ParA to hyphal tips and control polymerization of ParA. The Scy-ParA association

is assumed to be involved in the transition of hyphal elongation into sporulation

(Ditkowski et al., 2013). Thus these proteins need to be further analyzed for their

involvement in pellet formation. Including this, the presence of other proteins involved

in the growth and branching of hyphae into pellet formation need to be analyzed before

making any final conclusion.

3.4 Mechanism of pellet formation

In the previous study authors have discussed the process of pellet formation (Kumar and

Dubey, 2017) (Fig 3). DivIVA is found associated with cell wall synthesis, genetic

competence, and chromosome segregation during sporulation (Labajova et al., 2021). It

had been also suggested that DivIVA with some other proteins (Scy, ParA and FilP)

regulates formation of polarisome and forms branches in hyphae. Though continuous

agitation of culture medium enables clumping of mycelia, but there may be some

external biopolymer assisting the association of hyphae with each other resulting

compact pellet. To analyze this process, microscopic observation was conducted to

understand the pattern of pellet formation. It was revealed that during growth, hyphae

perform growth and branching behind apical tip. The growing hyphae come in close

proximity to form a network of hyphae which forms a reaction centre like structure.

This structure further grows and associates with other hyphae and finally converts into a

compact structure (Fig 3). This hypothesis is also supported by the study that apical

growth regulates cell polarity which determines the morphology of pellet (Hempel et

al., 2012; Holmes et al., 2013). Including this, researchers have demonstrated the

formation of cross-walls during growth of streptomycetes which compact the structure

into pellet (Flardh et al., 2012). Thus it may be suggested that pellet formation involves

the process of hyphae growth, development of polarity, branching and formation of

cross-walls which results into transformation of hyphae into compact pellet.

4 Conclusions

Streptomycetes are filamentous microorganisms which are reported for the production

of many valuable natural products including antibiotics and enzymes. The hyphae of

these microbes represent a distinct morphological form i.e. pellet during submerged

cultivation and undesirable for industrial process. Though, studies have reported

production of antibiotics using pellet morphology. Investigators have reported the

involvement of culture conditions and biomolecules controlling the morphology of

hyphae. In this study, microscopic analysis of morphological forms of *S. toxytricini* at

different time revealed that during submerged growth, the growth of hyphae took place

by tip extension, branching behind apical tip. The growing hyphae possibly attached

with each other and form cross-walls which further transformed into compact pellet

(Flardh et al., 2012; Hempel et al., 2012). The partial proteomic analysis in order to

understand the role of proteins in pellet formation revealed that there is presence of low

weigh proteins in pellet and culture medium which are possibly involved in pellet

formation but detailed study of such proteins like DivIVA, Scy, FilP, SsgA, ParA, Tat,

CslA, Afsk, CRP, HyaS, and Mat complex etc. is required.

Acknowledgements

Authors sincerely acknowledge the United States Department of Agriculture for

providing *S. toxytricini* NRRL B-5426 and Advanced Instrumentation Research

Facility, Jawaharlal Nehru University, New Delhi (India) for scanning electron

microscopy of pellets.

References

1. Böl, M., Schrinner, K., Tesche, S., & Krull, R. (2020). Challenges of influencing

cellular morphology by morphology engineering techniques and mechanical

induced stress on filamentous pellet systems-A critical review. *Engineering in*

life sciences, *21*(3-4), 51–67. https://doi.org/10.1002/elsc.202000060

2. Derouaux, A., Halici, S., Nothaft, H., Neutelings, T., Moutzourelis, G., Dusart,

J., Titgemeyer, F., Rigali, S., 2004. Deletion of a cyclic AMP receptor protein

homologue diminishes germination and affects morphological development of

Streptomyces coelicolor. J. Bacteriol. 186, 1893–1897. doi:

10.1128/JB.186.6.1893-1897.2004

3. Ditkowski, B., Holmes, N., Rydzak, J., Donczew, M., Bezulska, M., Ginda, K.,

et al. 2013. Dynamic interplay of ParA with the polarity protein, Scy,

coordinates the growth with chromosome segregation in *Streptomyces*

coelicolor. Open Biol. 3, 130006. doi: 10.1098/rsob.130006

4. Flardh, K., Richards, D.M., Hempel, A.M., Howard, M., Buttner, M.J., 2012.

Regulation of apical growth and hyphal branching in *Streptomyces*. Curr. Opin.

Microbiol. 15, 737–743. doi: 10.1016/j.mib.2012.10.012

5. Gomez-Escribano, J.P., Holmes, N.A., Schlimpert, S., Bibb, M.J., Chandra, G.,

Wilkinson, B., Buttner, M.J., Bibb, M.J., 2021. Streptomyces venezuelae NRRL

B-65442: genome sequence of a model strain used to study morphological

differentiation in filamentous actinobacteria. J. Ind. Microbiol. Biotechnol. doi:

10.1093/jimb/kuab035

6. Hempel, A.M., Cantlay, S., Molle, V., Wang, S.B., Naldrett, M.J., Parker, J.L.,

Richards, D.M., Jung, Y.G., Buttner, M.J., Flärdh, K., 2012. The Ser/Thr protein

kinase AfsK regulates polar growth and hyphal branching in the filamentous

bacteria *Streptomyces*. Proc. Natl. Acad. Sci. U.S.A. 109(35), E2371-E2379.

doi: 10.1073/pnas.1207409109

7. Hobbs, G., Frazer, C.M., Gardner, D.C.J., Cullum, J.A., Oliver, S.G., 1989.

Dispersed growth of *Streptomyces* in liquid culture. Appl. Microbiol.

Biotechnol. 31, 272-277. doi:10.1007/BF00258408

8. Holmes, N.A., Walshaw, J., Leggett, R.M., Thibessard, A., Dalton, K.A.,

Gillespie, M.D., Hemmings, A.M., Gust, B., Kelemen, G.H., 2013. Coiled-coil

protein Scy is a key component of a multiprotein assembly controlling polarized

growth in *Streptomyces*. Proc. Natl. Acad. Sci. U.S.A. 110(5), E397-406. doi:

10.1073/pnas.1210657110

9. Jakimowicz, D., van Wezel, G.P., 2012. Cell division and DNA segregation in

Streptomyces: How to build a septum in the middle of nowhere? Mol. Microbiol.

85, 393–404. doi: 10.1111/j.1365-2958.2012.08107.x

10. Kim, Y.M., Kim, J.H., 2004. Formation and dispersion of mycelial pellets of

Streptomyces coelicolor A3(2). J. Microbiol. 42(1), 64-7.

11. Koebsch, I., Overbeck, J., Piepmeyer, S., Meschke, H., Schrempf, H., 2009. A

molecular key for building hyphae aggregates: The role of the newly identified

Streptomyces protein HyaS. Microb. Biotechnol. 2(3), 343-360. doi:

10.1111/j.1751-7915.2009.00093.x

12. Kumar, P., Dubey, K.K., 2015. Current trends and future prospects of lipstatin: a

lipase inhibitor and pro-drug for obesity. RSC Adv. 5, 86954-66.

doi:10.1039/C5RA14892H

13. Kumar, P., Dubey, K.K., 2016. Modulation of fatty acid metabolism and

tricarboxylic acid cycle to enhance the lipstatin production through medium

engineering in *Streptomyces toxytricini*. Bioresour. Technol. 213, 64-8

doi:10.1016/j.biortech.2016.01.133

14. Kumar, P., Dubey, K.K., 2017. Mycelium transformation of Streptomyces

toxytricini into pellet: role of culture conditions and kinetics. Bioresour.

Technol. 228, 339-347. doi: 10.1016/j.biortech.2017.01.002

15. Labajová, N., Baranova, N., Jurásek, M., Vácha, R., Loose, M., Barák, I., 2021.

Cardiolipin-Containing Lipid Membranes Attract the Bacterial Cell Division

Protein DivIVA. Int. J. Mol. Sci. 22(15), 8350. doi: 10.3390/ijms22158350

16. Noens, E.E.E., Mersinias, V., Willemse, J., Traag, B.A., Laing, E., Chater, K.F.,

Smith, C.P., Koerten, H.K., van Wezel, G.P., 2007. Loss of the controlled

localization of growth stage-specific cell-wall synthesis pleiotropically affects

developmental gene expression in an ssgA mutant of *Streptomyces coelicolor*.

Mol. Microbiol. 64(5), 1244-1259

17. Ribeiro, R. M. M. G. P., Esperança, M. N., Sousa, A. P. A., Neto, Á. B., &

Cerri, M. O. (2021). Individual effect of shear rate and oxygen transfer on

clavulanic acid production by Streptomyces clavuligerus. *Bioprocess and*

biosystems engineering, *44*(8), 1721–1732. https://doi.org/10.1007/s00449-021-

02555-1

18. Rioseras, B., López-García, M.T., Yagüe, P., Sánchez, J., Manteca, A., 2014.

Mycelium differentiation and development of *Streptomyces coelicolor* in lab-

scale bioreactors: Programmed cell death, differentiation, and lysis are closely

linked to undecylprodigiosin and actinorhodin production. Bioresour. Technol.

151, 191-8. doi:10.1016/j.biortech.2013.10.068.

19. Ultee, E., van der Aart, L.T., Zhang, L., van Dissel, D., Diebolder, C.A., van

Wezel, G.P., Claessen, D., Briegel, A., 2020. Teichoic acids anchor distinct cell

wall lamellae in an apically growing bacterium. Commun. Biol. 3, 314. doi:

10.1038/s42003-020-1038-6

20. van Dissel, D., Willemse, J., Zacchetti, B., Claessen, D., Pier, G.B., van Wezel,

G.P., 2018. Production of poly-β-1, 6-N-acetylglucosamine by MatAB is

required for hyphal aggregation and hydrophilic surface adhesion by

Streptomyces. Microbial Cell 5(6), 269-279. doi: 10.15698/mic2018.06.635

21. van Veluw, G.J., Petrus, M.L., Gubbens, J., de Graaf, R., de Jong, I.P., van

Wezel, G.P., Wösten, H.A., Claessen, D., 2012. Analysis of two distinct

mycelial populations in liquid-grown *Streptomyces* cultures using a flow

cytometry-based proteomics approach. Appl. Microbiol. Biotechnol. 96(5),

1301–12. doi: 10.1007/s00253-012-4490-5

22. Vassallo, A., Palazzotto, E., Renzone, G., Botta, L., Faddetta, T., Scaloni, A.,

Puglia, A.M., Gallo, G., 2020. The Streptomyces coelicolor small ORF trpM

stimulates growth and morphological development and exerts opposite effects

on actinorhodin and calcium-dependent antibiotic production. Front. Microbiol.

11, 224. doi: 10.3389/fmicb.2020.00224

23. Wang, H., Zhao, G., Ding, X., 2017. Morphology engineering of Streptomyces

coelicolor M145 by sub-inhibitory concentrations of antibiotics. Sci. Rep. 7,

13226. doi:10.1038/s41598-017-13493-y

24. Willemse, J., Ruban-Osmialowska, B., Widdick, D., Celler, K., Hutchings, M.I.,

van Wezel, G.P., Palmer, T., 2012. Dynamic localization of Tat protein transport

machinery components in *Streptomyces coelicolor*. J. Bacteriol. 194(22), 6272-

6281. doi: 10.1128/JB.01425-12

25. Wu, Y., Kang, Q., Zhang, L. L., & Bai, L. (2020). Subtilisin-Involved

Morphology Engineering for Improved Antibiotic Production in

Actinomycetes. *Biomolecules*, *10*(6), 851.

https://doi.org/10.3390/biom10060851

26. Yue, C., Xu, H., Yu, Y., Yu, X., Yu, M., Zhang, C., You, Q., Xia, S., Ding, Z.,

Fu, H., Zeng, X. and Li, F., 2021. Improvement of natamycin production by

controlling the morphology of *Streptomyces gilvosporeus* Z8 with microparticle

talc in seed preculture. J Chem Technol Biotechnol, 96: 1533-

1542. https://doi.org/10.1002/jctb.6668

Figures

Fig. 1: Analysis of pellet morphology and mycelial structure by scanning electron

microscopy. **A**- Morphology of pellet at 10μM scale, **B-** Surface of pellet showing free

mycelia which show growth and branching at 10μM scale, **C**- Structure of hyphae

showing association with each other and branching at growing tips (2μM scale), **D**-

Apical growth and branching of hyphae (2μM scale).

Fig. 2: Analysis of extracellular and intracellular proteins of *S. toxytricini* by SDS-

PAGE. Lane 1 represents total intracellular proteins, lane 2,3 represents total

extracellular proteins isolated from culture medium. Marker lane is molecular weight

marker.

Fig. 3: Proposed mechanism for pellet formation. **A**-Diffused mycelia, **B**-Mycelia are coming together, **C**-mycelia are clumping, **D**-mycelial clumps are more visible and a reaction centre type structure is formed which possibly directs pellet formation, **E,F,G**-mycalial clumps are more condensed, **H**-Pellet formed and periphery having mycelia