Insights into the mechanism of mycelium transformation of *Streptomyces*

***toxytricini* into pellet**

Punit Kumar[1], Khusboo[2], Deepanshi Rajput[3], Kashyap Kumar Dubey[3]*

[1]Department of Morphology and Physiology, Karaganda Medical University,

Karaganda, Kazakhstan 100008

[2]Department of Biotechnology, Central University of Haryana, Mahendergarh, India

123031

[3]School of Biotechnology, Jawaharlal Nehru University, New Delhi, India 110067

*Corresponding author

E-mail: kashyapdubey@gmail.com (KKD)

Telephone No.: +91-9996122280

Abstract

Formation of the mycelial pellet in submerged cultivation of *Streptomycetes* is unwanted

in industrial fermentation processes as it imposes mass transfer limitations, changes in

the rheology of a medium, and affects the production of secondary metabolites. Though

detailed information is not available about the factors involved in regulating mycelial

morphology but it is studied that culture conditions and genetic information of strain play

a key role. Moreover, the proteomic study has revealed the involvement of low molecular

weight proteins such as; DivIVA, FilP, ParA, Scy, and SsgA proteins in apical growth

and branching of hyphae which results in the establishment of the mycelial network. The

present study proposes the mechanism of pellet formation of *Streptomyces toxytricini*

(NRRL B-5426) with the help of microscopic and proteomic analysis. The microscopic

analysis revealed that growing hyphae followed a certain organized path of growth and

branching, which was further converted into the pellet, and proteomic analysis revealed

the production of low molecular weight proteins, which possibly participate in the

regulation of pellet morphology.

Keywords: *Streptomyces*; pellet morphology; apical tip extension; mass transfer

limitations; hyphae branching; lipstatin; microscopic analysis of pellet

1. Introduction

During submerged fermentation conditions, filamentous microorganisms exhibit morphologies between diffused mycelia and pellets as per the culture conditions and type of microbial strain. It is evident that mycelial morphologies are associated with the production of the desired natural product. Moreover, the pellet morphologies affect the rheology of a medium, heterogeneous mass transfer, and downstream processing (Rioseras et al., 2014; Wang et al., 2017). Therefore, regulation of pellet morphology is necessary for industrial processes.

Studies conducted at the molecular level suggested the involvement of many genes and proteins that control the mycelium transformation or morphological differentiation of Streptomyces (Vassallo et al., 2020; Wu et al., 2020). The growth of *Streptomyces* takes place from spore into vegetative hyphae which is under the regulation of AMP receptor protein Crp (Derouaux et al., 2004). Further growth of the hyphae takes place by tip extension and branching, where polarity protein DivIVA plays a key role (Flardh et al., 2012). It was suggested that during the growth of hyphae, the development of cross-walls also takes place, which protects hyphae from fission and forms a multinucleated compartment (Jakimowicz and van Wezel, 2012). Moreover, many protein complexes of *Streptomyces* like Scy, Tat secretion system, SsgA, and CslA are also found to be associated with apical tip growth of *Streptomyces* (Hempel et al., 2012; Holmes et al., 2013; Noens et al., 2007; Willemse et al., 2012; Xu, 2008). It was reported that SsgA is involved in the identification of a location for the development of septum and germination site (Noens et al., 2007). Interestingly, Scy, ParA, and FilP proteins are assumed to interact with DivIVA to control the apical growth of mycelia (Ditkowski et al., 2013; Holmes et al., 2013). One more protein HyaS has been found conserved in streptomycetes

and found to be associated with the regulation of pellet morphology by maintaining

hyphal contacts (Koebsch et al., 2009).

Moreover, it was also suggested that some extracellular materials (proteins, sugars, and

DNA) work as adhesives and are involved in the formation of pellet-like structures and

provide protection. Extracellular DNA, hyaluronic acid, teichoic acids, and CslA have

been found to be associated with the pellet morphology of *Streptomyces* (Kim and Kim,

2004; Ultee et al., 2020). It was found that cellulose synthase-like protein CslA was also

identified near the hyphal tip which was involved in growth and structural transformation.

It was also studied that it interacts with DivIVA and regulate the formation of biopolymer

with glycosidic bonds near hyphal tips. The physical phenomenon, such as oxygen

transfer and shear rate, also influence the morphology of *Streptomyces* (Ribeiro et al.,

2021). Directive and quick-acting approaches, like the addition of microparticles and

macroparticles, have also been used to regulate mycelial morphology (Böl et al., 2020;

Yue et al., 2021).

Streptomyces genus is a group of Gram's positive bacteria which are one of the most

explored microorganisms at the commercial scale for the synthesis of natural products.

During submerged fermentation, these microorganisms represent pellet morphology

(Hobbs et al., 1989). *Streptomyces toxytricini* is recognized for the production of lipase-

inhibitory natural product lipstatin; which is used in an FDA-approved anti-obesity drug

(Kumar and Dubey, 2015,16). Like other species of the genus *Streptomyces toxytricini*

also exhibits pellet morphology in submerged fermentation (Kumar and Dubey, 2017).

In the previous report, authors have reported that culture conditions (pH of culture

medium, medium composition, agitation rate, and inoculum size) influence the pellet size

of *S. toxytricini* (Kumar and Dubey, 2017), but the mechanism of mycelial transformation

into pellet has not been elaborated. Though pellet size was reduced in with change in

culture conditions but biomass formation was also reduced. Thus in order to reduce the

pellet size or to maintain dispersed mycelial morphology without affecting the biomass,

mechanism of pellet formation must be studied. In the present study authors made an

attempt to microscopic analysis of hyphae growth and branching which further

transformed into the pellet. Further, partial analysis of proteins (produced by bacteria in

pellet and culture medium) was performed to understand the role of proteins in pellet

formation.

2. Materials and methods

2.1 Microorganism and growth conditions

S. toxytricini NRRL B-5426 strain was obtained from the Agricultural Research Service

(NRRL), Department of Agriculture, United States. The bacterium was grown in a shake

flask as per the conditions mentioned in (Kumar and Dubey, 2017). Incubation

temperature was maintained at 27.5°C till the appearance of visible colonies. Well-grown

colonies of *S. toxytricini* were pink in colour, elevated, circular in shape, and having a

specific odour.

2.2 Analysis of pellet morphology

For pellet morphology analysis, an inoculum of *S. toxytricini* was prepared by transferring

a loopful of bacteria (*S. toxytricini*) from a Petri plate into a shake flask, modified from

(Kumar and Dubey, 2017).

After incubation, pellets were settled down and centrifuged at 5000 rpm for 10 min to

remove the culture medium. Aggregated pellets were washed thrice by 0.1M sodium

phosphate buffer of pH 7.2±0.1 and stored for further processing. The Gram's staining

and methylene blues staining of pellets were performed to analyze morphology and shape.

Visualization was done in a compound microscope. For scanning electron microscopy

(SEM) of pellets, primary fixation of pellets was performed in 2.5% glutaraldehyde and

2% paraformaldehyde in 0.1M sodium phosphate buffer at pH 7.2±0.1. Further

dehydration, fixation and coating, and SEM analysis were done at Advanced

Instrumentation Research Facility, Jawaharlal Nehru University, New Delhi (India). For

biomass analysis, pellets were kept for drying at 50 ^{o}C in a hot air oven till the appearance

of constant weight.

2.3 Proteomic analysis of samples

Intracellular and extracellular proteins produced by *S. toxytricini* during growth were

extracted from the growth medium and pellets separately. For protein extraction from

culture broth, the acetone precipitation method was used and from protein isolation from

pellets, pellets were processed in lysis buffer at pH 7.0 [7,21]. Isolated proteins were

analyzed by 10% SDS PAGE.

3 Results and Discussion

3.1 Growth of *S. toxytricini*

The growth of *S. toxytricini* in submerged fermentation showed that if inoculum was

transferred from Petri plate to shake flask, it produced diffused mycelia and pellets of

smaller size while the transfer of inoculum from shake flask to shake flask, produced

visible pellets in the culture broth. Thus it can be assumed that in Petri plate bacterial

mycelium was not programmed for pellet formation, but in the culture broth, *S. toxytricini*

produces some chemical compound (intracellular or extracellular) that directs the

formation of pellets at the submerged level. Microscopic examination of *S. toxytricini,*

which was grown using inoculum from a Petri plate revealed that mycelial aggregation

took place, but the majority of these aggregations were clumps, loose mycelia, and

smaller pellets (upto 20μm). While shake flask to shake flask inoculation produced pellets

of larger size (30μm to 2mm) as major morphological form and very less loose mycelia.

Thus, it may be assumed that the transformation of mycelia into pellets is influenced by

culture conditions, and when *S. toxytricini* is grown in submerged conditions, it produced

some bio-chemicals or proteins which controlled pellet formation. Including this

heterogeneity of mycelial forms (loose, clump and pellet) is maintained in broth medium

which is according to previously reported study (Kumar and Dubey, 2017). Similarly,

some studies have reported that variation in culture conditions directly affected pellet

morphology of bacterial strain but such mycelial morphology is unwanted for industrial

processes (Flardh et al., 2012; Rioseras et al., 2014; van Veluw et al., 2012).

3.2 Morphology of pellets

For better insight into pellet structure, the morphology of pellets was analyzed by

scanning electron microscope (Fig 1). In this study, it was observed that pellet formation

takes place by interwoven hyphae (Fig 1A) where hyphae are joined with each other and

form a compact network (Kim and Kim, 2004; Koebsch et al., 2009). The surface of pellet

has loose mycelia which are sites of growth and these mycelia undergo fragmentation

during agitation to form new pellet (Fig 1B) (Koebsch et al., 2009). Further magnification

(2μm scale) revealed that hyphae are attached to each other and form tight junctions (Fig

1C). Microscopic analysis revealed that growing hyphae bears bud-like structures behind

the apical tip and these buds like structures form branches which were developed at

regular intervals (Fig 1D) (Hempel et al., 2012). Thus it may be assumed that branching

behind apical tip and tight junctions assist in compact pellet formation. As reported earlier

such types of association protect hyphae from fission and produce multinucleoid structure

(Jakimowicz and van Wezel, 2012). The microscopic analysis strongly supported the

observations of previous researchers such as; generation of multiple polarity centers

(Holmes et al., 2013), growth and development of pellet by tip extension, branching and

cross-wall formation (Hempel et al., 2012; Flardh et al., 2012).

3.3 Proteomics analysis

Researchers have reported many proteins (DivIVA, Scy, FilP, SsgA, ParA, Tat, CslA,

Afsk, CRP, HyaS, TrpM, and Mat complex etc) involved with apical growth, formation

of septa, branching in growing hyphae (Ditkowski et al., 2013; Hempel et al., 2012;

Holmes et al., 2013; Noens et al., 2007; van Dissel et al., 2018; Vassallo et al., 2020;

Willemse et al., 2012). To further elucidate the involvement of different proteins behind

pellet formation of *S. toxytricini*, secreted proteins in culture medium and intracellular

proteins were analyzed by electrophoresis. In this study, the partial analysis of proteins

(intracellular and extracellular) was conducted and It was observed that large numbers of

proteins of different molecular weight were produced at intracellular level and

extracellular level ranging from 95Kda to less than 20kDa (Fig 2). It is assumed that many

of these proteins are involved in pellet formation. It was observed in Fig 1 D, that growing

hyphae forms branch behind apical tip and studies have reported the expression of

DivIVA (21.5kDA) at growing tip which is suggested to interact with Scy, ParA and FilP

to regulate pellet formation (Ditkowski et al., 2013; Flardh et al., 2013; Holmes et al.,

2013). It has been suggested that Scy regulate number of polarity centers by associating

with DivIVA for new tip construction during branching (Holmes et al., 2013). Including

this, Scy was found to sequester DivIVA and initiate the formation of new polarity centers

(Holmes et al., 2013). The protein Scy is also assumed to associate with ParA to hyphal

tips and control polymerization of ParA. The Scy-ParA association is assumed to be

involved in the transition of hyphal elongation into sporulation (Ditkowski et al., 2013).

Thus these proteins need to be further analyzed for their involvement in pellet formation.

Including this, the presence of other proteins involved in the growth and branching of

hyphae into pellet formation need to be analyzed before making any final conclusion.

3.4 Mechanism of pellet formation

In the previous study authors have discussed the process of pellet formation (Kumar and

Dubey, 2017) (Fig 3). DivIVA is found associated with cell wall synthesis, genetic

competence, and chromosome segregation during sporulation (Labajova et al., 2021). It

had been also suggested that DivIVA with some other proteins (Scy, ParA and FilP)

regulates formation of polarisome and forms branches in hyphae. Though continuous

agitation of culture medium enables clumping of mycelia, but there may be some external

biopolymer assisting the association of hyphae with each other resulting compact pellet.

To analyze this process, microscopic observation was conducted to understand the pattern

of pellet formation. It was revealed that during growth, hyphae perform growth and

branching behind apical tip. The growing hyphae come in close proximity to form a

network of hyphae which forms a reaction centre like structure. This structure further

grows and associates with other hyphae and finally converts into a compact structure (Fig

3). This hypothesis is also supported by the study that apical growth regulates cell polarity

which determines the morphology of pellet (Hempel et al., 2012; Holmes et al., 2013).

Including this, researchers have demonstrated the formation of cross-walls during growth

of streptomycetes which compact the structure into pellet (Flardh et al., 2012). Thus it

may be suggested that pellet formation involves the process of hyphae growth,

development of polarity, branching and formation of cross-walls which results into

transformation of hyphae into compact pellet.

4 Conclusions

Streptomycetes are filamentous microorganisms which are reported for the production of

many valuable natural products including antibiotics and enzymes. The hyphae of these

microbes represent a distinct morphological form i.e. pellet during submerged cultivation

and undesirable for industrial process. Though, studies have reported production of

antibiotics using pellet morphology. Investigators have reported the involvement of

culture conditions and biomolecules controlling the morphology of hyphae. In this study,

microscopic analysis of morphological forms of *S. toxytricini* at different time revealed

that during submerged growth, the growth of hyphae took place by tip extension,

branching behind apical tip. The growing hyphae possibly attached with each other and

form cross-walls which further transformed into compact pellet (Flardh et al., 2012;

Hempel et al., 2012). The partial proteomic analysis in order to understand the role of

proteins in pellet formation revealed that there is presence of low weigh proteins in pellet

and culture medium which are possibly involved in pellet formation but detailed study of

such proteins like DivIVA, Scy, FilP, SsgA, ParA, Tat, CslA, Afsk, CRP, HyaS, and Mat

complex etc. is required.

Acknowledgements

Authors sincerely acknowledge the United States Department of Agriculture for

providing *S. toxytricini* NRRL B-5426 and Advanced Instrumentation Research Facility,

Jawaharlal Nehru University, New Delhi (India) for scanning electron microscopy of

pellets.

References

1. Böl, M., Schrinner, K., Tesche, S., & Krull, R. (2020). Challenges of influencing cellular morphology by morphology engineering techniques and mechanical induced stress on filamentous pellet systems-A critical review. *Engineering in life sciences*, *21*(3-4), 51–67. https://doi.org/10.1002/elsc.202000060

2. Derouaux, A., Halici, S., Nothaft, H., Neutelings, T., Moutzourelis, G., Dusart, J., Titgemeyer, F., Rigali, S., 2004. Deletion of a cyclic AMP receptor protein homologue diminishes germination and affects morphological development of *Streptomyces coelicolor*. J. Bacteriol. 186, 1893–1897. doi: 10.1128/JB.186.6.1893-1897.2004

3. Ditkowski, B., Holmes, N., Rydzak, J., Donczew, M., Bezulska, M., Ginda, K., et al. 2013. Dynamic interplay of ParA with the polarity protein, Scy, coordinates the growth with chromosome segregation in *Streptomyces coelicolor*. Open Biol. 3, 130006. doi: 10.1098/rsob.130006

4. Flardh, K., Richards, D.M., Hempel, A.M., Howard, M., Buttner, M.J., 2012. Regulation of apical growth and hyphal branching in *Streptomyces*. Curr. Opin. Microbiol. 15, 737–743. doi: 10.1016/j.mib.2012.10.012

5. Gomez-Escribano, J.P., Holmes, N.A., Schlimpert, S., Bibb, M.J., Chandra, G., Wilkinson, B., Buttner, M.J., Bibb, M.J., 2021. Streptomyces venezuelae NRRL B-65442: genome sequence of a model strain used to study morphological differentiation in filamentous actinobacteria. J. Ind. Microbiol. Biotechnol. doi: 10.1093/jimb/kuab035

6. Hempel, A.M., Cantlay, S., Molle, V., Wang, S.B., Naldrett, M.J., Parker, J.L., Richards, D.M., Jung, Y.G., Buttner, M.J., Flärdh, K., 2012. The Ser/Thr protein

kinase AfsK regulates polar growth and hyphal branching in the filamentous

bacteria *Streptomyces*. Proc. Natl. Acad. Sci. U.S.A. 109(35), E2371-E2379.

doi: 10.1073/pnas.1207409109

7. Hobbs, G., Frazer, C.M., Gardner, D.C.J., Cullum, J.A., Oliver, S.G., 1989.

Dispersed growth of *Streptomyces* in liquid culture. Appl. Microbiol.

Biotechnol. 31, 272-277. doi:10.1007/BF00258408

8. Holmes, N.A., Walshaw, J., Leggett, R.M., Thibessard, A., Dalton, K.A.,

Gillespie, M.D., Hemmings, A.M., Gust, B., Kelemen, G.H., 2013. Coiled-coil

protein Scy is a key component of a multiprotein assembly controlling polarized

growth in *Streptomyces*. Proc. Natl. Acad. Sci. U.S.A. 110(5), E397-406. doi:

10.1073/pnas.1210657110

9. Jakimowicz, D., van Wezel, G.P., 2012. Cell division and DNA segregation in

Streptomyces: How to build a septum in the middle of nowhere? Mol. Microbiol.

85, 393–404. doi: 10.1111/j.1365-2958.2012.08107.x

10. Kim, Y.M., Kim, J.H., 2004. Formation and dispersion of mycelial pellets of

Streptomyces coelicolor A3(2). J. Microbiol. 42(1), 64-7.

11. Koebsch, I., Overbeck, J., Piepmeyer, S., Meschke, H., Schrempf, H., 2009. A

molecular key for building hyphae aggregates: The role of the newly identified

Streptomyces protein HyaS. Microb. Biotechnol. 2(3), 343-360. doi:

10.1111/j.1751-7915.2009.00093.x

12. Kumar, P., Dubey, K.K., 2015. Current trends and future prospects of lipstatin: a

lipase inhibitor and pro-drug for obesity. RSC Adv. 5, 86954-66.

doi:10.1039/C5RA14892H

13. Kumar, P., Dubey, K.K., 2016. Modulation of fatty acid metabolism and

tricarboxylic acid cycle to enhance the lipstatin production through medium

engineering in *Streptomyces toxytricini*. Bioresour. Technol. 213, 64-8

doi:10.1016/j.biortech.2016.01.133

14. Kumar, P., Dubey, K.K., 2017. Mycelium transformation of Streptomyces

toxytricini into pellet: role of culture conditions and kinetics. Bioresour.

Technol. 228, 339-347. doi: 10.1016/j.biortech.2017.01.002

15. Labajová, N., Baranova, N., Jurásek, M., Vácha, R., Loose, M., Barák, I., 2021.

Cardiolipin-Containing Lipid Membranes Attract the Bacterial Cell Division

Protein DivIVA. Int. J. Mol. Sci. 22(15), 8350. doi: 10.3390/ijms22158350

16. Noens, E.E.E., Mersinias, V., Willemse, J., Traag, B.A., Laing, E., Chater, K.F.,

Smith, C.P., Koerten, H.K., van Wezel, G.P., 2007. Loss of the controlled

localization of growth stage-specific cell-wall synthesis pleiotropically affects

developmental gene expression in an ssgA mutant of *Streptomyces coelicolor*.

Mol. Microbiol. 64(5), 1244-1259

17. Ribeiro, R. M. M. G. P., Esperança, M. N., Sousa, A. P. A., Neto, Á. B., &

Cerri, M. O. (2021). Individual effect of shear rate and oxygen transfer on

clavulanic acid production by Streptomyces clavuligerus. *Bioprocess and

biosystems engineering*, *44*(8), 1721–1732. https://doi.org/10.1007/s00449-021-

02555-1

18. Rioseras, B., López-García, M.T., Yagüe, P., Sánchez, J., Manteca, A., 2014.

Mycelium differentiation and development of *Streptomyces coelicolor* in lab-

scale bioreactors: Programmed cell death, differentiation, and lysis are closely

288 linked to undecylprodigiosin and actinorhodin production. Bioresour. Technol.

289 151, 191-8. doi:10.1016/j.biortech.2013.10.068.

290 19. Ultee, E., van der Aart, L.T., Zhang, L., van Dissel, D., Diebolder, C.A., van

291 Wezel, G.P., Claessen, D., Briegel, A., 2020. Teichoic acids anchor distinct cell

292 wall lamellae in an apically growing bacterium. Commun. Biol. 3, 314. doi:

293 10.1038/s42003-020-1038-6

294 20. van Dissel, D., Willemse, J., Zacchetti, B., Claessen, D., Pier, G.B., van Wezel,

295 G.P., 2018. Production of poly-β-1, 6-N-acetylglucosamine by MatAB is

296 required for hyphal aggregation and hydrophilic surface adhesion by

297 Streptomyces. Microbial Cell 5(6), 269-279. doi: 10.15698/mic2018.06.635

298 21. van Veluw, G.J., Petrus, M.L., Gubbens, J., de Graaf, R., de Jong, I.P., van

299 Wezel, G.P., Wösten, H.A., Claessen, D., 2012. Analysis of two distinct

300 mycelial populations in liquid-grown *Streptomyces* cultures using a flow

301 cytometry-based proteomics approach. Appl. Microbiol. Biotechnol. 96(5),

302 1301–12. doi: 10.1007/s00253-012-4490-5

303 22. Vassallo, A., Palazzotto, E., Renzone, G., Botta, L., Faddetta, T., Scaloni, A.,

304 Puglia, A.M., Gallo, G., 2020. The Streptomyces coelicolor small ORF trpM

305 stimulates growth and morphological development and exerts opposite effects

306 on actinorhodin and calcium-dependent antibiotic production. Front. Microbiol.

307 11, 224. doi: 10.3389/fmicb.2020.00224

308 23. Wang, H., Zhao, G., Ding, X., 2017. Morphology engineering of Streptomyces

309 coelicolor M145 by sub-inhibitory concentrations of antibiotics. Sci. Rep. 7,

310 13226. doi:10.1038/s41598-017-13493-y

24. Willemse, J., Ruban-Osmialowska, B., Widdick, D., Celler, K., Hutchings, M.I.,

van Wezel, G.P., Palmer, T., 2012. Dynamic localization of Tat protein transport

machinery components in *Streptomyces coelicolor*. J. Bacteriol. 194(22), 6272-

6281. doi: 10.1128/JB.01425-12

25. Wu, Y., Kang, Q., Zhang, L. L., & Bai, L. (2020). Subtilisin-Involved

Morphology Engineering for Improved Antibiotic Production in

Actinomycetes. *Biomolecules*, *10*(6), 851.

https://doi.org/10.3390/biom10060851

26. Yue, C., Xu, H., Yu, Y., Yu, X., Yu, M., Zhang, C., You, Q., Xia, S., Ding, Z.,

Fu, H., Zeng, X. and Li, F., 2021. Improvement of natamycin production by

controlling the morphology of *Streptomyces gilvosporeus* Z8 with microparticle

talc in seed preculture. J Chem Technol Biotechnol, 96: 1533-

1542. https://doi.org/10.1002/jctb.6668

Figures

Fig. 1: Analysis of pellet morphology and mycelial structure by scanning electron

microscopy. **A**- Morphology of pellet at 10μM scale, **B-** Surface of pellet showing free

mycelia which show growth and branching at 10μM scale, **C**- Structure of hyphae

showing association with each other and branching at growing tips (2μM scale), **D**-

Apical growth and branching of hyphae (2μM scale).

Fig. 2: Analysis of extracellular and intracellular proteins of *S. toxytricini* by SDS-

PAGE. Lane 1 represents total intracellular proteins, lane 2,3 represents total

extracellular proteins isolated from culture medium. Marker lane is molecular weight

marker.

Fig. 3: Proposed mechanism for pellet formation. **A**-Diffused mycelia, **B**-Mycelia are coming together, **C**-mycelia are clumping, **D**-mycelial clumps are more visible and a reaction centre type structure is formed which possibly directs pellet formation, **E,F,G**-mycalial clumps are more condensed, **H**-Pellet formed and periphery having mycelia