The effectiveness of selection in a species affects the direction of amino acid frequency evolution

Hanon McShea[1], Catherine Weibel[2,3], Sawsan Wehbi [4], Peter Goodman[5], Jennifer E. James[2,6], Andrew L. Wheeler[4], Joanna Masel[2]

[1]Department of Earth System Science, Stanford University

[2]Department of Ecology & Evolutionary Biology, University of Arizona

[3]Department of Applied Physics, Stanford University

[4]Graduate Interdisciplinary Program in Genetics, University of Arizona

[5]Statistics & Data Science, University of Arizona

[6]Department of Ecology and Genetics, Uppsala University

Keywords

codon adaptation, amino acid cost, folding stability, thermostability, drift barrier, trait evolution,

mutation bias

Significance statement

According to the nearly neutral theory of molecular evolution, selection is less able to distinguish

between similar alleles in species with lower population size. We identify which amino acids are

subject to such weak preferences – these tend to be smaller, to be less costly to make, to promote

structural disorder of proteins, and to be enriched in thermophiles relative to mesophiles. The latter

agrees with theories of marginal protein stability under mutation-selection-drift balance.

Abstract

Nearly neutral theory predicts that species with higher effective population size (N_e) are better able

to purge slightly deleterious mutations. We compare evolution in high-N_e vs. low-N_e vertebrates

to reveal which amino acid frequencies are subject to subtle selective preferences. We take three

complementary approaches, two measuring flux and one measuring outcomes. First, we fit non-

stationary substitution models of amino acid flux using maximum likelihood, comparing the high-

N_e clade of rodents and lagomorphs to its low-N_e sister clade of primates and colugos. Second, we

compare evolutionary outcomes across a wider range of vertebrates, via correlations between

amino acid frequencies and N_e. Third, we dissect the details of flux in human, chimpanzee, mouse,

and rat, as scored by parsimony – this also enables comparison to a historical paper. All three

methods agree on which amino acids are preferred under more effective selection. Preferred amino

acids tend to be smaller, less costly to synthesize, and to promote intrinsic structural disorder.

Parsimony-induced bias in the historical study produces an apparent reduction in structural disorder, perhaps driven by slightly deleterious substitutions. Within highly exchangeable pairs of amino acids, arginine is strongly preferred over lysine, and valine over isoleucine, consistent with more effective selection preferring a marginally larger free energy of folding. These two preferences match differences between thermophiles and mesophilic relatives. These results reveal the biophysical consequences of mutation-selection-drift balance, and demonstrate the utility of nearly neutral theory for understanding protein evolution.

Introduction

Tomoko Ohta (1973) proposed that there is evolutionarily significant variation among species in the ability of selection to purge weakly deleterious mutations. According to Ohta's nearly neutral theory of evolution, species with small "effective population sizes" (N_e) are less able to purge slightly deleterious mutations and will thus retain more of them. By estimating N_e from neutral genetic diversity, Lynch and Conery (2003) used the nearly neutral theory to explain why species with smaller effective population sizes have more bloated genomes. Species with larger effective population sizes also show stronger synonymous codon usage bias due to selection (Akashi, 1996; Galtier et al., 2018; Subramanian, 2008; Weibel et al., 2024). Here we ask whether differences among species in amino acid frequencies can similarly be explained by the nearly neutral theory of evolution, and if so, what this tells us about the biophysical basis of intrinsic selective preferences.

We use three complementary approaches to test, for the first time, whether differences among species in the effectiveness of selection can predict amino acid frequencies. First and

primarily, we fit non-stationary amino acid substitution models on the basis of maximum

likelihood (ML). This is a major departure from assumptions of stationarity built into most models

of amino acid substitution (Bui et al., 2021; Dang et al., 2022; Le et al., 2012; Le & Gascuel, 2008;

Whelan & Goldman, 2001). To reduce the number of fitted parameters, we fit equilibrium amino

acid frequencies for different parts of the tree, while holding the exchangeability matrix

(Q.mammal; Bui et al., 2021) fixed. We fit all 20 amino acid frequencies, going beyond previous

work that used only one (Muñoz-Gómez et al., 2022) or two (Groussin et al., 2013) parameters to

summarize amino acid frequencies. We are able to get all amino acid frequencies because the

cogent3 code base (Kaehler et al., 2015; Schranz et al., 2008; Verbyla et al., 2013) enables us to

group branches together, avoiding the need to estimate 19 free parameters for every branch. We

then compare the fitted equilibrium amino acid frequencies within high-N_e (Phifer-Rixey et al.,

2012) Glires branches (including both Rodentia and Lagomorphs) to those within low-N_e (Tenesa

et al., 2007) Primatomorpha branches (including Primates plus colugos). A difference in the

effectiveness of selection between these two sister clades is well supported (Eőry et al., 2010;

Halligan et al., 2010; Keightley et al., 2005), and genome data is high-quality.

Second, we look at a broader species range (all vertebrates), to confirm that results from

our flux methods are driven by N_e rather than by other idiosyncrasies of Glires or Primatomorpha

biology. We examine whether final proteome-wide amino acid frequencies – rather than rates of

flux – are correlated with N_e. We quantify N_e using a recently developed metric, the Codon

Adaptation Index of Species (CAIS; Weibel et al., 2024). CAIS captures the fraction of

synonymous sites within the genome for which selection is able to overcome drift. Because the

distribution of selection coefficients is likely to be similar for different vertebrate species, CAIS

estimates N_e. CAIS uses Kullback-Leibler divergence to quantify the degree to which codon bias

departs from that expected from the %GC content of the species. It is calculated with respect to a

standardized set of amino acid frequencies and therefore, by construction, also avoids confounding

with amino acid frequencies. Weibel et al. (2024) found that high-CAIS species have higher

proteome-wide intrinsic structural disorder, demonstrating the metric's utility for revealing how

the effectiveness of selection shapes fundamental physical properties of macromolecules.

Conveniently, CAIS requires only a single complete genome for its estimation, unlike traditional

N_e estimates which require data on polymorphisms and mutation rate. CAIS directly measures the

property of interest, namely the degree to which weak selection is able to exert subtle selective

preferences within a species.

Third, noting that our ML method fits only enough free parameters to capture net amino

acid frequency change, we use parsimony to count instances of change, to determine which amino

acid pairs obey detailed balance vs. contribute to overall amino acid frequency change. This

parsimony approach is also partially motivated by resolving a historical controversy, after Jordan

et al. (2005) claimed to have discovered universal patterns of amino acid gains and losses across

the tree of life, dating back to the origins of the genetic code.

The findings of Jordan et al. (2005) were subject to intense criticism on three counts. First,

the method assumed that all derived alleles were substitutions, when some instead could be

polymorphisms – many of them slightly deleterious – segregating in the population (Hurst et al.,

2006; McDonald, 2006). Our analysis filters out polymorphic sites to obtain a dataset of true

substitutions (Figure 1A), something it was not possible to do at the time Jordan et al. (2005) was

published. Removing contamination by slightly deleterious polymorphisms should better reveal

any true universal trends. To ensure the availability of polymorphism data, we restrict our analysis

to two pairs of sister species. We compare recent fluxes in the primates *Homo sapiens* and *Pan*

troglodytes to a shared set of sequences in the rodents *Rattus norvegicus* and *Mus musculus*.

A second potential problem with this parsimony-based method is that even a set of true

substitutions will be enriched for slightly deleterious substitutions (McDonald, 2006). This is

because the brevity of their fixed status makes them more likely to survive ascertainment bias

when in the derived state than when in the ancestral (Figure 1B). According to this argument,

removing slightly deleterious polymorphisms should retain the direction of flux while reducing its

magnitude; slightly deleterious polymorphisms should mirror slightly deleterious fixations. A

similar argument applies with respect to equilibrium amino acid frequencies; a rare amino acid is

comparable to a deleterious allele (McDonald, 2006), explaining earlier, similar results by

Zuckerkandl et al. (1971), who observed an apparent increase in rare amino acids.

Goldstein and Pollock (2006) noted a third artifact that depends on the pattern of branch

lengths, combined with rate heterogeneity (Figure 1C). In their simulations, the length of the

branch between sister species relative to their branch with the outgroup affected which types of

substitutions were parsimony-informative and thus detectable. This bias in detection can produce

the appearance of strong net flux, even in simulations of a stationary and time-reversible model of

protein evolution, for which net flux is by definition zero. Because Jordan et al. (2005) selected

their triplets of species on the basis of similar branch length patterns (sister taxa pairwise amino

acid divergence <15%, outgroup <25%), as do we, the same branch-length artifact will appear

within any trends shared across different species triplets. Our approach is to statistically

differentiate taxonomically specific patterns in flux from "shared flux" values that capture both

this artifact and that from slightly deleterious fixations, as well as any true universal trends.

Figure 1: Artifacts of the parsimony net flux method and our approaches to them. Every color

change along a branch represents a substitution at a focal site. (A) We removed polymorphisms.

(B and C) Deleterious substitutions and branch length issues are expected to affect rodent and

primate branches similarly. We therefore subtracted flux that was seen in both rodent and primate

branches, leaving a metric that describes how rodent flux differs from primate flux (Figure 3).

V→C is used as an example of a slightly deleterious mutation prone to reversion, while V→L and

V→S are used as examples of effectively neutral substitutions. On the left side of (C), when sister

branch lengths are short, there will be greater sampling of fast-evolving sites, and when sister

branch lengths are long, it is easier to detect slow-evolving sites, because they are less prone to

multiple events, both within the long sister branches and in the still longer outgroup branch.

Here we fit non-stationary models of amino acid evolution to discover which amino acids

are favored by weak selection that is effective only in higher N_e species. We compare to parsimony-

based results to resolve a historic controversy, and to obtain a detailed picture of which amino acid

substitutions contribute to the non-stationarity. We confirm these flux results across a broader

taxonomic range by describing how amino acid frequency outcomes correlate with N_e.

Results

Low N_e results in costly amino acids produced by mutation bias

Cheap (Figure 2A), small (Figure 2B), and structural disorder-promoting (Figure 2C)

amino acids are at higher equilibrium frequency in the higher N_e clade, according to our maximum

likelihood flux method. Amino acids preferred in the higher N_e clade also tend to be dietarily non-

essential for animals (using Costa et al., 2015's conservative list of essential amino acids) (Figure

2A; t-test p = 0.006). They also have somewhat lower "stickiness," an empirical measure of an

amino acid's tendency to be in protein-protein interfaces rather than in non-interface surfaces

(Pearson's R^2 = 0.24, p = 0.03; Levy et al., 2012) and higher polarity (Pearson's R^2 = 0.22, p =

0.04). Interestingly, both disorder propensity and molecular volume also predict the marginal

fitness effects of each amino acid when expressed as part of a random peptide in the *Escherichia*

coli experiments of Neme et al. (2017), as inferred by Kosinski et al. (2022). Marginal fitness

effects in *E. coli* accordingly correlate even more strongly with amino acid preferences inferred

from non-stationary substitution models (Figure 2D). This is due to size and disorder; there is no

significant explanatory power left in the residual 48% of the marginal fitness effects calculated by

Kosinski et al. (2022) (Pearson's $R^2 = 0.05$, p = 0.3).

We note that the pervasive AT bias of mutation (Haag-Liautard et al., 2008; Hershberg &

Petrov, 2010; Lynch, 2010; Lynch et al., 2008), including but not limited to methyl-cytosine

deamination, makes the retention of GC-rich amino acids difficult. Figure 2E confirms that

retention of GC-rich amino acids is achieved more often under more effective selection. GC-rich

amino acids are also favored in highly expressed ($R^2 = 0.27$, p = 0.02; Jansen & Gerstein, 2000)

and slowly evolving ($R^2 = 0.20$, p = 0.05; Cherry, 2010) proteins. However, across all 20 amino

acids, we do not find a significant correlation between our ML flux difference scores and these

papers' estimates of amino acid enrichment as a function of expression or evolutionary rate.

Figure 2: The amino acids preferred in Glires vs. Primatomorpha evolution are cheaper, lower

%GC, smaller, more prone to structural disorder, and generally more benign. The y-axis "Flux

difference (ML)" indicates equilibrium Glires frequencies – Primatomorpha frequencies estimated

under a non-stationary model of amino acid substitution. This model was fitted by maximum

likelihood to an aligned set of orthologs, with one set of equilibrium amino acid frequencies fitted

to branches within a clade of 12 Glires species, another to branches within a sister clade of 16

Primatomorpha species, and a third to their shared root (see Methods). The time-reversible

exchangeability matrix (Q.mammal; Bui et al., 2021) was held constant. Error bars show ± 1

standard error; R and p-values on figure panels are for Pearson's R correlation. (A) Biosynthetic

cost in *Saccharomyces cerevisiae* under aerobic conditions, as a count of high-energy phosphate

bonds (Raiford et al., 2008); we obtain modestly lower $R^2 = 0.50$ and $p = 5e\text{-}04$ for aerobic cost

on a linear scale. Colors show essentiality in animals (Costa et al., 2015). (B) %GC, calculated as

average %GC for each amino acid's set of codons in the standard genetic code. (C) Mean

molecular volume, units in Angstroms cubed (Tsai et al., 1999). (D) Disorder propensity (Theillet

et al., 2013). (E) Marginal effects of amino acids when expressed as part of random peptides in *E.*

coli, in units of the effect on genotype frequency per cycle of growth and dilution (Kosinski et al.,

2022).

Parsimony fluxes shared by rodents and primates are likely driven by artifact

According to our parsimony method, mice and rats have very similar net amino acid fluxes

to each other (Supplementary Figure 1), as do humans and chimpanzees. This is expected, since

%GC content, effective population size, and many other possible influences are similar within

each closely related species pair. We therefore calculated "rodent" flux values by combining the

amino acid gain and loss counts for mouse and rat branches, and "primate" flux values by

combining the counts for human and chimpanzee branches (numbers in Supplementary Table 1).

While our rodent and primate net fluxes are similar to those of Jordan et al. (2005), they

are generally of smaller magnitude, particularly in primates (Figures 3A and 3B). We interpret this

as a weaker version of the same bias toward slightly deleterious mutations within our fluxes

compared to those of Jordan et al. (2005). As discussed in the Introduction (Figure 1B), fixation

events scored by parsimony are also expected to be biased toward slightly deleterious mutations.

This is because slightly deleterious mutations might sometimes reach fixation but will revert

relatively quickly to a less deleterious state along subsequent branches, and therefore will be scored

incorrectly.

The main difference between our method and that of Jordan et al. (2005) is that we exclude

polymorphisms. Our primate species pair is more closely related than our rodent species pair, i.e.

it has fewer true substitutions. If we assume similar levels of polymorphisms in both, the smaller

true divergence in primates means that removing polymorphism has a proportionately larger

impact on primate fluxes than rodent fluxes. This is compatible with the observed greater deflation

in net flux magnitudes that we see for primates (Fig. 3A) than for rodents (Fig. 3B).

Figure 3: Removing polymorphisms reduces the magnitude of parsimony flux, especially for primates (A and B). Net fluxes depend less on taxon when calculated by parsimony (C) than by maximum likelihood (D). The parsimony-based net flux for each amino acid = (# gains – # losses) / (# gains + # losses), making values greater than 0 indicate a tendency towards gain. Error bars show ± 1 standard error; R^2 and p-values on figure panels are for Pearson's R correlation. (A and B) Black lines show y = x. Shaded areas show sign reversal. (C) Red line shows the fit of a type II linear regression (y = 0.664x – 0.0154). Shared flux and flux difference are calculated from this regression line, as discussed in the main text. (D) The correlation between amino acid frequency

change from root in rodents vs. primates is attributable to one outlier, a low inferred serine

frequency at the root.

Remaining artifacts in our polymorphism-free version of the parsimony method should be

shared between rodents and primates (in addition to true universal trends, if any). In contrast,

differences in net parsimony fluxes may reflect the more effective selection in rodents relative to

the mutation bias that overwhelms selection in primates. By statistically isolating shared vs.

species-specific patterns in flux, any universal trend or artifact should be reflected in the shared

component, while the species-specific component will allow us to estimate the degree to which

each of the 20 amino acids is favored vs. disfavored in more effectively adapted rodents relative

to less well adapted primates. To do this, we fit a type-II linear regression model (Figure 3C, red

line) and calculated the projection along the model line ("shared flux") for each amino acid, as

well as the shortest distance from that line ("flux difference"). The shared component thus reflects

fluxes that are common to both rodents and primates, and the flux difference reflects subtle

differences between the patterns of fluxes in the two groups. The sign we give to the flux difference

allows it to be interpreted as a rodent-driven component, i.e., a positive value indicates that the

amino acid is gained in rodents to a greater degree (or lost to a lesser degree) than in primates.

Although Glires and Primatomorpha ML fluxes are also correlated (Figure 3D; numbers in

Supplementary Table 2), this correlation depends on a single outlier amino acid, serine. Serine

frequency appears to be abnormally low in the common ancestor, and thus regresses to the mean

(by increasing) in both clades. Serine is similarly abnormally low in the later roots of Glires and

of Primatomorpha, according to non-stationary substitution models fitted to each clade in isolation

(Supplementary Table 3). There is no significant correlation between Glires and Primatomorpha

ML fluxes once serine is excluded (without serine, Pearson's $R^2 = 0.11$, p = 0.2 and Spearman's

$R^2 = 0.11$, p = 0.2). Nor is the relationship statistically significant under a Spearman's ranked test,

even with the inclusion of serine (p = 0.06). This serine outlier at the root does not affect our ML

"flux difference" between Glires and Primatomorpha.

Figure 4: The slightly deleterious mutations artifactually captured by the parsimony method tend

to reduce intrinsic structural disorder. (A) Jordan et al. (2005) capture something similar to our

shared flux metric. (B) Shared flux describes an increasing frequency of low-frequency amino

acids. Frequency is measured as the average of human and chimpanzee amino acid frequencies,

with similar results (Pearson's $R^2 = 0.30$, p = 0.01) for the average of mouse and rat amino acid

frequencies. (C) Shared flux describes substitutions that reduce structural disorder. Disorder

propensities of amino acids were measured by Theillet et al. (2013) on the basis of

overrepresentation in disordered vs. well-folded proteins and protein regions. Error bars show ± 1

standard error; R^2 and p-values on figure panels are for Pearson's correlation.

As expected for a successful partition, the shared component of parsimony fluxes has a

strong positive relationship with Jordan et al.'s (2005) average flux (Figure 4A), while the flux

difference component has none (Pearson's R^2 = 0.03, p = 0.4). Results are similar albeit slightly

weaker for Jordan et al.'s (2005) more complex gain/loss rate metric. Shared flux describes the

gain of lower-frequency amino acids (Figure 4B) expected from regression to the mean, as

observed by Zuckerkandl et al. (1971).

From our Figure 2 results, if the shared flux reflects artifact, we expect the slightly

deleterious changes it represents to be reflected in disorder, size, cost, essentiality, %GC,

stickiness, and polarity. As predicted by the artifact hypothesis, amino acid changes described by

shared flux lead to decreased disorder (Figure 4C), increased stickiness (Pearson's R^2 = 0.40, p =

0.003), increased mean relative solvent accessibility (RSA) (Pearson's R^2 = 0.37, p = 0.004; Tien

et al., 2013), and increased polarity (R^2 = 0.31, p = 0.01), although they do not predict size, cost,

essentiality, or %GC. While results for Jordan et al.'s average flux are weaker than for our shared

flux (Pearson's R^2 = 0.35, p = 0.006 for Jordan et al.'s flux vs. disorder, relationships with

stickiness, polarity, and mean RSA are insignificant), in light of the strong correlation in Figure

4A, we nevertheless conclude that the results of Jordan et al. are likely dominated by artifact. The

species-specific component of flux does not correlate with any of these properties.

This leaves open the question of which artifact dominates – that toward slightly deleterious

substitutions in Fig. 1B or the more subtle branch length ascertainment bias illustrated in Fig. 1C.

If we assume that the shared component of parsimony flux preferentially represents fast-evolving

sites (Fig. 1C), we note that these are enriched in disordered regions, where slightly deleterious

mutations will tend to decrease disorder through regression to the mean. The fact that Jordan et

al.'s average flux corresponds to falling disorder is thus consistent with contributions from both

artifacts.

Three methods agree on which amino acids are preferred by selection

The slopes of amino acid frequencies as a function of effective population size describe

outcomes rather than fluxes, across a larger range of vertebrate species (numbers in Supplementary

Table 4). We use the Codon Adaptation Index of Species (CAIS) to measure effective population

size, capturing the proportion of codons subject to effective selection, while controlling for GC

and amino acid frequencies (see Methods).

Our three different metrics of selective preferences among amino acids are in good

agreement (Figure 5). They also reveal selective preferences between biophysically similar and

highly exchangeable amino acids (pairs joined by arrows in Figure 5, discussed in detail below).

As for our main metric of selective preference (ML flux difference; discussed above), codon %GC

also predicts parsimony flux difference (Pearson's R^2 = 0.23, p = 0.03) and preference in

vertebrates (Pearson's R^2 = 0.69, p = 5e-06). Preference in vertebrates is also predicted by mean

entropic penalty of folding (Pearson's R2 = 0.23, p = 0.03; Doig & Sternberg, 1995) and marginal

fitness in *E. coli* (Pearson's R^2 = 0.38, p = 0.004). However, the latter is only explained by

Kosinski's model residuals (Pearson's $R^2 = 0.22$, p = 0.04) – unlike ML flux difference, neither preference in vertebrates nor parsimony flux difference is significantly predicted by size, cost, polarity, disorder propensity, stickiness, or essentiality in animals. These results are consistent with our ML measure being superior to the other two, but given broad agreement, we nevertheless use the other two metrics to conduct follow-up analyses not easily possible via ML.

Figure 5: Three distinct methods with different biases recover similar selective preferences among amino acids. "Preference in vertebrates" indicates slopes from linear regressions of amino acid frequency on CAIS. "Flux difference (parsimony)" indicates the shortest distance of an amino acid to the model II fit shown in Figure 2. "Flux difference (ML)" indicates equilibrium Glires frequencies – Primatomorpha frequencies over the rodent tree estimated under a non-stationary model of protein evolution. Error bars show ± 1 standard error; R^2 and p-values are for Pearson's correlation. Connected pairs are biophysically similar, highly exchangeable amino acids for which all metrics show a marked preference in the same direction.

Different kinds of protein domains have mostly similar amino acid preferences

Vertebrate amino acid preferences are similar in ancient vs. recently born Pfam (Mistry et al., 2021) domains (Figure 6A). Partial exceptions are preferences against lysine and glutamic acid, and preferences for leucine and glycine, both of which are stronger in young than in old Pfams. Transmembrane vs. non-transmembrane domains also have similar amino acid preferences, despite operating in different biophysical contexts (Figure 6B). However, there are minor differences between the two with respect to the strength of preferences for leucine, valine, glycine, and arginine. In high CAIS species, transmembrane proteins also avoid phenylalanine more strongly than non-transmembrane domains do, consistent with its role in increasing both membrane permeability and heteromeric interactions in transmembrane domains (Kwon et al., 2016; Perkins & Vaida, 2017).

Figure 6: Vertebrate amino acid preferences are similar in different groups of protein domains. Error bars show ± 1 standard error; R^2 and p-values on figure panels are for Pearson's correlation.

Domain ages were taken from James et al. (2021), in which Pfam domains were assigned a date

of evolutionary origin using TimeTree (Kumar et al., 2017). (A) Vertebrate protein domains that

emerged prior to the last eukaryotic common ancestor (LECA) are categorized as "old", and

protein domains that emerged after the divergence of animals and fungi from plants as "young".

(B) Transmembrane vs non-transmembrane status was determined by James et al. (2021) using the

program tmhmm (Krogh et al., 2001). In both panels, the black y=x lines indicate equal effects in

the two sets of Pfam domains. Presence in an off-diagonal, shaded quadrant indicates that an amino

acid trend has different signs for the two subsets of the data. Error bars indicate +/- one standard

error.

Marked preferences between evolutionarily exchangeable pairs of amino acids

Selection on subtle differences between similar amino acids may be informative regarding

the biophysics of what is preferred. We selected amino acid pairs that have exchangeability values

greater than 1 in all BLOSUM (Henikoff & Henikoff, 1992) and PAM matrices (Dayhoff, 1972),

covering the range of branch lengths within our vertebrate dataset. These pairs are [R,K], [V,I],

[E,Q], and [M,L]. We further focus on the two pairs for which our three methods agree on the

direction of preference: [R,K] and [V,I]. These pairs are connected by grey lines in Figure 5. In

both cases, the strength of preference is large.

Arginine (R) is preferred over lysine (K). Notably, these are the only two amino acids with

sidechains that are positively charged at neutral pH. Both amino acids are basic and aliphatic, with

positively charged functional groups on an alkyl side chain, and both form salt bridges and other

non-covalent interactions in protein tertiary structure. The parsimony flux data shows not only that

rodents gain R and lose K, but more specifically that K residues are replaced with R (Figure 7, top

left). Out of all ancestral rodent K residues that we observed substitutions away from, 48% became

R (1452 of 3037 ancestral Ks), and 35% of derived rodent R residues came from K (1452 of 4126

derived Rs). This pattern is also found to a lesser degree in primates (113 of 526 and 705, or 21%

and 16%, respectively; Figure 7 top right). The next two most common substitutions from K in

rodents are to the polar uncharged amino acids N (261 of 3037) and T (188 of 3037). In addition

to K, R also comes from Q (609 of 4126), G (482 of 4126), S (274 of 4126), L (181 of 4126), and

others (Figure 7 top; Supplementary Table 5).

Chemical differences between the two otherwise similar amino acids may explain the shift

— K has the largest median hydrophobic water-accessible surface area in folded proteins, much

larger than R (47 vs. 20 Å^2; Lins et al., 2003). I.e., at sites where an exposed K residue is replaced

by an R, the exposed hydrophobic surface area is likely to be greatly reduced. This may increase

the marginal stability of the folded protein. In further support of selection for stability, R can form

more hydrogen bonds than K, which could have stabilizing effects enthalpically in the formation

of secondary and tertiary structure, and entropically in the ordering of water molecules at the

protein's surface. Taking into account protein interactions rather than folding alone, we note that

K is underrepresented in protein-protein interfaces relative to non-interface surfaces (Levy et al.,

2012), while R is about equally likely to be in either. Similar considerations might also apply to

protein-lipid and protein-nucleic acid interactions. Interface effects may explain the paucity of

X→K substitutions: introducing K can destroy an interface and thus be strongly deleterious. X→R

substitutions might be far more numerous because R is tolerated equally well in multiple contexts.

The small hydrophobic amino acid valine (V) is preferred over isoleucine (I), with I→V

substitutions again predominating in rodents but not primates (Figure 7 bottom). Isoleucine is very

harmful in random peptides in *E. coli* (Kosinski et al., 2022), where it is more than twice as

deleterious as the next most harmful amino acid. V and I have similar median hydrophobic surfaces

areas (5 and 4 Å^2, respectively). However, V side chains have a smaller entropic penalty upon

protein folding than I or the structurally similar alternative, L (-0.43 vs. -0.76 vs. -0.71 kcal/mol;

Doig & Sternberg, 1995). Such a small but fundamental effect on stability, given other similarities,

could explain the preference for V as an alternative to I. Considering the full set of substitutions

in Figure 7 (bottom), V shows general patterns of gain, with loss to M as an exception.

We note that in both [K,R] and [I,V] pairs, the less-preferred amino acid has an A or T

nucleotide whereas the more-preferred one has a G or C. Rodents have slightly higher %GC than

primates (42% vs. 41%; Weibel et al., 2024), so if the corresponding shift in mutation bias is still

in process of shifting amino acid frequencies, a recent increase in %GC could contribute to

preferences between similar amino acids inferred from flux. However, we observed the same

preferences using our CAIS approach, which is corrected for %GC.

To better understand the effects of mutation bias, we note that they should be most

pronounced at CpG sites. Four of the six R codons start with a CpG. C→T deamination at the first

position would yield a cysteine (C), tryptophan (W), or stop codon, none of these being aligned

with the direction of selective preference. In agreement with an expected greater relative role for

mutation bias over selection in low N_e species, we see more frequent R→C and R→W

substitutions in the raw primate parsimony flux data than in the rodent (Supplementary Table 6).

T→M, A→V, P→L, and S→L can also result from CpG deamination, and of these, T→M and

P→L are both more common in primates than in rodents, and both are counter to selective

preference (supported by all three metrics in Figure 5). There is no preference for A vs. V or S vs.

L under more effective selection, and these CpG substitutions are more common in rodents than

in primates. This demonstrates that although methyl-cytosine deamination at CpG sites is a

pervasive mutational force regardless of the effectiveness of selection, more effective selection is

needed to prevent the fixation of less-preferred amino acids resulting from CpG mutations.

Figure 7: Net substitutions involving biophysically similar pairs of amino acids in rodent and

primate lineages. Arrows represent all net substitutions involving K and R (top), or I and V

(bottom). Edge weights represent net flux between each pair (i.e. the number of K→R substitutions

minus the number of R→K substitutions), normalized according to the total number of

substitutions in that species pair (52,431 for rodents and 9,313 for primates). To give a sense of

scale, in rodents there is 1 net substitution from K to V, 179 from Q to R, and 793 from K to R. In

primates there is 1 net V→K substitution, 9 Q→R, and 57 K→R. Colors are for visual aid only

and represent amino acid chemical properties: hydrophobic amino acids are blue, positively

charged are red, polar non-charged are pink, negatively charged are gold, and other, "special case,"

amino acids are charcoal. Note that the time non-reversibility shown should not be taken as a

representation of the non-stationary amino acid frequencies – departures from stationarity are

represented in Figure 5.

Discussion

Smaller, cheaper, less sticky, more polar, more disorder-promoting amino acids are

selected for under higher N_e. These factors covary (i.e. disorder-promoting amino acids are

significantly cheaper, smaller, and less sticky; cheap amino acids are significantly smaller and less

sticky), making it hard to know their relative importance. Retaining amino acids that require high

GC, in the face of AT-biased mutation, also requires high N_e. There are particularly striking

preferences between highly exchangeable pairs of amino acids, with high N_e species preferring

arginine over lysine, and valine over isoleucine. We find similar preferences in old vs. young and

in transmembrane vs. non-transmembrane protein domains.

We used three complementary methods: two capturing how amino acid fluxes differ

between rodents vs. primates (based on maximum likelihood and on parsimony), and one capturing

amino acid frequency outcomes in a broader range of vertebrate species (based on correlation with

codon adaptation as a metric of N_e). Our correlation approach is vulnerable to different sites being

present in different taxa, while our flux approaches avoid this by using aligned orthologs. The

parsimony version of net flux is in theory highly vulnerable to the possibility that the effects of

high N_e of rodents and/or the low N_e of primates on amino acid frequencies have already reached

equilibrium. The fact that results agree with our maximum likelihood fluxes suggests that this is

not the case, in agreement with literature findings of recent changes in N_e (Deinum et al., 2015;

Jing et al., 2014; Lynch et al., 2023; Rajabi-Maham et al., 2008). Agreement between the methods,

with their different biases, gives us greater confidence. Our strongest results came from maximum

likelihood flux, with the parsimony approach providing more fine-grained information about

which substitutional paths drive amino acid frequency change, and the correlation approach

providing validation across a broader species range.

The parsimony flux method has been much criticized. We removed polymorphisms, which

are likely to contaminate substitution counts with non-fixed slightly deleterious mutations. This

improvement kept the direction but shrunk the magnitude of the flux patterns observed by Jordan

et al. (2005). This is compatible with shared patterns being artifacts from slightly deleterious

fixations and/or a phylogenetic tree that results in better ascertainment at faster evolving sites.

Whatever is responsible, the parsimony method universally produces gain of common amino acids

and loss of disorder-promoting amino acids. Slightly deleterious substitutions may be enriched for

those promoting order, and/or substitutions in rapidly evolving disordered regions may be subject

to a higher ascertainment rate.

Despite its weaknesses, the unique granularity of the parsimony flux method revealed that

within pairs of highly exchangeable amino acids, rodents exhibit a pronounced net flux from the

less favored member of the pair to the more favored, in a manner not found in primates (Figure 7).

It also revealed selection's ability to maintain CpG sites better in rodents than primates. Our

maximum likelihood method would have required an excessive number of free parameters to

achieve this; it instead held the exchangeability matrix constant at values estimated for mammals

as a whole. Improved methods for inferring individual substitutions (Monit & Goldstein, 2018)

can avoid parsimony biases, but are potentially vulnerable to a misspecified amino acid

substitution model.

A traditional view of structural biology might be skeptical of the value of information that

is restricted to amino acid frequencies. However, it is striking the degree to which amino acid

frequencies alone, absent information about amino acid ordering, are able to predict many

properties of a polypeptide, including translational efficiency, folding kinetics, and local protein

structure (Buhr et al., 2016; Saunders & Deane, 2010; Zarin et al., 2017, 2021). It is encouraging

that the direction of selective preference within [K, R] and [I, V] pairs is consistent with expected

selection on protein folding and stability, as supported by differences in median hydrophobic

surface area, hydrogen bonding capacity, and entropic penalty of folding. Note that the preferences

for R over K and V over I are statistical tendencies; the opposite might still hold at (a minority of)

individual sites. Because most proteins must fold (and unfold, and fold again), it is not surprising

that the thermodynamics of folding would be subject to selection detectable at the level of amino

acid frequencies. The specific biophysical hypotheses suggested by our observational work could

be experimentally tested by aggregating data on arginine vs. lysine or valine vs. isoleucine

mutations across a set of model proteins, where the effect of each mutation on marginal stability

and kinetics could be measured using multiplexed assays (Atsavapranee et al., 2021; Markin et al.,

2021; Tsuboyama et al., 2023).

Our observation of between-species variation in proteome-wide amino acid composition

shows different patterns from previous work on within-proteome variation. Some trends are

similar: like high-N_e species, highly expressed, slowly evolving proteins are enriched in amino

acids that are cheaper to synthesize (Akashi & Gojobori, 2002). However, more generally, amino

acids preferred in highly expressed proteins are not the same ones preferred under more effective

selection at the species level. Highly expressed proteins are enriched in A, G, V, R, and K (Jansen

& Gerstein, 2000), while slowly evolving proteins are enriched in A, G, V, D, I, M, N, Y and

depleted in R (Cherry, 2010), neither of which correlates with amino acid preference under more

effective selection. Disagreement may be because highly expressed proteins have different

structural and functional requirements than the proteome as a whole.

Amino acid substitutions during the evolution of thermophily do, however, parallel some

of our observations for more effective selection. Lecocq et al. (2021) found that thermophiles in

the archaeal order Methanococcales use R more often, and that K→R substitutions accompany

transitions to higher optimal growth temperature while R→K substitutions accompany transitions

to lower growth temperature. In thermophilic species of filamentous fungi, Van Noort et al. (2013)

found an excess of K→R substitutions using a similar method. Interestingly, Fontanillas et al. (2017) found an excess of R→K and V→I substitutions during a shift to cold adaptation of hydrothermal vent-dwelling polychaete worms, but did not find a corresponding excess of K→R or I→V substitutions in sister species that remained thermophiles throughout. The cold-adapted species had higher substitution rates and slightly higher dN/dS ratios (both <<<1), suggesting relaxed selection/reduced constraint relative to hot-adapted species. In all of these taxa, the amino acid preferred in hot conditions is the same as preferred by highly effective selection in our findings. Proteins tend to evolve marginal stability at mutation-selection-drift balance (Bloom, Raval, et al., 2007), with greater stability improving robustness to mistranscription and mistranslation (Drummond et al., 2005; Drummond & Wilke, 2008, 2009; Serohijos et al., 2012; Wilke et al., 2005), and potentially also mutation (Bloom, Lu, et al., 2007). Both more effective selection and hotter temperature will cause this stability to be slightly less marginal.

Pinney et al. (2021) identified protein sites at which amino acid identity correlated with optimal growth temperature across the bacterial domain. Opposite to our work and the studies reviewed above, at these sites, K and I are strongly associated with high growth temperature in bacterial enzymes, while R and V show no association with temperature. Interestingly, hot-adapted enzymes are also enriched in I-I, K-K, and I-K spatial interactions, and K-E salt bridges are also significantly enriched in hot-adapted enzymes, while R-D salt bridges are enriched in cold-adapted enzymes (K-D and R-E salt bridges were not significant). Perhaps against a background that is generally depleted in K and I by adaptation to hot conditions, the importance of the remaining K and I interactions becomes more important. In other words, the few K and I residues remaining may have important site-specific functions.

Proteome-wide amino acid frequencies are shaped by some combination of protein domain

birth, death, and tinkering. Differential rates of domain loss and duplication (James et al., 2023)

shift animal proteomes toward amino acid compositions favoring lower intrinsic structural disorder

(James et al., 2021). This is counterbalanced by *de novo* birth (Wilson et al., 2017). Our work here

instead focuses on descent with modification, which is likely to operate on faster timescales. This

focus is strictly enforced in both our parsimony and our likelihood-based flux analyses, by

considering only homologous sites shared by all four species. Our outcome results with CAIS also

suggest descent with modification, given that controlling for Pfam identity was not statistically

supported for individual amino acids (see Methods), and that similar results were previously found

for intrinsic structural disorder in which this control was included (Weibel et al., 2024).

Amino acids have distinctive biophysical properties, and selective preference among these

may be universal. The concordance of our mammalian study with previous results on random

peptides in *E. coli* supports the idea that for amino acid composition, "anything found to be true

of *E. coli* must also be true of Elephants" (Monod & Jacob, 1961). Sometimes the demands on a

position within a protein are different, e.g. for alpha helix vs. beta sheet vs. intrinsic disorder, for

surface vs. buried, for acid vs. base, or for lesser vs. greater rotameric degrees of freedom, but

given a particular demand, there are subtle preferences among amino acids that can deliver similar

things. This is what we have quantified. This is nearly neutral theory, as applied to the essence of

biochemistry.

Materials and Methods

Maximum likelihood flux estimates under a non-stationary amino acid substitution model

From the concatenated alignment of 5162 orthologous mammalian genes from (Wu et al.,

2018), we retained all 16 primates, all 12 rodents, two lagomorph species that are sister to rodents

(rabbit and pika) and one that is sister to primates (colugo). This results in one Glires clade

(Rodents plus Lagomorphs) and one Primatomorpha clade (Primates plus colugo), as illustrated in

Supplementary Figure 3. We used the cogent3 code base (Kaehler et al., 2015; Schranz et al., 2008;

Verbyla et al., 2013) to fit a non-stationary model of amino acid substitutions, given a fixed

topology of the species tree taken from the maximum clade credibility consensus tree from Vertlife

(Upham et al., 2019). Branch lengths were fitted by the model, but we did not model rate

heterogeneity among sites. We used the QMammal exchangeability matrix (Bui et al., 2021),

combined with three different equilibrium amino acid frequency vectors. One vector describes the

long-term equilibrium frequency along Primatomorpha branches, another along Glires branches,

and a third describes frequencies at the root. Inferred frequencies are given in Supplementary Table

2. The substitution rate matrix within a branch is then the matrix product of the exchangeability

matrix times the equilibrium frequencies.

To draw out the difference between high-N_e Glires and low-N_e Primatomorpha, we define

the "flux difference" for a given amino acid, as (Glires frequency – root frequency) –

(Primatomorpha frequency – root frequency) = Glires difference – Primatomorpha difference.

Parsimony approach to amino acid flux

We used the 65 amniote vertebrates multiple genome alignment, available on Ensembl (https://useast.ensembl.org/info/genome/compara/mlss.html?mlss=2041) (Yates et al., 2020). This alignment was generated using the Mercator-Pecan pipeline, which effectively handles rearrangements and duplications that have occurred during species' evolutionary history by organizing genomes into segments to create a homology map, prior to alignment (Paten et al., 2008). We downloaded the aligned coding regions for our four focal species from the Ensembl REST API server (https://rest.ensembl.org/, Ensembl 98, accessed during the months of November and December 2019) using our own Python scripts, using the mouse reference genome to retrieve alignment blocks based on mouse coding regions. To be included in our dataset, genomic regions for all four of our focal species had to be present in the multiple genome alignment.

The locations of polymorphic sites to be excluded were taken from the Ensembl variation database (Hunt et al., 2018). If a polymorphism reached a threshold minor allele frequency of 5% in any of the populations in which it was reported, that site was masked in the multiple species alignment. We note that due to the larger number of studies on human populations, there are more human polymorphisms recorded in Ensembl, and thus masked in our data, and fewer for rats and chimpanzees.

Aligned DNA sequences were then translated, and informative substitutions were identified, using our own Python scripts. To be included in our analysis, an amino acid site had to be identical in three species and different in the fourth, i.e., inferred by parsimony to occur on one of the branches between mouse-rat or human-chimp. For each species in our quartet, the

normalized gain/loss ratio (flux) for each amino acid was calculated by the same method as Jordan

et al. (2005), where flux = (# derived - #ancestral)/(#derived + # ancestral). Net flux values and

the shared and species-specific components are given in Supplementary Table 1.

Calculation of amino acid preference in vertebrates using CAIS

We use the CAIS calculations for 117 vertebrate species from Weibel et al. (2024). In that

paper, each species was assigned a fixed effect on the quantity of interest (in our case amino acid

frequency), while controlling for Pfam identity (Mistry et al., 2021) as a random effect. While

previously supported for intrinsic structural disorder, in our current analysis of amino acid

frequencies, controlling for Pfam identity was not supported ($p > 0.05$ for all 20 amino acids). We

therefore instead use the simple fractional content of each amino acid in the data for each species.

We did this both for the whole proteome, and for the set of annotated Pfam sequences. In both

cases, we then performed 20 weighted linear regressions to predict amino acid frequencies from

each species' CAIS value, and then used the regression slopes to summarize the degree of

preference for each amino acid, together with the associated standard errors on the slopes

(Supplementary Table 5). We do not correct for phylogeny (Felsenstein, 1985) before estimating

the slopes, because phylogenetic correction amplifies differences between closely related taxa,

whereas we are interested in differences that have might evolved slowly during longer periods of

relatively consistent effective population size. Phylogenetic correction via phylogenetically

independent contrasts (PIC) removes the relationship between CAIS slopes and flux metrics.

Linear models were implemented using the R package lme4 (Bates et al., 2015). Positive slopes

indicate that species with more effective selection (high CAIS) prefer that amino acid, negative

that they avoid it. We found that proteome-wide measurements yielded slopes with smaller total

standard error. Where not otherwise indicated, we therefore use the proteome-wide measurements. Comparison to CAIS slopes for the subset of the proteome that belongs to a Pfam (which are subsetted in Figure 6) is shown in Supplementary Figure 2.

Pfam subsets

In the vertebrate genomes, Pfams that emerged prior to the last eukaryotic common ancestor (LECA) are identified as "old", and Pfams that emerged after the divergence of animals and fungi from plants are identified as "young", as annotated by James et al. (2021). Pfams were assigned as either transmembrane or non-transmembrane by James et al. (2021) using the program Tmhmm (Krogh et al., 2001). Amino acid frequencies were calculated for each Pfam subset in each species.

Data availability

Full substitution tables used to calculate fluxes and CAIS slopes, as well as the values themselves and all code, calculations, and supplementary data are available at https://github.com/cyclase/aa_flux. All statistical modelling was done in R 3.5.1, 3.6.3, 4.2.1, or 4.3.1.

Acknowledgments

JM, CW, and PG were supported by the John Templeton Foundation (60814 and 62220). CW was supported by the Arnold and Mabel Beckman Foundation Scholars Program, the Western Alliance to Expand Student Opportunities (WAESO), Louis Stokes Alliance for Minority Participation (LSAMP) National Science Foundation (NSF) Cooperative Agreement (HRD-

1101728), and the UA/NASA Space Grant Undergraduate Research Internship program. HM was

supported by an NSF Graduate Research Fellowship. SW was supported in part by NASA

Astrobiology Program ICAR grant [80NSSC21K0592] to Betul Kacar. AW was supported by the

NIH [T32 GM132008]. We thank Minh Bui, Gavin Huttley, Luke Kosinski, Catherine DeBrule

Stark, Em Horst, and Bennett Kapili for helpful discussions. Work was partially motivated by a

bet with Eugene Koonin over a fine bottle of red wine regarding the reliability of the results of

Jordan et al.

Citations

Akashi, H. (1996). Molecular Evolution Between Drosophila melanogaster and D. simulans

Reduced Codon Bias, Faster Rates of Amino Acid Substitution, and Larger Proteins in D.

melanogaster. *Genetics*, *144*(3), 1297–1307. https://doi.org/10.1093/genetics/144.3.1297

Akashi, H., & Gojobori, T. (2002). Metabolic efficiency and amino acid composition in the

proteomes of Escherichia coli and Bacillus subtilis. *Proceedings of the National Academy

of Sciences*, *99*(6), 3695–3700. https://doi.org/10.1073/pnas.062526999

Atsavapranee, B., Stark, C. D., Sunden, F., Thompson, S., & Fordyce, P. M. (2021).

Fundamentals to function: Quantitative and scalable approaches for measuring protein

stability. *Cell Systems*, *12*(6), 547–560. https://doi.org/10.1016/j.cels.2021.05.009

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Bloom, J. D., Lu, Z., Chen, D., Raval, A., Venturelli, O. S., & Arnold, F. H. (2007). Evolution

favors protein mutational robustness in sufficiently large populations. *BMC Biology*, *5*(1),

29. https://doi.org/10.1186/1741-7007-5-29

Bloom, J. D., Raval, A., & Wilke, C. O. (2007). Thermodynamics of neutral protein evolution.

Genetics, *175*(1), 255–266. https://doi.org/10.1534/genetics.106.061754

Buhr, F., Jha, S., Thommen, M., Mittelstaet, J., Kutz, F., Schwalbe, H., Rodnina, M. V., &

Komar, A. A. (2016). Synonymous Codons Direct Cotranslational Folding toward

Different Protein Conformations. *Molecular Cell*, *61*(3), 341–351.

https://doi.org/10.1016/j.molcel.2016.01.008

Bui, M. Q., Dang, C. C., Vinh, L. S., & Lanfear, R. (2021). QMaker: Fast and Accurate Method

to Estimate Empirical Models of Protein Evolution. *Systematic Biology*, *70*(5), 1046–

1060. https://doi.org/10.1093/sysbio/syab010

Cherry, J. L. (2010). Highly expressed and slowly evolving proteins share compositional

properties with thermophilic proteins. *Molecular Biology and Evolution*, *27*(3), 735–741.

https://doi.org/10.1093/molbev/msp270

Costa, I. R., Thompson, J. D., Ortega, J. M., & Prosdocimi, F. (2015). Metazoan Remaining

Genes for Essential Amino Acid Biosynthesis: Sequence Conservation and Evolutionary

Analyses. *Nutrients*, *7*(1), Article 1. https://doi.org/10.3390/nu7010001

Dang, C. C., Minh, B. Q., McShea, H., Masel, J., James, J. E., Vinh, L. S., & Lanfear, R. (2022).

nQMaker: Estimating Time Nonreversible Amino Acid Substitution Models. *Systematic*

Biology, syac007. https://doi.org/10.1093/sysbio/syac007

Dayhoff, M. O. (1972). *Atlas of Protein Sequence and Structure*. National Biomedical Research

Foundation.

Deinum, E. E., Halligan, D. L., Ness, R. W., Zhang, Y.-H., Cong, L., Zhang, J.-X., & Keightley,

P. D. (2015). Recent Evolution in Rattus norvegicus Is Shaped by Declining Effective

Population Size. *Molecular Biology and Evolution*, *32*(10), 2547–2558.

https://doi.org/10.1093/molbev/msv126

Doig, A. J., & Sternberg, M. J. E. (1995). Side-chain conformational entropy in protein folding.

Protein Science, *4*(11), 2247–2251. https://doi.org/10.1002/pro.5560041101

Drummond, D. A., Bloom, J. D., Adami, C., Wilke, C. O., & Arnold, F. H. (2005). Why highly

expressed proteins evolve slowly. *Proceedings of the National Academy of Sciences*,

102(40), 14338–14343. https://doi.org/10.1073/pnas.0504070102

Drummond, D. A., & Wilke, C. O. (2008). Mistranslation-Induced Protein Misfolding as a

Dominant Constraint on Coding-Sequence Evolution. *Cell*, *134*(2), 341–352.

https://doi.org/10.1016/j.cell.2008.05.042

Drummond, D. A., & Wilke, C. O. (2009). The evolutionary consequences of erroneous protein

synthesis. *Nature Reviews Genetics*, *10*(10), Article 10. https://doi.org/10.1038/nrg2662

Eőry, L., Halligan, D. L., & Keightley, P. D. (2010). Distributions of Selectively Constrained

Sites and Deleterious Mutation Rates in the Hominid and Murid Genomes. *Molecular*

Biology and Evolution, *27*(1), 177–192. https://doi.org/10.1093/molbev/msp219

Felsenstein, J. (1985). Phylogenies and the Comparative Method. *The American Naturalist*,

125(1), 1–15. https://doi.org/10.1086/284325

Fontanillas, E., Galzitskaya, O. V., Lecompte, O., Lobanov, M. Y., Tanguy, A., Mary, J.,

Girguis, P. R., Hourdez, S., & Jollivet, D. (2017). Proteome Evolution of Deep-Sea

Hydrothermal Vent Alvinellid Polychaetes Supports the Ancestry of Thermophily and

Subsequent Adaptation to Cold in Some Lineages. *Genome Biology and Evolution*, *9*(2), 279–296. https://doi.org/10.1093/gbe/evw298

Galtier, N., Roux, C., Rousselle, M., Romiguier, J., Figuet, E., Glémin, S., Bierne, N., & Duret, L. (2018). Codon Usage Bias in Animals: Disentangling the Effects of Natural Selection, Effective Population Size, and GC-Biased Gene Conversion. *Molecular Biology and Evolution*, *35*(5), 1092–1103. https://doi.org/10.1093/molbev/msy015

Goldstein, R. A., & Pollock, D. D. (2006). Observations of Amino Acid Gain and Loss during Protein Evolution Are Explained by Statistical Bias. *Molecular Biology and Evolution*, *23*(7), 1444–1449. https://doi.org/10.1093/molbev/msl010

Groussin, M., Boussau, B., & Gouy, M. (2013). A branch-heterogeneous model of protein evolution for efficient inference of ancestral sequences. *Systematic Biology*, *62*(4), 523–538. https://doi.org/10.1093/sysbio/syt016

Haag-Liautard, C., Coffey, N., Houle, D., Lynch, M., Charlesworth, B., & Keightley, P. D. (2008). Direct Estimation of the Mitochondrial DNA Mutation Rate in Drosophila melanogaster. *PLoS Biology*, *6*(8), e204. https://doi.org/10.1371/journal.pbio.0060204

Halligan, D. L., Oliver, F., Eyre-Walker, A., Harr, B., & Keightley, P. D. (2010). Evidence for Pervasive Adaptive Protein Evolution in Wild Mice. *PLOS Genetics*, *6*(1), e1000825. https://doi.org/10.1371/journal.pgen.1000825

Henikoff, S., & Henikoff, J. G. (1992). Amino acid substitution matrices from protein blocks. *Proceedings of the National Academy of Sciences of the United States of America*, *89*(22), 10915–10919. https://doi.org/10.1073/pnas.89.22.10915

Hershberg, R., & Petrov, D. A. (2010). Evidence That Mutation Is Universally Biased towards

AT in Bacteria. *PLoS Genetics*, *6*(9), e1001115.

https://doi.org/10.1371/journal.pgen.1001115

Hunt, S. E., McLaren, W., Gil, L., Thormann, A., Schuilenburg, H., Sheppard, D., Parton, A.,

Armean, I. M., Trevanion, S. J., Flicek, P., & Cunningham, F. (2018). Ensembl variation

resources. *Database*, *2018*, bay119. https://doi.org/10.1093/database/bay119

Hurst, L. D., Feil, E. J., & Rocha, E. P. C. (2006). Causes of trends in amino-acid gain and loss.

Nature, *442*(7105), E11–E12. https://doi.org/10.1038/nature05137

James, J. E., Nelson, P., & Masel, J. (2023). Differential retention of Pfam domains creates long-

term evolutionary trends. *Molecular Biology and Evolution*, *2023*.

https://doi.org/10.1101/2022.10.27.514087

James, J. E., Willis, S. M., Nelson, P. G., Weibel, C., Kosinski, L. J., & Masel, J. (2021).

Universal and taxon-specific trends in protein sequences as a function of age. *eLife*, *10*,

e57347. https://doi.org/10.7554/eLife.57347

Jansen, R., & Gerstein, M. (2000). Analysis of the yeast transcriptome with structural and

functional categories: Characterizing highly expressed proteins. *Nucleic Acids Research*,

28(6), 1481–1488. https://doi.org/10.1093/nar/28.6.1481

Jing, M., Yu, H.-T., Bi, X., Lai, Y.-C., Jiang, W., & Huang, L. (2014). Phylogeography of

Chinese house mice (Mus musculus musculus/castaneus): Distribution, routes of

colonization and geographic regions of hybridization. *Molecular Ecology*, *23*(17), 4387–

4405. https://doi.org/10.1111/mec.12873

Jordan, I. K., Kondrashov, F. A., Adzhubei, I. A., Wolf, Y. I., Koonin, E. V., Kondrashov, A. S.,

& Sunyaev, S. (2005). A universal trend of amino acid gain and loss in protein evolution.

Nature, *433*(7026), 633–638. https://doi.org/10.1038/nature03306

Kaehler, B. D., Yap, V. B., Zhang, R., & Huttley, G. A. (2015). Genetic Distance for a General

Non-Stationary Markov Substitution Process. *Systematic Biology*, *64*(2), 281–293.

https://doi.org/10.1093/sysbio/syu106

Keightley, P. D., Lercher, M. J., & Eyre-Walker, A. (2005). Evidence for Widespread

Degradation of Gene Control Regions in Hominid Genomes. *PLOS Biology*, *3*(2), e42.

https://doi.org/10.1371/journal.pbio.0030042

Kosinski, L., Aviles, N., Gomez, K., & Masel, J. (2022). Random peptides rich in small and

disorder-promoting amino acids are less likely to be harmful. *Genome Biology and*

Evolution, evac085. https://doi.org/10.1093/gbe/evac085

Krogh, A., Larsson, B., von Heijne, G., & Sonnhammer, E. L. L. (2001). Predicting

transmembrane protein topology with a hidden markov model: Application to complete

genomes. *Journal of Molecular Biology*, *305*(3), 567–580.

https://doi.org/10.1006/jmbi.2000.4315

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: A Resource for

Timelines, Timetrees, and Divergence Times. *Molecular Biology and Evolution*, *34*(7),

1812–1819. https://doi.org/10.1093/molbev/msx116

Kwon, M. J., Park, J., Jang, S., Eom, C. Y., & Oh, E. S. (2016). The conserved phenylalanine in

the transmembrane domain enhances heteromeric interactions of syndecans. *Journal of*

Biological Chemistry, *291*(2), 872–881. https://doi.org/10.1074/jbc.M115.685040

Le, S. Q., Dang, C. C., & Gascuel, O. (2012). Modeling Protein Evolution with Several Amino

Acid Replacement Matrices Depending on Site Rates. *Molecular Biology and Evolution*,

29(10), 2921–2936. https://doi.org/10.1093/molbev/mss112

Le, S. Q., & Gascuel, O. (2008). An improved general amino acid replacement matrix.

Molecular Biology and Evolution, *25*(7), 1307–1320.

https://doi.org/10.1093/molbev/msn067

Lecocq, M., Groussin, M., Gouy, M., & Brochier-Armanet, C. (2021). The Molecular

Determinants of Thermoadaptation: Methanococcales as a Case Study. *Molecular*

Biology and Evolution, *38*(5), 1761–1776. https://doi.org/10.1093/molbev/msaa312

Levy, E. D., De, S., & Teichmann, S. A. (2012). Cellular crowding imposes global constraints on

the chemistry and evolution of proteomes. *Proceedings of the National Academy of*

Sciences, *109*(50), 20461–20466. https://doi.org/10.1073/pnas.1209312109

Lins, L., Thomas, A., & Brasseur, R. (2003). Analysis of accessible surface of residues in

proteins. *Protein Science*, *12*(7), 1406–1417. https://doi.org/10.1110/ps.0304803

Lynch, M. (2010). Rate, molecular spectrum, and consequences of human mutation. *Proceedings*

of the National Academy of Sciences, *107*(3), 961–968.

https://doi.org/10.1073/pnas.0912629107

Lynch, M., Ali, F., Lin, T., Wang, Y., Ni, J., & Long, H. (2023). The divergence of mutation

rates and spectra across the Tree of Life. *EMBO Reports*, *24*(10), e57561.

https://doi.org/10.15252/embr.202357561

Lynch, M., & Conery, J. S. (2003). The Origins of Genome Complexity. *Science*, *302*(5649),

1401–1404. https://doi.org/10.1126/science.1089370

Lynch, M., Sung, W., Morris, K., Coffey, N., Landry, C. R., Dopman, E. B., Dickinson, W. J.,

Okamoto, K., Kulkarni, S., Hartl, D. L., & Thomas, W. K. (2008). A genome-wide view

of the spectrum of spontaneous mutations in yeast. *Proceedings of the National Academy*

of Sciences of the United States of America, *105*(27), 9272.

https://doi.org/10.1073/pnas.0803466105

Markin, C. J., Mokhtari, D. A., Sunden, F., Appel, M. J., Akiva, E., Longwell, S. A., Sabatti, C.,

Herschlag, D., & Fordyce, P. M. (2021). Revealing enzyme functional architecture via

high-throughput microfluidic enzyme kinetics. *Science*, *373*(6553), eabf8761.

https://doi.org/10.1126/science.abf8761

McDonald, J. H. (2006). Apparent Trends of Amino Acid Gain and Loss in Protein Evolution

Due to Nearly Neutral Variation. *Molecular Biology and Evolution*, *23*(2), 240–244.

https://doi.org/10.1093/molbev/msj026

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L.,

Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A.

(2021). Pfam: The protein families database in 2021. *Nucleic Acids Research*, *49*(D1),

D412–D419. https://doi.org/10.1093/nar/gkaa913

Monit, C., & Goldstein, R. A. (2018). SubRecon: Ancestral reconstruction of amino acid

substitutions along a branch in a phylogeny. *Bioinformatics*, *34*(13), 2297–2299.

https://doi.org/10.1093/bioinformatics/bty101

Monod, J., & Jacob, F. (1961). Teleonomic mechanisms in cellular metabolism, growth, and

differentiation. *Cold Spring Harbor Symposia on Quantitative Biology*, *26*, 389–401.

https://doi.org/10.1101/sqb.1961.026.01.048

Muñoz-Gómez, S. A., Susko, E., Williamson, K., Eme, L., Slamovits, C. H., Moreira, D., López-

García, P., & Roger, A. J. (2022). Site-and-branch-heterogeneous analyses of an

expanded dataset favour mitochondria as sister to known Alphaproteobacteria. *Nature*

Ecology & Evolution, *6*(3), 253–262. https://doi.org/10.1038/s41559-021-01638-2

Neme, R., Amador, C., Yildirim, B., McConnell, E., & Tautz, D. (2017). Random sequences are

an abundant source of bioactive RNAs or peptides. *Nature Ecology & Evolution*, *1*(6), 1–

7. https://doi.org/10.1038/s41559-017-0127

Ohta, T. (1973). Slightly Deleterious Mutant Substitutions in Evolution. *Nature*, *246*(5428), 96–

98. https://doi.org/10.1038/246096a0

Paten, B., Herrero, J., Beal, K., Fitzgerald, S., & Birney, E. (2008). Enredo and Pecan: Genome-

wide mammalian consistency-based multiple alignment with paralogs. *Genome Research*,

18(11), 1814–1828. https://doi.org/10.1101/gr.076554.108

Perkins, R., & Vaida, V. (2017). Phenylalanine Increases Membrane Permeability. *Journal of the*

American Chemical Society, *139*(41), 14388–14391.

https://doi.org/10.1021/jacs.7b09219

Phifer-Rixey, M., Bonhomme, F., Boursot, P., Churchill, G. A., Piálek, J., Tucker, P. K., &

Nachman, M. W. (2012). Adaptive Evolution and Effective Population Size in Wild

House Mice. *Molecular Biology and Evolution*, *29*(10), 2949–2955.

https://doi.org/10.1093/molbev/mss105

Pinney, M. M., Mokhtari, D. A., Akiva, E., Yabukarski, F., Sanchez, D. M., Liang, R., Doukov,

T., Martinez, T. J., Babbitt, P. C., & Herschlag, D. (2021). Parallel molecular

mechanisms for enzyme temperature adaptation. *Science*, *371*(6533), eaay2784.

https://doi.org/10.1126/science.aay2784

Raiford, D. W., Heizer, E. M., Miller, R. V., Akashi, H., Raymer, M. L., & Krane, D. E. (2008).

Do Amino Acid Biosynthetic Costs Constrain Protein Evolution in Saccharomyces

cerevisiae? *Journal of Molecular Evolution*, *67*(6), 621–630.

https://doi.org/10.1007/s00239-008-9162-9

Rajabi-Maham, H., Orth, A., & Bonhomme, F. (2008). Phylogeography and postglacial

expansion of Mus musculus domesticus inferred from mitochondrial DNA coalescent,

from Iran to Europe. *Molecular Ecology*, *17*(2), 627–641. https://doi.org/10.1111/j.1365-

294X.2007.03601.x

Saunders, R., & Deane, C. M. (2010). Synonymous codon usage influences the local protein

structure observed. *Nucleic Acids Research*, *38*(19), 6719–6728.

https://doi.org/10.1093/nar/gkq495

Schranz, H. W., Yap, V. B., Easteal, S., Knight, R., & Huttley, G. A. (2008). Pathological rate

matrices: From primates to pathogens. *BMC Bioinformatics*, *9*(1), 550.

https://doi.org/10.1186/1471-2105-9-550

Serohijos, A. W. R., Rimas, Z., & Shakhnovich, E. I. (2012). Protein Biophysics Explains Why

Highly Abundant Proteins Evolve Slowly. *Cell Reports*, *2*(2), 249–256.

https://doi.org/10.1016/j.celrep.2012.06.022

Subramanian, S. (2008). Nearly neutrality and the evolution of codon usage bias in eukaryotic

genomes. *Genetics*, *178*(4), 2429–2432. https://doi.org/10.1534/genetics.107.086405

Tenesa, A., Navarro, P., Hayes, B. J., Duffy, D. L., Clarke, G. M., Goddard, M. E., & Visscher,

P. M. (2007). Recent human effective population size estimated from linkage

disequilibrium. *Genome Research*, *17*(4), 520–526. https://doi.org/10.1101/gr.6023607

Theillet, F.-X., Kalmar, L., Tompa, P., Han, K.-H., Selenko, P., Dunker, A. K., Daughdrill, G.

W., & Uversky, V. N. (2013). The alphabet of intrinsic disorder. *Intrinsically Disordered*

Proteins, *1*(1), e24360. https://doi.org/10.4161/idp.24360

Tien, M. Z., Meyer, A. G., Sydykova, D. K., Spielman, S. J., & Wilke, C. O. (2013). Maximum

Allowed Solvent Accessibilites of Residues in Proteins. *PLOS ONE*, *8*(11), e80635.

https://doi.org/10.1371/journal.pone.0080635

Tsai, J., Taylor, R., Chothia, C., & Gerstein, M. (1999). The packing density in proteins:

Standard radii and volumes. *Journal of Molecular Biology*, *290*(1), 253–266.

https://doi.org/10.1006/jmbi.1999.2829

Tsuboyama, K., Dauparas, J., Chen, J., Laine, E., Mohseni Behbahani, Y., Weinstein, J. J.,

Mangan, N. M., Ovchinnikov, S., & Rocklin, G. J. (2023). Mega-scale experimental

analysis of protein folding stability in biology and design. *Nature*, *620*(7973), Article

7973. https://doi.org/10.1038/s41586-023-06328-6

Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level sets

of phylogenies for questions in ecology, evolution, and conservation. *PLOS Biology*,

17(12), e3000494. https://doi.org/10.1371/journal.pbio.3000494

van Noort, V., Bradatsch, B., Arumugam, M., Amlacher, S., Bange, G., Creevey, C., Falk, S.,

Mende, D. R., Sinning, I., Hurt, E., & Bork, P. (2013). Consistent mutational paths

predict eukaryotic thermostability. *BMC Evolutionary Biology*, *13*(1), 7.

https://doi.org/10.1186/1471-2148-13-7

Verbyla, K. L., Yap, V. B., Pahwa, A., Shao, Y., & Huttley, G. A. (2013). The Embedding

Problem for Markov Models of Nucleotide Substitution. *PLOS ONE*, *8*(7), e69187.

https://doi.org/10.1371/journal.pone.0069187

Weibel, C. A., Wheeler, A. L., James, J. E., Willis, S. M., McShea, H., & Masel, J. (2024). *The*

protein domains of vertebrate species in which selection is more effective have greater

intrinsic structural disorder. 2023.03.02.530449.

https://www.biorxiv.org/content/10.1101/2023.03.02.530449v2

Whelan, S., & Goldman, N. (2001). A General Empirical Model of Protein Evolution Derived

from Multiple Protein Families Using a Maximum-Likelihood Approach. *Molecular*

Biology and Evolution, *18*(5), 691–699.

https://doi.org/10.1093/oxfordjournals.molbev.a003851

Wilke, C. O., Bloom, J. D., Drummond, D. A., & Raval, A. (2005). Predicting the Tolerance of

Proteins to Random Amino Acid Substitution. *Biophysical Journal*, *89*(6), 3714–3720.

https://doi.org/10.1529/biophysj.105.062125

Wilson, B. A., Foy, S. G., Neme, R., & Masel, J. (2017). Young genes are highly disordered as

predicted by the preadaptation hypothesis of de novo gene birth. *Nature Ecology &*

Evolution, *1*(6), Article 6. https://doi.org/10.1038/s41559-017-0146

Wu, S., Edwards, S., & Liu, L. (2018). Genome-scale DNA sequence data and the evolutionary

history of placental mammals. *Data in Brief*, *18*, 1972–1975.

https://doi.org/10.1016/j.dib.2018.04.094

Yates, A. D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M. R.,

Armean, I. M., Azov, A. G., Bennett, R., Bhai, J., Billis, K., Boddu, S., Marugán, J. C.,

Cummins, C., Davidson, C., Dodiya, K., Fatima, R., Gall, A., … Flicek, P. (2020).

Ensembl 2020. *Nucleic Acids Research*, *48*(D1), D682–D688.

https://doi.org/10.1093/nar/gkz966

Zarin, T., Strome, B., Peng, G., Pritišanac, I., Forman-Kay, J. D., & Moses, A. M. (2021).

Identifying molecular features that are associated with biological function of intrinsically

disordered protein regions. *eLife*, *10*, e60220. https://doi.org/10.7554/eLife.60220

Zarin, T., Tsai, C. N., Nguyen Ba, A. N., & Moses, A. M. (2017). Selection maintains signaling

function of a highly diverged intrinsically disordered region. *Proceedings of the National*

Academy of Sciences, *114*(8), E1450–E1459. https://doi.org/10.1073/pnas.1614787114

Zuckerkandl, E., Derancourt, J., & Vogel, H. (1971). Mutational trends and random processes in

the evolution of informational macromolecules. *Journal of Molecular Biology*, *59*(3),

473–490. https://doi.org/10.1016/0022-2836(71)90311-1