Ozanimod Attenuates Human Cerebrovascular Endothelial Derived MMP-9 Activity and

Preserves Barrier Properties Following In Vitro Acute Ischemic Injury

Trevor S. Wendt and Rayna J. Gonzales

Department of Basic Medical Sciences, University of Arizona College of Medicine,

Phoenix

Running title: Ozanimod Attenuates MMP-9 Activation in Brain Endothelial Cells

Corresponding Author:

Rayna J. Gonzales, Ph.D.

Department of Basic Medical Sciences

425 N. 5th Street, Building ABC1, Phoenix, AZ 85004-2157.

*E-mail: rjgonzal@email.arizona.edu

Telephone: 602-827-2143, Fax: 602-827-2127.

Abstract

Cerebrovascular endothelial integrity is critical in mitigating a vicious cascade of

secondary injuries following acute ischemic stroke (AIS). Matrix metalloproteinase-9

(MMP-9), a contributor to vascular integrity loss, is elevated during stroke and is

associated with worsened stroke outcome. Here we investigated the FDA approved

selective sphingosine-1-phosphate receptor 1 (S1PR1) ligand, ozanimod, on the

regulation/activity of MMP-9 as well as endothelial barrier components (PECAM-1,

claudin-5, and ZO-1) in human brain microvascular endothelial cells (HBMECs) following

acute hypoxia plus glucose deprivation (HGD; in vitro ischemic injury). We have

previously reported that selective S1PR1 activation improves HBMEC integrity; however,

specific mechanisms underlying S1PR1 involvement in barrier integrity have not been

clearly elucidated. We hypothesized that ozanimod would attenuate an HGD-induced

increase in MMP-9 activity which would concomitantly attenuate the loss of integral barrier

components. Primary male HBMECs were treated with ozanimod (0.5nM) or vehicle

(DMSO <0.1%) and exposed to either normoxia (21% O2) or HGD (1% O2) (3h).

Immunoblotting, zymography, qRT-PCR, H2O2 assay, and immunocytochemical labeling

techniques assessed processes related to MMP-9 and endothelial barrier markers. We

observed that HGD increased intracellular mechanisms associated with augmented

MMP-9 activation (VCAM-1, SOD1, H2O2, CD44, RECK, and TIMP1); ozanimod did not

alter these factors. HGD acutely increased MMP-9 activity and reduced claudin-5 and

PECAM-1 levels, and ozanimod attenuated these responses. In situ analysis via

PROSPER, suggested that attenuation of MMP-9 activity may be a primary factor in

maintaining these integral barrier proteins. Thus, we conclude that ozanimod has the

potential to attenuate an ischemic-like injury mediated decrease in HBMEC integrity in

part by decreasing MMP-9 activity as well as preserving levels of claudin-5 and PECAM-

1.

INTRODUCTION

Matrix metalloprotease-9 (MMP-9) is a key enzyme involved in the complex and multimodal pathological response during acute ischemic stroke (AIS). Upon endothelial activation, increased MMP-9 activity in-part, results in vascular integrity disruption and together can lead to a cascade of secondary injuries including barrier integrity loss, cerebrovascular inflammation, edema, infarct expansion, all contributing to increased hemorrhagic risk and worsened stroke outcomes if the occlusion is not resolved. (1, 2, 3, 4) Maintenance of neuronal survival is dependent upon the proper function of the blood brain barrier (BBB) which at the interface with the central nervous tissue is comprised of endothelial cells, astrocytes, pericytes, neurons, and extracellular matrix (ECM). [Reviewed in (5)]. Of these cells, the endothelium are primary regulators of permeability and if perturbed contributes significantly to the pathophysiological progression of AIS. (6, 7) At the level of the endothelium, a portion of its regulatory role is attributed to specific tight junction proteins and adhesion molecules such as claudin-5 (8), zonula occludens 1 (ZO-1) (9), PECAM-1 (10), and CD44 (11) which comprise critical aspects of barrier function and subsequent regulation of extracellular trafficking. In addition to tight junctions, the endothelium within the neurovascular unit has been shown to secrete ECM which plays an integral role in maintaining vessel and BBB integrity. [Reviewed in (12)] During the progression of AIS, tight junctions (13, 14, 15, 16, 17) and ECM are disrupted (18, 19, 20, 21, 22), which in part can be contributed to the notable acute activation of MMP-9. (23, 24, 25, 26) Moreover, MMP-9 is postulated to be one of the drivers responsible for the biphasic manner in which the BBB is disrupted, where the acute onset phase is classified as detrimental, and the late onset is beneficial in promoting vascular

remodeling. (26, 27, 28, 29) Targeting of MMP-9 activity during the acute phase of BBB

opening may serve as a likely potential therapeutic target to improve functional outcomes

following ischemic injury. (30)

In efforts to mitigate worsened stroke outcomes, the development of synthetic

compounds targeting selective sphingosine-1-phosphate receptors (S1PRs) as a novel

approach for stroke treatment has gained significant traction. (31, 32, 33, 34, 35, 36)

Ozanimod, an FDA approved compound, is currently used to manage multiple sclerosis

(37) and ulcerative colitis (38). This selective S1PR1 ligand has been further explored as

a potential therapeutic agent for stroke patients and has shown beneficial effects on

improving cerebrovascular integrity, neurological function, and reducing infarct volume

following ischemic injury. (39, 40, 41, 42, 43) S1PRs are a family of five G-protein coupled

receptors (S1PR1-5) connected with an array of intracellular second messengers and

signaling pathways. S1PR1 is a Gα_i protein (44) that ubiquitously and differentially

expressed across various tissues, and has been linked to multiple pathways including

PI3K/Akt, (45) Rac activation, (46) and MAPK. (47) Although studies have shown that

S1PR ligands contain lymphocyte modulatory properties, S1PR1 agents can directly elicit

effects at the level of the vasculature. This is supported by the findings that S1PR1 protein

is expressed in vascular smooth muscle (48) as well as endothelial cells. (49) In addition

to expression, a study by Grailhe et al. demonstrated maintenance of electrical

impedance via ozanimod, in a human umbilical vein endothelial cell line. (50) Moreover,

it has been observed that activation of endothelial S1PR1 promotes vascular integrity,

(51, 52, 53, 54) blood pressure homeostasis, (55) cell survival, (47, 56) and maintenance

of microvascular patency and perfusion following ischemic stroke. (41) These functional

outcomes secondary to endothelial S1PR1 activation showcase the potential of targeting

S1PR1 to attenuate infarct expansion, reduce occurrence of hemorrhagic transformation,

and mitigate stroke injury. (41)

We have previously demonstrated that ozanimod attenuates increased brain

endothelial cell barrier permeability via S1PR1 activation following acute exposure to

hypoxia plus glucose deprivation (HGD), an in vitro model of ischemic injury. (40) Thus,

specific mechanisms underlying S1PR1 involvement on barrier integrity merit further

investigation. However, to the best of our knowledge the role of S1PR1 on MMP-9 activity

using a stroke model has not been investigated. Therefore, in this study, we investigated

a potential underlying mechanism and functional outcomes behind the HGD-induced

endothelial cell permeability. Additionally, we addressed the beneficial potential of

ozanimod on attenuating HGD-mediated alterations in paracellular permeability related

structures such as tight junctions, adhesion molecules, and MMP-9 activity as well as

underlying MMP-9 activation processes. We hypothesized that ozanimod would attenuate

an HGD-induced increase in MMP-9 activity which would concomitantly attenuate the loss

of integral human brain endothelial barrier components related to endothelial barrier

permeability.

METHODS

Human Brain Microvascular Endothelial Cell Culture

Vendor-purchased cryopreserved subcultures of primary male human brain

microvascular endothelial cells (HBMECs; Cell Systems Kirkland, WA, Catalogue

number: ACBRI 376 Lot number: 376.02.03.01.2F) were received at *passage 3*, seeded

at a density of 2 to $3 \times 10^5/cm^2$, and cultured as previously described. (57) In brief, they

were grown in 5% CO_2 and room air at 37°C in phenol red-free Complete Classic Medium

(10% FBS; fetal bovine serum) supplemented with Bac-off Antibiotic (Cell Systems,

Catalogue number: 4Z0-643) and CultureBoost (Cell Systems, Catalogue number: 4CB-

500). Cells were cultured in sterile 100mm × 20mm petri dishes (Corning; Catalogue

number: 430591) which were coated with Attachment Factor (Cell Systems; Catalogue

number: 4Z0-201). Once cells reached 75-80% confluency, selected plates were

cryopreserved for future use, and the remaining cells were continued in culture and

studied at *passage 7*. For ICC and other cell imaging experiments, 100mm petri dishes

were split and seeded on sterilized round cover glasses (Fisherbrand Scientific;

Catalogue number: 12-545-100) treated with Attachment Factor within sterile 6-well

plates (Corning; Catalogue number: 3516) and grown to 75-80% confluency.

Drug Treatment and Hypoxia Plus Glucose Deprivation (HGD) Exposure

Ozanimod (RPC1063, 5-[3-[(1S)-2,3-dihydro-1-[(2-hydroxyethyl)amino]-1H-inden-

4-yl]-1,2,4-oxadiazol-5-yl]-2-(1-methylethoxy)-benzonitrile), a selective sphingosine-1-

phosphate receptor 1 (S1PR1) ligand (Cayman Chemical; Catalogue number: 19922),

was prepared fresh under sterile conditions on the same day of administration for planned

experiments as previously described. (48) Briefly, stock solutions (4mM) were made in

dimethyl sulfoxide (DMSO) and further diluted in 1X Dulbecco's phosphate buffer solution

(DPBS) (Corning; Catalogue number: 21-031-CV) to a final working concentration

(0.5nM). Similarly, W146 ([3R-amino-4-[(3-hexylphenyl)amino]-4-oxobutyl]-phosphonic

acid, mono(trifluoroacetate)), a selective S1PR1 antagonist (Cayman Chemical;

Catalogue number: 10009109), was freshly prepared under sterile conditions on the same

day of experiments. Stock solutions (10mM) were made in lab grade methanol and further

diluted in 1X DPBS to a final working concentration (10μM). This working solution and

preincubation period was chosen as it has been reported that W146 at a concentration of

10μM elicits no off-target S1PRs (i.e. S1PR2, S1PR3, and S1PR5) agonist nor antagonist

effects and has been previously utilized and shown to block the effects of a highly

selective S1PR1 ligand. (58) Vehicle solutions utilized in the control groups were

generated in tandem and consisted of a DMSO/DPBS solution as well as a

methanol/DMSO/DPBS solution prepared at a non-toxic concentration (DMSO <0.1%,

methanol <2.0%). These solutions were maintained at 37°C during preparation and

administration to HBMEC cultures. Prior to experimentation, designated plates were

divided into the following groups: normoxia or HGD. To then evaluate the effects of

ozanimod some plates were further designated into the following treatment groups:

normoxia plus vehicle (Normoxia + Vehicle), normoxia plus ozanimod (Normoxia + Ozd),

HGD plus vehicle (HGD + Vehicle), or HGD plus ozanimod (HGD + Ozd). To test whether

observed effects of ozanimod were S1PR1 dependent the following groups were added:

normoxia plus ozanimod plus W146 (Normoxia + Ozd + W146) and HGD plus ozanimod

plus W146 (HGD + Ozd + W146). Prior to HGD exposure, Complete Classic Medium was

replaced with modified DMEM (ThermoFisher Scientific; Catalogue number: A14430-01)

media containing no L-glutamine, phenol red, sodium pyruvate, or D-glucose. Next, cells

were pretreated with ozanimod or vehicle, and in some select plates the antagonist,

W146, was administered 0.5h prior to ozanimod treatment. Following drug treatment,

plates of cells designated for HGD were placed in a hypoxic sub chamber (BioSpherix) at

37°C and flushed with a medical grade gas mixture of 1% O_2, 5% CO_2, and nitrogen

balance and exposed for 0.5-3h depending on the experimental design. Normoxic control

plates containing Complete Classic Medium were replaced with fresh medium and

incubated in a separate CO_2 incubator at 37°C at 21% O_2.

Cellular Protein Isolation and Analysis

The isolation and quantification of protein performed is previously described. (48)

Briefly, culture plates were removed from either the hypoxic BioSpherix chamber or

normoxic incubator, placed on ice, washed twice with ice-cold phosphate buffered

solution (PBS), trypsinized (0.25%) for 3min at 37°C (ThermoFisher Scientific; Catalogue

number: 15090046) which was then neutralized with trypsin neutralizer (ThermoFisher

Scientific; Catalogue number: R002100), and pelleted (centrifuge: Thermo Sorvall

Legend RT+) at 1,900g (10 min, 4°C). The pellet was rinsed with ice-cold PBS,

centrifuged, supernatant discarded, and pellet resuspended in ice-cold lysis buffer

solution (50mM Tris-HCl, 150mM NaCl, 0.1% sodium dodecyl sulfate, 1% Nonidet P-40)

containing protease inhibitors (20μM pepstatin, 20μM leupeptin, 0.1U/mL aprotinin,

0.1mM PMSF, 1mM DTT), phosphatase inhibitors (25mM β-glycerophosphate, 0.1mM

sodium orthovanadate), 1mM EDTA, and 1mM EGTA for 15min. The resulting solution

was then homogenized, sonicated, and centrifuged (Fisher Accuspin Micro 17R) at

17,000g (10 min, 4°C). Following centrifugation, the supernatant (whole cell lysate) was

collected and stored at -80°C. Total protein was determined on the day of western blot

experimentation and determined by using a bicinchoninic acid protein assay kit

(ThermoFisher Scientific; Catalogue number: 23227) according to manufacturer's

protocol and measured on a Safire II (Tecan) plate reader.

Western Blot

Protein levels for MMP-9 as well as for endothelial barrier and activation markers

were examined following normoxic or HGD exposure using standard western blotting

described previously. (48, 59) Briefly, sample lysates were diluted in Tris-glycine SDS

sample buffer with the reducing agent, β-mercaptoethanol (10%) and boiled for 5min. The

equally diluted samples and fluorescent standards (LI-COR Biosciences) were loaded

onto 4–15% gradient Mini-PROTEAN TGX precast protein gels, (Bio-Rad Laboratories;

Catalogue number: 4561084). Protein separation occurred by SDS-PAGE using a mini

PROTEAN Tetra electrophoresis system (Bio-Rad Laboratories). The separated proteins

were then transferred to nitrocellulose membranes using the same apparatus and

membranes were then blocked in 3% nonfat dry milk rehydrated in Tris-Buffered saline

containing 1% Tween (TBST; 20mM Tris, 150mM NaCl) at room temperature for 1h to

alleviate nonspecific binding. Blots were then probed overnight at 4°C with MMP-9 (Cell

Signaling; Catalogue number: 3852S), PECAM-1 (Sigma Aldrich; Catalogue number:

SAB4502167), VCAM-1 (Santa Cruz Biotechnology; Catalogue number: sc-13160),

claudin-5 (ThermoFisher Scientific; Catalogue number: 34-1600), or ZO-1 (ThermoFisher

Scientific; Catalogue number: 33-9100) primary antibodies which were all diluted to 1:500

in Tris-buffered saline with 0.1% Tween 20 detergent (TBST). After TBST wash cycle (5×,

5min), membranes were incubated in goat anti-rabbit IRDye 800CW dye (LI-COR) or goat

anti-mouse IRDye 800CW dye (LI-COR) secondary antibodies at room temperature for

1h. Following another 5×, 5min TBST wash cycle, protein bands were visualized using

an Odyssey Classic infrared imager (LI-COR). Next, membranes were exposed to

overnight at 4°C with 1:15,000 anti-β-actin (Sigma Aldrich; Catalogue number: A5441)

and for 1h at RT goat anti-mouse IRDye 680LT (LI-COR) which served as a loading

control. Band densities were then analyzed using ImageStudio 3.0 software (LI-COR).

Polymerase Chain Reaction (PCR)

Reverse transcription PCR (RT-PCR) was used to determine the presence of

MMP-9, *RECK*, and *TIMP1* mRNA in HBMEC cultures. Briefly, RNA was extracted using

the Qiagen kit (Thermo Fisher, Cat. No. 74034) according to the manufacturer's

instructions. HBMECs were collected and centrifuged (Thermo Sorvall Legend RT +) at

3,000 rpm for 10min at 4°C. The subsequent cell pellets then underwent a series of

washes and centrifugations per the manufacturer's instructions which resulted in

collection of RNA in 14μL of diethyl pyrocarbonate (DEPC). Purity and concentration of

extracted RNA was confirmed using a Nanodrop 2000 (ThermoFisher Scientific). RNA

was reverse transcribed to make cDNA using the SuperScript III First-Strand Synthesis

System (ThermoFisher Scientific; Catalogue number: 18080051) according to the

manufacturer's instructions. Each reaction mixture contained equal concentration loading

of cDNA, as determined by cDNA concentration via Nanodrop 2000, 0.25μL of dNTP

(ThermoFisher Scientific; Catalogue number: 18427088), 1μL 10× Taq Buffer

(ThermoFisher Scientific; Catalogue number: EP0702), 0.25μL of Taq polymerase

(ThermoFisher Scientific; Catalogue number: EP0702), 9.25μL of DEPC, 0.25μL of

forward primer, and 0.25μL of reverse primer. All primers utilized in this study were

designed in house utilizing the NCBI and UCSC genome libraries and obtained from

Integrated DNA Technologies with standard desalting and at 25nM scale. The primer

sequences used to detect *MMP-9* expression were 5'-GTCGTGGTTCCAACTCGGTTTG-

3' (forward) and 5'-GTGGTACTGCACCAGGGCAAG-3' (reverse) and yielded an

expected product size of 131bp. The primer sequences used to detect *RECK* expression

were 5'-GCAGTGCGGGTGCATTGTGTTGTAATC-3' (forward) and 5'-

CAACCATTGTCTCTGGGCAATAATCTGGG-3' (reverse) and yielded an expected

product size of 147bp. The primer sequences used to detect *TIMP1* expression were 5'-

CTTCTGCAATTCCGACCTCGTCATCAGG-3' (forward) and 5'-

CGAACCGGATGTCAGCGGCATC-3' (reverse) and yielded an expected product size of

149bp. The reaction involved an initial melting step at 95°C for 3min followed by 35 cycles

of 95°C (denature) for 30s, primer-specific annealing temperature for 30s, and 72°C

(elongation) for 60s, then a 5min incubation at 72°C to terminate the reaction. Primer

annealing temperatures for *MMP-9*, *RECK*, and *TIMP1* are 64.1°C, 65.6°C, and 67.4°C,

respectively. Following the PCR reaction, samples were loaded alongside a UV standard

(Bioland Scientific; Catalogue number: DM01-01) onto a 1% agarose gel. Gels were then

run in a 1× tris-acetate-EDTA (TAE) (ThermoFisher Scientific; Catalogue number: B49)

solution for 15min at 75V following by 35min at 95V. Gels were finally visualized and

imaged using a ChemiDoc XRS UV gel imager to confirm the presence or absence of the

gene of interest.

Quantitative real time PCR

Quantitative real time PCR (qRT-PCR) was utilized to measure changes in mRNA

levels of *MMP-9*, *RECK*, *TIMP1*, *SOD1*, and *SOD2*. RNA extraction and generation of

cDNA was performed as described in the previous *polymerase chain reaction* section. All

primer efficiencies were assessed via serial dilution from 200ng/μL to 0.02ng/μL and were

found to be within 90-110% efficient. Following cDNA quantification via Nanodrop 2000

(ThermoFisher Scientific), cDNA from each sample was diluted to a working solution of

50ng/μL in DEPC. Duplicate wells were used for each target within the 96-well plate (USA

Scientific; Catalogue number: 1402-9100). Within each well, a mixture of cDNA (5μL),

target primers were diluted in DEPC (5μL; 0.4μM forward primer, 0.4μM reverse primer),

and POWER SYBR Green PCR Master Mix (ThermoFisher Scientific; Catalogue number:

4368706) was added in this order. The 96-well plates were then centrifuged (Thermo

Sorvall Legend RT+) at 1,000g for 2min and then immediately loaded into the

QuantStudio 6 and 7 Flex real-time PCR system (ThermoFisher Scientific). A two-step

cycling protocol was implemented to collect cycle threshold (Ct) values of specific target

amplicons and housekeeping gene *GAPDH* using the ΔΔCt relative quantification setting

within QuantStudio 6. The reaction involved 2min at 50°C, an initial denaturing step at

95°C for 10min followed by 40 cycles of 95°C (denature) for 15s and primer-specific

annealing temperature for 1min. Primer annealing temperatures for *MMP-9*, *RECK*,

TIMP1 SOD1, *SOD2*, and *GAPDH* are 64.1°C, 65.6°C, 67.4°C, 62.4°C, 64.6°C, and

64.4°C respectively. The primer sequences used to detect *MMP-9*, *RECK*, and *TIMP1*

expression are described in the previous *polymerase chain reaction* section. The primer

sequences used to detect *SOD1* were 5'- GGAGATAATACAGCAGGCTGTACCAG-3'

(forward) and 5'- CCACACCATCTTTGTCAGCAGTCAC-3' (reverse). The primer

sequences used to detect *SOD2* expression were 5'-

GTTTTGGGGTATCTGGGCTCCAG-3' (forward) and 5'-

GGTGACGTTCAGGTTGTTCACGTAG-3' (reverse). The primer sequences used to

detect *GAPDH* were 5'-GGTGTGAACCATGAGAAGTATGACAACAGC-3' (forward) and

5'- CCTTCCACGATACCAAAGTTGTCATGG-3' (reverse).

Immunocytochemistry

Immunocytochemical assessment of target proteins was previously described by

our lab(48) and implemented in this study to assess CD44. Briefly, HBMECs at density of

3 to 4 $\times$ 10^5/cm^2 were seeded onto glass coverslips coated with attachment factor (Cell

Systems; Catalogue number: 4Z0-201) within 6-well plates at *passage 7*, grown to 70-

80% confluency, and divided into test groups as described before. Following 3h HGD

exposure / drug treatment, plates were immediately removed from incubator/sub

chamber, placed on ice, media aspirated, and cells washed once with ice-cold DPBS.

HBMECs were then fixed with 4% formaldehyde (10 min, RT) and subsequently washed

twice for 4min with DPBS. HBMECs were then permeabilized with 0.1% TritonX-100

(Catalogue number: T9284) for 10 min and then blocked in 2% BSA (15min, RT). CD44

expression and nuclear localization was determined using anti-CD44 (Cell Signaling;

Catalogue number: 3570S) and PECAM-1 expression using anti-PECAM-1 (Sigma

Aldrich; Catalogue number: SAB4502167) at concentrations 1:100 in 2% BSA. Following

an overnight incubation at 4°C, HBMECs were washed by 2% BSA (3$\times$, 5min) followed

by secondary antibody incubation with either Alexa Fluor 488 goat anti-mouse

(ThermoFisher Scientific; Catalogue number: A11001) or Alexa Fluor 555 goat anti-rabbit

(Thermo Fisher Scientific; Cat. No. A21428) at 1:6,000 in 2% BSA, for 1h at RT protected

from light. A quick wash was then performed with the 2% BSA following by washes with

DPBS (3×, 5min). Two coverslips from each treatment group were incubated in

secondary antibody only and served as negative controls. Next, excess DPBS was

removed, and the coverslips were mounted onto slides utilizing vectashield mounting

medium with DAPI (Vector Laboratories; Catalogue number: H-1200). Coverslips were

then sealed onto slides using nail polish and images captured with an inverted digital

microscope (Keyence; Catalogue number: BZ-X800) following by area and localization

quantification using ImageJ software.

ImageJ Analysis Immunocytochemical Images

FIJI (NIH), an enhanced version of ImageJ2, was utilized to perform analysis on

the area of CD44 and PECAM-1 normalized to the number of HBMECs as well as nuclear

CD44 overlap using images from immunocytochemical experiments. Assessment of

immunocytochemical images via FIJI, again has been previously described and

implemented in this study with slight modifications. (48) Briefly, original

immunocytochemical images obtained from the automated inverted digital microscope

were inputted into FIJI and duplicated. The duplicate images were then split from RGB

into three individual 8-bit images corresponding to the original red, green, and blue

channels. From these 8-bit images, binary images were generated with a threshold

individualized for either CD44, PECAM-1 or DAPI. CD44 and PECAM-1 analyzation of

each treatment group was subjected to the automated IJ_IsoData threshold setting

whereas DAPI analyzation of each treatment group was subjected to the automated

Huang threshold setting. Next, the measurement parameters were set and redirected to

the corresponding channel of the original image and overlaid with the original image

inputted with constraints to not include any objects on the edge of the image. The

subsequent objects generated corresponded to either the labeled CD44, PECAM-1, or

nuclei and from which the total area of CD44 and PECAM-1 signal was quantized as μm^2

and normalized to the number of DAPI stained nuclei present. Additionally, the CD44

constrained binary images were utilized by the JACoP macro-plugin without Costes'

automatic threshold to obtain M1 and M2 coefficients which subsequently quantified the

colocalization of CD44 overlap with the nucleus and the nuclear overlap with CD44

respectively.

Zymography

Media was extracted from cultures following exposure/treatment: $\pm$HGD and

$\pm$ozanimod. The media was briefly centrifuged at 1,000g for 5min to remove any cellular

debris and the supernatant was removed utilized as conditioned media. Total protein

within conditioned media was determined on the day of zymography experimentation and

determined by using a bicinchoninic acid protein assay kit (ThermoFisher Scientific)

according to manufacturer's protocol and measured on a Safire II (Tecan) plate reader.

Conditioned media samples were diluted in Tris-glycine SDS sample buffer and the

equally diluted samples and fluorescent standards (LI-COR Biosciences) were loaded

onto Novex 10% zymogram plus gelatin protein gels (ThermoFisher Scientific; Catalogue

number: ZY00100BOX). Protein separation occurred by SDS-PAGE using a mini–

PROTEAN Tetra electrophoresis system (Bio-Rad Laboratories) at 100V for 2h and held at 4°C. After electrophoresis, the gel was incubated in 1× zymogram renaturing buffer (ThermoFisher Scientific; Catalogue number: LC2670) for 0.5h at RT with gentle agitation. Zymogram renaturing buffer was decanted and replaced with 1× zymogram developing buffer (ThermoFisher Scientific; Catalogue number: LC2671) and allowed to equilibrate the gel for 0.5h at RT with gentle agitation. Following equilibration, developing buffer was decanted and gels washed with nano-pure H_2O and replaced with fresh 1× zymogram developing buffer and developed for 24h at 37°C. Following developing, gels were briefly washed with nano-pure H_2O and stained with Coomassie blue staining solution (2.5g Coomassie Brilliant blue R-250, 450ml 100% methanol, 100ml 100% acetic acid, 400ml nano-pure H_2O) for 2h at RT with gentle agitation. Gels were then washed with nano-pure H_2O and de-stained with Coomassie blue de-staining solution (450ml 100% methanol, 100ml 100% acetic acid, 400ml nano-pure H_2O) for 40min at RT with gentle agitation. The area of gelatin degradation at approximately 82kDa and 63kDa was indicative of MMP-9 and MMP-2 activity respectively and appeared as distinct white bands. The intensities of both MMP-9 and MMP-2 were obtained by an Odyssey Classic infrared imager (LI-COR) and inverse bands densities were then analyzed using ImageStudio 3.0 software (LI-COR).

Extracellular H_2O_2 Quantification

Conditioned media (CM) was obtained as previously described in *zymography* section and utilized to ascertain concentration of hydrogen peroxide (H_2O_2). The aqueous pierce quantitative peroxide assay kit (ThermoFisher Scientific; Catalogue number:

23280) was utilized per manufacturer's instructions to measure levels of H_2O_2 across

treatment groups as previously described: ±HGD, ±ozanimod. In brief, equal loading of

conditioned media was added to 96-well plate followed by $200\mu L$ of reaction mixture

(1/100 v/v of Reagent A to Reagent B) for 20min at RT with agitation. Absorbance was

then measured on a Safire II (Tecan) plate reader at 595nm and colorimetric

quantification of H_2O_2 was compared to standards. Each sample was run in triplicate.

Reagents

All reagents were purchased from Sigma Aldrich (St. Louis, MO) unless otherwise noted.

Data and Statistical Analysis

Western blot and zymography analysis consisted of running lysates from each

treatment group, including vehicles, on the same western blot or zymogram gel for direct

comparison and homogeneity of data. Band densities during western blot analysis were

normalized to loading control β-actin band density and data expressed as fold change

relative to control group (i.e. normoxia or normoxia + vehicle). Experiments were repeated

for statistical analysis and data graphed using Prism 8 (GraphPad Software).

Confirmation of normal distribution was achieved with a Shapiro-Wilk test. *F*-test was

performed to confirm if data sets achieved equal variance among groups. For analysis of

datasets that achieved a normal distribution, direct comparisons between two groups

were made using an unpaired *t* test alone if equal variance was achieved or with Welch's

correction if equal variance was not achieved. Comparisons between three or more

groups were made using either one-way or two-way ANOVA with Tukey's multiple

comparisons post hoc test. Datasets that did not achieve a normal distribution would have

undergone direct comparisons via Mann-Whitney test. $P < 0.05$ was considered

significant. Values are expressed as means $\pm$ SEM.

RESULTS

HGD increased MMP-9 transcription which was not altered by ozanimod

In human epithelial cells exposed to chemical hypoxia conditions via $CoCl_2$, *MMP-*

9 mRNA levels were transiently elevated beginning at 2h and decreased at 6h. (60)

However, the expression of *MMP-9* mRNA in HBMECs following acute ischemic injury

remains unclear. Therefore, we hypothesized that acute HGD exposure would upregulate

MMP-9 mRNA expression and this increase would be attenuated by ozanimod. **Figure**

1A illustrates a representative image of a 2% agarose gel demonstrating confirmation of

MMP-9 gene expression within the HBMEC culture under both normoxic control

conditions and following exposure to 3h of HGD. We observed that under normoxic

conditions there was no detectable *MMP-9* gene expression following a 35 replication

PCR cycle, but HBMECs exposed to 3h of HGD contained a detectable amplicon. These

results were then substantiated following qRT-PCR analysis of the *MMP-9* gene graphed

in **Figure 1B**. Again, there was no detectable mRNA expression of *MMP-9* with a 40

replication PCR cycle under normoxic conditions, but there was a significant increase in

MMP-9 mRNA following HGD exposure for 3h. Interestingly, HBMECs treated with

ozanimod and exposed to HGD experienced a greater variability in *MMP-9* mRNA

expression compared to HGD alone, but there was no difference between the two groups,

thus suggesting that S1PR1 modulation does not alter acute HGD-mediated increases in

MMP-9 at the transcriptional level.

Intracellular pro-MMP-9 protein expression was decreased following HGD; ozd did

not alter levels of expression

Transcription and subsequent translation of MMP-9 leads to formation of a non-

enzymatically active pro-MMP-9 which has a molecular weight of 92kDa. Upon activation,

the pro-domain is removed resulting in a structure with a molecular weight of

approximately 82-84kDa. (61) In our next set of experiments we determined whether HGD

$\pm$ozanimod alters the intracellular MMP-9 protein levels of in HBMECs. **Figure 2A**

illustrates a representative western blot using anti-MMP-9 to detect pro- and active forms

of MMP-9 from HBMEC lysate. Densiometric analysis of intracellular MMP-9 revealed

that HGD exposure induced a decrease in pro-MMP-9 (92kDa) when compared to the

normoxic control, and ozanimod had no effect on this response (**Figure 2B**). In contrast,

the enzymatically active form of MMP-9 (84kDa) following HGD exposure was not

different compared to the normoxic control; however, there was a discernable increase in

the variability of the protein levels when compared to the pro-MMP-9 expression following

HGD exposure (**Figure 2C**). The impact of HGD in the conversion of pro-MMP-9 to MMP-

9 was assessed via the individual ratio expression of pro vs. active for each sample.

These data analysis demonstrated that HGD increased the ratio of active to inactive

MMP-9 protein levels compared to normoxic controls; however, ozanimod did not

attenuate this response (**Figure 2D**). These data viewed through the lens of previous data

shown in **Figure 1** suggests that 3h HGD exposure may be upregulating the synthesis

and then cleavage of MMP-9 which then becomes increasingly exported out of the

HBMECs into the extracellular space.

Ozanimod attenuated HGD-induced increases in extracellular MMP-9 activity

To address whether HGD increased HBMEC derived MMP-9 secretion and

subsequent activity, we next assessed extracellular MMP-9 activity via zymography from

conditioned media collected from HBMECs exposed to either normoxia or HGD for 3h in

the presence or absence of ozanimod. A representative zymography shown in **Figure 3A**

demonstrates the band migration and enzymatic activity of both MMP-9 and MMP-2 from

conditioned media of HBMECs. Inverse densiometric analysis of these bands are

graphically represented in **Figure 3B** and **Figure 3C** respectively. MMP-9 activity was

increased following HGD exposure and ozanimod reversed this response (**Figure 3B**). In

contrast, there was no difference in MMP-2 activity following HGD when compared to the

normoxic control (**Figure 3C**). The data shown in **Figure 3** demonstrate that 3h HGD

exposure increased HBMEC derived MMP-9 activity and that ozanimod treatment

attenuated this response. This suggests that although ozanimod had no effect at

preventing synthesis of MMP-9 at the level of intracellular HBMEC mRNA and protein,

the drug was effective at attenuating extracellular activity of the enzyme (**Figure 3D**). The

next step was to address how HGD exposure and treatment with ozanimod were

achieving these differential effects at an acute time point.

HGD decreased PECAM-1 levels which was attenuated by ozanimod

PECAM-1, also known as CD31, has been shown to be cleaved by MMP-9 within

the hepatic endothelium following ischemic injury. (16) Therefore, to validate the HGD-

mediated increase in MMP-9 activity, platelet endothelial cell adhesion molecule 1

(PECAM-1) levels were assessed. Immunofluorescent PECAM-1 labeling following 3h

HGD decreased and ozanimod attenuated this response (**Figure 4A**). Total area of anti-

PECAM-1 signal normalized to the number of nuclei demonstrated that there was a

decrease in labeled PECAM-1 protein following 3h HGD exposure and ozanimod

attenuated this response (**Figure 4B**). Similarly, when examining PECAM-1 via western

blot and under all treatment/exposure conditions we observed three distinct PECAM-1

bands that appeared to follow a similar trend of expression levels as the

immunocytochemical detection of PECAM-1 (**Figure 4C**). PECAM-1 has been described

as forming homophilic and heterophilic interactions which are both implicated in the

regulation of endothelial barrier integrity (62, 63, 64), thus suggesting that the distinct

PECAM-1/PECAM-1 bands detected at 240-260kDa are PECAM-1 dimers while the

detectable bands at 130kDa and 95kDa represent potential alternative splice forms of

PECAM-1. (65) The predicted molecular weight of PECAM-1 is approximately 80kDa

which corresponds closest to the most prominent form of PECAM-1 detected in HBMECs,

but the fully processed form corresponds to a molecular mass of 130kDa. (65) Further

investigation into the regulation of potential alternative splicing or post-translational

modification of PECAM-1 exceeds the scope of this study but opens the door for future

studies to delineate mechanisms behind this observed alternative splicing. These data

when viewed through the lens of MMP-9 activity suggest that ozanimod may be sheltering

PECAM-1 from MMP-9 mediated degradation and supports our previous observations

regarding MMP-9 activity in **Figure 3**. Together suggesting that ozanimod may be

attenuating HGD-induced endothelial barrier disruption at the level of PECAM-1.

Extracellular HBMEC derived H_2O_2 concentrations were increased following HGD

exposure

To address how acute HGD exposure increased MMP-9 activity, we next

examined levels hydrogen peroxide (H_2O_2) which is exported extracellularly (66, 67) and

been shown to activate human endothelial MMP-9 after 2h. (68) We concomitantly

choose to assess levels of vascular cell adhesion molecule 1 (VCAM-1) as it has been

previously found that VCAM-1, through an H_2O_2 mechanism, leads to endothelial MMP-

9 activation. (69) HGD induced an increase in VCAM-1 levels in HBMECs compared to

the normoxic controls; and ozanimod did not alter this response (**Figure 5A**). We next

examined mRNA levels of superoxide dismutase 1 (*SOD1*) and 2 (*SOD2*) in HBMECs

exposed to normoxia or HGD and treated with ozanimod or vehicle for 3h. HGD exposure

induced a significant increase in *SOD1* mRNA and ozanimod did not attenuate this

response (**Figure 5B**). This contrasts with the assessment of *SOD2* mRNA levels, which

revealed no difference across any treatment group (**Figure 5C**). We next examined the

concentration of extracellular H_2O_2 within the conditioned media extracted from HBMEC

cultures exposed to either normoxia or HGD and treated with either ozanimod or vehicle.

There was an observed temporal dependent and HGD mediated increase in HBMEC

derived extracellular H_2O_2 that peaked at 3h when compared to normoxic controls (**Figure**

5D). **Figure 5D** graphically illustrates extracellular H_2O_2 concentrations from conditioned

media at time points 1 and 3h and demonstrates no difference in extracellular H_2O_2 at 1h

but shows that HGD exposure significantly increased detectable H_2O_2 at 3h both

compared to the normoxic control at the same 3h time point as well as to the 1h HGD

plus vehicle time point. These findings corroborate our previous findings of increased

VCAM-1 protein and *SOD1* mRNA following 3h HGD exposure. Moreover, just as before

we did not observe any effect of either ozanimod in either the normoxic or HGD exposure

groups at any time point. There was also no discernible difference in extracellular H_2O_2

at 0.5h (data not shown) suggesting that H_2O_2 may be involved with an increase in MMP-

9 activity. These data together suggest that the temporally dependent HGD mediated

increases in VCAM-1 protein and *SOD1* mRNA may in part lead to an increase in

extracellular H_2O_2 concentrations. Moreover, these elevated extracellular H_2O_2

concentrations may partially be responsible for the previously observed acute HGD-

mediated increase in MMP-9 activity however does not address the mechanism as to how

ozanimod attenuated this response.

HGD decreased inhibitory mechanisms of MMP-9 export and activation

Although we observed that HGD acutely increased a positive mechanism of MMP-

9 activation, we did not observe an attenuative effect of ozanimod on this response

therefore we next addressed potential mechanisms leading to a decrease in MMP-9

activity. In human lung cancer cells, membrane anchored protein, reversion-inducing

cysteine-rich protein with Kazal motifs (RECK), negatively regulates MMP-9 secretion and

activity that culminated in an attenuation of tumor migration (70, 71) suggesting that its

upregulation during acute ischemic injury in the cerebrovasculature is indicative of an

increased probability of hemorrhagic transformation and increased negative outcomes.

Additionally, tissue inhibitors of metalloproteinases 1 (TIMP1) which is well-established

as capable of inhibiting almost all known MMPs with varying degrees of efficacy, has been

shown more to associate with MMP-9. (72) Therefore, in this study we hypothesized that

HGD exposure would decrease expression of these markers which would be attenuated

by treatment with ozanimod. A representative PCR gel confirmed that HBMEC *RECK*

mRNA expression under both normoxic and HGD conditions, and at a 35-replication cycle

limit it was visually observed that HGD decreased *RECK* mRNA expression (**Figure 6A**).

This was also confirmed using qRT-PCR measurement which is graphically shown and

demonstrates a significant decrease in *RECK* mRNA following 3h HGD exposure (**Figure**

6A). Ozanimod did not attenuate this effect (**Figure 6A**). Similarly, *TIMP1* mRNA

expression was visualized on an agarose gel that confirmed its expression in HBMECs

following both normoxic and HGD exposure for 3h. Moreover, qRT-PCR analysis of

TIMP1 revealed a significant decrease in HBMECs exposed to HGD, which again was

not attenuated by ozanimod (**Figure 6B**). These data suggest that in addition to

increasing a potential mechanism of activation (**Figure 5**), acute HGD exposure

decreased mechanisms related to inhibition of HBMEC MMP-9 activity; however, the

previously observed functional effects of ozanimod on MMP-9 activity remains to be

answered.

HGD exposure decreased HBMEC CD44 levels

To next address both a functional component integral to MMP-9 activity as well as

to endothelial barrier function, we investigated the levels and distribution of CD44. The

multimodal cell surface protein, CD44, has been implicated in many cellular roles such as

endothelial barrier function, (11) proliferation and survival, (73) and as an anchor point for

MMP-9. (17, 74) In this study we utilized an immunocytochemical approach to ascertain

the effects of HGD exposure as well as treatment with ozanimod on the levels and nuclear

overlap of CD44. **Figure 7A** demonstrates representative immunocytochemical images

of HBMECs with anti-CD44 (green) and DAPI stained nuclei (blue) following normoxia or

HGD exposure +/- ozanimod. Total area of anti-CD44 signal was measured and

normalized to the number of nuclei. There was a significant decrease in labeled CD44

protein following HGD when compared to the normoxic control and this response was not

attenuated by ozanimod (**Figure 7B**). Intriguingly however, within the normoxic control

groups we observed a noticeable impact of ozanimod on the distribution of CD44 with

regards to the nucleus. Quantification of the overlap of CD44 with the nucleus determined

with Mander's Coefficient via the JACoP macro-plugin within FIJI showed no difference

between HBMECs exposed to normoxia and HGD, but that treatment with ozanimod

under normoxic conditions significantly reduced the overlap of CD44 with the nucleus

(**Figure 7C**). There were no significant differences in CD44 overlap with the nucleus in

HBMECs treated with ozanimod under HGD conditions. As part of Mander's coefficient

we also obtained quantification of the inverse, which described the overlap of the nuclei

with CD44 (**Figure 7D**). Again, this analysis revealed no difference between HBMECs

exposed to normoxia and HGD for 3h but demonstrated that ozanimod exhibited nearly

identical effects with regards to overlap attenuation of the nuclei with CD44 under

normoxic conditions. The observed decreased levels of HBMEC CD44 following HGD

exposure suggests a decrease in not only CD44 mediated barrier integrity, survival, and

proliferation but also that secreted MMP-9 may be losing an anchor point from which to

perform its enzymatic activity and thusly might be free to expand beyond the confines of

the endothelial surface to perform degradation. Additionally, the observed decrease in

CD44 overlap with the nucleus following S1PR1 activation with ozanimod suggests

reduced translocation of the intracellular domain of CD44 to the nucleus, which has been

shown to induce transcriptional effects during cellular stress. (75, 76, 77) Since we

observed that ozanimod did not attenuate HGD mediated decreases in CD44 but did

previously reduce MMP-9 activity and preserved PECAM-1 expression, we hypothesized

that untethered MMP-9 would negatively impact other contributors of endothelial barrier

function and ozanimod would attenuate this response.

Ozd differentially attenuated HGD-mediated decreases in barrier proteins ZO-1 and

Claudin-5

Next, to test our hypothesis that the observed increased MMP-9 activity could

functionally be involved in endothelial barrier dysfunction, we assessed protein levels of

tight junction protein, claudin-5, and the intracellular anchoring protein integral for barrier

function, zonula occludens 1 (ZO-1). We chose to assess these markers as both claudin-

5 and ZO-1 have been shown to be decreased acutely in murine brain endothelial cells

following exposure to exogenous MMP-9. (13) **Figure 8A** demonstrates that HGD

exposure induced a decrease in claudin-5 protein levels and ozanimod attenuated this

response. Thus, suggesting that in addition to preserving PECAM-1 levels, ozanimod may

be capable of preserving endothelial barrier integrity by reducing the loss of claudin-5

following an acute ischemic-like injury. Finally, immunoblotting of ZO-1 protein showed

that HGD exposure significantly reduced the detectable levels of ZO-1 in HBMECs

compared to the normoxic control; however, ozanimod treatment did not exert any effects

on this response (**Figure 8C**). These data taken together with our previous observations

that ozanimod attenuated an HGD-mediated reduction in trans-endothelial electrical

resistance, (40) suggest that ozanimod, a selective S1PR1 ligand, may be functionally

attenuating a loss in endothelial barrier integrity in part by attenuating extracellular MMP-

9 activity and reducing the loss of PECAM-1 and claudin-5 proteins.

DISCUSSION:

In this study using an in vitro ischemic-like stroke model we have identified a

potential novel mechanism by which ozanimod, a selective S1PR1 ligand, attenuates

HGD-induced disruptions in human brain endothelial barrier function. We have previously

shown that our HGD model decreased HBMEC barrier integrity and S1PR1 activation

attenuated this response. (40) In this study we investigated a mechanism behind this

observation and our results demonstrate that HGD increased MMP-9 activation along

with altered endothelial barrier proteins and that S1PR1 may play a beneficial role in

attenuating this HGD induced response. In brief, HGD exposure 1) induced MMP-9

transcription, 2) increased extracellular MMP-9 enzymatic activity, 3) enhanced levels of

VCAM-1 and extracellular H_2O_2, 4) altered levels of tight junction, activation, and

regulatory/anchoring proteins, and 5) ozanimod differentially attenuated these responses.

To the best of our knowledge this is the first study to investigate the effect of a selective

S1PR1 compound following ischemic-like conditions on MMP-9 regulation and

endothelial barrier properties in a human brain endothelial cell model.

MMP-9 mRNA and Protein Regulation

Maintenance of endothelial barrier integrity both during physiological conditions

and following an ischemic event is important in the regulation of microvessel stability (78),

blood flow (79), and ultimately blood brain barrier integrity. (80, 81) Enzymatic activity of

MMP-9 is a key mediator of functional endothelial barrier disruption and serves clinically

as a marker of ischemic stroke. (82, 83) Thus attenuation of MMP-9 activity remains an

attractive therapeutic aim in the management of AIS. As such, studies implementing

genetic regulation of MMP-9 demonstrate reduced lesion volume and infarct expansion

following focal ischemia. (84, 85, 86) However, additional independent studies have

suggested that while attenuation of MMP-9 during the acute phase of ischemic stroke is

beneficial, this enzyme plays a key role in the later remodeling phase responsible for

angiogenesis and subsequently reperfusion of the damaged area. (26, 87) These insights

suggest that while genetic ablation of *MMP-9* may prove beneficial acutely, the long-term

effect of such approaches may prove untenable. In vascular smooth muscle cells *MMP-*

9 is upregulated via NFκB activation (88), which could potentially be mediating the drastic

increase in *MMP-9* mRNA following HGD exposure for 3h observed in this study (**Figure**

1B). Moreover, although levels of *MMP-9* mRNA were not changed with ozanimod during

HGD exposure, we observed a noticeable increase in variability between samples. This

could thusly be attributed to intercellular heterogeneity within the individual samples and

between different cultures and potentially be due to differential regulation of NFκB. This

postulation is supported by a study conducted in human pulmonary endothelial cells

which showed that overexpression of S1PR1 decreased p65 phosphorylation and

translocation to the nucleus. (89) Therefore, we conclude that 3h HGD exposure

significantly upregulates *MMP-9* mRNA levels which sets the stage for a potential increase in protein production.

The process of MMP-9 translation is tightly regulated and has previously been shown to be regulated by phosphorylation of eukaryotic initiation factor 4E (90) which when pharmacologically attenuated resulted in reduced *MMP-9* translation. (91) In this study we observed that HGD induced a significant increase in levels of *MMP-9* mRNA (**Figure 1**), as well as a decrease in the levels of pro-MMP-9 (**Figure 2B**). From these findings we hypothesized that either the translation of *MMP-9* mRNA to pro-MMP-9 was attenuated or that the inactive pro-MMP-9 protein was being increasingly converted to the enzymatically active MMP-9 form. We observed an increase in the ratio of the active MMP-9 to inactive pro-MMP-9 protein levels (**Figure 2D**) following HGD exposure which further confounded the potential variables at play and suggests that HGD exposure either induced 1) a decrease in pro-MMP-9 synthesis or 2) a conversion rate of pro-MMP-9 to MMP-9 that balances the rate by which MMP-9 is being transported extracellularly to ultimately match the levels of endogenously expressed MMP-9 protein under normoxic conditions. In support of the second potential conclusion, it has been previously reported that HUVECs exposed to angiogenic factors experienced increased rapid vesicle formation and secretion that contain the proteolytically active enzyme, MMP-9, which appears to maximize in the extracellular space at 3h. (92) Additionally, we have observed a significant increase in HBMEC vesicle formations following exposure to 3h of HGD compared to normoxic controls (Wendt and Gonzales, unpublished observation). To further address the questions posed by these findings of whether translation of *MMP-9* mRNA was being attenuated or if MMP-9 was being exported extracellularly following

HGD exposure, as well as the role of ozanimod, we assessed extracellular MMP-9.

Utilizing zymography, we observed increased extracellular MMP-9 activity following HGD

exposure and that ozanimod attenuated this response (**Figure 3B**). Although we

observed an increase in endothelial derived MMP-9 activity extracellularly, we detected

no difference in MMP-2 activity (**Figure 3C**). These data together thusly suggest that 3h

ischemia-like injury induces an increase in HBMEC MMP-9 transcription, protein

exportation, and extracellular activity and treatment with ozanimod attenuates the

functional extracellular enzymatic activity of MMP-9 (**Figure 3D**).

MMP-9 Activation

In efforts to identify a potential mechanism by which 3h HGD exposure induced

MMP-9 activity, we first measured VCAM-1 as it is a well-accepted marker of endothelial

cell activation and has been implicated upstream of MMP-9 activation through production

of H_2O_2. (69, 93, 94) Clinically, elevated intracranial levels of VCAM-1 correlated with

increased infarct and edema volumes in stroke patients that underwent mechanical

thrombectomies. (95) Moreover, Yousef et al demonstrated that both anti-VCAM-1

antibodies and genetic ablation of brain endothelial cell VCAM-1 reduced age-related

neurodegeneration in part through reduced microglial reactivity and cognitive defects.

(96) These studies highlight the significance of VCAM-1 in mediating both MMP-9

activation as well as mediating negative neurological outcomes. In our study we observed

a significant increase in VCAM-1 protein levels following acute HGD exposure; however,

there was no effect of ozanimod (**Figure 5A**). Previous work has demonstrated that

translocation of active NFκB to the nucleus plays a critical role in the rapid induction of

VCAM-1 in response to endothelial activation as well as inflammation. (97) This in part

supports the previous hypothesis that HGD exposure induced an increase in NFκB

activation. In addition to VCAM-1, we assessed mRNA levels of *SOD1* and *SOD2*

following HGD exposure in the presence or absence of ozanimod. Both enzymes are well

known and accepted for the ability to convert superoxide (O_2^-) to H_2O_2 [Reviewed in (98)]

and in this study were hypothesized to contribute to the activation of MMP-9.

Unexpectedly and counter to its consideration as a "housekeeping gene", we observed a

significant increase in mRNA levels of *SOD1* following HGD (**Figure 5B**). Although, *SOD1*

shares significant overlap in transcription factors associated with regulation of both *SOD2*

and *SOD3* [Reviewed in (99)], we observed no increase in *SOD2* mRNA levels following

HGD (**Figure 5C**). Further work is needed to identify a potential transcription factor

responsible for the upregulation of *SOD1* but not *SOD2* mRNA following acute HGD

exposure in HBMECs. Finally, and potentially as a consequence of both upregulated

VCAM-1 protein and *SOD1* mRNA expression, we observed a significant increase in

extracellular H_2O_2 following 3h HGD exposure (**Figure 5D**). Similarly, this increase in

H_2O_2 concentration was not changed with ozanimod treatment which mirrors our findings

at both the level of VCAM-1 and *SOD1*. Moreover, it is well known and accepted that

H_2O_2 due to its small non-polar nature and relatively long half-life can diffuse across

biological membranes serves a pleotropic role in cellular function as a second messenger

(100) as well as an activator of MMP-9. These findings partially reveal how HGD exposure

might be mediating an increase in MMP-9 activity but does not sufficiently elucidate the

attenuative role of ozanimod.

Inhibition of MMP-9

In addition to investigating a potential mechanism underlying MMP-9 activation, we interrogated inhibitory mechanisms of MMP-9 in hopes of identifying a means by which ozanimod exerts its attenuative effects. We demonstrated that mRNA levels of *RECK* and *TIMP1* decreased following HGD; however, ozanimod did not alter their expression (**Figure 6**). These endpoints were chosen to supplement the positive regulatory mechanisms identified in **Figure 5** demonstrating that in addition to increased H_2O_2 production, HGD mediated a downregulation in negative regulatory mechanisms associated with MMP-9. (70, 101, 102, 103, 104, 105) A previous study illustrated that RECK protein expression supported neuroprotection and reduced cell death in a mouse model of ischemic stroke (106) however, its neuro-focused viewpoint does not shed light on the impact of the cerebrovasculature. To this end and in support of our findings, an additional in vitro study conducted with human embryonic kidney epithelial cells demonstrated that beginning at 4h of hypoxic conditions, *RECK* mRNA levels were decreased compared to normoxic controls. (107) While RECK has been implicated in the structural inhibition and secretory regulation of MMP-9 (71), it is generally accepted that TIMP1 forms a complex with MMP-9 which prevents its enzymatic activity. (72) In contrast to our findings in **Figure 6B**, a study conducted in a murine model of 0.5h focal cerebral ischemia followed by reperfusion at various times found that the relative preischemic hemisphere concentration of TIMP1 was extremely low and was gradually upregulated beginning at 3h post-reperfusion. (108) However, in the same study the investigators demonstrated an increase in Evan's Blue extravasation in *TIMP1* ablated mice at 24h post-reperfusion, thus demonstrating its necessity in maintaining BBB integrity. To the best of our knowledge this is the first report to demonstrate a baseline level of *TIMP1*

mRNA under normoxic conditions that is acutely decreased following HGD exposure in

primary HBMECs; however further investigation is required to fully elucidate the

mechanism behind this observation. These data indicate that in addition to increased

positive regulation of MMP-9, HGD induces a decrease in the negative regulation of the

proteolytic enzyme; however, the regulatory mechanism behind ozanimod remains

unclear.

MMP-9 Membrane Docking

As previously mentioned, the transmembrane glycoprotein CD44 exhibits many

different cellular functions; however, in this study we viewed the expression of CD44

through the lens of MMP-9 and endothelial barrier integrity regulation. The relationship

between CD44 and MMP-9 was firmly established by a seminal study which illustrated

that docking of proteolytic active MMP-9 to surface CD44 enhanced tumor invasion and

growth as well as angiogenesis through a TGF-β mechanism. (109) This gave rise to a

multitude of additional studies which have primarily focused on the role of CD44 in tumor

microenvironments [Reviewed in (110)] which more recently has shifted to be in the

vascular spotlight. [Reviewed in (111)] Counter to our observations of decreased CD44

labeling following 3h HGD exposure (**Figure 7B**), a study in renal microvascular

endothelial cells demonstrated an exponential increase in CD44 expression at 1, 3, 7,

and 14 days following a 45 min ischemia reperfusion injury induced by bilateral renal

artery ligation. (112) Similarly, a model of permanent middle cerebral artery occlusion

performed in rats found that both mRNA and protein levels of CD44 in the ipsilateral

hemisphere of injury increased beginning at 6h post injury and was maximized at 24h.

(113) Additionally, these findings were recently functionally substantiated in a study with

iPSC-derived brain microvascular endothelial cells which demonstrated that prolonged

oxygen-glucose deprivation exposure for 24h increased CD44 expression and that

binding of hyaluronan induced a partial maintenance of endothelial barrier function

compared to oxygen-glucose deprivation alone. (114) While these studies do not coincide

with our findings here in this study, to the best of our knowledge this is the first report to

characterize the acute expression of CD44 in primary HBMECs following HGD exposure.

These data could suggest that CD44 undergoes a biphasic response in which expression

is acutely decreased and gradually increases following ischemic-like injury.

In addition to measuring levels of labeled CD44, the immunocytochemical staining

of this protein sheds light on its nuclear and non-nuclear overlap. Due to the

transmembrane nature of CD44, this protein is described as having extracellular,

transmembrane, and intracellular domains. [Reviewed in (75)] Soluble release of CD44

has been well accepted as a consequence of proteolytic cleavage of the extracellular

domain; however, further investigation into the regulation of CD44 has revealed that the

intracellular domain of CD44 is able to translocate to the nucleus and act as a transcription

factor following cleavage by membrane type 1-matrix metalloproteinase. (76, 115)

Moreover, in a separate study in human umbilical vein endothelial cells, induction of

endothelial migration and morphogenic differentiation via type 1-matrix metalloproteinase

was shown to be in part dependent on Gi proteins downstream of S1PR1 following

sphingosine-1-phosphate stimulation. (116) Suggesting that mechanisms downstream of

S1PR1 may also be consequently involved with the regulation of nuclear CD44

expression via membrane type 1-matrix metalloproteinase. The observations made in this

study indicate that ozanimod may be attenuating CD44 expression in the nucleus of

HBMECs; however, these observations were abolished following 3h HGD exposure

(**Figure 7C**). Thus, suggesting that HGD exposure may be differentially regulating various

intracellular signaling pathways that potentially negate the effects of ozanimod stimulated

S1PR1 signaling with relation to CD44 cleavage which merits further investigation.

Endothelial Barrier: PECAM-1

In a study by Kato et al (16), it was determined that MMP-9 induced cleavage of

PECAM-1 in murine hepatic endothelium following ischemia reperfusion injury. These

findings were expanded upon a year later which showcased that, deficiencies in MMP-2

expression resulted in upregulated MMP-9 activity and increased transmigration of

leukocytes, liver damage, and decreased survival following hepatic ischemic reperfusion

injury. (117) Intriguingly, these authors found a significant colocalization of MMP-2 with

PECAM-1 in naïve livers which suggests a potential regulatory role of MMP-2 on the

stability of PECAM-1 that is disrupted by MMP-9. These studies complement our data

which revealed a significant decrease in PECAM-1 staining following 3h HGD exposure

which was then attenuated by ozanimod (**Figures 4A & B**), mirroring the activity of MMP-

9. Although our data support the hypothesis of an MMP-9 dependent degradation of

PECAM-1 following ischemia reperfusion injury, another study conducted in primary rat

retinal endothelial cells under hyperglycemic conditions found that MMP-2 mediated

cleavage of PECAM-1. (118) While these interpretations contradict the hypothesis of

MMP-9 induced cleavage of PECAM-1, the authors found no difference in plasma MMP-

9 activity under diabetic conditions, but rather a significant increase MMP-2 activity.

Moreover, the diabetic conditions of the study, in which glucose is oversupplied, is almost

opposite in nature to the ischemic-like injury model utilized in this study, making a direct

comparison between the two outcomes difficult to justify. Additionally, although the

authors found multiple cleavage sites on rat PECAM-1 suitable for MMP-2, we utilized the

same in silico analysis via PROSPER (119) in this study to demonstrate that there are

significantly more sites on human PECAM-1 that can be cleaved by MMP-9 compared to

MMP-2 and MMP-3 (**Supplemental Figure 1**). Due to the overlapping substrate material

degraded by both MMP-9 and MMP-2 it is possible that human PECAM-1 is degraded by

MMP-2; however, in this study we did not observe an increase in MMP-2 activity and

surmise that MMP-9 is cleaving PECAM-1 and ozanimod may be attenuating this

response via blockade of MMP-9 activity.

In addition to being a marker of MMP-9 activity in this study, we utilized PECAM-1

levels as an indicator of endothelial barrier function. It has been previously observed that

the lack of PECAM-1 in primary mouse brain microvascular endothelial cells resulted in

an impaired endothelial barrier as measured by trans-endothelial electrical resistance

(TEER). (10) Moreover, gerbil brain microvascular endothelial cells exposed to 6h of

oxygen glucose deprivation and 12h of reperfusion, showed reduced levels of PECAM-1

as measured by both immunocytochemical and western blot analyses. (120) These

previous findings align with the observations in our study which demonstrate that 3h HGD

exposure reduced levels of PECAM-1 (**Figure 4B**). Additionally, our western blot

observations also shed light on the role of PECAM-1 in forming homophilic interactions

which was first hypothesized by Albelda et al (121) and later confirmed by a series of

studies to be necessary for its concentration at intercellular junctions. (122, 123, 124) This

subcellular localization in addition to the relative amount of PECAM-1 has been

postulated to regulate endothelial barrier integrity. (125) We hypothesized that the

subpopulation of PECAM-1/PECAM-1 detected in our studies at ~240-260kDa

corresponded to the dimerized PECAM-1 formed from homophilic interactions between

neighboring endothelial cells. In concordance with the previous observations, we

previously observed an HGD-mediated decrease in HBMEC TEER (40) and a

corresponding decrease in PECAM-1 (**Figure 4**), suggesting that the maintenance of

PECAM-1 is in part responsible for preserving endothelial barrier integrity following

ischemic-like injury potentially through the attenuation of MMP-9 activity. Overall, these

findings prompted a two-pronged interpretation. 1) The observed decrease in CD44

correlated with a decreased confinement of MMP-9 which 2) could have resulted in an

increased ability to degrade structures integral to endothelial barrier function.

Endothelial Barrier: Claudin-5 and ZO-1

To test the latter half of our conclusion we assessed levels of the integral

endothelial barrier protein, claudin-5. We chose this protein as it has been well

established as being highly expressed in brain endothelial cells at the BBB and shown to

be necessary for maintenance of BBB integrity. (8, 126) Moreover, as previously

conducted with PECAM-1 (**Supplemental Figure 1**), an in-silico prediction of claudin-5

protease cleavage sites was conducted via PROSPER revealing that MMP-9 is capable

of mediating degradation of both isoforms of claudin-5 protein (**Supplemental Figure 2**).

In line with our hypothesis and mirroring our findings of PECAM-1, we observed a notable

decrease in claudin-5 protein levels following HGD exposure that was then attenuated by

ozanimod, potentially mediated by attenuation of MMP-9 (**Figure 8A**). These results are

substantiated by findings in whole-cell extracts from rat brains that underwent a 1.5h

middle cerebral artery occlusion followed by 3h of reperfusion, which illustrated a

decrease in claudin-5 protein levels compared to sham animals and that administration

of a broad-spectrum MMP inhibitor reversed this response. (127) Moreover, in a separate

study at 6h post reperfusion following a 2h middle cerebral artery occlusion within a rat

model of ischemic stroke, a decrease in both claudin-5 and ZO-1 proteins was observed

and functionally correlated with both increased Evan's Blue extravasation and brain water

content. (128) These results correspond with the significant decrease in ZO-1 protein

levels detected in this study following HGD exposure (**Figure 8B**). However, unlike

claudin-5, the HGD-mediated decrease in ZO-1 levels was not attenuated by ozanimod

which further supports our hypothesis that ozanimod may be acting to decrease MMP-9

activity. While our findings support the aforementioned hypothesis, it is also possible that

ozanimod induces an S1PR1 mediated intracellular signaling cascade that stabilizes the

claudin-5 protein during ischemic-like injury which merits further investigation. Collectively

these data suggest that ozanimod, potentially through decreased MMP-9 activity (**Figure**

3B) leading to an attenuation of decreased claudin-5 protein levels (**Figure 8A**), lies at

the basis of the previously observed beneficial effects on endothelial barrier function. (40)

In summary, our results suggest that HGD exposure induces a concomitant

increase in MMP-9 activity and alteration of proteins associated with endothelial barrier

integrity in HBMECs and that ozanimod attenuates these HGD-induced responses, thus

adding to its therapeutic potential in cerebrovascular protection during the acute phase of

ischemic stroke. We acknowledge that the complex pathophysiological cascade shown

to occur during AIS cannot be exactly modeled in an in vitro setting and we acknowledge

that HBMECs in culture conditions do not exactly mimic the tubular characteristics similar

to those found in vivo. However, in vitro studies using human primary cells do allow the

investigation of specific basic cellular and molecular mechanisms under conditions of

hypoxia or oxygen and glucose deprivation which is similar to what is observed during

AIS. Based on our results in this study we posit that ozanimod is in part working at the

level of MMP-9 to attenuate HGD-induced decreases in HBMEC barrier integrity.

However, due to the complex and interconnectivity of GPCRs, it is also likely that

ozanimod elicits multiple responses in primary human brain microvascular endothelial

cells to attenuate ischemic stroke-like associated pathologies related to endothelial

barrier function. Thusly, future studies are planned to investigate receptor dependence

and the specific mechanism(s) of action in the cerebrovasculature including modulation

of Akt, PI3K, Rho, and upstream mediators as well as barrier proteins in freshly dissected,

uncultured, endothelial cells. Further understanding the role S1PRs play in maintaining

the integrity of the cerebrovasculature following an ischemic injury may provide a stronger

basis for the selective clinical targeting of S1PR1 with compounds such as ozanimod for

acute ischemic stroke.

FIGURE LEGENDS:

Figure 1. *MMP-9* mRNA levels in HBMECs following HGD. (A) Representative RT-

PCR gel for matrix metalloproteinase 9 (*MMP-9*) mRNA expression to verify expression

in HBMECs under normoxia or HGD conditions that revealed presence of *MMP-9* only

following HGD exposure. (B) Bar graph depicts qRT-PCR quantification of *MMP-9* mRNA

present within HBMECs exposed to either normoxia or HGD and treated with either

vehicle or ozanimod (Ozd; 0.5nM) normalized to the housekeeping gene (*GAPDH*) and

expressed as $2^{-\Delta\Delta Ct}$. DMSO was used as the vehicle for the delivery of ozanimod. n = 8-

12 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple comparisons post

hoc test. Normoxia + Vehicle vs. HGD + Vehicle **P = 0.0011, Normoxia + Ozd vs. HGD

+ Ozd ****P < 0.0001.

Figure 2. Effect of HGD on intracellular HBMEC MMP-9 protein expression. (A)

Representative western blot illustrating the band migration for two distinct MMP-9 bands,

pro-MMP-9 (92kDa; non-active form) and MMP-9 (84kDa; active form), as well as β-actin

in HBMECs following treatment with either vehicle or ozanimod (Ozd; 0.5nM) and

exposure to normoxia or HGD for 3h. Standard, SD. (B) Bar depicts pro-MMP-9 band

intensities normalized to β-actin (loading control) and expressed as fold change $\pm$ SEM.

n = 6-7 (data are mean $\pm$ SEM). Planned direct comparison was performed with unpaired

t test with Welch's correction. Normoxia + Vehicle vs. HGD + Vehicle *P = 0.0245. (C)

Bar graph illustrates band intensities of MMP-9 normalized to β-actin (loading control)

and expressed as fold change $\pm$ SEM. n = 6-7 (data are mean $\pm$ SEM). Two-way ANOVA

with Tukey's multiple comparisons post hoc test. (D) Bar graph demonstrates the ratio of

MMP-9 (C) to pro-MMP-9 (B) intensities normalized to β-actin and expressed as fold

change $\pm$ SEM. n = 6-7 (data are mean $\pm$ SEM). Planned direct comparison was

performed with unpaired t test with Welch's correction. Normoxia + Vehicle vs. HGD +

Vehicle *P = 0.0392.

Figure 3. Extracellular HBMEC MMP-9 and MMP-2 protein activity following HGD.

(A) Representative gelatin zymography illustrating the band migration and enzymatic

activity for two distinct matrix metalloproteinase bands, MMP-9 (82kDa) and MMP-2

(63kDa), in conditioned media collected from HBMECs following treatment with either

vehicle or ozanimod (Ozd; 0.5nM) and exposure to normoxia or HGD for 3h. Standard,

SD; SB, Sample Buffer. (B) Bar depicts MMP-9 activity by inverse densiometric analysis

and expressed as fold change $\pm$ SEM. n = 4 (data are mean $\pm$ SEM). Planned direct

comparisons were performed with unpaired t test with Welch's correction. Normoxia +

Vehicle vs. HGD + Vehicle *P = 0.0323, HGD + Vehicle vs HGD + Ozd **P = 0.0013. (C)

Bar graph illustrates quantification of MMP-2 activity by inverse densiometric analysis and

expressed as fold change $\pm$ SEM. n = 4 (data are mean $\pm$ SEM). Two-way ANOVA with

Tukey's multiple comparisons post hoc test. (D) Graphical illustration of observed HBMEC

MMP-9 regulation by HGD exposure and progression from intracellular mRNA and protein

to extracellular activity and the observed attenuative effects of ozanimod on MMP-9

activity.

Figure 4. HBMEC PECAM-1 protein expression following HGD. (A) Representative

immunocytochemical images demonstrating the labeling of platelet endothelial cell

adhesion molecule 1 (PECAM-1; red) in HBMECs following exposure to either HGD or

normoxia for 3h and treatment with vehicle or ozanimod (Ozd; 0.5nM). Imaged at 40X

magnification. Nucleus is labeled with 4',6-diamidine-2'-phenylindole dihydrochloride

(DAPI; blue). Scale bar, 50μM. (B) Bar graph depicts the relative quantification of

PECAM-1 in HBMECs normalized to the number of nuclei and expressed as a ratio of the

area occupied by PECAM-1 to number of nuclei (μm^2/nuclei). n = 4 (data are mean $\pm$

SEM). Planned direct comparisons were performed with unpaired t test. Normoxia +

Vehicle vs. HGD + Vehicle *P = 0.0131, HGD + Vehicle vs. HGD + Ozd *P = 0.0492. (C)

Representative western blot illustrating the band migration for three distinct sets of

platelet endothelial cell adhesion molecule 1 (PECAM-1) bands, PECAM-1/PECAM-1

(240-260kDa), PECAM-1 (130kDa), and a PECAM-1 Heavy Band (95kDa) in HBMECs

following treatment with either vehicle or ozanimod (Ozd; 0.5nM) and exposure to

normoxia or HGD for 3h. PECAM-1 expression was also demonstrated in mouse aortic

tissue lysate to serve as a positive control for the PECAM-1 antibody but was not included

in the interpretations of the study.

Figure 5. HGD induced alteration in HBMEC extracellular hydrogen peroxide (H_2O_2)

production. (A) Representative western blot demonstrates the band migration of

vascular cell adhesion molecule 1 (VCAM-1; 100kDa) in HBMECs following treatment

with either vehicle or ozanimod (Ozd; 0.5nM) and exposure to normoxia or HGD for 3h.

Bar graphs illustrate densiometric analysis of VCAM-1. n = 6 (data are mean $\pm$ SEM).

Planned direct comparison was performed with unpaired t test with Welch's correction.

Normoxia + Vehicle vs. HGD + Vehicle *P = 0.0199. Standard, SD; SB, Sample Buffer.

(B) Bar graph depicts qRT-PCR mediated quantitation of *SOD1* mRNA present within

HBMECs exposed to either normoxia or HGD and treated with either vehicle or ozanimod

(Ozd; 0.5nM) normalized to the housekeeping gene (*GAPDH*) and expressed as $2^{-\Delta\Delta Ct}$.

n = 10-15 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple comparisons

post hoc test. Normoxia + Vehicle vs. HGD + Vehicle ****P < 0.0001, HGD + Vehicle vs.

HGD + Ozd P = 0.0850. (C) Bar graph depicts qRT-PCR mediated quantitation of *SOD2*

mRNA present within HBMECs exposed to either normoxia or HGD and treated with

either vehicle or ozanimod (Ozd; 0.5nM) normalized to the housekeeping gene (*GAPDH*)

and expressed as $2^{-\Delta\Delta Ct}$. *n* = 8-12 (data are mean $\pm$ SEM). Two-way ANOVA with

Tukey's multiple comparisons post hoc test. (D) Graph demonstrates the quantification of

extracellular concentrations H_2O_2 from conditioned media extracted from HBMEC

cultures exposed to either normoxia or HGD for 1 and 3h and treated with vehicle (Veh)

or ozanimod (Ozd; 0.5nM). *n* = 3 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's

multiple comparisons post hoc test. 1h HGD + Vehicle vs. 3h HGD + Vehicle **P = 0.0015,

3h Normoxia + Vehicle vs. 3h HGD + Vehicle **P = 0.0014.

Figure 6. Extracellular transport, inhibition, and docking of MMP-9 in HBMECs

following HGD. (A) Bar graph depicts qRT-PCR mediated quantitation of *RECK* mRNA

present within HBMECs exposed to either normoxia or HGD and treated with either

vehicle or ozanimod (Ozd; 0.5nM) normalized to the housekeeping gene (*GAPDH*) and

expressed as $2^{-\Delta\Delta Ct}$. *n* = 8-12 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's

multiple comparisons post hoc test. Normoxia + Vehicle vs. HGD + Vehicle ****P <

0.0001. (B) Bar graph depicts qRT-PCR mediated quantitation of *TIMP1* mRNA present

within HBMECs exposed to either normoxia or HGD and treated with either vehicle or

ozanimod (Ozd; 0.5nM) normalized to the housekeeping gene (*GAPDH*) and expressed

as $2^{-\Delta\Delta Ct}$. *n* = 8-11 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple

comparisons post hoc test. Normoxia + Vehicle vs. HGD + Vehicle ***P = 0.0008.

Figure 7. CD44 protein expression and nuclear localization in HBMECs following

HGD. (A) Representative immunocytochemical images demonstrating the nuclear or non-

nuclear localization and labeling of anti-CD44 (CD44; green) in HBMECs following

exposure to either HGD or normoxia for 3h and treatment with vehicle or ozanimod (Ozd;

0.5nM). Imaged at 40X magnification. Nucleus is labeled with 4',6-diamidine-2'-

phenylindole dihydrochloride (DAPI; blue). Scale bar, 50μM. (B) Bar graph depicts the

relative quantification of CD44 in HBMECs normalized to the number of nuclei and

expressed as a ratio of the area occupied by CD44 to number of nuclei (μm^2/nuclei). n =

4 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple comparisons post hoc

test. Normoxia + Vehicle vs. HGD + Vehicle ***P = 0.0002. (C) Bar graph depicts

Mander's Coefficient (M1) of the overlap of CD44 with DAPI stained nuclei calculated

from the images shown in (A). n = 4 (data are mean $\pm$ SEM). Two-way ANOVA with

Tukey's multiple comparisons post hoc test. Normoxia + Vehicle vs. Normoxia + Ozd **P

= 0.0041, Normoxia + Ozd vs. HGD + Ozd **P = 0.0078. (D) Bar graph depicts Mander's

Coefficient (M2) of the overlap of DAPI stained nuclei with CD44 calculated from the

images shown in (A). n = 4 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's

multiple comparisons post hoc test. Normoxia + Vehicle vs. Normoxia + Ozd *P = 0.0435,

Normoxia + Ozd vs. HGD + Ozd *P = 0.0107.

1004

Figure 8. Impact of ozanimod on levels of HBMEC barrier proteins ZO-1 and

claudin-5 following. Representative western blots demonstrate the band migration of

(A) claudin-5 (35kDa), (B) zonula occludens 1 (ZO-1; 240kDa), and β-actin (42kDa) in

HBMECs following treatment with either vehicle or ozanimod (Ozd; 0.5nM) and exposure

to normoxia or HGD for 3h. Bar graphs illustrate densiometric analysis of claudin-5 (A),

and ZO-1 (B) expressed as fold change to normoxia plus vehicle. (A) n = 8-10 (data are

mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple comparisons post hoc test.

Normoxia + Vehicle vs. Normoxia + Ozd ***P = 0.0002. Planned direct comparison was

performed with unpaired t test with Welch's correction. HGD + Vehicle vs. HGD + Ozd

**P = 0.0093. (B) n = 6 (data are mean $\pm$ SEM). Two-way ANOVA with Tukey's multiple

comparisons post hoc test. Normoxia + Vehicle vs. HGD + Vehicle ****P < 0.0001.

Standard, SD; SB, Sample Buffer.

1017

Supplemental Figure 1. In silico analysis of cleavage sites within human PECAM-1

via PROSPER. (A) Whole sequence of human platelet endothelial cell adhesion molecule

1 (PECAM-1) highlighted at specific amino acid sites to indicate potential cleavage by

aspartic proteases (black), cysteine proteases (green), metalloproteases (yellow), serine

proteases (blue), and different multiple protease superfamilies (red). (B-D) Within the

PECAM-1 sequence highlighted yellow, specific sites with a cleavage probability score

greater than 0.8 are shown as red lines and represent independent, potential sites of

human PECAM-1 cleavage by MMP-9 (39 independent sites) (B), MMP-2 (3 independent

sites) (C), and MMP-3 (7 independent sites) (D).

1027

Supplemental Figure 2. In silico alignment of claudin-5 isoforms and analysis of

cleavage sites. (A) Whole proteomic sequence alignment of claudin-5 isoforms 1 and 2

demonstrating total length of isoform 1 (303 amino acids) and isoform 2 (218 amino

acids). Asterisks indicate overlapping amino acids between the two claudin-5 isoforms.

(B) Whole sequence of human claudin-5 isoform 1 highlighted at specific amino acid sites

to indicate potential cleavage by aspartic proteases (black), cysteine proteases (green),

metalloproteases (yellow), serine proteases (blue), and different multiple protease superfamilies (red). (C) Whole sequence of human claudin-5 isoform 2 highlighted at specific amino acid sites to indicate potential cleavage by aspartic proteases (black), cysteine proteases (green), metalloproteases (yellow), serine proteases (blue), and different multiple protease superfamilies (red). (D-E) Within the claudin-5 isoform 1 and 2 sequences highlighted yellow, specific sites with a cleavage probability score greater than 0.8 are shown as red lines and represent independent, potential sites of human claudin-5 isoform 1 cleavage by MMP-9 (20 independent sites) (D) and claudin-5 isoform 2 cleavage by MMP-9 (17 independent sites) (E).

1.   Kunz A, Iadecola C. Cerebral vascular dysregulation in the ischemic brain. Handb Clin Neurol. 2009;92:283-305.

2.   Hu X, De Silva TM, Chen J, Faraci FM. Cerebral Vascular Disease and Neurovascular Injury in Ischemic Stroke. Circ Res. 2017;120(3):449-71.

3.   Nelson CW, Wei EP, Povlishock JT, Kontos HA, Moskowitz MA. Oxygen radicals in cerebral ischemia. Am J Physiol. 1992;263(5 Pt 2):H1356-62.

4.   Abo-Ramadan U, Durukan A, Pitkonen M, Marinkovic I, Tatlisumak E, Pedrono E, et al. Post-ischemic leakiness of the blood-brain barrier: a quantitative and systematic assessment by Patlak plots. Exp Neurol. 2009;219(1):328-33.

5.   Jiang X, Andjelkovic AV, Zhu L, Yang T, Bennett MVL, Chen J, et al. Blood-brain barrier dysfunction and recovery after ischemic stroke. Prog Neurobiol. 2018;163-164:144-71.

6.   Henning EC, Latour LL, Warach S. Verification of enhancement of the CSF space, not parenchyma, in acute stroke patients with early blood-brain barrier disruption. J Cereb Blood Flow Metab. 2008;28(5):882-6.

7.   Nadareishvili Z, Simpkins AN, Hitomi E, Reyes D, Leigh R. Post-Stroke Blood-Brain Barrier Disruption and Poor Functional Outcome in Patients Receiving Thrombolytic Therapy. Cerebrovasc Dis. 2019;47(3-4):135-42.

8.   Greene C, Hanley N, Campbell M. Claudin-5: gatekeeper of neurological function. Fluids
Barriers CNS. 2019;16(1):3.

9.   Tornavaca O, Chia M, Dufton N, Almagro LO, Conway DE, Randi AM, et al. ZO-1 controls
endothelial adherens junctions, cell-cell tension, angiogenesis, and barrier formation. J Cell Biol.
2015;208(6):821-38.

10.   Wimmer I, Tietz S, Nishihara H, Deutsch U, Sallusto F, Gosselet F, et al. PECAM-1
Stabilizes Blood-Brain Barrier Integrity and Favors Paracellular T-Cell Diapedesis Across the
Blood-Brain Barrier During Neuroinflammation. Front Immunol. 2019;10:711.

11.   Flynn KM, Michaud M, Canosa S, Madri JA. CD44 regulates vascular endothelial barrier
integrity via a PECAM-1 dependent mechanism. Angiogenesis. 2013;16(3):689-705.

12.   Baeten KM, Akassoglou K. Extracellular matrix and matrix receptors in blood-brain barrier
formation and stroke. Dev Neurobiol. 2011;71(11):1018-39.

13.   Chen F, Ohashi N, Li W, Eckman C, Nguyen JH. Disruptions of occludin and claudin-5 in
brain endothelial cells in vitro and in brains of mice with acute liver failure. Hepatology.
2009;50(6):1914-23.

14.   Harkness KA, Adamson P, Sussman JD, Davies-Jones GA, Greenwood J, Woodroofe
MN. Dexamethasone regulation of matrix metalloproteinase expression in CNS vascular
endothelium. Brain. 2000;123 (Pt 4):698-709.

15.   Brilha S, Ong CWM, Weksler B, Romero N, Couraud PO, Friedland JS. Matrix
metalloproteinase-9 activity and a downregulated Hedgehog pathway impair blood-brain barrier
function in an in vitro model of CNS tuberculosis. Sci Rep. 2017;7(1):16031.

16.   Kato H, Kuriyama N, Duarte S, Clavien PA, Busuttil RW, Coito AJ. MMP-9 deficiency
shelters endothelial PECAM-1 expression and enhances regeneration of steatotic livers after
ischemia and reperfusion injury. J Hepatol. 2014;60(5):1032-9.

17.   Yosef G, Hayun H, Papo N. Simultaneous targeting of CD44 and MMP9 catalytic and
hemopexin domains as a therapeutic strategy. Biochem J. 2021;478(5):1139-57.

18.   Del Zoppo GJ, Milner R, Mabuchi T, Hung S, Wang X, Koziol JA. Vascular matrix adhesion
and the blood-brain barrier. Biochem Soc Trans. 2006;34(Pt 6):1261-6.

19.   Kwon I, Kim EH, del Zoppo GJ, Heo JH. Ultrastructural and temporal changes of the
microvascular basement membrane and astrocyte interface following focal cerebral ischemia. J
Neurosci Res. 2009;87(3):668-76.

20.   Jiao H, Wang Z, Liu Y, Wang P, Xue Y. Specific role of tight junction proteins claudin-5,
occludin, and ZO-1 of the blood-brain barrier in a focal cerebral ischemic insult. J Mol Neurosci.
2011;44(2):130-9.

21.   Liu P, Zhang R, Liu D, Wang J, Yuan C, Zhao X, et al. Time-course investigation of blood-
brain barrier permeability and tight junction protein changes in a rat model of permanent focal
ischemia. J Physiol Sci. 2018;68(2):121-7.

22.   Sifat AE, Vaidya B, Abbruscato TJ. Blood-Brain Barrier Protection as a Therapeutic
Strategy for Acute Ischemic Stroke. AAPS J. 2017;19(4):957-72.

23.   Romanic AM, White RF, Arleth AJ, Ohlstein EH, Barone FC. Matrix metalloproteinase
expression increases after cerebral focal ischemia in rats: inhibition of matrix metalloproteinase-
9 reduces infarct size. Stroke. 1998;29(5):1020-30.

24.   Zhong C, Yang J, Xu T, Peng Y, Wang A, Wang J, et al. Serum matrix metalloproteinase-
9 levels and prognosis of acute ischemic stroke. Neurology. 2017;89(8):805-12.

25.   Castellanos M, Leira R, Serena J, Pumar JM, Lizasoain I, Castillo J, et al. Plasma
metalloproteinase-9 concentration predicts hemorrhagic transformation in acute ischemic stroke.
Stroke. 2003;34(1):40-6.

26.   Abdelnaseer MM, Elfauomy NM, Esmail EH, Kamal MM, Elsawy EH. Matrix
Metalloproteinase-9 and Recovery of Acute Ischemic Stroke. J Stroke Cerebrovasc Dis.
2017;26(4):733-40.

27.   Dong X, Song YN, Liu WG, Guo XL. Mmp-9, a potential target for cerebral ischemic
treatment. Curr Neuropharmacol. 2009;7(4):269-75.

28.   Huang ZG, Xue D, Preston E, Karbalai H, Buchan AM. Biphasic opening of the blood-
brain barrier following transient focal ischemia: effects of hypothermia. Can J Neurol Sci.
1999;26(4):298-304.

29.   Bernardo-Castro S, Sousa JA, Brás A, Cecília C, Rodrigues B, Almendra L, et al.
Pathophysiology of Blood-Brain Barrier Permeability Throughout the Different Stages of Ischemic
Stroke and Its Implication on Hemorrhagic Transformation and Recovery. Front Neurol.
2020;11:594672.

30.   Sotgiu S, Zanda B, Marchetti B, Fois ML, Arru G, Pes GM, et al. Inflammatory biomarkers
in blood of patients with acute brain ischemia. Eur J Neurol. 2006;13(5):505-13.

31.   Fu Y, Zhang N, Ren L, Yan Y, Sun N, Li YJ, et al. Impact of an immune modulator
fingolimod on acute ischemic stroke. Proc Natl Acad Sci U S A. 2014;111(51):18315-20.

32.   Tran JQ, Hartung JP, Olson AD, Mendzelevski B, Timony GA, Boehm MF, et al. Cardiac
Safety of Ozanimod, a Novel Sphingosine-1-Phosphate Receptor Modulator: Results of a
Thorough QT/QTc Study. Clin Pharmacol Drug Dev. 2018;7(3):263-76.

33.   Cohen JA, Arnold DL, Comi G, Bar-Or A, Gujrathi S, Hartung JP, et al. Safety and efficacy
of the selective sphingosine 1-phosphate receptor modulator ozanimod in relapsing multiple
sclerosis (RADIANCE): a randomised, placebo-controlled, phase 2 trial. Lancet Neurol.
2016;15(4):373-81.

34.   Fu Y, Hao J, Zhang N, Ren L, Sun N, Li YJ, et al. Fingolimod for the treatment of
intracerebral hemorrhage: a 2-arm proof-of-concept study. JAMA Neurol. 2014;71(9):1092-101.

35.   Zhu Z, Fu Y, Tian D, Sun N, Han W, Chang G, et al. Combination of the Immune Modulator
Fingolimod With Alteplase in Acute Ischemic Stroke: A Pilot Trial. Circulation. 2015;132(12):1104-
12.

36.   Tian DC, Shi K, Zhu Z, Yao J, Yang X, Su L, et al. Fingolimod enhances the efficacy of
delayed alteplase administration in acute ischemic stroke by promoting anterograde reperfusion
and retrograde collateral flow. Ann Neurol. 2018;84(5):717-28.

37.   Lassiter G, Melancon C, Rooney T, Murat AM, Kaye JS, Kaye AM, et al. Ozanimod to
Treat Relapsing Forms of Multiple Sclerosis: A Comprehensive Review of Disease, Drug Efficacy
and Side Effects. Neurol Int. 2020;12(3):89-108.

38.   Sandborn WJ, Feagan BG, D'Haens G, Wolf DC, Jovanovic I, Hanauer SB, et al.
Ozanimod as Induction and Maintenance Therapy for Ulcerative Colitis. N Engl J Med.
2021;385(14):1280-91.

39.   Li W, Xu H, Testai FD. Mechanism of Action and Clinical Potential of Fingolimod for the
Treatment of Stroke. Front Neurol. 2016;7:139.

40.   Li YJ, Shi SX, Liu Q, Shi FD, Gonzales RJ. Targeted role for sphingosine-1-phosphate
receptor 1 in cerebrovascular integrity and inflammation during acute ischemic stroke. Neurosci
Lett. 2020;735:135160.

41.   Nitzsche A, Poittevin M, Benarab A, Bonnin P, Faraco G, Uchida H, et al. Endothelial S1P.
Circ Res. 2021;128(3):363-82.

42.   Zhang SQ, Xiao J, Chen M, Zhou LQ, Shang K, Qin C, et al. Sphingosine-1-Phosphate
Signaling in Ischemic Stroke: From Bench to Bedside and Beyond. Front Cell Neurosci.
2021;15:781098.

43.   Kim GS, Yang L, Zhang G, Zhao H, Selim M, McCullough LD, et al. Critical role of
sphingosine-1-phosphate receptor-2 in the disruption of cerebrovascular integrity in experimental
stroke. Nat Commun. 2015;6:7893.

44.   Lee MJ, Evans M, Hla T. The inducible G protein-coupled receptor edg-1 signals via the
G(i)/mitogen-activated protein kinase pathway. J Biol Chem. 1996;271(19):11272-9.

45.   Dantas AP, Igarashi J, Michel T. Sphingosine 1-phosphate and control of vascular tone.
Am J Physiol Heart Circ Physiol. 2003;284(6):H2045-52.

46.   Mehta D, Konstantoulaki M, Ahmmed GU, Malik AB. Sphingosine 1-phosphate-induced
mobilization of intracellular Ca2+ mediates rac activation and adherens junction assembly in
endothelial cells. J Biol Chem. 2005;280(17):17320-8.

47.   Lee MJ, Thangada S, Claffey KP, Ancellin N, Liu CH, Kluk M, et al. Vascular endothelial
cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. Cell.
1999;99(3):301-12.

48.   Wendt TS, Li YJ, Gonzales RJ. Ozanimod, an S1PR1 ligand, attenuates hypoxia plus
glucose deprivation induced autophagic flux and phenotypic switching in human brain VSM cells.
Am J Physiol Cell Physiol. 2021.

49.   Du J, Zeng C, Li Q, Chen B, Liu H, Huang X, et al. LPS and TNF-α induce expression of
sphingosine-1-phosphate receptor-2 in human microvascular endothelial cells. Pathol Res Pract.
2012;208(2):82-8.

50.   Grailhe P, Boutarfa-Madec A, Beauverger P, Janiak P, Parkar AA. A label-free impedance
assay in endothelial cells differentiates the activation and desensitization properties of clinical
S1P. FEBS Open Bio. 2020;10(10):2010-20.

51.   Lee JF, Zeng Q, Ozaki H, Wang L, Hand AR, Hla T, et al. Dual roles of tight junction-
associated protein, zonula occludens-1, in sphingosine 1-phosphate-mediated endothelial
chemotaxis and barrier integrity. J Biol Chem. 2006;281(39):29190-200.

52.   Akhter MZ, Chandra Joshi J, Balaji Ragunathrao VA, Maienschein-Cline M, Proia RL,
Malik AB, et al. Programming to S1PR1. Circ Res. 2021;129(2):221-36.

53.   Cartier A, Leigh T, Liu CH, Hla T. Endothelial sphingosine 1-phosphate receptors promote
vascular normalization and antitumor therapy. Proc Natl Acad Sci U S A. 2020;117(6):3157-66.

54.   Gaengel K, Niaudet C, Hagikura K, Laviña B, Siemsen BL, Muhl L, et al. The sphingosine-
1-phosphate receptor S1PR1 restricts sprouting angiogenesis by regulating the interplay between
VE-cadherin and VEGFR2. Dev Cell. 2012;23(3):587-99.

55.   Cantalupo A, Gargiulo A, Dautaj E, Liu C, Zhang Y, Hla T, et al. S1PR1 (Sphingosine-1-
Phosphate Receptor 1) Signaling Regulates Blood Flow and Pressure. Hypertension.
2017;70(2):426-34.

56.   Rutherford C, Childs S, Ohotski J, McGlynn L, Riddick M, MacFarlane S, et al. Regulation
of cell survival by sphingosine-1-phosphate receptor S1P1 via reciprocal ERK-dependent
suppression of Bim and PI-3-kinase/protein kinase C-mediated upregulation of Mcl-1. Cell Death
Dis. 2013;4:e927.

57.   Zuloaga KL, Swift SN, Gonzales RJ, Wu TJ, Handa RJ. The androgen metabolite, 5α-
androstane-3β,17β-diol, decreases cytokine-induced cyclooxygenase-2, vascular cell adhesion
molecule-1 expression, and P-glycoprotein expression in male human brain microvascular
endothelial cells. Endocrinology. 2012;153(12):5949-60.

58.   Gonzalez-Cabrera PJ, Jo E, Sanna MG, Brown S, Leaf N, Marsolais D, et al. Full
pharmacological efficacy of a novel S1P1 agonist that does not require S1P-like headgroup
interactions. Mol Pharmacol. 2008;74(5):1308-18.

59.   Raman P, Madhavpeddi L, Gonzales RJ. Palmitate induces glycosylation of
cyclooxygenase-2 in primary human vascular smooth muscle cells. Am J Physiol Cell Physiol.
2018;314(5):C545-C53.

60.   Hollborn M, Stathopoulos C, Steffen A, Wiedemann P, Kohen L, Bringmann A. Positive
feedback regulation between MMP-9 and VEGF in human RPE cells. Invest Ophthalmol Vis Sci.
2007;48(9):4360-7.

61.   Christensen J, Shastri VP. Matrix-metalloproteinase-9 is cleaved and activated by
cathepsin K. BMC Res Notes. 2015;8:322.

62.   Park S, Sorenson CM, Sheibani N. PECAM-1 isoforms, eNOS and endoglin axis in
regulation of angiogenesis. Clin Sci (Lond). 2015;129(3):217-34.

63.   Hu M, Zhang H, Liu Q, Hao Q. Structural Basis for Human PECAM-1-Mediated Trans-
homophilic Cell Adhesion. Sci Rep. 2016;6:38655.

64.   Newton JP, Hunter AP, Simmons DL, Buckley CD, Harvey DJ. CD31 (PECAM-1) exists
as a dimer and is heavily N-glycosylated. Biochem Biophys Res Commun. 1999;261(2):283-91.

65.   Wang Y, Su X, Sorenson CM, Sheibani N. Tissue-specific distributions of alternatively
spliced human PECAM-1 isoforms. Am J Physiol Heart Circ Physiol. 2003;284(3):H1008-17.

66.   Bienert GP, Schjoerring JK, Jahn TP. Membrane transport of hydrogen peroxide. Biochim
Biophys Acta. 2006;1758(8):994-1003.

67.   Gough DR, Cotter TG. Hydrogen peroxide: a Jekyll and Hyde signalling molecule. Cell
Death Dis. 2011;2:e213.

68.   Haorah J, Ramirez SH, Schall K, Smith D, Pandya R, Persidsky Y. Oxidative stress
activates protein tyrosine kinase and matrix metalloproteinases leading to blood-brain barrier
dysfunction. J Neurochem. 2007;101(2):566-76.

69.   Deem TL, Cook-Mills JM. Vascular cell adhesion molecule 1 (VCAM-1) activation of
endothelial cell matrix metalloproteinases: role of reactive oxygen species. Blood.
2004;104(8):2385-93.

70.   Takahashi C, Sheng Z, Horan TP, Kitayama H, Maki M, Hitomi K, et al. Regulation of
matrix metalloproteinase-9 and inhibition of tumor invasion by the membrane-anchored
glycoprotein RECK. Proc Natl Acad Sci U S A. 1998;95(22):13221-6.

71.   Chang CK, Hung WC, Chang HC. The Kazal motifs of RECK protein inhibit MMP-9
secretion and activity and reduce metastasis of lung cancer cells in vitro and in vivo. J Cell Mol
Med. 2008;12(6B):2781-9.

72.   Roderfeld M, Graf J, Giese B, Salguero-Palacios R, Tschuschner A, Müller-Newen G, et
al. Latent MMP-9 is bound to TIMP-1 before secretion. Biol Chem. 2007;388(11):1227-34.

73. Tsuneki M, Madri JA. CD44 regulation of endothelial cell proliferation and apoptosis via
modulation of CD31 and VE-cadherin expression. J Biol Chem. 2014;289(9):5357-70.

74. Yu Q, Stamenkovic I. Localization of matrix metalloproteinase 9 to the cell surface
provides a mechanism for CD44-mediated tumor invasion. Genes Dev. 1999;13(1):35-48.

75. Senbanjo LT, Chellaiah MA. CD44: A Multifunctional Cell Surface Adhesion Receptor Is
a Regulator of Progression and Metastasis of Cancer Cells. Front Cell Dev Biol. 2017;5:18.

76. Okamoto I, Kawano Y, Murakami D, Sasayama T, Araki N, Miki T, et al. Proteolytic release
of CD44 intracellular domain and its role in the CD44 signaling pathway. J Cell Biol.
2001;155(5):755-62.

77. Miletti-González KE, Murphy K, Kumaran MN, Ravindranath AK, Wernyj RP, Kaur S, et
al. Identification of function for CD44 intracytoplasmic domain (CD44-ICD): modulation of matrix
metalloproteinase 9 (MMP-9) transcription via novel promoter response element. J Biol Chem.
2012;287(23):18995-9007.

78. Claesson-Welsh L, Dejana E, McDonald DM. Permeability of the Endothelial Barrier:
Identifying and Reconciling Controversies. Trends Mol Med. 2021;27(4):314-31.

79. Andjelkovic AV, Xiang J, Stamatovic SM, Hua Y, Xi G, Wang MM, et al. Endothelial
Targets in Stroke: Translating Animal Models to Human. Arterioscler Thromb Vasc Biol.
2019;39(11):2240-7.

80. Nian K, Harding IC, Herman IM, Ebong EE. Blood-Brain Barrier Damage in Ischemic
Stroke and Its Regulation by Endothelial Mechanotransduction. Front Physiol. 2020;11:605398.

81. Krueger M, Mages B, Hobusch C, Michalski D. Endothelial edema precedes blood-brain
barrier breakdown in early time points after experimental focal cerebral ischemia. Acta
Neuropathol Commun. 2019;7(1):17.

82. Ramos-Fernandez M, Bellolio MF, Stead LG. Matrix metalloproteinase-9 as a marker for
acute ischemic stroke: a systematic review. J Stroke Cerebrovasc Dis. 2011;20(1):47-54.

83. Turner RJ, Sharp FR. Implications of MMP9 for Blood Brain Barrier Disruption and
Hemorrhagic Transformation Following Ischemic Stroke. Front Cell Neurosci. 2016;10:56.

84. Asahi M, Asahi K, Jung JC, del Zoppo GJ, Fini ME, Lo EH. Role for matrix
metalloproteinase 9 after focal cerebral ischemia: effects of gene knockout and enzyme inhibition
with BB-94. J Cereb Blood Flow Metab. 2000;20(12):1681-9.

85. Asahi M, Wang X, Mori T, Sumii T, Jung JC, Moskowitz MA, et al. Effects of matrix
metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter
components after cerebral ischemia. J Neurosci. 2001;21(19):7724-32.

86. Suofu Y, Clark JF, Broderick JP, Kurosawa Y, Wagner KR, Lu A. Matrix metalloproteinase-
2 or -9 deletions protect against hemorrhagic transformation during early stage of cerebral
ischemia and reperfusion. Neuroscience. 2012;212:180-9.

87.   Cai H, Ma Y, Jiang L, Mu Z, Jiang Z, Chen X, et al. Hypoxia Response Element-Regulated
MMP-9 Promotes Neurological Recovery via Glial Scar Degradation and Angiogenesis in Delayed
Stroke. Mol Ther. 2017;25(6):1448-59.

88.   Bond M, Chase AJ, Baker AH, Newby AC. Inhibition of transcription factor NF-kappaB
reduces matrix metalloproteinase-1, -3 and -9 production by vascular smooth muscle cells.
Cardiovasc Res. 2001;50(3):556-65.

89.   Wang L, Jiang H, Shen SM, Wen CX, Xing Z, Shi Y. Inhibition of autophagy and
chemokine induction by sphingosine 1-phosphate receptor 1 through NF-κB signaling in human
pulmonary endothelial cells infected with influenza A viruses. PLoS One. 2018;13(10):e0205344.

90.   Furic L, Rong L, Larsson O, Koumakpayi IH, Yoshida K, Brueschke A, et al. eIF4E
phosphorylation promotes tumorigenesis and is associated with prostate cancer progression.
Proc Natl Acad Sci U S A. 2010;107(32):14134-9.

91.   Gkogkas CG, Khoutorsky A, Cao R, Jafarnejad SM, Prager-Khoutorsky M, Giannakas N,
et al. Pharmacogenetic inhibition of eIF4E-dependent Mmp9 mRNA translation reverses fragile X
syndrome-like phenotypes. Cell Rep. 2014;9(5):1742-55.

92.   Taraboletti G, D'Ascenzo S, Borsotti P, Giavazzi R, Pavan A, Dolo V. Shedding of the
matrix metalloproteinases MMP-2, MMP-9, and MT1-MMP as membrane vesicle-associated
components by endothelial cells. Am J Pathol. 2002;160(2):673-80.

93.   Cook-Mills JM. Hydrogen peroxide activation of endothelial cell-associated MMPs during
VCAM-1-dependent leukocyte migration. Cell Mol Biol (Noisy-le-grand). 2006;52(4):8-16.

94.   Cook-Mills JM, Marchese ME, Abdala-Valencia H. Vascular cell adhesion molecule-1
expression and signaling during disease: regulation by reactive oxygen species and antioxidants.
Antioxid Redox Signal. 2011;15(6):1607-38.

95.   Maglinger B, Sands M, Frank JA, McLouth CJ, Trout AL, Roberts JM, et al. Intracranial
VCAM1 at time of mechanical thrombectomy predicts ischemic stroke severity. J
Neuroinflammation. 2021;18(1):109.

96.   Yousef H, Czupalla CJ, Lee D, Chen MB, Burke AN, Zera KA, et al. Aged blood impairs
hippocampal neural precursor activity and activates microglia via brain endothelial cell VCAM1.
Nat Med. 2019;25(6):988-1000.

97.   Milstone DS, Ilyama M, Chen M, O'Donnell P, Davis VM, Plutzky J, et al. Differential role
of an NF-κB transcriptional response element in endothelial versus intimal cell VCAM-1
expression. Circ Res. 2015;117(2):166-77.

98.   Wang Y, Branicky R, Noë A, Hekimi S. Superoxide dismutases: Dual roles in controlling
ROS damage and regulating ROS signaling. J Cell Biol. 2018;217(6):1915-28.

99.   Miao L, St Clair DK. Regulation of superoxide dismutase genes: implications in disease.
Free Radic Biol Med. 2009;47(4):344-56.

100.   Bretón-Romero R, Lamas S. Hydrogen peroxide signaling in vascular endothelial cells.
Redox Biol. 2014;2:529-34.

101.   Takagi S, Simizu S, Osada H. RECK negatively regulates matrix metalloproteinase-9
transcription. Cancer Res. 2009;69(4):1502-8.

102.   Drishya G, Nambiar J, Shaji SK, Vanuopadath M, Achuthan A, Kumar A, et al. RECK and
TIMP-2 mediate inhibition of MMP-2 and MMP-9 by. J Biosci. 2020;45.

103.   Chen G, Ge D, Zhu B, Shi H, Ma Q. Upregulation of matrix metalloproteinase 9
(MMP9)/tissue inhibitor of metalloproteinase 1 (TIMP1) and MMP2/TIMP2 ratios may be involved
in lipopolysaccharide-induced acute lung injury. J Int Med Res. 2020;48(4):300060520919592.

104.   Charzewski Ł, Krzyśko KA, Lesyng B. Structural characterisation of inhibitory and non-
inhibitory MMP-9-TIMP-1 complexes and implications for regulatory mechanisms of MMP-9. Sci
Rep. 2021;11(1):13376.

105.   Visse R, Nagase H. Matrix metalloproteinases and tissue inhibitors of metalloproteinases:
structure, function, and biochemistry. Circ Res. 2003;92(8):827-39.

106.   Wang H, Imamura Y, Ishibashi R, Chandana EP, Yamamoto M, Noda M. The Reck tumor
suppressor protein alleviates tissue damage and promotes functional recovery after transient
cerebral ischemia in mice. J Neurochem. 2010;115(2):385-98.

107.   Lee KJ, Lee KY, Lee YM. Downregulation of a tumor suppressor RECK by hypoxia through
recruitment of HDAC1 and HIF-1alpha to reverse HRE site in the promoter. Biochim Biophys Acta.
2010;1803(5):608-16.

108.   Fujimoto M, Takagi Y, Aoki T, Hayase M, Marumo T, Gomi M, et al. Tissue inhibitor of
metalloproteinases protect blood-brain barrier disruption in focal cerebral ischemia. J Cereb Blood
Flow Metab. 2008;28(10):1674-85.

109.   Yu Q, Stamenkovic I. Cell surface-localized matrix metalloproteinase-9 proteolytically
activates TGF-beta and promotes tumor invasion and angiogenesis. Genes Dev. 2000;14(2):163-
76.

110.   Chen C, Zhao S, Karnad A, Freeman JW. The biology and role of CD44 in cancer
progression: therapeutic implications. J Hematol Oncol. 2018;11(1):64.

111.   Chen L, Fu C, Zhang Q, He C, Zhang F, Wei Q. The role of CD44 in pathological
angiogenesis. FASEB J. 2020;34(10):13125-39.

112.   Rouschop KM, Roelofs JJ, Claessen N, da Costa Martins P, Zwaginga JJ, Pals ST, et al.
Protection against renal ischemia reperfusion injury by CD44 disruption. J Am Soc Nephrol.
2005;16(7):2034-43.

113.   Wang H, Zhan Y, Xu L, Feuerstein GZ, Wang X. Use of suppression subtractive
hybridization for differential gene expression in stroke: discovery of CD44 gene expression and
localization in permanent focal stroke in rats. Stroke. 2001;32(4):1020-7.

114.   Al-Ahmad AJ, Patel R, Palecek SP, Shusta EV. Hyaluronan impairs the barrier integrity of
brain microvascular endothelial cells through a CD44-dependent pathway. J Cereb Blood Flow
Metab. 2019;39(9):1759-75.

115.   Kajita M, Itoh Y, Chiba T, Mori H, Okada A, Kinoh H, et al. Membrane-type 1 matrix
metalloproteinase cleaves CD44 and promotes cell migration. J Cell Biol. 2001;153(5):893-904.

116.   Langlois S, Gingras D, Béliveau R. Membrane type 1-matrix metalloproteinase (MT1-
MMP) cooperates with sphingosine 1-phosphate to induce endothelial cell migration and
morphogenic differentiation. Blood. 2004;103(8):3020-8.

117.   Kato H, Duarte S, Liu D, Busuttil RW, Coito AJ. Matrix Metalloproteinase-2 (MMP-2) Gene
Deletion Enhances MMP-9 Activity, Impairs PARP-1 Degradation, and Exacerbates Hepatic
Ischemia and Reperfusion Injury in Mice. PLoS One. 2015;10(9):e0137642.

118.   Eshaq RS, Harris NR. Loss of Platelet Endothelial Cell Adhesion Molecule-1 (PECAM-1)
in the Diabetic Retina: Role of Matrix Metalloproteinases. Invest Ophthalmol Vis Sci.
2019;60(2):748-60.

119.   Song J, Tan H, Perry AJ, Akutsu T, Webb GI, Whisstock JC, et al. PROSPER: an
integrated feature-based tool for predicting protease substrate cleavage sites. PLoS One.
2012;7(11):e50300.

120.   Gao JQ, Wang P, Yan JW, Ba LN, Shi PL, Wu HM, et al. Shear Stress Rescued the
Neuronal Impairment Induced by Global Cerebral Ischemia Reperfusion via Activating PECAM-
1-eNOS-NO Pathway. Front Cell Dev Biol. 2020;8:631286.

121.   Albelda SM, Muller WA, Buck CA, Newman PJ. Molecular and cellular properties of
PECAM-1 (endoCAM/CD31): a novel vascular cell-cell adhesion molecule. J Cell Biol.
1991;114(5):1059-68.

122.   Sun QH, DeLisser HM, Zukowski MM, Paddock C, Albelda SM, Newman PJ. Individually
distinct Ig homology domains in PECAM-1 regulate homophilic binding and modulate receptor
affinity. J Biol Chem. 1996;271(19):11090-8.

123.   Sun J, Williams J, Yan HC, Amin KM, Albelda SM, DeLisser HM. Platelet endothelial cell
adhesion molecule-1 (PECAM-1) homophilic adhesion is mediated by immunoglobulin-like
domains 1 and 2 and depends on the cytoplasmic domain and the level of surface expression. J
Biol Chem. 1996;271(31):18561-70.

124.   Newton JP, Buckley CD, Jones EY, Simmons DL. Residues on both faces of the first
immunoglobulin fold contribute to homophilic binding sites of PECAM-1/CD31. J Biol Chem.
1997;272(33):20555-63.

125.   Privratsky JR, Newman PJ. PECAM-1: regulator of endothelial junctional integrity. Cell
Tissue Res. 2014;355(3):607-19.

126.   Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, et al. Size-selective loosening
of the blood-brain barrier in claudin-5-deficient mice. J Cell Biol. 2003;161(3):653-60.

127.   Yang Y, Rosenberg GA. MMP-mediated disruption of claudin-5 in the blood-brain barrier
of rat brain after cerebral ischemia. Methods Mol Biol. 2011;762:333-45.

128.   Huang L, Cao W, Deng Y, Zhu G, Han Y, Zeng H. Hypertonic saline alleviates
experimentally induced cerebral oedema through suppression of vascular endothelial growth
factor and its receptor VEGFR2 expression in astrocytes. BMC Neurosci. 2016;17(1):64.

1387

Figure 4.

A.

B. MMP-9

C. MMP-2

D. MMP-3

A.

CLUSTAL O(1.2.4) multiple sequence alignment

```
Isoform1     MTRARIGCFGPGGRARGTESAPEPSKRVPPGRSWQTQEVRQTRGANGLGPRAGSAGAKAP        60
Isoform2     ------------------------------------------------------------        0

Isoform1     GPAQGAAQHGLGGSAGLRVRVSPLAMGSAALEILGLVLCLVGWGGLILACGLPMWQVTAF       120
Isoform2                              MGSAALEILGLVLCLVGWGGLILACGLPMWQVTAF        35
                                      **********************************

Isoform1     LDHNIVTAQTTWKGLWMSCVVQSTGHMQCKVYDSVLALSTEVQAARALTVSAVLLAFVAL       180
Isoform2     LDHNIVTAQTTWKGLWMSCVVQSTGHMQCKVYDSVLALSTEVQAARALTVSAVLLAFVAL        95
             ***********************************************************

Isoform1     FVTLAGAQCTTCVAPGPAKARVALTGGVLYLFCGLLALVPLCWFANIVVREFYDPSVPVS       240
Isoform2     FVTLAGAQCTTCVAPGPAKARVALTGGVLYLFCGLLALVPLCWFANIVVREFYDPSVPVS       155
             ***********************************************************

Isoform1     QKYELGAALYIGWAATALLMVGGCLLCCGAWVCTGRPDLSFPVKYSAPRRPTATGDYDKK       300
Isoform2     QKYELGAALYIGWAATALLMVGGCLLCCGAWVCTGRPDLSFPVKYSAPRRPTATGDYDKK       215
             ***********************************************************

Isoform1     NYV       303
Isoform2     NYV       218
             ***
```

B. Isoform 1

C. Isoform 2

D. MMP-9 – Isoform 1

E. MMP-9 – Isoform 2