Discovery of NRG1-VII: a novel myeloid-derived class of NRG1 isoforms

Miguel Ángel Berrocal-Rubio[1], Yair David Joseph Pawer[1], Marija Dinevska[2], Ricardo De Paoli-Iseppi[1], Samuel S. Widodo[2], Josie Gleeson[1], Nadia Rajab[1], Will De Nardo[1], Jeannette Hallab[1], Anran Li[1], Theo Mantamadiotis[2,3], Michael B. Clark[1], Christine A. Wells[1#].

1. Department of Anatomy and Physiology, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Australia.
2. Department of Surgery, Royal Melbourne Hospital, The University of Melbourne, Australia.
3. Department of Microbiology and Immunology, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Australia

# Correspondence to CAW wells.c@unimelb.edu.au @Mincle (Twitter)

Key words: NRG1, myeloid, growth factor, isoforms, macrophages

**Abstract**

The growth factor Neuregulin-1 (NRG1) has pleiotropic roles in proliferation and differentiation of the stem cell niche in different tissues. It has been implicated in gut, brain and muscle development and repair. Six isoform classes of NRG1 and over 28 protein isoforms have been previously described. Here we report a new class of NRG1, designated NRG1-VII to denote that these NRG1 isoforms arise from a myeloid-specific transcriptional start site (TSS) previously uncharacterized. Long-read sequencing was used to identify eight high-confidence NRG1-VII transcripts. These transcripts presented major structural differences from one another, through the use of cassette exons and alternative stop codons. Expression of NRG1-VII was confirmed in primary human monocytes and tissue resident macrophages and iPSC-derived macrophages. Isoform switching via cassette exon usage and alternate polyadenylation was apparent during monocyte maturation and macrophage differentiation. NRG1-VII is the major class expressed by the myeloid lineage, including tissue-resident macrophages. Analysis of public gene expression data indicates that monocytes and macrophages are a primary source of NRG1, suggesting that NRG1-VII is the most common class of NRG1 in most adult human tissues, except brain. The size and structure of type VII isoforms suggests that they may be more diffusible through tissues than other NRG1 classes. However, the specific roles of type VII variants in tissue homeostasis and repair have not yet been determined.

## Introduction

Neuregulins (NRGs) are a family of highly pleiotropic growth factors derived from four paralogous genes (*NRG1-4*) (Figure 1A). NRGs are typically synthesized as transmembrane pro-peptides that are cleaved by metalloproteases in the extracellular space to form a bioactive peptide with an exposed epidermal growth factor-like (EGF) domain that can bind erythroblastic leukemia viral oncogene homolog (ERBB) receptors. The human *Neuregulin-1* (*NRG1*) locus, on Chromosome 8p12, generates numerous isoforms (Figure 1A, B) which are thought to be tissue-specific and functionally diverse (Falls, 2003). NRG1 has been implicated in the development of multiple tissues by promoting cell division within the stem cell niche and in differentiation trajectories (Yu et al., 2021, Wagner et al., 2007) including progenitor cells in the gut (Jardé et al., 2020), skeletal muscle (Gumà et al., 2010, Cheret et al., 2013) and cardiac cells (Wagner et al., 2007, Kramer et al., 1996), as well as nervous system development (Birchmeier 2009, Newbern et al., 2010). These studies demonstrate NRG1's key role in organogenesis and the importance of understanding how NRG1 isoforms exert tissue-specific effects to maintain the adult stem cell niche.

NRG1 and its receptors are involved in several diseases and targets for clinical research. Germline mutations in NRG1 are associated with developmental brain disorders such as schizophrenia (Craddock et al., 2005) as well as degenerative disorders such as amyotrophic lateral sclerosis (Sun et al., 2020) and Alzheimer's disease (Go et al., 2005). Constitutively activated isoforms of NRG1 are implicated in cancer, for which blocking the NRG1 isoform Heregulin or its receptors (ERBB) are effective clinical strategies against solid tumours (Sheng et al., 2010, Zhang et al., 2022). Deficiencies in NRG1 are associated with Hirshprung's disease, leading to poor innervation of the gut (Tang et al., 2012, Garcia-Barcelo et al., 2009), as well as abnormal brain development and mental disorders like bipolar disorder or schizophrenia (Georgieva et al., 2008, Marballi et al., 2012). In contrast, high levels of circulating NRG1 are associated with cardiac disease and morbidity following heart failure (Haller et al., 2022). NRG1 is also involved in modulating the immune response (Alizadeh et al., 2018, Ryzhov et al., 2017), controlling insulin related liver activity (Zhang et al., 2018), cell migration (Jumper et al., 2017) and cell-cell recognition and viability in the central nervous system (Garratt et al., 2000). Therefore, there are clear clinical benefits in understanding which cells express the different NRG1 isoforms in each tissue and how they play their tissue-specific roles in different biological processes.

The diversity of isoform functions in healthy development and disease across different organs is mirrored by the structural diversity and tissue specific transcriptional regulation of NRG1 products. The *NRG1* gene encodes at least 28 isoforms (Figure 1A) according to the ENTREZ Gene reference transcript list (Brown et al., 2015). Others have reported higher isoform heterogeneity from this locus (Mei 2008). This extraordinary transcript diversity is due to the use of alternate transcription initiation sites, cassette exon

usage, and alternate polyadenylation sequences. The locus is remarkably modular (Figure 1C), which

allows for different structural combinations of NRG1 domains depending on their start site (type I-VI), the

use of alternative linker domains, which are the protease targets for ectodomain shedding (1-4), the type

of EGF-like domain ($\alpha$, $\beta$ or $\gamma$), and whether the pro-peptide is membrane tethered or not (Figure 1C-E).

One well characterized isoform type, NRG1-III, contains a unique N-terminal domain that includes a

sequence that locates to the cell membrane (Nave et al., 2006). This means that upon NRG1 processing,

the growth factor domain will not be released from the source cell but will function as a juxtracrine signal

that allows neurons to establish key cell-cell interactions (with other neurons, oligodendrocytes, or muscle

fibers) and regulates the survival of both interacting cells (Garratt et al., 2000, Wolpowitz et al., 2000). In

rare circumstances, another cleavage site between the N-terminal domain and the EGF domain can also

be target by a protease, releasing the EGF domain to the ECM (Fleck et al., 2013).

**Figure 1. The human *NRG1* locus. A)** Screenshot of the NCBI RefSeq curated products from the *NRG1*
locus showing 28 protein coding transcriptional variants (NCBI Gene website accessed 26/1/2023). Left
and right labels correspond to transcript and protein accessions, respectively. **B)** Representation of the
NRG1 genomic locus (Chr 8p12). **C)** Schematic annotating exons in the human *NRG1* locus with modular
protein coding domains. **D)** Schematic showing combinatorial protein domains for previously annotated
NRG1 isoforms. Rows show the six major NRG1 classes, defined by alternate N-terminals. Columns
show alternate domains: domains within a column are mutually exclusive. Dotted lines show known
connections between domains in different NRG1 isoforms. Red stop symbols represent translation stop.
Red arrows represent a protease cleavage point. **E)** Representation of Neuregulin protein isoforms and
their domain distribution. Note that in NRG1 types IV, V and VI all isoforms that have been characterized
are transmembrane, while I, II and III also present isoforms that don't. Legend below the cell membrane
describes symbols for ADAM/BACE1 cleavage site (red arrow), glycosylation site (blue chain), kringle
domain (black V), IgG domain (purple circle), EGF domain (red oval), Transmembrane domain (orange
rectangle), intracellular domain (grey squiggle, with or without brown triangle). N-terminal domains of
NRG1 reference modularity shown in Fig. 1B and 1C, including the membrane-tethered N-terminal of
NRG1 type III.
106

More recently, NRG1 has been described as an important factor regulating the stem cell niche in the gut
(Jardé et al., 2020). In this context, NRG1 likely modulates stemness, proliferation and identity of
progenitor cells in the niche, and it is required to recapitulate certain secretory and absorptive functions in
human gut organoids (Kilik et al., 2021). Beyond development and tissue repair, NRG1 secreted by
macrophages has also been implicated in inflammation (Jardé et al., 2020, Garrido-Trigo et al., 2022).
However, the specific NRG1 isoforms expressed by macrophages in these contexts have not yet been
described.
115
Determining the sources and types of different NRG1 isoforms is an important part of understanding how
NRG1 directs these different developmental, reparative, and inflammatory outcomes. Here, data mining
to assess the expression profiles of *NRG1* led to the identification of a previously uncharacterized TSS
that appears to be used exclusively in cells of the myeloid lineage. We propose that transcripts generated
from this alternative TSS belong to a new NRG1 Class, NRG1-VII. Using Oxford Nanopore sequencing,
we identified eight class VII isoforms with distinct transcript structures and predicted the protein
characteristics of these isoforms. qRT-PCR targeting the unique first exon of *NRG1-VII* transcripts in
human cells confirmed that type VII isoforms are expressed by monocytes, infiltrating macrophages and
tissue resident macrophages. Immunohistochemistry using antibodies directed towards the EGF-like or
intracellular domains (ICD) demonstrated that tissue-resident macrophages are a major source of NRG1
in these tissues, an observation further supported by transcriptional evidence derived from single cell data
collated within the Human Protein Atlas. This study therefore contributes to untangling the complexity of
this already intricate locus by characterizing the structure and distribution of myeloid-specific NRG1
isoforms.
130
**Results**
NRG1-VII is defined by a novel TSS discovered in myeloid cells.
Since the different isoforms of NRG1 have varying structures and binding affinities to their receptors, we
sought to investigate which of these were expressed by myeloid cells. First, we used the FANTOM5
(Functional Annotation of the Mammalian Genome) database (The FANTOM Consortium; 2014), a
catalogue that maps the TSSs of genes expressed in 975 human samples (including cells, tissues, and
cell lines). Here, we found a previously uncharacterized TSS of NRG1 that was exclusively active in
myeloid cells, including monocytes, macrophages, and basophils (Figure 2A). A schematic of the locus
suggested that this represents a potential new class of *NRG1* transcripts which we prospectively named
*NRG1-VII* (Figure 2). The FANTOM TSS predicted that the *NRG1-VII* starting exon contained 139 base
pairs (bp) of mRNA sequence unique to transcripts originating from this site, and an in-frame methionine
was identified 115 bp downstream from the start of transcription (Figure 2B, C). Therefore, we aimed to
characterize whether this newly described TSS would lead to the expression of a new class of protein
coding transcripts.

The TSS for NRG1-VII initiates in the intron positioned 5' of ENSEMBL exon ENSE00003743434, which

encodes for part of the spacer sequence between the Ig and EGF-like domains in the canonical NRG1

protein (Figure 2B). The TSS adds a 5' untranslated region (UTR) to the transcript that extends the exon

from 51 bp to 190 bp (ENSE00002124728) and includes an initiating Methionine at nucleotide position

115-117 bp. This newly identified TSS generates a unique 5' UTR for its transcripts. Within this 5'-UTR, a

unique 20-nucleotide sequence was identified that allowed specific detection and amplification of the

NRG1-VII isoforms and primers were designed to target it (Figure 2C; Table 1). Isoforms arising from this

new TSS would present a unique sequence of 8 amino acids (MSYHLFFS), a type VII specific N-terminal

domain. This is followed by the 17 amino acids that are commonly present in this exon in other isoform

types (I, II, IV and V). This is the shortest known isoform specific N-terminal domain, and the only one that

is present within an exon that other isoforms may include.

We mapped the transcriptional activity arising from this TSS and identified 2 Expressed Sequence Tags

(ESTs) that had been sequenced from the 5' end (NCBI accessions BI908144 and BI907799) and

originated from the *NRG1-VII* TSS. These originated from the same library (SAMN00164230) made from

a pool of non-activated human leukocytes from anonymous donors. Both ESTs harbored evidence of an

open reading frame (ORF) in frame with other NRG1 isoforms, but both were 3' truncated. To further

validate the potential activity of this TSS we looked for evidence this isoform type was present in other

mammal species. Data available from 15 non-human primate species (Pipes et al., 2013), and cross

species alignment shows that the *NRG1-VII* TSS is present and highly specific to bone marrow and whole

blood, while isoforms isolated from other tissues use alternate TSSs (Sup. Figure 1A). We also found

transcriptional evidence of an equivalent TSS in *Mus musculus* and *Sus scrofa* derived from myeloid cells

(Sup. Figure 1B-C) confirming that the class VII TSS and its expression in myeloid cells are conserved

through evolution.

**Figure 2. Identification of a myeloid specific TSS in the human NRG1 locus. A)** Screenshot of the
ZENBU genome browser showing the Gencode v22 transcript track, (accessed 26/1/23) corresponding to
the FANTOM5 phase 1 and 2 TSS peaks overlapping the starting exon of NRG1 type VII. Red vertical
line encased in blue dashed box indicates the position for the TSS of interest. Tag selection (grey bar,
FANTOM5 CAGE track) indicates region for tag quantification in the FANTOM5 experiment table. The
table is truncated to show the top 6 samples are primary myeloid cells, representative of the top 57/115
samples with >1TPM) Experimental samples are ordered by highest number of CAGE tags counted in
this area on sense strand (green bars, showing counts as TPM). **B)** Schematic of the human *NRG1* locus
highlighting alternate transcriptional starting exons that correspond to seven isoform classes. The
schematic shows 5' exon composition until the first shared exon (containing the EGF domain). Exons are
numbered by ENSEMBL accessions. **C)** Alignment of mRNA and protein sequences of NRG1 type I and
type VII to show an in-frame translation initiation of NRG1 type VII. Initiating methionine (M) or start codon
(ATG) in red text. The blue arrow shows the target sequence of the forward primer for *NRG1-VII* mRNA's
unique 5' UTR. The *NRG1-VII* transcript start is conserved in other mammalian genomes (Sup. Figure 1).
186

*NRG1-VII* TSS transcribes at least 8 distinct high-confidence transcripts in myeloid cells.

To characterize the diversity of *NRG1* transcripts that use the *NRG1-VII* TSS, we performed Oxford

Nanopore long-read amplicon sequencing on *in vitro* and *in vivo* derived myeloid cells (Figure 3A-E). We

designed a forward primer that targets the 5' UTR of *NRG1-VII* transcripts, which is unique to this class of

NRG1 and does not overlap any other NRG1 isoform types; additionally, two reverse primers were

designed to target two of the known alternative transcriptional stops that we had previously validated as

active in myeloid cells through PCR (Figure 3C). We called amplicons "short" when the reverse primer

targeted exon "β" and "long" for the reverse primer in exon "α".

In total, we identified 8 novel high-confidence transcripts that were assigned at least 5% of the reads

present in a sample from either the short or long amplicons (Figure 3B, D), using the IsoLamp pipeline

(See Materials and Methods). Each amplicon was studied independently in *in vitro* progenitors and

differentiated macrophages, and in blood-isolated monocytes. Five of these transcripts had an open

reading frame (ORF) following a start codon (ATG), in frame with all other previously described NRG1

isoforms, and no premature stop codons in early exons. We predict that these five transcripts could be

protein coding (Figure 3B), including two short and three long isoforms. The three long isoforms would

include a transmembrane domain (NRG1-VII $\alpha$2a, $\alpha$2b and β2a) and likely undergo canonical processing

through metalloproteases for the EGF-containing peptide to be released. On the other hand, the two short

isoforms would lack the transmembrane and intracellular domains.

The short isoforms were detected in all myeloid libraries and lacked the transmembrane domain (Figure

3E). Of these, NRG1-VII $\alpha$β3 contained both $\alpha$ and β EGF-like exons, a combination that had been

previously captured in refseq NM_004495, which we now show was a truncated transcript as its TSS had

not been defined. This resolves the full-length sequence of the Class VII $\alpha$β isoforms. The presence of

both $\alpha$ and β domains in NRG1-VII $\alpha$β3 introduced a frame shift in the β exon, changing the peptide

sequence of this isoform to a unique C-terminal end. This feature might affect the binding dynamics of

these isoforms to ERBB receptors, making it of special interest.

Monocytes expressed both short isoforms, but only expressed one long isoform, NRG-VII $\alpha$2a. Two

additional isoforms were detected in the monocyte library that we predict are sensitive to nonsense

mediated decay (NMD). Transcripts NRG1-VII P$\alpha$ca and NRG1-VII Pβ3 include a previously

uncharacterized exon (Figure 3A) located in intron 2 of the other class VII transcripts. Additional support

for this exon can be found in isoforms present in smaller proportions like P- $\alpha$2a (Sup. Figure 2) This exon

spans 111 bases (chr8:32,676,084-32,676,195). We identified this as a poison exon, as it introduces an

amber stop codon (TAG), resulting in the following amino acid sequence:

MSYHLFFSEIITGMPASTEGAYVSSESPIRISVSTEGANTSSFITDECCHGGQYHNTAKSICLILMF-.

A third NMD candidate was identified in the iPSC-derived myeloid progenitors, NRG1-VII αβ2a. For

isoform αβ2a, this combination introduces an early stop codon that we theorize would lead the transcript

to nonsense mediated decay (NMD). All three NMD-candidate transcripts passed our high-confidence

transcript filters. We manually removed additional isoforms that passed our analysis threshold but had

less than 5% read coverage. These isoforms varied only in a few bases across a splice junction (Sup.

Figures 2-3) and were detected in only one library and were therefore considered as likely sequencing

artefacts. There are two possible exceptions: NRG1-VII β2b and P- α2a. Isoform β2b was detected in

both iPSC derived progenitors and iMACs, and P- α2a was found in monocytes, at levels between 1 and

5% (Sup. Figure 2).

**Figure 3. Sequence and structure of NRG1-VII isoforms A)** Representation of the different samples
analyzed in the study and the presence or absence of the identified high-confidence transcripts in each of
them (opaque = present, transparent = absent). **B)** Representation of the exon structure for 8 type VII
isoforms sequenced using Oxford Nanopore long-read sequencing and their designations based on exon
combinations according to the locus nomenclature. **C)** Human NRG1 locus reference with all known
exons. Arrows show target regions for the single forward primer and both reverse primers used in the
amplicon sequencing (See also Table 1). **D)** Predicted translation of protein sequences of the newly
characterized NRG1 type VII isoforms, compared to canonical NRG1-Iβ2a (top sequence). Text colour
indicates known protein motifs: purple Ig; red EGF; orange transmembrane domains. "/" represents
predicted pro-peptide cleavage points (green text represents ADAM/BACE1 proteolysis, blue represents
γ-secretase proteolysis). Dashed lines in all panels represent sequences that are not shown. **E)**
Schematic of predicted translated proteins for NRG1-VII isoforms compared to canonical NRG1-I (Left),
borrowing from domain schema shown in Figure 1. Alternate (α or β) EGF domains annotated in pink.
Intracellular triangles represent alternate a or b cytoplasmic tails. The yellow β domain represents the
peptide sequence that arises from the related exon but is translated to a different peptide sequence due
to a frame change.
252

<u>Macrophages are a major source of NRG1 in human tissue.</u>

We next assessed the distribution patterns of *NRG1* in single-cell RNA-seq experiments in the Human
Protein Atlas (Karlsson et al., 2021), which revealed that the *NRG1* locus was actively expressed in a
large variety of human tissues (Sup. Figure 4). Single cell expression data revealed NRG1 activity in
different cell types, such as neurons in the brain and eye; glandular stromal cells in colon, ovary, or
endometrium; endothelial cells in the heart and liver; and epithelial cells in the kidney and lung (Figure
4A). In most of these organs, macrophages are the main source of NRG1 (Figure 4A, Sup. Figure 4). One
outstanding exception is the brain, where neurons show the highest levels of NRG1 expression in the
human body. However, no isoform-specific single cell data is currently available to compare differential
isoform expression between the different macrophage types present across the tissues investigated.

<u>NRG1-VII expression in myeloid cells is affected by differentiation and maturation.</u>
We sought to confirm if isoform VII is the only or even primary TSS used by myeloid cells. Therefore,
using primers that could discriminate between each unique start exon (Table 1), we investigated the
patterns of the seven classes of NRG1 isoforms in different *in vivo* and *in vitro* myeloid cells (Figure 4B).
Expression of *NRG1-VII* was detected in all myeloid cells, and it was the isoform class showing highest
expression for cells that belong to this lineage (except for macrophages derived from a 2D epithelium). In
contrast, control cell types (hiPSCs and iPSC-derived cortical neurons) showed high expression of other
*NRG1* classes, but not *NRG1-VII.*

Having demonstrated that our iPSC derived myeloid cells can be used to model the use of this TSS, we
next investigated at which differentiation stage we could first detect the expression of NRG1-VII
transcripts. Thus, we investigated the NRG1-VII expression patterns during the differentiation process
towards the myeloid lineage, and its concurrence with specific myeloid markers (Figure 4C). Our results
show that NRG1-VII transcription in *in vitro* derived myeloid cells follows the upregulation of mature
monocyte markers like CD16, unlike what is observed in blood monocytes from *in vivo* samples (Figure
4C). This suggests that the sequence of events happening *in vivo* are not recapitulated in our *in vitro*
differentiation, and that a more mature myeloid identity is needed to activate transcription arising from the
*NRG1-VII* TSS in these cells.

<u>NRG1-VII expression in myeloid cells is affected by differentiation and maturation.</u>

To confirm that *NRG1* mRNAs are translated into proteins in myeloid cells, we then performed
immunohistochemical staining of NRG1 in human glioblastoma (GBM) tissue which is enriched in bone
marrow-derived macrophages (Klemm et al., 2020). Myeloid cells were identified using a CD68 antibody;
antibodies detecting the extracellular (EGF-like domain) and intracellular (ICD) domains of NRG1 were
used. NRG1 was detected in both myeloid and non-myeloid cells (Figure 4D-E). Not all CD68+ cells

expressed NRG1, and those cells that did exhibited different patterns of expression (Sup. Figure 2). Altogether, these results show that myeloid cells express NRG1 peptides in human tissues but are not the sole contributors to the NRG1 pool in the brain. They also reveal diverse NRG1 expression patterns in myeloid cells, which may be consistent with the different isoforms observed in our sequencing libraries, or consistent with active processing of membrane bound-NRG1.

**Figure 4. NRG1 expression profile in myeloid cells. A)** Quantification of NRG1 mRNA expression in different human tissues according to The Human Protein Atlas (Karlsson et al., 2021). A plethora of representative tissues that express NRG1 are shown here. The y axis represents nTPM (Number of transcripts per million). Neurons shown in yellow, myeloid cells in red, other cell types in grey. **B)** qRT-PCR data measuring presence of the different NRG1 classes in different cell types. Classes IV-VI were not detected in any of the samples. Expression ratios were calculated as described in Pfaffl, 2021. Mono= Monocyte, Mφ: macrophage, iPSC: Induced pluripotent stem cell. **C)** Time series showing qRT-PCR expression of typical myeloid markers in progenitor maturation and relationship with myeloid specific NRG1-VII transcripts. **D)** Immunostaining of GBM showing presence of CD68[+] cells that express NRG1 peptides *in vivo*. The left panel shows sample section stained with DAPI. The boxes indicate the regions shown in the right panel in D and the cell in E, respectively. Scale bar on left panel is 1 mm, scale bar on right panel represents 20 μm. **E)** Example of cell in which all markers (CD68, NRG1-EGF and NRG1-ICD) are expressed. Scale bars = 20 μm.

**Discussion**

Macrophages have been reported as a major source of NRG1 in different human tissues (Karlsson et al., 2021), including a recent report highlighting the importance of NRG1 in the developing gut (Jardé et al., 2020). Our investigation of *NRG1* expression in the public single cell atlases, including the Human Protein Atlas mRNA dataset, suggested that macrophages are a major source of NRG1 in most tissues. However, the specific isoforms involved in the macrophage mediated NRG1 secretion were uncharacterized. Here we report that myeloid cells preferentially use a novel TSS that is myeloid-specific and conserved which generates a previously uncharacterized class of NRG1 isoforms. The tissue variability and functional versatility of NRG1 are a consequence of the alternative promoter usage and exon retention that give rise to a high diversity of isoforms.

Transcripts arising from the human *NRG1* locus are differentially regulated by tissue type or developmental stage. Control over the six known isoform classes is achieved through differential proximal promoter usage at six unique TSSs, each of which is controlled by different transcription factors (Frensing et al., 2008). All available data indicates that the use of this TSS is exclusive to the myeloid lineage. The concrete mechanisms and transcription machinery involved in the process are yet to be described. Additional transcript processing can lead to alternative exon usage and define the functional modules in the protein, like the linker or the cytoplasmic tails (Figure 1 C, D), mechanisms that are tissue-specific (Wen et al., 1994). The diversity of NRG1 isoform classes and their specificity of expression in both a temporal anatomical and cell-specific manner, suggest distinct roles of different isoforms in tissue

patterning, especially in different brain regions (Liu et al., 2011). Further characterization of each of the
NRG1-VII isoforms is hence needed to elucidate their regulation and functional diversity.

It was previously reported that the highest NRG1 expression levels in human tissues was in blood plasma
(Pipes et al., 2013). This is unsurprising, as maturation of myeloid cells *in vivo* seems to cause the
downregulation of this gene across the monocyte-macrophage differentiation axis (Figure 2A), with a
subset of macrophages expressing NRG1 in tissues (Figure 4). Our review of the FANTOM and The
Human Protein Atlas suggests that monocytes are the main source of NRG1 in blood. The function of
monocyte-derived NRG1 has not yet been described, however, circulating levels of NRG1 correlate with
liver metabolic activity (Zhang et al., 2018) as well as post-infarct recovery and cardiovascular health
improvement (Mendes-Ferreira et al., 2013). This suggests that circulating NRG1 could play an important
role in hepatic and cardiac health.

We further found that monocytes express at least five high-confidence NRG1-VII isoforms, including
transcripts that contain a novel 'poison' exon (NRG1-VII P $\alpha$ca and P$\beta$3) predicted to prematurely
terminate translation (Figure 3A). We also note the unusual exon composition of isoform NRG1-VII P$\alpha$ca,
indicating it could be a PCR artifact; thus, further validation might be required on this isoform, despite its
amplicon constituting over 15% of the detected expression. Additional isoforms containing the monocyte
poison exon like NRG1-VII P$\alpha$2a were also found, but in much lower proportions (Sup. Figure 2).
However, this supports the use of the poison exon by monocytes. This exon introduces an early stop
codon and hence is likely to drive the transcript to nonsense mediated decay, generating a monocyte
specific transcriptional mechanism to regulate the levels of NRG1 synthesis. Due to the high levels of
expression of this gene in monocytes compared to other cell types, we hypothesize that these represent
regulatory mechanisms that allow control over the levels of NRG1 expression in circulation.

Neuregulins interact with monomeric ERBB proteins (ERBB1-4), via an EGF-like domain, promoting
dimerization and trans-phosphorylation of these receptors. Each ERBB monomer can bind different EGF
members with differential ligand binding affinities, but only ERBB 3 and 4 can bind Neuregulins. Though
NRG1 and NRG2 can bind to both receptors as monomers (Carraway III et al., 1997), NRG3 and NRG4
only bind ERBB4 monomers (Harari et al., 1999, Zhang et al., 1997). Dimerized receptors can
discriminate between different isoforms of NRG1, which is evidenced by differential phosphorylation on
receptor tyrosine residues (Pinkas-Kramarski et al., 1998, Sweeney et al., 2000). Dimerization leads to
the recruitment of different adaptor proteins, notably GRB or PI3K, which then trigger specific secondary
cascades and regulate cell activity (Buonanno et al., 2001). NRG1-VII isoforms present a very short N-
terminal domain compared to most other isoforms. It has been previously reported that the size and
structure of the NRG1 N-terminal domain can dictate the receptor availability and recycling, affecting
internal phosphorylation and internal signaling in the receiving cell (Warrant et al., 2006). Additionally, the

discovery of isoform NRG1-VII αβ3 as a coding isoform, adds a novel sequence possibility in the EGF

domain. This is likely to determine the protein's binding dynamics and internal phosphorylation, hence

changing its functional properties. Functional validation is necessary to confirm whether this isoform type

has unique properties given its EGF domain structure.

NRG1 is involved in many different functions, attributed to the many isoforms which are tissue-restricted

during different developmental stages. Further, while NRG1 is conserved at the protein level, the function

of specific isoforms has been shown to differ between species. For example, heart trabeculation failure,

which leads to a lethal phenotype in mice at day 10.5, is a phenotype common to NRG1-/-, ERBB2-/- and

ERBB4-/- mice (Kramer et al., 1996, Britsch et al., 1998). Although 11 different isoforms have been

identified in the heart at this developmental stage, the phenotype can be attributed specifically to the

absence of β-RGF types I and II, which are the isoforms containing the immunoglobulin (Ig) domain

(Pentassuglia et al., 2009). However, the opposite is observed in zebrafish where Ig-like neuregulin-1

isoforms are dispensable for this process (Samsa et al., 2016). Thus, while essential processes in

development can be attributed to specific subtypes of NRG1, differences between species showcase the

need to characterize the different human isoforms using appropriate cell type and developmental models.

NRG1 and the ERBB receptors are important clinical target in different cancer types. For example, NRG1

fusion proteins cause pathological activation of ERBB receptors (Laskin et al., 2020). NRG1 expression

correlates with a shorter survival in patients with glioma (Yin et al., 2015), which could be related to its

role in cell migration and proliferation. Glioblastomas particularly exhibit high macrophage infiltration (Wei

et al., 2019). Our results show that tumor-associated myeloid cells (CD68[+]) may contribute to the NRG1

pool seen in these tumors. Due to the lack of unique domains in type VII isoforms, the antibodies used in

this study targeted common NRG1 regions (the EGF-like and intracellular domains). Even though the

available antibodies lack specificity to prove that type VII isoforms are translated in this disease, the only

isoforms detected in primary monocytes or macrophages in this study belong to classes I and VII.

While NRG1 double positive myeloid cells may express newly synthesized pro-peptides, we also

observed CD68[+]/NRG1-ICD[+] cells indicating that the EGF-like domain had been cleaved. Alternatively,

the presence of CD68[+]/NRG1-EGF[+] cells suggests that some cells restrict their expression only to

isoforms that end transcription in linker 3 and lack a transmembrane domain. Therefore, we confirmed

that macrophages *in vivo* express NRG1 and that there are distinct populations of macrophages in tissue

based on their NRG1 expression patterns.

To determine whether *in vitro* derived myeloid cells can be used to model the activity of the NRG1-VII

TSS, we used primers designed to target specific TSS usage in different cell types. We confirmed that

NRG1-VII was expressed in all myeloid samples and in no others, suggesting the TSS is active. However,

iPSC-derived samples showed expression of NRG1-III. This could be a result of incomplete differentiation
of the culture, or retention of stem cell features in *in vitro* derived myeloid cells. Moreover, the time series
data obtained during the differentiation process shows that NRG1 expression follows mature markers like
CD16; *in vivo*, NRG1 expression precedes CD16, and as CD16 is upregulated, NRG1 expression
decreases. Hence, while we show activity of the TSS, the *in vitro* model may not recapitulate the
transcriptional and maturation sequence seen *in vivo* due to the differences in the differentiation process.
413
Description of a new TSS class, NRG1-VII, including at least five new protein-coding isoforms, has
expanded the known NRG1 protein coding isoforms from 28 to 33. This study adds eight new transcripts
that are specific to myeloid cells. However, it is likely that the full transcriptional profile of this locus has
not yet been described; for example, while long-read amplicon sequencing is a sensitive isoform recovery
method (Clark et al., 2020), it is limited by the primer set(s) used. Thus, only NRG1-VII isoforms utilizing
linker 3 or the "a tail" regions were amplifiable in this study. It is highly likely that NRG1-VII isoforms using
the 3' end present in exon c3 (ENSE00002109887) also exist, which should be validated with additional
work.
422
Here we showed that myeloid cells exhibit a unique regulation pattern of the NRG1 locus to generate cell
specific isoforms, potentially playing an important role in diverse diseases. Only through a thorough
investigation of this locus can we better understand each process and develop clinical strategies to
prevent or treat the different pathologies associated with this locus. Further detailed investigation on the
molecular genetic features and functions of these novel isoforms might uncover how NRG1-VII isoforms
elicit differential receptor activity and downstream effects as previously described for other NRG1
isoforms. This could lead to targeted therapies and an improved understanding of the complexity of the *in*
*vivo* system, helping us recreate the processes in which appropriate signals are essential to model the
desired biological mechanisms.
432
433  **Materials and methods**
434
Cell lines and ethics approvals.
Stem cell work was performed in accordance with The University of Melbourne ethics committee HREC
(approval 1851831). The line of human iPSCs used was: PB001.1 (Vlahos et al., 2019), obtained from the
Stem Cell Core Facility at the Murdoch Children's Research Institute. Kolf2.1 (hPSCReg accession
WTSIi018-B; Welcome Trust Sanger Institute) cells were used to differentiate cortical neurons (Ethics ID:
12374) and 2D macrophages. Monocytes were isolated from buffy coat, which was obtained from the
Australian Red Cross Blood Service in accordance with The University of Melbourne ethics committee
HREC (approval 1646608). Ethics for adipose tissue derived samples was obtained from the University of
Melbourne Human Ethics Committee (ethics ID 1851533) and approved by The Avenue Hospital Human
Research Ethics Committee (Ramsay Health; ethics ID WD00006, HREC reference number

2019/ETH/0050). For human glioblastoma samples, human ethics approval was covered by project
application 1853511, approved by the Medicine and Dentistry Human Ethics Sub-Committee, The
University of Melbourne.

<u>Stem cell culture.</u>
Cells were cultured in Gibco™ Essential 8™ media with Essential 8™ supplement (Thermo Fisher
Scientific; A1517001) on growth-factor reduced Matrigel® Matrix (Corning®; 356234) coated dishes. Cells
were cultured with daily changed fresh media. Cell culture was performed in an APT.line™ C150 (E2)
$CO_2$ manual incubator (BINDER; 7001-0172) in constant and stable conditions of humidity, temperature
(37°C) and $CO_2$ concentration in air (5%).

Cell passaging was performed routinely when cell confluency reached (70-80%). Cells were first washed
with Gibco™ PBS (Thermo Fisher Scientific; 10010023). Then, a dilution of sterile 0.5 M EDTA (Thermo
Fisher Scientific; 15575020) in said PBS at a concentration of 0.5 mM was used to detach the cells from
the plate. After 3-4 minutes in humid incubator conditions, cells were collected and replated with fresh
media.

<u>3D iPSC derived macrophages.</u>
As described in Rajab et al., 2021. Human iPSCs were differentiated into macrophages following (Joshi et
al., 2019), but with the following alterations to the protocol: harvested cells were cultured in MAGIC
media, which was changed as indicated in (Ng et al., 2008). During this process, cells were plated in
10cm Ø non-treated Petri dishes (IWAKI; 1020-100) and placed on an orbital shaker (N-Biotek orbital
shaker NB-T101SRC) in a humidified incubator with 5% $CO_2$ at 37°C.

After 11 days of culture, cells can be observed detaching from the embryoid bodies, remaining in
suspension in the media as non-adherent cells (which are characterized as myeloid progenitors). When
all cells are collected and allowed to settle in a 15 mL Falcon tube (Corning®; CLS431470-500EA),
embryoid bodies pelleted in the bottom, but progenitors stayed in suspension in the supernatant, and
could then be collected. The supernatant was then centrifuged (Heraeus Multifuge 1S-R) at 400 rpm for 5
minutes to separate the progenitors from the media. Progenitors were then resuspended in a 10% dilution
in volume of FBS and 100ng/mL CSF-1 (R&D Systems; 216-MC-500) in Gibco™ RPMI-1640 media
(Thermo Fisher Scientific; 11875093). Cells were plated in Costar® 6-well tissue-culture treated plates
(Corning®; 3516) for 4-7 days in stable incubator conditions (humid, 5% $CO_2$, 37°C), when cells showed
morphological and molecular features displayed by macrophages.

<u>iPSC derived cortical neurons</u>

Kolf2.1 hiPSCs (Kao et al., 2016) were cultured under xenogenic conditions as defined in Niclis et al.

(2017). The cells were then differentiated into cortical neurons following the protocol described by

(Gantner et al., 2021).

<u>Human samples and cell sorting.</u>

*Blood monocyte isolation.*

Buffy Coat was obtained from the Australian Red Cross Blood Service. The blood was diluted with PBS at

a 1:3 dilution and underlaid with Histopaque®-1077 (Sigma-Aldrich; Cat. No. 10771-100ml). The

underlaid blood samples were centrifuged (TECHCOMP CT1SRT) at 350g for 30 minutes at 24°C with no

brake. Peripheral blood mononuclear cells (PBMCs) were isolated from the interphase and washed twice

using MACs buffer (Gibco™ Dulbecco's phosphate-buffered saline (DPBS) (Ca2+Mg2+ free) (Thermo

Fisher Scientific; Cat. No. 14190144) with 0.5% heat inactivated Fetal Bovine Serum (FBS) (Thermo

Fisher Scientific; Cat. No. 10082147 or 10099141) and 2mM EDTA (Invitrogen™ UltraPure™ 0.5M EDTA

(Thermo Fisher Scientific; Cat. No. 15575020)) and centrifuging at 400g for 5 minutes at 4°C. Cell count

and viability were determined using 0.4% Gibco™ Trypan Blue (Thermo Fisher Scientific; Cat. No.

15250061) using a hemocytometer. Cells were centrifuged at 400g for 5 minutes at 4°C and resuspended

in 40μl MACs buffer per $10^7$ cells. Monocytes were positively selected by a magnetic field using Human

CD14 MicroBeads (Miltenyi Biotec; Cat. No. 130-050-201) and LS Columns (Miltenyi Biotec; Cat. No.

130-042-401).

*Adipose tissue samples.*

Adipose tissue was obtained and processed as described in (Raajendiran et al., 2019). Myeloid cells

were selected by FACS using a BD FACSAria™ III system (BD Biosciences). Cells that were positive for

markers CD45 and CD11b were isolated for this study.

<u>RNA extractions and cDNA synthesis.</u>

Total RNA was extracted using the RNeasy® Plus Mini Kit (Qiagen; 74134) according to manufacturer's

instructions. After final total RNA elution in RNase-free water, overall RNA quality and concentration were

measured using an RNA ScreenTape (Agilent Technologies; 5067-5576) in an Agilent 2200 Tapestation

System (Agilent Technologies; G2964-90003).

cDNA synthesis was then performed using the isolated total RNA and considering the concentration

values assigned to each of the samples for the coming steps. cDNA was synthesized using

SensiFASTTM cDNA Synthesis Kit (BioLine; BIO-65053) and following all protocol specifications from the

vendor. cDNA concentration and quality were checked using a D5000 ScreenTape (Agilent Technologies;

5067-5588) in an Agilent 2200 Tapestation System. Final cDNA samples were then stored at -20°C.

q-RT PCR.

For mRNA quantification, the Applied Biosystems Viia7 ™ real time system was used (Thermo Fisher
Scientific; 4453536) using a Fast 96 well plate hardware set up. The reactions in each well of the Micro
Amp Fast 96 well reaction plate 0.1 mL (Thermo Fisher Scientific; 4346907) were prepared as indicated
for the Fast SYBR™ Green Master Mix (Thermo Fisher Scientific; 4385612). Primers used in the reaction
were designed for each transcript class of interest (Table 1). Plates were then sealed using Optical
adhesive covers (Thermo Fisher Scientific; 436 0954). For quantification, n=3 technical replicates were
used.

Relative expression ratio was calculated as E(HKG)Average HKG Ct/ E(GOI)Average GOI Ct as
described in Pfaffle, 2021. Efficiencies for each primer pair were calculated from serial dilutions of
template and ranged between 95 and 98 % for all NRG1 isoforms classes reported in results. Isoforms IV,
V, VI were not detected in any samples. B2M was used as House Keeping gene (HKG) for all samples.

| Gene | F-Primer | Tm °C | R-primer | Tm °C |
|---|---|---|---|---|
| B2M | TAGCTGTGCTCGCGCTACT | 66.5 | TTCAATGTCGGATGGATGAA | 60.2 |
| NRG1 I | CAAAGAAGGCAGAGGCAAAG | 62.9 | AACTGGTTTCACACCGAAGG | 63.9 |
| NRG1 II | AACCTCAAGAAGGAGGTCAGC | 63.9 | AACTGGTTTCACACCGAAGG | 63.9 |
| NRG1 III | CCGACACCGAAGAATCGTAT | 63.8 | ACTCCCCTCCATTCACACAG | 64.0 |
| NRG1 IV | GCGACAGAGAGGGAGGA | 63.6 | AACTGGTTTCACACCGAAGG | 63.9 |
| NRG1 V | AATTCTTCTACGGAGTTTTAAGGTACAC | 62.6 | GCCGATTCCTGGCTTTTCAT | 67.2 |
| NRG1 VI | TCTTCAGGAACCACCTAAGCA | 63.6 | TCTCCTTCTCCGCACATTTT | 63.6 |
| NRG1 VII | TTAGTTCCAAGGCAGGCTGT | 63.7 | TTGCTCCTTCTGTGGATACTGA | 63.7 |
| CD11b | AGAACAACATGCCCAGAACC | 63.9 | GCGGTCCCATATGACAGTCT | 63.9 |
| CD14 | GCCGCTGTGTAGGAAAGAAG | 63.8 | ATCGTCCAGCTCACAAGGTT | 63.7 |
| CD16 | TGAGGTGTCACAGCTGGAAG | 64.3 | GGTTGACACTGCCAAACCTT | 63.9 |
| CD43 | AGTGCTGCGTCCTTATCAGC | 64.4 | CAAACAGGCAGGAGCAAGAG | 65.0 |
| CD45 | AGAATACTGGCCGTCAATGG | 63.8 | GCTGAAGGCATTCACTCTCC | 63.9 |
| SPL1 | CCAGCTCAGATGAGGAGGAG | 64.2 | CAGGTCCAACAGGAACTGGT | 64.0 |
| NRG1-VII Nanopore | Forward amplicon: TTTCTGTTGGTGCTGATATTGC TTAGTTCCAAGGCAGGCTGT | 84 | Reverse short amplicon: ACTTGCCTGTCGCTCTATCTTCG ATGCAGCAACAAGAAAGCA<br><br>Reverse long amplicon: ACTTGCCTGTCGCTCTATCTTCTT TCCTGTTTTCTATTTGCAGAAC | 79.4<br><br>83.9 |

**Table 1.** Primers designed for qRT-PCR and Nanopore Long Read Amplicon Sequencing. Note that
underlined regions of the primers correspond to the general ONT 5'mod sequence.

Nanopore Amplicon sequencing.

NRG1-VII was amplified using the LongAmp® Taq 2X Master Mix (New England Biolabs; Cat #: M0287S)
and the specified primers (Table 1) for either 30 cycled for the shorter amplicon or 40 cycles for the longer
amplicon. The samples used were a pool of monocytes from 3 different blood donors, a pool of myeloid
progenitors derived *in vitro* and macrophages differentiated from said *in vitro* derived progenitors. Target
sequences were then amplified using specific primers (Table 1) and purified using AMPure beads
(Beckman Coulter; A63880) at concentrations appropriate to each of the target sizes. Then, 2 ng of
purified cDNA from each sample were barcoded following the EXP-PBC096 protocol from Oxford
Nanopore Technologies. Samples were pooled (equimolar) and a sequencing library was prepared as
described in the SQK-LSK110 Oxford Nanopore protocol. Samples were then loaded on a Flongle flow
cell (FLO-FLG001) and sequenced using a GridION device.

Sequencing data analysis.
Basecalling was performed using Guppy (v6.3.8) using the super-accurate basecalling config file with a
Qscore        threshold      of       10.      To       identify      NRG1       Isoforms       we       used       IsoLamp       v1.0
(https://github.com/ClarkLaboratory/IsoLamp), a bash pipeline for the identification of known and novel
isoforms from targeted amplicon long-read sequencing data generated with Oxford Nanopore
technologies. Briefly, IsoLamp takes passed Nanopore reads and maps them to a reference genome
using Minimap2 (Li, 2018). Alignments that are highly accurate (>95%), are full-length and have high
accuracy splice junctions (>90%) are used for isoform discovery with bambu (v3.2.4). Isoform
quantification is then performed with Salmon v0.14.2 (ref). IsoLamp was run in de novo mode setting
BambuAnnotations=NULL. Each sample was run independently through the IsoLamp pipeline with
'downsample reads' parameter set to FALSE. All other parameters were set to default. Isoform annotation
files and count matrixes were visualized using IsoVis (https://isomix.org/isovis).

Automated multiplex immunohistochemistry.
Formalin-fixed paraffin-embedded (FFPE) GBM tissue sections were stained using the Bond RX
automated stainer (Leica Biosystems). Slides were deparaffinized in xylene followed by exposure to a
graded series of ethanol solutions for rehydration. Heat-induced epitope retrieval was performed with
either a Citrate pH 6 buffer or Tris Ethylenediaminietetraacetic acid (EDTA) pH 9 buffer. Slides were
blocked with 3% hydrogen peroxide ($H_2O_2$) to block endogenous peroxidase activity. For multiplexed IHC
staining the Opal 6-plex Detection Kit (Akoya Biosciences) was used. Serial multiplexing was performed
by repeating the sequence of antigen retrieval, primary antibody, and Opal polymer incubation, followed
by Opal fluorophore visualisation for all antibodies as follows. GBM tissue was stained with CD68 (Abcam
ab955, 1:100), NRG1 ICD (Abcam ab191139, 1:200) and NRG1 EGF-like domain (ThermoFisher; MA4-
12896, 1:100). Slides were incubated for 1 hour at room temperature with primary antibodies diluted in 1x
Opal blocking/antibody diluent (Akoya biosciences). Slides were subsequently incubated with the Opal
Polymer HRP Ms + Rb secondary polymer for 30 minutes prior to incubation with Opal fluorophores (Opal

520, 540, 570, 620, 650 and 690) diluted at 1:150 in 1x Plus Automation Amplification Diluent (Akoya

biosciences) for 10 minutes. Slides were counterstained with 10x Spectral DAPI and coverslipped with

ProLong Glass Antifade Mountant (Invitrogen). Multispectral images were acquired at 20x and 40x

magnification using PhenoImager ™ HT (Akoya Biosciences). inForm 2.4.8 software (Akoya Biosciences)

was used for spectral deconvolution. Deconvoluted multispectral images were subsequently fused in

HALO (Indica Labs).

**Supplementary data**

A

B

C

583
584
585

**Supplementary Figure 1. NRG1-VII TSS shows expression conservation in primate and non-**
**primate mammals. A)** NRG1 expression profiles of human and non-human primates showing that TSS
VII is conserved in bone marrow and whole blood **B)** Curated transcript in *Mus musculus* databases
showing conservation of TSS VII in blood cells. **C)** Predicted transcript in *Sus scrofa* based ETSs derived
from dendritic cells and other myeloid progenitor samples. Screenshot from https://www.ncbi.nlm.nih.gov1.
*Accessed on 15/12/2022.*

594
595 **Supplementary Figure 2. NRG1 long amplicon isoforms found through IsoLamp.** Each panel shows
596 the IsoVis (Ref) view for NRG1 amplicons in a different sample. The initial row shows the canonical
597 isoform structure chosen by Isomix. The rest show the resulting isoforms found by the IsoLamp analysis
598 pipeline. Isoforms that include the term "[alt]" show isoforms that are only different to the main isoform in
599 one of the splicing junctions by a few bases. The heat maps on the right-hand side describe relative

abundance of each transcript in that sample in nTPM. Panels correspond to **A)** iPSC derived myeloid progenitors; **B)** iPSC derived macrophages (iMACs); **C)** Monocytes.

**Supplementary Figure 3. NRG1 short amplicon isoforms found through IsoLamp.** Each panel shows the IsoVis (Ref) view for NRG1 amplicons in a different sample. The initial row shows the canonical isoform structure chosen by Isomix. The rest show the resulting isoforms found by the IsoLamp analysis pipeline. Isoforms that include the term "[alt]" show isoforms that are only different to the main isoform in one of the splicing junctions by a few bases. The heat maps on the right-hand side describe relative abundance of each transcript in that sample in nTPM. Panels correspond to **A)** iPSC derived myeloid progenitors; **B)** iPSC derived macrophages (iMACs); **C)** Monocytes. Different poison exons were detected for isoform NRG1-VII Pβ3 in each sample type, therefore further validation of these exons and isoform(s) is needed.

624
625
626

627
628
629 **Supplementary Figure 4. NRG1 expression levels in macrophages presen in a wide variety of**
630 **human tissues.** Quantification of NRG1 mRNA expression in macrophages present different human
631 tissues according to The Human Protein Atlas (Karlsson et al., 2021). The y axis shows the number of
632 transcripts per million. Yellow bars = neural tissue; Grey scale bars = levels of expression in all other
633 tissues.

656
657
658 **Supplementary Figure 5. Panel of marker combinations present in GBM samples.** CD68⁺ cells are
659 diverse in their NRG1 expression patterns. Panels show CD68⁺ cells that can be NRG1⁻, or positive for
660 either of the NRG1 antibodies independently. CD68⁻ cells were also observed to be positive for presence
661 NRG1. Scale bars = 20 μm.

## Availability of data and materials

665
FAST5 and BAM files are available from ENA under accession PRJEB62796
(https://www.ebi.ac.uk/ena/browser/home ). The different NRG1-VII isoform sequences and expected
proteins are available at NCBI under submission numbers OQ272754-OQ272776
(https://www.ncbi.nlm.nih.gov/ ). The processed data and scripts used in this study are available at
https://github.com/wellslab/NRG1-VII_Amplicons . IsoLamp is available on github at
https://github.com/ClarkLaboratory/IsoLamp.

**REFERENCES:**

Alizadeh, A., Santhosh, K.T., Kataria, H., Gounni, A.S. & Karimi-Abdolrezaee, S. 2018, "Neuregulin-1 elicits a regulatory immune response following traumatic spinal cord injury", *Journal of neuroinflammation,* vol. 15, no. 1, pp. 1-21.

Birchmeier, C. 2009, "ErbB receptors and the development of the nervous system", *Experimental cell research,* vol. 315, no. 4, pp. 611-618.

Britsch, S., Li, L., Kirchhoff, S., Theuring, F., Brinkmann, V., Birchmeier, C. & Riethmacher, D. 1998, "The ErbB2 and ErbB3 receptors and their ligand, neuregulin-1, are essential for development of the sympathetic nervous system", *Genes & development,* vol. 12, no. 12, pp. 1825-1836.

Brown, G.R., Hem, V., Katz, K.S., Ovetsky, M., Wallin, C., Ermolaeva, O., Tolstoy, I., Tatusova, T., Pruitt, K.D. & Maglott, D.R. 2015, "Gene: a gene-centered information resource at NCBI", Nucleic acids research, vol. 43, no. D1, pp. D36-D42.

Buonanno, A. & Fischbach, G.D. 2001, "Neuregulin and ErbB receptor signaling pathways in the nervous system", Current opinion in neurobiology, vol. 11, no. 3, pp. 287-296.

Carraway III, K.L., Weber, J.L., Unger, M.J., Ledesma, J., Yu, N., Gassmann, M. & Lai, C. 1997, "Neuregulin-2, a new ligand of ErbB3/ErbB4-receptor tyrosine kinases", Nature, vol. 387, no. 6632, pp. 512-516.

Cheret, C., Willem, M., Fricker, F.R., Wende, H., Wulf-Goldenberg, A., Tahirovic, S., Nave, K., Saftig, P., Haass, C. & Garratt, A.N. 2013, "Bace1 and Neuregulin-1 cooperate to control formation and maintenance of muscle spindles", *The EMBO journal,* vol. 32, no. 14, pp. 2015-2028.

Clark, M.B., Wrzesinski, T., Garcia, A.B., Hall, N.A., Kleinman, J.E., Hyde, T., Weinberger, D.R., Harrison, P.J., Haerty, W. & Tunbridge, E.M. 2020, "Long-read sequencing reveals the complex splicing profile of the psychiatric risk gene CACNA1C in human brain", *Molecular psychiatry,* vol. 25, no. 1, pp. 37-47.

Craddock, N., O'Donovan, M.C. & Owen, M.J. 2005, "The genetics of schizophrenia and bipolar disorder: dissecting psychosis", *Journal of medical genetics,* vol. 42, no. 3, pp. 193-204.

Falls, D.L. 2003, "Neuregulins: functions, forms, and signaling strategies", *The EGF Receptor Family*, pp. 15-31.

FANTOM Consortium and the RIKEN PMI and CLST (DGT). A promoter-level mammalian expression atlas. Nature 507, 462–470 (2014).

Frensing, T., Kaltschmidt, C. & Schmitt-John, T. 2008, "Characterization of a neuregulin-1 gene promoter: Positive regulation of type I isoforms by NF-κB", *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms,* vol. 1779, no. 2, pp. 139-144.

Fleck, D., van Bebber, F., Colombo, A., Galante, C., Schwenk, B.M., Rabe, L., Hampel, H., Novak, B., Kremmer, E., Tahirovic, S. and Edbauer, D., 2013. Dual cleavage of neuregulin 1 type III by BACE1 and ADAM17 liberates its EGF-like domain and allows paracrine signaling. *Journal of Neuroscience*, *33*(18), pp.7856-7869.

Gantner, C.W., Hunt, C.P., Niclis, J.C., Penna, V., McDougall, S.J., Thompson, L.H. & Parish, C.L. 2021, "FGF-MAPK signaling regulates human deep-layer corticogenesis", *Stem cell reports,* vol. 16, no. 5, pp. 1262-1275.

Garcia-Barcelo, M., Tang, C.S., Ngan, E.S., Lui, V.C., Chen, Y., So, M., Leon, T.Y., Miao, X., Shum, C.K. & Liu, F. 2009, "Genome-wide association study identifies NRG1 as a susceptibility locus for Hirschsprung's disease", *Proceedings of the National Academy of Sciences,* vol. 106, no. 8, pp. 2694-2699.

Garratt, A.N., Britsch, S. & Birchmeier, C. 2000, "Neuregulin, a factor with many functions in the life of a schwann cell", *Bioessays,* vol. 22, no. 11, pp. 987-996.

Garrido-Trigo, A., Corraliza, A., Veny, M., Dotti, I., Melon-Ardanaz, E., Rill, A., Crowell, H.L., Corbi, A., Gudino, V. & Esteller, M. 2022, "Macrophage and neutrophil heterogeneity at single-cell spatial resolution in inflammatory bowel disease", bioRxiv.

Georgieva, L., Dimitrova, A., Ivanov, D., Nikolov, I., Williams, N.M., Grozeva, D., Zaharieva, I., Toncheva, D., Owen, M.J. & Kirov, G. 2008, "Support for neuregulin 1 as a susceptibility gene for bipolar disorder and schizophrenia", *Biological psychiatry,* vol. 64, no. 5, pp. 419-427.

Go, R.C., Perry, R.T., Wiener, H., Bassett, S.S., Blacker, D., Devlin, B. & Sweet, R.A. 2005, "Neuregulin-1 polymorphism in late onset Alzheimer's disease families with psychoses", *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics,* vol. 139, no. 1, pp. 28-32.

Gumà, A., Martínez-Redondo, V., López-Soldado, I., Cantó, C. & Zorzano, A. 2010, "Emerging role of neuregulin as a modulator of muscle metabolism", *American Journal of Physiology-Endocrinology and Metabolism,* vol. 298, no. 4, pp. E742-E750.

Haller, P.M., Gonçalves, I.F., Acar, E., Jäger, B., Pilz, P.M., Wojta, J., Huber, K., Kiss, A. & Podesser, B.K. 2022, "Relationship between plasma Neuregulin-1 and cardiac function in patients with ST-elevation myocardial infarction", *Reviews in* Cardiovascular Medicine, vol. 23, no. 2, pp. 63.

Harari, D., Tzahar, E., Romano, J., Shelly, M., Pierce, J., Andrews, G. & Yarden, Y. 1999, "Neuregulin-4: a novel growth factor that acts through the ErbB-4 receptor tyrosine kinase", Oncogene, vol. 18, no. 17, pp. 2681-2689.

Jardé, T., Chan, W.H., Rossello, F.J., Kahlon, T.K., Theocharous, M., Arackal, T.K., Flores, T., Giraud, M., Richards, E. & Chan, E. 2020b, "Mesenchymal niche-derived neuregulin-1 drives intestinal stem cell proliferation and regeneration of damaged epithelium", *Cell Stem Cell,* vol. 27, no. 4, pp. 646-662. e7.

Joshi, K., Elso, C., Motazedian, A., Labonne, T., Schiesser, J.V., Cameron, F., Mannering, S.I., Elefanty, A.G. & Stanley, E.G. 2019, "Induced pluripotent stem cell macrophages present antigen to proinsulin-specific T cell receptors from donor-matched islet-infiltrating T cells in type 1 diabetes", *Diabetologia,* vol. 62, no. 12, pp. 2245-2251.

Jumper, N., Hodgkinson, T., Paus, R. & Bayat, A. 2017, "A Role for Neuregulin-1 in Promoting Keloid Fibroblast Migration via ErbB2-mediated Signaling.", *Acta Dermato-Venereologica,* vol. 97, no. 6.

Kao, T., Labonne, T., Niclis, J.C., Chaurasia, R., Lokmic, Z., Qian, E., Bruveris, F.F., Howden, S.E., Motazedian, A. & Schiesser, J.V. 2016, "GAPTrap: a simple expression system for pluripotent stem cells and their derivatives", *Stem Cell Reports,* vol. 7, no. 3, pp. 518-526.

Karlsson, M., Zhang, C., Méar, L., Zhong, W., Digre, A., Katona, B., Sjöstedt, E., Butler, L., Odeberg, J. & Dusart, P. 2021a, "A single–cell type transcriptomics map of human tissues", *Science Advances,* vol. 7, no. 31, pp. eabh2169.

Kilik, U., Yu, Q., Holtackers, R., Seimiya, M., dos Santos, Aline Xavier da Silveira, Treutlein, B., Spence, J.R. & Camp, J.G. 2021, "Maturation of human intestinal epithelium from pluripotency in vitro", *bioRxiv*.

Klemm, F., Maas, R.R., Bowman, R.L., Kornete, M., Soukup, K., Nassiri, S., Brouland, J., Iacobuzio-Donahue, C.A., Brennan, C. & Tabar, V. 2020, "Interrogation of the microenvironmental landscape in brain tumors reveals disease-specific alterations of immune cells", *Cell,* vol. 181, no. 7, pp. 1643-1660. e17.

Kramer, R., Bucay, N., Kane, D.J., Martin, L.E., Tarpley, J.E. & Theill, L.E. 1996, "Neuregulins with an Ig-like domain are essential for mouse myocardial and neuronal development", *Proceedings of the National Academy of Sciences of the United States of America,* vol. 93, no. 10, pp. 4833-4838.

Laskin, J., Liu, S., Tolba, K., Heining, C., Schlenk, R., Cheema, P., Cadranel, J., Jones, M., Drilon, A. & Cseh, A. 2020, "NRG1 fusion-driven tumors: biology, detection, and the therapeutic role of afatinib and other ErbB-targeting agents", *Annals of Oncology,* vol. 31, no. 12, pp. 1693-1703.

Li, H. 2018, "Minimap2: pairwise alignment for nucleotide sequences", *Bioinformatics,* vol. 34, no. 18, pp. 3094-3100.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R. & 1000 Genome Project Data Processing Subgroup 2009, "The Sequence Alignment/Map format and SAMtools", *Bioinformatics (Oxford, England),* vol. 25, no. 16, pp. 2078-2079.

Liu, X., Bates, R., Yin, D.M., Shen, C., Wang, F., Su, N., Kirov, S.A., Luo, Y., Wang, J.Z., Xiong, W.C. & Mei, L. 2011, "Specific regulation of NRG1 isoform expression by neuronal activity", *The Journal of neuroscience: the official journal of the Society for Neuroscience,* vol. 31, no. 23, pp. 8491-8501.

Marballi, K., Cruz, D., Thompson, P. & Walss-Bass, C. 2012, "Differential neuregulin 1 cleavage in the prefrontal cortex and hippocampus in schizophrenia and bipolar disorder: preliminary findings", *PloS one,* vol. 7, no. 5, pp. e36431.

Mei, L. & Xiong, W. 2008, "Neuregulin 1 in neural development, synaptic plasticity and schizophrenia", *Nature Reviews Neuroscience,* vol. 9, no. 6, pp. 437-452.

Nave, K. & Salzer, J.L. 2006, "Axonal regulation of myelination by neuregulin 1", *Current opinion in neurobiology,* vol. 16, no. 5, pp. 492-500.

784    NCBI Gene [Internet]. Bethesda (MD): National Library of Medicine (US), National Center for
785           Biotechnology Information; 2004 – [2022, Aug, 10]. Available from:
786           https://www.ncbi.nlm.nih.gov/gene/3084
787    Newbern, J. & Birchmeier, C. 2010, "Nrg1/ErbB signaling networks in Schwann cell development and
788           myelination", *Seminars in cell & developmental biology*Elsevier, , pp. 922.
789    Ng, E.S., Davis, R., Stanley, E.G. & Elefanty, A.G. 2008, "A protocol describing the use of a recombinant
790           protein-based, animal product-free medium (APEL) for human embryonic stem cell differentiation as
791           spin embryoid bodies", *Nature protocols,* vol. 3, no. 5, pp. 768-776.
792    Niclis, J.C., Gantner, C.W., Alsanie, W.F., McDougall, S.J., Bye, C.R., Elefanty, A.G., Stanley, E.G.,
793           Haynes, J.M., Pouton, C.W. & Thompson, L.H. 2017, "Efficiently specified ventral midbrain
794           dopamine neurons from human pluripotent stem cells under xeno-free conditions restore motor
795           deficits in parkinsonian rodents", *Stem cells translational medicine,* vol. 6, no. 3, pp. 937-948.
796    Pentassuglia, L. & Sawyer, D.B. 2009, "The role of Neuregulin-1β/ErbB signaling in the heart",
797           *Experimental cell research,* vol. 315, no. 4, pp. 627-637.
798    Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT–PCR. Nucleic
799           acids research, 29(9), pp.e45-e45.
800    Pinkas-Kramarski, R., Shelly, M., Guarino, B.C., Wang, L.M., Lyass, L., Alroy, I., Alamandi, M., Kuo, A.,
801           Moyer, J.D. & Lavi, S. 1998, "ErbB tyrosine kinases and the two neuregulin families constitute a
802           ligand-receptor network", Molecular and cellular biology, vol. 18, no. 10, pp. 6090-6101.
803    Pipes, L., Li, S., Bozinoski, M., Palermo, R., Peng, X., Blood, P., Kelly, S., Weiss, J.M., Thierry-Mieg, J. &
804           Thierry-Mieg, D. 2013, "The non-human primate reference transcriptome resource (NHPRTR) for
805           comparative functional genomics", *Nucleic acids research,* vol. 41, no. D1, pp. D906-D914.
806    Raajendiran, A., Ooi, G., Bayliss, J., O'Brien, P.E., Schittenhelm, R.B., Clark, A.K., Taylor, R.A.,
807           Rodeheffer, M.S., Burton, P.R. & Watt, M.J. 2019, "Identification of metabolically distinct adipocyte
808           progenitor cells in human adipose tissues", *Cell reports,* vol. 27, no. 5, pp. 1528-1540. e7.
809    Rajab, N., Angel, P.W., Deng, Y., Gu, J., Jameson, V., Kurowska-Stolarska, M., Milling, S., Pacheco,
810           C.M., Rutar, M. & Laslett, A.L. 2021, "An integrated analysis of human myeloid cells identifies gaps
811           in in vitro models of in vivo biology", Stem cell reports, vol. 16, no. 6, pp. 1629-1643.
812    Rajab, N., Gearing, L.J., Prawer, Y.D., Angel, P.W., Grimmond, S.M., Laslett, A.L., Choi, J. & Wells, C.A.
813           2022, "Proliferation is a driver of quorum sensing in an in vitro model of tissue resident
814           macrophages", *bioRxiv*.
815    Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G. & Mesirov, J.P.
816           2011, "Integrative genomics viewer", *Nature biotechnology,* vol. 29, no. 1, pp. 24-26.
817    Ryzhov, S., Matafonov, A., Galindo, C.L., Zhang, Q., Tran, T., Lenihan, D.J., Lenneman, C.G., Feoktistov,
818           I. & Sawyer, D.B. 2017, "ERBB signaling attenuates proinflammatory activation of nonclassical
819           monocytes", *American Journal of Physiology-Heart and Circulatory Physiology,* vol. 312, no. 5, pp.
820           H907-H918.
821    Samsa, L.A., Ito, C.E., Brown, D.R., Qian, L. & Liu, J. 2016, "IgG-containing isoforms of Neuregulin-1 are
822           dispensable for cardiac trabeculation in zebrafish", *PLoS One,* vol. 11, no. 11, pp. e0166734.
823    Sheng, Q., Liu, X., Fleming, E., Yuan, K., Piao, H., Chen, J., Moustafa, Z., Thomas, R.K., Greulich, H. &
824           Schinzel, A. 2010, "An activated ErbB3/NRG1 autocrine loop supports in vivo proliferation in ovarian
825           cancer cells", *Cancer cell,* vol. 17, no. 3, pp. 298-310.
826    Sun, L., Cheng, B., Zhou, Y., Fan, Y., Li, W., Qiu, Q., Fang, Y., Xiao, S., Zheng, H. & Li, X. 2020, "ErbB4
827           mutation that decreased NRG1-ErbB4 signaling involved in the pathogenesis of amyotrophic lateral
828           sclerosis/frontotemporal dementia", *Journal of* Alzheimer's Disease, vol. 74, no. 2, pp. 535-544.
829    Sweeney, C., Lai, C., Riese, D.J.,2nd, Diamonti, A.J., Cantley, L.C. & Carraway, K.L.,3rd 2000, "Ligand
830           discrimination in signaling through an ErbB4 receptor homodimer", The Journal of biological
831           chemistry, vol. 275, no. 26, pp. 19803-19807.
832    Tang, C.S., Ngan, E.S., Tang, W., So, M., Cheng, G., Miao, X., Leon, T.Y., Leung, B.M., Hui, K.W. & Lui,
833           V.H. 2012, "Mutations in the NRG1 gene are associated with Hirschsprung disease", *Human*
834           *genetics,* vol. 131, no. 1, pp. 67-76.
835    Vlahos, K., Sourris, K., Mayberry, R., McDonald, P., Bruveris, F.F., Schiesser, J.V., Bozaoglu, K.,
836           Lockhart, P.J., Stanley, E.G. & Elefanty, A.G. 2019, "Generation of iPSC lines from peripheral blood
837           mononuclear cells from 5 healthy adults", *Stem Cell Research,* vol. 34, pp. 101380.

Wagner, M. & Siddiqui, M. 2007, "Signal transduction in early heart development (II): ventricular chamber
specification, trabeculation, and heart valve formation", *Experimental biology and medicine,* vol. 232,
no. 7, pp. 866-880.
Warren CM, Kani K, Landgraf R. The N-terminal domains of neuregulin 1 confer signal attenuation. J Biol
Chem. 2006;281(37):27306-27316.
Wen, D., Suggs, S.V., Karunagaran, D., Liu, N., Cupples, R.L., Luo, Y., Janssen, A.M., Ben-Baruch, N.,
Trollinger, D.B. & Jacobsen, V.L. 1994, "Structural and functional aspects of the multiplicity of Neu
differentiation factors", *Molecular and cellular biology,* vol. 14, no. 3, pp. 1909-1919.
Wolpowitz, D., Mason, T.B., Dietrich, P., Mendelsohn, M., Talmage, D.A. & Role, L.W. 2000, "Cysteine-
rich domain isoforms of the neuregulin-1 gene are required for maintenance of peripheral synapses",
*Neuron,* vol. 25, no. 1, pp. 79-91.
Yu, Q., Kilik, U., Holloway, E.M., Tsai, Y., Harmel, C., Wu, A., Wu, J.H., Czerwinski, M., Childs, C.J. & He,
Z. 2021, "Charting human development using a multi-endodermal organ atlas and organoid models",
*Cell,* vol. 184, no. 12, pp. 3281-3298. e22.
Zhang, C., Mei, W. & Zeng, C. 2022, "Oncogenic Neuregulin 1 gene (NRG1) fusions in cancer: A
potential new therapeutic opportunities", Biochimica et Biophysica Acta (BBA)-Reviews on Cancer,
pp. 188707.
Zhang, D., Sliwkowski, M.X., Mark, M., Frantz, G., Akita, R., Sun, Y., Hillan, K., Crowley, C., Brush, J. &
Godowski, P.J. 1997, "Neuregulin-3 (NRG3): a novel neural tissue-enriched protein that binds and
activates ErbB4", Proceedings of the National Academy of Sciences, vol. 94, no. 18, pp. 9562-9567.
Zhang, P., Kuang, H., He, Y., Idiga, S.O., Li, S., Chen, Z., Yang, Z., Cai, X., Zhang, K., Potthoff, M.J., Xu,
Y. & Lin, J.D. 2018, "NRG1-Fc improves metabolic health via dual hepatic and central action", *JCI
insight,* vol. 3, no. 5, pp. 10.1172/jci.insight.98522.
861
862

**Authorship contributions**
Conceptualization MABR, CAW, Data curation MABR, RDP, YDJP, Formal analysis MABR, RDP, YDJP,
Funding acquisition CAW, MBC, Investigation MABR, CAW, Methodology MABR, RDP, YDJP, MD, SSW,
JG, WD, NR, AL, JH, Project administration CAW, Resources CAW, MBC, TM, Software MABR, RDP,
YDJP, JG Supervision CAW, MBC, Validation MABR, RDP, YDJP, MD, SSW, Visualization MABR, MD,
SSW, writing original draft MABR, Writing – review and editing MABR, RDP, YDJP, MD, SSW, WD, NR,
AL, TM, MBC, CAW.
870

871  **Funding sources**
872  MABR is funded by a University of Melbourne Research Scholarship. MD is supported by the Australian
873  Government Research Training Program Scholarship. MBC is supported by an NHMRC Investigator
874  Fellowship (APP11968410). The project was supported by NHMRC Synergy grant APP 1186371 to CAW.
875

876  **Conflicts of Interest**
877  YDJP, RDP and MBC have received support from Oxford Nanopore Technologies (ONT) to present their
878  findings at scientific conferences. However, ONT played no role in study design, execution, analysis or
879  publication. No conflicts are declared for MABR, MD, SSW, NR, WDN, TM or CAW.
880

**Acknowledgements.**
The authors thank Professor Matt Watt (Department of Anatomy and Physiology, University of
Melbourne), for gifting adipose tissue samples. The authors thanks Dr Le Christy Ying and Prof Paul
Hertzog (Hudson Institute, Clayton, Melbourne) for gifting 2D macrophage RNA for this study. The
authors thank Mrs Zahra Elahi (Department of Anatomy and Physiology, University of Melbourne) for her
logistic and emotional support, crucial to this study. The authors thank Mr Yidi Deng (Department of
Maths and Statistics, University of Melbourne) for discussions about statistical approaches.
888

889  **Sequence Accessions**: All mRNA sequences have been deposited at NCBI GenBank nucleotide
890  database, Accessions assigned 18/1/23: OQ272754, OQ272755, OQ272756, OQ272757, OQ272758,
891  OQ272759, OQ272760, OQ272761, OQ272762, OQ272763, OQ272764, OQ272765, OQ272766,

892    OQ272767, OQ272768, OQ272769, OQ272770, OQ272771, OQ272772, OQ272773, OQ272774,
893    OQ272775, OQ272776.
894