Individual prediction tendencies are related but not the same across modalities

Juliane Schubert[1], Nina Suess[1], Nathan Weisz[1,2]

[1]Centre for Cognitive Neuroscience and Department of Psychology, University of Salzburg, Austria.
[2]Neuroscience Institute, Christian Doppler University Hospital, Paracelsus Medical University, Salzburg, Austria.

Address for correspondence

Juliane Schubert

University of Salzburg

Center for Cognitive Neuroscience

Hellbrunnerstr. 34

A-5020 SALZBURG

Austria/Europe

Tel. 0043 662 8044 5135

Email: juliane.schubert@plus.ac.at

ABSTRACT

Predictive processing theories, which model the brain as a "prediction machine", explain a wide range of cognitive functions, including learning, perception and action. Furthermore, it is increasingly accepted that aberrant prediction tendencies play a crucial role in psychiatric disorders. Given this explanatory value for clinical psychiatry, prediction tendencies are often implicitly conceptualised as individual traits or as tendencies that generalise across situations. As this has not yet explicitly been shown, in the current study, we quantify to what extent the individual tendency to anticipate sensory features of high probability generalises across modalities. Using magnetoencephalography (MEG), we recorded brain activity while participants were presented a sequence of four different (either visual or auditory) stimuli, which changed according to predefined transitional probabilities of two entropy levels: ordered vs. random. Our results show that under conditions of low entropy, stimulus features of high probability are preactivated in the visual and auditory modalities. Crucially, however, the magnitude of the tendency to predict sensory events seems to correlate only weakly between the two modalities. Furthermore, reliability statistics indicate poor internal consistency, suggesting that the measures from the different modalities are unlikely to reflect a single, common cognitive process. In sum, our findings suggest that quantification and interpretation of individual prediction tendencies cannot be generalised across modalities.

Keywords: auditory processing, visual processing, inter-individual differences, magnetoencephalography, predictive processing

INTRODUCTION

In our complex sensory environments, the amount, as well as the level of ambiguity, of information entering our senses requires a system that can filter and integrate information efficiently. As such a system, our brain has to rely fully on its sensory input to infer the true states of objects that cause such input. This inference results in the formation of a so-called internal model, which exists so that predictions can be drawn from it. Even though the assumed implementations of these predictions vary across different "predictive brain" theories (Friston, 2010; Knill & Pouget, 2004; Yon et al., 2019), most agree on the importance of predictions for perception in general. Bottom-up-driven sensory representations in the brain inevitably lag behind the events that caused them, and therefore compensatory mechanisms such as the prediction of future events are highly beneficial for our everyday lives (e.g. to correctly allocate moving objects). Indeed, such anticipation and pre-activation of sensory input has been found for visual perception (Hogendoorn & Burkitt, 2018), as well as for language processing (Dikker & Pylkkänen, 2013). Furthermore, we recently showed that "prediction tendency", which we defined as the tendency to anticipate auditory features of high probability before their occurrence, contributes to explaining individual differences in speech tracking (Schubert et al., 2023).

Complementary to our results, various research has pointed out that individuals seem to vary in the extent to which they rely on top-down priors compared to bottom-up sensory signals for perception. Crucially, these differences in handling predictions or in overall "prediction tendency" have been associated with clinical psychological conditions such as autism (Sinha et al., 2014), schizophrenia (Sterzer et al., 2018), depression (Barrett et al., 2016), PTSD (Kube et al., 2020) and tinnitus (Partyka et al., 2019; Sedley et al., 2016). Taking its explanatory value to clinical psychiatry into account, prediction tendency is often implicitly conceptualised as an individual trait or as a tendency that generalises across situations. Indeed, in line with predictive brain theories for psychosis (Sterzer et al., 2018), it has been found that strong predictive tendencies can promote phantom perception in the auditory modality (Powers et al., 2017). Together with our findings, which state that the aforementioned internal mechanisms are related to differences in speech tracking (Schubert et al., 2023), this strongly suggests that individual prediction tendencies are generalisable across different listening situations. However, the extent to which they generalise across different modalities remains unclear.

Furthermore, there are considerable differences between different modalities (e.g. auditory vs. visual) in the predictability of sensory information and, therefore, also in the way

predictions should best be applied. Visual information (e.g. a single object or a whole scene) naturally unfolds across space, and sensory inference is often drawn from spatially surrounding information. In audition, however, this is not necessarily the case. Considering speech and music, for example, whose predictability mainly unfolds over time, which seems to be a core characteristic of auditory objects. It follows that, in the auditory domain, two types of predictions are usually distinguished, carrying spectral (*what is going to happen*) or temporal (*when is it going to happen*) information (Auksztulewicz et al., 2018; Wollman & Morillon, 2018). In the visual modality, on the other hand, the focus is often on object-based (*what is it*) and spatial (*where is it*) predictions, based on dual-stream theories of vision (e.g. Rao & Ballard, 2005). Nevertheless, a correct anticipation of sensory events should include all of these aspects (*what, when and where*) in both modalities, especially since predictions that generalise and complement across modalities are highly beneficial in natural settings (integrating, for example, lip movements and acoustic speech; Crosse et al., 2016; Sumby & Pollack, 1954). If "prediction tendency" can indeed be conceptualized as an individual trait, it should be similar across situations, as well as across modalities. Complementing correlation estimates, standard measures of internal consistency should help in evaluating whether measures from the auditory and visual modality reflect a putatively common construct. In such a case, internal consistency measures should be sufficiently high (at least >= .7; Tavakol & Dennick, 2011). Inherent differences between auditory and visual perception, however, emphasise the importance of comparative research in the field of predictive processing.

In the current study, we used magnetoencephalographic (MEG) data in visual and auditory versions of an entropy modulation paradigm (see also Demarchi et al., 2019; Schubert et al, 2023) to capture individual prediction tendencies for both modalities. In this paradigm, participants were presented with a sequence of four different (either visual or auditory) stimuli, which changed according to predefined transitional probabilities of two entropy levels: ordered vs. random. In order to optimise comparability, we kept constant spatial predictability (visual stimuli were presented at screen center and auditory stimuli were presented binaurally at phantom center), as well as temporal predictability (with a fixed stimulation rate of 3 Hz), modulating only one stimulus feature in each modality. In the visual version, participants were presented with a sequence of gabor patches that changed in orientation, whereas in the auditory version, participants listened to pure tones of different frequencies. For both modalities, individual prediction tendency was defined as the tendency to anticipate and pre-activate stimulus features of high probability.

Fig. 1: *Quantification of individual prediction tendency.* ***A)*** *Participants were presented with sequences of four different stimuli. Transitional probabilities varied according to two entropy conditions (ordered vs. random).* ***C)*** *Visual stimulation consisted of gabor patches in four different orientations, auditory stimulation consisted of pure tones at four different frequencies. An LDA classifier was used to decode stimulus feature from brain activity across time, trained on (high-probability) ordered forward transition trials and tested on all repetition trials.* ***B)*** *Expected classifier decision values contrasting the brain's pre-stimulus tendency to predict a forward transition between the two entropy levels. Individual prediction tendency quantification results from the summed difference between conditions (ordered > random) over pre-stimulus time.*

METHODS

Subjects

In total, 35 subjects (16 male, mean age = 32.5, range = 19 - 57) were recruited to participate in the experiment. All participants reported normal hearing and had normal, or corrected to normal, vision. They gave written, informed consent and reported that they had no previous neurological or psychiatric disorders. The experimental procedure was approved by the ethics committee of the University of Salzburg and was carried out in accordance with the declaration of Helsinki. All participants received either a reimbursement of 10 € per hour or course credits for their participation.

Experimental Procedure

Before the start of the experiment, participants' individual head shapes were assessed using cardinal head points (nasion and pre-auricular points), digitised with a Polhemus Fastrak Digitiser (Polhemus), as well as around 300 points on the scalp. For every participant, MEG sessions started with a 5-minute resting-state recording, after which individual hearing threshold was determined, using a pure tone of 1043 Hz. This was followed by 4 experimental blocks (2 blocks per modality) of an entropy modulation paradigm. Participants started with either the auditory or visual blocks (chosen randomly for each participant). In the auditory paradigm (see also Schubert et al., 2023), participants passively listened to sequences of four different pure tones (1: 440 Hz, 2: 587 Hz, 3: 782 Hz, 4: 1043 Hz) while watching a landscape movie (LoungeV Films, 2017). All tones were presented binaurally at equal volume for the left and right ears (i.e. at phantom center) at 40 db above the individual hearing threshold. In the visual paradigm, a gabor patch (spatial frequency: 0.01 cycles/pixel, sigma: 60 pixels, phase: 90°) was presented at four perceptually different orientations (1: 0°, 2: 45°,3: 90°, 4: 135°) in the center of the screen while participants listened to podcasts from "Bayern 2 - radioWissen": "Kakao - Das braune Gold" (Cocoa - the brown gold) and "Zimt - Die Würze des Lebens" (Cinnamon - the spice of life). There was no task, and participants were just instructed to relax and to move as little as possible. To ensure temporal predictability, all stimuli were presented at a fixed stimulation rate of 3 Hz, and each stimulus presentation (i.e. pure tone or gabor patch) lasted for 100ms. One block consisted of a sequence of 2800 trials, totalling 5600 trials per modality. Transitional probabilities between stimuli (1, 2, 3, 4) were determined by two different entropy levels (ordered vs. random see **Fig. 1A**) . Entropy changed pseudorandomly within each block every 700 trials. While in an "ordered" context, certain transitions (hereinafter referred to as forward transitions, i.e. 1→2, 2→3, 3→4, 4→1) were to be expected, with a high probability of 75%, self repetitions (e.g., 1→1, 2→2,...) were rather unlikely, with a probability of 25%. However, in a "random" context, all possible transitions (including forward transitions and self repetitions) were equally likely, with a probability of 25% (see **Fig. 1A** and see also Schubert et al., 2023). Furthermore, a pseudorandom 10% of the trials (i.e. ~ 70 trials per stimulus, per entropy, per modality) were omitted (as in Demarchi et al., 2019). Between each of the four blocks (~15 min each), there was a short break. In total, the experiment lasted approximately 2h per participant (including MEG preparation time). The experiment was coded and conducted with the Psychtoolbox-3 (Brainard, 1997; Kleiner et al., 2007), with an additional class-based library ('Objective Psychophysics Toolbox', o_ptb) on top of it (Hartmann & Weisz, 2020).

MEG Data Acquisition and Preprocessing

A whole-head MEG system (Elekta Neuromag Triux, Elekta Oy, Finland), placed within a standard, passive magnetically shielded room (AK3b, Vacuumschmelze, Germany), was used to capture magnetic brain activity with a sampling frequency of 1 kHz (hardware filters: 0.1 - 330 Hz). The signal was recorded with 102 magnetometers and 204 orthogonally placed planar gradiometers at 102 different positions. In a first step, a signal space separation algorithm, implemented in the Maxfilter program (version 2.2.15) provided by the MEG manufacturer, was used to clean the data of external noise and realign data from different blocks to a common standard head position. Data preprocessing was performed using Matlab R2020b (The MathWorks, Natick, Massachusetts, USA) and the FieldTrip Toolbox (Oostenveld et al., 2010). To identify eye blinks and heartbeat artifacts, 50 independent components were identified from filtered (0.1 - 100 Hz) continuous data of the first experimental block of both modalities (auditory + visual). On average, 2.6 (range = 2 - 5) components were removed for each subject. All data was filtered between 0.1 Hz and 30 Hz (Kaiser windowed finite impulse response filter) and downsampled to 100 Hz. Then, the data of each block was epoched into segments of 1200 ms (from 400 ms before stimulus onset to 800 ms after onset) for further analysis (as in Schubert et al., 2023).

Decoding Analysis

Multivariate pattern analyses were carried out using the MVPA-Light package (Treder, 2020) and were conducted separately for each modality. A multi-class linear discriminant analyser (LDA) was used to classify stimulus feature (i.e. 1 - 4, sound frequency or gabor-patch orientation) from brain activity between -0.3 and 0.3 ms in a time-resolved manner. Based on resulting classifier decision values (i.e. d1 - d4 for every test-trial and time-point), we calculated the individual prediction tendency within each modality (as in Schubert et al., 2023): We define individual prediction tendency as the tendency to pre-activate sound frequencies of high probability (i.e. a forward transition from one stimulus to another: 1→2, 2→3, 3→4, 4→1). In order to capture any prediction-related neural activity, we trained the classifier exclusively on ordered forward trials. Afterwards, the classifier was tested on all self-repetition trials, providing classifier decision values for every stimulus feature, which were then transformed into corresponding transitions (e.g. d1(t) | 1(t-1) "dval for 1 at trial t, given that 1 was presented at trial t-1" → repetition, d2(t) | 1(t-1) → forward,...). The tendency to represent a forward vs. repetition transition was contrasted for both ordered and random trials. Using self-repetition trials for testing, we ensured a fair comparison between the ordered and random contexts (with an equal number of trials and the same preceding

bottom-up input). Thus, we quantified "prediction tendency" as the classifier's pre-stimulus tendency to a forward transition in an ordered context exceeding the same tendency in a random context (which can be attributed to carry-over processing of the preceding stimulus). Then, using the summed difference across pre-stimulus times, one value can be extracted per subject, per modality.

Statistical Analysis

The classifier's tendency toward a forward transition in an ordered context was compared to the same tendency in a random context, using a cluster-based permutation t-test (using 10,000 random permutations, a cluster-alpha of 0.025 and a monte-carlo critical alpha of 0.025) over a time window from -0.3 s to 1.5 s. This contrast was calculated for both modalities separately. Afterwards, the summed differences across pre-stimulus time was extracted for each subject and modality, resulting in individual prediction tendency values. These single values were then used to calculate the pearson's correlation coefficient between auditory and visual prediction tendency. To extend our findings by classical estimates of internal consistency, we also calculated Cronbach's alpha and the Spearman-Brown reliability estimates, assuming that prediction tendency can be understood as a multi-dimensional concept, here captured with a two-item (auditory, visual) scale. Additionally, we also calculated the reliability of predictions between modalities (Spearman-Brown's cofficient: rho = 2*r/(1+r); formula taken from Eisinga et al., 2013) for all possible pre-stimulus time-by-time combinations. All analyses were done in MATLAB version R2021a.

RESULTS

Anticipatory predictions can be found in both modalities

To investigate anticipatory prediction, participants were presented a sequence of four different (either visual or auditory) stimuli, which changed according to predefined transitional probabilities of two entropy levels: ordered vs. random. In a first step, we were interested if we would see a pre-activation of high-probability stimulus features in brain activity across subjects. Cluster-based permutation showed an overall auditory prediction tendency (i.e. ordered > random) in a sustained pre-stimulus cluster from -0.25 s to -0.02 s and others that extended into the post-stimulus period (from 0 s to 0.05 s and from 0.07 s to 0.09 s). For visual prediction tendency, we found 3 pre-stimulus clusters: from -0.25 s to -0.21 s, from -0.17 s to -0.12 s and from -0.04 s to -0.02 s. These results confirm that,

irrespective of individual differences, an overall tendency to anticipate stimulus features of increased probability can be found in both modalities (see Fig. 2A).

Individual prediction tendency is weakly correlated between modalities

In a second step, we wanted to further investigate prediction tendency as a subject-specific, potentially unified concept. For this purpose, we looked particularly into the generalisation between auditory and visual modalities. Using the summed prediction tendency values across pre-stimulus time (-0.3 s - 0 s), one value was extracted per subject, per modality. Even though individual prediction tendency values were, on average, higher for the auditory (mean = 0.127, sd = 0.123) than the visual (mean = 0.091, sd = 0.119) modality, there was no significant difference in a one-sample t-test (t = 1.574, 95% CI = [-0.011, 0.085]). In total, 23 subjects showed a stronger auditory prediction tendency, compared to 12 subjects with stronger prediction tendency in the visual domain (see Fig. 2B). Crucially, we found a weak, but significant, correlation of individual prediction tendency between modalities, using pearson's correlation (r = 0.34, p = 0.044; see Fig. 2B). Assuming that both measures represent the same concept of individual prediction tendency, we further calculated Cronbach's alpha and Spearman-Brown's rho to estimate internal consistency (see Table 1). Because there seems to be some disagreement as to which reliablity statistic should be used for measures comprising two items, we decided to report all three here (Eisinga et al., 2013). In an attempt to account for individual temporal dynamics of each modality, we also estimated the reliability for all possible pre-stimulus time-by-time combinations. It seems that early auditory predictions generalise best to late visual predictions, with overall low reliability (see Fig. 2C). Crucially, based on guidelines from classical test theory, our results indicate that the effect is too weak to provide a reliable generalisation from one modality to another (Tavakol & Dennick, 2011). In sum, these findings suggest that individual prediction tendencies don't necessarily generalise across different modalities but are related in a way that needs to be investigated further to draw unambiguous conclusions.

Table 1: Comparison of correlation coefficient, cronbach alpha and spearman brown coefficient

	Pearson's r	Cronbach's alpha	Spearman-Browns rho
Reliability estimation	0.343	0.510	0.511

Note: Reliability was calculated treating *prediction tendency* as a two-item (*auditory*, *visual*) scale.

Fig. 2: A) *Time-resolved prediction tendency for auditory and visual modalities: In both modalities, there seems to be an anticipatory tendency to represent stimulus features of high probability (i.e. forward transition) in a predictable context (y-axis represents the difference in dvals (ordered - random) for a forward transition; solid lines on x-axis indicate significant time-points (based on a random permutation test using an alpha level of 0.025)).* **B)** *Correlation of individual prediction tendency between modalities: There is a weak generalisation from auditory to visual prediction tendency (the solid line represents a perfect generalisation from one modality to another; the dashed line indicates the least-squared regression line).* **C)** *Temporal generalisation of forward predictions across modalities: early anticipatory predictions in the auditory modality seem to generalise best to late anticipatory predictions in the visual modality, however only poor reliability is indicated overall (colormap represents Spearman-Browns coefficient, * taken from Tavakol & Dennick, 2011; N = 35).*

DISCUSSION

Predictive processing has been found throughout the range of human cognition: in language perception and production (e.g. Pickering & Garrod, 2013), object recognition (e.g. Oliva & Torralba, 2007), motor perception and action (e.g. Shipp et al., 2013), emotional awareness (e.g. Smith et al., 2019), and decision making (e.g. Summerfield & de Lange, 2014). However, the extent to which all these predictions can be attributed to a collective mechanism remains unclear. Beyond that, several conceptual and mathematical predictive brain models have been proposed to explain cognitive and perceptual pathologies in clinical neuroscience (for a review see Smith et al., 2021). Considering that psychiatric conditions such as depression, autism or schizophrenia are assumed to be highly individualized and stable across time and situations, estimations of prediction tendency are likely to be interpreted as a "trait-dependent" rather than a "state-dependent" measure. In this regard, prediction tendency should a) vary considerably across individuals, b) show long-term stability within an individual and c) generalise across different situations. Yet research that validates this claim is scarce. In the current study, we aimed to investigate to what extent the individual tendency to anticipate sensory features of high probability generalises across modalities. Even though aberrant predictive tendencies, especially linked to psychosis, have been proposed for auditory (Corlett et al., 2019) as well as visual perception (Adams et al., 2012), this is, to our knowledge, the first study to directly compare individual prediction tendencies between two modalities. Our results show that under conditions of low entropy, stimulus features of high probability are preactivated in the visual modality as well as the auditory. Crucially, however, this tendency to predict sensory events seems to correlate only weakly between the two modalities, suggesting that the quantification and interpretation of prediction tendencies cannot be generalised.

From our evidence, we can conclude that around 12% of total variance in visual prediction tendency can be explained by auditory prediction tendency and vice versa. Even though the correlation between modalities provided moderate evidence for a relationship, we argue that this seems too weak to support the assumption that individual prediction tendency can be generalised from one modality to another. Considering that the same metric was applied in both cases, we orient ourselves primarily on standards that are also used to assess reliability, for which acceptable statistics usually range between 0.7 and 0.95 (Tavakol & Dennick, 2011). Additionally, it should be noted that our prediction tendency quantification was derived from the decoding of stimulus features from brain activity, a measure that is highly affected by a subject-specific signal-to-noise ratio (as a result of anatomical differences or individual distributive properties, a problem often discussed when calculating

source localisation; Puce & Hämäläinen, 2017). It can therefore be assumed that decoding accuracies are correlated within subjects, and it should be considered that a certain proportion of the relationship is driven by individual differences in signal-to-noise ratio alone. In addition, it should be considered that our quantification might not be a "pure" measure of prediction tendency, which reduces internal consistency, even within a modality. In sum, we support the notion that scientific evidence should not be judged solely on the basis of its significance, but that the strength of effects should be evaluated within the framework of underlying assumptions. We therefore conclude that our estimates of reliability are too weak to allow a generalisation across modalities.

Our results indicate that individuals who show a stronger auditory prediction tendency don't necessarily show a similarly strong tendency for visual predictions. There are different possible explanations for these findings. In the current paradigm, stimulus predictability was modulated via transitional probabilities in a sequence of events, a feature that is quite characteristic for auditory stimuli, such as music and speech, but maybe less so for visual stimulation (ten Oever et al., 2014). To make optimal use of top-down inferences in auditory processing, sensory information has to be integrated over (past) time (and extrapolated into the future) and, crucially, has to be accurately located within time. Although visual processing has to meet similar requirements, particularly when confronted with moving objects, the motivation for anticipatory representations might be different (e.g. to facilitate suitable motor action in time), potentially resulting in increasingly goal-driven predictions in the visual modality (for a review see Fiehler et al., 2019). Such differences in motivation could very well lead to modality-specific implementations of anticipatory predictions, which might explain the weak generalisation between modalities in our results.

Indeed, it seems that the extent to which an observer (versus a listener) has to be actively engaged in the sampling of information is different between vision and audition. For example, information that gradually unfolds over time, such as a spoken sentence, can (under optimal noise conditions) be received without any active adjustments from the listener (although see, for example, Gehmacher et al., 2022, for top-down influence on auditory nerve activity). Brain responses to unexpected changes in auditory stimulation (the classic "mismatch negativity"), for example, have even been found during sleep (Loewy et al., 1996). Meanwhile visual information, which is usually spread throughout space, extending the foveal field of the observer, requires active effort (e.g. eye, head or body movements) to be sampled. According to some theories, however, active sampling of sensory evidence is crucial for the formation of internal models and the predictions that come from them (Friston et al., 2017). Since our paradigm was entirely passive, no beneficial gain was to be expected from focusing on the sequences. Thus, participants likely did not allocate their attentional

resources to them. If the proposed link between predictions and active engagement is stronger in vision, a lack of attention and goal-directedness could explain why prediction tendency did not generalise from the auditory to the visual modality. We suggest that future research should investigate the (potentially different) roles of attention for predictive processing in vision versus audition.

Conclusions and implications for future research

Predictive processing theories, which model the brain as a "prediction machine", are able to explain a wide range of cognitive functions, including learning, perception and action. Furthermore, it seems to be widely accepted that aberrant prediction tendencies play a crucial role in long-term psychiatric disorders. Crucially, in clinical neuroscience, prediction tendency is often implicitly conceptualized as an individual trait or as a tendency that generalises across situations, yet research to validate this claim is scarce. In the current study, we found evidence for anticipatory predictions in the auditory modality as well as the visual. However, our results suggest that quantification and interpretation of individual prediction tendency cannot necessarily be generalised across modalities. This emphasises the importance of research, further investigating and validating the implicit assumption of "prediction tendency" as a trait, since this assumption bears concrete implications for the growing field of computational modeling in neuroscience and psychiatry. We propose that future research focuses on the validation of state- vs. trait-dependent assumptions in predictive processing theories. Furthermore, we strongly suggest investigation of the intraindividual stability of prediction tendency within and across different modalities.

REFERENCES

Adams, R. A., Perrinet, L. U., & Friston, K. (2012). Smooth Pursuit and Visual Occlusion: Active Inference and Oculomotor Control in Schizophrenia. *PLOS ONE*, *7*(10), e47502. https://doi.org/10.1371/journal.pone.0047502

Auksztulewicz, R., Schwiedrzik, C. M., Thesen, T., Doyle, W., Devinsky, O., Nobre, A. C., Schroeder, C. E., Friston, K. J., & Melloni, L. (2018). Not All Predictions Are Equal: "What" and "When" Predictions Modulate Activity in Auditory Cortex through Different Mechanisms. *The Journal of Neuroscience*, *38*(40), 8680–8693. https://doi.org/10.1523/JNEUROSCI.0369-18.2018

Barrett, L. F., Quigley, K. S., & Hamilton, P. (2016). An active inference theory of allostasis and interoception in depression. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1708), 20160011. https://doi.org/10.1098/rstb.2016.0011

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Corlett, P. R., Horga, G., Fletcher, P. C., Alderson-Day, B., Schmack, K., & Powers, A. R. (2019). Hallucinations and Strong Priors. *Trends in Cognitive Sciences*, *23*(2), Article 2. https://doi.org/10.1016/j.tics.2018.12.001

Crosse, M. J., Liberto, G. M. D., & Lalor, E. C. (2016). Eye Can Hear Clearly Now: Inverse Effectiveness in Natural Audiovisual Speech Processing Relies on Long-Term Crossmodal Temporal Integration. *Journal of Neuroscience*, *36*(38), 9888–9895. https://doi.org/10.1523/JNEUROSCI.1396-16.2016

Demarchi, G., Sanchez, G., & Weisz, N. (2019). Automatic and feature-specific prediction-related neural activity in the human auditory system. *Nature Communications*, *10*. https://doi.org/10.1038/s41467-019-11440-1

Dikker, S., & Pylkkänen, L. (2013). Predicting language: MEG evidence for lexical preactivation. *Brain and Language*, *127*(1), 55–64. https://doi.org/10.1016/j.bandl.2012.08.004

Eisinga, R., Grotenhuis, M. te, & Pelzer, B. (2013). The reliability of a two-item scale:

Pearson, Cronbach, or Spearman-Brown? *International Journal of Public Health*, *58*(4), 637–642. https://doi.org/10.1007/s00038-012-0416-3

Fiehler, K., Brenner, E., & Spering, M. (2019). Prediction in goal-directed action. *Journal of Vision*, *19*(9), 10. https://doi.org/10.1167/19.9.10

Friston, K. (2010). The free-energy principle: A unified brain theory? *Nature Reviews Neuroscience*, *11*(2), 127–138. https://doi.org/10.1038/nrn2787

Friston, K., FitzGerald, T., Rigoli, F., Schwartenbeck, P., & Pezzulo, G. (2017). Active Inference: A Process Theory. *Neural Computation*, *29*(1), 1–49. https://doi.org/10.1162/NECO_a_00912

Gehmacher, Q., Reisinger, P., Hartmann, T., Keintzel, T., Rösch, S., Schwarz, K., & Weisz, N. (2022). Direct Cochlear Recordings in Humans Show a Theta Rhythmic Modulation of Auditory Nerve Activity by Selective Attention. *Journal of Neuroscience*, *42*(7), 1343–1351. https://doi.org/10.1523/JNEUROSCI.0665-21.2021

Hartmann, T., & Weisz, N. (2020). An Introduction to the Objective Psychophysics Toolbox. *Frontiers in Psychology*, *11*, 585437. https://doi.org/10.3389/fpsyg.2020.585437

Hogendoorn, H., & Burkitt, A. N. (2018). Predictive coding of visual object position ahead of moving objects revealed by time-resolved EEG decoding. *NeuroImage*, *171*, 55–61. https://doi.org/10.1016/j.neuroimage.2017.12.063

Kleiner, M., Brainard, D. H., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's new in Psychtoolbox-3. *Perception*, *36*, 1–16. https://doi.org/10.1068/v070821

Knill, D. C., & Pouget, A. (2004). The Bayesian brain: The role of uncertainty in neural coding and computation. *Trends in Neurosciences*, *27*(12), Article 12. https://doi.org/10.1016/j.tins.2004.10.007

Kube, T., Berg, M., Kleim, B., & Herzog, P. (2020). Rethinking post-traumatic stress disorder – A predictive processing perspective. *Neuroscience & Biobehavioral Reviews*, *113*, 448–460. https://doi.org/10.1016/j.neubiorev.2020.04.014

Loewy, D. H., Campbell, K. B., & Bastien, C. (1996). The mismatch negativity to frequency deviant stimuli during natural sleep. *Electroencephalography and Clinical*

Neurophysiology, *98*(6), 493–501. https://doi.org/10.1016/0013-4694(96)95553-4

Partyka, M., Demarchi, G., Roesch, S., Suess, N., Sedley, W., Schlee, W., & Weisz, N.

(2019). *Phantom auditory perception (tinnitus) is characterised by stronger*

anticipatory auditory predictions (p. 869842). bioRxiv. https://doi.org/10.1101/869842

Powers, A. R., Mathys, C., & Corlett, P. R. (2017). Pavlovian Conditioning-Induced

Hallucinations Result from Overweighting of Perceptual Priors. *Science (New York,*

N.Y.), *357*(6351), 596–600. https://doi.org/10.1126/science.aan3458

Puce, A., & Hämäläinen, M. S. (2017). A Review of Issues Related to Data Acquisition and

Analysis in EEG/MEG Studies. *Brain Sciences*, *7*(6), Article 6.

https://doi.org/10.3390/brainsci7060058

Rao, R. P. N., & Ballard, D. H. (2005). CHAPTER 91—Probabilistic Models of Attention

Based on Iconic Representations and Predictive Coding. In L. Itti, G. Rees, & J. K.

Tsotsos (Eds.), *Neurobiology of Attention* (pp. 553–561). Academic Press.

https://doi.org/10.1016/B978-012375731-9/50095-1

Schubert, J., Schmidt, F., Gehmacher, Q., Bresgen, A., & Weisz, N. (2023). Cortical speech

tracking is related to individual prediction tendencies. *Cerebral Cortex*, bhac528.

https://doi.org/10.1093/cercor/bhac528

Sedley, W., Friston, K. J., Gander, P. E., Kumar, S., & Griffiths, T. D. (2016). An Integrative

Tinnitus Model Based on Sensory Precision. *Trends in Neurosciences*, *39*(12),

799–812. https://doi.org/10.1016/j.tins.2016.10.004

Sinha, P., Kjelgaard, M. M., Gandhi, T. K., Tsourides, K., Cardinaux, A. L., Pantazis, D.,

Diamond, S. P., & Held, R. M. (2014). Autism as a disorder of prediction. *Proceedings*

of the National Academy of Sciences, *111*(42), 15220–15225.

https://doi.org/10.1073/pnas.1416797111

Smith, R., Badcock, P., & Friston, K. J. (2021). Recent advances in the application of

predictive coding and active inference models within clinical neuroscience. *Psychiatry*

and Clinical Neurosciences, *75*(1), 3–13. https://doi.org/10.1111/pcn.13138

Sterzer, P., Adams, R. A., Fletcher, P., Frith, C., Lawrie, S. M., Muckli, L., Petrovic, P.,

Uhlhaas, P., Voss, M., & Corlett, P. R. (2018). The Predictive Coding Account of

Psychosis. *Biological Psychiatry*, *84*(9), 634–643.

https://doi.org/10.1016/j.biopsych.2018.05.015

Sumby, W. H., & Pollack, I. (1954). Visual Contribution to Speech Intelligibility in Noise. *The*

Journal of the Acoustical Society of America, *26*(2), 212–215.

https://doi.org/10.1121/1.1907309

Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International Journal*

of Medical Education, *2*, 53–55. https://doi.org/10.5116/ijme.4dfb.8dfd

ten Oever, S., Schroeder, C. E., Poeppel, D., van Atteveldt, N., & Zion-Golumbic, E. (2014).

Rhythmicity and cross-modal temporal cues facilitate detection. *Neuropsychologia*,

63, 43–50. https://doi.org/10.1016/j.neuropsychologia.2014.08.008

Treder, M. S. (2020). MVPA-Light: A Classification and Regression Toolbox for

Multi-Dimensional Data. *Frontiers in Neuroscience*, *14*, 289.

https://doi.org/10.3389/fnins.2020.00289

Wollman, I., & Morillon, B. (2018). Organizational principles of multidimensional predictions

in human auditory attention. *Scientific Reports*, *8*(1), Article 1.

https://doi.org/10.1038/s41598-018-31878-5

Yon, D., de Lange, F. P., & Press, C. (2019). The Predictive Brain as a Stubborn Scientist.

Trends in Cognitive Sciences, *23*(1), Article 1.

https://doi.org/10.1016/j.tics.2018.10.003