Title: **Erasable Hippocampal Neural Signatures Predict Memory Discrimination**

Authors: Kinsky, Nathaniel R.,[1,3,5,6] Orlin, Daniel J.,[3,4,5] Ruesch, Evan A.[3], Diba, Kamran[1,2],
Ramirez, Steve[3,6]

1 – Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI 48109

2 – Neuroscience Graduate Program, University of Michigan, Ann Arbor, MI 48109

3 – Center for Systems Neuroscience, Boston University, Boston, MA 02451

4 – Neuroscience Graduate Program, Oregon Health & Science University, Portland, OR 97239

5 – These authors contributed equally to this work

6 – Corresponding authors. Correspondence should be addressed to N.R.K
(nkinsky@umich.edu) or S.R. (dvsteve@bu.edu)

Abstract: We leveraged the spatial-temporal resolution of in vivo calcium imaging in freely
moving mice to interrogate how blocking memory consolidation impacted learning-related
hippocampal dynamics following contextual fear conditioning. We found that memory specificity
correlated with the amount of cell turnover between a shock and neutral arena and that learning
caused context-specific remapping. We induced amnesia by blocking protein synthesis following
learning, which promoted an acute suppression of neural activity, and arrested learning-related
place field remapping. Finally, freeze-predicting neural ensembles emerged following learning,
and their coordinated activity required protein synthesis. We conclude that context-specific
place field remapping and the development of coordinated ensemble activity underlie contextual
fear memory consolidation.

Main Text:

The process of memory consolidation requires protein synthesis following learning to produce
stable long-term memories (Barondes & Cohen, 1968; Davis & Squire, 1984; Squire &
Barondes, 1974). At the synaptic level, protein synthesis is necessary for the formation of late-
phase but not early-phase LTP in HPC neurons (Frey & Morris, 1997; Huang et al., 1994;
Nguyen et al., 1994). Likewise, blocking protein synthesis following learning impairs long-term
memory, but not short-term memory, for a hippocampus(HPC)-dependent contextual fear
conditioning (CFC) task (Ryan et al., 2015; Schafe et al., 1999). These studies suggest that
memory consolidation requires protein-synthesis to reinforce learning-related connections which
are potentiated during CFC learning. Despite this evidence linking synaptic potentiation *in vitro*
to consolidation, little is known about how protein synthesis influences the functional coding
properties of neurons *in vivo* before, during, and after learning to support memory consolidation.
To causally test the hypothesis that learning-related place-cell remapping (Muller & Kubie,
1987) is necessary for consolidation of a CFC memory (Moita et al., 2004; Wang et al., 2012)
we combined in vivo calcium imaging in mice with systemic administration of the protein
synthesis inhibitor, anisomycin, to track the evolution, remapping, and stabilization of HPC place
fields during and after CFC. We further explored how between-animal variability in HPC
dynamics influenced memory specificity and how blocking protein-synthesis impacted the
development of HPC ensemble activity during freezing behavior.

Weeks prior to training, all mice received viral infusions of the genetically encoded calcium
indicator GCaMP6f (Chen et al., 2013) in region CA1 of the dorsal hippocampus. Subsequently,

following two days of pre-exposure ("Before" learning, days -2 and -1) to an operant chamber
(shock arena) and an open field (neutral arena), mice received a mild foot-shock on day 0
(training), after which they were moved to their home cage and immediately given systemic
injections of either anisomycin (ANI group) or vehicle (CTRL group). We then performed a test
of short-term memory 4 hours later and three tests of long-term memory 1, 2, and 7 days
following training (Figure 1A), each time by measuring freezing behavior. We titrated the shock
level during training such that the mice froze significantly more in the shock arena relative to the
neutral arena following learning while still exploring the majority of both arenas. We observed a
range of freezing levels during the day 1 and 2 memory tests (Figure S1A) and sub-divided
CTRL mice into two groups: Learners, who froze significantly more in the shock arena, and
Non-Learners, who either showed generalized freezing or froze at low levels in both arenas
(Figures 1B, 1G, Figure S1B). Learners exhibited reduced freezing on day 7, indicative of
extinction (Figure 1B). All subsequent analyses related to long-term memory therefore utilized
the day 1 and 2 recall sessions ("After" consolidation). In contrast, mice in the ANI group
exhibited no difference in freezing between arenas at any time point, indicating that blocking
protein synthesis impaired context-specific fear memory (Figure 1C). Both the CTRL and ANI
groups exhibited significant freezing in the shock arena during the 4 hour test (Figure 1D, Figure
S1A), though we note that the behavior in the ANI group at this time point could arise from
either contextual fear or non-specific aversive effects of anisomycin treatment (Figure S2) since
these mice froze at high levels in both arenas.

We visualized the activity of pyramidal neurons in region CA1 of the dorsal hippocampus
(Figure 1C) using a miniaturized epifluorescence microscope (Ghosh et al., 2011; Ziv et al.,
2013). We identified a large number of neurons in each 10 minute session (n = 128 to 1216),
extracted their corresponding calcium traces, and tracked them between sessions throughout
the CFC task, which allowed us to determine the long-term evolution of the HPC neural code
(Figure 1C,F). We next assessed behavioral expression of a CFC memory by calculating a
behavioral discrimination index (DI_{beh}) to quantify how much each animal froze in the shock vs.
neutral arenas. Negative DI values indicated higher freezing in the shock arena. Learners
exhibited negative DI_{beh} levels on days 1 and 2 that were significantly different from both Non-
Learners (by definition) and from the ANI group as well (Figure 1G). We noticed that many
neurons exhibited strong changes in mean event rate between arenas (Figure 1F) and
calculated a neural discrimination index (DI_{neural}) to quantify the distinctiveness of neural activity
between arenas (0 = similar, 1 = distinct). Same-day neural discrimination between arenas
correlated strongly with across-day neural discrimination in the same arena (Figure 1E, Figure
S1C). This indicates that mice exhibit natural variability in neural discrimination which is
invariant between different arenas and across time. Interestingly, we noticed that DI_{neural} was
significantly higher for Learners vs. Non-Learners in the sessions *prior to* the shock (Figure 1H),
indicating that inherent differences in neural discrimination influence how well mice learn the
CFC task.

Previous work has demonstrated that higher overlap between HPC neurons active in two
distinct arenas correlates with increased generalization or linking of a contextual fear memory
across arenas (Cai et al., 2016). Accordingly, we hypothesized that the distinctiveness of the
HPC neural code between arenas would predict how much mice froze in the neutral arena. We
therefore utilized a regression analysis to determine if neural discrimination during memory
recall *after* shock likewise correlated with memory discrimination. We utilized two metrics

Figure 1: Neural discrimination between arenas predicts specificity of fear learning. A) Schematic of the behavioral paradigm. Mice freely explored two distinct arenas (neutral and shock) for 10 minutes each day. Mice underwent mild contextual fear conditioning on day 0 in the shock arena followed by immediate I.P. administration of anisomycin or vehicle in their home cage. Memory recall tests were conducted 4 hours and 1, 2, and 7 days post-shock. The time of each session is referenced to the shock session. **B)** (left) Learner (CTRL) mice freezing on all days. Red = shock arena, blue = neutral arena. *p=1.3e-0.5 shock - neutral freezing from days -2/-1 to days 1 one-sided t-test (n=4 mice). (right) Same but for Non-Learner (CTRL) mice (n=3 mice). **C)** (left) Neural overlap plots between neutral and shock arenas for an example Learner mouse on day -1, before shock. Green = cells active in the shock arena only, yellow = cells active in the neutral arena only, orange = cells active in both arenas. (right) Same for example Non-Learner on day -2 showing higher overlap of active cells between arenas. **D)** Same as B but for ANI group **E)** Mean neural discrimination between arenas (same day, days -2 to -1 and 1 to 2) vs. within arenas (same arena, days -2 and -1, days 1 and 2). *p=2.35e-8 (ρ=-0.56) Spearman correlation. **F)** Example calcium activity from the Learner mouse shown in C (left) for cells active in both arenas. Black = calcium trace, red = putative spiking activity during transient rises. Top row shows shock arena preferring cells, bottom row shows neutral arena preferring cells. **G)** Behavioral discrimination between arenas after shock (Days 1-2) shows formation of a specific fear memory for Learners only, by definition (positive = more freezing in neutral arena, negative = more freezing in shock arena, 0 = equal freezing in both arenas). *p=0.022, **p=0.00038 1-sided t-test **H)** Neural discrimination between arenas BEFORE shock indicates Learners formed more distinct representations of each arena prior to learning. Same conventions as F. *p=0.030, +p=0.059 two-sided t-test **I)** Neural overlap between arenas correlates with specificity of fear memory on days 1-2 for Learners but not Non-Learners. *p=0.033 (ρ=0.61). **J)** Same as I) but plotting neural overlap vs. behavioral discrimination between arenas. +p=0.059 (ρ=-0.56) for Learners.

to quantify the distinctiveness of the HPC neural code between arenas: 1) the overlap ratio of neurons active in both arenas divided by the total number of cells active in either arena, which accounts for cells that may be silent in one of the arenas, and 2) the aforementioned DI_{neural} metric, which is based on the cells that are active in both arenas. We found that overlap ratio

was significantly correlated with DI_{beh} for Learners, but not for Non-Learners, on days 1-2
(Figure 1I); likewise, the correlation between DI_{beh} and DI_{neural} approached significance for
Learners but not for Non-Learners on days 1-2 (Cai et al., 2016; Figure 1J). Together, our
results indicate that mice with segregated experiences in the HPC neural code were more
capable of forming discrete long-term memories and that the distinctiveness of the HPC calcium
activity predicts the specificity of memory for CFC.

Next, we probed how arresting protein synthesis impacted HPC dynamics. We hypothesized
that, by preventing plasticity, ANI administration would slow or stop the normal rate of cell
turnover observed across days (Figure 2A). Surprisingly, we found that ANI administration
rapidly accelerated the turnover in which cells are active at any given time. This "cell turnover"
was indicated by lower overlap of active cells from experiment start to the 4 hour session in the
ANI group compared to the CTRL group (Figure 2A-C). This effect appeared to be driven by a
sharp decrease in the number of active neurons in the ANI group at the 4 hour session following
administration of ANI, despite these mice displaying comparable levels of freezing to CTRL
mice at this time point (Figure 2D). Interestingly, this accelerated turnover in cells was only
temporary and was followed by a decrease in the cell turnover rate from the 4 hour to day 1
session for ANI compared to CTRL mice (Figure 2B-C). Importantly, we observed no difference
in the mean height of calcium transients for neurons active before, during, or after ANI
administration, indicating that the observed decrease in the number of active neurons was not
due to differences in Ca2+ levels or depletion of the GCaMP protein (Figure S7D-E). To confirm
that this decrease in activity was not due to suppression of network activities, such as those
reported for intracranial infusions of protein synthesis inhibitors in anesthetized or immobile
rodents (Barondes & Cohen, 1966; Sharma et al., 2012), we performed extracellular recordings
in a freely-moving rat implanted with a linear probe in the CA1 region of the hippocampus
following administration of anisomycin. We observed preserved theta activity, theta modulation
of spiking, as well as robust sharp wave ripple activity for up to 5 hours following anisomycin
administration (Figure S7A-C). These observations therefore indicate that blocking protein
synthesis following learning does not abolish all neuronal activity but instead reduces of the
number of active neurons, which may be those specifically involved in learning.

We next hypothesized that arresting protein synthesis, which disrupts the permanence of newly
formed place fields (Agnihotri et al., 2004), would likewise impair the long-term stability of CFC-
related remapping (Moita et al., 2004; Wang et al., 2012). We therefore assessed place field
remapping within and across epochs by correlating event rate maps for all neurons active
between two sessions (Figure 2E-F) following administration of anisomycin vs. saline. We
noticed that place fields recorded in the Learners group exhibited very low across-session
correlations in the neutral arena throughout the experiment (Figure S3), which could indicate
remapping. However, in a separate study, we observed what appeared to be remapping but
what was actually a coherent rotation of all place fields between two recording sessions (Kinsky
et al., 2018). This manifests when all place fields rotate to the same degree around a singular
point, indicative of confusion in an animal's axis of orientation, such as between west and north
(Keinath et al., 2017). Importantly, such angular disorientation does not impair the ability of mice
to identify specific places (Julian et al., 2015), and between-cell firing relationships are
maintained during coherent rotations whereas they are scrambled following remapping.
Coherent rotations match the geometry of the environment, e.g. producing rotations in 60
degree increments in a triangle versus 90 degrees in a square. Therefore, since both shock and
neutral arenas were square, we rotated maps from each session in 90 degree increments and

Figure 2: Anisomycin accelerates cell turnover and stifles learning-related place field remapping A) Cell overlap ratio with Day -2 session, CTRL group. Blue = within shock arena, red = shock v. neutral arena. **B)** Same as A) but for ANI group. **C)** Change in overlap ratios from a and b. *p=0.028, **p=0.00024 two-sided t-test. **D)** Number of active neurons observed each day, normalized to day -1. *p=0.039 two-sided t-test. **E)** Stable place field. (top) Example mouse trajectory (black) with calcium activity (red) overlaid for the same cell from day -2 to -1 in shock arena, (bottom) occupancy normalized rate maps for the same cells. **F)** Remapping place field. Same as E) but for a different cell that remaps from day -2 to day 1 in the shock arena. **G)** Place field correlations for all mice before shock (Days -2 and -1). *p<0.0032 Learners vs ANI and Non-Learners vs ANI, **p=1.9e-6 Non-Learners vs ANI, k-s test after Bonferroni correction. **H)** Same as G) but for days after shock. *p=0.045 Learners vs Non-Learners and p=0.0059 Non-Learners vs ANI, +p=0.1 Learners vs ANI. **I)** Same as G) but to assess learning-related remapping from before to after shock. *p=0.00021 Learners vs Non-Learners and p=3.4e-11 Non-Learners vs ANI, +p=0.12 Learners vs ANI. **J)** Place field correlations before shock broken down by mouse. **K)** Same as J) but for sessions after shock. **L)** Same as J) but from before to after shock. p=0.002 mixed ANOVA, group x arena interaction. *p=0.025 Non-Learners vs ANI, **p=0.013 Learners vs Non-Learners, ***p=0.034 Learners vs ANI, post-hoc pairwise t-test after Bonferroni correction.

used only the orientation that produced the highest correlation between sessions. We found
generally higher correlations compared to using non-rotated maps (Figure 2G-H, J-K vs. Figure
S3F-G), confirming that coherent rotations occurred in many recording sessions. In particular,
place field correlations were high both before learning and after consolidation (Figure 2J-K),
though Learner correlations trended toward lower stability after shock (Figure 2H,K). We then
compared place fields from before to after learning to assess learning-related remapping. In
agreement with previous studies (Moita et al., 2004; Wang et al., 2012), Learner place fields
remapped, as indicated by lower correlations in the shock arena compared to the other groups
(Figure 2I,L). Interestingly, compared to the other groups, Non-Learners exhibited lower place
map correlations in the neutral arena, indicating paradoxically stable place fields in the shock
arena but remapping in the neutral arena (Figure 2I,L). If learning causes remapping, this
double dissociation indicates that Non-Learner memory deficits might result from improperly
associating the neutral arena with shock. In contrast, the place maps of mice in the ANI group

displayed high correlations in all arenas, including after learning. This deficit in remapping
following anisomycin therefore indicates that protein synthesis is required to stabilize the set of
place fields which emerge following learning to support memory consolidation. Overall, these
observations indicate that remapping is necessary for the creation of context-specific memories
and that a lack of remapping or improperly remapping may underlie the memory deficits
observed in Non-Learners and in the ANI group.

Figure 3: ANI administration suppresses the development of coordinated freeze-related neural activity. A) and B) Example traces from two freeze-cells which exhibit coordinated activity prior to freezing event during the day 1 memory recall session in the shock arena. Red = putative spiking activity, pink = cell shown in C, blue = cell shown in E. **C)** Proportion of freeze-tuned cells detected each day across all groups. Green = freezing epochs. **D) and E)** Example Learner freeze-tuned cells identified during the 4 hour (D) or day 1 (E) memory test tracked across sessions. Peri-event calcium activity rasters are centered on freeze onset time (solid green). Dashed green = baseline calcium event probability, red solid = peri-freeze calcium event probability, bins with p<0.01 (circular permutation test) noted with red bars at top. D/E corresponds to pink/blue cells shown in A-B. Bold = session with significant freeze-tuning. **F)** Same as D and E but for ANI mouse shock cell identified during the 4 hour session. **G)** Change in peak peri-freeze calcium event probability for all freeze-tuned cells detected during the 4 hour session. **H)** Same as G but for freeze-tuned cells detected during Day 1 recall session. p < 0.02 1-way ANOVA each day separately, *p=0.02, **p=0.001, ***p=0.0006 post-hoc Tukey test. **I)** Freeze-tuned cells exhibit increased covariance in the CTRL compared to the ANI group. Mean covariance of freeze-tuned cells from each session shown. p=0.016 two-way ANOVA (Time). *p=0.018 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **J)** Small but significant increase in covariance of all cells for CTRL mice during the 1 day recall session. p=0.0015 (Time), 0.005 (Group), 0.036 (Group x Time) two-way ANOVA. *p=0.014, **p=0.008, ***p=0.004 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **K)** Mean covariance of freeze-tuned cells detected during the 4 hour session tracked across sessions. P=0.005 (Group), 0.04 (Group x Time) two-way ANOVA. *p=0.014 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **L)** Same as K but for freeze-tuned cells detected during the day 1 recall session. p=0.0003 (Group) two-way ANOVA. *p=0.016 post-hoc pairwise t-test (two-sided) after Bonferroni correction.

In addition to the spatial code, hippocampal neural activity also reflects non-spatial variables in
a task (McKenzie et al., 2014; Muzzio et al., 2009; Wood et al., 1999). We noticed that many
hippocampal neurons exhibited calcium transients around the time that mice froze (Figure 3A-B,
Figure S6A-C) in line with recent studies (Lee & Han, 2022; Schuette et al., 2020). Neurons
which exhibited a significant increase in calcium event probability +/- 2 sec from a freezing
epoch we therefore dubbed freeze-tuned cells (Figure 3D-F). We observed similar proportions
of freeze-tuned cells across all groups and recording sessions, even before CFC (Figure 3C,
Figure S6D). However, we noticed that freeze-tuned cells appeared to activate more reliably
around freezing epochs following CFC for Learners compared to similar immobility epochs in
Non-Learners and in the ANI group (Figure 3D-E vs. Figure 3F and Figure S6B-C). Therefore,
CFC learning may cause an increase in the reliable firing of previously identified immobility
signaling cells, as primarily seen in CA2, but also in CA1 (Kay et al., 2016).

To quantify changes in within-cell freeze-turning reliability, we tracked the peak, peri-freeze
calcium event probability of each freeze cell backwards and forwards in time from the 4 hour
recall session. We hypothesized that freeze-tuning reliability would increase immediately
following learning at the 4 hour session and remain stable thereafter (see Figure 3D).
Surprisingly, freeze-tuning reliability did not significantly change from the day before to the day
after the 4 hour session for any group (Figure 3G). However, when we tracked cells backward
and forward from the subsequent day 1 recall session, we found that freeze cells exhibited
much higher tuning stability spanning from the 4 hour recall to the day 2 recall sessions in
Learners than did these cells in other groups (Figure 3H). This indicates that in Learners,
reliable freeze-tuning begins to emerge between 4 hours and 1 day following learning and was
maintained thereafter (see Figure 3E for exemplar cell). In contrast, freeze-tuning in mice from
the ANI group and Non-Learners was more transient and unreliable across all time spans.

Lastly, we investigated whether the higher reliability of freeze-cells in Learners translated to
increased co-activity (see Figure 3A-B) in this neuronal subpopulation. Freeze-cell covariance
increased incrementally starting at the 4 hour session for all mice; covariance remained high on
days 1 and 2 for the CTRL group (Learner and Non-Learners combined) but returned to
baseline for mice in the ANI group (Figure 3I). A similar increase in covariance for the CTRL
group was also observed across all cells (Figure 3J). These effects were driven by increased
covariance for Learners but not Non-Learners (Figure S6J-K). Importantly, the ANI group also
displayed increased covariance between all cells at the 4 hour session, but not at subsequent
time points which have been shown to require post-learning protein synthesis for consolidation.
We observed consistent results when we downsampled the number of freezing events following
learning to match that on days -2 and -1 (Figure S6F-G). For freeze cells, but not all cells, the
increased covariance was driven by peri-freeze neural activity (Figure S6H-I). In these analyses
we compared freeze-cells identified independently on each day of the experiment. We next
performed additional analyzes focused on the same set of freeze-tuned cells conditioned on
their activity in specific sessions. Freeze-cells identified at the 4 hour session displayed
increased covariance from the days before training for the CTRL, but not the ANI group (Figure
3K). Freeze-cells that were identified on the day 1 recall test exhibited increased covariance on
days 1 and 2 for the CTRL, but not the ANI group (Figure 3L). However, these cells did not
exhibit increased covariance during the previous day's 4 hour session, indicating that even
though freeze-tuning begins to emerge immediately following learning, cell connections continue
to reorganize further to form coordinated ensembles. These results indicate that freeze-related
tuning emerges following learning and continues to take shape in the ensuing hours until it

stabilizes one day later. Importantly, the coordination of these freeze-tuned cells into neuronal
ensembles requires protein synthesis.

Conclusions:

The results of our study provide evidence that HPC spatial representations support contextual
memory formation and consolidation. We found that between-arena neural overlap correlated
strongly with across-day neural overlap (Figure 1E), suggesting that the neural basis for
contextual discrimination and representational drift may share the same mechanism
(Hainmueller & Bartos, 2018; Kinsky et al., 2020; Mankin et al., 2012). This inherent neural
variability in HPC neuron dynamics *prior to learning* impacts the specificity of contextual fear
memory (Figure 1H). Based on these observations, we propose that the ability of the HPC to
form distinct representations of two arenas despite overlapping contextual qualities allows for
the formation of discrete contextual memories that could underlie why some mice learn while
others do not. Conversely, the similarity of neural representations *after consolidation* could
serve as a biological signature of memory generalization (Figure 1I). We saw that anisomycin
acutely suppressed the number of active cells immediately following learning (Figure 2D). Given
that remapping is observed in Learners during this time window, we speculate that arresting
protein synthesis may temporarily block the formation of new place-fields in the subset of cells
that undergo remapping following CFC, potentially by preventing the formation and
strengthening of new neuronal connections. This idea is resonant with the notion that more
excitable/active neurons are preferentially involved in memory trace formation (Rashid et al.,
2016; Sweis et al., 2021). While such reduced activity could also potentially result from the
blockage of constitutive protein translation (Scavuzzo et al., 2019), our study does not support
this view as we did not observe a global shutdown of neural activity (Figure S7). In addition to
this suppression, anisomycin's amnestic effects coincided with a reduction in learning-related
remapping, effectively halting HPC contextual representations in their prior state (Figure 2L).

Lastly, we confirm that a subset of hippocampal cells exhibits freeze tuning following fear
conditioning (Schuette et al., 2020) and that the organization of these cells into reliable, co-
active neuronal ensembles increases following learning (Lee & Han, 2022) and requires protein
synthesis (Figure 3H-I). Importantly, covariance increased incrementally for both ANI and CTRL
groups at the 4 hour recall session, a time point at which protein synthesis is not required for the
induction of (early-phase) long-term potentiation *in vitro* (Frey & Morris, 1997; Nguyen et al.,
1994)*. However, at later time points, when protein synthesis *is* required to maintain (late-phase)
long-term potentiation (Frey et al., 1993; Nguyen et al., 1994), between-cell covariance
remained high for the CTRL group only (Figure 3G-J). Therefore, in addition to maintaining
learning-related synaptic changes over hours to days, protein synthesis might also be
instrumental for producing coordinated firing of large groups of neurons. A previous study
demonstrated that synchronous optogenetic stimulation of engram neurons tagged during
learning could artificially reactivate a fear memory even when normal long-term recall of the fear
memory was blocked by anisomycin (Ryan et al., 2015). Our results provide a parsimonious
explanation for these results by demonstrating that anisomycin halts the co-firing of freeze-
tuned cells, weakening the capacity of these neurons to transmit behavior-related information to
downstream regions through coincident firing (Lisman, 1997) and impairing their ability to trigger
memory recall (Ryan & Frankland, 2022). Future work will disentangle the differential
contributions that halting protein synthesis versus suppressing neural activity play in disrupting
memory consolidation.

Overall, our results indicate that protein synthesis is necessary for forming new stable spatial
representations of an aversive context following learning and for producing coordinated activity
of freeze-tuned neurons.

Acknowledgments: First and foremost, we would like to thank Howard Eichenbaum who helped
conceive and design this study before his unfortunate passing in 2017. We would also like to
thank Sam McKenzie for his help during early experimental design. We would like to thank
Michael Hasselmo and Ian Davison for their support and feedback while performing the
recordings for this study. Next, we thank Sam Levy, Dave Sullivan, and Will Mau for their
assistance in all phases of calcium imaging throughout. We would like to thank Zach
Pennington for valuable feedback concerning anisomycin preparation and administration, and
Denise Cai and Lucas Carstensen for analysis suggestions. We would like to thank Pho Hale,
Rachel Wahlberg, and Utku Kaya for feedback on the manuscript. We would like to
acknowledge the GENIE Program, specifically Vivek Jayaraman, PhD, Douglas S. Kim, PhD,
Loren L. Looger, PhD, Karel Svoboda, PhD from the GENIE Project, Janelia Research Campus,
Howard Hughes Medical Institute, for providing the GCaMP6f virus. Finally, we would like to
acknowledge Inscopix, Inc. for making single-photon calcium imaging miniscopes widely
available, and specifically Lara Cardy and Vardhan Dani for all their technical support
throughout the experiment. This work was supported by NIH Grants R01 MH052090, R01
MH051570, R01MH117964, NIH NRSA Fellowship 1F32NS117732-01, NIH Early
Independence Award DP5 OD023106-01, an NIH Transformative R01 Award, a Young
Investigator Grant from the Brain and Behavior Research Foundation, a Ludwig Family
Foundation grant, and the McKnight Foundation Memory and Cognitive Disorders award, and
Boston University's Neurophotonics Center,

References

Agnihotri, N. T., Hawkins, R. D., Kandel, E. R., & Kentros, C. G. (2004). The long-term stability
of new hippocampal place fields requires new protein synthesis. *Proceedings of the
National Academy of Sciences*, *101*(10), 3656–3661.
https://doi.org/10.1073/pnas.0400385101

Barondes, S. H., & Cohen, H. D. (1966). Puromycin Effect on Successive Phases of Memory
Storage. *Science*, *151*(3710), 594–595. https://doi.org/10.1126/science.151.3710.594

Barondes, S. H., & Cohen, H. D. (1967). Delayed and Sustained Effect of Acetoxycycloheximide
on Memory in Mice. *Proceedings of the National Academy of Sciences of the United States
of America*, *58*, 157–164.

Barondes, S. H., & Cohen, H. D. (1968). Memory impairment after subcutaneous injection of
acetoxycycloheximide. *Science (New York, N.Y.)*, *160*(3827), 556–557.
http://www.ncbi.nlm.nih.gov/pubmed/5689416

Cai, D. J., Aharoni, D., Shuman, T., Shobe, J., Biane, J., Lou, J., Kim, I., Baumgaertel, K.,
Levenstain, A., Tuszynski, M., Mayford, M., & Silva, A. J. (2016). A shared neural
ensemble links distinct contextual memories encoded close in time. *Nature*, *534*, 115–118.
https://doi.org/10.1038/nature17955

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter, E. R.,
Kerr, R. a, Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K., & Kim, D. S. (2013).

Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295–
300. https://doi.org/10.1038/nature12354

Davis, H. P., & Squire, L. R. (1984). Protein synthesis and memory: A review. *Psychological*
Bulletin, *96*(3), 518–559. https://doi.org/10.1037/0033-2909.96.3.518

Frey, U., Huang, Y.-Y., & Kandel, E. R. (1993). Effects of cAMP Simulate a Late Stage of LTP in
Hippocampal CA1 Neurons. *Science*, *260*(5114), 1661–1664.
https://doi.org/10.1126/science.8389057

Frey, U., & Morris, R. G. M. (1997). Synaptic tagging and long-term potentiation. *Nature*,
385(6616), 533–536. https://doi.org/10.1038/385533a0

Frey, U., & Morris, R. G. M. (1998). Synaptic tagging: Implications for late maintenance of
hippocampal long- term potentiation. *Trends in Neurosciences*, *21*(5), 181–188.
https://doi.org/10.1016/S0166-2236(97)01189-2

Ghosh, K. K., Burns, L. D., Cocker, E. D., Nimmerjahn, A., Ziv, Y., Gamal, A. el, & Schnitzer, M.
J. (2011). Miniaturized integration of a fluorescence microscope. *Nature Methods*, *8*(10),
871–878. https://doi.org/10.1038/nmeth.1694

Hainmueller, T., & Bartos, M. (2018). Parallel emergence of stable and dynamic memory
engrams in the hippocampus. *Nature*, *558*(7709), 292–296. https://doi.org/10.1038/s41586-
018-0191-2

Huang, Y. Y., Li, X. C., & Kandel, E. R. (1994). cAMP contributes to mossy fiber LTP by
initiating both a covalently mediated early phase and macromolecular synthesis-dependent
late phase. *Cell*, *79*(1), 69–79. https://doi.org/10.1016/0092-8674(94)90401-4

Julian, J. B., Keinath, A. T., Muzzio, I. A., & Epstein, R. A. (2015). Place recognition and
heading retrieval are mediated by dissociable cognitive systems in mice. *Proceedings of*
the National Academy of Sciences, *112*(20), 6503–6508.
https://doi.org/10.1073/pnas.1424194112

Kay, K., Sosa, M., Chung, J., Karlsson, M. P., Larkin, M., Grossrubatscher, I., & Frank, L. M.
(2016). A hippocampal network for spatial coding during immobility. *Nature*, *531*, 185–190.
https://doi.org/10.1038/nature17144

Keinath, A. T., Julian, J. B., Epstein, R. A., & Muzzio, I. A. (2017). Environmental Geometry
Aligns the Hippocampal Map during Spatial Reorientation. *Current Biology*, *27*(3).
https://doi.org/http://dx.doi.org/10.1016/j.cub.2016.11.046

Kinsky, N. R., Mau, W., Sullivan, D. W., Levy, S. J., Ruesch, E. A., & Hasselmo, M. E. (2020).
Trajectory-modulated hippocampal neurons persist throughout memory-guided navigation.
Nature Communications, 1–14. https://doi.org/10.1038/s41467-020-16226-4

Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E., & Eichenbaum, H. B. (2018).
Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales.
Current Biology, *28*(22), 1–11. https://doi.org/10.1016/J.CUB.2018.09.037

Lee, H.-S., & Han, J.-H. (2022). Activity patterns of individual neurons and ensembles correlated
with retrieval of a contextual memory in the dorsal CA1 of mouse hippocampus. *The*

Journal of Neuroscience, *43*(1), JN-RM-1407-22. https://doi.org/10.1523/jneurosci.1407-
22.2022

Lisman, J. E. (1997). Bursts as a unit of neural information: Making unreliable synapses reliable.
Trends in Neurosciences, *20*(1), 38–43. https://doi.org/10.1016/S0166-2236(96)10070-9

Mankin, E. A., Sparks, F. T., Slayyeh, B., Sutherland, R. J., Leutgeb, S., & Leutgeb, J. K.
(2012). Neuronal code for extended time in the hippocampus. *Proceedings of the National*
Academy of Sciences, *109*(47), 19462–19467. https://doi.org/10.1073/pnas.1214107109/-
/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1214107109

Mau, W., Sullivan, D. W., Kinsky, N. R., Hasselmo, M. E., Howard, M. W., & Eichenbaum, H. B.
(2018). The Same Hippocampal CA1 Population Simultaneously Codes Temporal
Information over Multiple Timescales. *Current Biology*, *28*(10), 1499-1508.e4.
https://doi.org/10.1016/j.cub.2018.03.051

McKenzie, S., Frank, A. J., Kinsky, N. R., Porter, B., Rivière, P. D., & Eichenbaum, H. B. (2014).
Hippocampal representation of related and opposing memories develop within distinct,
hierarchically organized neural schemas. *Neuron*, *83*(1), 202–215.
https://doi.org/10.1016/j.neuron.2014.05.019

Moita, M. A. P., Rosis, S., Zhou, Y., LeDoux, J. E., & Blair, H. T. (2004). Putting Fear in Its
Place: Remapping of Hippocampal Place Cells during Fear Conditioning. *Journal of*
Neuroscience, *24*(31), 7015–7023. https://doi.org/10.1523/JNEUROSCI.5492-03.2004

Muir, D. R., & Kampa, B. M. (2015). FocusStack and StimServer: a new open source MATLAB
toolchain for visual stimulation and analysis of two-photon calcium neuronal imaging data.
Frontiers in Neuroinformatics, *8*(January), 1–13. https://doi.org/10.3389/fninf.2014.00085

Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial
firing of hippocampal complex-spike cells. *The Journal of Neuroscience*, *7*(7), 1951–1968.
http://www.ncbi.nlm.nih.gov/pubmed/3612226

Muzzio, I. A., Levita, L., Kulkarni, J., Monaco, J. D., Kentros, C. G., Stead, M., Abbott, L. F., &
Kandel, E. R. (2009). Attention enhances the retrieval and stability of visuospatial and
olfactory representations in the dorsal hippocampus. *PLoS Biology*, *7*(6).
https://doi.org/10.1371/journal.pbio.1000140

Nguyen, P., Abel, T., & Kandel, E. R. (1994). Requirement of a critical period of transcription for
induction of a late phase of LTP. *Science*, *265*(5175), 1104–1107.
https://doi.org/10.1126/science.8066450

Olypher, A. v, Lánský, P., Muller, R. U., & Fenton, A. A. (2003). Quantifying location-specific
information in the discharge of rat hippocampal place cells. *Journal of Neuroscience*
Methods, *127*(2), 123–135. https://doi.org/10.1016/S0165-0270(03)00123-7

Rashid, A. J., Yan, C., Mercaldo, V., Hsiang, H. L., Park, S., Cole, C. J., Cristofaro, A. de, Yu,
J., Ramakrishnan, C., Lee, S. Y., Deisseroth, K., Frankland, P. W., & Josselyn, S. A.
(2016). Competition between engrams influences fear memory formation and recall.
Science, *353*(6297), 383–388. https://doi.org/10.1126/science.aaf0594

Resendez, S. L., Jennings, J. H., Ung, R. L., Namboodiri, V. M. K., Zhou, Z. C., Otis, J. M., Nomura, H., McHenry, J. A., Kosyk, O., & Stuber, G. D. (2016). Visualization of cortical, subcortical and deep brain neural circuit dynamics during naturalistic mammalian behavior with head-mounted microscopes and chronically implanted lenses. *Nature Protocols*, *11*(3), 566–597. https://doi.org/10.1038/nprot.2016.021

Ryan, T. J., & Frankland, P. W. (2022). Forgetting as a form of adaptive engram cell plasticity. *Nature Reviews Neuroscience*, *23*(3), 173–186. https://doi.org/10.1038/s41583-021-00548-3

Ryan, T. J., Roy, D. S., Pignatelli, M., Arons, A., & Tonegawa, S. (2015). Engram cells retain memory under retrograde amnesia. *Science*, *348*(6238), 1007–1013. https://doi.org/10.1126/science.aaa5542

Scavuzzo, C. J., LeBlancq, M. J., Nargang, F., Lemieux, H., Hamilton, T. J., & Dickson, C. T. (2019). The amnestic agent anisomycin disrupts intrinsic membrane properties of hippocampal neurons via a loss of cellular energetics. *Journal of Neurophysiology*, *122*(3), 1123–1135. https://doi.org/10.1152/jn.00370.2019

Schafe, G. E., Nadel, N. v., Sullivan, G. M., Harris, A., & LeDoux, J. E. (1999). Memory consolidation for contextual and auditory fear conditioning is dependent on protein synthesis, PKA, and MAP kinase. *Learning and Memory*, *6*(2), 97–110. https://doi.org/10.1101/lm.6.2.97

Schuette, P. J., Reis, F. M. C. V., Maechler, P., Chakerian, M., Torossian, A., Blair, G., Wang, W., Blair, H. T., Fanselow, M. S., Kao, J. C., & Adhikari, A. (2020). Long-term characterization of hippocampal remapping during contextual fear acquisition and extinction. *The Journal of Neuroscience*, *September*, JN-RM-1022-20. https://doi.org/10.1523/jneurosci.1022-20.2020

Sharma, A. v., Nargang, F. E., & Dickson, C. T. (2012). Neurosilence: Profound Suppression of Neural Activity following Intracerebral Administration of the Protein Synthesis Inhibitor Anisomycin. *The Journal of Neuroscience*, *32*(7), 2377–2387. https://doi.org/10.1523/JNEUROSCI.3543-11.2012

Squire, L. R., & Barondes, S. H. (1973). Memory impairment during prolonged training in mice given inhibitors of cerebral protein synthesis. *Brain Research*, *56*(C), 215–225. https://doi.org/10.1016/0006-8993(73)90336-3

Squire, L. R., & Barondes, S. H. (1974). Anisomycin, like other inhibitors of cerebral protein synthesis, impairs 'long-term' memory of a discrimination task. *Brain Research*, *66*(2), 301–308. https://doi.org/10.1016/0006-8993(74)90148-6

Sweis, B. M., Mau, W., Rabinowitz, S., & Cai, D. J. (2021). Dynamic and heterogeneous neural ensembles contribute to a memory engram. *Current Opinion in Neurobiology*, *67*, 199–206. https://doi.org/10.1016/j.conb.2020.11.017

Vöröslakos, M., Petersen, P. C., Vöröslakos, B., & Buzsáki, G. (2021). Metal microdrive and head cap system for silicon probe recovery in freely moving rodent. *ELife*, *10*. https://doi.org/10.7554/eLife.65859

Wang, M. E., Wann, E. G., Yuan, R. K., Ramos Álvarez, M. M., Stead, S. M., & Muzzio, I. A.
(2012). Long-term stabilization of place cell remapping produced by a fearful experience.
The Journal of Neuroscience, *32*(45), 15802–15814.
https://doi.org/10.1523/JNEUROSCI.0480-12.2012

Wood, E. R., Dudchenko, P. A., & Eichenbaum, H. B. (1999). The global record of memory in
hippocampal neuronal activity. *Nature*, *397*(6720), 613–616. https://doi.org/10.1038/17605

Yger, P., Spampinato, G. L. B., Esposito, E., Lefebvre, B., Deny, S., Gardella, C., Stimberg, M.,
Jetter, F., Zeck, G., Picaud, S., Duebel, J., & Marre, O. (2018). A spike sorting toolbox for
up to thousands of electrodes validated with ground truth recordings in vitro and in vivo.
eLife, *7*, 1–23. https://doi.org/10.7554/eLife.34518

<u>Methods:</u>

Animals

Sixteen (n = 10 controls, 6 anisomycin) male C57/BL6 mice (Jackson Laboratories), age 16 to
22 weeks during behavioral and imaging experiments and weighing 25-32g were used in this
study. Three mice were excluded after performing this study: one mouse after histology
revealed the GRIN lens implant and viral expression to be medial to the intended imaging, while
the other two were excluded due to unstable/overexpression of GCaMP that produced aberrant
calcium activity which emerged toward the end of the experiment. After exclusion of these mice,
we retained 8 mice in the control (CTRL) group and 5 mice in the anisomycin group (ANI) group.
Additionally, behavioral video tracking files for one CTRL mouse were corrupted during
recording during all neutral field recordings from day 0 on: this mouse was excluded from all
analyses which required using behavior in the neutral arena (e.g., place field correlations and
any analyses where the CTRL group was split into Learners and Non-Learners). Mice were
socially housed in a vivarium on a 12 hour light-dark cycle with 1-3 other mice prior to surgery
and were housed singly thereafter. Mice were given free access to food and water throughout
the study. All procedures were performed in compliance with the guidelines of the Boston
University Animal Care and Use Committee.

One male Long Evans rat, 10 months old and weighing ~480g, was used for the
electrophysiological recording in this study. Rats were socially housed in a vivarium on an
adjusted 12 hour light-dark cycle (lights on at noon, off at midnight) with 1-3 other rats prior to
surgery and given free access to food and water throughout the study. All procedures were
performed in compliance with the guidelines of the University of Michigan Animal Care and Use
Committee.

Viral Constructs

For mice experiments we used an AAV9.*Syn*.GcaMP6f.WPRE.SV40 virus from the University of
Pennsylvania Vector Core/Addgene with an initial titer of ~$4x10^{12}$ GC/mL and diluted it into
sterilized potassium phosphate buffered saline (KPBS) to a final titer of ~$2\text{-}4x10^{12}$ GC/mL for
injection.

For rat experiments, we used an pGP.AAV9.*Syn*.GcaMP7f.WPRE.SV40 virus from the
University of Pennsylvania Vector Core/Addgene with an initial titer of $2.6x10^{13}$ GC/mL and
diluted it into sterilized phosphate buffered saline (PBS) to a final titer of $2.6x10^{12}$ GC/mL for
injection. Due to poor expression no imaging was performed.

Stereotactic Surgery

We performed two stereotactic surgeries and one base-plate implant on naïve mice, aged 3-8
months, according to previously published procedures (Kinsky et al., 2018; Resendez et al.,
2016). Both surgeries were performed under 1-2% isoflurane mixed with oxygen. Mice were
given 0.05mL/kg buprenorphine (Buprenex) for analgesia (subcutaneously, SC), 5.0mL/kg of
the anti-inflammatory drug Rimadyl (Pfizer, SC), and 400mL/kg of the antibiotic Cefazolin
(Pfizer, SC) immediately after induction. They were carefully monitored to ensure they never
dropped below 80% of their pre-operative weight during convalescence and received the same
dosage of Buprenex, Cefazolin, and Rimadyl twice daily for three days following surgery. In the

first surgery, a small craniotomy was performed at AP -2.0, ML +1.5 (right) and 250nL of
GcaMP6f virus (at the titer noted below) was injected 1.5mm below the brain surface at
40nL/min using a 1µL Hamilton syringe and infusion pump. The needle remained in place a
minimum of 10 minutes after the infusion finished at which point it was slowly removed, the
mouse's scalp was sutured, and the mouse was removed from anesthesia and allowed to
recover.

3-4 weeks after viral infusion, mice underwent second surgery to attach a gradient index (GRIN)
lens (GRINtech, 1mm x 4mm). After performing an ~2mm craniotomy around the implant area,
we carefully aspirated cortex using blunted 25ga and 27ga needles under constant irrigation
with cold, sterile saline until we visually identified the medial-lateral striations of the corpus
callosum. We carefully removed these striations using a blunted 31ga needle while leaving the
underlying anterior-posterior striations intact, after which we applied gelfoam for 5-10 minutes to
stop any bleeding. We then lowered the GRIN lens to 1.1mm below bregma. Note that this
entailed pushing down ~50-300µm to counteract brain swelling during surgery. We then applied
Kwik-Sil (World Precision Instruments) to provide a seal between skull and GRIN lens and then
cemented the GRIN lens in place with Metabond (Parkell), covered it in a layer of Kwik-Cast
(World Precision Instruments), and then removed the animal from anesthesia and allowed him
to recover after removing any sharp edges remaining from dried Metabond with a dental drill
and providing any necessary sutures.

Finally, after ~2-4 weeks we performed a procedure in which the mouse was put under
light anesthesia to attach a base plate for easy future attachment of a miniature epifluorescence
microscope (Ghosh et al., 2011, Inscopix, Inc.). Importantly, no tissue was cut during this
procedure. After induction, we attached the base plate to the camera via a set screw, set the
camera's focus level at ~1/3 from the bottom of its range, and carefully lowered the camera
objective and aligned it to the GRIN lens by eye, and visualized fluorescence via nVistaHD until
we observed clear vasculature and putative cell bodies expressing GcaMP6f (Resendez et al.,
2016). To counteract downward shrinking during curing, we then raised the camera up ~50µm
before applying Flow-It ALC Flowable Composite (Pentron) between the underside of the
baseplate and the cured Metabond on the mouse's skull. After light curing, we applied opaque
Metabond over the Flow-It ALC epoxy to the sides of the baseplate to provide additional
strength and to block ambient light infiltration. Mice were allowed to recover for several days
prior to habituation to camera attachment and performance of the behavioral task outlined
below. In the event that we did not observe clear vasculature and cell bodies when we first
visualized fluorescence we covered the GRIN lens with Kwik-Cast and removed the mouse from
anesthesia without attaching the baseplate. We then waited an additional week and repeated
the steps above.

For rats, we performed two surgeries in a similar manner as described above for mice.
However, rats were administered pre-operative and post-operative Meloxicam orally for
analgesia (in lieu of Buprenex) and triple-antibiotic was applied locally (in lieu of Cefazolin
injections) to the incision at the end of surgery. Meloxicam was additionally administered for two
days post-surgery during recovery, and animals were monitored daily for a minimum of seven
days during recovery. 0.4mL of a lidocaine/bupivacaine cocktail were given under the scalp to
provide local anesthesia at the incision site. In the first surgery, 1000nL of GcaMP7f virus was
infused in the prelimbic cortex at the center of a 1mm craniotomy (AP + 2.9, ML + 3.6, from
Bregma, DV -3.0 at an 18 degree angle from top of brain). Following infusion, ~1.5 mm of

overlying cortex was removed and a 23ga needle was lowered to ~500μm above the target site.
Then, a 0.6 x 7 mm GRIN lens was lowered to 3.0mm below the top of the brain, the area
between the skull and lens was sealed with Kwik-Sil, and the lens was affixed to the skull with
Vivid-Flow light-curable composite (Pearson Dental) and Metabond (Parkell). The lens was then
covered in Kwik-Sil for protection. During this surgery, ground and reference screws were also
placed over the cerebellum and a 3d printed crown base was attached to the rat's skull
(Vöröslakos et al., 2021) to which crown walls and top were connected and to further protect the
lens and future microdrive/probe implant. The rat was screened for fluorescence 8-12 weeks
later, but no cell dynamics were observed so no imaging equipment was implanted for this rat.

16 weeks later, the rat was again given pre-operative Meloxicam and anesthetized under
isoflurane. The crown walls were removed and a 1.0mm craniotomy was performed at AP-4.8,
ML+3.6 from bregma. After removing dura and stopping bleeding with cold, sterile saline, a
NeuroNexus A1x32-5mm-50-177 probe, attached to a metal microdrive, was implanted at 2.3
mm below the brain surface and the metal drive base was attached to the skull with Unifast light
cured dental epoxy (Henry-Schein). The craniotomy was sealed with Dow-Sil, the probe was
protected with dental wax, and the ground and reference wires were connected to the probe
electronic interface board (EIB). The crown walls were re-attached, the EIB was connected to
the crown walls, and the rat was removed from isoflurane and allowed to recovery. The rat was
monitored daily for 7 days prior to recording, during which the probe was lowered ~1mm until
sharp wave ripples and spiking activity were visualized indicating localization of the probe in the
CA1 cell layer.

Histology procedures

Hippocampal slices were prepared following extraction from mice in accordance with the
standard methods and guidelines of the Boston University Animal Care and Use Committee. In
brief, mice were euthanized with Euthasol (Virbac), transcardially perfused with
paraformaldehyde (PFA), and decapitated. Following extraction, brains were placed in PFA for
approximately 48 hours before undergoing sectioning. Brains that were sliced using a Cryostat
underwent an additional step of sucrose cryoprotection and subsequent freezing in -80C. Brains
were mounted to the slicing platform using Tissue Tek O. C. T. (Sakura) and kept at -30C
throughout sectioning. 50μm slices were collected across the entire aspiration site in the dorsal
hippocampus region. Brains that were sliced using a vibratome were stabilized using super glue
and submerged in 1% PBS. A Leica VT1000 S vibratome was equipped with a platinum coated
double edged blade (Electron Microscopy Sciences , Cat. #72003-01) and set to a maximal
speed of 0.9mm/s for collecting 50 μm slices. Slices prepared from both the cryostat and
microtome were directly mounted onto (type of slides go here) and cover-slipped using DAPI
following sectioning. No histology was performed in the rat study.

Behavioral Paradigm

Prior to surgery mice were handled to habituate them subsequent camera attachment. 3-7 days
following base plate attachment surgery we conditioned mice to the imaging procedures by
further handling them for 5-10 minutes for a minimum three days. During this handling a plastic
"dummy" microscope (Inscopix) of approximately the same size/weight as the imaging camera
was attached to each mouse's head and remained on his head for 1-2 hours in his home cage.
When it became easy to attach the scope to the mouse's head a real imaging miniscope was
attached to head and an optimal focus plane chosen. We then recorded three 5 minute imaging

videos at this focus and +/- ¼ turn (~25μm) in the mouse's home cage. These movies were
processed as described in the Image Acquisition and Processing section and an optimal zoom
was chosen based on whichever focus plane maximized cell yield and produced clear looking
cell bodies. Animals were then placed in a novel environment with a different size and shape
compared to the experimental environments for a 10 minute session to habituation them to the
general experimental outline and ensure that they explored novel arenas.

Following habituation to the imaging procedures mice performed a contextual fear
conditioning (CFC) task with simultaneous imaging of hippocampus neurons over the course of
10 days. Note that all recording sessions are referred to by their time relative to applying the
mild foot-shock and the arena in which the recording occurred: e.g., Shock Day -2 occurred in
the shock arena two days prior to foot-shock while Neutral 4 hours occurred four hours after
foot-shock in the neutral arena. A typical day (Days -2, -1, 1, 2, and 7) consisted of two separate
10 minute recording blocks/sessions: one in the Neural arena and one in the shock arena. Mice
first explored a square (neutral) arena, placed in the center of a well-lit room, for 10 minutes.
The neutral arena was a square constructed of 3/8" plywood (25cm x 25cm x 15 cm), which was
painted yellow with sealable paint. Additionally, one wall was painted with black horizontal
stripes for visual orientation purposes. The neutral arena was wiped down with 70% ethanol ~10
minutes prior to recording. After 10 minutes of exploration the experimenter took the mouse out
of the arena, leaving the miniscope camera on their head and placed the mouse in its home
cage on a moveable cart upon which it was immediately transported down a short hallway to
second room.

The second room was dimly lit and contained the fear conditioning (shock) arena. The
shock arena (Coulbourn Instruments, Whitehall, PA, USA) consisted of metal-panel side walls,
Plexiglas front and rear walls, and a stainless-steel grid floor composed of 16 grid bars (22cm x
22cm). Following 10 minutes of exploration of the shock arena, mice were removed from the
arena, the camera was removed, and mice were returned to their home cage. Both arenas were
wiped down with 70% ethanol ~10min prior to recording to eliminate any odor cues. Note that
mice always explored the neutral arena first and the shock arena second. For the Day 0
sessions, mice first explored the neutral arena for 10 minutes and were transported to the shock
arena as usual. However, during this session (Shock Day 0) the mouse was immediately given
a single 0.25mA shock and allowed to explore the arena for an additional 60 seconds only
before being removed and returned to his cage. Efficacy of shock was confirmed post-hoc by
eye by the presence of jumping/darting behavior immediately post-shock. The 4 hour session
was identical to the Day -2, 1, 1, 2, and 7 sessions. With the exception of the 4 hour session, all
recording sessions were performed in the first half of the mouse's life cycle while the 4 hour
session occurred in the second half of the light cycle.

On day zero, after the camera was removed and prior to returning to their home cage, mice
received an intraperitoneal injection of either anisomycin (150 mg/kg, Sigma-Aldrich A9789) or
the equivalent amount of vehicle. After injection, they were returned to their cage for 4 hours
until the next recording session began.

Following extensive habituation to a rest box during the seven day recovery period, rat neural
activity and behavior was recorded across ~ 5 hours. Following a 15 minute baseline recording
(PRE) in the rest box, the animal was given an I.P. injection of anisomycin and then immediately
placed on a novel linear track which he explored for 45 minutes (TRACK). The rat was then
placed back into the rest box for 3.5 hours (POST). Following that, the animal was placed on a

second novel track for 45 minutes (TRACK) followed by a brief recording in the rest box
(POST2). Data was acquired continuously throughout with the exceptions of periodic cable
disconnections to perform the I.P. injection, start a new recording epoch, and
disconnect/reconnect cables that became twisted.

Anisomycin

For mice recordings, 25 mg of anisomycin was dissolved into 50 µL of 6N HCl and 500 µL of
1.8%NaCl. ~125 µL of 1N NaOH was then added to the solution followed by 0.1-0.5 µL of 1N
NaOH, testing pH after each addition until a final pH of 7.0 to 7.5 was reached, with a final
concentration of 24-27 mg/mL. In the case that pH rose above 7.5 during titration and and/or the
anisomycin went back into precipitate, small amounts (10-20 µL) of 6N HCl were added until
particles were no longer visible and the titration with 1N NaOH was restarted. Mice were
administered 150mg/kg of anisomycin solution via intraperitoneal injection, or ~0.15-0.18mL for
a typical 30g mouse.

For rat recordings, 100mg of anisomycin was dissolved into 1.6mL of 0.1N HCl (in 0.9% saline).
~240 µL of 1N HCl was added, then 10-12 µL of NaOH was added in 1-2 µL amounts, testing
pH between each step until a pH of 7-7.5 was reached. 0.9% Saline was then added until a final
concentration of 33 mg/mL was reached. Due to a small amount of waste, the final amount
injected was 50mg (1.5 mL) which corresponds to 100 mg/kg for the rat.

Behavioral Tracking and Fear Metrics

We utilized two different camera/software configurations for tracking animal behavior. Both
configurations generated a TTL pulse at the beginning of behavioral tracking to synchronize with
image acquisition. We utilized Cineplex software (v2, Plexon) to track animal location at 30Hz in
the neutral arena. We used FreezeFrame (Actimetrics) to track animal location in the shock
arena at 3.75Hz. Animal location was obtained post-hoc via custom-written, freely available
Python software (www.github.com/wmau/FearReinstatement). We observed inconsistent frame
rates and inaccurate acquisition of behavioral video frames for one mouse in the neutral arena
during the day 0, 4 hour, and day 1-2 sessions. These sessions were excluded from analysis.

Freezing was calculated by first dowsampling neutral position data to 3.75 Hz to match the
sample rate used in the shock arena. We then identified freezing epochs as any periods of 10
consecutive frames (2.67 seconds) or more where the mouse's velocity was less than
1.5cm/second.

Neural Discrimination

631   We evaluated the extent to which each animal's behavior reflected the expression of a context-
632   specific fear memory through a behavioral discrimination index (DI_{beh}), calculated as follows:

$$DI_{beh} = \frac{Frz_{Neutral} - Frz_{Shock}}{Frz_{Neutral} + Frz_{Shock}}$$

Where $Frz_{Neutral}$ and Frz_{Shock} are the percentages of time spent freezing in the neutral and shock
arenas, respectively. Thus, a negative DI_{beh} value indicated more freezing behavior in the shock
arena (indicating successful encoding of a context-specific fear memory), a positive DI_{beh} value
indicated more freezing behavior in the neutral arena, and a DI_{beh} value around zero indicated

equal/low freezing behavior in each arena (indicating the formation of a non-specific or weak
fear memory).

Imaging Acquisition and Processing

Brain imaging data was obtained using nVista HD (Inscopix) at 720 x 540 pixels and a 20 Hz
sample rate. Note that imaging data for one mouse was obtained at 10 Hz. Prior to
neuron/calcium event identification we first pre-processed each movie using Inscopix Imaging
Suite (Inscopix) software. Preprocessing entailed three steps a) motion corrections, and b)
cropping the motion-corrected movie to eliminate any dead pixels or areas with no calcium
activity, and c) extracting a minimum projection of the pre-processed movie for later neuron
registration. We did not analyze one imaging session in which we had to reconnect the camera
cable mid-session and could not synchronize the imaging data with behavioral data. Maximum
projections of imaging movies were made using the Inscopix Imaging Suite or custom-written
functions based off of an open-source MATLAB library (Muir & Kampa, 2015).

Electrophysiological Recordings

652 Data was acquired using an Intan 1028 channel recording system through OpenEphys software
653 into binary format and behavior was tracked via Omnitrack high resolution cameras.

Data Analysis

655 Data analysis was performed in both Python and MATLAB software. Python analysis code is
656 available at https://github.com/nkinsky/Eraser.

Spike sorting and analysis

Electrophysiological recordings were automatically clustered using SpyKING CIRCUS software
(Yger et al., 2018) and units were manually curated in phy. Units were grouped into single units
if they exhibited a clear refractory period and were well-isolated from other putative spikes.
Other units which exhibited a clear waveform but were either poorly isolated or exhibited
refractory period violations were classified as multi-unit activity (MUA). All single units and MUA
were combined and cross-correlograms for the combined activity were created for each epoch
of the recording separately.

Tenaspis

Neuron regions-of-interest (ROIs) and calcium events were identified using a custom written,
open source algorithm employed in MATLAB 2016b called A Technique for Extracting Neuronal
Activity from Single Photon Neuronal Image Sequences (Tenaspis) (Mau et al., 2018). This
procedure was comprehensively documented in Kinsky et al., 2018:

"Tenaspis is open-source and available at: https://github.com/SharpWave/TENASPIS. First,
Tenaspis filters each calcium imaging movie with a band-pass filter per (Kitamura et al., 2015)
to accentuate the separation between overlapping calcium events. Specifically, Tenaspis
smooths the movie with a 4.5 μm disk filter and divides it by another movie smoothed with a
23.6 μm disk filter. Second, it adaptively thresholds each imaging frame to identify separable
pockets of calcium activity, designated as blobs, on each frame. Blobs of activity are accepted
at this stage of processing only if they approximate the size and shape of a mouse hippocampal
neuron, as measured by their radius (min = ~6μm, max = ~11μm), the ratio of long to short axes
(max = 2), and solidity (min = 0.95), a metric used by the *regionprops* function of MATLAB we

employ to exclude jagged/strange shaped blobs. Third, Tenaspis strings together blobs on
successive frames to identify potential calcium transients and their spatial activity patterns.
Fourth, Tenaspis searches for any transients that could results from staggered activity of two
neighboring neurons. It rejects any transients whose centroid travels more than 2.5µm between
frames and whose duration is less than 0.20 seconds. Fifth, Tenaspis identifies the probable
spatial origin of each transient by constructing putative regions-of-interest (ROIs), defined as all
connected pixels that are active on at least 50% of the frames in the transient. Sixth, Tenaspis
creates initial neuron ROIs by merging putative transient ROIs that are discontinuous in time but
occur in the same location. Specifically, it first attempts to merge all ROIs whose centroids are
less than a distance threshold of ~0.6µm from each other. In order to merge two transient ROIs,
the two-dimensional Spearman correlation between the ROIs must yield $r^2 > 0.2$ and $p < 0.01$.
Tenaspis then successively increases the distance threshold and again attempts to merge ROIs
until no more valid merges occur (at a distance threshold of ~3µm, typically). Seventh, Tenaspis
integrates the fluorescence value of each neuron ROI identified in the previous step across all
frames to get that neuron's calcium trace, and then identifies putative spiking epochs for each
neuron. Specifically, it first identifies the rising epochs of any transients identified in earlier
steps. Then, it attempts to identify any missed transients as regions of the calcium trace that
have a) a minimum peak amplitude > 1/3 of the transients identified in step 3, b) a high
correlation ($p < 0.00001$) between active pixels and the pixels of the average neuron ROI
identified in step 6, and b) a positive slope lasting at least 0.2 seconds. Last, Tenaspis searches
for any neuron ROIs that overlap more than 50% and whose calcium traces are similar and
merges their traces and ROIs."

Between Session Neuron Registration

We utilized custom-written, freely available MATLAB code (available at
https://github.com/nkinsky/ImageCamp) to perform neuron registration across sessions in
accordance with previous results (Kinsky et al., 2018). The details of this procedure described in
Kinsky et al. (2018) are reproduced here:

"Neuron registration occurred in two steps: session registration and neuron registration.

Session registration – Prior to mapping neurons between sessions, we determined how
much the imaging window shifted between sessions. In order to isolate consistent features of
the imaging plane for each mouse (such as vasculature or coagulated blood), we created a
minimum projection of all of the frames of the motion-corrected and cropped brain imaging
movie for each recording session. One session ("registered session") was then registered to a
base session using the "imregtform" function from the MATLAB Image Processing Toolbox,
assuming a rigid geometric transform (rotation and translation only) between images, and the
calculated transformation object was saved for future use.

Neuron Registration – Next, each ROI in the registered session was transformed to its
corresponding location in the base session. Each neuron in the base session was then mapped
to the neuron with the closest center-of-mass in the registered session, unless the closest
neuron exceeded our maximum distance threshold of 3 pixels (3.3 µm). In this case the base
session neuron was designated to map to no other neurons in the registered session. If, due to
high density of neurons in a given area, we found that multiple neurons from the base session
mapped to the same neuron in the registered session, we then calculated the spatial correlation

(Spearman) between each pair of ROIs and designated the base session ROI with the highest
correlation as mapping to the registered session ROI.

For multiple session registrations, the same procedure as above was performed for each
session in two different ways. First, we registered each session directly to the first session in the
experiment and updated ROI locations/added new ROIs to set of existing ROIs with each
registration. This helped account for slight day-to-day drift in neurons ROIs due to shifts in
vasculature, build-up of fluid underneath the viewing window, creep/shrinkage of dental cement,
etc. Second, to ensure that neuron ROIs did not drift excessively across sessions we also
performed all the above steps but did NOT update ROI locations allowing us to register each set
of ROIs to those furthest away chronologically. The resulting mappings were then compared
across all sessions, and any neuron mappings that differed between the two methods (e.g.,
ROIs that moved excessively across the duration of the experiment) were excluded from
analysis. Those that remained in the same location were included."

The procedure to assess the quality of across session registration was described in
Kinsky et al. (2018) and is reproduced here: "We checked the quality of neuron registration
between each session-pair in two ways: 1) by plotting the distribution of changes in ROI
orientations between session and comparing it to chance, calculated by shuffling neuron identity
between session 1000 times, and 2) plotting ROIs of all neurons between two sessions and
looking for systematic shifts in neuron ROIs that could lead to false negatives/positives in the
registration." All session-pairs (except those few in which we could not synchronized imaging
and behavioral data as noted above) met the above two criteria and were thus included in our
analysis.

Cells that had calcium activity in the first session (neutral arena) for which we did not identify a
matching neuron in the second session (shock arena) were classified as OFF cells. Likewise,
neurons active in the shock arena with no matching partner in the neutral arena were classified
as ON cells.

All neuron registrations were cross validated by overlaying ROIs from each session and
evaluating their match by eye. In a few cases, we noticed erroneous registrations and adjusted
our between-session neuron alignment by calculating the rigid geometric transformation using
4-5 cell ROIs active in both sessions.

Neural Discrimination Metrics

The extent to which gross hippocampal ensemble activity differed between arenas was
calculated in two ways. First, we calculated the proportion of cells that turned ON and OFF
between arenas divided by the total number of cells active in either arena.

Next, we calculated the extent to which each neuron active in both arenas distinguished
between arenas by changing its event rate in a manner analogous to $DI_{Fr.}$ However, we took the
absolute value to account for the fact that both positive and negative event rate changes could
reflect neural differentiation between arenas. Then, we took the mean across all neurons to
obtain a neural discrimination index (DI_{Neuron}):

761
$$DI_{Neuron} = \left| \frac{ER_{Neutral} - ER_{Shock}}{ERz_{Neutral} + ER_{Shock}} \right|$$

Placefield Analysis

Calcium transients occurring when the mouse was running greater than or equal to
1.5cm/second were spatially binned (1cm by 1cm) and occupancy normalized following which
place fields were identified and quantified in a manner similar to Kinsky et al. (2018),
reproduced here:

"Spatial mutual information (SI) was computed from the following equations, adapted from
(Olypher et al., 2003)

770
$$I_{pos}(x_i) = \sum_{k=0}^{1} P_{k|x_i} \, log \left(\frac{P_{k|x_i}}{P_k} \right)$$

771
$$SI = \sum_{i=1} P_{x_i} I_{pos}(x_i)$$

where:

- P_{xi} is the probability the mouse is in pixel x_i

- P_k is the probability of observing k calcium events (0 or 1)

- $P_{k|xi}$ is the conditional probability of observing k calcium events in pixel $x_{i.}$

The SI was then calculated 1000 times using shuffled calcium event timestamps, and a neuron
was classified as a place cell if it 1) had at least 5 calcium transients during the session, and 2)
the neuron's SI exceeded 95% of the shuffled Sis…We defined the extent of a place field as all
connected occupancy bins whose smoothed event rate exceeded 50% of the peak event rate
occupancy bin."

Since spatial mutual information is biased by the number of samples (Olypher et al., 2003), we
re-sampled the behavioral tracking data to match that of the imaging data (20Hz). This required
up-sampling the shock arena data (3.75Hz->20Hz) and down sampling the neutral arena data
(30Hz->20Hz).

Placefield similarity between sessions was assessed by first smoothing the 2-d occupancy
normalized event rate maps with a gaussian kernel (2.5cm std), flattening the smoothed maps
into a vector, and then performing a Spearman correlation between all neurons active in both
sessions. To quantify chance-level place field similarity we randomly shuffled the mapping
between neurons from the first to the second session before performing the Spearman
correlation. We then repeated this procedure 100 times.

To assess the possibility that the configuration of place fields rotated together coherently
between sessions (Kinsky et al., 2018), we again performed a Spearman correlation but after
rotating the 2-d occupancy map in the second session 90 degrees. Since, due to small camera
distortions, some 2-d occupancy maps were not square, one some occasions we resized
(minimally) the second map to match the size/shape of the first map using the reshape function
in Python's numpy package prior to correlate the two maps. We repeated this in successive 90
degree increments and then took the mean correlation of all neurons that were active in both

sessions to determine the optimal/"best" rotation of the place field map as that which maximized
the correlation between sessions.

We also performed a "center-out" rotation analysis to assess coherent place field rotations
between sessions. First, the angle to the pixel with the maximum occupancy normalized event
rate was identified for each cell. Second, this angle was recalculated for the same cell in a
different session in the same box. These two angles were subtracted to get the "center-out"
rotation between sessions. Sessions which exhibited a coherent rotation displayed a peak in a
histogram of center-out angles at 0, 90, 180, or 270 degrees, while sessions which exhibited
global remapping exhibited a uniform distribution of rotation angles.

Freeze-tuned Cell Analysis

Freeze onset and offset times were first identified for each mouse/session as noted in the
Behavioral Tracking and Fear Metrics section above. We then formed calcium event rasters
using the neural activity for each cell +/- 2 seconds from freeze onset, organizing the data into a
nfreeze_onsets x ntime_bins array. We then summed this raster along the 0^{th} dimension to get
a freeze tuning curve. To calculated significance, we randomly, circularly shifted the putative
spiking activity for a cell and calculated a shuffled tuning curve in a similar manner to the actual
data. We repeated this procedure 1000 times, and calculated significance for each time bin as
the number of shuffles where the shuffled tuning exceeded the actual tuning curve divided by
1000. Last, we designated cells as significantly freeze-tuned if they had 3 or more bins with $p <$
0.01 and were active on at least 25% of freezing events.

Covariance Analysis

Putative spiking activity for each cell was first binned into 0.5 second windows and z-scored
after binning, forming a *ncells x nbins* array. The covariance of this array was then calculated
using the *cov* function in numpy. For between-session comparisons, cells active in both
sessions were matched up and a new array was formed with the base (1^{st}) session covariance
in the lower diagonal and the registered (2^{nd}) session in the upper diagonal. All entries along the
main diagonal were ignored.

<u>Author Contributions</u>

Conceptualization: N.R.K with the help of Howard Eichenbaum; Methodolgy: N.R.K; Software:
N.R.K., E.A.R; Validation: N.R.K; Formal Analysis: N.R.K, D.O.O., E.A.R; Investigation: N.R.K.,
D.O.O., E.A.R.; Data Curation: N.R.K., D.O.O., E.A.R.; Writing – original draft preparation:
N.R.K.; Writing – review and editing; N.R.K, D.O.O, E.A.R., K.D., S.R.; Visualization: N.R.K.;
Project Administration; N.R.K, S.R.; Funding Acquisition: N.R.K., K.D., S.R.

<u>Competing Interests</u>

The authors declare no competing interests.

 FIGURE LEGENDS

Figure 1: Neural discrimination between arenas predicts specificity of fear learning. A) Schematic of the behavioral paradigm. Mice freely explored two distinct arenas (neutral and shock) for 10 minutes each day. Mice underwent mild contextual fear conditioning on day 0 in the shock arena followed by immediate I.P. administration of anisomycin or vehicle in their home cage. Memory recall tests were conducted 4 hours and 1, 2, and 7 days post-shock. The time of each session is referenced to the shock session. **B)** (left) Learner (CTRL) mice freezing on all days. Red = shock arena, blue = neutral arena. *p=1.3e-0.5 shock – neutral freezing from days -2/-1 to days ½ one-sided t-test (n=4 mice). (right) Same but for Non-Learner (CTRL) mice (n=3 mice). **C)** (left) Neural overlap plots between neutral and shock arenas for an example Learner mouse on day -1, before shock. Green = cells active in the shock arena only, yellow = cells active in the neutral arena only, orange = cells active in both arenas. (right) Same for example Non-Learner on day -2 showing higher overlap of active cells between arenas. **D)** Same as B but for ANI group **E)** Mean neural discrimination between arenas (same day, days -2 to -1 and 1 to 2) vs. within arenas (same arena, days -2 and -1, days 1 and 2). *p=2.35e-8 (ρ=-0.56) Spearman correlation**. F)** Example calcium activity from the Learner mouse shown in C (left) for cells active in both arenas. Black = calcium trace, red = putative spiking activity during transient rises. Top row shows shock arena preferring cells, bottom row shows neutral arena preferring cells. **G)** Behavioral discrimination between arenas after shock (Days 1-2) shows formation of a specific fear memory for Learners only, by definition (positive = more freezing in neutral arena, negative = more freezing in shock arena, 0 = equal freezing in both arenas). *p=0.022, **p=0.00038 1-sided t-test **H)** Neural discrimination between arenas BEFORE shock indicates Learners formed more distinct representations of each arena prior to learning. Same conventions as F. *p=0.030, +p=0.059 two-sided t-test. **I)** Neural overlap between arenas correlates with specificity of fear memory on days 1-2 for Learners but not Non-Learners. *p=0.033 (ρ=0.61). **J)** Same as I) but plotting neural overlap vs. behavioral discrimination between arenas. +p=0.059 (ρ=-0.56) for Learners.

Figure 2: Anisomycin accelerates cell turnover and stifles learning-related place field remapping A) Cell overlap ratio with Day -2 session, CTRL group. Blue = within shock arena, red = shock v. neutral arena. **B)** Same as A) but for ANI group. **C)** Change in overlap ratios from a and b. *p=0.028, **p=0.00024 two-sided t-test. **D)** Number of active neurons observed each day, normalized to day -1. *p=0.039 two-sided t-test. **E)** Stable place field. *(top)* Example mouse trajectory (black) with calcium activity (red) overlaid for the same cell from day -2 to -1 in shock arena, *(bottom)* occupancy normalized rate maps for the same cells. **F)** Remapping place field. Same as E) but for a different cell that remaps from day -2 to day 1 in the shock arena. **G)** Place field correlations for all mice before shock (Days -2 and -1). *p<0.0032 Learners vs ANI and Non-Learners vs ANI, **p=1.9e-6 Non-Learners vs ANI, k-s test after Bonferroni correction. **H)** Same as G) but for days after shock. *p=0.045 Learners vs Non-Learners and p=0.0059 Non-Learners vs ANI, +p=0.1 Learners vs ANI. **I)** Same as G) but to assess learning-related remapping from before to after shock. *p=0.00021 Learners vs Non-Learners and p=3.4e-11 Non-Learners vs ANI, +p=0.12 Learners vs ANI. **J)** Place field correlations before shock broken down by mouse. **K)** Same as J) but for sessions after shock. **L)** Same as J) but from before to after shock. p=0.002 mixed ANOVA, group x arena interaction. *p=0.025 Non-Learners vs ANI, **p=0.013 Learners vs Non-Learners, ***p=0.034 Learners vs ANI, post-hoc pairwise t-test after Bonferroni correction.

Figure 3: ANI administration suppresses the development of coordinated freeze-related neural activity. **A) and B)** Example traces from two freeze-cells which exhibit coordinated activity prior to freezing event during the day 1 memory recall session in the shock arena. Red = putative spiking activity, pink = cell shown in C, blue = cell shown in E. **C)** Proportion of freeze-tuned cells detected each day across all groups. Green = freezing epochs. **D) and E)** Example Learner freeze-tuned cells identified during the 4 hour (D) or day 1 (E) memory test tracked across sessions. Peri-event calcium activity rasters are centered on freeze onset time (solid green). Dashed green = baseline calcium event probability, red solid = peri-freeze calcium event probability, bins with p<0.01 (circular permutation test) noted with red bars at top. D/E corresponds to pink/blue cells shown in A-B. Bold = session with significant freeze-tuning. **F)** Same as D and E but for ANI mouse shock cell identified during the 4 hour session. **G)** Change in peak peri-freeze calcium event probability for all freeze-tuned cells detected during

the 4 hour session. **H)** Same as G but for freeze-tuned cells detected during Day 1 recall session. p < 0.02 1-way ANOVA each day separately, *p=0.02, **p=0.001, ***p=0.0006 post-hoc Tukey test. **I)** Freeze-tuned cells exhibit increased covariance in the CTRL compared to the ANI group. Mean covariance of freeze-tuned cells from each session shown. p=0.016 two-way ANOVA (Time). *p=0.018 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **J)** Small but significant increase in covariance of all cells for CTRL mice during the 1 day recall session. p=0.0015 (Time), 0.005 (Group), 0.036 (Group x Time) two-way ANOVA. *p=0.014, **p=0.008, ***p=0.004 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **K)** Mean covariance of freeze-tuned cells detected during the 4 hour session tracked across sessions. P=0.005 (Group), 0.04 (Group x Time) two-way ANOVA. *p=0.014 post-hoc pairwise t-test (two-sided) after Bonferroni correction. **L)** Same as K but for freeze-tuned cells detected during the day 1 recall session. p=0.0003 (Group) two-way ANOVA. *p=0.016 post-hoc pairwise t-test (two-sided) after Bonferroni correction.

Figure S1: **CTRL animals exhibit behavioral variability following learning and similar rates of cell turnover between arenas vs. across days.**

A) CTRL mice freezing on all days. Red = shock arena, blue = neutral arena *p=0.025 shock - neutral freezing from days -2/-1 to days 1/2, one-sided t-test (n=7 mice). **B)** Distribution of DI$_{beh}$ scores for all mice in CTRL group on days 1 and 2. Dashed line indicates cutoff between Learners and Non-Learners. **C)** Cell overlap 1 day apart in the same arena (days -2 to -1 and 1 to 2) vs. cell overlap between arenas on the same day (days -2, -1, 1, 2) for all mice in CTRL group. *p=1.7e-5, r=0.74 Spearman correlation.

Figure S2: Non-specific effects of Anisomycin include a reduction in locomotion

A) 4 mice were given I.P injections of anisomycin only (no shock) and their locomotion was tracked over 24 hours. Normal activity did not return to baseline until between 6 and 24 hours later.

Figure S3: **Coherent Place Field Rotations Observed Between Sessions**

A) Example animal trajectories from neutral arena day -2 (top row) and day -1 (middle row) with calcium activity overlaid (red). Each column corresponds to one cell. Bottom row shows data rotated 90 degrees, demonstrating a coherent rotation of spatial activity for all neurons. **B)** Smoothed, occupancy normalized calcium event maps corresponding to data shown in A). **C)** The angle from the center of the arena to each cell's maximum intensity place field center was calculated for each session (center-out angle). The distribution of center-out angles plotted, demonstrating a coherent rotation of place fields from Day -2 to Day -1 by 90 degrees. **D)** Place field correlations (smoothed event maps) between sessions indicate apparently low stability across days without considering rotations, giving the false impression that the place field map randomly reorganizes between sessions. **E)** High correlations were observed after considering a coherent 90 degree rotation between sessions, indicating that place fields retain the same relative structure but rotate together as a whole. **F)** Mean correlations for each mouse and **G)** combined correlations for all neurons calculated without considering rotations gives the impression of instability before/after shock and heightened remapping for all groups from before to after learning.

Figure S4: Place field correlations with STM (4 hour) session

A) Distribution of place field correlations for all mice combined for sessions before (Day -2 and Day -1) vs 4 hour session. **B)** Same as A) but for 4 hour vs days 1-2 sessions. *p=0.0217 Non-Learners v ANI in shock arena, **p=0.00014 Non-Learners v ANI neutral arena, ***p=3.6e-8 Learners v Non-Learners neutral Arena. All p-values after Bonferroni correction for 3 comparisons. **C)** Before v STM mean place field correlations for each mouse/session-pair. **D)** Same as C but for STM v After.

Figure S5: Population Vector (PV) correlations indicate that anisomycin disrupts cell turnover

A)-E) 1D PV correlations between sessions including only cells active in BOTH sessions. **A)** Before (Days -2 and -1), *p=0.006 shock v neutral arena, mixed ANOVA. **B)** After (Days 1 and 2) *p=0.024 shock v neutral arena, mixed ANOVA. **C)** Before v After, p < 0.003 neutral v shock arena and Group x Arena interaction, mixed ANOVA. *p<0.001, +p=0.056 t-test (two-sided) after Bonferroni correction. **D)** Before v STM (4 hour), p=0.052 Group x Arena interaction, mixed ANOVA. **E)** STM v After *p=0.004 Neutral v Shock arena, mixed ANOVA. **F)-J)** 1D PV correlations including cells active in EITHER session (includes new and silent cells). **F)** Before *p<0.001 shock v neutral arena, mixed ANOVA. **G)** After, p > 0.12 all comparisons mixed ANOVA. **H)** Before v After, p<0.003 all comparisons mixed ANOVA *p=0.023 Learners v ANI, **p=0.007 Non-Learners v ANI, +p=0.083 Non-Learners v ANI t-test (two-sided) after Bonferroni correction. **I)** Before v STM p<0.005 Arena and Group comparisons, p=0.059 Arena x Group interaction, *p=0.004 Learners v ANI, **p=0.033 Learners v ANI t-test (two-sided) after Bonferroni correction. **J)** STM v After p<0.022 Arena and Group comparisons, p=0.103 Arena x Group interaction, *p=0.041 Learners v ANI, **p=0.003 Non-Learners v ANI, ***p=0.003 Non-Learners v ANI t-test (two-sided) after Bonferroni correction. Green = Learners, Orange = Non-Learners, Blue = ANI.

Figure S6: Freeze-cell covariance increases are driven by Learners and not purely a result of less freezing in Non-Learners and the ANI group.

A)-C) Example freeze-tuned cells tracked across sessions forward and backward in time from the day indicated in bold. Peri-event calcium activity rasters are centered on freeze onset time (solid green). Dashed green = baseline calcium event probability, red solid = peri-freeze calcium event probability, bins with p<0.01 (circular permutation test) noted with red bars at top. D/E corresponds to pink/blue cells shown in A-B. **A)** Example cell from Non-Learner **B)-C)** Example cells from two different Learners. **D)** Proportion freeze-tuned cells detected in neutral arena across days. **E)** Mean covariance of all cells in neutral arena prior to learning exhibit small changes, compare y-axis to Figure 3J. p=0.012 two-way ANOVA, *p=0.04 post-hoc pairwise two-sided t-test after Bonferroni correction. **F)** Mean covariance freeze-tuned cells after randomly downsampling the number of freeze events to match the average number observed during days -2 and -1. p=0.02 two-way ANOVA, +p= 0.055 post-hoc pairwise two-sided t-test after Bonferroni correction, mean of 100 downsampling iterations. **G)** Same as F) but for all cells p0.0015 two-way ANOVA, *p=0.02, **p=0.008, ***p=0.004 post-hoc pairwise two-sided t-test after Bonferroni correction. **H)** Mean covariance of freeze cells excluding peri-freeze times (freeze start +/- 2 sec) from neural activity. P=0.012 (Time) two-way ANOVA. **I)** Same as H) but for all cells. p=0.001 (Time) two-way ANOVA *p=0.045, **p=0.006, ***p=0.004 post-hoc pairwise t-test (two-sided). **J)** Mean covariance of freeze cells before, 4 hours after, and 1-2 days after learning/ANI injection broken down by learning group. p=0.014 (Time) two-way ANOVA. *p=0.04 post-hoc pairwise t-test (two-sided). **K)** Same as J) but for all cells. p=0.0008 (Time) two-way ANOVA. *p=0.012, **p=0.0028 post-hoc pairwise t-test (two-sided). **L)** Same as J) but tracking freeze-tuned cells from the 4 hour session forward/backward in time. p=0.006 (Time) two-way ANOVA. +p=0.13 post-hoc pairwise t-test. **M)** Same as L) but for 1 day freeze cells tracked forward/backward in time.

Figure S7: Reduced activity following anisomycin administration is not an imaging artifact and does not result from global disruption of electrical neural activity in hippocampal neurons.

A-C) Neural activity was tracked across ~5 hours before and after systemic administration of anisomycin. **A)** Cross correlograms for all single and multi-unit activity combined are shown from the pre epoch in a rest box (15 minutes), running on a novel track immediately following anisomycin injection (45 minutes), post epoch in the rest box (3.5 hours), running on a second novel track (45 minutes), and a second post epoch in the rest box (15 minutes). Clear modulation of firing at the theta timescale is observed. **B)**

Example trace from electrode in pyramidal cell layer of CA1 showing theta activity 10 minutes and 4
hours 15 minutes post injection anisomycin injection. **C)** Example sharp wave ripple events occurring from
25 minutes to 4 hours 15 minutes post anisomycin injection.

D-E) The signal-to-noise ratio of all neurons captured using calcium imaging and active between sessions
was tracked between sessions. **D)** Mean height of calcium transient peaks for all cells matched from day -
1 to 4 hour session. $p > 0.63$ both groups, two-sided t-test. **E)** Same as D) but tracking cells from day -1 to
day 1, $p > 0.68$ both groups.