<u>Title</u>: **Erasable Hippocampal Neural Signatures Predict Memory Discrimination**

<u>Authors</u>: *Kinsky, Nathaniel R.[1,3,5], Orlin, Daniel J.[3,4,5], Ruesch, Evan A.[3], Kim, Brian[1,2], Coello,
Siria[3], Diba, Kamran[1,2], *Ramirez, Steve[3]

<u>Summary</u>

Memories involving the hippocampus can take several days to consolidate, challenging efforts
to uncover the neuronal signatures underlying this process. Using calcium imaging in freely
moving mice, we tracked the hippocampal dynamics underlying memory formation across a ten-
day contextual fear conditioning (CFC) task. Following learning, context-specific place field
remapping correlated with memory performance. To causally test whether these hippocampal
dynamics support memory consolidation, we induced amnesia in a group of mice by
pharmacologically blocking protein synthesis immediately following learning. We found that
halting protein synthesis following learning paradoxically accelerated cell turnover and also
arrested learning-related remapping, paralleling the absence of remapping observed in
untreated mice that exhibited poor memory expression. Finally, coordinated neural activity that
emerged following learning was dependent on intact protein synthesis and predicted memory-
related freezing behavior. We conclude that context-specific place field remapping and the
development of coordinated ensemble activity require protein synthesis and underlie contextual
fear memory consolidation.

<u>Introduction</u>

The process of memory consolidation requires protein synthesis following learning to produce
stable long-term memories (Barondes & Cohen, 1968; Davis & Squire, 1984; Squire &
Barondes, 1974). At the synaptic level, protein synthesis is necessary for the formation of late-
phase but not early-phase long-term potentiation (LTP) in hippocampal (HPC) neurons (Frey &
Morris, 1997; Huang et al., 1994; Nguyen et al., 1994). At the behavioral level, blocking protein
synthesis following learning impairs long-term memory for hippocampus-dependent contextual
fear conditioning (CFC) while leaving short term memory intact (Ryan et al., 2015; Schafe et al.,
1999). These studies suggest that memory consolidation requires new proteins to reinforce
learning-related connections which are potentiated during learning. Despite the evidence linking
in vitro synaptic potentiation to memory consolidation, little is known about how protein
synthesis *in vivo* influences the functional coding properties of neurons before, during, and after
learning to support memory consolidation. Experience-dependent remapping of place fields in
the hippocampus, which requires NMDA receptor dependent-plasticity (Dupret et al., 2010), is
thought to reflect learning-related reorganization of the hippocampal network (Bostock et al.,

[1]Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI 48109,
USA
[2]Neuroscience Graduate Program, University of Michigan, Ann Arbor, MI 48109, USA
[3]Center for Systems Neuroscience, Boston University, Boston, MA 02451, USA
[4]Neuroscience Graduate Program, Oregon Health & Science University, Portland, OR 97239,
USA
[5]These authors contributed equally to this work
*Corresponding authors. Correspondence should be addressed to N.R.K (nkinsky@umich.edu)
or S.R. (dvsteve@bu.edu)

1991; Colgin et al., 2008). Therefore, to causally test the hypothesis that lasting, learning-
related place-cell remapping (Muller & Kubie, 1987) is necessary for consolidation of a CFC
memory (Moita et al., 2004; Wang et al., 2012) we combined *in vivo* calcium imaging in mice
with systemic administration of the protein synthesis inhibitor, anisomycin, and tracked the
evolution, remapping, and stabilization of HPC place fields during and after CFC. Finally, we
leveraged the ability to record from large cell ensembles to investigate whether blocking protein
synthesis disrupted the development of coordinated neural activity predictive of freezing
behavior. Our results indicate that context-specific remapping and coordinated, freeze-
predicting ensemble activity emerge in a protein-synthesis dependent manner following learning
to support CFC memory consolidation.

Results

Hippocampal neural dynamics between arenas predicts memory specificity

Weeks prior to training, all mice received viral infusions of the genetically encoded calcium
indicator GCaMP6f (Chen et al., 2013) in region CA1 of the dorsal hippocampus. Subsequently,
following two days of pre-exposure ("Before" learning, days -2 and -1) to an open field (neutral
arena) and an operant chamber (shock arena) mice received a mild foot-shock on day 0
(training), after which they were moved to their home cage and immediately given systemic
injections of either anisomycin (ANI group) or vehicle (CTRL group). The shock arena recording
immediately followed each neutral arena recording, separated by approximate 5 minutes to
disconnect the recording camera and move the mouse the other room/arena. We then
performed a test of short-term memory 4 hours later and three tests of long-term memory 1, 2,
and 7 days following training (Figure 1A), each time by measuring freezing behavior. We titrated
the shock level during training such that the mice froze significantly more in the shock arena
relative to the neutral arena following learning while still exploring the majority of both arenas.
Nonetheless, we observed a range of freezing levels during the day 1 and 2 memory tests
(Figure S1A) and subsequently sub-divided CTRL mice into two groups: Learners, who froze
significantly more in the shock arena, and Non-Learners, who either showed generalized
freezing or froze at low levels in both arenas (Figures 1B and Figure S1B,E). Learners exhibited
reduced freezing on day 7, indicative of extinction (Figure 1B). All subsequent analyses related
to long-term memory therefore utilized the day 1 and 2 recall sessions ("After" learning and
consolidation). In contrast, mice in the ANI group exhibited no difference in freezing between
arenas at any time point, indicating that blocking protein synthesis impaired context-specific fear
memory (Figure 1C). To assess memory specificity, we calculated a behavioral discrimination
index (DI_{beh}) which quantified how much each animal froze in the shock vs. neutral arenas.
Negative DI values indicated higher freezing in the shock arena. Learners exhibited negative
DI_{beh} levels on days 1 and 2 that were significantly different from both Non-Learners (by
definition) and from the ANI group as well (Figure 1D). Variability in freezing on the days prior to
shock could indicate heightened anxiety by some mice, which could in turn influence contextual
fear learning. However, we found no difference between groups in thigmotaxis, a metric of
anxiety (Figure S1F-G). Both the CTRL and ANI groups exhibited significant freezing in the
shock arena during the 4 hour test, though we note that the behavior in the ANI group at this
time point could arise from either contextual fear or non-specific aversive effects of anisomycin
treatment (Figure S2) since these mice froze at high levels in both arenas.

We visualized the activity of pyramidal neurons in region CA1 of the dorsal hippocampus
(Figure 1E) using a miniaturized epifluorescence microscope (Ghosh et al., 2011; Ziv et al.,

2013). We identified a large number of neurons in each 10 minute session (n = 128 to 1216,
Figure S3), extracted their corresponding calcium traces (Figure 1F), and tracked them between
sessions throughout the CFC task, which allowed us to determine the long-term evolution of the
HPC neural code. We noticed that many neurons exhibited strong changes in mean event rate
between arenas (Figure 1F) and calculated a neural discrimination index (DI_{neural}) to quantify the
distinctiveness of neural activity between arenas (0 = similar, 1 = distinct). Same-day neural
discrimination between arenas correlated strongly with across-day neural discrimination in the
same arena (Figure S1D). This indicates that mice exhibit natural variability in neural
discrimination which is invariant between different arenas and across time.

Figure 1: Mice exhibit variability in memory recall and neural activity prior to learning in a protein-synthesis dependent contextual
fear conditioning task. A) Schematic of the behavioral paradigm. Mice freely explored two distinct arenas (neutral and shock) for 10
minutes each day. Mice underwent mild contextual fear conditioning on day 0 in the shock arena followed by immediate I.P. administration of
anisomycin or vehicle in their home cage. Memory recall tests were conducted 4 hours and 1, 2, and 7 days post-shock. The time of each
session is referenced to the shock session. **B)** (left) Learner (CTRL) mice freezing on all days. Red = shock arena, blue = neutral arena.
*p=4.5e-0.4 shock – neutral freezing from day-1 to day 1 one-sided paired t-test (n=4 mice, t=13.4). (right) Same but for Non-Learner (CTRL)
mice (n=3 mice, p=0.249, t=0.819). **C)** Same as B but for ANI group (n=5 mice, p=0.219, t=0.859). **D)** Behavioral discrimination between
arenas after shock (Days 1-2) shows formation of a specific fear memory for Learners only, by definition (positive = more freezing in neutral
arena, negative = more freezing in shock arena, 0 = equal freezing in both arenas). *p= $3.78e^{-4}$(t=4.48), **p=0.0229 (t=2.17) 1-sided t-test of
mean DI value from Days 1 & 2, n=8/6/10 sessions for Learners/Non-Learners/ANI group, **E)** (left) Neural overlap plots between neutral and

Surprisingly, we noticed that Learners exhibited a trend toward higher DI_{neural} values compared to Non-Learners in the sessions prior to the shock that did not reach significance (Figure 1G). This was not due to higher activity rates for Non-Learners since we found no differences between groups in mean event rate and or distribution of event rates on the days prior to shock (Figure 1H-I). Finally, we found that freezing behavior the day of training correlated with Neutral arena freezing during the 4 Hour and Day 2 recall sessions and exhibited a trend toward correlation on Days 1 and 7 (Figure 1J and Figure S1H). These results indicate that the animal's behavioral state the day of conditioning, and potentially neural discrimination between arenas prior to learning, can influence memory specificity.

Blocking protein synthesis disrupts persistent neural activity following learning and arrests learning-related place field remapping

Next, we probed how arresting protein synthesis impacted HPC dynamics. Previous studies by us and others have found that hippocampal cells exhibit constant turnover over time such that the subset of active cells slowly changes over time (Kinsky et al., 2018; Ziv et al., 2013). We hypothesized that, by preventing plasticity, ANI administration would slow or stop the normal rate of cell turnover, measured as the overlap of active cells at each timepoint with the first day recording day (Figure 2A). Surprisingly, ANI administration temporarily accelerated cell turnover between the Day -1 and 4 hour session after which turnover rate returned to normal (Figure 2B, C). We noticed that the normalized number of active neurons also appeared to be lower for the ANI group compared to the CTRL group at the 4 hour session when anisomycin was still on board, and potentially the following sessions, which could contribute to increased turnover (Figure 2D). However, diminished cell activity could also result from reduced locomotion (Rich et al., 2014), which we observed in both groups following conditioning / anisomycin. To disentangle the effects of anisomycin and locomotion, we therefore fit the normalized number of cells observed in each recording session to a generalized linear model with arena, freezing ratio, anisomycin status (acute = 4 hr session, after = Day 1, 2, and 7), experimental group, and anisomycin status x experimental group as covariates (Methods). Under this framework, we found a highly significant influence of freezing ratio on active cell number ($p=2.3x10^{-5}$) and a trend toward fewer active cells for the ANI group at the 4 hour session (p=0.056) and following sessions (p=0.094) that did not reach significance (Figure 2D). These results indicate that reduced locomotion is the primary driver of reduced cell activity observed following anisomycin administration. To test whether reduced activity resulted from cell death, we injected a separate cohort of mice with either saline or anisomycin and then performed immunohistochemistry to stain for apoptosis 4 hours after injection. We found negligible levels of apoptosis overall with no difference between groups (Figure S4). We also observed no difference in the mean height of calcium transients (a measure of signal-to-noise ratio) for neurons active before, during, or after ANI administration, indicating that the observed decrease in the number of active neurons was not due to differences in Ca2+ levels or depletion of the GCaMP protein (Figure S5D-E). To confirm that anisomycin administration did not suppress network activities, which was reported

for intracranial infusions (but not systemic injections) of protein synthesis inhibitors in
anesthetized or immobile rodents (Barondes & Cohen, 1966; Sharma et al., 2012: Park et al.,
2023), we performed extracellular recordings in freely moving rats implanted with a linear probe
in the CA1 region of the hippocampus following administration of anisomycin. We observed
preserved theta activity, theta modulation of spiking, and robust sharp wave ripple activity for up
to 5 hours following anisomycin administration (Figure S5A-C). These observations indicate that
blocking protein synthesis following learning does not simply abolish all neuronal activity but
instead acutely accelerates the turnover of cells which may be specifically involved in learning.

Figure 2: Preventing protein synthesis accelerates cell turnover and stifles learning-related place field remapping. A) Cell
overlap ratio with Day -2 session, CTRL group. Blue = within shock arena, red = shock v. neutral arena. **B)** Same as A) but for ANI
group. **C)** Change in overlap ratios from A) and B), dots show values from both arenas for each mouse, **p=0.00174 two-sided t-test
of mean value for each mouse (t=4.11, n=7 Ctrl mice and 5 ANI mice). **D)** Number of active neurons observed each day, normalized
to day -1. p=2.e-5 freeze-ratio, #p=0.056 group x 4 hr session interaction, p=0.094 group x after interaction, generalized linear
model. **E)** and **F)** Example place fields exhibiting learning-related remapping. **E)** Place field in shock arena from Learner mouse.
(*top*) Example mouse trajectory (black) with calcium activity (red) overlaid for the same cell from day -2 to -1 in shock arena,
(*bottom*) occupancy normalized rate maps for the same cells with warm colors indicating areas of high calcium activity. **F)** Same as
E) but for Non-Learner mouse in Neutral arena. **G)** and **H)** Example stable place fields. **G)** Same as E) but for a different cell from
same mouse in the shock arena prior to conditioning. **H)** Same as F) but for a different cell from the same mouse in the neutral
arena after conditioning. **I)** Place field correlations for all mice before shock (Days -2 and -1), boxplots show population median and
1st/3rd quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with session means overlaid in dots.
Dashed line and grey shading show mean and 95% CI of correlations calculated from shuffling cell identify 1000 times between
sessions. **J)** Same as I) but for Day -1 to Day 1, *p=0.0496, **p=0.0034. **K)** Same as I) but for Day 1 to Day 2. **L)** Same as I) but for
Day 2 to Day 7. Statistics for I-L: un-paired one-sided HB test after Bonferroni correction, n=10,000 bootstraps.

We next hypothesized that arresting protein synthesis, which disrupts the permanence of newly
formed place fields (Agnihotri et al., 2004), would likewise impair the long-term stability of CFC-
related remapping (Moita et al., 2004; Wang et al., 2012). We therefore assessed place field

remapping within and across epochs by correlating event rate maps for all neurons active
between two sessions (Figure 2E-L) following administration of anisomycin, compared with
saline. We noticed that place fields recorded in the Learners group exhibited very low across-
session correlations in the neutral arena throughout the experiment (Figure S6), which could
indicate remapping. However, in a separate study, we observed what appeared to be
remapping between two recording sessions, but which was actually a coherent rotation of all
place fields around a singular point (Kinsky et al., 2018), indicative of confusion in an animal's
axis of orientation, such as between west and north (Keinath et al., 2017). Importantly, such
angular disorientation does not impair the ability of mice to identify specific places (Julian et al.,
2015), and between-cell firing relationships are maintained during coherent rotations whereas
they are scrambled following remapping. Coherent rotations match the geometry of the
environment, e.g. producing rotations in 60 degree increments in a triangle versus 90 degrees in
square environments such as our neutral and shock arenas. Therefore, to ensure we were not
mistaking coherent rotations for remapping, we rotated all place field maps from each session in
90 degree increments and used only the orientation that produced the highest correlation
between sessions.

We found generally higher correlations compared to using non-rotated maps (Figure 2I-L vs.
Figure S6F), confirming that coherent rotations occurred in many recording sessions. In
particular, place field correlations were high both before learning and after consolidation for all
groups (Figure 2I, K, L and Figure S7), though Learner correlations trended toward lower
stability after shock (Figure 2K, L) which could indicate extinction (Wang et al., 2015). We then
compared place fields between session to assess short-term (Figure S8, 4 hour session to Days
-1 and 1) and long-term (Figure 2J, Day -1 to Day 1) learning-related remapping. In agreement
with previous studies (Moita et al., 2004; Wang et al., 2012), Learner place fields remapped
from Day -1 to Day 1, as indicated by lower correlations in the shock arena compared to the
other groups (Figure 2J, Figure S7B). Interestingly, Non-Learners exhibited lower place map
correlations from Day -1 to Day 1 in the neutral arena compared to the other groups, indicating
paradoxically stable place fields in the shock arena but remapping in the neutral arena (Figure
2J, Figure S7C, Figure S8C). We observed similar results when examining population vector
correlations (Figure S9). If learning causes remapping, this double dissociation indicates that
Non-Learner memory deficits might result from improperly associating the neutral arena with
shock. In contrast, the place maps of mice in the ANI group displayed high correlations in both
arenas across all time points, including after learning (Figure 2I-L, Figure S7E-F). This deficit in
remapping following anisomycin therefore indicates that protein synthesis is required to stabilize
the set of place fields which emerge following learning to support memory consolidation.
Overall, these observations indicate that remapping is necessary for the creation of context-
specific memories and that a lack of remapping, or improperly remapping, may underlie the
memory deficits observed in Non-Learners and in the ANI group.

Coordinated freeze-predictive neural activity emerges following consolidation

In addition to the spatial code, hippocampal neural activity can also reflect non-spatial variables
in a task (McKenzie et al., 2014; Muzzio et al., 2009; Wood et al., 1999). We noticed that, in line
with recent studies (Lee & Han, 2022; Schuette et al., 2020), many hippocampal neurons
reliably produced calcium transients around the time that mice froze (Figure 3A-B, D-F and
S10A-C). We observed similar proportions of peri-freeze tuned cells across all groups and
recording sessions, even before CFC (Figure S10D-E). Examining the +/- 4 seconds around

freezing events for Days 1-2 (after learning) revealed that population-level neural activity
peaked in the 2 seconds prior to freezing for control mice compared to the ANI group (Figure
3C) but did not differ between Learners and Non-Learners (Figure S10F). This pre-freeze
increase in activity for the CTRL vs. the ANI group was not present before learning or at the 4
hour session (Figure S10L-M). After learning, neurons with significant peri-freeze tuning tiled the
entire +/- 4 seconds around freeze onset (Figure 3G). However, the average peak timing of cells
with significant peri-freeze activity shift shifted to significantly early time points for Learners vs.
Non-Learners and ANI mice after learning (Figure 3H) with ~70% of cells active before freeze
onset. We therefore refer to these neurons as freeze-predictive cells, noting that the predictive
nature of their activity is pronounced in Learners compared to Non-Learners and the ANI group.

To quantify the reliability of freeze-predictive turning within cells, we tracked the peak, peri-
freeze calcium event probability of each freeze cell backwards and forwards in time from the
Day 1 recall session, after putative memory consolidation had occurred. We found that freeze-
predictive cells exhibited much higher tuning stability spanning from the 4 hour recall to the day
1 recall sessions in Learners than did these cells in other groups (Figure 3I). This indicates that
in Learners, reliable freeze-predictive tuning began to emerge between 4 hours and 1 day
following learning and was maintained thereafter (see Figure 3E for exemplar cell). In contrast,
peri-freezing activities in mice from the ANI group and in Non-Learners were more transient and
unreliable during this time span.

Next, we investigated whether the higher reliability of freeze-predicting cells in Learners was
related to increased co-activity (see Figure 3A-B) in this neuronal subpopulation by calculating
pairwise coactivation of neurons +/- 4 seconds from freeze onset. We found that pairwise
coactivation increased significantly from 0-2 seconds prior to freezing during the 4 hour and Day
1-2 sessions for Learners but not for Non-Learners or for ANI group mice (Figure 3J-L). We
observed a similar, but much less pronounced, increase for Learners in the Neutral arena for
the after learning recall session (Figure 3M). To account for the overall increase in population-
level activity prior to freezing (Figure 3C), we calculated chance level pairwise coactivation by
shuffling the order of freeze onsets for one cell in each pair 1000 times and determined the
probability that the actual pairwise chance exceeded shuffle for each bin. We found that the
proportion of freeze-predicting cells with significant coactivation (> 3 consecutive bins with
coactivity exceeding 95% of shuffles) increased for Learners, but not Non-Learners or ANI
group mice, from before learning to the 4 hour recall session and before learning to after
learning recall sessions (Figure 3N).

We then examined whether we observed similar increases in co-activity in the overall neural
population. To quantify this, we calculated the covariance between all cell pairs. We found that
the covariance of all cell-pairs increased significantly following CFC for Learners for both the 4
hour and after recall sessions (Figure S10H). Furthermore, the observed covariance increase
was not driven by freeze-predictive activity (Figure S10J). We observed a similar increase in
covariance for the ANI group at the 4hr session and a much smaller, but significant, increase
during the after recall sessions (Figure S10H). We observed no such increase for the ANI group
when we excluded freezing times from the covariance calculation (Figure S10I). These
increases were not observed prior to shock in the neutral arena (Figure S10G), suggesting that
the heightened covariance we observed was related to CFC learning. Consistent with our
pairwise coactivity analyses, we found a trend toward increase covariance of freeze-predicting
cells during the after recall sessions for Learners, but not Non-Learners or ANI mice (Figure

3O). We observed a similar trend in freeze-predicting cell covariance when we downsampled
the number of freezing events following learning to match that on days -2 and -1 (Figure S10I).

Figure 3: Arresting protein synthesis suppresses the development of coordinated freeze-predicting neural activity. **A) and B)** Example traces from two freeze-predicting cells which exhibit coordinated activity prior to freezing event during the day 1 memory recall session in the shock arena. Red = putative spiking activity, pink = cell shown in C, blue = cell shown in E. **C)** After learning (Days 1 and 2), z-scored population level calcium activity peaks between 0 and 2 seconds prior to freezing for CTRL relative to the ANI group. Line/shading = mean +/- 95% CI. Red: bins with p < 0.05, independent t-test (one-sided, n=7 CTRL mice and 5 ANI mice). **D) and E)** Example Learner freeze-predicting cells identified during the 4 hour (D) or day 1 (E) memory test tracked across

Finally, we performed additional analyzes focused on the same set of freeze-predictive cells
conditioned on their activity during the Day 1 recall test immediately after putative memory
consolidation was finalized. Freeze-predicting cells identified on the 1 day recall session
displayed a trend toward increased covariance following CFC in Learners but not in Non-
Learners or in the ANI group (Figure S10K). However, these cells did not exhibit increased
covariance during the previous day's 4 hour session, suggesting that even though freeze-
predictive activity begins to emerge immediately following learning, functional cell connections
continue to reorganize up to one day later to form coordinated ensembles. These results
demonstrate that freeze-predictive tuning emerges following learning and continues to take
shape in the ensuing hours until it stabilizes one day later. Importantly, the coordination of these
freeze-predictive cells into neuronal ensembles requires protein synthesis, and they fail to
coactivate under memory consolidation failures.

<u>Discussion</u>

The results of our study provide evidence that HPC spatial and non-spatial representations
support contextual memory formation and consolidation. We first probed how protein-synthesis
influenced learning-related hippocampal dynamics following CFC. We saw that anisomycin
acutely accelerate cell turnover in the 4 hours following learning. In addition to this turnover
acceleration, anisomycin's amnestic effects coincided with a reduction in learning-related
remapping, effectively halting HPC contextual representations in their prior state. This paralleled
the absence of remapping in the shock arena for Non-Learners, providing strong evidence for
the persistence of remapping as a mechanism underlying spatial memory consolidation. Given
that remapping *was* observed in Learners during this time window, we speculate that arresting
protein synthesis temporarily blocked the formation of new place-fields in the subset of cells that
would otherwise undergo remapping following CFC. By this conjecture, existing synapses were
weakened to enable the formation of new connections between cells, and these new
connections were stifled by anisomycin, diminishing the excitatory drive to these cells to the
point where calcium activity was no longer observable. This idea is resonant with the notion that
more excitable/active neurons are preferentially involved in memory trace formation (Rashid et
al., 2016; Sweis et al., 2021). It also provides an explanation for why we observed accelerated
cell turnover following ANI administration as the cells involved in learning are effectively

silenced, allowing for a new set of cells to become active. Reduced activity – which may arise
from non-specific effects of ANI, or the blockage of constitutive protein translation (Scavuzzo et
al., 2019) – could also underlie increased cell turnover. However, our study does not support
this view. Consistent with a recent paper which found that ANI did not silence hippocampal
neuron activity (Park et al., 2023), we did not observe a widespread shutdown of neural activity
as we found that both theta oscillations and sharp-wave ripples were intact in ANI-treated mice.

Surprisingly, remapping occurred in untreated mice that exhibited poor memory recall (Non-
Learners); however, this remapping was limited to the neutral arena. This indicates that ANI
induced memory failures and poor learning manifest via the same underlying mechanism – a
failure to remap in the shock arena – but with important differences. ANI impairs memory by
preventing learning-related plasticity which stifles remapping, while poor learning can occur
when the shock is improperly associated with the neutral arena, causing remapping therein. In
line with previous studies, we found that fear conditioning induced place field remapping in
Learners in the shock arena (Wang et al., 2012; Moita et al., 2004). Surprisingly, however, Non-
Learners also exhibited remapping, but in the neutral arena only. If remapping is a neural
substrate for spatial learning, we speculate that this might explain the increased freezing
behavior observed for Non-Learners in the neutral arena: they are improperly associating the
neutral arena with shock. Alternatively, the observed remapping could be an indicator of low
hippocampal engagement during the task, which has been shown to decrease discrimination
between two similar arenas (Wiltgen et al., 2010). However, under either of these situations, we
would also expect to observe remapping in the shock arena since Non-Learners freeze equally
in both contexts. Future studies could shed further light on this phenomenon. Overall, our
findings support the view that the long-term stability of learning-related remapping requires
protein synthesis and underlies CFC memory consolidation.

Memory generalization or linking occurs when mice freeze at high levels in a neutral arena. Cai
et al. (2016) demonstrated that exposing mice to two arenas five hours apart compared to
seven days apart increases the likelihood of memory linking. Despite the close temporal
proximity between arena exposures in this experiment, our intent was not to increase contextual
fear memory/generalization. Instead, our protocol was designed to produce a moderate level of
freezing following conditioning, specifically in the shock arena, as too much freezing would
cause insufficient exploration to observe place fields. While close temporal exposure between
the conditioning and neutral arena can increase the likelihood of memory linking, it does not
guarantee it. Many other factors, such as the amount of pre-exposure time (Frankland et al.,
2004) and familiarity with the general experimental procedure can influence contextual memory
generalization. It should also be noted that, even procedures designed specifically to create
linked/generalized memories produce large variability in freezing levels, with many mice
exhibiting low levels of memory linking despite a 5 hour separation between arena exposures
(Shen et al., 2022). In contrast, some mice can exhibit generalized freezing in a neutral arena
which they never seen before (Wiltgen et al., 2010). The observed variability in memory
linking/generalization we observed in our control mice therefore agrees with the high levels of
variability observed in previous studies.

Next, we found that a subset of hippocampal cells reliably activated near freezing events during
fear memory recall, consistent with previous reports (Schuette et al., 2020; Mocle et al., 2024).
These cells activities were particularly heightened 1-2 seconds prior to freezing epochs in
animals that exhibited a strong, specific CFC memory but not in the ANI group or untreated

mice that exhibited poor memory. The predictive nature of these cells distinguishes them from
previously reported immobility cells which occur primarily in CA2 but also in CA1 (Kay et al.,
2016). We further found that freeze-predictive cells organized into reliable, co-active neuronal
ensembles following learning (Lee & Han, 2022; Rajasethupathy et al., 2015). Importantly, the
covariance of all cells increased incrementally for both ANI and Learner groups at the 4 hour
recall session, a time point at which protein synthesis is not considered to be required for the
induction of (early-phase) long-term potentiation *in vitro* (Frey & Morris, 1997; Nguyen et al.,
1994)*. However, at later time points, when protein synthesis *is* required to maintain (late-phase)
long-term potentiation (Frey et al., 1993; Nguyen et al., 1994), between-cell covariance
remained high for Learners only, consistent with a recent study demonstrating that the strength
of correlated activity in ventral CA1 neurons responsive to a foot shock predicted the strength of
contextual fear memory retrieval (Jimenez et al., 2020). It is important to note that all Control
mice exhibited heightened calcium activity 1-2 seconds prior to freezing epochs; however,
freeze-predictive neurons exhibited higher stability and coactivity for Learners compared to Non-
Learners and ANI mice. Therefore, in addition to maintaining learning-related synaptic changes
over hours to days, protein synthesis might also be instrumental for producing coordinated firing
of large ensembles of neurons to support memory recall.

The synchronous bursts of neuron ensembles we observed prior to freezing may provide a
neuronal basis for memory recall. For example, they could indicate the activation of sharp-wave
ripples (SWRs): transient, high frequency oscillations observable in the hippocamp local field
potential which can facilitate transmitting information throughout the cortex. SWRs are primarily
observed during periods of awake immobility and sleep and are hypothesized support memory
consolidation and planning through reactivation of neuronal activity patterns observed during
learning and active exploration (Buzsáki, 2015). However, SWRs can also occur during periods
of movement; these exploratory SWRs (eSWRs) are posited to strengthen connections between
place cells for reactivation and subsequent stabilization during sleep (O'Neill et al., 2006). In line
with this view, synchronous neuronal activation during eSWRs could strengthen connections
between freeze-predicting cells and stabilize their activity patterns to guide future memory
recall. Alternatively, coordinated activation of these cells could cooccur with other prominent
hippocampal oscillations such as theta or gamma (Colgin, 2016). Whether it occurs with SWRs,
theta, or gamma, coincident firing of freeze-predictive ensembles could be important for guiding
memory recall. In line with this idea, one study demonstrated that synchronous optogenetic
stimulation of engram neurons tagged during learning could artificially reactivate a fear memory
even when normal long-term recall of the fear memory was hindered by arresting protein
synthesis after learning, suggesting that a memory trace still resided in the network (Ryan et al.,
2015). Our findings provide a parsimonious explanation for this previous result by demonstrating
that anisomycin halts the co-activation of freeze-predictive cells, weakening the capacity of
these neurons to transmit behavior-related information to downstream regions through
coincident firing (Lisman, 1997) and impairing their ability to trigger memory recall (Ryan &
Frankland, 2022).

Overall, our results provide a bridge between the neuronal activities that underly memory
formation and the protein signaling events that are critical for plastic modification of synapses,
indicating that protein synthesis is necessary for the formation of new stable spatial
representations of an aversive context following learning and for producing coordinated activity
of freeze-predictive neurons.

Limitations of the study

In this study, we casually test how post-learning protein synthesis impacts the neural dynamics related to contextual memory consolidation and recall in the hippocampus. We find that protein synthesis is required for place cells to remap following learning and for freeze-predicting neurons to develop lasting functional connections. We utilized systemic injections of anisomycin to pharmacologically arrest protein synthesis. While anisomycin has been used for decades to block memory consolidation and disrupt long-term potentiation, it produces many off target effects including malaise, which we show in Figure S2, and which could contribute to the accelerated cell turnover and lower cell activity observed in the ANI group following administration. Moreover, it lacks temporal, regional, and cell-type specificity. Therefore, we are unable to determine whether the pharmacologic memory consolidation failure we induce is due to blocking protein synthesis or due to other, non-specific effects, including an acceleration of cell turnover or due to disruptive effects in non-hippocampal brain regions. Future studies could utilize more specific approaches (Shrestha et al., 2020) to disentangle this question. Nonetheless, this study's finding that anisomycin blocks the coactivity of freeze-predictive cell-pairs is consistent with previous literature demonstrating that protein synthesis is required to produce lasting long-term potentiation between different groups of neurons (Frey and Morris, 1997). We also utilize systemic injections rather than local infusions of anisomycin due to location of a GRIN lens immediately over the hippocampus. The neural and behavioral effects of anisomycin could therefore be due to disruption of protein synthesis in other, non-hippocampal, regions. Regardless, our results are consistent with previous behavioral studies using local hippocampal infusions to block contextual fear memory consolidation and lesion studies demonstrating that the hippocampus is necessary for contextual fear memory formation and consolidation (Kim & Fanselow, 1992; Debiec et al., 2002; Rossatto et al., 2007; Ocampo et al., 2017). Last, we utilize calcium imaging to capture neural activity, which allows for tracking of individual neurons across all stages of the experiment. However, due to the slower temporal dynamics of GCaMP, we are unable to capture fast time-scale hippocampal activity, such as SWRs, reactivation, replay, and theta sequences, that are important for memory formation and consolidation. Future studies could use electrophysiology to explore how these phenomena are likewise impacted by protein synthesis.

Acknowledgments: First and foremost, we would like to thank Howard Eichenbaum who helped conceive and design this study before his unfortunate passing in 2017. We would also like to thank Sam McKenzie for his help during early experimental design. We would like to thank Michael Hasselmo and Ian Davison for their support and feedback while performing the recordings for this study. Next, we thank Sam Levy, Dave Sullivan, and Will Mau for their assistance in all phases of calcium imaging throughout. We would like to thank Zach Pennington for valuable feedback concerning anisomycin preparation and administration, and Denise Cai and Lucas Carstensen for analysis suggestions. We would like to thank Pho Hale, Rachel Wahlberg, and Utku Kaya for feedback on the manuscript. We would like to acknowledge the GENIE Program, specifically Vivek Jayaraman, PhD, Douglas S. Kim, PhD, Loren L. Looger, PhD, Karel Svoboda, PhD from the GENIE Project, Janelia Research Campus, Howard Hughes Medical Institute, for providing the GCaMP6f virus. Finally, we would like to acknowledge Inscopix, Inc. for making single-photon calcium imaging miniscopes widely available, and specifically Lara Cardy and Vardhan Dani for all their technical support throughout the experiment. This work was supported by NIH Grants R01 MH052090, R01 MH051570, R01MH117964, NIH NRSA Fellowship 1F32NS117732-01, NIH Early Independence Award DP5 OD023106-01, an NIH Transformative R01 Award, a Young

Investigator Grant from the Brain and Behavior Research Foundation, a Ludwig Family
Foundation grant, and the McKnight Foundation Memory and Cognitive Disorders award, and
Boston University's Neurophotonics Center.

Author Contributions

Conceptualization: N.R.K with the help of Howard Eichenbaum; Methodolgy: N.R.K; Software:
N.R.K., E.A.R; Validation: N.R.K; Formal Analysis: N.R.K, D.J.O., E.A.R, B.K; Investigation:
N.R.K., D.J.O., E.A.R., B.K, S.C.; Data Curation: N.R.K., D.J.O., E.A.R., B.K.; Writing – original
draft preparation: N.R.K.; Writing – review and editing; N.R.K, D.J.O, E.A.R., B.K., K.D., S.R.;
Visualization: N.R.K.; Project Administration; N.R.K, S.R.; Funding Acquisition: N.R.K., K.D.,
S.R.

Competing Interests

The authors declare no competing interests.

Materials & Correspondence

All requests for materials and correspondence should be directed to Nat Kinsky
(nkinsky@umich.edu) or Steve Ramirez (dvsteve@bu.edu).

References

Agnihotri, N. T., Hawkins, R. D., Kandel, E. R., & Kentros, C. G. (2004). The long-term stability
of new hippocampal place fields requires new protein synthesis. *Proceedings of the
National Academy of Sciences*, *101*(10), 3656–3661.
https://doi.org/10.1073/pnas.0400385101

Barondes, S. H., & Cohen, H. D. (1966). Puromycin Effect on Successive Phases of Memory
Storage. *Science*, *151*(3710), 594–595. https://doi.org/10.1126/science.151.3710.594

Barondes, S. H., & Cohen, H. D. (1968). Memory impairment after subcutaneous injection of
acetoxycycloheximide. *Science (New York, N.Y.)*, *160*(3827), 556–557.
http://www.ncbi.nlm.nih.gov/pubmed/5689416

Bostock, E., Muller, R. U., & Kubie, J. L. (1991). Experience-dependent modifications of
hippocampal place cell firing. *Hippocampus*, *1*(2), 193–205.
https://doi.org/10.1002/hipo.450010207

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory
and planning. *Hippocampus*, *25*(10), 1073–1188. https://doi.org/10.1002/hipo.22488

Cai, D. J., Aharoni, D., Shuman, T., Shobe, J., Biane, J., Lou, J., Kim, I., Baumgaertel, K.,
Levenstain, A., Tuszynski, M., Mayford, M., & Silva, A. J. (2016). A shared neural
ensemble links distinct contextual memories encoded close in time. *Nature*, *534*, 115–118.
https://doi.org/10.1038/nature17955

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter, E. R.,
Kerr, R. a, Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K., & Kim, D. S. (2013).
Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295–
300. https://doi.org/10.1038/nature12354

Colgin, L. L. (2016). Rhythms of the hippocampal network. *Nature Reviews Neuroscience*, *17*(4), 239–249. https://doi.org/10.1038/nrn.2016.21

Colgin, L. L., Moser, E. I., & Moser, M.-B. (2008). Understanding memory through hippocampal remapping. *Trends in Neurosciences, 31*(9), 469–477. https://doi.org/10.1016/j.tins.2008.06.008

Debiec, J., LeDoux, J. E., & Nader, K. (2002). Cellular and systems reconsolidation in the hippocampus. *Neuron, 36*(3), 527–538. http://www.ncbi.nlm.nih.gov/pubmed/12408854

Davis, H. P., & Squire, L. R. (1984). Protein synthesis and memory: A review. *Psychological Bulletin*, *96*(3), 518–559. https://doi.org/10.1037/0033-2909.96.3.518

Dupret, D., O'Neill, J., Pleydell-Bouverie, B., & Csicsvari, J. (2010). The reorganization and reactivation of hippocampal maps predict spatial memory performance. *Nature Neuroscience*, *13*(8), 995–1002. https://doi.org/10.1038/nn.2599

Frey, U., Huang, Y.-Y., & Kandel, E. R. (1993). Effects of cAMP Simulate a Late Stage of LTP in Hippocampal CA1 Neurons. *Science*, *260*(5114), 1661–1664. https://doi.org/10.1126/science.8389057

Frey, U., & Morris, R. G. M. (1997). Synaptic tagging and long-term potentiation. *Nature*, *385*(6616), 533–536. https://doi.org/10.1038/385533a0

Geva, N., Deitch, D., Rubin, A., & Ziv, Y. (2023). Time and experience differentially affect distinct aspects of hippocampal representational drift. *Neuron*, *111*(15), 2357-2366.e5. https://doi.org/10.1016/j.neuron.2023.05.005

Ghosh, K. K., Burns, L. D., Cocker, E. D., Nimmerjahn, A., Ziv, Y., Gamal, A. El, & Schnitzer, M. J. (2011). Miniaturized integration of a fluorescence microscope. *Nature Methods*, *8*(10), 871–878. https://doi.org/10.1038/nmeth.1694

Huang, Y. Y., Li, X. C., & Kandel, E. R. (1994). cAMP contributes to mossy fiber LTP by initiating both a covalently mediated early phase and macromolecular synthesis-dependent late phase. *Cell*, *79*(1), 69–79. https://doi.org/10.1016/0092-8674(94)90401-4

Julian, J. B., Keinath, A. T., Muzzio, I. A., & Epstein, R. A. (2015). Place recognition and heading retrieval are mediated by dissociable cognitive systems in mice. *Proceedings of the National Academy of Sciences*, *112*(20), 6503–6508. https://doi.org/10.1073/pnas.1424194112

Kay, K., Sosa, M., Chung, J., Karlsson, M. P., Larkin, M., Grossrubatscher, I., & Frank, L. M. (2016). A hippocampal network for spatial coding during immobility. *Nature*, *531*, 185–190. https://doi.org/10.1038/nature17144

Keinath, A. T., Julian, J. B., Epstein, R. A., & Muzzio, I. A. (2017). Environmental Geometry Aligns the Hippocampal Map during Spatial Reorientation. *Current Biology*, *27*(3). https://doi.org/http://dx.doi.org/10.1016/j.cub.2016.11.046

Kim, J. J., & Fanselow, M. S. (1992). Modality-specific retrograde amnesia of fear. *Science*, *256*(5057), 675–677. http://www.ncbi.nlm.nih.gov/pubmed/1585183

Kinsky, N. R., Mau, W., Sullivan, D. W., Levy, S. J., Ruesch, E. A., & Hasselmo, M. E. (2020).
Trajectory-modulated hippocampal neurons persist throughout memory-guided navigation.
Nature Communications, 1–14. https://doi.org/10.1038/s41467-020-16226-4

Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E., & Eichenbaum, H. B. (2018).
Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales.
Current Biology, *28*(22), 1–11. https://doi.org/10.1016/J.CUB.2018.09.037

Kinsky, N. R., Vöröslakos, M., Lopez Ruiz, J. R., Watkins de Jong, L., Slager, N., McKenzie, S.,
Yoon, E., & Diba, K. (2023). Simultaneous electrophysiology and optogenetic perturbation
of the same neurons in chronically implanted animals using μLED silicon probes. *STAR*
Protocols, *4*(4). https://doi.org/10.1016/j.xpro.2023.102570

Lee, H.-S., & Han, J.-H. (2022). Activity patterns of individual neurons and ensembles correlated
with retrieval of a contextual memory in the dorsal CA1 of mouse hippocampus. *The*
Journal of Neuroscience, *43*(1), JN-RM-1407-22. https://doi.org/10.1523/jneurosci.1407-
22.2022

Levy, S. J., Kinsky, N. R., Mau, W., Sullivan, D. W., & Hasselmo, M. E. (2021). Hippocampal
spatial memory representations in mice are heterogeneously stable. *Hippocampus*, *31*(3),
244–260. https://doi.org/10.1002/hipo.23272

Lisman, J. E. (1997). Bursts as a unit of neural information: Making unreliable synapses reliable.
Trends in Neurosciences, *20*(1), 38–43. https://doi.org/10.1016/S0166-2236(96)10070-9

Mankin, E. A., Sparks, F. T., Slayyeh, B., Sutherland, R. J., Leutgeb, S., & Leutgeb, J. K.
(2012). Neuronal code for extended time in the hippocampus. *Proceedings of the National*
Academy of Sciences, *109*(47), 19462–19467. https://doi.org/10.1073/pnas.1214107109

Mau, W., Sullivan, D. W., Kinsky, N. R., Hasselmo, M. E., Howard, M. W., & Eichenbaum, H. B.
(2018). The Same Hippocampal CA1 Population Simultaneously Codes Temporal
Information over Multiple Timescales. *Current Biology*, *28*(10), 1499-1508.e4.
https://doi.org/10.1016/j.cub.2018.03.051

McKenzie, S., Frank, A. J., Kinsky, N. R., Porter, B., Rivière, P. D., & Eichenbaum, H. B. (2014).
Hippocampal representation of related and opposing memories develop within distinct,
hierarchically organized neural schemas. *Neuron*, *83*(1), 202–215.
https://doi.org/10.1016/j.neuron.2014.05.019

Mocle, A. J., Ramsaran, A. I., Jacob, A. D., Rashid, A. J., Luchetti, A., Tran, L. M., Richards, B.
A., Frankland, P. W., & Josselyn, S. A. (2024). Excitability mediates allocation of pre-
configured ensembles to a hippocampal engram supporting contextual conditioned threat
in mice. Neuron, 1–11. https://doi.org/10.1016/j.neuron.2024.02.007

Moita, M. A. P., Rosis, S., Zhou, Y., LeDoux, J. E., & Blair, H. T. (2004). Putting Fear in Its
Place: Remapping of Hippocampal Place Cells during Fear Conditioning. *Journal of*
Neuroscience, *24*(31), 7015–7023. https://doi.org/10.1523/JNEUROSCI.5492-03.2004

Muir, D. R., & Kampa, B. M. (2015). FocusStack and StimServer: a new open source MATLAB
toolchain for visual stimulation and analysis of two-photon calcium neuronal imaging data.
Frontiers in Neuroinformatics, *8*(January), 1–13. https://doi.org/10.3389/fninf.2014.00085

Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *The Journal of Neuroscience*, *7*(7), 1951–1968. http://www.ncbi.nlm.nih.gov/pubmed/3612226

Muzzio, I. A., Levita, L., Kulkarni, J., Monaco, J. D., Kentros, C. G., Stead, M., Abbott, L. F., & Kandel, E. R. (2009). Attention enhances the retrieval and stability of visuospatial and olfactory representations in the dorsal hippocampus. *PLoS Biology*, *7*(6). https://doi.org/10.1371/journal.pbio.1000140

Nguyen, P., Abel, T., & Kandel, E. R. (1994). Requirement of a critical period of transcription for induction of a late phase of LTP. *Science*, *265*(5175), 1104–1107. https://doi.org/10.1126/science.8066450

Ocampo, A. C., Squire, L. R., & Clark, R. E. (2017). Hippocampal area CA1 and remote memory in rats. *Learning & Memory*, *24*, 563–569. https://doi.org/10.1101/lm.045781.117.24

Olypher, A. V, Lánský, P., Muller, R. U., & Fenton, A. A. (2003). Quantifying location-specific information in the discharge of rat hippocampal place cells. *Journal of Neuroscience Methods*, *127*(2), 123–135. https://doi.org/10.1016/S0165-0270(03)00123-7

O'Neill, J., Senior, T., & Csicsvari, J. (2006). Place-selective firing of CA1 pyramidal cells during sharp wave/ripple network patterns in exploratory behavior. *Neuron*, *49*(1), 143–155. https://doi.org/10.1016/j.neuron.2005.10.037

Park, E. H., Kao, H. Y., Jourdi, H., van Dijk, M. T., Carrillo-Segura, S., Tunnell, K. W., Gutierrez, J., Wallace, E. J., Troy-Regier, M., Radwan, B., Lesburguères, E., Alarcon, J. M., & Fenton, A. A. (2023). Phencyclidine Disrupts Neural Coordination and Cognitive Control by Dysregulating Translation. *Biological Psychiatry Global Open Science*, *4*(1), 252–263. https://doi.org/10.1016/j.bpsgos.2023.04.009

Rashid, A. J., Yan, C., Mercaldo, V., Hsiang, H. L., Park, S., Cole, C. J., Cristofaro, A. De, Yu, J., Ramakrishnan, C., Lee, S. Y., Deisseroth, K., Frankland, P. W., & Josselyn, S. A. (2016). Competition between engrams influences fear memory formation and recall. *Science*, *353*(6297), 383–388. https://doi.org/10.1126/science.aaf0594

Rajasethupathy, P., Sankaran, S., Marshel, J. H., Kim, C. K., Ferenczi, E., Lee, S. Y., Berndt, A., Ramakrishnan, C., Jaffe, A., Lo, M., Liston, C., & Deisseroth, K. (2015). Projections from neocortex mediate top-down control of memory retrieval. *Nature*, *526*(7575), 653–659. https://doi.org/10.1038/nature15389

Resendez, S. L., Jennings, J. H., Ung, R. L., Namboodiri, V. M. K., Zhou, Z. C., Otis, J. M., Nomura, H., McHenry, J. A., Kosyk, O., & Stuber, G. D. (2016). Visualization of cortical, subcortical and deep brain neural circuit dynamics during naturalistic mammalian behavior with head-mounted microscopes and chronically implanted lenses. *Nature Protocols*, *11*(3), 566–597. https://doi.org/10.1038/nprot.2016.021

Rossato, J. I., Bevilaqua, L. R. M., Myskiw, J. C., Medina, J. H., Izquierdo, I., & Cammarota, M. (2007). On the role of hippocampal protein synthesis in the consolidation and reconsolidation of object recognition memory. *Learning & Memory*, *14*(1–2), 36–46. https://doi.org/10.1101/lm.422607

Rubin, A., Geva, N., Sheintuch, L., & Ziv, Y. (2015). Hippocampal ensemble dynamics timestamp events in long-term memory. *ELife*, *4*(December), 1–16. https://doi.org/10.7554/eLife.12247

Ryan, T. J., & Frankland, P. W. (2022). Forgetting as a form of adaptive engram cell plasticity. *Nature Reviews Neuroscience, 23*(3), 173–186. https://doi.org/10.1038/s41583-021-00548-3

Ryan, T. J., Roy, D. S., Pignatelli, M., Arons, A., & Tonegawa, S. (2015). Engram cells retain memory under retrograde amnesia. *Science*, *348*(6238), 1007–1013. https://doi.org/10.1126/science.aaa5542

Saravanan, V., Berman, G. J., & Sober, S. J. (2020). Application of the hierarchical bootstrap to multi-level data in neuroscience. *Neurons, Behavior, Data Analysis and Theory*, *3*(5), 1–25. http://www.ncbi.nlm.nih.gov/pubmed/33644783%0Ahttp://www.pubmedcentral.nih.gov/articl erender.fcgi?artid=PMC7906290

Scavuzzo, C. J., LeBlancq, M. J., Nargang, F., Lemieux, H., Hamilton, T. J., & Dickson, C. T. (2019). The amnestic agent anisomycin disrupts intrinsic membrane properties of hippocampal neurons via a loss of cellular energetics. *Journal of Neurophysiology*, *122*(3), 1123–1135. https://doi.org/10.1152/jn.00370.2019

Schafe, G. E., Nadel, N. V., Sullivan, G. M., Harris, A., & LeDoux, J. E. (1999). Memory consolidation for contextual and auditory fear conditioning is dependent on protein synthesis, PKA, and MAP kinase. *Learning and Memory*, *6*(2), 97–110. https://doi.org/10.1101/lm.6.2.97

Schuette, P. J., Reis, F. M. C. V., Maechler, P., Chakerian, M., Torossian, A., Blair, G. J., Wang, W., Blair, H. T., Fanselow, M. S., Kao, J. C., & Adhikari, A. (2020). Long-term characterization of hippocampal remapping during contextual fear acquisition and extinction. *The Journal of Neuroscience*, *September*, JN-RM-1022-20. https://doi.org/10.1523/jneurosci.1022-20.2020

Sharma, A. V., Nargang, F. E., & Dickson, C. T. (2012). Neurosilence: Profound Suppression of Neural Activity following Intracerebral Administration of the Protein Synthesis Inhibitor Anisomycin. *The Journal of Neuroscience*, *32*(7), 2377–2387. https://doi.org/10.1523/JNEUROSCI.3543-11.2012

Shrestha, P., Ayata, P., Herrero-Vidal, P., Longo, F., Gastone, A., LeDoux, J. E., Heintz, N., & Klann, E. (2020). Cell-type-specific drug-inducible protein synthesis inhibition demonstrates that memory consolidation requires rapid neuronal translation. *Nature Neuroscience*, *23*(2), 281–292. https://doi.org/10.1038/s41593-019-0568-z

Squire, L. R., & Barondes, S. H. (1974). Anisomycin, like other inhibitors of cerebral protein synthesis, impairs 'long-term' memory of a discrimination task. *Brain Research*, *66*(2), 301–308. https://doi.org/10.1016/0006-8993(74)90148-6

Sweis, B. M., Mau, W., Rabinowitz, S., & Cai, D. J. (2021). Dynamic and heterogeneous neural ensembles contribute to a memory engram. *Current Opinion in Neurobiology*, *67*, 199–206. https://doi.org/10.1016/j.conb.2020.11.017

Vöröslakos, M., Petersen, P. C., Vöröslakos, B., & Buzsáki, G. (2021). Metal microdrive and
head cap system for silicon probe recovery in freely moving rodent. *ELife*, *10*(x), 1–21.
https://doi.org/10.7554/eLife.65859

Wang, M. E., Wann, E. G., Yuan, R. K., Ramos Álvarez, M. M., Stead, S. M., & Muzzio, I. A.
(2012). Long-term stabilization of place cell remapping produced by a fearful experience.
The Journal of Neuroscience, *32*(45), 15802–15814.
https://doi.org/10.1523/JNEUROSCI.0480-12.2012

Wang, M. E., Yuan, R. K., Keinath, A. T., Ramos Alvarez, M. M., & Muzzio, I. A. (2015).
Extinction of Learned Fear Induces Hippocampal Place Cell Remapping. *Journal of*
Neuroscience, *35*(24), 9122–9136. https://doi.org/10.1523/JNEUROSCI.4477-14.2015

Wood, E. R., Dudchenko, P. A., & Eichenbaum, H. B. (1999). The global record of memory in
hippocampal neuronal activity. *Nature*, *397*(6720), 613–616. https://doi.org/10.1038/17605

Yger, P., Spampinato, G. L. B., Esposito, E., Lefebvre, B., Deny, S., Gardella, C., Stimberg, M.,
Jetter, F., Zeck, G., Picaud, S., Duebel, J., & Marre, O. (2018). A spike sorting toolbox for
up to thousands of electrodes validated with ground truth recordings in vitro and in vivo.
ELife, *7*, 1–23. https://doi.org/10.7554/eLife.34518

Ziv, Y., Burns, L. D., Cocker, E. D., Hamel, E. O., Ghosh, K. K., Kitch, L. J., El Gamal, A., &
Schnitzer, M. J. (2013). Long-term dynamics of CA1 hippocampal place codes. *Nature*
Neuroscience, *16*(3), 264–266. https://doi.org/10.1038/nn.3329

STAR★Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Other		
pAAV (AAV9).Syn.GCaMP6f.WPRE.SV40	Addgene	100837
1 µL syringe	Hamilton	65458-02
Micro Injection pump	WPI	UMP3 + Micro4 Controller
Microendoscope + Data Acquisition Box	Inscopix	V2
1028 ch Recording system	Intan	RHS2000
32 channel headstage	Intan	C3314
32 channel linear probe	NeuroNexus	A1x32-5mm-50-177
32 channel uLED probe	Neurolight	Chronic 32ch Optoelectrode
Recording cable	Intan	Standard SPI cable
Metal microdrive	3D Neuro	R2Drive
Plastic microdrive and rat protective crown	Vöröslakos et al. (2021)[3]	https://doi.org/10.5281/zenodo.8209229
Software		
Mosaic 1.2 Image Post-processing software	Inscopix	V1.2
nVista Recording software	Inscopix	nVista
OSC1lite Software	This paper	https://doi.org/10.5281/zenodo.8209373
RHX Recording Software	Intan	RHX
OpenEphys GUI	OpenEphys	GUI
MATLAB	The MathWorks	2019b
TENASPIS: Custom MATLAB code for cell identification and Ca2+ transient extraction	Mau et al. (2018)	https://github.com/SharpWave/TENASPIS (v4)
Custom MATLAB code for cell registration and data analysis	Kinsky et al. (2018)	https://github.com/nkinsky/ImageCamp
Custom Python code for Ca2+ data analysis	This paper	https://github.com/nkinsky/Eraser

• Python	Python.org	3.9.18
• Numpy	Numpy.org	1.26.3
• Matplotlib	Matplotlib.org	3.8.2
• Seaborn	Seaborn.pydata.org	0.13.2
• Pingouin	Pingouin-stats.org	0.5.4
Custom Python code for electrophysiological data analysis	Kinsky et al. (2024)	https://github.com/diba-lab/NeuroPy
• Python	Python.org	3.11.6
• Numpy	Numpy.org	1.24.4
• Matplotlib	Matplotlib.org	3.8.0
• Seaborn	Seaborn.pydata.org	0.13.0
• Pingouin	Pingouin-stats.org	0.5.3

Experimental model and study participant details

Sixteen (n = 10 controls, 6 anisomycin) male C57/BL6 mice (Jackson Laboratories), age 16 to 22 weeks during behavioral and imaging experiments and weighing 25-32g were used in this study. Three mice were excluded after performing this study: one mouse after histology revealed the GRIN lens implant and viral expression to be medial to the intended imaging, while the other two were excluded due to unstable/overexpression of GCaMP that produced aberrant calcium activity which emerged toward the end of the experiment. After exclusion of these mice, we retained 8 mice in the control (CTRL) group and 5 mice in the anisomycin group (ANI) group. Additionally, behavioral video tracking files for one CTRL mouse were corrupted during recording during all neutral field recordings from day 0 on: this mouse was excluded from all analyses which required using behavior in the neutral arena (e.g., place field correlations and any analyses where the CTRL group was split into Learners and Non-Learners). Mice were socially housed in a vivarium on a 12 hour light-dark cycle with 1-3 other mice prior to surgery and were housed singly thereafter. Mice were given free access to food and water throughout the study. All procedures were performed in compliance with the guidelines of the Boston University Animal Care and Use Committee.

One male Long Evans rat, 10 months old and weighing 480g, and one female Long Evans rat, 5 months old and weighing 240g, were used for the electrophysiological recordings in this study. Rats were socially housed in a vivarium on an adjusted 12 hour light-dark cycle (lights on at noon, off at midnight) with 1-3 other rats prior to surgery and given free access to food and water throughout the study. One week following recovery from the second surgery (see below), and prior to performing the experiments shown in Figure S5, the second rat was water restricted and performed a separate set of experiments in which she ran for water reward on a linear track during which cells were focally inhibited via delivery of light from one of 12 μLEDs on the implanted silicon probe. During water restriction, her health was monitored and she was weighed daily to ensure she maintained at least 80% of her pre-restriction weight. Following

709 completion of these recordings, the second rat was kept on water restriction while she
710 performed the experiments outlined below in the "Behavioral Paradigm" section under
711 administration of saline or anisomycin. All procedures were performed in compliance with the
712 guidelines of the University of Michigan Animal Care and Use Committee.

Method details

Viral Constructs

For mice experiments we used an AAV9.*Syn*.GcaMP6f.WPRE.SV40 virus from the University of
Pennsylvania Vector Core/Addgene with an initial titer of ~4x10^{12} GC/mL and diluted it into
sterilized potassium phosphate buffered saline (KPBS) to a final titer of ~2-4x10^{12} GC/mL for
injection.

For rat experiments, we used an pGP.AAV9.*Syn*.GcaMP7f.WPRE.SV40 virus from the
University of Pennsylvania Vector Core/Addgene with an initial titer of 2.6x10^{13} GC/mL and
diluted it into sterilized phosphate buffered saline (PBS) to a final titer of 2.6x10^{12} GC/mL for
injection. Due to poor expression no imaging was performed.

Stereotactic Surgery

We performed two stereotactic surgeries and one base-plate implant on naïve mice, aged 3-8
months, according to previously published procedures (Kinsky et al., 2018; Resendez et al.,
2016). Both surgeries were performed under 1-2% isoflurane mixed with oxygen. Mice were
given 0.05mL/kg buprenorphine (Buprenex) for analgesia (subcutaneously, SC), 5.0mL/kg of
the anti-inflammatory drug Rimadyl (Pfizer, SC), and 400mL/kg of the antibiotic Cefazolin
(Pfizer, SC) immediately after induction. They were carefully monitored to ensure they never
dropped below 80% of their pre-operative weight during convalescence and received the same
dosage of Buprenex, Cefazolin, and Rimadyl twice daily for three days following surgery. In the
first surgery, a small craniotomy was performed at AP -2.0, ML +1.5 (right) and 250nL of
GcaMP6f virus (at the titer noted below) was injected 1.5mm below the brain surface at
40nL/min using a 1µL Hamilton syringe and infusion pump. The needle remained in place a
minimum of 10 minutes after the infusion finished at which point it was slowly removed, the
mouse's scalp was sutured, and the mouse was removed from anesthesia and allowed to
recover.

3-4 weeks after viral infusion, mice underwent second surgery to attach a gradient index (GRIN)
lens (GRINtech, 1mm x 4mm). After performing an ~2mm craniotomy around the implant area,
we carefully aspirated cortex using blunted 25ga and 27ga needles under constant irrigation
with cold, sterile saline until we visually identified the medial-lateral striations of the corpus
callosum. We carefully removed these striations using a blunted 31ga needle while leaving the
underlying anterior-posterior striations intact, after which we applied gelfoam for 5-10 minutes to
stop any bleeding. We then lowered the GRIN lens to 1.1mm below bregma. Note that this
entailed pushing down ~50-300µm to counteract brain swelling during surgery. We then applied
Kwik-Sil (World Precision Instruments) to provide a seal between skull and GRIN lens and then
cemented the GRIN lens in place with Metabond (Parkell), covered it in a layer of Kwik-Cast
(World Precision Instruments), and then removed the animal from anesthesia and allowed him

to recover after removing any sharp edges remaining from dried Metabond with a dental drill
and providing any necessary sutures.

Finally, after ~2-4 weeks we performed a procedure in which the mouse was put under
light anesthesia to attach a base plate for easy future attachment of a miniature epifluorescence
microscope (Ghosh et al., 2011, Inscopix, Inc.). Importantly, no tissue was cut during this
procedure. After induction, we attached the base plate to the camera via a set screw, set the
camera's focus level at ~1/3 from the bottom of its range, and carefully lowered the camera
objective and aligned it to the GRIN lens by eye, and visualized fluorescence via nVistaHD until
we observed clear vasculature and putative cell bodies expressing GcaMP6f (Resendez et al.,
2016). To counteract downward shrinking during curing, we then raised the camera up ~50µm
before applying Flow-It ALC Flowable Composite (Pentron) between the underside of the
baseplate and the cured Metabond on the mouse's skull. After light curing, we applied opaque
Metabond over the Flow-It ALC epoxy to the sides of the baseplate to provide additional
strength and to block ambient light infiltration. Mice were allowed to recover for several days
prior to habituation to camera attachment and performance of the behavioral task outlined
below. In the event that we did not observe clear vasculature and cell bodies when we first
visualized fluorescence we covered the GRIN lens with Kwik-Cast and removed the mouse from
anesthesia without attaching the baseplate. We then waited an additional week and repeated
the steps above.

For rats, we performed two surgeries in a similar manner as described above for mice.
However, rats were administered pre-operative and post-operative Meloxicam orally for
analgesia (in lieu of Buprenex) and triple-antibiotic was applied locally (in lieu of Cefazolin
injections) to the incision at the end of surgery. Meloxicam was additionally administered for two
days post-surgery during recovery, and animals were monitored daily for a minimum of seven
days during recovery. 0.4mL of a lidocaine/bupivacaine cocktail were given under the scalp to
provide local anesthesia at the incision site. In the first surgery, 1000nL of GCaMP7f virus was
infused in the prelimbic cortex at the center of a 1mm craniotomy (AP + 2.9, ML + 3.6, from
Bregma, DV -3.0 at an 18 degree angle from top of brain). Following infusion, ~1.5 mm of
overlying cortex was removed and a 23ga needle was lowered to ~500µm above the target site.
Then, a 0.6 x 7 mm GRIN lens was lowered to 3.0mm below the top of the brain, the area
between the skull and lens was sealed with Kwik-Sil, and the lens was affixed to the skull with
Vivid-Flow light-curable composite (Pearson Dental) and Metabond (Parkell). The lens was then
covered in Kwik-Sil for protection. During this surgery, ground and reference screws were also
placed over the cerebellum and a 3d printed crown base was attached to the rat's skull
(Vöröslakos et al., 2021) to which crown walls and top were connected and to further protect the
lens and future microdrive/probe implant. The rat was screened for fluorescence 8-12 weeks
later, but no cell dynamics were observed so no imaging equipment was implanted for this rat.

16 weeks later, the rat was again given pre-operative Meloxicam and anesthetized under
isoflurane for probe implant (Kinsky et al., 2023). The crown walls were removed and a 1.0mm
craniotomy was performed at AP-4.8, ML+3.6 from bregma. After removing dura and stopping
bleeding with cold, sterile saline, a NeuroNexus A1x32-5mm-50-177 probe, attached to a metal
microdrive, was implanted at 2.3 mm below the brain surface and the metal drive base was
attached to the skull with Unifast light cured dental epoxy (Henry-Schein). The craniotomy was
sealed with Dow-Sil, the probe was protected with dental wax, and the ground and reference
wires were connected to the probe electronic interface board (EIB). The crown walls were re-

attached, the EIB was connected to the crown walls, and the rat was removed from isoflurane
and allowed to recovery. The rat was monitored daily for 7 days prior to recording, during which
the probe was lowered ~1mm until sharp wave ripples and spiking activity were visualized
indicating localization of the probe in the CA1 cell layer. Full details including videos
demonstrating the implant process are also documented in Kinsky et al., (2023).

Histology procedures

Hippocampal slices were prepared following extraction from mice in accordance with the
standard methods and guidelines of the Boston University Animal Care and Use Committee. In
brief, mice were euthanized with Euthasol (Virbac), transcardially perfused with
paraformaldehyde (PFA), and decapitated. Following extraction, brains were placed in PFA for
approximately 48 hours before undergoing sectioning. Brains that were sliced using a Cryostat
underwent an additional step of sucrose cryoprotection and subsequent freezing in -80C. Brains
were mounted to the slicing platform using Tissue Tek O. C. T. (Sakura) and kept at -30C
throughout sectioning. 50μm slices were collected across the entire aspiration site in the dorsal
hippocampus region. Brains that were sliced using a vibratome were stabilized using super glue
and submerged in 1% PBS. A Leica VT1000 S vibratome was equipped with a platinum coated
double edged blade (Electron Microscopy Sciences , Cat. #72003-01) and set to a maximal
speed of 0.9mm/s for collecting 50 μm slices. Slices prepared from both the cryostat and
microtome were directly mounted onto (type of slides go here) and cover-slipped using DAPI
following sectioning. No histology was performed in the rat study.

For cell death (apoptosis) experiments, adult male mice weighing 20.1-25g were injected with
either anisomycin (150mg/kg, IP, ~0.11mL) or saline (IP, ~0.11mL) and perfused 4hr later with
PBS followed by 4% paraformaldehyde. Brain tissue was removed and fixed for 48hr in PFA.
Fixed tissue was sliced into 50μm thick free-floating sections and stored into 0.01% azide.
Tissue was sliced coronally (50um thickness) on a vibratome and stained using Biosensis
(Ready-to-Dilute) Fluoro-Jade C (FJC) staining kit. Hippocampal sections were selected and
mounted on charged slides and left to dry overnight for proper adhesion. Following kit
instructions, slides incubated in a coplin jar with 9 parts of 80% ethanol and 1 part of sodium
hydroxide for 5 minutes. Slides were then rinsed in 70% ethanol for 2 minutes followed by 2
minutes in distilled water. To minimize background fluorescence, slides were submerged in 9
part distilled water and 1 part potassium permanganate for 5 minutes, followed by 2 x 2 minute
rinses in distilled water. Slides were then incubated in 9 parts distilled water and 1 part FJC in
the dark for 15 minutes, followed by 3 x 1 minute rinses in distilled water. Finally, tissue dried for
10-20 minutes before being submerged in xylene for 10 minutes and mounted with DPX.
Fluorescence was visualized using a confocal microscope.

Behavioral Paradigm

Prior to surgery mice were handled to habituate them subsequent camera attachment. 3-7 days
following base plate attachment surgery we conditioned mice to the imaging procedures by
further handling them for 5-10 minutes for a minimum three days. During this handling a plastic
"dummy" microscope (Inscopix) of approximately the same size/weight as the imaging camera
was attached to each mouse's head and remained on his head for 1-2 hours in his home cage.
When it became easy to attach the scope to the mouse's head a real imaging miniscope was
attached to head and an optimal focus plane chosen. We then recorded three 5 minute imaging
videos at this focus and +/- ¼ turn (~25μm) in the mouse's home cage. These movies were

processed as described in the Image Acquisition and Processing section and an optimal zoom
was chosen based on whichever focus plane maximized cell yield and produced clear looking
cell bodies. Animals were then placed in a novel environment with a different size and shape
compared to the experimental environments for a 10 minute session to habituation them to the
general experimental outline and ensure that they explored novel arenas.

Following habituation to the imaging procedures mice performed a contextual fear
conditioning (CFC) task with simultaneous imaging of hippocampus neurons over the course of
10 days. Note that all recording sessions are referred to by their time relative to applying the
mild foot-shock and the arena in which the recording occurred: e.g., Shock Day -2 occurred in
the shock arena two days prior to foot-shock while Neutral 4 hours occurred four hours after
foot-shock in the neutral arena. A typical day (Days -2, -1, 1, 2, and 7) consisted of two separate
10 minute recording blocks/sessions: one in the Neural arena and one in the shock arena. Mice
first explored a square (neutral) arena, placed in the center of a well-lit room, for 10 minutes.
The neutral arena was a square constructed of 3/8" plywood (25cm x 25cm x 15 cm), which was
painted yellow with sealable paint. Additionally, one wall was painted with black horizontal
stripes for visual orientation purposes. The neutral arena was wiped down with 70% ethanol ~10
minutes prior to recording. After 10 minutes of exploration the experimenter took the mouse out
of the arena, leaving the miniscope camera on their head and placed the mouse in its home
cage on a moveable cart upon which it was immediately transported down a short hallway to
second room. Therefore, the time between the end of the neutral and shock arena recordings
was approximately five minutes.

The second room was dimly lit and contained the fear conditioning (shock) arena. The
shock arena (Coulbourn Instruments, Whitehall, PA, USA) consisted of metal-panel side walls,
Plexiglas front and rear walls, and a stainless-steel grid floor composed of 16 grid bars (22cm x
22cm). Following 10 minutes of exploration of the shock arena, mice were removed from the
arena, the camera was removed, and mice were returned to their home cage. Both arenas were
wiped down with 70% ethanol ~10min prior to recording to eliminate any odor cues. Note that
mice always explored the neutral arena first and the shock arena second. For the Day 0
sessions, mice first explored the neutral arena for 10 minutes and were transported to the shock
arena as usual. However, during this session (Shock Day 0) the mouse was immediately given
a single 0.25mA shock and allowed to explore the arena for an additional 60 seconds only
before being removed and returned to his cage. Efficacy of shock was confirmed post-hoc by
eye by the presence of jumping/darting behavior immediately post-shock. The 4 hour session
was identical to the Day -2, 1, 1, 2, and 7 sessions. With the exception of the 4 hour session, all
recording sessions were performed in the first half of the mouse's life cycle while the 4 hour
session occurred in the second half of the light cycle.

On day zero, after the camera was removed and prior to returning to their home cage, mice
received an intraperitoneal injection of either anisomycin (150 mg/kg, Sigma-Aldrich A9789) or
the equivalent amount of vehicle. After injection, they were returned to their cage for 4 hours
until the next recording session began.

Following extensive habituation to a rest box during the seven day recovery period, rat neural
activity and behavior was recorded across ~ 5 hours. Following a 15-30 minute baseline
recording (PRE) in the rest box, the animal was given an I.P. injection of anisomycin and then
immediately placed on a novel linear track which he explored for 45 minutes (TRACK). The rat
was then placed back into the rest box for 3.5 hours (POST). Following that, the animal was

placed on a second novel track for 45 minutes (TRACK2) followed by a brief recording in the
rest box (POST2). The second rat underwent a similar procedure, except that there was a 2
hour POST recording prior to the TRACK recording and no POST2 recording. Similar
procedures were followed the day before and after anisomycin injection but using saline for
injection instead of anisomycin. Data was acquired continuously throughout with the exceptions
of periodic cable disconnections to perform the I.P. injection, start a new recording epoch, and
disconnect/reconnect cables that became twisted.

Anisomycin

For mice recordings, 25 mg of anisomycin (Sigma Aldrich) was dissolved into 50 μL of 6N HCl
and 500 μL of 1.8%NaCl. ~125 μL of 1N NaOH was then added to the solution followed by 0.1-
0.5 μL of 1N NaOH, testing pH after each addition until a final pH of 7.0 to 7.5 was reached, with
a final concentration of 24-27 mg/mL. In the case that pH rose above 7.5 during titration and
and/or the anisomycin went back into precipitate, small amounts (10-20 μL) of 6N HCl were
added until particles were no longer visible and the titration with 1N NaOH was restarted. Mice
were administered 150mg/kg of anisomycin solution via intraperitoneal injection, or ~0.15-
0.18mL for a typical 30g mouse.

For rat recordings, 100mg of anisomycin was dissolved into 1.6mL of 0.1N HCl (in 0.9% saline).
~240 μL of 1N HCl was added, then 10-12 μL of NaOH was added in 1-2 μL amounts, testing
pH between each step until a pH of 7-7.5 was reached. 0.9% Saline was then added until the
appropriate concentration was reached to inject 1.5mL of anisomycin at 150mg/kg. In one rat
(Rat1 in Figure S5), due to a small amount of waste, the final amount injected corresponded to
100 mg/kg.

Quantification and statistical analysis

Behavioral Tracking and Fear Metrics

We utilized two different camera/software configurations for tracking animal behavior. Both
configurations generated a TTL pulse at the beginning of behavioral tracking to synchronize with
image acquisition. We utilized Cineplex software (v2, Plexon) to track animal location at 30Hz in
the neutral arena. We used FreezeFrame (Actimetrics) to track animal location in the shock
arena at 3.75Hz. Animal location was obtained post-hoc via custom-written, freely available
Python software (www.github.com/wmau/FearReinstatement). We observed inconsistent frame
rates and inaccurate acquisition of behavioral video frames for one mouse in the neutral arena
during the day 0, 4 hour, and day 1-2 sessions. These sessions were excluded from analysis.

Freezing was calculated by first downsampling neutral position data to 3.75 Hz to match the
sample rate used in the shock arena. We then identified freezing epochs as any periods of 10
consecutive frames (2.67 seconds) or more where the mouse's velocity was less than
1.5cm/second.

Neural Discrimination

922 We evaluated the extent to which each animal's behavior reflected the expression of a context-
923 specific fear memory through a behavioral discrimination index (DI_{beh}), calculated as follows:

$$DI_{beh} = \frac{Frz_{Neutral} - Frz_{Shock}}{Frz_{Neutral} + Frz_{Shock}}$$

Where $Frz_{Neutral}$ and Frz_{Shock} are the percentages of time spent freezing in the neutral and shock arenas, respectively. Thus, a negative DI_{beh} value indicated more freezing behavior in the shock arena (indicating successful encoding of a context-specific fear memory), a positive DI_{beh} value indicated more freezing behavior in the neutral arena, and a DI_{beh} value around zero indicated equal/low freezing behavior in each arena (indicating the formation of a non-specific or weak fear memory).

Imaging Acquisition and Processing

Brain imaging data was obtained using nVista HD (Inscopix) at 720 x 540 pixels and a 20 Hz sample rate. Note that imaging data for one mouse was obtained at 10 Hz. Prior to neuron/calcium event identification we first pre-processed each movie using Inscopix Imaging Suite (Inscopix) software. Preprocessing entailed three steps a) motion corrections, and b) cropping the motion-corrected movie to eliminate any dead pixels or areas with no calcium activity, and c) extracting a minimum projection of the pre-processed movie for later neuron registration. We did not analyze one imaging session in which we had to reconnect the camera cable mid-session and could not synchronize the imaging data with behavioral data. Maximum projections of imaging movies were made using the Inscopix Imaging Suite or custom-written functions based off of an open-source MATLAB library (Muir & Kampa, 2015).

Electrophysiological Recordings

Data was acquired using an Intan 1028 channel recording system through OpenEphys software into binary format and behavior was tracked using Optitrack high resolution cameras and Motive image acquisition software.

Data Analysis

Data analysis was performed in both Python and MATLAB software. Python analysis code is available at https://github.com/nkinsky/Eraser.

Spike sorting and analysis

Electrophysiological recordings were automatically clustered using SpyKING CIRCUS software (Yger et al., 2018) and units were manually curated in phy (https://github.com/cortex-lab/phy/). Units were grouped into single units if they exhibited a clear refractory period and were well-isolated from other putative spikes. Other units which exhibited a clear waveform but were either poorly isolated or exhibited refractory period violations were classified as multi-unit activity (MUA). All single units and MUA were combined and cross-correlograms for the combined activity were created for each epoch of the recording separately.

Tenaspis

Neuron regions-of-interest (ROIs) and calcium events were identified using a custom written, open source algorithm employed in MATLAB 2016b called A Technique for Extracting Neuronal Activity from Single Photon Neuronal Image Sequences (Tenaspis) (Mau et al., 2018). This procedure was comprehensively documented in Kinsky et al., 2018:

"Tenaspis is open-source and available at: https://github.com/SharpWave/TENASPIS. First,
Tenaspis filters each calcium imaging movie with a band-pass filter per (Kitamura et al., 2015)
to accentuate the separation between overlapping calcium events. Specifically, Tenaspis
smooths the movie with a 4.5 µm disk filter and divides it by another movie smoothed with a
23.6 µm disk filter. Second, it adaptively thresholds each imaging frame to identify separable
pockets of calcium activity, designated as blobs, on each frame. Blobs of activity are accepted
at this stage of processing only if they approximate the size and shape of a mouse hippocampal
neuron, as measured by their radius (min = ~6µm, max = ~11µm), the ratio of long to short axes
(max = 2), and solidity (min = 0.95), a metric used by the *regionprops* function of MATLAB we
employ to exclude jagged/strange shaped blobs. Third, Tenaspis strings together blobs on
successive frames to identify potential calcium transients and their spatial activity patterns.
Fourth, Tenaspis searches for any transients that could results from staggered activity of two
neighboring neurons. It rejects any transients whose centroid travels more than 2.5µm between
frames and whose duration is less than 0.20 seconds. Fifth, Tenaspis identifies the probable
spatial origin of each transient by constructing putative regions-of-interest (ROIs), defined as all
connected pixels that are active on at least 50% of the frames in the transient. Sixth, Tenaspis
creates initial neuron ROIs by merging putative transient ROIs that are discontinuous in time but
occur in the same location. Specifically, it first attempts to merge all ROIs whose centroids are
less than a distance threshold of ~0.6µm from each other. In order to merge two transient ROIs,
the two-dimensional Spearman correlation between the ROIs must yield $r^2 > 0.2$ and $p < 0.01$.
Tenaspis then successively increases the distance threshold and again attempts to merge ROIs
until no more valid merges occur (at a distance threshold of ~3µm, typically). Seventh, Tenaspis
integrates the fluorescence value of each neuron ROI identified in the previous step across all
frames to get that neuron's calcium trace, and then identifies putative spiking epochs for each
neuron. Specifically, it first identifies the rising epochs of any transients identified in earlier
steps. Then, it attempts to identify any missed transients as regions of the calcium trace that
have a) a minimum peak amplitude > 1/3 of the transients identified in step 3, b) a high
correlation ($p < 0.00001$) between active pixels and the pixels of the average neuron ROI
identified in step 6, and b) a positive slope lasting at least 0.2 seconds. Last, Tenaspis searches
for any neuron ROIs that overlap more than 50% and whose calcium traces are similar and
merges their traces and ROIs."

Between Session Neuron Registration

We utilized custom-written, freely available MATLAB code (available at
https://github.com/nkinsky/ImageCamp) to perform neuron registration across sessions in
accordance with previous results (Kinsky et al., 2018). The details of this procedure described in
Kinsky et al. (2018) are reproduced here:

"Neuron registration occurred in two steps: session registration and neuron registration.

Session registration – Prior to mapping neurons between sessions, we determined how
much the imaging window shifted between sessions. In order to isolate consistent features of
the imaging plane for each mouse (such as vasculature or coagulated blood), we created a
minimum projection of all of the frames of the motion-corrected and cropped brain imaging
movie for each recording session. One session ("registered session") was then registered to a
base session using the "imregtform" function from the MATLAB Image Processing Toolbox,
assuming a rigid geometric transform (rotation and translation only) between images, and the
calculated transformation object was saved for future use.

Neuron Registration – Next, each ROI in the registered session was transformed to its
corresponding location in the base session. Each neuron in the base session was then mapped
to the neuron with the closest center-of-mass in the registered session, unless the closest
neuron exceeded our maximum distance threshold of 3 pixels (3.3 µm). In this case the base
session neuron was designated to map to no other neurons in the registered session. If, due to
high density of neurons in a given area, we found that multiple neurons from the base session
mapped to the same neuron in the registered session, we then calculated the spatial correlation
(Spearman) between each pair of ROIs and designated the base session ROI with the highest
correlation as mapping to the registered session ROI.

For multiple session registrations, the same procedure as above was performed for each
session in two different ways. First, we registered each session directly to the first session in the
experiment and updated ROI locations/added new ROIs to set of existing ROIs with each
registration. This helped account for slight day-to-day drift in neurons ROIs due to shifts in
vasculature, build-up of fluid underneath the viewing window, creep/shrinkage of dental cement,
etc. Second, to ensure that neuron ROIs did not drift excessively across sessions we also
performed all the above steps but did NOT update ROI locations allowing us to register each set
of ROIs to those furthest away chronologically. The resulting mappings were then compared
across all sessions, and any neuron mappings that differed between the two methods (e.g.,
ROIs that moved excessively across the duration of the experiment) were excluded from
analysis. Those that remained in the same location were included."

The procedure to assess the quality of across session registration was described in
Kinsky et al. (2018) and is reproduced here: "We checked the quality of neuron registration
between each session-pair in two ways: 1) by plotting the distribution of changes in ROI
orientations between session and comparing it to chance, calculated by shuffling neuron identity
between session 1000 times, and 2) plotting ROIs of all neurons between two sessions and
looking for systematic shifts in neuron ROIs that could lead to false negatives/positives in the
registration." All session-pairs (except those few in which we could not synchronized imaging
and behavioral data as noted above) met the above two criteria and were thus included in our
analysis.

Cells that had calcium activity in the first session (neutral arena) for which we did not identify a
matching neuron in the second session (shock arena) were classified as OFF cells. Likewise,
neurons active in the shock arena with no matching partner in the neutral arena were classified
as ON cells.

All neuron registrations were cross validated by overlaying ROIs from each session and
evaluating their match by eye. In a few cases, we noticed erroneous registrations and adjusted
our between-session neuron alignment by calculating the rigid geometric transformation using
4-5 cell ROIs active in both sessions. Figure S11 provides examples of our between-session
neuron registration results and quantification procedures.

Neural Discrimination Metrics

The extent to which gross hippocampal ensemble activity differed between arenas was
calculated in two ways. First, we calculated the proportion of cells that turned ON and OFF
between arenas divided by the total number of cells active in either arena.

Next, we calculated the extent to which each neuron active in both arenas distinguished
between arenas by changing its event rate in a manner analogous to $DI_{beh.}$ However, we took
the absolute value to account for the fact that both positive and negative event rate changes
could reflect neural differentiation between arenas. Then, we took the mean across all neurons
to obtain a neural discrimination index (DI_{Neuron}):

1054
$$DI_{Neuron} = \left| \frac{ER_{Neutral} - ER_{Shock}}{ERz_{Neutral} + ER_{Shock}} \right|$$

1055

1056 Generalized Linear Model (GLM) to Assess anisomycin effects on the number of active
1057 cells

All GLM analyses were performed using the *GLM* function in the Python *statsmodels* package.
The dependent variable was the number of neurons recorded in each session, normalized to the
amount recorded on Day -2. Covariates included were a constant/intercept term, group (ANI
vs. CTRL), arena (Shock vs. Neutral), anisomycin stage (acute = 4 hour, after = Days 1, 2, and
7), freezing ratio, and two interaction terms: group x acute and group x after. The GLM was fit
assuming a gaussian distribution with the identify link function and using the iteratively
reweighted least squares method.

1065 Placefield Analysis

Calcium transients occurring when the mouse was running greater than or equal to
1.5cm/second were spatially binned (1cm by 1cm) and occupancy normalized following which
place fields were identified and quantified in a manner similar to Kinsky et al. (2018),
reproduced here:

"Spatial mutual information (SI) was computed from the following equations, adapted from
(Olypher et al., 2003)

1072
$$I_{pos}(x_i) = \sum_{k=0}^{1} P_{k|x_i} \, log \left(\frac{P_{k|x_i}}{P_k} \right)$$

1073
$$SI = \sum_{i=1} P_{x_i} I_{pos}(x_i)$$

where:

-   P_{x_i} is the probability the mouse is in pixel x_i

-   P_k is the probability of observing k calcium events (0 or 1)

-   $P_{k|x_i}$ is the conditional probability of observing k calcium events in pixel x_i.

The SI was then calculated 1000 times using shuffled calcium event timestamps, and a neuron
was classified as a place cell if it 1) had at least 5 calcium transients during the session, and 2)
the neuron's SI exceeded 95% of the shuffled Sis…We defined the extent of a place field as all
connected occupancy bins whose smoothed event rate exceeded 50% of the peak event rate
occupancy bin."

Since spatial mutual information is biased by the number of samples (Olypher et al., 2003), we
re-sampled the behavioral tracking data to match that of the imaging data (20Hz). This required
up-sampling the shock arena data (3.75Hz->20Hz) and down sampling the neutral arena data
(30Hz->20Hz).

Placefield similarity between sessions was assessed by first smoothing the 2-d occupancy
normalized event rate maps with a gaussian kernel (2.5cm std), flattening the smoothed maps
into a vector, and then performing a Spearman correlation between all neurons active in both
sessions. To quantify chance-level place field similarity we randomly shuffled the mapping
between neurons from the first to the second session before performing the Spearman
correlation. We then repeated this procedure 100 times.

To assess the possibility that the configuration of place fields rotated together coherently
between sessions (Kinsky et al., 2018), we again performed a Spearman correlation but after
rotating the 2-d occupancy map in the second session 90 degrees. Since, due to small camera
distortions, some 2-d occupancy maps were not square, one some occasions we resized
(minimally) the second map to match the size/shape of the first map using the reshape function
in Python's *numpy* package prior to correlate the two maps. We repeated this in successive 90
degree increments and then took the mean correlation of all neurons that were active in both
sessions to determine the optimal/"best" rotation of the place field map as that which maximized
the correlation between sessions.

We also performed a "center-out" rotation analysis to assess coherent place field rotations
between sessions. First, the angle to the pixel with the maximum occupancy normalized event
rate was identified for each cell. Second, this angle was recalculated for the same cell in a
different session in the same box. These two angles were subtracted to get the "center-out"
rotation between sessions. Sessions which exhibited a coherent rotation displayed a peak in a
histogram of center-out angles at 0, 90, 180, or 270 degrees, while sessions which exhibited
global remapping exhibited a uniform distribution of rotation angles.

Freeze-predicting Cell Analysis

Freeze onset and offset times were first identified for each mouse/session as noted in the
Behavioral Tracking and Fear Metrics section above. We then formed calcium event rasters
using the neural activity for each cell +/- 4 seconds from freeze onset, organizing the data into a
nfreeze_onsets x ntime_bins array. We then summed this raster along the 0^{th} dimension to get
a freeze tuning curve. To calculated significance, we randomly, circularly shifted the putative
spiking activity for a cell and calculated a shuffled tuning curve in a similar manner to the actual
data. We repeated this procedure 1000 times, and calculated significance for each time bin as
the number of shuffles where the shuffled tuning exceeded the actual tuning curve divided by
1000. Last, we designated cells as significantly freeze-tuned if they had 3 or more bins with $p <$
0.01 and were active on at least 25% of freezing events.

Covariance Analysis

Putative spiking activity for each cell was first binned into 0.5 second windows and z-scored
after binning, forming a *ncells x nbins* array. The covariance of this array was then calculated
using the *cov* function in numpy, returning a *ncells x ncells* array. For between-session
comparisons, cells active in both sessions were registered and a new array was formed with the
base (1^{st}) session covariance in the lower diagonal and the registered (2^{nd}) session in the upper

diagonal. All entries along the main diagonal were ignored. This analysis was also performed
including only peri-freeze times (to assess peri-freeze covariance), after randomly
downsampling the number of freeze epochs to match the average from days -2 and -1 (to
control for increased sampling on days 1 and 2), and after excluding all peri-freeze times (to
assess non peri-freeze covariance).

Coactivation Analysis

1131

Putative spiking activity was first binned relative to the start of each freezing epoch ("freeze
onset"), yielding a *nfreeze_onsets x nbins* array of binarized neural activity (1 = active, 0 =
inactive). Pairwise coactivation was then calculated by taking the dot product of the binarized
neural activity arrays between each pair of cells and taking the mean along the first axis,
yielding the probability that both cells in a given cell-pair were active in each sampling bin. To
calculate population-level coactivity, the number of active cells was first calculated for each
recording session. The average number of active cells was then calculated across all peri-
freeze trials (+/- 4 seconds) and this number was z-scored relative to activity across the entire
session. To determine which cell-pairs exhibited significant pairwise coactivity beyond chance
level, we first formed a *nfreeze_onsets x nbins* array of neural activity for each cell in a pair. We
then randomly permuted the rows of the array for the second cell in the pair and calculated a
shuffled level coactivation probability for that pair as described above. We repeated this process
100 times for each cell pair, and then calculated a p-value for each bin as 1 – (# of bins where
shuffle exceeds actual coactivity) / 100. A cell-pair was designated to have significant pairwise
coactivation if it had > 3 consecutive bins with $p < 0.05$.

Hierarchical Bootstrapping

1147

We utilized hierarchical bootstrapping (HB, Saravanan et al., 2020) to estimate confidence
intervals and P-values for different metrics in line with code found at
https://github.com/soberlab/Hierarchical-Bootstrap-Paper. For each metric we generated a
population of 10,000 values by resampling with replacement at each level of the data hierarchy
(first mouse, then for each mouse, one session, then the variable measured, e.g. place field
correlations or freeze-predicting cell covariance). We then pooled all the values and calculated
the test statistic (typically the mean) and corresponding confidence interval / interquartile range.
One or two-tailed tests were used with $\alpha = 0.05$, and the p-value was calculated by using the
joint probability distribution of the bootstrapped samples to determine that the mean of one
group was greater than the mean of the other group. All hierarchical bootstrapped data was
visualized using boxplots (mean and 1st/3rd quartiles) with whiskers extending to the 95%
confidence intervals created using the boxplot function from the Python package *Seaborn*.
Chance level was calculated by shuffling cell identity before calculating each metric, and the
mean and 95% confidence intervals were visualized using the *matplotlib* package.

<u>Parametric statistics</u>

1162

1163   <u>Where applicable, all parametric statistics used are noted in the corresponding figure legend.</u>

1164

 FIGURE LEGENDS

Figure 1: Mice exhibit variability in memory recall and neural activity prior to learning in a
contextual fear conditioning task . A) Schematic of the behavioral paradigm. Mice freely explored two
distinct arenas (neutral and shock) for 10 minutes each day. Mice underwent mild contextual fear
conditioning on day 0 in the shock arena followed by immediate I.P. administration of anisomycin or
vehicle in their home cage. Memory recall tests were conducted 4 hours and 1, 2, and 7 days post-shock.
The time of each session is referenced to the shock session. **B)** (left) Learner (CTRL) mice freezing on all
days. Red = shock arena, blue = neutral arena. *p=4.5e-0.4 shock – neutral freezing from day-1 to day 1
one-sided paired t-test (n=4 mice, t=13.4). (right) Same but for Non-Learner (CTRL) mice (n=3 mice,
p=0.249, t=0.819). **C)** Same as B but for ANI group (n=5 mice, p=0.219, t=0.859). **D)** Behavioral
discrimination between arenas after shock (Days 1-2) shows formation of a specific fear memory for
Learners only, by definition (positive = more freezing in neutral arena, negative = more freezing in shock
arena, 0 = equal freezing in both arenas). *p=0.009 (t=3.56), #p=0.06 (t=1.83) 1-sided t-test of mean DI
value from Days 1 & 2, n=8/6/10 sessions for Learners/Non-Learners/ANI group. **E)** (left) Neural overlap
plots between neutral and shock arenas for an example Learner mouse on day -1, before shock. Green =
cells active in the shock arena only, yellow = cells active in the neutral arena only, orange = cells active in
both arenas. (right) Same for example Non-Learner on day -2 showing higher overlap of active cells
between arenas. **F)** Example calcium activity from the Learner mouse shown in C (left) for cells active in
both arenas. Black = calcium trace, red = putative spiking activity during transient rises. Top row shows
shock arena preferring cells, bottom row shows neutral arena preferring cells. **G)** Neural discrimination
index (DIneural) between groups on Days -2 and -1. Boxplots show population median and 1st/3rd
quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with session means
overlaid in dots, #p=0.09 after Bonferroni correction for multiple comparisons**. H)** Same as G) but for
event rate in Shock arena**, I)** Same as G) but for event rate interquartile range (IQR) in Shock arena. **J)**
Freezing in Neutral arena on Day 2 vs. Neutral arena on Day 0. Pearson correlation value and p-value
(two-sided) shown on plot. Statistics for G-J: un-paired one-sided HB test for days -2 and -1 after
Bonferroni correction, n=10,000 bootstraps.

Figure 2: Preventing protein synthesis accelerates cell turnover and stifles learning-related place
field remapping. A) Cell overlap ratio with Day -2 session, CTRL group. Blue = within shock arena, red =
shock v. neutral arena. **B)** Same as A) but for ANI group. **C)** Change in overlap ratios from A) and B), dots
show values from both arenas for each mouse, **p=0.00174 two-sided t-test of mean value for each
mouse (t=4.11, n=7 CTRL mice and 5 ANI mice). **D)** Number of active neurons observed each day,
normalized to day -1. p=2.e-5 freeze-ratio, #p=0.056 group x 4 hr session interaction, p=0.094 group x
after interaction, generalized linear model. **E)** and **F)** Example place fields exhibiting learning-related
remapping. **E)** Place field in shock arena from Learner mouse. (*top)* Example mouse trajectory (black)
with calcium activity (red) overlaid for the same cell from day -2 to -1 in shock arena, *(bottom)* occupancy
normalized rate maps for the same cells with warm colors indicating areas of high calcium activity. **F)**
Same as E) but for Non-Learner mouse in Neutral arena. **G)** and **H)** Example stable place fields. **G)** Same
as E) but for a different cell from same mouse in the shock arena prior to conditioning. **H)** Same as F) but
for a different cell from the same mouse in the neutral arena after conditioning. **I)** Place field correlations
for all mice before shock (Days -2 and -1), boxplots show population median and 1st/3rd quartiles
(whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with session means overlaid in
dots. Dashed line and grey shading show mean and 95% CI of correlations calculated from shuffling cell
identify 1000 times between sessions. **J)** Same as I) but for Day -1 to Day 1, *p=0.0496, **p=0.0034. **K)**
Same as I) but for Day 1 to Day 2. **L)** Same as I) but for Day 2 to Day 7. Statistics for I-L: un-paired one-
sided HB test after Bonferroni correction, n=10,000 bootstraps.

Figure 3: Arresting protein synthesis suppresses the development of coordinated freeze-
predicting neural activity. **A) and B)** Example traces from two freeze-predicting cells which exhibit
coordinated activity prior to freezing event during the day 1 memory recall session in the shock arena.
Red = putative spiking activity, pink = cell shown in C, blue = cell shown in E. **C)** After learning (Days 1
and 2), z-scored population level calcium activity peaks between 0 and 2 seconds prior to freezing for
CTRL relative to the ANI group. Line/shading = mean +/- 95% CI. Red: bins with p < 0.05, independent t-
test (one-sided, n=7 CTRL mice and 5 ANI mice). **D) and E)** Example Learner freeze-predicting cells

identified during the 4 hour (D) or day 1 (E) memory test tracked across sessions. Peri-event calcium
activity rasters are centered on freeze onset time (solid green). Dashed green = baseline calcium event
probability, red solid = peri-freeze calcium event probability, bins with p<0.01 (circular permutation test,
n=1000) noted with red bars at top. D/E corresponds to pink/blue cells shown in A-B. Bold = session with
significant freeze-tuning. **F)** Same as D and E but for ANI mouse freeze-predicting cell identified during
the 4 hour session. **G)** Peri-freeze calcium event probability for all freeze-predicting cells detected for
each group after learning (Days 1-2), sorted by time of peak activation. **H)** (left) Cumulative distribution of
peak peri-freeze activation times before learning. *p=0.49, **p=1e-5 two-sided Wilcoxon rank-sum test , c:
p=0.005 Non-Learners v. ANI, 1-sided Mann-Whitney U-test (n = 329/458/543 neurons for Learners/Non-
Learners/ANI group) (right) same as left but for after learning. *p < 0.022, **p=2e-7 two-sided Wilcoxon
rank-sum test. a: p=0.022 Learners v. Non-Learners, b: p=0.029 Learners v. ANI 1-sided Mann-Whitney
U-test (n=194/315/366 neurons for Learners/Non-Learners/ANI group). **I)** Change in peak peri-freeze
calcium event probability for all freeze-predicting cells detected during the Day 1 session and either the 4
hour or Day 2 session. p < 0.02 1-way ANOVA each day separately, *p=0.02, **p=0.001, ***p=0.0006
post-hoc Mann-Whiney U-test (n=30/35/29 4h to Day 1 cells and n = 35/37/45 Day 1 to Day 2 cells for
Learners/Non-Learners/ANI group. **J)** Pairwise coactivation probability of all freeze-predicting cells for
Learners during Before, 4 hour, and After sessions in Shock arena. Maroon/Black bars at top indicate
significant increases in coactivation at 4 hour / After time points compared to before, p< 0.05 1-sided
Mann-Whitney U-test (n= 4).   **K)** Same as J) but for Non-Learners (n=3). **L)** Same as K) but for ANI group
(n= 5). **M)** Same as K) but for Learners in Neutral Arena (n=4). **N)** Proportion of freeze-predictive cells
with significant pairwise coactivation compared to chance (trial shuffle). *p=4e-8, **p=3.6e-7, #p=0.093.
boxplots show population median and 1st/3rd quartiles (whiskers, 95% CI) estimated using hierarchical
bootstrapping (HB) data with session means overlaid in dots. **O)** Freeze-predicting cells exhibit a trend
toward increased peri-freeze covariance (z-scored relative to the Day -2 and -1 covariance values for all
cells) for Learners but not Non-Learners or ANI group mice. Mean covariance of freeze-predictive cells
from each session shown. #p=0.06. Statistics for K and O: un-paired one-sided HB test after Bonferroni
correction, n=10,000 shuffles.

Figure S1: **CTRL animals exhibit behavioral variability following learning and similar rates of cell**
turnover between arenas vs. across days.

A) CTRL mice freezing on all days. Red = shock arena, blue = neutral arena. **B)** Distribution of DI$_{beh}$
scores for all mice in CTRL group on days 1 and 2. Dashed line indicates cutoff between Learners and
Non-Learners. **C)** Cell overlap 1 day apart in the same arena (days -2 to -1 and 1 to 2) vs. cell overlap
between arenas on the same day (days -2, -1, 1, 2) for all mice. *p=4.05e-7, ρ=0.81 Spearman
correlation, n=26 overlap values across 13 mice. **D)** Same as C) but for neural discrimination index.
*p=6.29e-7, ρ=0.82 Spearman correlation. **E)** Freeze ratio plots with each mouse's value connected by a
line. **F)** No difference in thigmotaxis prior to conditioning in Neutral arena, dots show mean thigmotaxis
ratio for each mouse from Days -2 and -1, p=0.23 ANOVA. **G)** Same as F) but in Shock arena, p=0.95
ANOVA. **H)** Neutral arena freezing ratio for each session after shock plotted versus Neutral arena
freezing on day of training (day 0) with Pearson correlation and associated p-value (two-sided) shown, n
= 7 mice. **I)** Same as H) but for Shock arena freezing versus Neutral arena freezing on day of training.

Figure S2: Non-specific effects of Anisomycin include a reduction in locomotion

A) 4 mice were given I.P injections of anisomycin only (no shock) and their locomotion was tracked over
24 hours. Normal activity did not return to baseline until between 6 and 24 hours later. Black solid/dashed
lines = 4 hour mean +/- std freezing ratio for non-ANI fear conditioned mice shock arena (see B). **B)**
Freezing ratios for all mice in the Shock arena prior to conditioning and 4 hours after conditioning shown
for reference. **C)** Freezing ratios in Neutral arena immediately before and one day after conditioning.
Lines show mean +/- std.

Figure S3: Absolute cell numbers recorded across all sessions.

A) Total number of neurons recorded across all sessions in Control group. **B)** Same as A but for ANI group. **C)** Same as A but for Learners. **D)** Same as A but for Non-Learners. **E)** Histogram of cell counts across all sessions with each group mean shown with dashed lines.

Figure S4: Anisomycin does not cause neuronal cell death at 4hr post injection in the hippocampus.

A) Experiment schematic. Animals were given I.P. injections of either ANI (n=8) or saline (n=7). 4 hours later, brains were extracted and processed for Fluoro-Jade C. **B)** Fluoro-Jade C positive neuronal count normalized to area; **C-H)** Representative images of whole hippocampus (HPC), CA1, and dentate gyrus (DG) in **C-E)** anisomycin-injected mice or **F-H)** saline- injected mice. Scale bar equals 200µm.

Figure S5: Reduced activity following anisomycin administration is not an imaging artifact and does not result from global disruption of electrical neural activity in hippocampal neurons.

A-C) Neural activity was tracked across ~5 hours before and after systemic administration of anisomycin in a rat. **A)** Cross correlograms for all single and multi-unit activity combined are shown from the pre epoch in a rest box (15 minutes), running on a novel track immediately following anisomycin injection (45 minutes), post epoch in the rest box (3.5 hours), running on a second novel track (45 minutes), and a second post epoch in the rest box (15 minutes). Clear modulation of firing at the theta timescale is observed. **B)** Example trace from electrode in pyramidal cell layer of CA1 showing theta activity 10 minutes and 4 hours 15 minutes post injection anisomycin injection. **C)** Example sharp wave ripple events occurring from 25 minutes to 4 hours 15 minutes post anisomycin injection across 9 channels of a linear probe spanning above to below the pyramidal cell layer. **D-F)** Same as A-C but for a different rat following systemic anisomycin injection. F) Shows one trace from an electrode in the cell layer with a raster of spiking activity from all units recorded shown below the trace, demonstrating a population burst coincident with each sharp wave ripple. **G-I)** Same as D-F but the following day after systemic saline (control) injection demonstrating no lasting effects of anisomycin 24 hours after injection. **J-K)** The signal-to-noise ratio of all mouse neurons captured using calcium imaging and active between sessions was tracked between sessions. **J)** Mean height of calcium transient peaks for all cells matched from day -1 to 4 hour session. p>0.63 both groups, two-sided t-test (n=8 CTRL and 7 ANI, 1 additional CTRL mouse included whose video tracking behavioral data was corrupted and could therefore could not be classified as a Learner or Non-Learner). **K)** Same as J) but tracking cells from day -1 to day 1, p>0.68 both groups, two-sided t-test.

Figure S6: **Coherent Place Field Rotations Observed Between Sessions**

A) Example animal trajectories from neutral arena day -2 (top row) and day -1 (middle row) with calcium activity overlaid (red). Each column corresponds to one cell. Bottom row shows data rotated 90 degrees, demonstrating a coherent rotation of spatial activity for all neurons. **B)** Smoothed, occupancy normalized calcium event maps corresponding to data shown in A). **C)** The angle from the center of the arena to each cell's maximum intensity place field center was calculated for each session (center-out angle). The distribution of center-out angles plotted, demonstrating a coherent rotation of place fields from Day -2 to Day -1 by 90 degrees. **D)** Place field correlations (smoothed event maps) between sessions indicate apparently low stability across days without considering rotations, giving the false impression that the place field map randomly reorganizes between sessions. **E)** High correlations were observed after considering a coherent 90 degree rotation between sessions, indicating that place fields retain the same relative structure but rotate together as a whole. **F)** Mean correlations for each mouse without considering rotations gives the impression of instability before/after shock and heightened remapping for all groups from before to after learning. Boxplots show population median and $1^{st}/3^{rd}$ quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with session means overlaid in dots. Dashed line and grey shading show mean and 95% CI of correlations calculated from shuffling cell identify 1000 times between sessions.

Figure S7: Example stable and remapping place fields across sessions

A) Example place field plots for Learner mouse in Neutral arena Before conditioning (Day -2 to Day -1),
from Before to After conditioning (Day -1 to Day 1), and After consolidation (Day 1 to Day 2)
demonstrating high stability in Neutral arena across all time points. **(top row)** Mouse trajectory in black
with cell calcium activity overlaid in red. **(bottom row)** Smoothed, occupancy normalized calcium event
spatial maps corresponding to raw trajectory and event data shown in top row, with warmer colors
indicating high event rates and cool colors indicating low event rates. Spearman correlation value
between event rate maps shown at top. Dashed lines denote separate different cells at each time point,
and solid lines separate different comparison times. **B)** Same as A) but for different Learner in Shock
arena demonstrating remapping. Red box shows two example remapping cells in the Shock arena. **C)**
and **D)** Same and A) and B) but for two different Non-Learners with red box showing remapping cells in
the Neutral Arena. **E)** and **F)** Same as A) and B) but for two different ANI group mice showing stable
place fields between sessions at all time points.

Figure S8: Place field correlations with STM (4 hour) session

A) Place field correlations for all mice combined for Day -1 vs 4 hour session. Boxplots show population
median and $1^{st}/3^{rd}$ quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with
session means overlaid in dots. Dashed line and grey shading show mean and 95% CI of correlations
calculated from shuffling cell identity 1000 times between sessions. **B)** Same as A) but for 4 hour vs Day
1 session. All hierarchical bootstrap test comparisons are ns. **C)** Place field correlations for all mice in
each group from before to after shock (Day -1 to Day 1). Boxplots show population median and $1^{st}/3^{rd}$
quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with session means
overlaid in dots. Significant p-values calculated using a one-sided HB permutation test are shown directly
on each panel.

Figure S9: Population Vector (PV) correlations.

A)-E) 1D PV correlations between sessions including only cells active in BOTH sessions. **A)** Before (Day
-2 vs -1), *p=0.042, ***p<0.0006. **B)** After (Day 1 vs 2) *p=0.034, #p=0.072. **C)** Before v After (Day -1 vs
1), *p<0.039, **p=0.0098. **D)** Day -1 vs 4 hr session. **E)** 4 hr session vs Day 1 *p=0.0049, #p=0.076. **F)**
Day 2 vs Day 7, *p=0.0189, **p=0.0164, #p=0.064. Green = Learners, Orange = Non-Learners, Blue =
ANI. Boxplots show population median and $1^{st}/3^{rd}$ quartiles (whiskers, 95% CI) estimated using
hierarchical bootstrapping (HB) data with session means overlaid in dots. Dashed line and grey shading
show mean and 95% CI of correlations calculated from shuffling cell identify 1000 times between
sessions. Statistics: un-paired one-sided HB test after Bonferroni correction.

Figure S10: Freeze-cell covariance increases are driven by Learners and not purely a result of less
freezing in Non-Learners and the ANI group.

A)-C) Example freeze-predicting cells tracked across sessions forward and backward in time from the day
indicated in bold. Peri-event calcium activity rasters are centered on freeze onset time (solid green).
Dashed green = baseline calcium event probability, red solid = peri-freeze calcium event probability, bins
with p<0.01 (circular permutation test) noted with red bars at top. D/E corresponds to pink/blue cells
shown in A-B. **A)** Example freeze-predictive cell from Non-Learner **B)-C)** Example freeze-predicting cells
from two different Learners. **D)** Proportion freeze-predicting cells detected in Neutral arena across days.
Bars=mean, line=std. **E)** Same as D) but for Shock arena. **F)** After learning (Days 1 and 2), z-scored
population level calcium activity peaks between 0 and 2 seconds prior to freezing for both Learners and
Non-Learners. Line/shading = mean +/- 95% CI. Red: bins with p < 0.05, independent t-test (one-sided,
n=4 Learners and 3 Non-Learners). **G)** Mean covariance of all cells in neutral arena prior to learning
exhibit small changes, compare y-axis to Figure 3O and S10H, H.**p=0.0048, ***p<1e-8, #p=0.06. **H)**
Same as G) but for all cells in shock arena. *p=0.026, **p=0.00052, ***p<2.5e-6. **I)** Same as G) but for
freeze predictive cells only, peri-freeze times only, and after randomly downsampling the number of
freeze events to match the average number observed during days -2 and -1. #p=0.056 **J)** Same as H) but
excluding all peri-freeze times. *p=0.0018**p=1.2e-4, ***p=2.1e-5. **K)** Mean covariance of freeze-
predicting cells detected on Day 1 tracked across time. #p=0.068. For G-K, boxplots show population
median and $1^{st}/3^{rd}$ quartiles (whiskers, 95% CI) estimated using hierarchical bootstrapping (HB) data with

session means overlaid in dots. Dashed line and grey shading show mean and 95% CI of correlations calculated from shuffling cell identify 1000 times between sessions. Statistics: un-paired one-sided HB test after Bonferroni correction. **L-M)** There are no significant differences between the proportion of active units (z-scored) peaks for Control animals compared to the ANI group between 0 and 2 seconds prior to freezing before conditioning (L) and at the 4 hour session (M). Line/shading = mean +/- 95% CI. Red: bins with $p < 0.05$, independent t-test (two-sided).

Figure S11: Between Session Neuron Registration Metrics

A) Example neuron registration between the Day -1 and 4 Hour session for one mouse in the shock arena depicting cell ROIs active during the Day -1 session only (blue), the 4 Hour session only (teal), and both the Day -1 and 4 Hour session (yellow) with ROIs which were successfully registered across sessions outlined in red. Insets show magnified ROIs and demonstrate that cells registered across days exhibit similar shape and orientation. **B)** and **C)** Minimum projections of the imaging movie from the sessions shown in A demonstrating high day-to-day stability, evidenced by clear alignment of landmarks (e.g. vasculature) between sessions. **D)** Change in ROI orientation for all neurons registered to the Day -2 session for the mouse shown in A). Note that the majority of registered neurons exhibit very small changes in orientation between sessions, even up to 9 days later (Day -2 to Day 7). **E)** and **F)** Similar to D) but for all mice in the CTRL and ANI groups separately for Day -1 to Day 4 (from before to after anisomycin administration). Note a similar distribution of ROI orientation changes, indicating that the observed acceleration of cell turnover following anisomycin administration is not due to poor neuron registration. **G)** Side-by-side comparison of all neuron ROI changes for each group shown as a cumulative density function.