Maternal hormones engage a dormant monomorphic aggression circuit, leading to the introduction of an innate behavior in adulthood in females

Stefanos Stagkourakis[1],[¶], Paul Williams[2], Giada Spigolon[1, ¶], Shreya Khanal[1], Katharina Ziegler[1], Laura Heikkinen[1,2], Gilberto Fisone[1] and Christian Broberger[1,2],**

[1]Department of Neuroscience, Biomedicum B:4, Karolinska Institutet, Solnavägen 9, 171 65 Stockholm, Sweden and

[2]Department of Biochemistry and Biophysics, Stockholm University, Svante Arrhenius väg 16C, 104 05 Stockholm, Sweden

[¶]Current address: Division of Biology and Biological Engineering 156-29, Tianqiao and Chrissy Chen Institute for Neuroscience, California Institute of Technology, Pasadena, CA 91125, USA

*Correspondence: stefanos.stagkourakis@caltech.edu, christian.broberger@dbb.su.se

Acknowledgments

Members of the Broberger laboratory are thanked for discussion during the preparation of this manuscript. This study was made possible by a European Research Council starting grant (ENDOSWITCH 261286), the Swedish Research Council (2018-02480), the Strategic Research Program in Diabetes at Karolinska Institutet, *Hjärnfonden* (the Swedish Brain Foundation), Novo Nordisk Fonden, Wenner-Gren Foundation funding, and internal funds from Karolinska Institutet.

Author contributions

S.S. and C.B. conceived the study and wrote the manuscript. S.S. designed, performed, and analyzed experiments. P.W., G.S., G.F., and C.B. designed experiments. P.W., G.S., S.K., K.Z., and L.H. performed experiments. All authors reviewed the manuscript.

Competing interests

The authors declare no competing financial interests.

In Brief

Maternal hormones activate a subset of hypothalamic "aggression neurons", which are required for the transient expression of maternal aggression during lactation.

Highlights

- Quiescent PMv^{DAT} neurons in female mice become hyperexcitable in lactating dams.
- Manipulation of PMv^{DAT} cell activity bidirectionally regulates maternal aggression.
- The maternal hormones prolactin and oxytocin excite PMv^{DAT} cells.
- Activation of PMv^{DAT} neurons impairs the expression of non-aggression-related behavioral programs.

Summary

Aggression - a sexually dimorphic behavior, is prevalent in males and typically absent in virgin females. Following parturition, however, the transient expression of aggression in adult females ensures the pups' defense from predators and infanticide. While maternal hormones are known to elicit nursing, whether they play a role in the expression of maternal aggression remains unknown. Here we show that a molecularly defined subset of ventral premammillary (PMv[DAT]) neurons, instrumental for intermale aggression, switch into a hyperexcitable state during lactation. We identify that the maternal hormones prolactin and oxytocin excite these cells, an effect mediated through T-type Ca^{2+} channels. Optogenetic manipulation or genetic ablation of PMv[DAT] neurons profoundly affects maternal aggression, while activation of these neurons impairs the expression of non-aggression-related behaviors. This work identifies a monomorphic neural substrate that incorporates hormonal cues to enable the transient expression of a dormant behavioral program in adult females.

Introduction

Aggression is a near-ubiquitous adaptive behavior in animals and man that serves multiple purposes and takes several forms, depending on agent and context (*1-8*). Intermale aggression is the most prevalent form of aggression found in nature, serving to establish intermale hierarchy, and consequently, access to resources and mating opportunities. In contrast, aggressive encounters in female mammals are mostly absent, with some notable exceptions (*9-11*). Female interactions with conspecifics, however, change dramatically after pregnancy and parturition, when lactating dams typically exhibit profound aggressive behavior towards both conspecifics of both sexes (*12*). Indeed, maternal aggression is as indelible of a component as other maternal behaviors, such as nursing and nest-building. But this dramatic – yet reversible – phenotypical metamorphosis also presents a neurobiological enigma, specifically; How does the brain reconfigure to transiently enable the expression of a behavior that is otherwise outside of the animal's repertoire? The brain loci and mechanisms underlying maternal aggression are largely unknown (*13*), and the signals that drive this transition, alongside their site of action, remain ambiguous (*14*). Importantly, it remains unknown whether the neural modules that lead to the expression of aggression are shared between the two sexes, or exhibit sexual dimorphism. Finally, a question that stands out is whether brain nodes that elicit certain maternal behaviors, facilitate or impair the expression of other maternal behaviors. Addressing this question would suggest a principle for the neural encoding of maternal behaviors, in *general* vs. *specialized* behavioral repertoire-eliciting nuclei.

Here we shed light on the above questions by focusing on the ventral premammillary nucleus (PMv), a node extending within the caudal two-thirds of the classically delineated "hypothalamic attack area" (*15*), and which has been recently implicated in social behavior (*16, 17*) and inter-male aggression (*18-21*). This nucleus receives dense inputs from areas associated with sensory processing (*22*), and projects (*23*) to nuclei associated with social behavior (*24-27*), aggression (*19, 28-30*), and reward (*20, 31-34*). Within the PMv, a subset of glutamatergic cells that express the dopamine transporter (PMv^{DAT} neurons) has been specifically implicated in male agonistic behaviors (*18, 20, 21*).

Here, we identify PMv^{DAT} cells as quiescent in the virgin female mouse brain, with a dramatic switch in their excitability during motherhood – excited by the hormones that surge during nursing and which facilitate the expression of maternal care. Specifically, prolactin leads to excitatory pre- and post-synaptic changes, an effect also mimicked by another maternal hormone – oxytocin. Optogenetic control of these neurons provides a bidirectional handle on maternal aggression, while partial genetic ablation hinders the expression of the behavior. Finally activation of PMv^{DAT} neurons during a maternal care paradigm, impairs the expression of pups retrieval and nursing, suggestive of a specialized role of PMv cells at eliciting a dedicated behavioral expression.

The present study suggests how the sex-specific expression of an adaptive behavior can result from state-dependent tuning of neuronal excitability.

Results

Low *vs* high baseline activity of PMv^{DAT} cells in virgin *vs* postpartum females

While maternal aggression is widespread in the animal kingdom, recent studies suggest a high degree of inter-individual variability in the expression of aggressive behavior ([20, 34, 35]). Previous work has also shown that a number of variables impact on the expression of maternal aggression, including litter size ([36]), pup presence ([37]), the intruder's sex ([38]) and behavior ([39]). To control for these factors, we used a three consecutive day (p3-p5) resident-intruder (RI) paradigm to screen lactating dams for aggressive behavior, using C57 or BALB/c juvenile female virgin mice (p40-50) as intruders and restricting litter size to four to six pups *post partum* (Figure 1A). Under this paradigm, 42% of multiparous lactating dams expressed maternal aggression (Figure 1A), with increased duration and decreased latency upon repeated experience in aggressive encounters (Figure 1B), similar to what is observed in intermale aggression using the same paradigm ([20, 34, 40]).

A subpopulation of male laboratory rodents have consistently been shown to be non-aggressive in the RI paradigm ([20, 34, 40]). Strikingly, PMv^{DAT} neurons in male non-aggressor mice are significantly less electrically excitable than in aggressors ([41]). This observation prompted the question if a similar phenotype-correlated electrical profile could be identified in lactating (aggressive) mice, *vs*. virgin female mice ([42, 43]) who are non-aggressive. Brain slice whole-cell patch-clamp recordings were performed on PMv^{DAT} cells, which were filled with 0.2% Neurobiotin for post-recording identification (Figure 1E, F). Notably, PMv^{DAT} cells from lactating dams were more depolarized compared to cells from virgin females (Figure 1G), exhibiting significantly higher firing frequency (Figure 1H, Table S1) and a more depolarized resting membrane potential (Figure 1I, Table S1).

PMv neurons have been reported to activate in conjunction with an aggressive encounter in lactating rat dams ([44]). To determine if this includes the PMv^{DAT} neurons, we examined c-Fos immunoreactivity. Both the total number of c-Fos^{+} neurons, and the proportion of DAT^{+} cells containing c-Fos^{+} immunoreactivity, were increased following the expression of maternal aggression (Figure 1C, D, Table S1).

Having established that PMv^{DAT} neurons are in a hyperexcitable state in lactating dams compared to the non-aggressive virgin female mouse, we next applied genetic activation, inactivation and deletion of these cells in order to investigate the effect of such manipulations *in vivo* on the expression of maternal aggression. Genetic modifications were introduced into PMv^{DAT} neurons by stereotactic injection of Cre-dependent constructs within an AAV vector.

Bidirectional control of maternal aggression through PMv^{DAT} neurons

PMv^{DAT} neurons in lactating dams were transduced with ChR2 (Figure 2B) and baseline behavior was recorded in trials that were characterized by close investigation, but rare expressions of aggression against male and female conspecifics (Figure 2A). Optogenetic stimulation resulted in significantly prolonged attack bouts appearing within seconds of initiation of photostimulation (Figure 2C-E, Video S1, Table S1).

To determine the behavioral effect of inhibition of the PMv^{DAT} population, juvenile female intruders were used to facilitate high levels of maternal aggression in baseline conditions(*38, 45*) (Figure 2F). Indeed, under control conditions, lactating dams (Figure 2G) frequently engaged in sustained attacks. However, application of optogenetic inhibition in these mice was followed by decreased attack bout duration and delayed initiation of subsequent attack episodes (Figure 2H-J, Table S1).

Lastly, genetic deletion of the PMv^{DAT} neurons in lactating dams (Figure 2K, L) - sparing neighbouring DAT neuron populations (Figure S1A, B and Table S1) – dramatically decreased the attack frequency and duration and increased latency to attack (Figure 2M-O, Table S1). Together, these experiments establish a causal role for PMv^{DAT} neurons in the expression of maternal aggression.

Maternal hormones excite PMv^{DAT} neurons

Endocrine hallmarks of the *post partum* female include a surge in the serum levels of specific hormones (*46-50*). Among these hormones, the most well-documented maternal functions are served by prolactin (Prl) (*51*), which rises in late pregnancy and remains elevated after birth in rodents (*52-54*), facilitating maternal pup care behavior and offspring survival (*55-57*). Oxytocin (OT), which shows similar dynamics, plays a parallel and complementary role in the mother's care for offspring (*58-61*). We next explored if these maternal hormones can affect the activity of PMv^{DAT} neurons that the data above implicate in puerperal aggression.

Serum levels of Prl were found ten-fold higher levels in lactating mouse dams compared to virgin females at any point across the estrous cycle (Figure 3A, Table S1), in agreement with earlier literature (*62*). We next determined if this difference in circulating Prl corresponds to differential Prl receptor (Prl-R) activation in the PMv of virgin *vs* lactating female mice. Indeed, phosphorylation of the Signal Transducer and Activator of Transcription 5 (pSTAT5), a key and sensitive downstream mediator of Prl-R activation (*63-74*), was significantly elevated in PMv^{DAT} neurons in lactating females (Figure 3B-D, Table S1). These findings prompted the question if PMv^{DAT} cells can be directly modulated by Prl.

Application of Prl to PMv^{DAT} neurons during *in vitro* patch clamp recording yielded several electrophysiological changes, which can increase the excitability and output of these cells. Firstly, a prominent and reversible depolarization, paralleled by increased discharge frequency, was seen (Figure 3E, Table S1), similar to observations from genetically unidentified PMv neurons (*75*).

Secondly, action potential duration was significantly prolonged (Figure 3F, Table S1), a change that can increase neurotransmitter output by over a magnitude (*76*). Thirdly, the capacity for and strength of post-inhibitory rebound were augmented (Figure 3G, Table S1), key features of PMv[DAT] neurons and implicated in their propensity for regenerative discharge (*20*). The post-synaptic Prl-induced current in PMv[DAT] cells consisted of a low- and a high-voltage component (Figure 3H, Table S1). Application of the selective T-type low-threshold Ca^{2+} current blocker, ML-218, abolished the low-voltage component (Figure 3I, Table S1). Lastly we tested whether application of the T-type Ca^{2+} channel blocker impaired the post-inhibitory rebound, and the effect of prolactin specific to the potentiation of the rebound excitation. Indeed, ML218 abolished the post-inhibitory rebound and the effect of prolactin (Figure 3J, Table S1).

Interestingly, the application of Prl also affected synaptic input into PMv[DAT] neurons, with an increase of both spontaneous excitatory and inhibitory post-synaptic currents (data not shown). A likely source of the increased excitatory synaptic currents is recurrent axon collaterals, as the glutamatergic PMv[DAT] population is densely interconnected (*20*). These data reveal a wide repertoire of Prl actions on the membrane properties of PMv[DAT] neurons that together converge to increase excitability.

Oxytocin, beyond its role in milk ejection (*77-79*), is most commonly discussed as an agent of maternal care and social bonding (*80, 81*). Previous studies have, however, also correlated increased serum OT levels to maternal aggression (*82-84*). Yet, the neural substrates underlying this effect remain unknown. Immunofluorescence staining of the PMv identified close appositions between OT positive fibers and PMv[DAT] cell somata (Figure 3K). This finding, and the presence of OT receptors (OT-R's) in the PMv (*85*), led us to evaluate responsiveness of PMv[DAT] neurons to OT. Application of OT led to a reversible depolarization of PMv[DAT] neurons (Figure 3L, Table S1), similar to the effect of Prl (*vide supra*), as did application of the selective OT-R agonist, TGOT (*86, 87*)(Figure 3M, Table S1). However, the effect of OT persisted also in the presence of an OT-R antagonist (vasotocin [d(CH2)$_5^1$, Tyr(Me)^2, Thr^4, Orn^8, des-Gly NH$_2^9$]), indicating that vasopressin receptors (*88-93*) contribute to the peptide's action on PMv[DAT] neurons (Figure 3M, Table S1). The OT effect depended on a post-synaptic depolarization, composed of low- and high-voltage components similar to the Prl-induced current (Figure 3N, Table S1).

To investigate whether maternal hormones acting in the PMv are sufficient to trigger aggression in an inherently non-aggressive female, we performed a 28-day localized application of Prl, OT, or both peptide hormones (co-administration) unilaterally into the nucleus of virgin female adult mice. Two weeks after implantation we performed a panel of tests, to investigate changes in aggressive and/or maternal behavior, as well as other behavioral features. No attacks or changes in the degree of close investigation were observed in the RI test (Figure S2A, B and Table S1). Notably, however, local co-administration of maternal hormones in PMv decreased sociability in the group (Figure S2C,

Table S1) and, importantly, impaired maternal behavior (Figure S2D, Table S1). Modest changes in locomotion, but not anxiety parameters, were identified (Figure S2E-F, Table S1).

Together these experiments identify that PMv-targetted application of maternal hormones in non-lactating females does not lead to the expression of aggression, suggestive of necessary upstream and/or downstream changes implemented in the brains of lactating dams. However, of interest are the observations that administration of maternal hormones in PMv in non-lactating mice, leads to a decrease in sociability (Figure S2C) and impairs performance in the pup-retrieval test (Figure S2D).

Activation of PMv^{DAT} neurons impairs maternal care

Aggression in dams is part of a behavioral program that turns on at parturition and declines by weaning (*38, 94*). This program includes a coincident initiation of pup care – such as nursing, grooming and crouching, and maternal aggression – aimed to inflict injury and deter infanticidal conspecifics or predators from attacking the pups (*94*). In the final experiment, we tested whether *pup care* and *maternal aggression* can be triggered by common anatomical substrates – such as PMv, or whether activation of defined brain loci/circuitry leads to the execution of dedicated/non-overlapping behavioral programs in expense of other behaviors.

To investigate this issue, we tested the effect of optogenetic activation of PMv^{DAT} neurons in the absence of a conspecific – during a pup retrieval test (*95, 96*). The pup retrieval test was performed in lactating dams expressing ChR2 or eYFP in PMv^{DAT} neurons (Figure 4A, B). Control (eYFP-expressing) dams retrieved all pups within a minute of the task initiation during photostimulation (Figure 4B, D, E, Video S2, Table S1). In contrast, photostimulated ChR2-expressing dams exhibited severely impaired pup-retrieval behavior, and in many cases, failed to retrieve any pups over the maximum test period of three minutes (Figure 4B-E, Video S3, Table S1). No aggression towards the pups was observed in any of the trials.

These data suggest that activation of these neurons promotes a specialized behavioral program – the one of maternal aggression, in expense of other behaviors. This is illustrated in the data discussed here, where even in the absence of a competing intruder stimulus, normal nursing behavior is inhibited. It is likely that PMv^{DAT} neuron activity serves multiple functions, facilitating the expression of maternal aggression and inhibiting the expression of antagonistic behaviors.

Discussion

The data presented here, identify a PMv neuron subpopulation in the hypothalamus as a key node in orchestrating maternal aggression but not maternal care. The identification of the node's activation post a maternal aggression episode together with the finding that maternal hormones act upon the PMv[DAT] cells in an excitatory manner provide a long-missing link on the neurobiological understanding of aggressive behavior in females.

These data suggest that the sexual dimpoirtphism in the expression of aggressive behavior likely does not stem from a fundamentally different wiring between the male and the female brain. Rather, a common framework for aggression exists across the two sexes that can be recruited for adaptive purposes (e.g. to establish social rank among males or to secure the survival of pups in post-partum females). Remains a conundrum how this in the end yields different motor expression – pattern of behavior – and what the underlying switch consists of.

It's important to note here, that PMv[DAT] neurons have been associated with a number of social behaviors in male mice (*18-20*), primarily with aggressive behavior. Together these findings suggest that at least part of the aggression circuit active in males, is recruited via the increase of maternal hormones in the serum of lactating females. Of particular interest is the observation that the increase in serum Prl likely leads to activation of numerous nuclei in the CNS to induce the collective behavioral repertoire known as maternal behavior, with classically defined adaptations including energy homeostasis (*97*), nursing behavior (*57*), and aggression (*98-100*). While recent work has provided evidence for the role of Prl-R activation in the medial preoptic area of the hypothalamus specifically in nursing behavior (*57*), here we suggest that Prl-R mediated activation of PMv and of the downstream circuit recruits the maternal aggression behavioral repertoire, while suppressing competing behavioral programs.

Finally, application of maternal hormones locally in the PMv of adult females was shown to be insufficient to induce aggressive behavior in female mice indicative of the necessary circuit adaptations upstream to PMv. Future work is needed in order to further identify the necessary sites of action of maternal hormones, whose activation enables the expression of aggression in females.

References

1. D. Maestripieri, Costs and Benefits of Maternal Aggression in Lactating Female Rhesus Macaques. *Primates; journal of primatology* **35**, 443-453 (1994).
2. J. P. Kent, Maternal Aggression and Interindividual Distance in the Broody Hen (Gallus-Gallus). *Behav Process* **27**, 37-44 (1992).
3. A. Troisi, F. R. Damato, A. Carnera, L. Trinca, Maternal Aggression by Lactating Group-Living Japanese Macaque Females. *Hormones and behavior* **22**, 444-452 (1988).
4. B. Olivier *et al.*, Maternal Aggression in Rats. *Aggressive Behav* **11**, 169-169 (1985).
5. E. R. Stockman, Aggressive Experience and Maternal Aggression in the Mongolian Gerbil. *Physiology & behavior* **30**, 319-321 (1983).
6. H. I. Siegel, A. L. Giordano, C. M. Mallafre, J. S. Rosenblatt, Maternal Aggression in Hamsters - Effects of Stage of Lactation, Presence of Pups, and Repeated Testing. *Hormones and behavior* **17**, 86-93 (1983).
7. R. D. Stjohn, P. A. Corning, Maternal Aggression in Mice. *Behavioral biology* **9**, 635-639 (1973).
8. M. H. Figler, G. S. Blank, H. V. S. Peeke, Maternal aggression and post-hatch care in red swamp crayfish, Procambarus clarkii (Girard): The influences of presence of offspring, fostering, and maternal molting. *Mar Freshw Behav Phy* **30**, 173-194 (1997).
9. K. Hashikawa *et al.*, Esr1(+) cells in the ventromedial hypothalamus control female aggression. *Nature neuroscience* **20**, 1580-1590 (2017).
10. N. M. Rendon, G. E. Demas, Bi-directional actions of dehydroepiandrosterone and aggression in female Siberian hamsters. *Journal of experimental zoology. Part A, Ecological genetics and physiology* **325**, 116-121 (2016).
11. L. E. Been, K. M. Moore, B. C. Kennedy, R. L. Meisel, Metabotropic Glutamate Receptor and Fragile X Signaling in a Female Model of Escalated Aggression. *Biological psychiatry* **79**, 685-692 (2016).
12. J. S. Lonstein, S. C. Gammie, Sensory, hormonal, and neural control of maternal aggression in laboratory rodents. *Neuroscience and biobehavioral reviews* **26**, 869-888 (2002).
13. M. Liu, D. W. Kim, H. Zeng, D. J. Anderson, Make war not love: The neural substrate underlying a state-dependent switch in female social behavior. *Neuron* **110**, 841-856 e846 (2022).
14. T. Georgescu, Z. Khant Aung, D. R. Grattan, R. S. E. Brown, Prolactin-mediated restraint of maternal aggression in lactation. *Proceedings of the National Academy of Sciences of the United States of America* **119**, (2022).
15. M. R. Kruk *et al.*, Discriminant analysis of the localization of aggression-inducing electrode placements in the hypothalamus of male rats. *Brain research* **260**, 61-79 (1983).
16. J. C. Cavalcante, J. C. Bittencourt, C. F. Elias, Female odors stimulate CART neurons in the ventral premammillary nucleus of male rats. *Physiology & behavior* **88**, 160-166 (2006).
17. M. Yokosuka *et al.*, Female-soiled bedding induced Fos immunoreactivity in the ventral part of the premammillary nucleus (PMv) of the male mouse. *Physiology & behavior* **68**, 257-261 (1999).
18. M. E. Soden *et al.*, Genetic Isolation of Hypothalamic Neurons that Regulate Context-Specific Male Social Behavior. *Cell reports* **16**, 304-313 (2016).
19. D. Y. Lin *et al.*, Functional identification of an aggression locus in the mouse hypothalamus. *Nature* **470**, 221-+ (2011).
20. S. Stagkourakis *et al.*, A neural network for intermale aggression to establish social hierarchy. *Nature neuroscience* **21**, 834-+ (2018).
21. A. X. Chen *et al.*, Specific Hypothalamic Neurons Required for Sensing Conspecific Male Cues Relevant to Inter-male Aggression. *Neuron*, (2020).
22. J. C. Cavalcante, J. C. Bittencourt, C. F. Elias, Distribution of the neuronal inputs to the ventral premammillary nucleus of male and female rats. *Brain research* **1582**, 77-90 (2014).
23. N. S. Canteras, R. B. Simerly, L. W. Swanson, Projections of the Ventral Premammillary Nucleus. *Journal of Comparative Neurology* **324**, 195-212 (1992).

24. M. M. Sanchez, F. Aguado, F. Sancheztoscano, D. Saphier, Effects of Prolonged Social-Isolation on
Responses of Neurons in the Bed Nucleus of the Stria Terminalis, Preoptic Area, and Hypothalamic
Paraventricular Nucleus to Stimulation of the Medial Amygdala. *Psychoneuroendocrinology* **20**, 525-
541 (1995).
25. N. Duque-Wilckens, M. Steinman, V. Grinevich, B. Trainor, The Role of Oxytocin Neurons in the Bed
Nucleus of the Stria Terminalis in Mediating Social Withdrawal. *Biological psychiatry* **81**, S44-S45
(2017).
26. K. Lei, B. S. Cushing, S. Musatov, S. Ogawa, K. M. Kramer, Estrogen Receptor-alpha in the Bed Nucleus
of the Stria Terminalis Regulates Social Affiliation in Male Prairie Voles (Microtus ochrogaster). *PloS*
one **5**, (2010).
27. A. M. Jasnow, M. Davis, K. L. Huhman, Involvement of central amygdalar and bed nucleus of the stria
terminalis corticotropin-releasing factor in behavioral responses to social defeat. *Behavioral*
neuroscience **118**, 1052-1061 (2004).
28. D. J. Albert, M. L. Walsh, B. B. Gorzalka, S. Mendelson, C. Zalys, Intermale Social Aggression -
Suppression by Medial Preoptic Area Lesions. *Physiology & behavior* **38**, 169-173 (1986).
29. J. S. Rosenblatt, S. Hazelwood, J. Poole, Maternal behavior in male rats: Effects of medial preoptic area
lesions and presence of maternal aggression. *Hormones and behavior* **30**, 201-215 (1996).
30. N. Scott, M. Prigge, O. Yizhar, T. Kimchi, A sexually dimorphic hypothalamic circuit controls maternal
care and oxytocin secretion. *Nature* **525**, 519-+ (2015).
31. H. Vogel *et al.*, GLP-1 and estrogen conjugate acts in the supramammillary nucleus to reduce food-
reward and body weight. *Neuropharmacology* **110**, 396-406 (2016).
32. R. Shin, S. Ikemoto, Administration of the GABA(A) receptor antagonist picrotoxin into rat
supramammillary nucleus induces c-Fos in reward-related brain structures. Supramammillary
picrotoxin and c-Fos expression. *BMC neuroscience* **11**, (2010).
33. S. Ikemoto, B. M. Witkin, A. Zangen, R. A. Wise, Rewarding effects of AMPA administration into the
supramammillary or posterior hypothalamic nuclei but not the ventral tegmental area. *Journal of*
Neuroscience **24**, 5758-5765 (2004).
34. S. A. Golden *et al.*, Basal forebrain projections to the lateral habenula modulate aggression reward.
Nature **534**, 688-692 (2016).
35. S. A. Golden *et al.*, Compulsive Addiction-like Aggressive Behavior in Mice. *Biological psychiatry* **82**,
239-248 (2017).
36. D. Maestripieri, E. Alleva, Maternal Aggression and Litter Size in the Female House Mouse. *Ethology*
84, 27-34 (1990).
37. K. J. Flannelly, E. D. Kemble, The Effect of Pup Presence and Intruder Behavior on Maternal Aggression
in Rats. *B Psychonomic Soc* **25**, 133-135 (1987).
38. M. Haney, J. F. Debold, K. A. Miczek, Maternal Aggression in Mice and Rats Towards Male and Female
Conspecifics. *Aggressive Behav* **15**, 443-453 (1989).
39. J. Mos, B. Olivier, R. Vanoorschot, Maternal Aggression Towards Different Sized Male Opponents -
Effect of Chlordiazepoxide Treatment of the Mothers and D-Amphetamine Treatment of the Intruders.
Pharmacol Biochem Be **26**, 577-584 (1987).
40. S. Stagkourakis, G. Spigolon, G. Liu, D. J. Anderson, Experience-dependent plasticity in an innate social
behavior is mediated by hypothalamic LTP. *Proceedings of the National Academy of Sciences of the*
United States of America **117**, 25789-25799 (2020).
41. S. Stagkourakis *et al.*, A neural network for intermale aggression to establish social hierarchy. *Nature*
neuroscience **21**, 834-842 (2018).
42. S. M. Howard, R. Gandelman, C. Rosenthal, Isolation Potentiates the Aggression-Activating Property
of Testosterone in Female Mice. *Physiology & behavior* **26**, 971-972 (1981).
43. M. S. Barkley, B. D. Goldman, Testosterone-Induced Aggression in Adult Female Mice. *Hormones and*
behavior **9**, 76-84 (1977).

44.   S. C. Motta *et al.*, Ventral premammillary nucleus as a critical sensory relay to the maternal aggression
355   network. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 14438-
356   14443 (2013).

357   45.   K. J. Flannelly, L. Flannelly, Opponents Size Influences Maternal Aggression. *Psychological reports* **57**,
358   883-886 (1985).

359   46.   M. D. M. Hallak, J. W. Hotra, Maternal serum hormone concentrations in Long-Evans rats. *J Trace
360   Microprobe T* **21**, 103-110 (2003).

361   47.   C. M. Dwyer, A. B. Lawrence, Induction of maternal behaviour in non-pregnant, hormone-primed
362   ewes. *Anim Sci* **65**, 403-408 (1997).

363   48.   J. H. Duenhoelter, Maternal Hormone Measurements. *Jama-J Am Med Assoc* **248**, 1109-1109 (1982).

364   49.   C. D. Hansult, S. Kessler, Role of Hormones in Maternal Aggression and Infant Care. *Behavior genetics*
365   **7**, 65-66 (1977).

366   50.   M. X. Zarrow, V. H. Denenberg, R. Gandelman, Prolactin - Is It an Essential Hormone for Maternal
367   Behavior in Mammal. *Hormones and behavior* **2**, 343-+ (1971).

368   51.   O. Riddle, Prolactin in Vertebrate Function and Organization. *Jnci-J Natl Cancer I* **31**, 1039-+ (1963).

369   52.   M. Lhermite, Gonadotropin and Prolactin Function during Pregnancy and Postpartum. *Rev Fr Gynecol
370   Obste* **71**, 159-165 (1976).

371   53.   H. Moltz, R. Levin, M. Leon, Prolactin in Postpartum Rat - Synthesis and Release in Absence of Suckling
372   Stimulation. *Science* **163**, 1084-& (1969).

373   54.   J. W. Riesen, Saiduddi.S, W. E. Graves, W. J. Tyler, L. E. Casida, Pituitary Prolactin Activity in Postpartum
374   Dairy Cows. *Journal of animal science* **25**, 929-& (1966).

375   55.   M. Numan, H. Moltz, M. Leon, Interference with Prolactin Release and Maternal Behavior of Female
376   Rats. *Hormones and behavior* **3**, 29-& (1972).

377   56.   R. S. Bridges, B. D. Goldman, L. P. Bryant, Serum Prolactin Concentrations and Initiation of Maternal-
378   Behavior in Rat. *Hormones and behavior* **5**, 219-226 (1974).

379   57.   R. S. E. Brown *et al.*, Prolactin action in the medial preoptic area is necessary for postpartum maternal
380   nursing behavior. *Proceedings of the National Academy of Sciences of the United States of America*
381   **114**, 10779-10784 (2017).

382   58.   C. A. Pedersen, A. J. Prange, Jr., Induction of maternal behavior in virgin rats after
383   intracerebroventricular administration of oxytocin. *Proceedings of the National Academy of Sciences
384   of the United States of America* **76**, 6661-6665 (1979).

385   59.   C. A. Pedersen, J. A. Ascher, Y. L. Monroe, A. J. Prange, Jr., Oxytocin induces maternal behavior in virgin
386   female rats. *Science* **216**, 648-650 (1982).

387   60.   A. Watarai *et al.*, The blockade of oxytocin receptors in the paraventricular thalamus reduces maternal
388   crouching behavior over pups in lactating mice. *Neuroscience letters* **720**, 134761 (2020).

389   61.   F. Champagne, J. Diorio, S. Sharma, M. J. Meaney, Naturally occurring variations in maternal behavior
390   in the rat are associated with differences in estrogen-inducible central oxytocin receptors. *Proceedings
391   of the National Academy of Sciences of the United States of America* **98**, 12736-12741 (2001).

392   62.   A. Guillou *et al.*, Assessment of lactotroph axis functionality in mice: longitudinal monitoring of PRL
393   secretion by ultrasensitive-ELISA. *Endocrinology* **156**, 1924-1930 (2015).

394   63.   F. Gouilleux, H. Wakao, M. Mundt, B. Groner, Prolactin Induces Phosphorylation of Tyr694 of Stat5
395   (Mgf), a Prerequisite for DNA-Binding and Induction of Transcription. *Embo Journal* **13**, 4361-4369
396   (1994).

397   64.   L. DaSilva *et al.*, Prolactin recruits STAT1, STAT3 and STAT5 independent of conserved receptor
398   tyrosines TYR402, TYR479, TYR515 and TYR580. *Molecular and cellular endocrinology* **117**, 131-140
399   (1996).

400   65.   S. Ali, S. Ali, Prolactin receptor regulates Stat5 tyrosine phosphorylation and nuclear translocation by
401   two separate pathways. *Journal of Biological Chemistry* **273**, 7709-7716 (1998).

402   66.   A. Lerant, B. Kanyicska, M. E. Freeman, Nuclear translocation of STAT5 and increased expression of
403   Fos related antigens (FRAs) in hypothalamic dopaminergic neurons after prolactin administration.
404   *Brain research* **904**, 259-269 (2001).

67. J. L. Brockman, L. A. Schuler, Prolactin signals via Stat5 and Oct-1 to the proximal cyclin D1 promoter.
Molecular and cellular endocrinology **239**, 45-53 (2005).

68. B. J. Cave *et al.*, Prolactin-induced activation of STAT5 within the hypothalamic arcuate nucleus.
Neuroreport **16**, 1423-1426 (2005).

69. S. Jaroenporn, K. Nagaoka, R. Ohta, G. Watanabe, K. Taya, Prolactin induces phosphorylation of the
STAT5 in adrenal glands of Hatano rats during stress. *Life sciences* **85**, 172-177 (2009).

70. J. D. Buntin, L. Buntin, Increased STAT5 signaling in the ring dove brain in response to prolactin
administration and spontaneous elevations in prolactin during the breeding cycle. *General and*
comparative endocrinology **200**, 1-9 (2014).

71. R. S. E. Brown, I. C. Kokay, A. E. Herbison, D. R. Grattan, Distribution of Prolactin-Responsive Neurons
in the Mouse Forebrain. *Journal of Comparative Neurology* **518**, 92-102 (2010).

72. R. A. Augustine *et al.*, Prolactin regulation of oxytocin neurone activity in pregnancy and lactation. *J*
Physiol-London **595**, 3591-3605 (2017).

73. R. S. E. Brown, A. E. Herbison, D. R. Grattan, Effects of Prolactin and Lactation on A15 Dopamine
Neurones in the Rostral Preoptic Area of Female Mice. *Journal of neuroendocrinology* **27**, 708-717
(2015).

74. R. S. E. Brown, R. Piet, A. E. Herbison, D. R. Grattan, Differential Actions of Prolactin on Electrical
Activity and Intracellular Signal Transduction in Hypothalamic Neurons. *Endocrinology* **153**, 2375-2384
(2012).

75. M. A. Silveira *et al.*, Acute effects of somatomammotropin hormones on neuronal components of the
hypothalamic-pituitary-gonadal axis. *Brain research* **1714**, 210-217 (2019).

76. B. L. Sabatini, W. G. Regehr, Control of neurotransmitter release by presynaptic waveform at the
granule cell to Purkinje cell synapse. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **17**, 3425-3435 (1997).

77. A. Kamikawa, J. Seko, Physiological and pharmacological evaluation of oxytocin-induced milk ejection
in mice. *Experimental animals* **69**, 345-353 (2020).

78. Y. F. Wang, G. I. Hatton, Milk ejection burst-like electrical activity evoked in supraoptic oxytocin
neurons in slices from lactating rats. *Journal of neurophysiology* **91**, 2312-2321 (2004).

79. J. D. Donker, J. H. Koshi, W. E. Petersen, The effect of exogenous oxytocin in blocking the normal
relationship between endogenous oxytocic substance and the milk ejection phenomenon. *Science*
119, 67-68 (1954).

80. C. Finkenwirth, E. Martins, T. Deschner, J. M. Burkart, Oxytocin is associated with infant-care behavior
and motivation in cooperatively breeding marmoset monkeys. *Hormones and behavior* **80**, 10-18
(2016).

81. H. E. Ross, L. J. Young, Oxytocin and the neural mechanisms regulating social cognition and affiliative
behavior. *Frontiers in neuroendocrinology* **30**, 534-547 (2009).

82. J. P. Stohn *et al.*, Increased aggression and lack of maternal behavior in Dio3-deficient mice are
associated with abnormalities in oxytocin and vasopressin systems. *Genes Brain and Behavior* **17**, 23-
35 (2018).

83. O. J. Bosch, Maternal aggression in rodents: brain oxytocin and vasopressin mediate pup defence.
Philos T R Soc B **368**, (2013).

84. O. J. Bosch, S. L. Meddle, D. I. Beiderbeck, A. J. Douglas, I. D. Neumann, Brain oxytocin correlates with
maternal aggression: Link to anxiety. *Journal of Neuroscience* **25**, 6807-6815 (2005).

85. P. Kremarik, M. J. Freund-Mercier, M. E. Stoeckel, Oxytocin and vasopressin binding sites in the
hypothalamus of the rat: histoautoradiographic detection. *Brain research bulletin* **36**, 195-203 (1995).

86. M. Zaninetti, M. Raggenbass, Oxytocin receptor agonists enhance inhibitory synaptic transmission in
the rat hippocampus by activating interneurons in stratum pyramidale. *The European journal of*
neuroscience **12**, 3975-3984 (2000).

87. J. Elands, C. Barberis, S. Jard, [3H]-[Thr4,Gly7]OT: a highly selective ligand for central and peripheral
OT receptors. *The American journal of physiology* **254**, E31-38 (1988).

88. J. Gruskos, A. Goldwaser, D. A. Guarracino, Competition of peptide hormone mimics with vasopressin and oxytocin for binding their target membrane receptors. *Abstr Pap Am Chem S* **244**, (2012).

89. M. Hibert *et al.*, Functional architecture of vasopressin/oxytocin receptors. *Journal of receptor and signal transduction research* **19**, 589-596 (1999).

90. H. H. Zingg, Vasopressin and oxytocin receptors. *Bailliere Clin Endoc* **10**, 75-96 (1996).

91. M. Manning, S. Stoev, W. Y. Chan, W. H. Sawyer, Receptor-Specific Antagonists of Vasopressin and Oxytocin - a Current Perspective. *Neurohypophysis : A Window on Brain Function* **689**, 219-232 (1993).

92. S. Jard, J. Elands, A. Schmidt, C. Barberis, Vasopressin and Oxytocin Receptors - an Overview. *Progress in Endocrinology 1988, Vol 1 and 2* **799**, 1183-1188 (1988).

93. P. J. Martin, H. O. Schild, Interaction of Oxytocin and Vasopressin with Tissue Receptors. *Biochemical pharmacology* **12**, 139-& (1963).

94. R. D. St John, P. A. Corning, Maternal aggression in mice. *Behavioral biology* **9**, 635-639 (1973).

95. A. Zimprich *et al.*, Assessing Sociability, Social Memory, and Pup Retrieval in Mice. *Current protocols in mouse biology* **7**, 287-305 (2017).

96. A. Bendesky *et al.*, The genetic basis of parental care evolution in monogamous mice. *Nature* **544**, 434-439 (2017).

97. S. R. Ladyman *et al.*, Prolactin receptors in Rip-cre cells, but not in AgRP neurones, are involved in energy homeostasis. *Journal of neuroendocrinology* **29**, (2017).

98. J. A. Barry *et al.*, Prolactin and aggression in women with fertility problems. *Journal of Obstetrics and Gynaecology* **34**, 605-610 (2014).

99. A. R. Consiglio, R. S. Bridges, Circulating prolactin, MPOA prolactin receptor expression and maternal aggression in lactating rats. *Behavioural brain research* **197**, 97-102 (2009).

100. P. E. Gleason, S. D. Michael, J. J. Christian, Prolactin-Induced Aggression in Female Peromyscus-Leucopus. *Behavioral and neural biology* **33**, 243-248 (1981).

101. M. I. Ekstrand *et al.*, Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 1325-1330 (2007).

102. K. B. J. Franklin, G. Paxinos, *THE MOUSE BRAIN IN STEREOTACTIC COORDINATES*. (2008), vol. Third edition.

103. R. Behringer, M. Gertsenstein, K. V. Nagy, A. Nagy, Selecting Female Mice in Estrus and Checking Plugs. *Cold Spring Harb Protoc* **2016**, (2016).

104. A. X. Chen *et al.*, Specific Hypothalamic Neurons Required for Sensing Conspecific Male Cues Relevant to Inter-male Aggression. *Neuron* **108**, 763-774 e766 (2020).

105. T. Itakura *et al.*, A single vomeronasal receptor promotes intermale aggression through dedicated hypothalamic neurons. *Neuron* **110**, 2455-2469 e2458 (2022).

Materials and methods

Animals. All animal experiments had received approval from the local ethical board, *Stockholms Norra Djurförsöksetiska Nämnd*, and were performed in accordance with the European Communities Council Directive of November 24, 1986 (86/609/EEC). Wild-type mice with C57BL/6J and BALB/c background were used, in addition to previously generated C57BL/6J *Slc6a*[Cre] (DAT-Cre) knock-in(*101*) and floxed-tdTomato mice (The Jackson Laboratory, strain datasheet – 007909). Animals were group-housed, up to five per cage, in a temperature (23°C) and humidity (55%) controlled environment, in a 12h light, 12h dark cycle with *ad libitum* access to food and water. Cages were changed on a weekly basis.

Adult female virgin, and lactating female mice used in experiments were single housed for a period of two to four weeks, depending on the experimental design. Only multiparous lactating mice were used in the experimental group of *lactating dams*, typically following their third or fourth parturition.

Viral vectors. For channelrhodopsin optogenetic studies animals were injected in the PMv with 400 nL of AAV5-EF1a-DIO-hChR2(H134R)-eYFP-WPRE-hGH (Addgene20298) 8.41×10^{12} genomic copies per mL. For halorhodopsin-mediated neuronal silencing, animals were injected with 400 nL of AAV5-EF1a-DIO-eNpHR3.0-eYFP-WPRE-hGH (Addgene26966) 7.02×10^{12} genomic copies per mL. The control groups in optogenetic were injected with 400 nL of AAV5-EF1a-DIO-eYFP-WPRE-hGH (Addgene27056) 5.82×10^{12} genomic copies per mL. The ChR2, eNpHR3 and eYFP AAV5 were prepared by the University of Pennsylvania Vector Core. For targeted cell ablation of DAT$^+$ PMv neurons animals were injected in PMv with 300 nL of AAV5-flex-taCasp3-TEVp 2.9×10^{12} genomic copies per mL, and the AAV was prepared by the viral vector core at the University of North Carolina. Viral injections were performed bilaterally.

Stereotactic surgery and viral gene transfer. Adult DAT-Cre mice of 5 to 8 months age (sexually inexperienced) were stereotactically injected with a virus, and implanted with fiber implants when purposed for *in vivo* optogenetic behavior experiments, and were individually housed for 2 weeks post-surgery. Animals were anaesthetized with isoflurane (1–5%) and placed in a stereotaxic frame (David Kopf Instruments). Virus was injected into the PMv bilaterally using a pulled glass capillary (World Precision Instruments) by nanolitre pressure injection at a flow rate of 50 nL per min (Micro4 controller, World Precision Instruments; Nanojector II, Drummond Scientific). Stereotactic injection coordinates to target the PMv were obtained from the Paxinos and Franklin atlas (*102*) (Bregma: -2.45 mm, midline ±0.6 mm, dorsal surface -5.5 mm). Ferrules and fiber-optic patch cords were purchased from Thorlabs and Doric Lenses, respectively. The virus-injected animals were housed individually during a 2-week recovery period, and then examined behaviorally and histologically.

In the case of the *lactating dams* group, sexually experienced female mice with prior maternal experience were placed in a cage with a male mouse, and were checked for plugs in the morning hours of the next four days (*103*). Following identification of plugs, female mice were single housed, and surgeries were performed for AAV injections, and/or optic fiber implants. The surgical procedures occurred within a week of plug identification. Female mice in which plug identification did not occur or was ambiguous were excluded from the study.

Optogenetics. In optogenetic experiments, subjects were coupled via a ferrule patch cord to a ferrule on the head of the mouse using a zirconia split sleeve (Doric Lenses). The optical fiber was connected to a laser (447 nm for ChR2; 635 nm for eNpHR3; CNI-MLL-III-447-200-5-LED and CNI-MLL-III-635-200-5-LED, CNI lasers 200 mW) via a fiber-optic rotary joint (FRJ_1x1_FC-FC, Doric Lenses) to avoid twisting of the cable caused by the animal's movement. After a testing session, DAT-Cre animals were uncoupled from the fiber-optic cable and returned to a housing room. The frequency and duration of photostimulation were controlled using custom written LabView software. Laser power was controlled by dialling an analogue knob on the power supply of the laser sources. Light power was measured from the tip of the ferrule in the patch cord before being installed in the brain (the ferrule-connector system) at different laser output settings, using an optical power and energy meter and a photodiode power sensor (Thorlabs). Upon identification of the fiber placement coordinates in brain tissue slides, irradiance (light intensity) was calculated using the brain tissue light transmission calculator based on (http://www.stanford.edu/group/dlab/cgi-bin/graph/chart.php) using laser power measured at the tip and the distance from the tip to the target brain region measured by histology. Animals showing no detectable viral expression in the target region and/or ectopic fiber placement were excluded from analysis. In some experiments, animals were photostimulated with a train of 473 nm light (20 Hz, 5 msec, 5 min) 45 min before perfusion in the absence of an intruder at an intensity which had evoked a behavioral phenotype in the final testing session. Then, brain sections were immunohistochemically labelled for c-Fos to identify optogenetically activated cells.

Behavioral tests. Behavioral tests were performed at 2h post-initiation of the light phase and 2h prior to initiation of the dark phase. Mice were acclimated to the testing facility for 1 h prior to testing. Behaviors were recorded using a digital video recording unit. Behavioral annotations were performed manually, with conditions blinded to the experimenter performing the behavioral scoring.

Resident-intruder (RI) test. RI tests were initiated at 2 to 4 weeks post-surgery and repeated weekly for 2 to 5 weeks. All dams used in the present study were multiparous and were used in behavioral experiments during their third or fourth litter. Mouse cages were not cleaned for a minimum of 3 days prior to the behavioral test. Intruders were individually introduced to a DAT-Cre mouse in a testing session in a random order with respect to sex, with a 5 min interval between intruders. The

strains of intruders were: juvenile females and adult females. Subjects involved in the RI test were exposed to 3 experimental days of the test, resulting in multiple measurements per resident.

ChR2-mediated activation in RI. After the introduction of an intruder, a virus-injected animal was recorded for 3 min to assess baseline behavior towards each intruder. The baseline recording was followed by photostimulation trials (3 min in duration) with varying irradiance (intensity), stimulation intervals, distance, orientation and recent behavior history of the two animals at the onset of photostimulation. The AAV5-DIO-eYFP and AAV5-DIO-ChR2 injected experimental animals were processed in random order.

eNpHR3-mediated silencing in RI. DAT-Cre lactating females expressing eNpHR3 or control eYFP were introduced to one to four juvenile female intruders, in 3 acclimation sessions without photostimulation to assess baseline aggression as well as to augment aggressiveness. Testing sessions initiated with recordings in the absence of laser stimulation to assess baseline behavior towards each intruder on the day of the experiment. In photostimulation trials, irradiance (intensity) ranging from 2.1 – 11.6 mW/mm^2 was delivered continuously in varying intervals (typically 2-10 sec) depending on the phase of the resident-intruder interaction. To examine whether photostimulation during charging stopped escalation to attack, residents were photostimulated during body realignment to gain access to the back of the intruder.

Pup retrieval test. Adult virgin female or lactating mice 5-8 months old were exposed to foster pups or to their own litter, respectively. The litter size per test is described in the individual tests/figures. Nesting material was provided in each cage for building a nest. The activity in the cage was video recorded, and maternal behaviors were scored by an experimenter blinded to the experimental conditions.

Three-chambered sociability test. The social approach apparatus was an open-topped box made of acrylic (60 cm in length × 40 cm in width × 20 cm in height), and divided into three chambers with two clear acrylic walls. Dividing walls had retractable doorways allowing access into each chamber. The wire cup used to contain the stranger mice was made of cylindrical chrome bars spaced 1 cm apart (10 cm H; bottom diameter: 10 cm). Test mice were confined in the center chamber at the beginning of each phase. Exploration of an enclosed mouse or a wire cup was defined as when a test mouse oriented toward the cup with the distance between the nose and the cup less than 1 cm, or as climbing on the cup. The time spent in each chamber and time spent exploring enclosed novel mice or empty cups (novel objects) were recorded from a camera mounted overhead, and analyzed by Ethovision.

Open field test (OFT). OFTs were performed in a white acrylic glass box (45 × 45 × 40 cm) with an overhead lamp directed to the centre of the field, providing 120 lux of illumination on the floor of the arena. Each mouse was placed in the corner of the apparatus and locomotion parameters were recorded for 20 min.

Elevated plus maze (EPM). The EPM test was performed using a polyvinyl chloride maze comprising a central part (5x5 cm), two opposing open arms (32.5x5 cm), and two opposing closed arms (32.5x5x32.5 cm). The apparatus was set to a height of 50 cm, and the open arms were provided with uniform overhead illumination of 6 lux. Mice were placed in the open arm point close to the centre facing the closed arms, and video recordings were performed for a total duration of 20 min.

Chronic administration of maternal hormones in the PMv. For localized chronic administration of maternal hormones in the PMv, we performed cannula implantations using the Paxinos and Franklin atlas (*102*) coordinates (Bregma: -2.45 mm, midline ±0.6 mm, dorsal surface -5.5 mm). Cannulae (0008663 Brain Infusion Kit 2) and osmotic minipumps (0000298 ALZET Model 2004) were purchased from ALZET, and prepared according to protocol.

Brain slice electrophysiology. Acute slices of the mediobasal hypothalamus were prepared from adult DAT-tdTomato mice (own breeding). Slices were cut on a vibratome (Leica VT1000S) to 250 μm thickness and continuously perfused with oxygenated aCSF containing (in millimolar): NaCl (127), KCl (2.0), NaH_2PO_4 (1.2), $NaHCO_3$ (26), $MgCl_2$ (1.3), $CaCl_2$ (2.4), and D-glucose (10), at room temperature during recording. Each slice was exposed only to a single bath application of pharmacological compounds and was used for a single experiment. Whole-cell current- and voltage-clamp recordings were performed with micropipettes filled with intracellular solution containing (in millimolar), K-gluconate (140), KCl (10), HEPES (10), EGTA (10), and Na_2ATP (2) (pH 7.3 with KOH). Recordings were performed using a Multiclamp 700B amplifier, a DigiData 1440 digitizer, and pClamp 10.2 software (Molecular Devices, San Jose, CA, USA). Slow and fast capacitative components were semi-automatically compensated. Access resistance was monitored throughout the experiments, and neurons in which the series resistance exceeded 15 MΩ or changed ≥20% were excluded from the statistics. Liquid junction potential was 16.4 mV and not compensated. The recorded current was sampled at 20 kHz.

Reagents used in slice electrophysiology experiments; Neurobiotin™ tracer (Vector laboratories) was used in combination with Streptavidin, DyLight™ 405 conjugated (21831 Thermoscientific) or Avidin-FITC (43-4411 Invitrogen). TTX was purchased from Alomone Labs. Oxytocin nonapeptide, the OT-receptor (OT-R) agonist, (Thr4, Gly7)-oxytocin (TGOT), and the oxytocin receptor antagonist, vasotocin [d(CH2)$_5$^{1}, Tyr(Me)^2, Thr^4, Orn^8, des-Gly NH$_2$^{9}] (OR-A), were purchased from Bachem. Prolactin CYT-321 was purchased from PROSPEC, and SR 95531 (Gabazine) and ML-218

638 hydrochloride were purchased from Tocris. Matlab and OriginPro8 were used for electrophysiological
639 data analysis.
640

Immunofluorescence. Mice were anaesthetised with sodium pentobarbital (200 mg/kg, i.p.,
Sanofi-Aventis, France), then transcardially perfused with 10 mL Ca^{2+}-free Tyrode's solution (37°C)
containing 0.2% heparin, followed by 10 mL fixative (4% paraformaldehyde and 0.4% picric acid in
0.16 M phosphate buffer (PBS), 37°C), then 50 mL ice cold fixative. Whole brains were dissected,
immersed in ice cold fixative for 90 min then stored in 0.1M PBS (pH 7.4) containing 20% sucrose,
0.02% bacitracin and 0.01% sodium azide for three days, before freezing with CO_2. Coronal sections
were cut at a thickness of 14 μm on a cryostat (Microm, Walldorf) and thaw-mounted onto gelatin-
coated glass slides. For indirect immunofluorescence staining (performed at room temperature unless
otherwise specified), air-dried sections were washed in 0.01 M PBS for 30 min before incubation with
primary antisera diluted in PBS containing 0.3% Triton X-100 and 1% BSA for 16 hours at 4°C. The
slides were then washed for 30 min in PBS followed by 2h incubation with Alexa-488-conjugated
donkey anti-rabbit secondary antisera (1:500; Invitrogen). Slides went through a final wash for 30 min
in PBS and mounted with glycerol containing (2.5% DABCO; Sigma, St Louis). This method was used
with the following antibodies: NeuN was detected with primary antibody rabbit anti-NeuN (1:500; Cell
Signalling, D4G40), eYFP was detected with chicken anti-GFP (1:500; Aves Labs, GFP-1020).
Immunohistochemistry for pSTAT5: Prior to immunofluorescence staining, antigen retrieval was
performed by incubating sections for 15 min in citric acid (pH 7.4) at 80°C, then cooled at room
temperature for a further 30 min. After a 1% H_2O_2 Tris-buffered wash, sections were incubated in
rabbit pSTAT5 primary antibody (pSTAT5 Tyr 694, Cat#: C11C5, 1:500; Cell Signaling Technology)
for 72 hours at 4°C. Primary antibody incubation was followed by Alexa-488-conjugated donkey anti-
rabbit secondary antisera (1:500; Invitrogen).

To perform c-fos immunostaining, mice were deeply anesthetized with sodium pentobarbital (as
described above) and perfused transcardially with 4% (weight/vol) ice-cold paraformaldehyde in 0.1M
PBS. Brains were post-fixed in the same solution and 40 μm-thick coronal slices were cut at the
vibratome (Leica, Germany). Two PMv coronal sections per animal were selected to perform c-fos
immunostaining as follows. Sections were washed in TBS (100 mM Tris-Cl, 150 mM NaCl, pH 7.5),
incubated 1h at 25°C in 1% BSA-0.3% Triton X-100-TBS solution and then kept at 4°C in rabbit anti-
c-fos antibody (sc-52 LotG1108, Santa Cruz Biotechnology) solution (1:200 in 1% BSA-TBS). After
TBS washing, sections were incubated for 1h at 25°C in Alexa Fluor® 647 (Invitrogen) goat anti-rabbit
secondary antibody (1:500 in 1% BSA-TBS).
671

Confocal microscopy and cell-counting. All brain slices were imaged by epifluorescence
microscopy (ZEISS Imager M1) or confocal microscopy (Zeiss, LSM 800) for subsequent analysis.
Brain areas were determined according to their anatomy using Paxinos and Franklin Brain Atlas(*102*).

For PMv^{DAT} cell counts the entire PMv was cut, stained and counted. Quantification of c-fos staining
was obtained by averaging the number of positive cells of right and left PMv in two brain sections (-
2.46 and -2.54 from Bregma) (*102*). All counts were done manually using Image J software and blind
to test conditions.

Randomization and blinding. Behavioral data collection and analysis was performed blind to
experimental conditions. Anatomy data analysis but not tissue collection was blinded.
Electrophysiological data sampling and analysis was not blinded, with the exception of all whole-cell
patch clamp datasets presented in Figure 1. Mice were first screened for expressing maternal
aggression and then further assigned to groups for behavioral experiments.

Statistical analysis. No statistical methods were used to pre-determine sample sizes but our
sample sizes are similar to those reported in previous publications (*20, 104, 105*). Data met the
assumptions of the statistical tests used, and were tested for normality and equal variance. All *t*-tests
and one-way ANOVAs were performed using Graph Pad Prism software (Graphpad Software Inc.).
The Tukey and Bonferroni *posthoc* tests were used as appropriate for one-way ANOVAs. Normality
was determined by D'Agostino–Pearson normality test. Statistical significance was set at $P < 0.05$.

Figure 1. PMvDAT neurons activate in lactating dams following an aggressive encounter and are electrically hyperexcitable during nursing state.

(A) Schematic representation of the experimental design used to identify lactating dams expressing maternal aggression (n = 126 lactating mice). Lactating dams/mice were between five and eight months of age.

(B) Quantification of maternal aggression traits identify increased attack duration (left; shown as the percentage of total session time) and decreased attack latency (right) with consecutive trials in the RI paradigm shown in Figure 1A (n = 15 mice per group, RM one-way ANOVA, p = 0.0010).

(C) Schematic representation of the experimental design used to perform c-Fos studies following RI testing.

(D) Confocal images of c-Fos immunoreactivity in PMv, in the absence (top panels) or presence (bottom panels) of a juvenile female conspecific post-RI testing (n = 12 sections from 6 mice without intruder exposure, and n = 10 sections from 5 mice with intruder exposure; comparisons performed using two-tailed unpaired *t*-test, p < 0.0001).

(E) Identification of PMvDAT neurons using the DAT-tdTomato mouse line to perform whole-cell patch clamp slice recordings from age-matched adult virgin female and lactating mice (five to eight months old).

(F) PMvDAT neurons were identified and reconstructed post the recording session using Neurobiotin-FITC.

(G) Example recordings from PMvDAT neurons (left) and group data in pie charts (right) from virgin female (blue) and lactating (purple) mice, exhibiting low *vs* high baseline spike discharge, respectively (n = 53 and 21 recorded cells from 26 virgin female and 11 lactating dams, respectively).

(H) Quantification of firing frequency in PMvDAT cells in virgin females *vs* lactating dams (n = 21-53 cells, two-tailed unpaired t-test, p < 0.0001).

717   (I) Quantification of membrane potential in PMv^{DAT} cells in virgin females *vs* lactating dams (n = 21-
718   53 cells, two-tailed unpaired t-test, p < 0.0001).

719   See also Table S1.

Figure 2. Manipulation of the PMv^{DAT} neuron activity allows bidirectional control of maternal aggression.

(A) Schematic of the experimental design used to perform ChR2 PMv^{DAT} neuron photoactivation in the resident lactating dam during resident-intruder (RI) testing. Adult male and female intruders were used to decrease the levels of maternal aggression expressed in baseline conditions (*38, 45*).

(B) Bilateral Cre-dependent ChR2 transfection and fiber implants in the PMv of lactating dams (left) and confocal image of a sample section counterstained with the pan-neuronal marker NeuN, used to validate ChR2 expression and stereotactic implantation coordinates (right).

(C) Sample behavior raster plots at baseline (top) and during ChR2 stimulation (bottom) in a lactating dam during a RI test.

(D) Attack duration with and without photostimulation in lactating dams injected with eYFP or ChR2, during the RI test against adult male or adult female intruders (n = 5-9 mice per group, ordinary one-way ANOVA, p < 0.0001).

(E) Attack latency with and without photostimulation in lactating dams injected with eYFP or ChR2, during the RI test against adult male or adult female intruders (n = 5-9 mice per group, ordinary one-way ANOVA, p = 0.0027).

(F) Schematic of the experimental design used to perform eNpHR3 PMv^{DAT} neuron photoinhibition studies in the resident lactating dam during RI testing.

(G) Bilateral Cre-dependent eNpHR3 transfection and fiber implants in the PMv of lactating dams (left) and confocal image of a sample section counterstained with the pan-neuronal marker NeuN, used to validate eNpHR3 expression and stereotactic implantation coordinates (right).

(H) Sample behavior raster plots during baseline and eNpHR3 mediated photoinhibition in a lactating dam during a RI test.

(I) Quantification of aggression parameters (n = 8 mice per group, two-tailed paired t-test).

(J) Quantification of the latency to an attack bout following a photoinhibition episode (n = 8 mice per group, two-tailed paired t-test, p = 0.0131).

(K) Schematic of the experimental design used to perform RI tests in lactating dams with complete or partial lesion of the PMv^{DAT} cells.

(L) Bilateral Cre-dependent taCasp3 transfection in the PMv of lactating dams (left) and confocal image of an eYFP (middle) *vs* taCasp3 (right) sample section, used to validate the extent of the PMv^{DAT} cell lession.

(M) Sample behavior raster plots in control *vs* PMv^{DAT} neuron lesioned lactating dams during a RI test.

(N) Quantification of aggression parameters (n = 34 trials per group, 12 mice per group, two-tailed unpaired t-test). All but 2 mice were tested three times in the RI test. Two mice, one from each group, were tested two times in the RI test.

(O) Quantification of PMv^{DAT} neurons in eYFP and taCasp3 injected mice (n = 24 sections per group, duplicates from 12 mice per group, two-tailed unpaired t-test, p < 0.0001).

See also Table S1, and Figure S1.

Figure 3. Maternal hormones activate PMv^{DAT} neurons.

(A) Serum prolactin levels in virgin females across the estrous cycle (proestrus, estrus, met-/diestrus) *vs* lactating dams (n = 4-9 mice per group, ordinary one-way ANOVA, p = 0.0007).

(B) Confocal images of pSTAT5 immunoreactivity in PMv, in virgin females in estrus *vs* lactating dams (left). Overlap of pSTAT5 and DAT-tdTomato fluorescence along the Y axis, with 0 being the most ventral point of the confocal image and 1 the most dorsal (right).

(C) Quantification of pSTAT5 immunoreactive cells in the PMv of virgin females in estrus *vs* lactating dams (n = 10 sections per group, 5 mice per group, 2 sections per mouse, two-tailed unpaired t-test, p = 0.0011).

(D) Quantification of DAT-tdTomato colocalizing pSTAT5 immunoreactive cells in the PMv of virgin

females in estrus *vs* lactating dams (n = 10 sections per group, 5 mice per group, 2 sections per

mouse, two-tailed unpaired t-test, p = 0.0015).

(E) Prolactin leads to a reversible depolarization of PMv^{DAT} neurons (n = 7 cells from 5 mice, RM one-

way ANOVA, p = 0.0016).

(F) Prolactin induced changes in the action potential waveform of PMv^{DAT} neurons (left). Quantification

of action potential (AP) half-width and firing threshold in control (black) *vs* prolactin (orange, n = 15

action potentials in control *vs* bath application of prolactin per cell, 7 cells from 5 mice, two-tailed

paired t-test).

(G) (left) Comparison of post-inhibitory rebound bouts in control (TTX) *vs* prolactin application (TTX +

prolactin). Black traces represent the hyperpolarization induced by the highest absolute amplitude

negative current pulse, with the effect of a smaller amplitude pulse illustrated in dark gray and the

absence of any pulse in light gray. (middle) Quantification of the 1st post-inhibitory rebound in control

(black) *vs* prolactin (orange, n = 11 cells per group collected from 6 mice, two-tailed paired t-test, p =

0.0098). (right) Quantification of a 2st post-inhibitory rebound probability in control (black) *vs* prolactin

(orange, n = 11 cells per group collected from 6 mice, two-tailed paired t-test, p = 0.0061).

(H) Identification of a prolactin induced postsynaptic depolarization on PMv^{DAT} cells in current clamp

(left, n=13 cells per group collected from 8 mice, two-tailed paired t-test, p < 0.0001) and voltage

clamp (middle, n = 5 cells per group collected from 5 mice, two-tailed paired t-test, p = 0.0078).

Characterization of the prolactin induced current across the membrane voltage spectrum in PMv^{DAT}

neurons, with a low- and high-voltage component (right, n = 7 cells collected from 4 mice).

(I) Application of the selective t-type Ca^{2+} current, blocker ML218, blocks the low-voltage component

of the prolactin induced current (Prl$_I$). Application of prolactin does not induce an inward current in

PMv^{DAT} neurons pre-exposed to ML218 (left, n = 6 cells per group collected from 4 mice, two-tailed

paired t-test, p = 0.5129). In the presence of ML218 the prolactin induced current is composed only

of a high-voltage component (middle, n = 6 cells collected from 4 mice). Comparison of the low- (right,

n=6-7 cells per group, two-tailed unpaired t-test, p = 0.0042) and high-voltage (right, n=6-7 cells per

group, two-tailed unpaired t-test, p = 0.1384) component of the prolactin induced current in the

absence *vs* presence of ML218.

(J) (top) Electrophysiology recordings showing that the T-type Ca^{2+} channel blocker ML218, blocks

the effect of prolactin at potentiating rebound excitation bouts. (bottom left) Quantification of the

presence of post-inhibitory rebound in the three conditions (n = 7 cells per group collected from 3

mice, ordinary one-way ANOVA, p < 0.0001). (bottom right) Quantification of post-inhibitory rebound

duration in the three conditions (n = 7 cells per group collected from 3 mice, ordinary one-way ANOVA,

p < 0.0001).

(K) Oxytocin-immunoreactive fibers forming close appositions to PMv^{DAT} neurons (n = 5 mice).

(L) Oxytocin leads to a reversible depolarization of PMv^{DAT} neurons (n = 7 cells per group, collected

from 5 mice, RM one-way ANOVA, p < 0.0001).

(M) Application of the selective oxytocin receptor agonist, TGOT, leads to depolarization of PMv^{DAT}

neurons (top, n = 5 cells per group, collected from 5 mice, two-tailed paired t-test, p = 0.0274).

Application of oxytocin under oxytocin receptor blockade, leads to depolarization of PMv^{DAT} neurons

(bottom, n = 5 cells per group, collected from 5 mice, two-tailed paired t-test, p = 0.0008).

(N) Identification of an oxytocin induced postsynaptic depolarization on PMv^{DAT} cells in current clamp

(left, n=8 cells per group collected from 4 mice, two-tailed paired t-test, p < 0.0001). Characterization

of the oxytocin induced current across the membrane voltage spectrum in PMv^{DAT} neurons, with a

low- and high-voltage component (right, n = 8 cells collected from 4 mice).

See also Table S1.

Figure 4. Activation of PMv^DAT neurons suppresses non-aggression-related behavioral programs.

(A) Schematic of the experimental design used to perform ChR2 PMv^{DAT} neuron photoactivation during the pup retrieval test.

(B) Images from two photostimulation trials at the start (0 s) and 50 s *post*-trial initiation, with lactating dams expressing eYFP (left) and ChR2 (right) in PMv^{DAT} neurons.

(C) Sample behavior raster plots of two sequential trials without and with photostimulation in a ChR2-expressing lactating dam during the pup retrieval test.

(D) Pup retrieval duration with and without photostimulation in lactating dams injected with eYFP or ChR2, during the pup retrieval test (n = 5 mice per group, ordinary one-way ANOVA, p < 0.0001).

(E) Percentage of pups retrieved with and without photostimulation in lactating dams injected with eYFP or ChR2, during the pup retrieval test (n = 5 mice per group, ordinary one-way ANOVA, p < 0.0001).

See also Table S1.

Figure S1. Injection of AAV-flex-taCasp3 in the PMv of DAT-Cre animals does not extend across the rostrocaudal axis, leaving the midbrain and arcuate nucleus dopamine neurons unaffected, Related to Figure 2.

(A) Confocal image of an eYFP- (left) *vs* taCasp3- (middle) PMv injected sample sections of midbrain (substantia nigra, SN, and ventral tegmental area, VTA) DAT-tdTomato neurons, and quantification of cell number per section (n = 24-26 sections per group, duplicates from 12-13 mice per group, two-tailed unpaired t-test, p = 0.1141).

(B) Confocal image of an eYFP- (left) *vs* taCasp3- (middle) PMv injected sample sections of arcuate DAT-tdTomato neurons, and quantification of cell number per section (n = 24-26 sections per group, duplicates from 12-13 mice per group, two-tailed unpaired t-test, p = 0.5650). 3V, third ventricle.

See also Table S1.

Figure S2. Chronic administration of maternal hormones in PMv is not sufficient to evoke aggression in virgin females, Related to Figure 3.

(A) Schematic of the experimental design followed to infuse maternal hormones into the PMv parenchyma via cannulas for a period of 28 days, and perform a panel of behavioral tests.

(B) Quantification of social behaviors in the RI test, with implanted resident adult virgin females *vs* juvenile virgin female intruders. No changes in close investigation or mounting duration were recorded (n = 21-24 trials per group, triplicates from 7-8 mice per group, RM one-way ANOVA). No episodes of aggression occurred in any of the groups.

(C) Design and results from the sociability test, where implanted adult virgin female mice can choose between an empty chamber and one with a juvenile virgin female conspecific present. (n = 13-24 trials per group, duplicates or triplicates from 7-8 mice per group, ordinary one-way ANOVA).

(D) Design and results from the pup retrieval test, with implanted adult virgin female mice (n = 21-24 trials per group, triplicates from 7-8 mice per group, ordinary one-way ANOVA).

(E) Results from the open field test, with implanted adult virgin female mice (n = 21-24 trials per group, triplicates from 7-8 mice per group, ordinary one-way ANOVA).

(F) Results from the elevated plus maze test, with implanted adult virgin female mice (n = 21-24 trials per group, triplicates from 7-8 mice per group, ordinary one-way ANOVA). See also Table S1.

Table S1

Figures	Test	n (per group)	KW statistic or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure 1B, left – attack duration (%)	RM one-way ANOVA with Kruskal-Wallis post-hoc test	12	KW statistic = 18.18	11	0.0001	day 1 *vs.* day 2	0.0051	**
						day 1 *vs.* day 3	0.0001	***
						day 2 *vs.* day 3	>0.9999	ns
Figure 1B, right – attack latency (s)	RM one-way ANOVA with Kruskal-Wallis post-hoc test	12	KW statistic = 7.182	11	0.0276	day 1 *vs.* day 2	0.2988	ns
						day 1 *vs.* day 3	0.0238	*
						day 2 *vs.* day 3	0.9411	ns
Figure 1D, c-Fos cells (#)	Two-tailed unpaired t-test	10-12	t = 9.661	20	< 0.0001	c-Fos in PMv without *vs.* with intruder	N/A	****
Figure 1D, DAT/c-Fos cells (#)			t = 11.72			DAT/c-Fos colocalization without *vs.* with intruder	N/A	****
Figure 1H, firing frequency (Hz)	Two-tailed unpaired t-test	21-53	t = 4.540	72	< 0.0001	virgin *vs.* lactating	N/A	****

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure 2D, attack duration (s)	Ordinary one-way ANOVA with uncorrected Fisher's LSD test as multiple comparisons test	5-9	$F_{(7, 48)} = 11.55$	7	< 0.0001	AAV-eYFP dam, against adult virgin male intruder – baseline vs. photostimulation	> 0.9999	ns
						AAV-eYFP dam, against adult virgin female intruder – baseline vs. photostimulation	> 0.9999	ns
						AAV-ChR2 dam, against adult virgin male intruder – baseline vs. photostimulation	> 0.9999	ns
						AAV-ChR2 dam, against adult virgin female intruder – baseline vs. photostimulation	< 0.0001	****
Figure 2E, attack latency (s)	Ordinary one-way ANOVA with uncorrected Fisher's LSD test as multiple comparisons test	5-9	$F_{(7, 48)} = 3.732$	7	0.0027	AAV-eYFP dam, against adult virgin male intruder – baseline vs. photostimulation	0.7149	ns
						AAV-eYFP dam, against adult virgin female intruder – baseline vs. photostimulation	0.9468	ns
						AAV-ChR2 dam, against adult virgin male intruder – baseline vs. photostimulation	0.1043	ns
						AAV-ChR2 dam, against adult virgin female intruder – baseline vs. photostimulation	0.0266	*

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure 2I, left attack duration/trial (s)	Two-tailed paired t-test	8	t = 9.280	7	< 0.0001	AAV-ChR2 dam, against a juvenile virgin female intruder – baseline *vs.* photostimulation	N/A	****
Figure 2I, middle attack bout (s)	Two-tailed paired t-test	8	t = 5.356	7	0.0011	AAV-ChR2 dam, against a juvenile virgin female intruder – baseline *vs.* photostimulation	N/A	**
Figure 2I, right attack frequency/trial (#)	Two-tailed paired t-test	8	t = 3.752	7	0.0071	AAV-ChR2 dam, against a juvenile virgin female intruder – baseline *vs.* photostimulation	N/A	**
Figure 2J, latency to first attack following a photoinhibition episode (s)	Two-tailed paired t-test	8	t = 3.300	7	0.0131	AAV-ChR2 dam, against a juvenile virgin female intruder – baseline *vs.* photostimulation	N/A	*
Figure 2N, left attack duration (%)	Two-tailed unpaired t-test	34	t = 8.827	54	< 0.0001	against a juvenile virgin female intruder – eYFP *vs.* taCasp3 injected lactating dams	N/A	****
Figure 2N, middle attack frequency (#)	Two-tailed unpaired t-test	34	t = 5.348	66	< 0.0001	against a juvenile virgin female intruder – eYFP *vs.* taCasp3 injected lactating dams	N/A	****
Figure 2N, right latency to first attack (min)	Two-tailed unpaired t-test	34	t = 6.506	41	< 0.0001	against a juvenile virgin female intruder – eYFP *vs.* taCasp3 injected lactating dams	N/A	****
Figure 2O, PMv^{DAT} neurons (#)	Two-tailed unpaired t-test	24	t = 22.78	48	< 0.0001	eYFP *vs.* taCasp3 injected lactating dams	N/A	****

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure 3A, Serum Prolactin (ng/mL)	Ordinary one-way ANOVA with Dunnett's multiple comparisons test	4-9	$F_{(3, 19)} = 8.797$	3	0.0007	proestrus *vs.* lactating	0.0022	**
						estrus *vs.* lactating	0.0199	*
						met-/diestrus *vs.* lactating	0.0011	**
Figure 3C, pSTAT5 cells in PMv (#)	Two-tailed unpaired t-test	10	t = 3.878	18	0.0011	Virgin in estrus *vs.* lactating	N/A	**
Figure 3D, PMv^{DAT}/pSTAT5 cell colocalization (#)	Two-tailed unpaired t-test	10	t = 3.737	18	0.0015	Virgin in estrus *vs.* lactating	N/A	**
Figure 3E, firing frequency (Hz)	RM one-way ANOVA with Bonferroni's multiple comparisons test	7	$F_{(1.723, 10.34)} = 13.73$	6	0.0016	control *vs.* prolactin	0.0069	**
						control *vs.* wash	> 0.9999	ns
						prolactin *vs.* wash	0.0097	**
Figure 3F, action potential half width (ms)	Two-tailed paired t-test	15	t = 8.847	14	< 0.0001	control *vs.* prolactin	N/A	****
Figure 3F, action potential firing threshold (mV)	Two-tailed paired t-test	15	t = 1.926	14	0.0746	control *vs.* prolactin	N/A	ns
Figure 3G, duration of 1st rebound depolarization (s)	Two-tailed paired t-test	11	t = 3.183	10	0.0098	control *vs.* prolactin	N/A	**
Figure 3G, 2nd rebound depolarization probability	Two-tailed paired t-test	11	t = 3.464	10	0.0061	control *vs.* prolactin	N/A	**
Figure 3H, left membrane potential (mV)	Two-tailed paired t-test	13	t = 9.753	12	< 0.0001	control *vs.* prolactin	N/A	****
Figure 3H, middle Holding I (pA)	Two-tailed paired t-test	5	t = 4.941	4	0.0078	control *vs.* prolactin	N/A	**
Figure 3I, left Holding I (pA)	Two-tailed paired t-test	6	t = 0.7040	5	0.5129	control *vs.* prolactin	N/A	ns

Figure 3I, top right Prl$_I$ LV(pA)	Two-tailed unpaired t-test	5-7	t = 3.685	10	0.0042	control *vs.* prolactin	N/A	**
Figure 3I, bottom right Prl$_I$ LV(pA)	Two-tailed unpaired t-test	5-7	t = 1.611	10	0.1384	control *vs.* prolactin	N/A	ns
Figure 3J, bottom left rebound presence	Ordinary one-way ANOVA with Dunnett's multiple comparisons test	7	t = 3.429	18	< 0.0001	TTX *vs.* ML218	N/A	****
Figure 3J, bottom left rebound presence	Ordinary one-way ANOVA with Dunnett's multiple comparisons test	7	t = 3.429	18	< 0.0001	ML218 *vs.* prolactin	N/A	ns
Figure 3J, bottom left rebound duration	Ordinary one-way ANOVA with Dunnett's multiple comparisons test	7	t = 5.135	18	< 0.0001	TTX *vs.* ML218	N/A	****
Figure 3J, bottom left rebound duration	Ordinary one-way ANOVA with Dunnett's multiple comparisons test	7	t = 5.135	18	< 0.0001	ML218 *vs.* prolactin	N/A	ns
Figure 3J, bottom right sEPSC amplitude (pA)	Two-tailed paired t-test	8	t = 4.069	7	0.0048	control *vs.* prolactin	N/A	**
Figure 3L, firing frequency (Hz)	RM one-way ANOVA with Bonferroni's multiple comparisons test	7	F $_{(1.114, 6.682)}$ = 75.40	6	< 0.0001	control *vs.* oxytocin	< 0.0001	****
						control *vs.* wash	> 0.9999	ns
						oxytocin *vs.* wash	< 0.0001	****
Figure 3M, top firing frequency (Hz)	Two-tailed paired t-test	5	t = 3.393	4	0.0274	control *vs.* TGOT	N/A	*
Figure 3M, bottom membrane pot. (mV)	Two-tailed paired t-test	5	t = 9.253	4	0.0008	control *vs.* TGOT	N/A	***
Figure 3N, left membrane pot. (mV)	Two-tailed paired t-test	8	t = 8.040	7	< 0.0001	control *vs.* TGOT	N/A	****

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure 4D, pup retrieval duration (s)	Ordinary one-way ANOVA with Bonferroni's multiple comparisons test	5	$F_{(3, 16)} = 25.59$	3	< 0.0001	AAV-eYFP dam – baseline *vs.* photostimulation	> 0.9999	ns
						AAV-ChR2 dam – baseline *vs.* photostimulation	< 0.0001	****
Figure 4E, pup retrieved (%)	Ordinary one-way ANOVA with Bonferroni's multiple comparisons test	5	$F_{(3, 16)} = 49.80$	3	< 0.0001	AAV-eYFP dam – baseline *vs.* photostimulation	> 0.9999	ns
						AAV-ChR2 dam – baseline *vs.* photostimulation	< 0.0001	****
Figure S1A, SN&VTA DAT neurons (#)	Two-tailed unpaired t-test	24-26	$t = 1.609$	48	0.1141	eYFP *vs.* taCasp3 injected lactating dams	N/A	ns
Figure S1B, arcuate DAT neurons (#)	Two-tailed unpaired t-test	24-26	$t = 0.5794$	48	0.5650	eYFP *vs.* taCasp3 injected lactating dams	N/A	ns
Figure S2B, close investigation (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 3.760$	3	0.0138	saline *vs.* prolactin (Prl)	0.8239	ns
						saline *vs.* oxytocin (OT)	> 0.9999	ns
						saline *vs.* Prl & OT	0.1259	ns
Figure S2B, mounting (s)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 0.9736$	3	0.4092	saline *vs.* prolactin (Prl)	> 0.9999	ns
						saline *vs.* oxytocin (OT)	> 0.9999	ns
						saline *vs.* Prl & OT	0.3291	ns
Figure S2C, time spent moving (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	13-24	$F_{(3, 75)} = 1.698$	3	0.1747	saline *vs.* prolactin (Prl)	> 0.9999	ns
						saline *vs.* oxytocin (OT)	> 0.9999	ns
						saline *vs.* Prl & OT	0.4069	ns
Figure S2C, time in social zone (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	13-24	$F_{(3, 75)} = 3.953$	3	0.0114	saline *vs.* prolactin (Prl)	> 0.9999	ns
						saline *vs.* oxytocin (OT)	0.2401	ns
						saline *vs.* Prl & OT	0.0346	*

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure S2D, latency to first pup retrieval (min)	RM one-way ANOVA with Bonferroni's multiple comparisons test	15-20	$F_{(3, 65)} = 2.150$	3	0.1025	saline vs. prolactin (Prl)	> 0.9999	ns
						saline vs. oxytocin (OT)	0.8639	ns
						saline vs. Prl & OT	0.0418	*
Figure S2D, pups retrieved (#)	RM one-way ANOVA with Bonferroni's multiple comparisons test	20-24	$F_{(3, 82)} = 3.452$	3	0.0202	saline vs. prolactin (Prl)	0.0124	*
						saline vs. oxytocin (OT)	0.8639	ns
						saline vs. Prl & OT	0.0432	*
Figure S2D, proximity to pups (min)	RM one-way ANOVA with Bonferroni's multiple comparisons test	20-24	$F_{(3, 82)} = 4.716$	3	0.0044	saline vs. prolactin (Prl)	0.0330	*
						saline vs. oxytocin (OT)	0.3891	ns
						saline vs. Prl & OT	0.0017	**
Figure S2D, nesting (min)	RM one-way ANOVA with Bonferroni's multiple comparisons test	20-24	$F_{(3, 82)} = 3.819$	3	0.0129	saline vs. prolactin (Prl)	0.2276	ns
						saline vs. oxytocin (OT)	0.0669	ns
						saline vs. Prl & OT	0.0051	**
Figure S2E, time spent moving (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 18.17$	3	< 0.0001	saline vs. prolactin (Prl)	0.0007	***
						saline vs. oxytocin (OT)	0.0009	***
						saline vs. Prl & OT	> 0.9999	ns
Figure S2E, time spent in the center (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 10.59$	3	< 0.0001	saline vs prolactin (Prl)	0.1397	ns
						saline vs. oxytocin (OT)	0.0025	**
						saline vs. Prl & OT	0.1915	ns
Figure S2F, time spent moving (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 7.991$	3	< 0.0001	saline vs. prolactin (Prl)	0.0697	ns
						saline vs. oxytocin (OT)	0.0578	ns
						saline vs. Prl & OT	0.4917	ns

Figures	Test	n (per group)	F or t values	df	p value	Comparisons	Adjusted p value	p value summary
Figure S2F, time spent in the open arms (%)	RM one-way ANOVA with Bonferroni's multiple comparisons test	21-24	$F_{(3, 83)} = 2.163$	3	0.0985	saline *vs.* prolactin (Prl)	> 0.9999	ns
						saline *vs.* oxytocin (OT)	0.1189	ns
						saline *vs.* Prl & OT	> 0.9999	ns