1

Predicting the Sequence-Dependent Backbone Dynamics of Intrinsically Disordered

Proteins

Short title: Predictor for IDP sequence-dependent dynamics

Sanbo Qin[1] and Huan-Xiang Zhou[1,2,*]

[1]Department of Chemistry and [2]Department of Physics, University of Illinois Chicago,

Chicago, IL 60607, USA

*Corresponding author. E-mail: hzhou43@uic.edu

Abstract

How the sequences of intrinsically disordered proteins (IDPs) code for functions is still an enigma. Dynamics, in particular residue-specific dynamics, holds crucial clues. Enormous efforts have been spent to characterize residue-specific dynamics of IDPs, mainly through NMR spin relaxation experiments. Here we present a sequence-based method, SeqDYN, for predicting residue-specific backbone dynamics of IDPs. SeqDYN employs a mathematical model with 21 parameters: one is a correlation length and 20 are the contributions of the amino acids to slow dynamics. Training on a set of 45 IDPs reveals aromatic, Arg, and long-branched aliphatic amino acids as the most active in slow dynamics whereas Gly and short polar amino acids as the least active. SeqDYN predictions not only provide an accurate and insightful characterization of sequence-dependent IDP dynamics but may also serve as indicators in a host of biophysical processes, including the propensities of IDP sequences to undergo phase separation.

Introduction

Intrinsically disordered proteins (IDPs) or regions (IDRs) do not have the luxury of a three-dimensional structure to help decipher the relationship between sequence and function. Instead, dynamics has emerged as a crucial link between sequence and function for IDPs (1). Nuclear magnetic resonance (NMR) spin relaxation is a uniquely powerful technique for characterizing IDP dynamics, capable of yielding residue-specific information (2). Backbone ^{15}N relaxation experiments typically yield three parameters per residue: transverse relaxation rate (R_2), longitudinal relaxation rate (R_1), and steady-state heteronuclear Overhauser enhancement (NOE). While all three parameters depend on ps-ns dynamics, R_2 is the one most affected by slower dynamics (10s of ns to 1 μs). An increase in either the timescale or the amplitude of slower dynamics results in higher R_2. For IDPs, R_2 is also the parameter that exhibits the strongest dependence on sequence (1, 2).

R_2 was noted early on as an important indicator of residual structure in the unfolded state of the structured protein lysozyme (3). This property has since been measured for many IDPs to provide insight into various biophysical processes. Just as the residual structure in the unfolded state biases the folding pathway of lysozyme (3), a nascent α-helix in the free state of Sendai virus nucleoprotein C-terminal domain (Sev-NT), as indicated by highly elevated R_2 (4), biases the coupled binding and folding pathway in the presence of its target phosphoprotein (5). Local secondary structure preformation also facilitates the binding of yes-associated protein (YAP) with its target transcription factor (6). Likewise a correlation has been found between R_2 in the free state and the membrane binding propensity of synaptobrevin-2: residues with elevated R_2 have increased propensity for membrane binding (7). R_2 in the free state has also been used to uncover factors that promote liquid-liquid phase separation of IDPs. For example, a nascent α-helix (shown by elevated R_2) is important for the phase separation of the TDP-43 low-complexity domain, as both the deletion of the helical region and a helix-breaking mutation (Ala to Pro) abrogates phase separation (8). Similarly, nascent α-helices in the free state of cytosolic abundant heat-soluble 8 (CAHS-8), upon

raising concentration and lowering temperature stabilize to form the core of fibrous gels (9).

For the hnRNPA1 low-complexity domain (A1-LCD), aromatic residues giving rise to local

peaks in R_2 also mediate phase separation (10).

Both NMR relaxation data and molecular dynamics (MD) simulations have revealed

determinants of R_2 for IDPs. It has been noted that the flexible Gly tends to lower R_2 whereas

secondary structure and contact formation tend to raise R_2 (11). This conclusion agrees well

with recent MD simulations (1, 12-14). These MD studies, using IDP-specific force fields, are

able to predict R_2 in quantitative agreement with NMR measurements, without ad hoc

reweighting as done in earlier studies. According to MD, most contact clusters are formed by

local sequences, within blocks of up to a dozen or so residues (1, 12, 14). Tertiary contacts

can also form but are relatively rare; as such their accurate capture requires extremely

extensive sampling and still poses a challenge for MD simulations. Contrary to Gly, aromatic

residues have been noted as mediators of contact clusters (3, 10).

Schwalbe et al. (15) introduced a mathematical model to describe the R_2 profile along

the sequence for lysozyme in the unfolded state. The R_2 value of a given residue was

expressed as the sum of contributions from this residue and its neighbors. This model yields a

mostly flat profile across the sequence, except for falloff at the termini, resulting in an overall

bell shape. Klein-Seetharaman et al. (3) then fit peaks above this flat profile as a sum of

Gaussians. Cho et al. (16) proposed bulkiness as a qualitative indicator of backbone

dynamics. Recently Sekiyama et al. (17) calculated R_2 as the geometric mean of "indices of

local dynamics"; the latter were parameterized by fitting to the measured R_2 for a single IDP.

All these models merely describe the R_2 profile of a given IDP, and none of them is

predictive.

Here we present a method, SeqDYN, for predicting R_2 of IDPs. Using a mathematical

model introduced by Li et al. (18) to predict propensities for binding nanoparticles and also

adapted for predicting propensities for binding membranes (19), we express the R_2 value of a

residue as the product of contributing factors from all residues. The contributing factor

attenuates as the neighboring residue becomes more distant from the central residue. The

model, after training on a set of 45 IDPs, has prediction accuracy that is competitive against

that of the recent MD simulations using IDP-specific force fields (1, 12-14). For lysozyme,

the SeqDYN prediction agrees remarkably well with R_2 measured in the unfolded state.

Results

The data set of IDPs with R_2 rates

We collected R_2 data for a total of 54 nonhomologous IDPs or IDRs (Table 1; Figure 1).

According to indicators from NMR properties, including low or negative NOEs, narrow

dispersion in backbone amide proton chemical shifts, and small secondary chemical shifts

(SCSs), most of the proteins are disordered with at most transient α-helices. A few are

partially folded, including Sev-NT with a well-populated (~80%) long helix (residues 478-

491) (20), CREB-binding protein fourth intrinsically disordered linker (CBP-ID4) with > 50%

propensities for two long helices (residues 2-25 and 101-128) (21), HOX transcription factor

DFD (HOX-DFD) with a well-folded domain comprising three helices (22), and Hahellin

(apo form) as a molten globule (23). In Figure 2, we display representative conformations of

five IDPs, ranging from fully disordered MAPK kinase 4 (MKK4) (24) and α-synuclein (25)

to Measles virus phosphoprotein N-terminal domain (Mev-P$_{NTD}$) (26) with transient short

helices to Sev-NT and CBP-ID4 with stable long helices. The sequences of all the IDPs are

listed in Table 2.

We used 45 of the 54 IDPs to train and validate SeqDYN and reserved the remaining 9

for testing. The sequence lengths of the training set range from 39 to 406 residues, with an

average of 125.3 residues. Altogether R_2 data are available for 3966 residues. A large

majority (35 out of 45) of the 45 IDPs have mean R_2 values ($\bar{R}_2$, calculated among all the

residues in a protein) between 2.5 and 5.5 s^{-1} (Table 1 and Figure 3A). This $\bar{R}_2$ range is much

lower than that of structured proteins with similar sequence lengths. The low $\bar{R}_2$ values and

lack of dependence on sequence length (Figure 3-figure supplement 1A) suggest that R_2 of

the IDPs is mostly dictated by local sequence instead of tertiary interaction.

The most often used temperature for acquiring the R_2 data was 298 K, but low

temperatures (277 to 280 K) were used in a few cases (Table 1 and Figure 3-figure

supplement 1B). Of the seven IDPs with $\bar{R}_2 > 6.4$ s^{-1}, four can be attributed to low

temperatures (27-30), one is due to a relatively low temperature (283 K) as well as the

presence of glycerol (20% v/v) (31), and two can be explained by tertiary structure formation

[a folded domain (22) or molten globule (23)]. A simple reason for higher R_2 values at lower

temperatures is the higher water viscosity, resulting in a slowdown in molecular tumbling; a

similar effect is achieved by adding glycerol. In some cases, R_2 was measured at both low and

room temperatures (4, 10). To a good approximation, the effect of lowering temperature is a

uniform scaling of R_2 across the IDP sequence. For Sev-NT, downscaling of the R_2 values at

278 K by a factor of 2.0 brings them into close agreement with those at 298 K (Figure 3-

figure supplement 1C), with a root-mean-square-deviation (RMSD) of 0.5 s^{-1} among all the

residues. Likewise, for A1-LCD, downscaling by a factor of 2.4 brings the R_2 values at 288 K

into good match with those at 298 K (Figure 3-figure supplement 1D), with an RMSD of 0.4

s^{-1}. Because SeqDYN is concerned with the sequence dependence of R_2, a uniform scaling

has no effect on model parameter or prediction; therefore mixing the data from different

temperatures is justified. The same can be said about the different magnetic fields in acquiring

the R_2 data (Table 1 and Figure 3-figure supplement 1E). Increasing the magnetic field raises

R_2 values, and the effect is also approximated well by a uniform scaling (4, 8, 29).

One measure on the level of sequence dependence of R_2 is the standard deviation, σ_{R_2},

calculated among the residues of an IDP. Among the training set, the R_2 values of 30 IDPs

have moderate sequence variations, with σ_{R_2} ranging from 0.5 to 1.5 s^{-1} (Table 1); the

histogram of σ_{R_2} calculated for the entire training set peaks around 0.75 s^{-1} (Figure 3A).

There is a moderate correlation between σ_{R_2} and $\bar{R}_2$ (Figure 3A, inset), reflecting in part the

fact that σ_{R_2} can be raised simply by a uniform upscaling, e.g., as a result of lowering

temperature. Still, only two of the five IDPs with high $\bar{R}_2$ attributable to lower temperature or

presence of glycerol are among the seven IDPs with high sequence variations ($\sigma_{R_2} > 2$ s^{-1}).

Therefore the sequence variation of R_2 as captured by σ_{R_2} manifests mostly the intrinsic effect

of the IDP sequence, not the influence of external factors such as temperature or magnetic

field strength. The mean σ_{R_2} value among the training set is 1.24 s^{-1}.

One way to eliminate the influence of external factors is to scale the R_2 values of each

IDP by its $\bar{R}_2$; we refer to the results as scaled R_2, or sR_2. We then pooled the sR_2 values for

all residues in the training set, and separated them according to amino-acid types. The amino

acid type-specific mean sR_2 values, or msR_2, are displayed in Figure 3B. The seven amino

acids with the highest msR_2 in descending order are Trp, Arg, Tyr, Phe, Ile, His, and Leu.

The presence of all the four aromatic amino acids in this "high-end" group immediately

suggests π-π stacking as important for raising R_2; the presence of Arg further implicates

cation-π interactions. In the other extreme, the seven amino acids with the lowest msR_2 in

ascending order are Gly, Cys, Val, Asp, Ser, Thr, and Asn. Gly is well-known as a flexible

residue; it is also interesting that all the four amino acids with short polar sidechains are found

in this "low-end" group. Pro has an excessively low msR_2 [with data from only two IDPs (32,

33)], but that is due to the absence of an amide proton.

The SeqDYN model and parameters

The null model is to assume a uniform R_2 for all the residues in an IDP. The root-mean-

square-error (RMSE) of the null model is equal to the standard deviation, σ_{R_2}, of the

measured R_2 values. The mean RMSE, $\overline{\text{RMSE}}$, of the null model, equal to 1.24 s^{-1} for the

training set, serves as the upper bound for evaluating the errors of R_2 predictors. The next

improvement is a one-residue predictor, where first each residue (with index n) assumes its

amino acid-specific mean sR_2 (msR_2) and then a uniform scaling factor Υ is applied:

$$R_2(n) = \Upsilon \cdot \text{ms}R_2(n) \tag{1}$$

163   This one-residue model does only minutely better than the null model, with a $\overline{\mathrm{RMSE}}$ of 1.22 s⁻

164   ¹.

165   In SeqDYN, we account for the influence of neighboring residues. Specifically, each

166   residue i contributes a factor $f(i; n)$ to the R_2 value of residue n. Therefore

$$R_2(n) = \Upsilon \prod_{i=1}^{N} f(i; n) \tag{2a}$$

167   where N is the total number of residues in the IDP. The contributing factor depends on the

168   sequence distance $s = |i - n|$ and the amino-acid type of residue i:

$$f(i; n) = 1 + \frac{q(i) - 1}{1 + bs^2} \tag{2b}$$

There are 21 global parameters. The first 20 are the q values, one for each of the 20 types of

amino acids; the last parameter is b, appearing in the Lorentzian form of the sequence-

distance dependence. We define the correlation length, L_{corr}, as the sequence distance at

which the contributing factor is midway between the values at $s = 0$ and ∞. It is easy to

verify that $L_{\mathrm{corr}} = b^{-1/2}$. Note that the single-residue model can be seen as a special case of

SeqDYN, with L_{corr} set to 0 and q set to msR_2.

The functional forms of Eqs (2a,b) were adapted from Li et al. (18); we also used them

for predicting residue-specific membrane association propensities of IDPs (19). In these

previous applications, a linear term was also present in the denominator of Eq (2b). In our

initial training of SeqDYN, the coefficient of the linear term always converged to near zero.

We thus eliminated the linear term. In addition to the Lorentzian form, we also tested a

Gaussian form for the sequence-distance dependence and found somewhat worse

performance. The more gradual attenuation of the Lorentzian form with increasing sequence

distance evidently provides an overall better model for the R_2 data in the entire training set.

Others (16, 17, 24) have modeled R_2 as the average of some parameters over a window; a

window has an extremely abrupt sequence-distance dependence (1 for $s < L_{\mathrm{corr}}$ and 0 for

$s > L_{\mathrm{corr}}$).

We parametrized the SeqDYN model represented by Eqs (2a,b) on the training set of

45 IDPs. In addition to the 21 global parameters noted above, there are also 45 local

parameters, namely one uniform scaling factor (Υ) per IDP. The parameter values were

selected to minimize the sum of the mean-square-errors for the IDPs in the training set,

calculated on R_2 data for a total of 3924 residues. We excluded the 42 Pro residues in the

training set because, as already noted, their R_2 values are lower for chemical reasons. We will

present validation and test results below, but first let us look at the parameter values.

The q values are displayed in Figure 3B alongside msR_2. In descending order, the

seven amino acids with the highest q values are Trp, Ile, Tyr, Arg, His, Phe, and Leu. These

are exactly the same amino acids in the high-end group for msR_2, though their order there is

somewhat different. In ascending order, the seven amino acids (excluding Pro) with the

lowest q values are Gly, Asn, Ser, Asp, Val, Thr, and Cys. The composition of the low-end

group is also identical to that for msR_2. The q values thus also suggest that π-π and cation-π

interactions in local sequences may raise R_2, whereas Gly and short-polar residues may lower

R_2.

Given the common amino acids at both the high and low ends for msR_2 and q, it is not

surprising that these two properties exhibit a strong correlation, with a coefficient of

determination (R^2; excluding Pro) at 0.92 (Figure 4A). Also, because the high-end group

contains the largest amino acids (e.g., Trp and Tyr) whereas the low-end group contains the

smallest amino acids (e.g., Gly and Ser), we anticipated some correlation of msR_2 and q with

amino-acid size. We measure the latter property by the molecular mass (m). As shown in

Figure 4B, both msR_2 and q indeed show a medium correlation with m, with $R^2 = 0.67$

(excluding Pro) and 0.61, respectively. A bulkiness parameter was proposed as an indicator of

sequence-dependent backbone dynamics of IDPs (16, 24). Bulkiness was defined as the

sidechain volume-to-length ratio, and identified amino acids with aromatic or branched

aliphatic sidechains as bulky (34). We found only modest correlations between either msR_2 or

q and bulkiness, with R^2 just below 0.4 (Figure 4C).

The optimized value of b is 3.164×10^{-2}, corresponding to an L_{corr} of 5.6 residues.

The resulting optimized $\overline{RMSE}$ is 0.95 s^{-1}, a clear improvement over the value 1.24 s^{-1} of the

null model. To check the sensitivity of prediction accuracy to b, we set b to values

corresponding to $L_{corr} = 0, 1, 2, ..$, and retrained SeqDYN for b fixed at each value. Note that

the null-model $\overline{RMSE}$, 1.24 s^{-1}, sets an upper bound. This upper bound is gradually reached

when L_{corr} is increased from the optimal value. In the opposite direction, when L_{corr} is

decreased from the optimal value, $\overline{RMSE}$ rises quickly, reaching 1.22 s^{-1} at $L_{corr} = 0$. The

latter $\overline{RMSE}$ is the same as that of the single-residue model. Lastly we note that there is a

strong correlation between the uniform scaling factors and $\bar{R}_2$ values among the 45 IDPs ($R^2 =$

0.77), as to be expected. For 39 of the 45 IDPs, Y values fall in the range of 0.8 to 2.0 s^{-1}.

As presented next, we evaluate the performance of SeqDYN by leave-one-out cross

validation, where each IDP in turn was left out of the training set and the model was trained

on the remaining 44 IDPs to predict R_2 for the IDP that was left out. The parameters from the

leave-one-out (also known as jackknife) training sessions allow us to assess the potential bias

of the training set. For this purpose, we compare the values of the 21 global parameters, either

from the full training set or from taking the averages of the jackknife training sessions. For

each of the q parameters, the values from these two methods differ only in the fourth digit;

e.g., for Leu, they are both 1.1447 from full training and from jackknife training. The values

for b are 3.164×10^{-2} from full training as stated above and 3.163×10^{-2} from jackknife

training. The close agreement in parameter values between full training and jackknife training

suggests no significant bias in the training set.

Another question of interest is whether the difference between the q parameters of two

amino acids is statistically significant. To answer this question, we carried out five-fold cross-

validation training, resulting in five independent estimates for each parameter. For example,

the mean ± standard deviation of the q parameter is 1.1405 ± 0.0066 for Leu and $1.2174 \pm$

0.0211 for Ile. A t-test shows that their difference is extremely statistically significant ($P <$

0.0001). In contrast, the difference between Leu and Phe ($q = 1.1552 \pm 0.0304$) is not

significant. t-test results for other pairs of amino acids are found in Figure 4-figure

supplement 1.

Validation of SeqDYN predictions

We now present leave-one-out cross-validation results. We denote the RMSE of the R_2

prediction for the left-out IDP as RMSE(-1). As expected, RMSE(-1) is higher than the

RMSE obtained with the IDP kept in the training set, but the increases are generally slight.

Specifically, all but eight of the IDPs have increases < 0.1 s^{-1}; the largest increase is 0.35 s^{-1},

for CBP-ID4. The mean RMSE(-1), or $\overline{\text{RMSE}}(-1)$, for the 45 IDPs is increased by 0.05 s^{-1}

over $\overline{\text{RMSE}}$, to 1.00 s^{-1}. The latter value is still a distinct improvement over the mean RMSE

1.24 s^{-1} of the null model. The histogram of RMSE(-1) for the 45 IDPs is shown in Figure

5A. It peaks at 0.5 s^{-1}, which is a substantial downshift from the corresponding peak at 0.75 s^{-}

^{1} for σ_{R_2} (Figure 3A). Thirty-four of the 45 IDPs have RMSE(-1) values lower than the

corresponding σ_{R_2}.

To further illustrate the performance of SeqDYN, we present the comparison of

predicted and measured R_2 values for five IDPs: MKK4, α-synuclein, Mev-P$_{NTD}$, Sev-NT,

and CBP-ID4 (Figure 5B-F). A simple common feature is the falloff of R_2 at the N- and C-

termini, resulting from missing upstream or downstream residues that otherwise would be

coupled to the terminal residues, as first recognized by Schwalbe et al. (15). Representative

conformations of the five IDPs are displayed in Figure 2, with residues colored according to

the predicted R_2 values. For four of these IDPs, the RMSE(-1) values range from 0.44 to 0.76

s^{-1} and are scattered around the peak of the histogram, while the RMSE(-1) for the fifth IDP,

namely CBP-ID4, the RMSE(-1) value is 2.01 s^{-1} and falls on the tail of the histogram (Figure

5A). Figure 5B displays the measured and predicted R_2 for MKK4. SeqDYN correctly

predicts higher R_2 values in the second half of the sequence than in the first half. It even

correctly predicts the peak around residue Arg75. The sequence in this region is

H$_{72}$IERLRTH$_{79}$; six of these eight residues belong to the high-end group. In contrast, the

lowest R_2 values occur in the sequence $S_7GGGGSGGGSGSG_{19}$, comprising entirely of two

amino acids in the low-end group.

R_2 values for α-synuclein are shown in Figure 5C. Here SeqDYN correctly predicts

higher R_2 near the C-terminus and a dip around Gly68. However, it misses the R_2 peaks

around Tyr39 and Asp121. MD simulations (1) have found that these R_2 peaks can be

explained by a combination of secondary structure formation (β-sheet around Tyr39 and

polyproline II helix around Asp121) and local (between Tyr39 and Ser42) or long-range

(between Asp121 and Lys96) interactions. SeqDYN cannot account for long-range

interactions (e.g., between β-strands and between Asp121 and Lys96). Figure 5D shows that

SeqDYN gives excellent R_2 predictions for Mev-P$_{NTD}$. It correctly predicts the high peaks

around Arg17, Glu31, Leu193, and lower peaks around Arg235 and Trp285, but does

underpredict the narrow peak around Tyr113.

The overall R_2 profile of Sev-NT is predicted well by SeqDYN, but the peak in the

long helical region (residues 478-491) is severely underestimated (green curve in Figure 5E).

A similar situation occurs for CBP-ID4, where the peak in the second long helical region

(around Glu113) is underpredicted (green curve in Figure 5F). While the measured R_2

exhibits a higher peak in the second helical region than in the first helical region (around

Arg16), the opposite is predicted by SeqDYN. When the R_2 data were included in the training

set (i.e., full training), the second peak is higher than the first one, but that is not a real

prediction because the R_2 data themselves were used for training the model. It merely means

that the SeqDYN functions can be parameterized to produce any prescribed R_2 profile along

the sequence. Indeed, when the R_2 data of CBP-ID4 alone were used to parameterize

SeqDYN, the measured R_2 profile is closely reproduced (Figure 5-figure supplement 1). The

reversal in R_2 peak heights between the two helical regions is the reason for the

aforementioned unusual increase in RMSE when CBP-ID4 was left out of the training set.

R2 boost in long helical regions

It is apparent that SeqDYN underestimates the R_2 of stable long helices. Transient short

helices does not seem to be a problem, since these are present, e.g., in Mev-P_{NTD}, where

transient helix formation in the first 37 residues and between residues 189-198 (26) coincides

with R_2 peaks that are correctly predicted by SeqDYN. SeqDYN can treat coupling between

residues within the correlation length of 5.6 residues, but a much longer helix would tumble

more slowly than implied by an L_{corr} of 5.6, and thus it makes sense that SeqDYN would

underestimate R_2 in that case.

Our solution then is to apply a boost factor to the long helical region. To do so, we

have to know whether an IDP does form long helices and if so what the constituent residues

are. Secondary structure predictors tend to overpredict α-helices and β-strands for IDPs, as

they are trained on structured proteins. One way to counter that tendency is to make the

criteria for α-helices and β-strands stricter. We found that, by filtering PsiPred

(http://bioinf.cs.ucl.ac.uk/psipred) (35) helix propensity scores (p_{Hlx}) with a very high cutoff

of 0.99, the surviving helix predictions usually correspond well with residues identified by

NMR as having high helix propensities. For example, for Mev-P_{NTD}, PsiPred plus filtering

predicts residues 14-17, 28-33, and 191-193 as helical; all of them are in regions that form

transient helices according to chemical shifts (26). Likewise long helices are also correctly

predicted for Sev-NT (residues 477-489) and CBP-ID4 (residues 6-17 and 105-116) (20, 21).

We apply a boost factor, B_{Hlx}, to helices with a threshold length of 12:

$$B_{Hlx} = 1 + \alpha p_{Hlx}\Theta(p_{Hlx} \geq 0.99; L_{Hlx} \geq 12) \qquad (3)$$

The Θ function is 1 if the helix propensity score is above the filtering cutoff and the helix

length (L_{Hlx}) is above the threshold, and 0 otherwise. With a boost amplitude α at 0.5, the

boosted SeqDYN prediction for Sev-NT reaches excellent agreement with the measured R_2

(Figure 5E, red curve). The RMSE(-1) is reduced from 0.76 s^{-1} to 0.38 s^{-1} upon boosting.

Applying the same helix boost to CBP-ID4 also results in a modest reduction in RMSE(-1),

from 2.01 to 1.90 s^{-1} (Figure 5F, red curve). The only other IDP for which PsiPred plus

filtering predicts a long helix is the N-terminal region of lysyl-tRNA synthetase (KRS-NT).

The authors who studied this protein did not report on secondary structure (36), but feeding

their reported chemical shifts to the TALOS+ server

(https://spin.niddk.nih.gov/bax/nmrserver/talos/) (37) found only short stretches of residues

that fall into the helical region of the Ramachandran map. The SeqDYN prediction for KRS-

NT is already good [RMSE(-1) = 0.83 s^{-1}]; applying a helix boost would deteriorate the

RMSE(-1) to 1.16 s^{-1}.

Further test on a set of nine IDPs

We have reserved nine IDPs for testing SeqDYN (parameterized on the training set of 45

IDPs). The level of disorder in these test proteins also spans the full range, from absence of

secondary structures [ChiZ N-terminal region (12), Pdx1 C-terminal region (11), and TIA-1

prion-like domain (17)] to presence of transient short helices [synaptobrevin-2 (7), α-

endosulfine (38), YAP (6), angiomotin-like 1 (AMOTL1) (39)] to formation of stable long

helices [FtsQ (14) and CAHS-8 (9)]. For eight of the nine test IDPs, the RMSEs of SeqDYN

predictions are lower than the experimental σ_{R_2} values, by an average of 0.66 s^{-1}. For the

ninth IDP (Pdx1), the SeqDYN RMSE is slightly higher, by 0.06 s^{-1}, than the experimental

σ_{R_2}. Together, the nine test IDPs have a mean RMSE of 1.13 s^{-1}, close to the $\overline{\text{RMSE}}(-1)$ of

1.00 s^{-1} for the training set in the leave-one-out cross-validation.

The comparison of predicted and measured R_2 profiles along the sequence is presented

in Figure 6A-I. For ChiZ, SeqDYN correctly predicts the major peak around Arg25 and the

minor peak around Arg46 (Figure 6A). The R_2 profile of Pdx1 is largely featureless, except

for a dip around Gly216, which is correctly predicted by SeqDYN (Figure 6B). Correct

prediction is also obtained for the higher R_2 in the first half of TIA-1 prion-like domain than

in the second half (Figure 6C). SeqDYN gives an excellent prediction for synaptobrevin-2,

including a linear increase up to Arg56 and the major peak around Trp89 (Figure 6D).

The prediction is also very good for α-endosulfine, including elevated R_2 around

Glu34, which coincides with the presence of a transient helix, and depressed R_2 in the last 40

residues (Figure 6E). The only miss is an underprediction for the peak around Lys74.

SeqDYN also predicts well the overall shape of the R_2 profile for YAP, including peaks

around Asn70, Leu91, Arg124, and Arg161, but severely underestimates the peak height

around Asn70 (Figure 6F).NOE signals indicate contacts between Met86, Leu91, Fhe95, and

Fhe96 (6); evidently this type of local contacts is captured well by SeqDYN. The R_2 elevation

around Asn70 is mostly due to helix formation: residues 61-74 have helix propensities up to

40% (6). PsiPred predicts helix for residues 62-73, but only residues 65-68 survive the

filtering that we impose, resulting in a helix that is too short to apply a helix boost. The

prediction for AMOTL1 is mostly satisfactory, including peaks around Phe200 and Arg264

and a significant dip around Gly292 (Figure 6G). However, whereas the two peaks have

approximately equal heights in the measured R_2 profile, the predicted peak height around

Phe200 is too low. SCSs indicate helix propensity around both R_2 peaks (39). PsiPred also

predicts helix in both regions, but only five and two residues, respectively, survive after

filtering, and are too short for applying a helix boost.

For FtsQ, SeqDYN correctly predicts elevated R_2 for the long helix [residues 46-74

(14)] but underestimates the magnitude (RMSE = 2.32 s^{-1}; green curve in Figure 6H). PsiPred

plus filtering predicts a long helix formed by residues 47-73. Applying the helix boost

substantially improves the agreement with the measured R_2, with RMSE reducing to 1.71 s^{-1}

(red curve in Figure 6H). SeqDYN also gives a qualitatively correct R_2 profile for CAHS-8,

with higher R_2 for the middle section (residues 95-190) (RMSE = 2.36 s^{-1}; green curve in

Figure 6I). However, it misses the extra elevation in R_2 for the first half of the middle section

(residues 95-145). According to SCS, the first and second halves have helix propensities of

60% and 30%, respectively (9). PsiPred plus filtering predicts helices for residues 96-121,

124-141, 169-173, and 179-189. Only the first two helices, both in the first half of the middle

section, are considered long according to our threshold. Once again, applying the helix boost

leads to marked improvement in the predicted in R_2, with RMSE reducing to 1.92 s^{-1} (red

curve in Figure 6I).

Inputting the sequences of structured proteins predicts R_2 in the unfolded state

SeqDYN is trained on IDPs, what if we feed it with the sequence of a structured protein? The

prediction using the sequence of hen egg white lysozyme, a well-studied single-domain

protein, is displayed in Figure 7A. It shows remarkable agreement with the R_2 profile

measured by Klein-Seetharaman et al. (3) in the unfolded state (denatured by 8 M urea at pH

2 and reduced to break disulfide bridges), including a major peak around Trp62, a second

peak around Trp111, and a third peak around Trp123. Klein-Seetharaman et al. mutated

Trp62 to Gly and the major peak all but disappeared. This result is also precisely predicted by

SeqDYN with the mutant sequence (Figure 7B).

SeqDYN also predicts well the R_2 profiles of other proteins in the unfolded state. For

unfolded apomyoglobin (8 M urea; pH 2.3), Schwarzinger et al. (40) claimed that depressed

R_2 corresponded to stretches of small amino acids (Gly and Ala) whereas elevated

corresponded to local hydrophobic interactions. SeqDYN reproduces all the observed peaks

and valleys in the R_2 profile (Figure 7C). The deepest valley indeed occurs over a Gly/Ala-

rich stretch, G$_{125}$ADAQGA$_{131}$, but the highest peak occurs over a stretch, I$_{102}$KYLEFI$_{108}$, that

contains both hydrophobic and charged residues, all of which are on the high end of the q

parameters (Figure 3B). The R_2 profile of unfolded ubiquitin (8 M urea; pH 2) is relatively

flat, which Wirmer et al. (41) attributed to lack of residual secondary structure, based on the

assumption that β-sheets (major elements of folded ubiquitin) are less resistant to denaturation

than α-helices. SeqDYN predicts a relatively flat R_2 profile (Figure 7D), but the reason is that

the ubiquitin sequence lacks a contiguous stretch of high-q amino acids.

Discussion

We have developed a powerful method, SeqDYN, that predicts the backbone amide

transverse relaxation rates (R_2) of IDPs. The method is based on IDP sequences, is extremely

fast, and available as a web server at https://zhougroup-uic.github.io/SeqDYNidp/. The

excellent performance supports the notion that the ns-dynamics reported by R_2 is coded by the

local sequence, comprising up to 6 residues on either side of a given residue. The amino-acid

types that contribute the most to coupling within a local sequence are aromatic (Trp, Tyr, Phe,

and His), Arg, and long branched aliphatic (Ile and Leu), suggesting the importance of π-π,

cation-π, and hydrophobic interactions in raising R_2. These interactions are interrupted by Gly

and amino acids with short polar sidechains (Ser, Thr, Asn, and Asp), leading to reduced R_2.

Transient short helices produce moderate elevation in R_2, whereas stable long helices result in

a big boost in R_2. Tertiary contacts can also raise R_2, but appear to be infrequent in most IDPs

(1).

It is also possible that R_2 reported by backbone amide ^{15}N relaxation (as is the case for

most of the IDPs studied here) may not be particularly sensitive to exchange effects, which

likely involve tertiary contact formation. For the D2 domain of p27$^{\text{Kip1}}$, the exchange

contributions measured using ^{15}N relaxation were small (< 2.5 s^{-1}) but were as large as 25 s^{-1}

when measured by high-power ^{1}H relaxation dispersion (42). This experiment measures the

effective transverse relation rate, $R_{2,\text{eff}}$, over a range of effective radiofrequency ω_{eff}. The

exchange contribution is maximal for the value $R_{2,\text{eff}}^{\text{low } \omega_{\text{eff}}}$ at low ω_{eff} but is largely quenched

for the value $R_{2,0}^{\text{app}}$ in the high-ω_{eff} limit. The SeqDYN prediction for this IDP matches much

better with $R_{2,0}^{\text{app}}$ than with $R_{2,\text{eff}}^{\text{low } \omega_{\text{eff}}}$ (Figure 8). It is not clear whether this IDP is unique in

forming persistent tertiary contacts that give rise to substantial exchange contributions or the

^{1}H relaxation dispersion experiment is unique in reporting the exchange contributions. At the

minimum, SeqDYN yields the exchange-free portion of the transverse relaxation rate,

enabling easy identification of residues that potentially participate in tertiary contacts. For the

D2 domain of p27$^{\text{Kip1}}$, SeqDYN correctly predicts the $R_{2,0}^{\text{app}}$ local maxima at W76 and Y88. It

is these same two residues that show substantial exchange contributions and putatively

participate in tertiary contact (42). Therefore local contacts may seed tertiary contacts. If $R_{2,\text{eff}}$

data with substantial exchange contributions become available for more IDPs, SeqDYN may

be retrained to make predictions for IDPs forming persistent tertiary contacts.

The q parameters, while introduced here to characterize the propensities of amino

acids to participate in local interactions, appear to correlate with the tendencies of amino acids

to drive liquid-liquid phase separation. Consistent with the rank order of q, Trp, Tyr, and Arg

have been reported to be strong drivers of phase separation, Lys is a moderate driver, whereas

Gly and Ser suppress phase separation (10, 33, 43). Recent measurements of the threshold

concentration produced the following order for the propensity of phase separation by eight

nonpolar amino acids in homotetrapeptides of the form XXssXX (ss: backbone disulfide

bond): Trp > Phe > Leu > Met > Ile > Val > Ala > Pro (44). This order is the same as that of

the q parameters, except that the q values of Ile and Val are in the second and last places,

respectively. Threshold concentrations of IDPs are now predicted reasonably well by coarse-

grained simulations where each amino acid is modeled by a single bead with a Lennard-Jones

diameter d_0 and a stickiness parameter λ (45). Our q parameter shows a good correlation (R^2

= 0.59) with the compound parameter $d_0^3 \lambda$ (Figure 9). Therefore the q parameter may serve as

a predictor for the tendency of an amino acid to drive phase separation. In essence, the same

ability of an amino acid, e.g., Trp, to form interactions with neighboring residues of an IDP in

the free state also applies when it comes to interactions with residues on neighboring chains in

a dense phase.

Our method incorporates ideas from a number of previous efforts at describing R_2.

The first serious effort was by Schwalbe et al. (15), who accounted for contributions from

neighboring residues as additive terms, instead of multiplicative factors as in SeqDYN. Cho et

al. (16) and Delaforge et al. (24) used the running average of the bulkiness parameter over a

window of five to nine residues as a qualitative indicator of R_2. Here again the calculation

was based on an additive model. Sekiyama et al. (17) employed a multiplicative model, with

R_2 calculated as a geometric mean of "indices of local dynamics" over a five-residue window.

These indices, akin to our q parameters, were trained on a single IDR (TIA-1 prion-like

domain) and used to reproduce the measured R_2 for the same IDR. As we have illustrated on

CBP-ID4 (Figure 5-figure supplement 1), training on a single protein merely biases the

parameters to that model and has little value in predicting R_2 for other proteins. In

comparison, SeqDYN is trained on 45 IDPs and its predictions are robust and achieve

quantitative agreement with measured R_2.

Ten of the IDPs tested here have been studied recently by MD simulations using IDP-

specific force fields (1, 12-14). In Table 3 we compare the RMSEs of SeqDYN predictions

with those for R_2 calculations from MD simulations. For five of these IDPs: A1-LCD, Aβ40,

α-synuclein, tau K18, and FtsQ, RMSEs of SeqDYN and MD are remarkably similar. Four of

these IDPs lack significant population of α-helices or β-sheets, but FtsQ forms a stable long

helix. For one other IDP, namely HOX-DFD, MD, by explicitly modeling its folded domain,

does a much better job in predicting R_2 than SeqDYN (RMSEs of 1.40 s^{-1} vs 1.99 s^{-1}).

However, for the four remaining IDPs: p53TAD, Pup, Sev-NT, and ChiZ, SeqDYN

significantly outperforms MD, with RMSEs averaging only 0.47 s^{-1}, compared to the MD

counterpart of 1.14 s^{-1}. Overall, SeqDYN is very competitive against MD in predicting R_2,

but without the significant computational cost. While MD simulations can reveal details of

local interactions, as noted for α-synuclein, and capture tertiary interactions if they occur,

they still suffer from perennial problems of force-field imperfection and inadequate sampling.

SeqDYN provides an accurate description of IDP dynamics at a "mean-field" level, but could

miss idiosyncratic behaviors of specific local sequences.

Deep-learning models have become very powerful, but they usually have millions of

parameters and require millions of protein sequences for training (46). In contrast, SeqDYN

employs a mathematical model with dozens of parameters and requires only dozens of

proteins for training. Reduced models (by collapsing amino acids into a small number of

distinct types) have even been trained on < 10 IDPs to predict propensities for binding

nanoparticles (18) or membranes (19). The mathematical model-based approach may be

useful in other applications where data, similar to R_2, are limited, including predictions of

IDP secondary chemical shifts or residues that bind drug molecules (47) or protein targets, or

even in protein design, e.g., for recognizing an antigenic site or a specific DNA site.

Methods

Collection of IDPs with measured R_2

Starting from six nonhomologous IDPs in our previous MD study (1), we obtained R_2 data for

eight IDPs from the Bimolecular Magnetic Resonance Data Bank (BMRB; https://bmrb.io);

data for two other IDPs were from our collaborators (12, 14). Most of the 54 IDPs studied

here were from searching the literature. Disorder was judged by dispersion in backbone amide

proton chemical shifts, NOE, and SCS. R_2 data that were not available from the authors or

BMRB were obtained by digitizing R_2 plots presented in figures of published papers, using

WebPlotDigitizer (https://automeris.io/WebPlotDigitizer) (48) and further inspected visually.

Homology of IDPs was checked by sequence alignment using Clustal W

(http://www.clustal.org/clustal2) (49), and presented as a clock-like tree using the "ape"

package (http://ape-package.ird.fr) (50). IDPs that had discernible homology with the selected

training set were removed. Removed IDPs included HOX-SCR and β-synuclein from our

previous MD study (1), due to homology with HOX-DFD and α-synuclein, respectively.

Coding for SeqDYN

The training of SeqDYN was coded in python, similar to our previous work for predicting

residue-specific membrane association propensities (ReSMAP; https://zhougroup-

uic.github.io/ReSMAPidp/) (19). The cost function was the sum of mean-squared-errors for

the IDPs in the training set. We used the least_squares function in scipy.optimize, with Trust

Region Reflective as the minimization algorithm and all parameters restricted to the positive

range. For the web server (https://zhougroup-uic.github.io/SeqDYNidp/), we rewrote the prediction code javascript.

ACKNOWLEDGEMENTS

This work was supported by Grant GM118091 from the National Institutes of Health.

REFERENCES

1. S. Dey, M. MacAinsh, H. X. Zhou, Sequence-Dependent Backbone Dynamics of Intrinsically Disordered Proteins. *J Chem Theory Comput* **18**, 6310-6323 (2022).

2. A. R. Camacho-Zarco *et al.*, NMR Provides Unique Insight into the Functional Dynamics and Interactions of Intrinsically Disordered Proteins. *Chem Rev* **122**, 9331-9356 (2022).

3. J. Klein-Seetharaman *et al.*, Long-Range Interactions Within a Nonnative Protein. *Science* **295**, 1719-1722 (2002).

4. A. Abyzov *et al.*, Identification of Dynamic Modes in an Intrinsically Disordered Protein Using Temperature-Dependent NMR Relaxation. *J Am Chem Soc* **138**, 6240-6251 (2016).

5. R. Schneider *et al.*, Visualizing the Molecular Recognition Trajectory of an Intrinsically Disordered Protein Using Multinuclear Relaxation Dispersion NMR. *J Am Chem Soc* **137**, 1220-1229 (2015).

6. M. Feichtinger *et al.*, Long-range structural preformation in yes-associated protein precedes encounter complex formation with TEAD. *iScience* **25**, 104099 (2022).

7. N.-A. Lakomek, H. Yavuz, R. Jahn, Á. Pérez-Lara, Structural dynamics and transient lipid binding of synaptobrevin-2 tune SNARE assembly and membrane fusion. *Proc Natl Acad Sci U S A* **116**, 8699-8708 (2019).

8. A. E. Conicella, G. H. Zerze, J. Mittal, N. L. Fawzi, ALS Mutations Disrupt Phase

Separation Mediated by alpha-Helical Structure in the TDP-43 Low-Complexity C-

Terminal Domain. *Structure* **24**, 1537-1549 (2016).

9. A. Malki *et al.*, Intrinsically Disordered Tardigrade Proteins Self-Assemble into Fibrous

Gels in Response to Environmental Stress. *Angew Chem Int Ed Engl* **61**, e202109961

(2022).

10.   E. W. Martin *et al.*, Valence and patterning of aromatic residues determine the phase

behavior of prion-like domains. *Science* **367**, 694-699 (2020).

11.   E. C. Cook, D. Sahu, M. Bastidas, S. A. Showalter, Solution Ensemble of the C-

Terminal Domain from the Transcription Factor Pdx1 Resembles an Excluded Volume

Polymer. *J Phys Chem B* **123**, 106-116 (2019).

12.   A. Hicks, C. A. Escobar, T. A. Cross, H. X. Zhou, Sequence-dependent correlated

segments in the intrinsically disordered region of ChiZ. *Biomolecules* **10**, 1-23 (2020).

13.   L. Yu, R. Bruschweiler, Quantitative prediction of ensemble dynamics, shapes and

contact propensities of intrinsically disordered proteins. *PLoS Comput Biol* **18**, e1010036

(2022).

14.   S. T. Smrt, C. A. Escobar, S. Dey, T. A. Cross, H. X. Zhou, An Arg/Ala-rich helix in

the N-terminal region of M. tuberculosis FtsQ is a potential membrane anchor of the Z-

ring. *Commun Biol* **6**, 311 (2023).

15.   H. Schwalbe *et al.*, Structural and dynamical properties of a denatured protein.

Heteronuclear 3D NMR experiments and theoretical simulations of lysozyme in 8 M urea.

Biochemistry **36**, 8977-8991 (1997).

16.   M. K. Cho *et al.*, Amino acid bulkiness defines the local conformations and dynamics

of natively unfolded alpha-synuclein and tau. *J Am Chem Soc* **129**, 3032-3033 (2007).

17.   N. Sekiyama *et al.*, ALS mutations in the TIA-1 prion-like domain trigger highly

condensed pathogenic structures. *Proc Natl Acad Sci U S A* **119**, e2122523119 (2022).

18.   D. W. Li, M. Xie, R. Bruschweiler, Quantitative cooperative binding model for

intrinsically disordered proteins interacting with nanomaterials. *J Am Chem Soc* **142**,

10730-10738 (2020).

19.   S. Qin, A. Hicks, S. Dey, R. Prasad, H. X. Zhou, ReSMAP: Web Server for Predicting

Residue-Specific Membrane-Association Propensities of Intrinsically Disordered Proteins.

Membranes **12**, 773 (2022).

20.   M. R. Jensen *et al.*, Quantitative conformational analysis of partially folded proteins

from residual dipolar couplings: application to the molecular recognition element of Sendai

virus nucleoprotein. *J Am Chem Soc* **130**, 8055-8061 (2008).

21.   A. Piai *et al.*, Just a Flexible Linker? The Structural and Dynamic Properties of CBP-

ID4 Revealed by NMR Spectroscopy. *Biophys J* **110**, 372-381 (2016).

22.   S. Maiti *et al.*, Dynamic Studies on Intrinsically Disordered Regions of Two

Paralogous Transcription Factors Reveal Rigid Segments with Important Biological

Functions. *J Mol Biol* **431**, 1353-1369 (2019).

23.   S. Patel, V. Ramanujam, A. K. Srivastava, K. V. Chary, Conformational propensities

and dynamics of a betagamma-crystallin, an intrinsically disordered protein. *Phys Chem

Chem Phys* **16**, 12703-12718 (2014).

24. E. Delaforge *et al.*, Deciphering the Dynamic Interaction Profile of an Intrinsically

Disordered Protein by NMR Exchange Spectroscopy. *J Am Chem Soc* **140**, 1148-1158

(2018).

25. Y. H. Sung, D. Eliezer, Residual structure, backbone dynamics, and interactions

within the synuclein family. *J Mol Biol* **372**, 689-707 (2007).

26. S. Milles *et al.*, An ultraweak interaction in the intrinsically disordered replication

machinery is essential for measles virus function. *Sci Adv* **4**, eaat7778 (2018).

27. Z. Solyom *et al.*, The Disordered Region of the HCV Protein NS5A: Conformational

Dynamics, SH3 Binding, and Phosphorylation. *Biophys J* **109**, 1483-1496 (2015).

28. E. W. Martin *et al.*, Sequence Determinants of the Conformational Properties of an

Intrinsically Disordered Protein Prior to and upon Multisite Phosphorylation. *J Am Chem*

Soc **138**, 15323-15335 (2016).

29. A. M. Janke *et al.*, Lysines in the RNA Polymerase II C-Terminal Domain Contribute

to TAF15 Fibril Recruitment. *Biochemistry* **57**, 2549-2563 (2018).

30. M. Jenner *et al.*, Mechanism of intersubunit ketosynthase-dehydratase interaction in

polyketide synthases. *Nat Chem Biol* **14**, 270-275 (2018).

31. E. M. Bafaro, M. W. Maciejewski, J. C. Hoch, R. E. Dempski, Concomitant disorder

and high-affinity zinc binding in the human zinc- and iron-regulated transport protein 4

intracellular loop. *Protein Sci* **28**, 868-880 (2019).

32. M. G. Murrali, A. Piai, W. Bermel, I. C. Felli, R. Pierattelli, Proline Fingerprint in

Intrinsically Disordered Proteins. *ChemBioChem* **19**, 1625-1629 (2018).

33.   L. E. Wong, T. H. Kim, D. R. Muhandiram, J. D. Forman-Kay, L. E. Kay, NMR

Experiments for Studies of Dilute and Condensed Protein Phases: Application to the

Phase-Separating Protein CAPRIN1. *J Am Chem Soc* **142**, 2471-2489 (2020).

34.   J. M. Zimmerman, N. Eliezer, R. Simha, The characterization of amino acid sequences

in proteins by statistical methods. *J Theor Biol* **21**, 170-201 (1968).

35.   L. J. McGuffin, K. Bryson, D. T. Jones, The PSIPRED protein structure prediction

server. *Bioinformatics* **16**, 404-405 (2000).

36.   H. Y. Cho *et al.*, Characterization of the interaction between lysyl-tRNA synthetase

and laminin receptor by NMR. *FEBS Lett* **588**, 2851-2858 (2014).

37.   Y. Shen, F. Delaglio, G. Cornilescu, A. Bax, TALOS+: a hybrid method for predicting

protein backbone torsion angles from NMR chemical shifts. *J Biomol NMR* **44**, 213-223

(2009).

38.   C. Thapa, P. Roivas, T. Haataja, P. Permi, U. Pentikainen, Interaction mechanism of

endogenous PP2A inhibitor protein ENSA with PP2A. *Febs J* **289**, 519-534 (2022).

39.   A. Vogel, A. Crawford, A. Nyarko, Multivalent Angiomotin-like 1 and Yes-associated

protein form a dynamic complex. *Protein Sci* **31**, e4295 (2022).

40.   S. Schwarzinger, P. E. Wright, H. J. Dyson, Molecular hinges in protein folding: the

urea-denatured state of apomyoglobin. *Biochemistry* **41**, 12681-12686 (2002).

41.   J. Wirmer, W. Peti, H. Schwalbe, Motional properties of unfolded ubiquitin: a model

for a random coil protein. *J Biomol NMR* **35**, 175-186 (2006).

42.   D. Ban, L. I. Iconaru, A. Ramanathan, J. Zuo, R. W. Kriwacki, A Small Molecule

Causes a Population Shift in the Conformational Landscape of an Intrinsically Disordered

Protein. *J Am Chem Soc* **139**, 13692-13700 (2017).

43.   J. Wang *et al.*, A Molecular Grammar Governing the Driving Forces for Phase

Separation of Prion-like RNA Binding Proteins. *Cell* **174**, 688-699 e616 (2018).

44.   Y. Zhang, R. Prasad, S. Su, D. Lee, H. X. Zhou, Amino Acid-Dependent Phase

Equilibrium and Material Properties of Tetrapeptide Condensates. *Cell Rep Phys Sci* **5**,

102218 (2024).

45.   G. Tesei, K. Lindorff-Larsen, Improved predictions of phase behaviour of intrinsically

disordered proteins by tuning the interaction range. *Open Res Eur* **2**, 94 (2022).

46.   A. Rives *et al.*, Biological structure and function emerge from scaling unsupervised

learning to 250 million protein sequences. *Proc Natl Acad Sci U S A* **118**, e2016239118

(2021).

47.   P. Robustelli *et al.*, Molecular Basis of Small-Molecule Binding to alpha-Synuclein. *J*

Am Chem Soc **144**, 2501-2510 (2022).

48.   A. Rohatgi, *Webplotdigitizer: Version 4.6* (2022).

49.   M. A. Larkin *et al.*, Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947-

2948 (2007).

50.   E. Paradis, J. Claude, K. Strimmer, APE: Analyses of Phylogenetics and Evolution in

R language. *Bioinformatics* **20**, 289-290 (2004).

51.   N. Rezaei-Ghaleh, G. Parigi, M. Zweckstetter, Reorientational Dynamics of Amyloid-

beta from NMR Spin Relaxation and Molecular Simulation. *J Phys Chem Lett* **10**, 3369-

3375 (2019).

52.   S. Yao *et al.*, Characterisation of the conformational preference and dynamics of the

intrinsically disordered N-terminal region of Beclin 1 by NMR spectroscopy. *Biochim*

Biophys Acta **1864**, 1128-1137 (2016).

53.   B. Szalaine Agoston, D. Kovacs, P. Tompa, A. Perczel, Full backbone assignment and

dynamics of the intrinsically disordered dehydrin ERD14. *Biomol NMR Assign* **5**, 189-193

(2011).

54.   Z. Zheng, D. Ma, T. L. Yahr, L. Chen, The transiently ordered regions in intrinsically

disordered ExsE are correlated with structural elements involved in chaperone binding.

Biochem Biophys Res Commun **417**, 129-134 (2012).

55.   C. W. Lawrence, S. A. Showalter, Carbon-Detected (15)N NMR Spin Relaxation of

an Intrinsically Disordered Protein: FCP1 Dynamics Unbound and in Complex with

RAP74. *J Phys Chem Lett* **3**, 1409-1413 (2012).

56.   K. A. Burke, A. M. Janke, C. L. Rhine, N. L. Fawzi, Residue-by-Residue View of In

Vitro FUS Granules that Bind the C-Terminal Domain of RNA Polymerase II. *Mol Cell*

60, 231-241 (2015).

57.   T. Gruber *et al.*, Macromolecular Crowding Induces a Binding Competent Transient

Structure in Intrinsically Disordered Gab1. *J Mol Biol* **434**, 167407 (2022).

58.   M. O. Ebert, S. H. Bae, H. J. Dyson, P. E. Wright, NMR relaxation study of the

complex formed between CBP and the activation domain of the nuclear hormone receptor

coactivator ACTR. *Biochemistry* **47**, 1299-1308 (2008).

59.   R. Kiss, D. Kovacs, P. Tompa, A. Perczel, Local structural preferences of calpastatin,

the intrinsically unstructured protein inhibitor of calpain. *Biochemistry* **47**, 6936-6945

(2008).

60.   F. C. Lopes *et al.*, Pliable natural biocide: Jaburetox is an intrinsically disordered

insecticidal and fungicidal polypeptide derived from jack bean urease. *Febs J* **282**, 1043-

1064 (2015).

61. M. De Avila, K. A. Vassall, G. S. Smith, V. V. Bamm, G. Harauz, The proline-rich

region of 18.5 kDa myelin basic protein binds to the SH3-domain of Fyn tyrosine kinase

with the aid of an upstream segment to form a dynamic complex in vitro. *Biosci Rep* **34**,

e00157 (2014).

62. S. Mokhtarzada, C. Yu, A. Brickenden, W. Y. Choy, Structural characterization of

partially disordered human Chibby: insights into its function in the Wnt-signaling pathway.

Biochemistry **50**, 715-726 (2011).

63. M. Schiavina *et al.*, Ensemble description of the intrinsically disordered N-terminal

domain of the Nipah virus P/V protein from combined NMR and SAXS. *Sci Rep* **10**, 19574

(2020).

64. J. L. Neira *et al.*, Dynamics of the intrinsically disordered protein NUPR1 in isolation

and in its fuzzy complexes with DNA and prothymosin alpha. *Biochim Biophys Acta*

Proteins Proteom **1867**, 140252 (2019).

65. B. Mateos *et al.*, The Ambivalent Role of Proline Residues in an Intrinsically

Disordered Protein: From Disorder Promoters to Compaction Facilitators. *J Mol Biol* **432**,

3093-3111 (2020).

66. M. Xie, D. W. Li, J. Yuan, A. L. Hansen, R. Bruschweiler, Quantitative Binding

Behavior of Intrinsically Disordered Proteins to Nanoparticle Surfaces at Individual

Residue Level. *Chemistry* **24**, 16997-17001 (2018).

67. J. Song *et al.*, Intrinsically disordered gamma-subunit of cGMP phosphodiesterase

encodes functionally relevant transient secondary and tertiary structure. *Proc Natl Acad Sci*

U S A **105**, 1505-1510 (2008).

68.   C. Olivieri *et al.*, Multi-state recognition pathway of the intrinsically disordered

protein kinase inhibitor by protein kinase A. *Elife* **9** (2020).

69.   A. Borgia *et al.*, Extreme disorder in an ultrahigh-affinity protein complex. *Nature*

555, 61-66 (2018).

70.   M. A. Ahmed, V. V. Bamm, G. Harauz, V. Ladizhansky, The BG21 isoform of Golli

myelin basic protein is intrinsically disordered with a highly flexible amino-terminal

domain. *Biochemistry* **46**, 9700-9712 (2007).

71.   V. Csizmok, I. C. Felli, P. Tompa, L. Banci, I. Bertini, Structural and dynamic

characterization of intrinsically disordered human securin by NMR spectroscopy. *J Am*

Chem Soc **130**, 16873-16879 (2008).

72.   T. Mittag *et al.*, Structure/function implications in a dynamic complex of the

intrinsically disordered Sic1 with the Cdc4 subunit of an SCF ubiquitin ligase. *Structure*

18, 494-506 (2010).

73.   X. Wang *et al.*, A large intrinsically disordered region in SKIP and its disorder-order

transition induced by PPIL1 binding revealed by NMR. *J Biol Chem* **285**, 4951-4963

(2010).

74.   R. Thapar, G. A. Mueller, W. F. Marzluff, The N-terminal domain of the Drosophila

histone mRNA binding protein, SLBP, is intrinsically disordered with nascent helical

structure. *Biochemistry* **43**, 9390-9400 (2004).

75.   N. Rezaei-Ghaleh *et al.*, Local and global dynamics in intrinsically disordered

synuclein. *Angew Chem Int Ed Engl* **57**, 15262-15266 (2018).

76.   I. R. Chandrashekaran *et al.*, Structure and Functional Characterization of the

Conserved JAK Interaction Region in the Intrinsically Disordered N-Terminus of SOCS5.

Biochemistry **54**, 4672-4682 (2015).

77.   P. Barre, D. Eliezer, Structural transitions in tau k18 on micelle binding suggest a

hierarchy in the efficacy of individual microtubule-binding repeats in filament nucleation.

Protein Sci **22**, 1037-1048 (2013).

78.   E. A. Cino, M. Karttunen, W. Y. Choy, Effects of molecular crowding on the

dynamics of intrinsically disordered proteins. *PLoS One* **7**, e49876 (2012).

79.   J. Harris, M. Shadrina, C. Oliver, J. Vogel, A. Mittermaier, Concerted millisecond

timescale dynamics in the intrinsically disordered carboxyl terminus of gamma-tubulin

induced by mutation of a conserved tyrosine residue. *Protein Sci* **27**, 531-545 (2018).

80.   A. Hicks, M. MacAinsh, H. X. Zhou, Removing thermostat distortions of protein

dynamics in constant-temperature molecular dynamics simulations. *J Chem Theory*

Comput **17**, 5920-5932 (2021).

81.   H. J. Feldman, C. W. Hogue, Probabilistic sampling of protein conformations: new

hope for brute force? *Proteins* **46**, 8-23 (2002).

82.   P. Bernado, M. Blackledge, A self-consistent description of the conformational

behavior of chemically denatured proteins from NMR and small angle scattering. *Biophys*

J **97**, 2839-2845 (2009).

Table 1. Experimental conditions, means and standard deviations of measured R_2, and

SeqDYN prediction RMSEs

Protein name	# of res	Temp (K)	B_0 (MHz)	$\bar{R}_2$ (s^{-1})	σ_{R_2} (s^{-1})	RMSE (s^{-1})	PMID(ref)
Training set (45)[a]							
A1-LCD	131	298	800	2.68	0.46	0.60	32029630(10)
Aβ40	40	278	600	3.40	0.92	0.38	31181936(51)
Ash1	83	278	800	9.80	1.40	1.41	27807972(28)
Beclin1	165	288	800	5.37	1.03	1.14	27288992(52)
CAPRIN1	103	303	600	5.34	0.88	0.72	31898464(33)
CBP-ID4	207	283	700	5.45	2.55	2.01;1.90[b]	29790640(32)
GbnD4-DHD	91	280	700	6.81	1.55	1.28	29309054(30)
ERD14	185	288	600	3.96	0.87	0.54	21336827(53)
ExsE	88	298	600	3.18	0.88	0.76	22138394(54)
FCP1	85	298	500	2.94	0.54	0.43	26286791(55)
FUS	163	298	850	3.48	0.51	0.54	26455390(56)
GAb1	82	298	500	3.99	0.88	0.89	34929201(57)
hACTR	69	304	600	3.26	0.47	0.49	18177052(58)
Hahellin	92	298	800	9.94	2.69	2.85	24671380(23)
hCSD1	141	298	500	3.56	0.93	0.99	18537264(59)
HOX-DFD	90	298	600	6.98	3.15	1.99	30802457(22)
hZIP4-ICL2	100	283	800	9.54	2.37	1.58	30793391(31)
Jaburetox	94	298	800	6.01	2.30	2.27	25605001(60)
KRS-NT	72	303	600	3.26	0.93	0.83	24983501(36)
MBP-xα2	70	295	600	3.83	0.60	0.54	25343306(61)
MKK4	86	278	850	4.49	1.42	0.63	29276882(24)
N-Cbγ	63	298		4.19	1.20	1.25	21182262(62)
Niv-P$_{NTD}$	406	288	700	5.41	1.82	1.66	33177626(63)
NS5A-D2D3	268	278	800	8.62	3.85	2.14	26445449(27)
NUPR1	93	298	600	2.98	0.82	0.76	31325636(64)
OPN	220	310	800	2.59	0.82	0.54	31794728(65)
p53TAD	73	298	850	2.72	0.66	0.33	30240067(66)
PDEγ	87	298		3.96	1.05	0.71	18230733(67)
PKIα	75	300	900	3.41	0.87	0.52	32338601(68)
Mev-P$_{NTD}$	304	298	950	2.92	0.59	0.48	30140745(26)
ProTα	113	283	800	3.40	0.56	0.43	29466338(69)
Pup	64	298	850	2.66	0.51	0.43	30240067(66)
rmBG21	199	300	600	4.06	0.90	0.63	17676872(70)
RPB1	201	277	850	6.48	1.74	1.33	28945358(29)
securin	202	283	500	5.49	1.13	1.08	19053469(71)
Sev-NT	124	298	600	3.20	1.42	0.76;0.38[b]	27112095(4)
Sic1	92	278	500	3.34	0.59	0.48	20399186(72)
SKIPN	71	298		5.64	1.05	1.46	20007319(73)
SLBP-NT	113	298	600	3.96	1.40	1.61	15260482(74)
α-synuclein	140	298	600	2.96	0.53	0.44	30184304(75)
SOCS5-JIR	70	303	800	4.32	2.36	1.91	26173083(76)
tau K18	129	283	700	4.12	0.95	0.83	23740819(77)
TC1	106	298	600	4.65	1.61	1.24	23189168(78)
TDP-43	151	283	500	4.07	1.51	0.96	27545621(8)
γ-tubulin-CT	39	288	500	2.23	0.35	0.27	29127738(79)
Test set (9)							
AMOTL1	207	283	800	8.45	2.55	2.04	35481651(39)
CAHS-8	233	303	850	4.43	3.25	2.36;1.92[b]	34750927(9)
ChiZ	64	298	800	4.33	0.89	0.74	32585849(12)

α-endosulfine	121	298	800	3.21	0.81	0.48	34346186(38)
FtsQ	99	305	800	6.44	3.78	2.32;1.71[b]	36959324(14)
Pdx1	83	298	500	2.98	0.70	0.76	30525611(11)
synaptobrevin-2	96	278	600	5.54	1.80	0.72	30975750(7)
TIA-1	91	310	800	4.01	0.89	0.55	36112647(17)
YAP	122	298	800	3.19	1.44	1.23	35378854(6)

[a]For training set, RMSE is calculated for prediction based on leave-one-out training (using 44 IDPs).

[b]First number is for SeqDYN prediction; second number is after applying a helix boost.

Table 2. Sequences of 54 IDPs[a]

Training set (45)
>A1-LCD (Uniprot ID P04256, residues 186-320; deletion of 258-263)
<u>GS</u>MASASSSQRGRSGSGNFGGGRGGGFGGNDNFGRGGNFSGRGGFGGSRGGGGYGGSGDGYNGFGNDGSNFGGGG
NYNNQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNQGGYGGSSSSSSYGSGRRF
>Aβ40 (Uniprot ID Q28053, residues 7-46)
DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV
>Ash1 (Uniprot ID P34233, residues 420-500)
<u>GA</u>SASSSPSPSTPTKSGKMRSRSSSPVRPKAYTPSPRSPNYHRFALDSPPQSPRRSSNSSITKKGSRRSSGSSPT
RHTTRVCV
>Beclin1 (Uniprot ID Q14457, residues 1-150)
<u>MGSSHHHHHHSQDP</u>MEGSKTSNNSTMQVSFVSQRSSQPLKLDTSFKILDRVTIQELTAPLLTTAQAKPGETQEEE
TNSGEEPFIETPRQDGVSRRFIPPARMMSTESANSFTLIGEASDGGTMENLSRRLKVTGDLFDIMSGQTDVDHPL
SEESTDTLLDQLDT<u>Y</u>
>CAPRIN1 (Uniprot ID Q14444, residues 607-709)
SRGVSRGGSRGARGLMNGYRGPANGFRGGYDGYRPSFSNTPNSGYTQSQFSAPRDYSGYQRDGYQQNFKRGSGQS
GPRGAPRGRGGPPRPNRGMPQMNTQQVN
>CBP-ID4 (Uniprot ID Q92793, residues 1852-2057)
<u>M</u>QQQIQHRLQQAQLMRRRMATMNTRNVPQQSLPSPTSAPPGTPTQQPSTPQTPQPPAQPQPSPVSMSPAGFPSVA
RTQPPTTVSTGKPTSQVPAPPPPAQPPPAAVEAARQIEREAQQQQHLYRVNINNSMPPGRTGMGTPGSQMAPVSL
NVPRPNQVSGPVMPSMPPGQWQQAPLPQQQPMPGLRPVISMQAQAAVAGPRMPSVQ
>GbnD4-DHD (Uniprot ID A0A808VWJ6, residues 412-482)
<u>MKHHHHHHHHGGLVPRGSHGS</u>DEGVPDALRADTVPRAGPVRYARRRYWIGEARSDALAPAAPLEREPLPAEAMGA
YFAIRRTDADDTVAAH
>ERD14 (Uniprot ID P42763, residues 1-185)
MAEEIKNVPEQEVPKVATEESSAEVTDRGLFDFLGKKKDETKPEETPIASEFEQKVHISEPEPEVKHESLLEKLH
RSDSSSSSSSEEEGSDGEKRKKKKEKKKPTTEVEVKEEEKKGFMEKLKEKLPGHKKPEDGSAVAAAPVVVPPPVE
EAHPVEKKGILEKIKEKLPGYHPKTTVEEEKKDKE
>ExsE (Uniprot ID Q9I322, residues 1-81)
MKIESIPPVQPSQDAGAEAVGHFEGRSVTRAAVRGDDRSSVAGLARWLARNVAGDPRSEQALQRLADGDGTPLEA
RTVRRR<u>EFLEGSS</u>
>FCP1 (Uniprot ID Q9Y5B0, residues 879-961)
<u>PG</u>PEEQEEEPQPRKPGTRRERTLGAPASSERSAAGGRGPRGHKRKLNEEDAASESSRESSNEDEGSSSEADEMAK
ALEAELNDLM
>FUS (Uniprot ID P35637, residues 1-163)
MASNDYTQQATQSYGAYPTQPGQGYSQQSSQPYGQQSYSGYSQSTDTSGYGQSSYSSYGQSQNTGYGTQSTPQGY
GSTGGYGSSQSSQSSYGQQSSYPGYGQQPAPSSTSGSYGSSSQSSSYGQPQSGSYSQQPSYGGQQQSYGQQQSYN
PPQGYGQQNQYNS
>GAb1 (Uniprot ID B7Z3B9, residues 510-591)
SSPMIKPKGDKQVEYLDLDLDSGKSTPPRKQKSSGSGSSVADERVDYVVVDQQKTLALKSTREAWTDGRQSTESE
TPAKSVK
>hACTR (Uniprot ID Q9Y6Q9, residues 1023-1091)
GTQNRPLLRNSLDDLVGPPSNLEGQSDERALLDQLHTLLSNTDATGLEEIDRALGIPELVNQGQALEPK
>Hahellin (Uniprot ID Q2SHN6, residues 162-252)
<u>M</u>GEKTVKLYEDTHFKGYSVELPVGDYNLSSLISRGALNDDLSSARVPSGLRLEVFQHNNFKGVRDFYTSDAAELS
RDNDASSVRVSKMETTN
>hCSD1 (Uniprot ID P20810, residues 137-277)
AVPVESKPDKPSGKSGMDAALDDLIDTLGGPEETEEENTTYTGPEVSDPMSSTYIEELGKREVTIPPKYRELLAK
KEGITGPPADSSKPIGPDDAIDALSSDFTCGSPTAAGKKTEKEESTEVLKAQSAGTVRSAAPPQEK
>HOX-DFD (Uniprot ID P07548, residues 337-426)
TDGERIIYPWMKKIHVAGVANGSYQPGMEPKRQRTAYTRHQILELEKEFHYNRYLTRRRRIEIAHTLVLSERQIK
IWFQNRRMKWKKDNK
>hZIP4-ICL2 (Uniprot ID Q6P5W5, residues 424-498)
<u>GDRGPEFELGT</u>LPRDPEDLEDGPCGHSSHSHGGHSHGVSLQLAPSELRQPKPPHEGSRADLVAEESPELLNPEPR
RLSPELRLLPY<u>GHGLSAWSHPQFEK</u>
>Jaburetox (Uniprot ID I1K3K3, residues 230-320)
<u>M</u>GPVNEANCKAAMEIVCRREFGHKEEEDASEGVT<u>T</u>GDPDCPFTKAIPREEYANKYGPTIGDKIRLGDTDLIAEIE
KDFALYGDESVFGGGKVI<u>H</u>
>KRS-NT (Uniprot ID Q15046, residues 1-72)
MAAVQAAEVKVDGSEPKLSKNELKRRLKAEKKVAEKEAKQKELSEKQLSQATAAATNHTTDNGVGPEEESVD
>MBP-xα2 (Uniprot ID P04370, residues 172-237)

```
792    SIGRFFSGDRGAPKRGSGKDSHTRTTHYGSLPQKSQHGRTQDENPVVHFFKNIVTPRTPPPSQGKGRGLS
793    >MKK4 (Uniprot ID P45985, residues 1-86)
794    MAAPSPSGGGGSGGGSGSGTPGPVGSPAPGHPAVSSMQGKRKALKLNFANPPFKSTARFTLNPNPTGVQNPHIER
795    LRTHSIESSGK
796    >N-Cby (Uniprot ID B0QY54, residues 1-63)
797    MPFFGNTFSPKKTPPRKSASLSNLHSLDRSTREVELGLEYGSPTMNLAGQSLKFENGQWIAET
798    >Niv-P_NTD (Uniprot ID P0C1C7, residues 1-406)
799    MDKLELVNDGLNIIDFIQKNQKEIQKTYGRSSIQQPSIKDQTKAWEDFLQCTSGESEQVEGGMSKDDGDVERRNL
800    EDLSSTSPTDGTIGKRVSNTRDWAEGSDDIQLDPVVTDVVYHDHGGECTGYGFTSSPERGWSDYTSGANNGNVCL
801    VSDAKMLSYAPEIAVSKEDRETDLVHLENKLSTTGLNPTAVPFTLRNLSDPAKDSPVIAEHYYGLGVKEQNVGPQ
802    TSRNVNLDSIKLYTSDDEEADQLEFEDEFAGSSSEVIVGISPEDEEPSSVGGKPNESIGRTIEGQSIRDNLQAKD
803    NKSTDVPGAGPKDSAVKEEPPQKRLPMLAEEFECSGSEDPIIRELLKENSLINCQQGKDAQPPYHWSIERSISPD
804    KTEIVNGAVQTADRQRPGTPMPKSRGIPIKK
805    >NS5A-D2D3 (Uniprot ID O92972, residues 2163-2419)
806    GHMASGSLRGGEPEPDVTVLTSMLTDPSHITAETAKRRLARGSPPSLASSSASQLSAPSLKATCTTHHDSPDADL
807    IEANLLWRQEMGGNITRVESENKVVILDSFEPLHADGDEREISVAAEILRKSRKFPSALPIWARPDYNPPLLESW
808    KDPDYVPPVVHGCPLPPTKAPPIPPPRRKRTVVLTESNVSSALAELATKTFGSSGSSAVDSGTATALPDQASDDG
809    DKGSDVESYSSMPPLEGEPGDPDLSDGSWSTVSEEASEDVVCC
810    >NUPR1 (Uniprot ID O60356, residues 2-82)
811    MRGSHHHHHHGSATFPPATSAPQQPPGPEDEDSSLDESDLYSLAHSYLGGGGRKGRTKREAAANTNRPSPGGHER
812    KLVTKLQNSERKKRGARR
813    >OPN (Uniprot ID F1NSM8, residues 46-264)
814    MHQDHVDSQSQEHLQQTQNDLASLQQTHYSSEENADVPEQPDFPDVPSKSQETVDDDDDDDNDSNDTDESDEVFT
815    DFPTEAPVAPFNRGDNAGRGDSVAYGFRAKAHVVKASKIRKAARKLIEDDATTEDGDSQPAGLWWPKESREQNSR
816    ELPQHQSVENDSRPKFDSREVDGGDSKASAGVDSRESQGSVPAVDASNQTLESAEDAEDRHSIENNEVTR
817    >p53TAD (Uniprot ID P04637, residues 1-71)
818    MEEPQSDPSVEPPLSQETFSDLWKLLPENNVLSPLPSQAMDDLMLSPDDIEQWFTEDPGPDEAPRMPEAAPRV
819    >PDEγ (Uniprot ID P61248, residues 1-87)
820    MNLEPPKAEIRSATRVMGGPVTPRKGPPKFKQRQTRQFKSKPPKKGVQGFGDDIPGMEGLGTDITVIAPWEAFNH
821    LELHELAQYGII
822    >PKIα (Uniprot ID P61925, residues 2-76)
823    TDVETTYADFIASGRTGRRNAIHDILVSSASGNSNELALKLAGLDINKTEGEEDAQRSSTEQSGEAQGEAAKSES
824    >Mev-P_NTD (Uniprot ID P03422, residues 1-304)
825    MAEEQARHVKNGLECIRALKAEPIGSLAIEEAMAAWSEISDNPGQERATCREEKAGSSGLSKPCLSAIGSTEGGA
826    PRIRGQGPGESDDDAETLGIPPRNLQASSTGLQCYYVYDHSGEAVKGIQDADSIMVQSGLDGDSTLSGGDNESEN
827    SDVDIGEPDTEGYAITDRGSAPISMGFRASDVETAEGGEIHELLRLQSRGNNFPKLGKTLNVPPPPDPGRASTSG
828    TPIKKGTERRLASFGTEIASLLTGGATQCARKSPSEPSGPGAPAGNVPECVSNAALIQEWTPESGTTISPRSQNN
829    EEGG
830    >ProTα (Uniprot ID P06454, residues 1-111)
831    GPMSDAAVDTSSEITTKDLKEKKEVVEEAENGRDAPANGNAENEENGEQEADNEVDEEEEGGEEEEEEEEGDGE
832    EEDGDEDEEAESATGKRAAEDDEDDDVDTKKQKTDEDD
833    >Pup (Uniprot ID P9WHN4, residues 1-64)
834    MAQEQTKRGGGGGDDDDIAGSTAAGQERREKLTEETDDLLDEIDDVLEENAEDFVRAYVQKGGQ
835    >rmBG21 (Uniprot ID P04370, residues 2-190)
836    GNHSGKRELSAEKASKDGEIHRGEAGKKRSVGKLSQTASEDSDVFGEADAIQNNGTSAEDTAVTDSKHTADPKNN
837    WQGAHPADPGNRPHLIRLFSRDAPGREDNTFKDRPSESDELQTIQEDPTAASGGLDVMASQKRPSQRSKYLATAS
838    TMDHARHGFLPRHRDTGILDSIGRFFSGDRGAPKRGSGKVSLEHHHHHH
839    >RPB1 (Uniprot ID P24928, residues 1773-1970)
840    GHMSPNYTPTSPNYSPTSPSYSPTSPSYSPTSPSYSPSSPRYTPQSPTYTPSSPSYSPSSPSYSPASPKYTPTSP
841    SYSPSSPEYTPTSPKYSPTSPKYSPTSPKYSPTSPTYSPTTPKYSPTSPTYSPTSPVYTPTSPKYSPTSPTYSPT
842    SPKYSPTSPTYSPTSPKGSTYSPTSPGYSPTSPTYSLTSPAISPDDSDEEN
843    >securin (Uniprot ID O95997, residues 1-202)
844    MATLIYVDKENGEPGTRVVAKDGLKLGSGPSIKALDGRSQVSTPRFGKTFDAPPALPKATRKALGTVNRATEKSV
845    KTKGPLKQKQPSFSAKKMTEKTVKAKSSVPASDDAYPEIEKFFPFNPLDFESFDLPEEHQIAHLPLSGVPLMILD
846    EERELEKLFQLGPPSPVKMPSPPWESNLLQSPSSILSTLDVELPPVCCDIDI
847    >Sev-NT (Uniprot ID Q07097, residues 401-524)
848    LSGGDGAYHEPTGGGAIEVALDNADIDLETEAHADQDARGWGGESGERWARQVSGGHFVTLHGAERLEEETNDED
849    VSDIERRIAMRLAERRQEDSATHGDEGRNNGVDHDEDDDAAAVAGIGGI
850    >Sic1 (Uniprot ID P38634, residues 1-90)
851    GSMTPSTPPRSRGTRYLAQPSGNTSSSALMQGQKTPQKPSQNLVPVTPSTTKSFKNAPLLAPPNSNMGMTSPFNG
852    LTSPQRSPFPKSSVKRT
853    >SKIPN (Uniprot ID G3V5R3, residues 59-129)
854    GDGGAFPEIHVAQYPLDMGRKKKMSNALAIQVDSEGKIKYDAIARQGQSKDKVIYSKYTDLVPKEVMNADD
855    >SLBP-NT (Uniprot ID Q9VAN6, residues 17-108)
```

<u>MGSSHHHHHHSSGLVPRGSHM</u>GSGSLNSSASSISIDVKPTMQSWAQEVRAEFGHSDEASSSLNSSAASCGSLAKK
ETADGNLESKDGEGREMAFEFLDGVNEVKFERLVKEEK
>α-synuclein (Uniprot ID P37840, residues 1-140)
MDVFMKGLSKAKEGVVAAAEKTKQGVAEAAGKTKEGVLYVGSKTKEGVVHGVATVAEKTKEQVTNVGGAVVTGVT
AVAQKTVEGAGSIAAATGFVKKDQLGKNEEGAPQEGILEDMPVDPDNEAYEMPSEEGYQDYEPEA
>SOCS5-JIR (Uniprot ID A0A5E4BAI0, residues 12-81)
RSLRQRLQDTVGLCFPMRTYSKQSKPLFSNKRKIHLSELMLEKCPFPAGSDLAQKWHLIKQHTAPVSPHS
>tau K18 (Uniprot ID Q9MYX8, residues 186-314)
QTAPVPMPDLKNVKSKIGSTENLKHQPGGGKVQIINKKLDLSNVQSKCGSKDNIKHVPGGGSVQIVYKPVDLSKV
TSKAGSLGNIHHKPGGGQVEVKSEKLDFKDRVQSKIGSLDNITHVPGGGNKKIE
>TC1 (Uniprot ID Q9NR00, residues 1-106)
MKAKRSHQAIIMSTSLRVSPSIHGYHFDTASRKKAVGNIFENTDQESLERLFRNSGDKKAEERAKIIFAIDQDVE
EKTRALMALKKRTKDKLFQFLKLRKYSIKVH
>TDP-43 (Uniprot ID Q13148, residues 267-414)
<u>GHM</u>NRQLERSGRFGGNPGGFGNQGGFGNSRGGGAGLGNNQGSNMGGGMNFGAFSINPAMMAAAQAALQSSWGMMG
MLASQQNQSGPSGNNQNQGNMQREPNQAFGSGNNSYSGSNSGAAIGWGSASNAGSGSGFNGGFGSSMDSKSSGWG
M
>γ-tubulin-CT (Uniprot ID P53378, residues 439-473)
<u>LLRG</u>AAEQDSYLDDVLVDDENMVGELEEDLDADGDHKLV

Test set (9)
>AMOTL1 (Uniprot ID Q8IY63, residues 178-384)
STQPQQNNEELPTYEEAKAQSQFFRGQQQQQQQQGAVGHGYYMAGGTSQKSRTEGRPTVNRANSGQAHKDEALKE
LKQGHVRSLSERIMQLSLERNGAKQHLPGSGNGKGFKVGGGPSPAQPAGKVLDPRGPPPEYPFKTKQMMSPVSKT
QEHGLFYGDQHPGMLHEMVKPYPAPQPVRTDVAVLRYQPPPEYGVTSRPCQLPFPST
>CAHS-8 (Uniprot ID P0CU50, residues 1-227)
MSGRNVESHMERNEKVVVNNSGHADVKKQQQQVEHTEFTHTEVKAPLIHPAPPIISTGAAGLAEEIVGQGFTASA
ARISGGTAEVHLQPSAAMTEEARRDQERYRQEQESIAKQQEREMEKKTEAYRKTAEAEAEKIRKELEKQHARDVE
FRKDLIESTIDRQKREVDLEAKMAKRELDREGQLAKEALERSRLATNVEVNFDSAAGHTVSGGTTVSTSDKMEIK
RN<u>ENLYFQ</u>
>ChiZ (Uniprot ID I6YA32, residues 1-64)
MTPVRPPHTPDDPLNLRGPLDGPRWRRAEPAQSRRPGRSRPGGAPLRYHRTGVGMSRTGHGSRPV
>α-endosulfine (Uniprot ID O43768, residues 1-121)
MSQKQEEENPAEETGEEKQDTQEKEGILPERAEEEAKLKAKYPSLGQKPGGSDFLMKRLQKGQKYFDSGDYNMAKA
KMKNKQLPSAGPDKNLVTGDHIPTPQDLPQRKSSLVTSKLAGGQVE
>FtsQ (Uniprot ID Q8IY63, residues 1-99)
MTEHNEDPQIERVADDAADEEAVTEPLATESKDEPAEHPEFEGPRRRARRERAERRAAQARATAIEQARRAAKRR
ARGQIVSEQNPAKPAARGVVRGLK
>Pdx1 (Uniprot ID P52945, residues 204-283)
<u>GPG</u>EEDKKRGGGTAVGGGGVAEPEQDCAVTSGEELLALPPPPPPGGAVPPAAPVAAREGRLPPGLSASPQPSSVA
PRRPQEPR
>synaptobrevin-2 (Uniprot ID P63027, residues 1-96)
MSATAATAPPAAPAGEGGPPAPPPNLTSNRRLQQTQAQVDEVVDIMRVNVDKVLERDQKLSELDDRADALQAGAS
QFETSAAKLKRKYWWKNLKMM
>TIA-1 (Uniprot ID P31483, residues 320-386)
<u>MGSSHHHHHHHHHHHHSENLYFQ</u>GGQYVPNGWQVPAYGVYGQPWSQQGFNQTQSSAPWMGPNYSVPPPQGQNGSM
LPSQPAGYRVAGYETQ
>YAP (Uniprot ID P46937, residues 50-171)
AGHQIVHVRGDSETDLEALFNAVMNPKTANVPQTVPMRLRKLPDSFFKPPEPKSHSRQASTDAGTAGALTPQHVR
AHSSPASLQLGAVSPGTLTPTGVVSGPAATPTAQHLRQSSFEIPDDV

[a]Terminal tags and other insertions are underlined.
906

907 Table 3. RMSEs (s^{-1}) of R_2 predictions by SeqDYN and MD for 10 IDPs

IDP name	SeqDYN	MD
A1-LCD	0.60[a]	0.59[d, e]
Aβ40	0.38[a]	0.38[d]
HOX-DFD	1.99[a]	1.40[d]
α-synuclein	0.44[a]	0.50[d]
p53TAD	0.33[a]	1.04[f]
Pup	0.43[a]	1.00[f]
Sev-NT	0.38[a, b]	1.10[d, g]
tau K18	0.83[a]	0.80[d]
ChiZ	0.74[c]	1.40[h]
FtsQ	1.71[c, b]	1.70[i]

[a]Based on leave-one-out training (using 44 IDPs).

[b]Helix boost applied.

[c]Based on training by the full training set (45 IDPs).

[d]From ref (1).

[e]RMSE is scaled down by a factor of 2.39, to correct for the effect of temperature (MD at 288

K; see Figure 3-figure supplement 1C).

[f]From ref (13).

[g]RMSE is scaled down by a factor of 2.99, to correct for the effects of temperature and

magnetic field (MD at 274 K and 850 MHz; see Figure 3-figure supplement 1B).

[h]Originally calculated in ref (12) with correction in ref (80).

[i]From ref (14).

FIGURE CAPTIONS

Figure 1. Clock-like tree plot showing lack of homology among the 45 IDPs. The level of homology between two sequences is measured by the distance from their convergence point to the center of the clock. The highest level of apparent identity is between A1-LCD and TDP-43, at 25%, but these two proteins differ in both secondary structure formation and R_2 characteristics. There is, however, a 20-residue overlap between the N-terminus of MBP-xα2 and the C-terminus of rmBG21.

Figure 2. Representative conformations of five IDPs. (A-E) MKK4, α-synuclein, Mev-P$_{\text{NTD}}$, Sev-NT, and CBP-ID4. Conformations were initially generated using TraDES (http://trades.blueprint.org) (81), selected to have radius of gyration close to predicted by a scaling function $R_{\text{g}} = 2.54N^{0.522}$ (Å) (82). Conformations for residues predicted as helical by PsiPred plus filtering were replaced by an ideal helix. Finally residues are colored according to a scheme ranging from green for low predicted R_2 to red for high predicted R_2.

Figure 3. Properties of the 45 IDPs in the training set. (A) Histograms of means and standard deviations, calculated for individual proteins. Curves are drawn to guide the eye. Inset: correlation between $\bar{R}_2$ and σ_{R_2}. (B) Experimental mean scaled R_2 (msR_2) and SeqDYN q parameters, for the 20 types of amino acids. Note that Pro residues have low msR_2 for the lack of backbone amide proton. Amino acids are in descending order of q.

Figure 3-figure supplement 1. Possible effects of sequence length, temperature, and magnetic field on R_2. (A) Lack of dependence of $\bar{R}_2$ on sequence length. (B) Counts of IDPs with R_2 measured at various temperatures. (C) Matching of R_2 profiles of Sev-NT measured at two temperatures after uniform scaling. (D) Matching of R_2 profiles of A1-LCD measured at two temperatures after uniform scaling. (E) Counts of IDPs with R_2 measured at various magnetic fields.

Figure 4. SeqDYN model parameters. (A) Correlation between msR_2 and q. The values are also displayed as bars in Figure 3B. (B) Correlation of msR_2 and q with amino-acid molecular mass. (C) Correlation of msR_2 and q with bulkiness. (D) The optimal correlation

length and deterioration of SeqDYN prediction as the correlation length is moved away from

the optimal value.

Figure 4-figure supplement 1. T-test of on the q parameters for pairs of amino acids. q

parameters were obtained from five-fold cross-validation training, resulting in five

independent values for each q parameter. Mean presented as red bars; standard deviation

presented as error bar. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ns, not significant. q

parameters for all neighboring pairs not explicitly indicated are not significantly different.

Figure 5. Quality of SeqDYN predictions. (A) Histogram of RMSE(-1). Letters indicate

RMSE(-1) values of the IDPs to be presented in panels (B-F). (B-F) Measured (bars) and

predicted (curves) R_2 profiles for MKK4, α-synuclein, Mev-P_{NTD}, Sev-NT, and CBP-ID4. In

(E) and (F), green curves are SeqDYN predictions and red curves are obtained after a helix

boost.

Figure 5-figure supplement 1. Close reproduction (curve) of the measured R_2 profile (bars) of

CBP-ID4 when that set of data alone was used to parameterize SeqDYN. The resulting model

has no value for predicting R_2 for other proteins.

Figure 6. Measured (bars) and predicted (curves) R_2 profiles for ChiZ N-terminal region,

TIA1 prion-like domain, Pdx1 C-terminal region, synaptobrevin-2, α-endosulfine, YAP,

AMOTL1, FtsQ, and CAHS-8. In (C), R_2 does not fall off at the N-terminus because the

sequence is preceded by an expression tag MGSSHHHHHHHHHHHHS. In (H) and (I), green

curves are SeqDYN predictions and red curves are obtained after a helix boost.

Figure 7. R_2 profiles predicted (curves) by SeqDYN show close agreement with those

measured (bars) on structured proteins in the unfolded state. (A) Wild-type lysozyme (8 M

urea; pH 2; cysteine-methylated). (B) Lysozyme with Trp62 to Gly mutation (pH 2).

Methylated cysteines were treated as Ala in the SeqDYN predictions. (C) Apomyoglogin (8

M urea; pH 2.3). (D) Ubiquitin (8 M urea; pH 2).

Figure 8. Comparison between SeqDYN prediction (curves) and effective transverse

relaxation rate (bars) from ^{1}H dispersion relaxation experiment. (A) $R_{2,\text{eff}}$ in the high-ω_{eff}

limit. (B) $R_{2,\text{eff}}$ at low ω_{eff}.

Figure 9. Correlation between the stickiness parameters (λ) and the NMR relaxation

parameters (q). The regression line is shown as dashes.

Dynamics is a crucial link between sequence and function for intrinsically disordered proteins

(IDPs). Qin and Zhou developed a sequence-based method to predict IDP dynamics. The

predictions provide deep physicochemical insight and can also serve as indicators of

properties including phase-separation propensities of IDP sequences.

Keywords: NMR spectroscopy; phase separation; intrinsically disordered proteins; backbone

dynamics