Leptin antagonism improves Rett syndrome phenotype in symptomatic male *Mecp2-null* mice.

Yasmine Belaïdouni[1], Diabe Diabira[1], Mélanie Brosset-Heckel[1], Victoria Valsamides[1], Jean-Charles Graziano[1#], Catarina Santos[3], Clément Menuet[1], Gary A. Wayman[2] and Jean-Luc Gaiarsa[1*].

[1]INMED, INSERM, Aix Marseille Univ, France.

[2]Program in Neuroscience, Department of Integrative Physiology and Neuroscience, Washington State University, Pullman, WA, United States.

[3]Phenotype Expertise, Marseille, France.

[*]Correspondence to: jean-luc.gaiarsa@inserm.fr

#present address: Aix-Marseille Univ, Institut Paoli Calmettes, INSERM, CNRS, CRCM, Marseille, France.

Running head: Rett syndrome and leptin.

Key words: leptin, breathing, Rett syndrome, hippocampus, E/I balance, Mecp2.

ABSTRACT

Rett syndrome (RTT) is a severe neurodevelopmental disorder that arise from *de novo* mutations in the X-linked gene *MECP2* (methyl-CpG-binding protein 2). Circulating levels of the adipocyte hormone leptin are elevated in RTT patients and rodent models of the disease. Leptin targets a large number of brain structures and regulates a wide range of developmental and physiological functions which are altered in RTT. We hypothesized that elevated leptin levels might contribute to RTT pathogenesis. Accordingly, we show that pharmacological antagonism of leptin or genetic reduction of leptin production prevents the degradation of health status, weight loss and the progression of breathing and locomotor deficits. At the neuronal level, the anti-leptin strategies rescue the hippocampal excitatory/inhibitory imbalance and synaptic plasticity impairment. Targeting leptin might therefore represent a new approach for RTT treatment.

INTRODUCTION

Rett syndrome (RTT; OMIM identifier #312750) is a rare non-inherited genetic neurodevelopmental disorder characterized by severe impairments affecting nearly every aspect of child's life. After an apparently normal development until about 6–18 months of age, RTT patients undergo a period of rapid regression including, but not limited to, loss of purposeful hand movements, loss of acquired speech, breathing irregularities, seizures, and severe cognitive deficits (Sandweiss et al., 2020). Typical RTT cases arise from *de novo* mutations in the X-linked gene *MECP2* (methyl-CpG-binding protein 2). The MECP2 protein binds to methylated DNA to modulate the expression of thousands of genes (Lewis et al., 1992) and plays an important role in brain development and functioning throughout the lifespan (Cheval et al., 2012) .

Several clinical features of the human disorder are recapitulated in *Mecp2*-deficient mice

(Ricceri et al., 2008). These mouse models therefore represent an essential tool for deciphering

the cellular mechanisms of the disease and for testing potential therapeutic treatments. The

demonstration that phenotypic rescue is possible in *Mecp2*-deficient mice upon re-expressing

Mecp2 gene in adult mice (Guy et al., 2007) has indicated that RTT is not an irrevocable disease,

giving hope for therapeutic interventions.

Many aspects of neurochemistry are altered in RTT, including the levels of neurotrophic

factors and neuromodulators, some of which holding therapeutic promises (Ehinger et al., 2018).

Among these deregulated factors, epidemiological and animals studies showed that the

circulating levels of leptin are elevated in RTT patients and rodent models of the disease (Blardi

et al., 2007, 2009; Acampa et al., 2008; Park et al., 2014; Fukuhara et al., 2019). Leptin, the

product of the obese (*ob*) gene, is a circulating hormone secreted mainly from the white

adipocytes and transported across the blood brain barrier to the hypothalamus, to suppress food

intake (Ahima and Flier, 2000). It is increasingly clear that the hypothalamus is not the only site

of leptin action, and that food intake is not its only biological role. Leptin is a pleiotropic

hormone that modulates motivation (Davis et al., 2010), cognitive functions (Harvey, 2022),

anxiety (Guo et al., 2012), the excitability of neuronal networks (Shanley et al., 2002),

epileptiform activities (Mora-Muñoz et al., 2018), breathing activity (Gauda et al., 2020), and

more (Signore et al., 2008; Lim et al., 2009; Li et al., 2018). Large bodies of evidence also

indicate that leptin acts as an important neurotrophic factor during perinatal periods promoting

neuritic growth and synaptogenesis in a variety of brain structures including the hypothalamus

(Bouret et al., 2004; Pinto et al., 2004) and hippocampus (O'Malley et al., 2007; Moult and

Harvey, 2008; Dhar et al., 2014; Dumon et al., 2018; Sahin et al., 2020, 2021). Due to these

important physiological and developmental functions, dysregulation – either excess or deficiency

– in leptin availability or signaling can have far reaching effects. Accordingly, animal studies

showed that abnormal leptin levels at any time of lifespan can lead to adverse outcomes including

social and cognitive impairments, altered breathing activity, seizures susceptibility and abnormal

brain development (Harvey, 2003; Greco et al., 2010; Dumon et al., 2018; Mora-Muñoz et al.,

2018; Gauda et al., 2020). Thus, excess levels of leptin in RTT patients (Blardi et al., 2007, 2009;

Acampa et al., 2008) might contribute to the disorders associated with Rett syndrome.

To address this hypothesis, we confirmed an excess of serum leptin in male $Mecp^{2tm1-1Bird}$

mice (hereafter referred to as $Mecp2^{-/y}$ mice) and assessed the effects of "anti-leptin" strategies in

ameliorating deficits typically displayed by this RTT mouse model. We show that

pharmacological antagonism of leptin or genetic reduction of leptin production prevents the

degradation of health status, weight loss as well as the progression of breathing and locomotor

deficits in male $Mecp2^{-/y}$ mice and, at the neuronal level, rescues the hippocampal

excitatory/inhibitory imbalance and synaptic plasticity impairment.

RESULTS

Serum leptin levels and leptin mRNA expressions are altered in *Mecp2$^{-/y}$* mice.

The aim of this study was to assess the contribution of leptin in RTT. Therefore, as a prerequisite for this study, we evaluated the circulating levels of leptin by ELISA, and the *Lep* mRNA expression from two main sources of leptin, the white adipose tissues (WAT) and gastrocnemius muscle, by qRT-PCR in male *Mecp2$^{-/y}$* and WT littermates. We found that *Mecp2$^{-/y}$* mice exhibited elevated circulating leptin levels compared to WT littermates, and that this difference reach significance at postnatal day (P) 30 and 50 (p=0.045 and 0.0002 respectively, Two-tailed unpaired *t*-test, **Figure 1A, supplementary table 1**). Interestingly, the elevated level of leptin in *Mecp2$^{-/y}$* mice was observed despite their decrease in body weight (19.8 ± 0.6 vs 13.6 ± 0.5g, p<0.0001, two-tailed unpaired *t*-test, **Figures 1B**). *Mecp2$^{-/y}$* mice also exhibited higher expression of leptin mRNA in gastrocnemius muscle (**Figure 1C**), as well as in visceral and inguinal WAT (**Figures 1D and E**) compared to their WT littermates. The difference reaches significance at P30 in inguinal WAT (p=0.03, Two-tailed unpaired *t*-test, **supplementary table 1)** and at P50 in gastrocnemius muscle (p=0.0015), visceral WAT (p=0.03) and inguinal WAT (p=0.03, Two-tailed unpaired *t*-test, **supplementary table 1**). Altogether, these data show that circulating leptin levels and leptin mRNA expression are elevated in male *Mecp2$^{-/y}$* mice compared to their age-matched WT littermates, providing a framework to assess the role of leptin in RTT-associated symptoms.

Leptin antagonism during the symptomatic stage prevents the progression of breathing deficits in *Mecp2$^{-/y}$* mice.

Respiratory disorders have been well described in patients and rodent models of RTT (Ramirez et

al., 2013). Leptin has been reported to regulate breathing acting on brainstem respiratory

networks and carotid body (Gauda et al., 2020). We therefore investigate the possible

contribution of leptin to RTT-associated breathing disorders. We first characterized the

respiratory activity of *Mecp2*$^{-/y}$ mice from our breeding colony. Consistent with previous studies

(Katz et al., 2009), when compared with their age-matched WT littermates, P40 *Mecp2*$^{-/y}$ mice

developed respiratory disorders characterized by increased number of apneas (p<0.0001, Two-

tailed unpaired *t*-test, **Figure 2A, supplementary table 2**), cycle duration variability expressed as

breathing irregularity score (14.6$\pm$0.7 vs 21.5$\pm$0.8, 28 and 29 mice, p<0.0001, Two-tailed

unpaired *t*-test, **data not shown**), and minute ventilation (5.5$\pm$0.4 vs 9.4$\pm$0,7 ml g^{-1} min^{-1}, 28 and

29 mice, p<0.0001, Two-tailed unpaired *t*-test, **data not shown**). These breathing distresses

worsen with age in *Mecp2*$^{-/y}$ mice, while the respiratory parameters remained constant in WT

(**Figures 2B and C, supplementary table 2**). Next, to test the possible role of leptin in RTT-

associated breathing disorders, P40 WT and *Mecp2*$^{-/y}$ mice received daily sub-cutaneous injection

of leptin recombinant (5µg/g) or every other day sub-cutaneous injection of a pegylated leptin

antagonist (5µg/g) (Gertler and Elinav, 2014), during 10 days. Sham mice received the same

volume of vehicle. We used plethysmography to record breathing activity of each mouse at P40,

the day before the start of treatment, and at P50, the last day of treatment. Leptin treatment of WT

mice led to a significant increase in the number of apneas (p=0.017 when compared to P40, Two-

tailed paired *t*-test, **Figure 2B,** p=0.012 when compared to P50 sham WT mice, Two-tailed

unpaired *t*-test, **Figure 2D, supplementary table 2**) and breathing irregularity score (p=0.04

when compared to P50 sham WT mice, Two-tailed unpaired *t*-test**, Figures 2E, supplementary**

table 2), while in sham WT mice these parameters remained unchanged (**Figures 2B, D and E,**

supplementary table 2). The mean values of tidal volume (**data not shown**), respiratory

frequency (**data not shown**) and minute ventilation (p=0.72 when compared to P50 treated WT

mice, Two-tailed unpaired *t*-test, **Figure 2F, supplementary table 2**) were not affected by the

leptin-treatment.

We next investigated the effect of anti-leptin treatment (5µg/g, every other day during 10 days)

on the breathing activity of the *Mecp2*$^{-/y}$ mice. While the sham and anti-leptin treated *Mecp2*$^{-/y}$

mice show similar phenotypic profile at the beginning of the treatment (p=0.43, Two-tailed

unpaired *t*-test, **Figure 2C**), anti-leptin treated *Mecp2*$^{-/y}$ mice show better breathing parameters

(number of apneas and breathing irregularity score) compared to sham *Mecp2*$^{-/y}$ mice at the end

of the treatment (p=0.015 and 0.0015, Two-way ANOVA followed by a Bonferroni's multiple

comparison, **Figures 2C, D and E, supplementary table 2**). The mean values of tidal volume

(**data not shown**), respiratory frequency (**data not shown**) and minute ventilation (p>0.999,

Two-way ANOVA followed by a Bonferroni's multiple comparison, **Figure 2F, supplementary**

table 5) of P50 *Mecp2*$^{-/y}$ mice were not affected by the anti-leptin-treatment. We also tested the

effect of leptin treatment and found no effect on the breathing parameters in *Mecp2*$^{-/y}$ mice

(**Figures 2D, E and F, supplementary table 2**).

Altogether, these data suggest a contribution of leptin to the progression of breathing difficulties

in symptomatic *Mecp2*$^{-/y}$ mice.

Leptin antagonism during the symptomatic stage prevents the worsening of health

condition of *Mecp2*$^{-/y}$ mice.

We next assessed the efficacy of the anti-leptin treatment on the overall health of the *Mecp2*$^{-/y}$

mice. The thoroughly described loss in body weight in *Mecp2*$^{-/y}$ mice (p=0.00001, Two-tailed

unpaired *t*-test **Figures 3A and B**, **supplementary table 3**) was abolished by the anti-leptin

treatment (5µg/g, every other day during 10 days, from P40 to P50) while the leptin-treated

Mecp2$^{-/y}$ mice (5µg/g, 10 daily injections from P40 to P50) showed no difference compared to

sham *Mecp2$^{-/y}$* littermates (p=0.0001 and 0.59 respectively at P50, Two-way ANOVA followed

by a Bonferroni's multiple comparison, **Figures 3B and C**, **supplementary table 3**). The leptin

treatment also did not induce significant change in the body weight of the WT mice (p=0.06 at

P50, Two-tailed unpaired *t*-test, **Figures 3C**, **supplementary table 3).** We also performed a

general scoring of the health condition of the mice, including tremor, general aspect, spontaneous

activity, limb grasp and posture. WT and *Mecp2$^{-/y}$* mice showed significant difference at P40

(p=0.000001, two-tailed unpaired *t*-test, **Figure 3D, supplementary table 3**). The general health

scoring performed on sham- and anti-leptin treated *Mecp2$^{-/y}$* mice the day before the start of

treatment (P40) and the last day of treatment (P50) revealed a deterioration of the health

condition in sham *Mecp2$^{-/y}$* mice that was not observed in anti-leptin treated *Mecp2$^{-/y}$* mice

(p=0.03 and 0.16, two-tailed paired *t*-test, **Figure 3E, supplementary table 3**). Improved

symptoms are posture, tremor and general aspect while spontaneous activity and limb grasp

showed no significant improvement (**Figure 3F, supplementary table 3**). Although it does not

reach significance, this improvement in overall health of the anti-leptin treated *Mecp2$^{-/y}$* mice was

accompanied by a decrease of early lethality, with 50% of the sham- and anti-leptin treated

Mecp2$^{-/y}$ mice surviving after the postnatal day 48.5 and 59 respectively (p=0.1, two-tailed

unpaired *t*-test, p=0.16, Kaplan- Meier log rank test, **Figure 3G**). No lethality nor change in the

health scoring were observed in WT mice treated with either vehicle or leptin (n=10 for both,

data not shown).

We also assess the effect of the anti-leptin treatment on the locomotor activity and motor

coordination of *Mecp2$^{-/y}$* mice in the open field and the accelerating rotarod tests. In the open

filed test, the distance traveled by the *Mecp2$^{-/y}$* mice was reduced compare to their WT littermates

(p=0.016, Two-tailed unpaired *t*-test **Figures 4A**, **supplementary table 7**). This locomotor

deficit worsens with age in *Mecp2$^{-/y}$* mice (p=0.03, P40 vs P50 *Mecp2$^{-/y}$* sham, Two-tailed paired

t-test, **Figures 4B**, **supplementary table 4**). The anti-leptin treatment (5µg/g, every other day

during 10 days, from P40 to P50) prevented the degradation of locomotor activity of the *Mecp2$^{-/y}$*

mice (p=0.66, P40 vs P50 *Mecp2$^{-/y}$* anti-leptin, Two-tailed paired *t*-test and p=0.04, P50 *Mecp2$^{-/y}$*

sham vs P50 *Mecp2$^{-/y}$* anti-leptin, Two-tailed unpaired *t*-test, **Figures 4B and C**, **supplementary**

table 4). Analysis of the behavior of the leptin-treated WT mice indicates that they performed

similarly to their sham littermates (p=0.4, Two-tailed unpaired *t*-test **Figure 4C**, **supplementary**

table 4). In the accelerating rotarod test, *Mecp2$^{-/y}$* mice showed a significant decrease in the

latency to fall compared to WT littermates (p<0.0001, Mann Whitney *U* test, **Figure 4D**,

supplementary table 7). This phenotype did not worsen with age (p=0.29, P40 vs P50 *Mecp2$^{-/y}$*

sham, Wilcoxon matched pairs signed rank test, **Figures 4E**, **supplementary table 4**). Neither

the leptin- nor the anti-leptin treatment affected the performances of respectively WT and *Mecp2^{-}*

$^{/y}$ mice on the accelerating rotarod test (p=0.18 and 0.42 respectively, two-tailed unpaired *t*-test,

Figure 4F, supplementary table 4).

Overall, these observations show that the anti-leptin treatment ameliorates the overall health

condition, i.e., posture, tremor and general aspect and locomotor activity of the symptomatic

Mecp2$^{-/y}$ mice.

Leptin antagonism during the symptomatic stage does not affect anxiety in *Mecp2$^{-/y}$* mice.

Leptin has been reported to exert anti-depressant and pro-cognitive like effects in rodents (Guo et

al., 2012; Harvey, 2022). We therefore investigate the possible contribution of leptin to cognitive

behavioral deficits repeatedly reported in RTT models (Kerr et al., 2010). Our first set of

behavioral tests examined anxiety, cognition and social interaction in P50 WT and *Mecp2*$^{-/y}$ mice

from our breeding colony. We first compared the spontaneous exploration in the elevated plus

maze (EPM) as a measure of anxiety and found that *Mecp2*$^{-/y}$ mice spent more time in the open

arms than their WT littermate (p<0.0001, Two-way ANOVA followed by a Bonferroni's multiple

comparison, **Figure 4G**, **supplementary table 4**). The mean dipping time was also increased in

Mecp2$^{-/y}$ mice (p=0.086, Two-way ANOVA followed by a Bonferroni's multiple comparison,

Figure 4H, **supplementary table 4**) a behavior also consistent with decreased anxiety. However,

no genotype effects were found in tests used to evaluate object recognition and social behavior

(**supplementary Figure S1, supplementary table S1**). We therefore limited the study of the role

of leptin to the EPM. Analysis of the behavior of the leptin-treated WT and anti-leptin treated-

Mecp2$^{-/y}$ mice in the EPM indicates that they performed similarly to their vehicle-treated WT

littermate, i.e. they stayed similar time in the open arms and exhibited similar mean dipping time

(**Figures 4H and G**, **supplementary table 4**).

Overall, these observations show that the anti-leptin treatment does not improve anxiety

impairment in the symptomatic *Mecp2*$^{-/y}$ mice.

Symptomatic leptin antagonism does not improve seizures susceptibility in *Mecp2*$^{-/y}$ mice.

Seizures are predominant features in RTT patients and abnormal cortical activities have been

observed in *Mecp2*-deficient mice (Glaze, 2002; D'Cruz et al., 2010). Leptin modulates the

excitability of neuronal networks and exerts both anti-convulsant and pro-convulsant effects

(Mora-Muñoz et al., 2018). To gain insight into the possible contribution of leptin in RTT-

associated epileptic activity, we challenged P50 *Mecp2*$^{-/y}$ mice and their WT littermates with the

volatile convulsant agent flurothyl (2,2,2-trifluoroethyl ether). Flurothyl induced severe tonic-

clonic seizures with a loss of posture in both WT- and *Mecp2*$^{-/y}$ mice. The onset of seizures was

more rapid in *Mecp2*$^{-/y}$ mice as compared to their WT littermates (p=0.0007, Two-tailed unpaired

t-test, **Figure 5A, supplementary table 5**). Moreover, all *Mecp2*$^{-/y}$ mice died after the tonic-

clonic seizures, while lethality was observed in 44% of WT mice (p=0.0029, Fisher exact test,

Figure 5C). We next investigated the effect of daily leptin treatment (0.5µg/g during 10

consecutive days) on the seizure latency in WT mice and found no significant difference in

seizure latency (p=0.34, Two-tailed unpaired *t*-test, **Figure 5B, supplementary table 5**) and

lethality (p>0.999, Fisher exact test, **Figure 5D**) compared to WT sham WT. Similarly, neither

the leptin- nor the anti-leptin (0.5µg/g, every other day during 10 days) treatments affected

flurothyl-induced seizures latency (p>0.999 for both, two-way ANOVA followed by a

Bonferroni's multiple comparison) and lethality (p>0.999 for both, Fisher exact test) in *Mecp2*$^{-/y}$

mice (**Figures 5B and D, supplementary table 5**).

Altogether, these data suggest that leptin does not contribute to seizure susceptibility in

symptomatic *Mecp2*$^{-/y}$ mice.

Leptin antagonism during the symptomatic stage restores the E/I balance in CA3

hippocampal neurons in *Mecp2*$^{-/y}$ mice.

We then assessed the potential benefits of anti-leptin treatment at the neuronal level. Alteration in

the synaptic excitation/inhibition (E/I) balance is a widespread feature of neuronal networks in

Mecp2-deficient mice (Li, 2022). Leptin has been reported to modulate the development and

functioning of both glutamatergic and GABAergic hippocampal synapses (Moult and Harvey,

2009; Dhar et al., 2014; Guimond et al., 2014; Dumon et al., 2018; Sahin et al., 2020, 2021;

Harvey, 2022), raising the possibility that elevated leptin levels might contribute to the altered E/I

balance in $Mecp2^{-/y}$ mice. We therefore performed whole cell patch clamp recordings of CA3

pyramidal neurons performed on acute hippocampal slices obtained from WT and $Mecp2^{-/y}$ mice

and found that, as already reported (Calfa et al., 2015; Lozovaya et al., 2019), the E/I balance was

increased at both presymptomatic (0.9 ± 0.2 vs 4.5 ± 0.9 in respectively P30 WT (10 cells, 3 mice)

and $Mecp2^{-/y}$ (13 cells, 3 mice) mice littermates, p=0.014, Two-tailed Mann-Whitney test, **data**

not shown) and symptomatic stages (p<0.0001 at P50, Two-tailed Mann Whitney test, **Figures**

6A, B and D, supplementary table 6). We next assed the effect of the leptin and leptin

antagonism treatment on the E/I balance. The leptin treatment (0.5µg/g during 10 consecutive

days from P40 to P50) led to a significant increase in the E/I balance in P50 WT mice, thus

phenocopying the $Mecp2^{-/y}$ mice (p=0.014, Two-tailed Kruskal-Wallis test followed by Dunn's

post hoc comparison, **Figure 6C, supplementary table 2).** This increase in the E/I balance is

mainly accounted for by a reduction of GABAergic activity (p=0.002, **Figure 6E,**

supplementary table 6). When administered to *db* mice (mice deficient for the long-form leptin

receptor, the only receptor capable of activating intracellular pathways (Ahima and Flier, 2000)),

leptin had no effect on the E/I balance (1.57 ± 0.3 vs 1.58 ± 0.3 for respectively *db*-sham (13 cells,

2 mice) and *db*-lep (9 cells, 2 mice), p=0.787, Two-tailed Mann-Whitney test) and GABAergic

activity (26.5 ± 6.1 vs 20.6+4.1 Hz, p=0.638, Two-tailed Mann-Whitney test, **data not shown**). In

contrast to leptin treatment, the anti-leptin treatment (0.5µg/g every other day from P40 to P50)

had no effect on the E/I balance of WT mice (p>0.999 when compared to WT-sham, Two-tailed

Kruskal-Wallis test followed by Dunn's post hoc comparison, **supplementary table 6**).

Treatment of $Mecp2^{-/y}$ mice revealed that leptin had no significant effect on the E/I balance

(p=>0.999, Two-tailed Kruskal-Wallis test followed by Dunn's post hoc comparison, **Figure 6C,**

supplementary table 6). In contrast, the anti-leptin treatment restored the E/I balance in *Mecp2$^{-/y}$*

mice (p=0.005 when compared to Mecp2$^{y/-}$-sham, p>0.9999 when compared to WT-sham, Two-

tailed Kruskal-Wallis test followed by Dunn's post hoc comparison, **Figure 6C, supplementary**

table 6) and increased the GABAergic activity (p=0.028 when compared to *Mecp2$^{-/y}$*-sham,

p>0.9999 when compared to WT-sham, Two-tailed Kruskal-Wallis test followed by Dunn's post

hoc comparison, **Figure 6E, supplementary table 6**).

Altogether, these data show that the leptin-treatment increases the E/I balance in WT mice

phenocopying the *Mecp2$^{-/y}$* mice, while the anti-leptin treatment rescues the E/I balance in

Mecp2$^{-/y}$ mice. These observations suggest a contribution of leptin to the hippocampal E/I

imbalance in symptomatic *Mecp2$^{-/y}$* mice.

Leptin antagonism during the symptomatic stage restores the hippocampal synaptic

plasticity in *Mecp2$^{-/y}$* mice.

Impairments of synaptic plasticity are common features of mice RTT models (Asaka et al., 2006;

Moretti et al., 2006; Lozovaya et al., 2019) and leptin has been reported to modulate the strength

of glutamatergic synapses (McGregor and Harvey, 2019). These observations prompted us to

examine the effect of leptin and leptin antagonist treatment on long term potentiation (LTP) at the

hippocampal Schaffer collaterals-CA1 synapses of P50 WT and *Mecp2$^{-/y}$* mice. Field EPSPs were

recorded in the stratum radiatum and evoked by stimuli of increasing strength. The slope of the

input-output curves obtained from *Mecp2$^{-/y}$* mice were not significantly different from WT

littermates showing that the basal neurotransmission at the Schaffer collateral-CA1 synapses is

unaltered in symptomatic *Mecp2$^{-/y}$* mice (the slope of curve afferent volley amplitude vs field

EPSP slope was 5.1$\pm$0.7 vs 5.2$\pm$0.7 in respectively WT and *Mecp2*$^{-/y}$ mice, p=0.8, Mann Whitney

U test, data not shown). Next, LTP was induced by two trains of 100 Hz tetanic stimulation (1 sec

duration, 20 sec interval). As already reported (Asaka et al., 2006), the magnitude of LTP was

reduced in *Mecp2*$^{-/y}$ compared to WT littermates (p=0.012 at 45-50 min post-tetanus, repeated

measure ANOVA, **Figure 7A, supplementary table 7**). The magnitude of potentiation in the

first minutes following the tetanic stimulation was also reduced (p=0.08 at 0-2 min post-tetanus,

repeated measure ANOVA, **Figure 7A, supplementary table 7**). Leptin treatment of WT mice

(5µg/g during 10 consecutive days from P40) attenuated the magnitude of late and early LTP

compared to sham-treated but these differences did not reach significance (p=0.16 and 0.27 for

the early and late LTP, repeated measure ANOVA, **Figure 7B, supplementary table 7**).

Conversely, the anti-leptin treatment (0.5µg/g every other day during 10 days from P40) restored

the magnitude of early and late LTP in *Mecp2*$^{-/y}$ mice (p=0.08 and 0.03 respectively, repeated

measure ANOVA, **Figure 7C, supplementary table 7**).

Altogether, these data suggest that leptin contributes to the impairment of hippocampal LTP in

symptomatic *Mecp2*$^{-/y}$ mice.

RTT-associated symptoms are improved in leptin haplo insufficient mice.

To further evaluate the possible contribution of leptin in RTT phenotype, we generated *Mecp2*$^{-/y}$

mice that are haplo insufficient for leptin. For this, female *Mecp2*$^{+/-}$ mice were backcrossed with

heterozygous leptin deficient male mice (*ob/+*). The resulting double transgenic *Mecp2*$^{-/y;ob/+}$

mice showed normalized leptin levels at P50-60 (p=0.8 compared to WT mice, Two-way

ANOVA followed by a Bonferroni's multiple comparison, **Figure 8A**, **supplementary table 8**).

We next asked whether the RTT associated deficits were ameliorated in these leptin haplo

insufficient mice at P50-60, an advanced symptomatic stage. Whole cell recordings performed on

hippocampal slices revealed that the E/I balance and sGABA-PSCs frequency are improved in

CA3 hippocampal neurons of *Mecp2*$^{-/y;ob/+}$ mice (p=0.39 and 0.02 respectively when compared to

WT mice, **Figure 8A and C**, **supplementary table 8**). Field recordings on hippocampal slices

further revealed that the magnitude of early and late LTP at the Schaffer collateral-CA1 synapses

was increased in *Mecp2*$^{-/y;ob/+}$ mice (p=0.08 and 0.05 respectively when compared to *Mecp2*$^{-/y}$

mice, **Figure 8D**, **supplementary table 8**).   The *Mecp2*$^{-/y;ob/+}$ mice also showed improved

breathing activity (p=0.002 when compared to WT, **Figure 8E**, **supplementary table 8)** and less

weight loss (p=0.05 when compared to WT, **Figure 8F**, **supplementary table 8**).

Overall, these data show that reduced leptin production improves the symptoms of *Mecp2*$^{-/y}$ mice.

DISCUSSION

Here, we show that *Mecp2*$^{-/y}$ mice have elevated circulating leptin levels and leptin

mRNA expression compared to their age-matched WT littermates. Knowing the pleiotropic

central and peripheral roles of leptin, this finding provided a framework for investigating the

possible contribution of this adipocyte hormone in RTT pathogenesis. We thus tested the rescue

potential of leptin antagonism approaches. We found that 10-day treatment from P40 of male

Mecp2-null mice with a leptin antagonist prevents the worsening of several symptoms including

breathing difficulty, locomotor deficit, weight loss and degradation of general health condition.

At the neuronal level, the anti-leptin treatment restores the E/I balance and synaptic plasticity in

the hippocampus. A further validation of our hypothesis is the benefits observed in mice haplo

insufficient for leptin (*Mecp2*$^{-/y;ob/+}$ mice). The leptin antagonism strategy was however

ineffective against seizure susceptibility, anxiety and motor coordination deficit. In line with the

contribution of leptin to RTT pathogenesis, we showed that leptin treatment of WT mice induces

hippocampal E/I imbalance, reduces hippocampal synaptic plasticity and causes breathing

abnormality, thus phenocopying some RTT symptoms.

Previous studies found that leptin concentrations were increased in patients with Rett

syndrome, but not associated with increases in weight gain or body mass index (Blardi et al.,

2007, 2009; Acampa et al., 2008). Elevated leptin level could be related to chronic inflammatory

state due to frequent respiratory infections (Cortelazzo et al., 2014) as well as hormonal and/or

metabolic deregulations (Motil et al., 2011; Stagi et al., 2015). Medications, such as

anticonvulsants, can also alter circulating leptin levels in patients (Belcastro et al., 2013).

However, according to previous studies (Park et al., 2014; Fukuhara et al., 2019), we showed that

Mecp2-null mice also present elevated levels of circulating leptin, even though they did not

receive any medication. This leptin elevation was observed in males in different models and

genetic backgrounds as well as in female *Mecp2*$^{+/-}$ mice (Park et al., 2014; Torres-Andrade et al.,

2014; Fukuhara et al., 2019) (present study and unpublished observations) highlighting the

robustness of this finding and its possible relevance to the clinical situation. The mechanism

underlying the increase in leptin protein and mRNA levels observed in *Mecp2*-deficient mice is

presently unknown but likely rely on non-cell autonomous mechanisms since (i) neuron-specific

Mecp2 knockout mice (*Mecp2*$^{tm1.1Jae}$ KO mice) showed increased circulating leptin levels and

WAT leptin mRNA expression (Park et al., 2014) and (ii) adipocyte-specific Mecp2 knockout

mice (*Mecp2*^{Adi} KO mice) showed down regulated WAT leptin mRNA expression and similar

circulating leptin levels compared to WT mice (Liu et al., 2019). An increase in WAT mass

could explain the elevated leptin levels in *Mecp2*-deficient mice. Previous studies have shown

that Lep mRNA expression and circulating leptin levels are increased in *Mecp2*-deficient mice

(Torres-Andrade et al., 2014) and POMC neuron-specific Mecp2 knockout mice (Wang et al.,

2014) in association with an increase in body weight. Two other studies reported elevated leptin

levels in *Mecp2*-deficient mice without weight change but with increased WAT mass (Park et al.,

2014; Fukuhara et al., 2019). In the present study, WAT mass was not measured, but it is worth

noting that the progressive increase in leptin levels and leptin mRNA expression were observed

despite a decline in body weight. Adipose tissue is also present within skeletal muscle.

Intramuscular adipose tissue (IMAT) accumulation is observed in pathological conditions such as

muscle dystrophy, obesity and metabolic diseases (Sciorati et al., 2015). IMAT accumulation

may underlie elevated leptin mRNA expression in muscle of symptomatic male *Mecp2*-null mice,

which exhibit hypotrophic fibres and tissue fibrosis (Conti et al., 2015).

We have shown that pharmacological leptin antagonism and genetic reduction of leptin production ameliorate some RTT phenotype and prevent the degradation of others in symptomatic male *Mecp2*-null mice. We used a pegylated super active mouse leptin antagonist (Peg-SMLA). This antagonist poorly penetrated in the central nervous system, and inhibits leptin activity by two mechanisms: (i) by blocking the transport of circulating leptin across the blood brain barrier (BBB), thus reducing central leptin levels, and (ii) by blocking the binding of leptin to its peripheral receptors, thus reducing peripheral leptin signaling (Elinav et al., 2009). Likewise, haplo insufficiency for leptin will reduce both central and peripheral actions of leptin in *Mecp2*$^{-/y;ob/+}$ mice. Several observations support the hypothesis that the beneficial action of leptin antagonism relies on a reduction in the central action of leptin. First, functional leptin receptors are expressed throughout the brain (Caron et al., 2010). Then, *in vitro* studies indicate that leptin directly regulates the excitability and synaptic function of neurons within different brain regions including the hippocampus (Shanley et al., 2002; Harvey et al., 2006; Solovyova et al., 2009; Dhar et al., 2014; Guimond et al., 2014; Dumon et al., 2018; Sahin et al., 2021) and nucleus tractus solitarii, a crucial site for the modulation of breathing, cardiovascular control and food intake (Williams and Smith, 2006; Neyens et al., 2020; Yu et al., 2022). Lastly, *in vivo* experiments show that local/systemic injections of leptin, as well as chemogenetic/optogenetic activation of leptin receptor expressing neurons, modulate physiological and behavioral responses, and that targeted deletion of central leptin receptors prevent these effects (Oomura et al., 2006; Inyushkin et al., 2009; Vong et al., 2011; Guo et al., 2013; Zuure et al., 2013; Bassi et al., 2014, 2015; Yu et al., 2022). The respiratory network is complex and the effects of leptin on respiratory control are still poorly understood. However, alterations in the E/I balance have been

reported in respiratory nuclei that express leptin receptor such as the nucleus tractus solitarius and

the rostro ventrolateral medulla (Medrihan et al., 2008; Chen et al., 2018). The latter could

contribute to apneustic and irregular breathing in *Mecp2*-deficient mice (Ramirez et al., 2013).

Additional work is required to determine whether and how leptin antagonisms affect the E/I

balance in these brain regions, but also in cortical areas where the E/I balance is reduced (Li,

2022). Finally, we cannot completely rule out a peripheral effect of treatments in addition to the

central effect, acting for example on carotid bodies.

Knowing the facilitatory role of leptin on breathing and synaptic function, the present

finding might seem counter-intuitive. It is however worth noting that an excess of leptin can lead

to a blunted response to leptin, a condition known as "leptin resistance". Thus, rodents fed with a

high-fat diet that causes elevated leptin levels and leptin resistance also show altered

hippocampal synaptic function, breathing difficulties and changes in anxiety level (Grillo et al.,

2011; Valladolid-Acebes et al., 2012; Farr et al., 2015; Gauda et al., 2020). It cannot be totally

excluded that the anti-leptin treatment or the haplo insufficiency for leptin, by decreasing the

central levels of leptin, reduces the phenomenon of resistance and allows leptin to exert its

beneficial effects. Similarly, leptin treatment of WT mice could lead to leptin resistance, which

would be responsible for the symptoms observed in these mice. Elevated leptin levels have also

been observed in children with autism spectrum disorders (Ashwood et al., 2008; Blardi et al.,

2010; Raghavan et al., 2018). Yet, the clinical features seen in ASD do not completely overlap

those broader of Rett syndrome, suggesting that the functional and physiological actions of leptin

may depend, among other things, on the mutations associated with the etiology of the disease, in

the present situation of the Mecp2 mutations.

The success of most genetic and pharmacological therapeutic approaches relies on the

ability of the virus or drugs to cross the BBB. To overcome this difficulty, we have used a

competitive antagonist of leptin whose main action is to block the transport of leptin across the

BBB (Elinav et al., 2009). We have shown that the treatment is well tolerated (no premature

death during the treatment, no overt side effect) holding potential therapeutic application.

Moreover leptin antagonist therapy has been reported to be efficient for the prevention or

treatment of diseases in which high leptin levels is part of the picture (Singh et al., 2013; Fisch et

al., 2020). It is noteworthy that leptin antagonist treatment also improves bone mineralization

(Chapnik et al., 2013) and muscle function (Gonzalez et al., 2021), two peripheral symptoms

observed in RTT. Furthermore, hyperleptinemia has been proposed to play a role in sympathetic

over activity and cardiac abnormalities observed in RTT (Acampa et al., 2008). Overall, these

observations suggest that leptin might participate to RTT clinical manifestations other than

central synaptic activity, breathing and weight loss, highlighting the global therapeutic interest of

this approach.

Given the progressive nature of Rett symptoms and the high inter-individual variability,

we assessed their evolution (i.e., the difference between the values at the beginning and at the end

of treatment) to estimate the potential efficacy of the leptin antagonist treatment. Our data show

that treatment with the leptin antagonist in the symptomatic period delays the progression of

certain symptoms (i.e. breathing, locomotor activity, weight loss, general health status) rather

than curing them. The performance to rotarod, which does not show any degradation in the

studied period, was not modified. The notion of a "pre-symptomatic" period in RTT has recently

been questioned by studies showing subtle symptoms in RTT girls and early alterations in animal

models before the appearance of overt symptoms (Cosentino et al., 2019). Therefore, even if

translational applications would be difficult, it will be interesting in the future to determine

whether pre-symptomatic treatments have greater positive effects.

MATERIALS and METHODS

Animals

All animal procedures were carried out in accordance with the European Union Directive of 22 September (2010/63/EU) and have been approved by the Ethical Committee for Animal Experimentation (APAFIS-Number 17605-2018-1119-1115-7094) delivered by the French Ministry of Education and Research.

Experiments were performed on male wild type and $Mecp^{2tm1-1Bird}$ mice (Guy et al., 2001) C57BL/6 genetic background (Jackson Laboratory, stock number 003890). Hemizygous mutant males (hereafter referred as to $Mepc2^{-/y}$ mice) were generated by crossing heterozygous females ($Mecp2^{+/-}$) with C57BL/6 wild type males. $Mecp2^{-/y}$ mice haplo insufficient for leptin (hereafter referred as to $Mecp2^{-/y;ob/+}$) were generated by crossing heterozygous females $Mecp2^{+/-}$ with C57BL/6 heterozygous leptin deficient males mice (Jackson Laboratory genotyping protocol, strains B6.Cg-Lepob/J, ID 000632). Animals were housed in a temperature-controlled environment with a 12h light/dark cycle and free access to food and water. Genotyping was performed by PCR techniques according to Jackson Laboratory protocols.

Leptin and anti-leptin injection

Recombinant murine leptin (Peprotech, New Jersey, USA) was reconstituted in sterile water and daily injected (5 µg/g) sub-cutaneous at 12-14 hr a.m during 10 days starting at postnatal (P) 40. The pegylated super active mouse leptin antagonist (Peg-SMLA, Protein Laboratories, Rehovot, Israel) was reconstituted in sterile water (Elinav et al., 2009; Gertler and Elinav, 2014). As the

half-life of the Peg-SMLA was about 24 hours, mice were injected (5 µg/g) sub-cutaneous at 12-

14 hr a.m every other day during 10 days starting at P40. Control mice received same volume

injections of vehicle.

Leptin immunoassay

Blood samples were centrifuged (10.000 rpm, 10 min, 4°C) immediately after collection at 10–11

hr a.m. Quantification of endogenous leptin was performed with Mouse Leptin ELISA Kit

(BioVendor R&D^R, Brno, Czech Republic) following the manufacturer's protocol. The measured

concentration of samples was calculated from the standard curve and expressed as ng/ml.

Real-time reverse transcription quantitative polymerase chain reaction

Visceral and inguinal fat pads and gastrocnemius muscles were dissected, rapidly frozen in liquid

nitrogen and stored at -80°C. Total RNAs from fat pads and gastrocnemius muscle were isolated

using a RNeasy Mini kit (74106, Qiagen, Germany) and RNA Fibrous Tissue Mini kit (74704,

Qiagen) respectively, following the manufacturer's instructions and quantified by reading the

absorbance at 260 nm (NanoPhotometer, IMPLEN). RNAs were converted to cDNA using 1 µg

RNA and a QuantiTect Reverse Transcription kit (Qiagen) according to manufacturer's

instructions. Quantitative RT-PCR was performed with a Light Cycler 480 SYBR Green IMaster

(Roche Applied Science) following the manufacturer's instructions, using the following

oligonucleotides (QuantiTect Primer Assays, Qiagen): Leptin (Mm_Lep_1_SG QT00164360

(NM_008493)), and glyceraldehyde-3-phosphate dehydrogenase (GAPDH, QT001199633).

Relative mRNA values were calculated using the LC480 software and GAPDH as the

housekeeping gene. PCR was performed in triplicate.

Hippocampal slice preparation and electrophysiological recordings

Brains were removed and immersed into ice-cold (2° to 4°C) artificial cerebrospinal fluid

(ACSF) with the following composition: 126 mM NaCl, 3.5 mM KCl, 2 mM CaCl2, 1.3 mM

MgCl2, 1.2 mM NaH2PO4, 25 mM NaHCO3, and 11 mM glucose (pH 7.4) equilibrated with

95% O2 and 5% CO2. Hippocampal slices (350μm thick) were cut using a vibrating microtome

(Leica VT1000S, Germany) in ice-cold oxygenated choline-replaced ACSF and were allowed to

recover at least 90 min in ACSF at room temperature (25°C). Slices were then transferred to a

submerged recording chamber perfused with oxygenated (95% O2 and 5% CO2) ACSF (3

ml/min) at 34°C.

Whole cell recordings were performed from CA3 pyramidal neurons in the voltage-clamp mode

using an axopatch 200B amplifier (Molecular Devices LLC, San Jose, USA). To assess the

excitatory/inhibitory balance, the glass recording electrodes (4-7 MΩ) were filled with a solution

containing 100 mM K-gluconate, 13 mM KCl, 10 mM Hepes, 1.1 mM EGTA, 0.1 mM CaCl2, 4

mM Mg–adenosine 5′-triphosphate, and 0.3 mM Na–guanosine 5′-triphosphate. The pH of the

intracellular solution was adjusted to 7.2, and the osmolality was adjusted to 280 mOsmol liter−1.

With this solution, $GABA_A$-receptor mediated postsynaptic currents ($GABA_A$-PSCs) reversed at

−70 mV. $GABA_A$-PSCs and Glut-PSCs were recorded at a holding potential of −45 and -70 mV

respectively. Spontaneous synaptic currents, were recorded with Axoscope software version 8.1

(Molecular Devices LLC, San Jose, USA) and analyzed offline with Mini Analysis Program

version 6.0 (Synaptosoft).

Field recordings were performed in area CA1. A cut was made between CA1 and CA3. The Schaffer collaterals/commissural fibers were stimulated using a bipolar tungsten electrode (66 μm; A-M Systems, WA, USA) placed on the surface of the stratum radiatum of CA1 (10–50 μs, 5–15 V, 0.03 Hz). Extracellular tungsten electrodes (California Fine Wire,CA, USA) were used to record dendritic field excitatory postsynaptic potentials (fEPSP) from the stratum radiatum of the CA1 region. The signals were amplified using a DAM80 amplifier (WPI, UK), digitized with an Axon Digidata 1550B (Molecular Devices, CA, USA), recorded with Axoscope software version 8.1 (Molecular Devices, CA, USA) and analyzed offline with Mini Analysis Program version 6.0 (Synaptosoft, GA, USA) by measuring the onset (a 30–70% rising phase) slope of the fEPSP. LTP was induced by two 100 Hz trains of 100 stimuli 20 sec apart at 50% of the maximal intensity. The slope of the fEPSP was measured and expressed relative to the preconditioning baseline.

Plethysmography.

The breathing activity of non-anesthetized freely moving mice was recorded using a constant flow whole body plethysmograph (EMKA technologies, Paris, France) with 200ml animal chambers maintained at 25±0.5 $^{\circ}$C and ventilated with air (600ml min-1). The breathing activity of pairs of WT and *Mecp2$^{-/y}$* mice littermates was simultaneously recorded. Mice were allowed to acclimate to the experimental room for 1 hour and to the plethysmography chamber and airflow for approximatively 30 mins before breathing measurement. Breathing activity was recorded during 1 hour. A differential pressure transducer measured the changes in pressure in the body chamber resulting from the animal's respiration. Signals from the pressure transducer were stored on a personal computer and analyzed offline via the Spike 2 interface and software (Cambridge

Electronic Design, Cambridge, UK). Only periods of quiet breathing without body movements

were analyzed, during which the number of apneas (> two respiratory cycles) per hour was

quantified.

Fluorothyl-induced seizures

To test seizure susceptibility, we used the convulsant agent flurothyl (2,2,2-trifluoroethyl ether).

P50-55 male mice were placed in a transparent, ventilated but hermetically sealed, cage, into

which the epileptic agent flurothyl was delivered at a rate of 25nl/min using a nano-pump

(Harvard Apparatus) and homogeneously distributed using a mini-ventilator incorporated into the

chamber. Behavioral responses were recorded using a video camera. The latency of tonic-clonic

seizures was determined post hoc. The progressive injection of flurothyl into the cage produced a

stereotypical behavioral manifestation of limbic seizure episodes that started from animal

immobility, followed by intermediate stages (rigid posture, tranquility period, jiggles, and jerks)

and ended by severe tonic-clonic seizures (stage 4). The injection of flurothyl was stopped ten

seconds after beginning of tonic-clonic seizures and the cage was ventilated with fresh air. Some

mice regain motor and exploratory activity after 10 mins. Others die during the tonic seizures.

The percentage of surviving/dying mice varies according to the genotype and the treatment (see

results section).

General health scoring and lifespan

Weight was measured every day from the beginning of the treatment. The phenotypic score was

evaluated the day before treatment and the last day of treatment. Severity score, typically used in

RTT phenotypic assessments (Guy et al., 2001; Scaramuzza et al., 2021), was used to group

animals into 4 severity classes: absence of phenotype (4) to severe phenotype (0). The parameters

scored are: tremor, posture, limb grasp, spontaneous activity in the home cage and general aspect.

To be noticed, mice that rapidly lost weight were euthanized for ethical reasons. The day of the

sacrifice was considered as the endpoint of lifespan assessment.

Accelerating rotarod

A rotarod apparatus (Biological Research Apparatus, Gemonio, Italy) was used to measure the

motor coordination. After a 5 mins habituation session (4 r.p.m), each mouse was given three

trials with the rate of rotation increasing from 4 to 40 r.p.m. over 5 mins. The trial ended when

the mouse fell from the rod or remained on the rotarod for at least 5 mins. The time spent on the

rotarod was recorded by an automated unit, which stopped as the mouse fell. The mouse was

placed back in its home cage for 10 mins between each trial. The latency to fall was determined

as the best of the 3 trials.

Behavior

All the behavioral tests were performed by Phenotype Expertise, Inc. (France). For all tests,

animals were first acclimated to the behavioral room for 30 minutes. WT and $Mecp2^{-/y}$ mice

underwent elevated plus maze, open field, three chamber test and spontaneous social interaction

test at P50. WT and $Mecp2^{-/y}$ treated mice underwent elevated plus maze and open field at P40,

before the beginning of the treatment, and at P50, at the end of the treatment.

Elevated-Plus Maze. The EPM was used to assess anxiety state of animals. The device consists of

a labyrinth of 4 arms 5 cm wide located 80 cm above the ground. Two opposite arms are open

(without wall) while the other two arms are closed by side walls. The light intensity was adjusted

to 20 Lux on the open arms. Mice were initially placed on the central platform and left free to

explore the cross-shaped labyrinth for 5 minutes. Maze was cleaned and wiped with H2O and

with 70% ethanol between each mouse. Animal movement was video-tracked using Ethovision

software 11.5 (Noldus). Time spent in open and closed arms, the number of entries in open arms,

as well as the distance covered, are directly measured by the software.

Open-field. Open field was used to evaluate both the locomotor activity of the animals. Mice

were individually placed in a 40 x 40 cm square arena with an indirect illumination of 60 lux.

Mouse movement was video-tracked using Ethovision software 11.5 (Noldus) for 10 minutes.

Total distance traveled and time in center (exclusion of a 5 cm border arena) are directly

measured by the software. Grooming (time and events) and rearing were manually counted in live

using manual functions of the software, by an experimented behaviorist. The open-field arena

was cleaned and wiped with H2O and with 70% ethanol between each mouse.

Three-chamber social preference test. The test was performed as described previously (Bertoni et

al., 2021) with the following modifications to accommodate for the lower mobility of the

animals. A square arena 40 x 40 cm was used, with the wired cages placed in opposite diagonal

corners. The task was carried out in four trials. The three-chambers apparatus was cleaned and

wiped with water and with 70% ethanol between each trial and each mouse. In the first trial

(habituation), a test mouse was placed in the center of the arena and was allowed to freely

explore each chamber. The mouse was video-tracked for 5 min using Ethovision software. At the

end of the trial, the animal was briefly removed from the arena to allow for the preparation of the

following trial. In the second trial (social exploration), a 8- weeks old C57BL/6J congener mouse

(S1) was placed randomly in one of the two wire cages to avoid a place preference. The second

wire cage remained empty (E). Then the test mouse was placed in the center of the arena and

allowed to freely explore for 10 min. A second 8-weeks old C57BL/6J congener mouse (S2) was

placed in the second wire cage for the third trial (social discrimination). Thus, the tested mouse

had the choice between a familiar mouse (S1) and a new stranger mouse (S2) for 10 min. At the

end of the trial, the mouse was returned to home-cage for 30 min. For the fourth trial (short-term

social memory), S2 was replaced by a new stranger mouse (S3), the familiar mouse (S1) staying

the same. Then tested mouse was allowed to freely explore the arena for 10 min. Time spent in

each chamber and time of contact with each wire cage (with a mouse or empty) were calculated

using Ethovision software. The measure of the real social contact is represented by the time spent

in nose-to-nose interactions with the unfamiliar or familiar mouse. This test was performed using

grouped house mice of 4 months old.

Spontaneous social interaction test. The tested mouse was placed into the same OF arenas that

were used for the OF test and allowed to explore this empty arena for 2.5 min. Immediately

following this initial stage, the mouse was again allowed to explore the OF for an additional 10

min with the target (Swiss mouse) present. Mouse movement was video-tracked using Ethovision

software 11.5 (Noldus) and the time spent in nose-to-nose, nose-to-body, and nose-to-genitals

interactions was measured.

Statistics

Statistical analyses were conducted with GraphPad Prism (GraphPad software 5.01). Shapiro-

Wilk normality test was used to determine the normality of distributions. We used a Two-tailed

Mann-Whitney U test or Two-tailed unpaired t-test for comparison between two independent

groups, a Wilcoxon matched pairs signed rank test or Two-tailed paired t-test for comparison

between two matched data, a Two-tailed Kruskal-Wallis test followed by a Dunn's multiple

comparison or a two-way ANOVA followed by a Bonferroni's multiple comparison for

comparison between three or more independent groups, and a Fisher exact test for nominal

comparison between two independent groups. The effect of tetanic stimulation of the slope of

field EPSPs was analyzed using repeated measure ANOVA. The survival analysis was performed

using a Kaplan-Meier log-rank test. Possible outliers within an experimental group were

identified with Grubb's test and discarded from the final analysis. To ensure the consistency and

reproducibility of our results, we conducted repeated trials in different acute hippocampal slices

prepared from at least three different animals from three different littermates. All data are

expressed as mean $\pm$ standard error to the mean (S.E.M.). For results expressed as percent of WT

(i.e. serum and mRNA leptin, Figure 1, all values (WT and $Mecp2^{-/y}$ mice) were normalized to

the mean WT value. In the figures, box plots represent the 1^{rst} and 3^{rd} quartiles; whiskers show

data range; horizontal lines show the median.

ACKNOWLEDGMENTS

We thank the members of the Molecular and Cellular Biology facility and Animal facility at INMED. We thank Drs F. Muscatelli and I. Medyna for critical reading of the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY

All datasets generated for this study are available on request to the corresponding author.

AUTHOR CONTRIBUTIONS

J-LG, YB, CM and GAW conceived the experiments and wrote the manuscript. YB, J-LG, DD, MB-H and VV performed the experiments. CS performed the behavioral analysis. J-CG bred the colony. All authors approved the final version of the manuscript.

FUNDING

This work was supported by the National Institute of Health (Grant 1R01HD092396), the Association Française du Syndrome de Rett, INSERM Transfert, Fondation Lejeune, and the Excellence Initiative of Aix-Marseille University-A*midex, a french "Investissements d'Avenir programme", MarMara AMX-19-IET-007.

    FIGURE LEGENDS

    Figure 1: Circulating leptin levels and leptin mRNA expressions are elevated in
    symptomatic *Mecp2*$^{-/y}$ mice.

    A) Box plots of serum leptin levels determined in blood samples taken at postnatal day (P) 15, 30
    and 50, using ELISA kit and expressed as percentage of age matched wild type (WT) mice mean.
    B) Scatter plots of serum leptin levels as a function of mice body weight of P30-P50 WT and
    Mecp2$^{-/y}$ mice. **C)** Box plots of normalized Leptin mRNA expression in gastrocnemius muscle of
    P30 and P50 WT and *Mecp2*$^{-/y}$ littermates. **D, E)** Box plots of normalized Leptin mRNA
    expression in the visceral (V) and inguinal (In) white adipose tissue (WAT) of P30 and P50 WT
    and *Mecp2*$^{-/y}$ littermates. In this and following figures, box plots represent quartiles, whiskers
    show data range, lozenges represent arithmetic means. Numbers in parentheses indicate the
    number of mice used. *P < 0.05, **P < 0.01, ***P < 0.001, two-tailed unpaired student *t*-test.

    Figure 2: Leptin antagonism prevents the progression of breathing deficits of symptomatic
    ***Mecp2*$^{-/y}$ mice.**
    A) Box plots of apnea frequency in P40 untreated WT and *Mecp2*$^{-/y}$ mice. **B, C)** Plots of apnea
    frequency in untreated and treated WT (**B**) and *Mecp2*$^{-/y}$ (**C**) mice. Lines indicate individual mice
    recorded the day before the treatment (P40) and at the end of the treatment (P50). Connected
    scares indicate mean±SEM. **D-F)** Box plots of the percentage of change of apnea frequency (**D**),
    breathing irregularity score (**E**) and minute ventilation (**F**) in sham- and treated- WT and *Mecp2*$^{-/y}$
    mice. Numbers in parenthesis indicate the number of mice used. **P < 0.01; ***P < 0.001, Two-
    tailed paired t-test (B, C), Two-tailed unpaired t-test (D-F), and Two-way ANOVA followed by a
    Bonferroni's multiple comparison (D-F).

    Figure 3: Leptin antagonism ameliorates the general health and lifespan of symptomatic
    ***Mecp2*$^{-/y}$ mice.**
    A) Box plots of the body weight of P40 WT and *Mecp2*$^{-/y}$ mice. **B)** Body weight change (% of
    P40) as a function of age in sham-, leptin- and anti-leptin treated *Mecp2*$^{-/y}$ mice. The treatment
    started at P40 and ended at P50. **C)** Box plot of the body weight change (% of P40) at P50 in
    sham and treated mice at the indicated genotype and treatment. **D)** Severity score of P40 WT and
    Mecp2$^{-/y}$ mice. mean±SEM. **E)** Plots of the severity score in sham- and anti-leptin treated *Mecp2*$^{-/y}$
    mice. Connected points indicate individual mice recorded the day before the treatment (P40)
    and at the end of the treatment (P50). Connected squares indicate mean±SEM. **F)** Radar plot of
    the symptoms scored in sham and anti-leptin treated *Mecp2*$^{-/y}$ mice. **G)** Plot of the percentage of
    surviving mice as a function of age. The treatment started at P40 and was maintained until the
    death of the animal. Numbers in parenthesis indicate the number of mice used. *P < 0.05; **P <
    0.01; ***P < 0.001, Two-way ANOVA followed by a Bonferroni's multiple comparison (C),
    two-tailed unpaired *t*-test (A, C,), two-tailed paired *t*-test (D, E).

Figure 4: Leptin antagonism prevents the worsening of locomotor deficit but does not improve motor coordination nor anxiety alterations of symptomatic *Mecp2*$^{-/y}$ mice.

A) Box plots of the distance traveled by P40 WT and *Mecp2*$^{-/y}$ mice in an open field. **B**) Plots of distance traveled by sham and anti-leptin *Mecp2*$^{-/y}$ mice in an open field. Connected points indicate individual mice recorded the day before the treatment (P40) and at the end of the treatment (P50). Connected squares indicate mean±SEM. **C**) Box plots of the distance traveled in an open field by P50 sham and treated mice at the indicated genotype and treatment. **D**) Mean ± SEM plots of the latency to fall (best performance of 3 trials) of P40 WT and *Mecp2*$^{-/y}$ mice in the accelerating rotarod. **E**) Plots of latency to fall (best performance) by sham and anti-leptin *Mecp2*$^{-/y}$ mice. Connected points indicate individual mice recorded the day before the treatment (P40) and at the end of the treatment (P50). Connected squares indicate mean±SEM. **F**) Box plots of the latency to fall (best performance) by P50 sham and treated mice at the indicated genotype and treatment. **G, H**) Box plots of the time spent in open arms (**G**) and dipping (**H**) by P40 WT and *Mecp2*$^{-/y}$ mice, and P50 WT- and *Mecp2*$^{-/y}$-sham and treated mice in an elevated plus maze. mean±SEM. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$, two-tailed unpaired *t*-test (A), two-tailed paired *t*-test (B), Wilcoxon matched pairs rank test (E), Mann Whitney test (D), Two-way ANOVA followed by a Bonferroni's multiple comparison (C, F, G, H).

Figure 5: Leptin antagonism during the symptomatic stage does not improve seizure susceptibility of *Mecp2*$^{-/y}$ mice.

A, B) Box plots of tonic-clonic seizures onset latency of P50 untreated (**A**) and treated (**B**) WT and *Mecp2*$^{-/y}$ mice. **C, D**) Flurothyl-induced lethality of P50 untreated (**C**) and treated (**D**) WT and *Mecp2*$^{-/y}$ mice. Numbers in parenthesis indicate the number of mice used. **P<0.05, ***$P < 0.001$, Two-tailed unpaired *t*-test (**A, B**), Fisher exact test (**C**).

Figure 6: Leptin contributes to the E/I imbalance in the hippocampus of symptomatic *Mecp2*$^{-/y}$ mice.

A) Examples of whole-cell recordings performed on CA3 pyramidal neurons at a holding potential of −45 and -70 mV at P50. The glutamatergic synaptic currents are inwards and the GABAergic synaptic currents are outwards. **B-D)** Box plots of the ratio of spontaneous glutamatergic and GABAergic postsynaptic currents (**B, C**) and frequency of spontaneous GABAergic postsynaptic currents (**D, E**) recorded from P50 untreated- (**B, D**) and treated- (**C, E**) WT and *Mecp2*$^{-/y}$ CA3 pyramidal neurons. Scale bars: 100pA, 2 sec. Numbers in parenthesis indicate the number of cells recorded and mice used. **$P < 0.01$; ***$P < 0.001$, Two-tailed Mann-Whitney test (**B, D**). Two-tailed Kruskal-Wallis test followed by Dunn's post hoc comparison (**C, E**).

Figure 7: Leptin contributes to hippocampal synaptic plasticity impairment of symptomatic *Mecp2*$^{-/y}$ mice.

LTP was recorded for 50 min following two trains of 100 Hz tetanic stimulation (1sec duration,
20 sec interval) at the Schaffer collateral-CA1 synapses of acute hippocampal slices from
symptomatic (P50) untreated and treated $Mecp2^{-/y}$ and WT mice. Insets show examples of fEPSP.
Scale bar: 100μV, 3ms. Numbers in parenthesis indicate the number of slices recorded and mice
used. *$P < 0.05$, Repeated measure ANOVA.

Figure 8: RTT-associated symptoms are improved in leptin haplo insufficient mice.

A) Serum leptin levels in P50 WT, $Mecp2^{-/y}$ and $Mecp2^{-/y;ob/+}$ littermates. **B,C**) Box plots of the
frequency ratio of spontaneous glutamatergic and GABAergic postsynaptic currents (B) and
frequency of spontaneous GABAergic postsynaptic currents (C) recorded from CA3 pyramidal
neurons at P50. **D**) LTP at the Schaffer collateral-CA1 synapses of acute hippocampal slices
from symptomatic (P50-60) $Mecp2^{-/y}$ and $Mecp2^{-/y;ob/+}$ mice. **E,F**) Box plots of apnea frequency
(E) and body weight (F) in P50 WT, $Mecp2^{-/y}$ and $Mecp2^{-/y;ob/+}$ littermates. Numbers in
parenthesis indicate the number of mice used and number of cells recorded. *$P < 0.05$; **$P <$
0.01$; ***$P < 0.001$. Two-way ANOVA followed by a Bonferroni's multiple comparison (A, D,
E), two-tailed Kruskal-Wallis followed by a Dunn's multiple comparison (B,C), repeated
measure ANOVA (F).

REFERENCES

Acampa, M., Guideri, F., Hayek, J., Blardi, P., De Lalla, A., Zappella, M., et al. (2008). Sympathetic overactivity and plasma leptin levels in Rett syndrome. *Neurosci. Lett.* 432, 69–72. doi: 10.1016/j.neulet.2007.12.030.

Ahima, R. S., and Flier, J. S. (2000). Leptin. *Annual Review of Physiology* 62, 413–437. doi: 10.1146/annurev.physiol.62.1.413.

Asaka, Y., Jugloff, D. G. M., Zhang, L., Eubanks, J. H., and Fitzsimonds, R. M. (2006). Hippocampal synaptic plasticity is impaired in the Mecp2-null mouse model of Rett syndrome. *Neurobiol Dis* 21, 217–227. doi: 10.1016/j.nbd.2005.07.005.

Ashwood, P., Kwong, C., Hansen, R., Hertz-Picciotto, I., Croen, L., Krakowiak, P., et al. (2008). Brief report: plasma leptin levels are elevated in autism: association with early onset phenotype? *J Autism Dev Disord* 38, 169–175. doi: 10.1007/s10803-006-0353-1.

Bassi, M., Furuya, W. I., Menani, J. V., Colombari, D. S. A., do Carmo, J. M., da Silva, A. A., et al. (2014). Leptin into the ventrolateral medulla facilitates chemorespiratory response in leptin-deficient (ob/ob) mice. *Acta Physiol (Oxf)* 211, 240–248. doi: 10.1111/apha.12257.

Bassi, M., Nakamura, N. B., Furuya, W. I., Colombari, D. S. A., Menani, J. V., do Carmo, J. M., et al. (2015). Activation of the brain melanocortin system is required for leptin-induced modulation of chemorespiratory function. *Acta Physiol (Oxf)* 213, 893–901. doi: 10.1111/apha.12394.

Belcastro, V., D'Egidio, C., Striano, P., and Verrotti, A. (2013). Metabolic and endocrine effects of valproic acid chronic treatment. *Epilepsy Res* 107, 1–8. doi: 10.1016/j.eplepsyres.2013.08.016.

Bertoni, A., Schaller, F., Tyzio, R., Gaillard, S., Santini, F., Xolin, M., et al. (2021). Oxytocin administration in neonates shapes hippocampal circuitry and restores social behavior in a mouse model of autism. *Mol Psychiatry*. doi: 10.1038/s41380-021-01227-6.

Blardi, P., de Lalla, A., Ceccatelli, L., Vanessa, G., Auteri, A., and Hayek, J. (2010). Variations of plasma leptin and adiponectin levels in autistic patients. *Neurosci. Lett.* 479, 54–57. doi: 10.1016/j.neulet.2010.05.027.

Blardi, P., de Lalla, A., D'Ambrogio, T., Vonella, G., Ceccatelli, L., Auteri, A., et al. (2009). Long-term plasma levels of leptin and adiponectin in Rett syndrome. *Clin. Endocrinol. (Oxf)* 70, 706–709. doi: 10.1111/j.1365-2265.2008.03386.x.

Blardi, P., de Lalla, A., D'Ambrogio, T., Zappella, M., Cevenini, G., Ceccatelli, L., et al. (2007). Rett syndrome and plasma leptin levels. *J Pediatr* 150, 37–39. doi: 10.1016/j.jpeds.2006.10.061.

Bouret, S. G., Draper, S. J., and Simerly, R. B. (2004). Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science* 304, 108–110. doi: 10.1126/science.1095004.

Calfa, G., Li, W., Rutherford, J. M., and Pozzo-Miller, L. (2015). Excitation/inhibition imbalance and impaired synaptic inhibition in hippocampal area CA3 of Mecp2 knockout mice. *Hippocampus* 25, 159–168. doi: 10.1002/hipo.22360.

Caron, E., Sachot, C., Prevot, V., and Bouret, S. G. (2010). Distribution of leptin-sensitive cells in the
postnatal and adult mouse brain. *J. Comp. Neurol.* 518, 459–476. doi: 10.1002/cne.22219.

Chahrour, M., and Zoghbi, H. Y. (2007). The story of Rett syndrome: from clinic to neurobiology. *Neuron*
56, 422–437. doi: 10.1016/j.neuron.2007.10.001.

Chapnik, N., Solomon, G., Genzer, Y., Miskin, R., Gertler, A., and Froy, O. (2013). A superactive leptin
antagonist alters metabolism and locomotion in high-leptin mice. *J Endocrinol* 217, 283–290. doi:
10.1530/JOE-13-0033.

Chen, C.-Y., Di Lucente, J., Lin, Y.-C., Lien, C.-C., Rogawski, M. A., Maezawa, I., et al. (2018).
Defective GABAergic neurotransmission in the nucleus tractus solitarius in Mecp2-null mice, a model of
Rett syndrome. *Neurobiol Dis* 109, 25–32. doi: 10.1016/j.nbd.2017.09.006.

Cheval, H., Guy, J., Merusi, C., De Sousa, D., Selfridge, J., and Bird, A. (2012). Postnatal inactivation
reveals enhanced requirement for MeCP2 at distinct age windows. *Hum Mol Genet* 21, 3806–3814. doi:
10.1093/hmg/dds208.

Conti, V., Gandaglia, A., Galli, F., Tirone, M., Bellini, E., Campana, L., et al. (2015). MeCP2 Affects
Skeletal Muscle Growth and Morphology through Non Cell-Autonomous Mechanisms. *PLoS One* 10,
e0130183. doi: 10.1371/journal.pone.0130183.

Cortelazzo, A., De Felice, C., Guerranti, R., Signorini, C., Leoncini, S., Pecorelli, A., et al. (2014).
Subclinical inflammatory status in Rett syndrome. *Mediators Inflamm* 2014, 480980. doi:
10.1155/2014/480980.

Davis, J. F., Choi, D. L., and Benoit, S. C. (2010). Insulin, leptin and reward. *Trends Endocrinol Metab*
21, 68–74. doi: 10.1016/j.tem.2009.08.004.

D'Cruz, J. A., Wu, C., Zahid, T., El-Hayek, Y., Zhang, L., and Eubanks, J. H. (2010). Alterations of
cortical and hippocampal EEG activity in MeCP2-deficient mice. *Neurobiol Dis* 38, 8–16. doi:
10.1016/j.nbd.2009.12.018.

Dhar, M., Wayman, G. A., Zhu, M., Lambert, T. J., Davare, M. A., and Appleyard, S. M. (2014). Leptin-
induced spine formation requires TrpC channels and the CaM kinase cascade in the hippocampus. *J.*
Neurosci. 34, 10022–10033. doi: 10.1523/JNEUROSCI.2868-13.2014.

Dumon, C., Diabira, D., Chudotvorova, I., Bader, F., Sahin, S., Zhang, J., et al. (2018). The adipocyte
hormone leptin sets the emergence of hippocampal inhibition in mice. *Elife* 7, e36726. doi:
10.7554/eLife.36726.

Ehinger, Y., Matagne, V., Villard, L., and Roux, J.-C. (2018). Rett syndrome from bench to bedside:
recent advances. *F1000Res* 7, 398. doi: 10.12688/f1000research.14056.1.

Elinav, E., Niv-Spector, L., Katz, M., Price, T. O., Ali, M., Yacobovitz, M., et al. (2009). Pegylated leptin
antagonist is a potent orexigenic agent: preparation and mechanism of activity. *Endocrinology* 150, 3083–
3091. doi: 10.1210/en.2008-1706.

Farr, O. M., Tsoukas, M. A., and Mantzoros, C. S. (2015). Leptin and the brain: influences on brain
development, cognitive functioning and psychiatric disorders. *Metabolism* 64, 114–130. doi:
10.1016/j.metabol.2014.07.004.

Fisch, S., Bachner-Hinenzon, N., Ertracht, O., Guo, L., Arad, Y., Ben-Zvi, D., et al. (2020). Localized
Antileptin Therapy Prevents Aortic Root Dilatation and Preserves Left Ventricular Systolic Function in a
Murine Model of Marfan Syndrome. *J Am Heart Assoc* 9, e014761. doi: 10.1161/JAHA.119.014761.

Fukuhara, S., Nakajima, H., Sugimoto, S., Kodo, K., Shigehara, K., Morimoto, H., et al. (2019). High-fat
diet accelerates extreme obesity with hyperphagia in female heterozygous Mecp2-null mice. *PLoS One* 14,
e0210184. doi: 10.1371/journal.pone.0210184.

Gauda, E. B., Conde, S., Bassi, M., Zoccal, D. B., Almeida Colombari, D. S., Colombari, E., et al. (2020).
Leptin: Master Regulator of Biological Functions that Affects Breathing. *Compr Physiol* 10, 1047–1083.
doi: 10.1002/cphy.c190031.

Gertler, A., and Elinav, E. (2014). Novel superactive leptin antagonists and their potential therapeutic
applications. *Curr Pharm Des* 20, 659–665. doi: 10.2174/13816128113199990014.

Glaze, D. G. (2002). Neurophysiology of Rett syndrome. *Mental Retardation and Developmental
Disabilities Research Reviews* 8, 66–71. doi: 10.1002/mrdd.10024.

Gonzalez, A., Cheung, W. W., Perens, E. A., Oliveira, E. A., Gertler, A., and Mak, R. H. (2021). A Leptin
Receptor Antagonist Attenuates Adipose Tissue Browning and Muscle Wasting in Infantile Nephropathic
Cystinosis-Associated Cachexia. *Cells* 10, 1954. doi: 10.3390/cells10081954.

Greco, S. J., Bryan, K. J., Sarkar, S., Zhu, X., Smith, M. A., Ashford, J. W., et al. (2010). Leptin reduces
pathology and improves memory in a transgenic mouse model of Alzheimer's disease. *J Alzheimers Dis*
19, 1155–1167. doi: 10.3233/JAD-2010-1308.

Grillo, C. A., Piroli, G. G., Evans, A. N., Macht, V. A., Wilson, S. P., Scott, K. A., et al. (2011).
Obesity/hyperleptinemic phenotype adversely affects hippocampal plasticity: effects of dietary restriction.
Physiol Behav 104, 235–241. doi: 10.1016/j.physbeh.2010.10.020.

Guimond, D., Diabira, D., Porcher, C., Bader, F., Ferrand, N., Zhu, M., et al. (2014). Leptin potentiates
GABAergic synaptic transmission in the developing rodent hippocampus. *Frontiers in Cellular
Neuroscience* 8, 235. doi: 10.3389/fncel.2014.00235.

Guo, M., Huang, T.-Y., Garza, J. C., Chua, S. C., and Lu, X.-Y. (2013). Selective deletion of leptin
receptors in adult hippocampus induces depression-related behaviours. *Int. J. Neuropsychopharmacol.* 16,
857–867. doi: 10.1017/S1461145712000703.

Guo, M., Lu, Y., Garza, J. C., Li, Y., Chua, S. C., Zhang, W., et al. (2012). Forebrain glutamatergic
neurons mediate leptin action on depression-like behaviors and synaptic depression. *Transl Psychiatry* 2,
e83. doi: 10.1038/tp.2012.9.

Guy, J., Gan, J., Selfridge, J., Cobb, S., and Bird, A. (2007). Reversal of neurological defects in a mouse
model of Rett syndrome. *Science* 315, 1143–1147. doi: 10.1126/science.1138389.

Guy, J., Hendrich, B., Holmes, M., Martin, J. E., and Bird, A. (2001). A mouse Mecp2-null mutation
causes neurological symptoms that mimic Rett syndrome. *Nat Genet* 27, 322–326. doi: 10.1038/85899.

Harvey, J. (2003). Leptin: a multifaceted hormone in the central nervous system. *Mol Neurobiol* 28, 245–
258. doi: 10.1385/MN:28:3:245.

Harvey, J. (2022). Food for Thought: Leptin and Hippocampal Synaptic Function. *Front Pharmacol* 13,
882158. doi: 10.3389/fphar.2022.882158.

Harvey, J., Solovyova, N., and Irving, A. (2006). Leptin and its role in hippocampal synaptic plasticity.
Prog Lipid Res 45, 369–378. doi: 10.1016/j.plipres.2006.03.001.

Inyushkin, A. N., Inyushkina, E. M., and Merkulova, N. A. (2009). Respiratory responses to
microinjections of leptin into the solitary tract nucleus. *Neurosci Behav Physiol* 39, 231–240. doi:
10.1007/s11055-009-9124-8.

Katz, D. M., Dutschmann, M., Ramirez, J.-M., and Hilaire, G. (2009). Breathing disorders in Rett
syndrome: progressive neurochemical dysfunction in the respiratory network after birth. *Respir Physiol*
Neurobiol 168, 101–108. doi: 10.1016/j.resp.2009.04.017.

Kerr, B., Silva, P. A., Walz, K., and Young, J. I. (2010). Unconventional Transcriptional Response to
Environmental Enrichment in a Mouse Model of Rett Syndrome. *PLOS ONE* 5, e11534. doi:
10.1371/journal.pone.0011534.

Lewis, J. D., Meehan, R. R., Henzel, W. J., Maurer-Fogy, I., Jeppesen, P., Klein, F., et al. (1992).
Purification, sequence, and cellular localization of a novel chromosomal protein that binds to methylated
DNA. *Cell* 69, 905–914. doi: 10.1016/0092-8674(92)90610-o.

Li, L.-L., Jin, M.-F., and Ni, H. (2018). Zinc/CaMK II Associated-Mitophagy Signaling Contributed to
Hippocampal Mossy Fiber Sprouting and Cognitive Deficits Following Neonatal Seizures and Its
Regulation by Chronic Leptin Treatment. *Front Neurol* 9, 802. doi: 10.3389/fneur.2018.00802.

Li, W. (2022). Excitation and Inhibition Imbalance in Rett Syndrome. *Front Neurosci* 16, 825063. doi:
10.3389/fnins.2022.825063.

Lim, G., Wang, S., Zhang, Y., Tian, Y., and Mao, J. (2009). Spinal leptin contributes to the pathogenesis
of neuropathic pain in rodents. *J Clin Invest* 119, 295–304. doi: 10.1172/JCI36785.

Liu, C., Wang, J., Wei, Y., Zhang, W., Geng, M., Yuan, Y., et al. (2019). Fat-Specific Knockout of Mecp2
Upregulates Slpi to Reduce Obesity by Enhancing Browning. *Diabetes* 69, 35–47. doi: 10.2337/db19-
0502.

Lozovaya, N., Nardou, R., Tyzio, R., Chiesa, M., Pons-Bennaceur, A., Eftekhari, S., et al. (2019). Early
alterations in a mouse model of Rett syndrome: the GABA developmental shift is abolished at birth. *Sci*
Rep 9, 9276. doi: 10.1038/s41598-019-45635-9.

McGregor, G., and Harvey, J. (2019). Leptin Regulation of Synaptic Function at Hippocampal TA-CA1
and SC-CA1 Synapses: Implications for Health and Disease. *Neurochem Res* 44, 650–660. doi:
10.1007/s11064-017-2362-1.

Medrihan, L., Tantalaki, E., Aramuni, G., Sargsyan, V., Dudanova, I., Missler, M., et al. (2008). Early
defects of GABAergic synapses in the brain stem of a MeCP2 mouse model of Rett syndrome. *J*
Neurophysiol 99, 112–121. doi: 10.1152/jn.00826.2007.

Mora-Muñoz, L., Guerrero-Naranjo, A., Rodríguez-Jimenez, E. A., Mastronardi, C. A., and Velez-van-
Meerbeke, A. (2018). Leptin: role over central nervous system in epilepsy. *BMC Neurosci* 19, 51. doi:
10.1186/s12868-018-0453-9.

Moretti, P., Levenson, J. M., Battaglia, F., Atkinson, R., Teague, R., Antalffy, B., et al. (2006). Learning
and memory and synaptic plasticity are impaired in a mouse model of Rett syndrome. *J Neurosci* 26, 319–
327. doi: 10.1523/JNEUROSCI.2623-05.2006.

Motil, K. J., Barrish, J. O., Lane, J., Geerts, S. P., Annese, F., McNair, L., et al. (2011). Vitamin D
deficiency is prevalent in girls and women with Rett syndrome. *J Pediatr Gastroenterol Nutr* 53, 569–
574. doi: 10.1097/MPG.0b013e3182267a66.

Moult, P. R., and Harvey, J. (2008). Hormonal regulation of hippocampal dendritic morphology and
synaptic plasticity. *Cell Adh Migr* 2, 269–275. doi: 10.4161/cam.2.4.6354.

Moult, P. R., and Harvey, J. (2009). Regulation of glutamate receptor trafficking by leptin. *Biochem Soc*
Trans 37, 1364–1368. doi: 10.1042/BST0371364.

Neyens, D., Zhao, H., Huston, N. J., Wayman, G. A., Ritter, R. C., and Appleyard, S. M. (2020). Leptin
Sensitizes NTS Neurons to Vagal Input by Increasing Postsynaptic NMDA Receptor Currents. *J Neurosci*
40, 7054–7064. doi: 10.1523/JNEUROSCI.1865-19.2020.

O'Malley, D., MacDonald, N., Mizielinska, S., Connolly, C. N., Irving, A. J., and Harvey, J. (2007).
Leptin promotes rapid dynamic changes in hippocampal dendritic morphology. *Mol Cell Neurosci* 35,
559–572. doi: 10.1016/j.mcn.2007.05.001.

Oomura, Y., Hori, N., Shiraishi, T., Fukunaga, K., Takeda, H., Tsuji, M., et al. (2006). Leptin facilitates
learning and memory performance and enhances hippocampal CA1 long-term potentiation and CaMK II
phosphorylation in rats. *Peptides* 27, 2738–2749. doi: 10.1016/j.peptides.2006.07.001.

Park, M. J., Aja, S., Li, Q., Degano, A. L., Penati, J., Zhuo, J., et al. (2014). Anaplerotic triheptanoin diet
enhances mitochondrial substrate use to remodel the metabolome and improve lifespan, motor function,
and sociability in MeCP2-null mice. *PLoS One* 9, e109527. doi: 10.1371/journal.pone.0109527.

Pinto, S., Roseberry, A. G., Liu, H., Diano, S., Shanabrough, M., Cai, X., et al. (2004). Rapid rewiring of
arcuate nucleus feeding circuits by leptin. *Science* 304, 110–115. doi: 10.1126/science.1089459.

Raghavan, R., Zuckerman, B., Hong, X., Wang, G., Ji, Y., Paige, D., et al. (2018). Fetal and Infancy
Growth Pattern, Cord and Early Childhood Plasma Leptin, and Development of Autism Spectrum
Disorder in the Boston Birth Cohort. *Autism Res* 11, 1416–1431. doi: 10.1002/aur.2011.

Ramirez, J.-M., Ward, C. S., and Neul, J. L. (2013). Breathing challenges in Rett syndrome: lessons
learned from humans and animal models. *Respir Physiol Neurobiol* 189, 280–287. doi:
10.1016/j.resp.2013.06.022.

Ricceri, L., De Filippis, B., and Laviola, G. (2008). Mouse models of Rett syndrome: from behavioural
phenotyping to preclinical evaluation of new therapeutic approaches. *Behav Pharmacol* 19, 501–517. doi:
10.1097/FBP.0b013e32830c3645.

Sahin, G. S., Dhar, M., Dillon, C., Zhu, M., Shiina, H., Winters, B. D., et al. (2020). Leptin stimulates
synaptogenesis in hippocampal neurons via KLF4 and SOCS3 inhibition of STAT3 signaling. *Mol Cell*
Neurosci 106, 103500. doi: 10.1016/j.mcn.2020.103500.

Sahin, G. S., Luis Rodriguez-Llamas, J., Dillon, C., Medina, I., Appleyard, S. M., Gaiarsa, J.-L., et al.
(2021). Leptin increases GABAergic synaptogenesis through the Rho guanine exchange factor β-PIX in
developing hippocampal neurons. *Sci Signal* 14, eabe4111. doi: 10.1126/scisignal.abe4111.

Sandweiss, A. J., Brandt, V. L., and Zoghbi, H. Y. (2020). Advances in understanding of Rett syndrome
and MECP2 duplication syndrome: prospects for future therapies. *Lancet Neurol* 19, 689–698. doi:
10.1016/S1474-4422(20)30217-9.

Scaramuzza, L., De Rocco, G., Desiato, G., Cobolli Gigli, C., Chiacchiaretta, M., Mirabella, F., et al.
(2021). The enhancement of activity rescues the establishment of Mecp2 null neuronal phenotypes. *EMBO*
Mol Med, e12433. doi: 10.15252/emmm.202012433.

Sciorati, C., Clementi, E., Manfredi, A. A., and Rovere-Querini, P. (2015). Fat deposition and
accumulation in the damaged and inflamed skeletal muscle: cellular and molecular players. *Cell Mol Life*
Sci 72, 2135–2156. doi: 10.1007/s00018-015-1857-7.

Shanley, L. J., Irving, A. J., Rae, M. G., Ashford, M. L. J., and Harvey, J. (2002). Leptin inhibits rat
hippocampal neurons via activation of large conductance calcium-activated K+ channels. *Nat Neurosci* 5,
299–300. doi: 10.1038/nn824.

Signore, A. P., Zhang, F., Weng, Z., Gao, Y., and Chen, J. (2008). Leptin neuroprotection in the CNS:
mechanisms and therapeutic potentials. *J. Neurochem.* 106, 1977–1990. doi: 10.1111/j.1471-
4159.2008.05457.x.

Singh, U. P., Singh, N. P., Guan, H., Busbee, B., Price, R. L., Taub, D. D., et al. (2013). Leptin antagonist
ameliorates chronic colitis in IL-10$^{-/-}$ mice. *Immunobiology* 218, 1439–1451. doi:
10.1016/j.imbio.2013.04.020.

Solovyova, N., Moult, P. R., Milojkovic, B., Lambert, J. J., and Harvey, J. (2009). Bi-directional
modulation of fast inhibitory synaptic transmission by leptin. *J. Neurochem.* 108, 190–201. doi:
10.1111/j.1471-4159.2008.05751.x.

Stagi, S., Cavalli, L., Congiu, L., Scusa, M. F., Ferlini, A., Bigoni, S., et al. (2015). Thyroid function in
Rett syndrome. *Horm Res Paediatr* 83, 118–125. doi: 10.1159/000370066.

Torres-Andrade, R., Moldenhauer, R., Gutierrez-Bertín, N., Soto-Covasich, J., Mancilla-Medina, C.,
Ehrenfeld, C., et al. (2014). The increase in body weight induced by lack of methyl CpG binding protein-2
is associated with altered leptin signalling in the hypothalamus. *Exp. Physiol.* 99, 1229–1240. doi:
10.1113/expphysiol.2014.079798.

Valladolid-Acebes, I., Merino, B., Principato, A., Fole, A., Barbas, C., Lorenzo, M. P., et al. (2012). High-
fat diets induce changes in hippocampal glutamate metabolism and neurotransmission. *Am J Physiol*
Endocrinol Metab 302, E396-402. doi: 10.1152/ajpendo.00343.2011.

Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., and Lowell, B. B. (2011). Leptin action on GABAergic
neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron* 71, 142–154. doi:
10.1016/j.neuron.2011.05.028.

Wang, X., Lacza, Z., Sun, Y. E., and Han, W. (2014). Leptin resistance and obesity in mice with deletion
of methyl-CpG-binding protein 2 (MeCP2) in hypothalamic pro-opiomelanocortin (POMC) neurons.
Diabetologia 57, 236–245. doi: 10.1007/s00125-013-3072-0.

Williams, K. W., and Smith, B. N. (2006). Rapid inhibition of neural excitability in the nucleus tractus
solitarii by leptin: implications for ingestive behaviour. *J Physiol* 573, 395–412. doi:
10.1113/jphysiol.2006.106336.

Yu, H., Shi, L., Chen, J., Jun, S., Hao, Y., Wang, S., et al. (2022). A Neural Circuit Mechanism
Controlling Breathing by Leptin in the Nucleus Tractus Solitarii. *Neurosci Bull* 38, 149–165. doi:
10.1007/s12264-021-00742-4.

Zuure, W. A., Roberts, A. L., Quennell, J. H., and Anderson, G. M. (2013). Leptin signaling in GABA
neurons, but not glutamate neurons, is required for reproductive function. *J Neurosci* 33, 17874–17883.
doi: 10.1523/JNEUROSCI.2278-13.2013.