Root cap cell corpse clearance limits microbial colonization in *Arabidopsis thaliana*

Nyasha M. Charura[1+], Ernesto Llamas[1+], Concetta De Quattro[1], David Vilchez[2,3,4], Moritz

K. Nowack[5,6#] and Alga Zuccaro[1#*]

[1] Cluster of Excellence on Plant Sciences (CEPLAS), Institute for Plant Sciences,

University of Cologne, D-50674 Cologne, Germany.

[2] Cologne Excellence Cluster for Cellular Stress Responses in Aging-Associated

Diseases (CECAD), University of Cologne, 50931 Cologne, Germany.

[3] Center for Molecular Medicine Cologne (CMMC), University of Cologne, 50931 Cologne,

Germany.

[4] Faculty of Medicine, University Hospital Cologne, 50931 Cologne, Germany.

[5] Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent,

Belgium

[6] VIB Center for Plant Systems Biology, 9052 Ghent, Belgium

[#]shared last author

[+]shared first author

*Corresponding author: azuccaro@uni-koeln.de

Abstract

Programmed cell death occurring during plant development (dPCD) is a fundamental

process integral for plant growth and reproduction. Here, we investigate the connection

between developmentally controlled PCD and fungal accommodation in *Arabidopsis*

thaliana roots, focusing on the root cap-specific transcription factor

ANAC033/SOMBRERO (SMB) and the senescence-associated nuclease BFN1.

Mutations of both dPCD regulators increase colonization by the beneficial fungus

Serendipita indica, primarily in the differentiation zone. *smb-3* mutants additionally exhibit

hypercolonization around the meristematic zone and a delay of *S. indica*-induced root-

growth promotion. This demonstrates that root cap dPCD and rapid post-mortem

clearance of cellular corpses represent a physical defense mechanism restricting

microbial invasion of the root. Additionally, reporter lines and transcriptional analysis

revealed that *BFN1* expression is downregulated during *S. indica* colonization in mature

root epidermal cells, suggesting a transcriptional control mechanism that facilitates the accommodation of beneficial microbes in the roots.

Key findings in bullet points:

- The process of programmed cell death in root development (dPCD) influences the extent and outcomes of fungal symbiosis
- Fungal colonization of the root tip and differentiation zone is restricted by SMB-mediated clearance of dead cells, which preserves the meristem and regulates symbiosis
- Expression of plant nuclease *BFN1*, which is associated with senescence, is modulated to facilitate root accommodation of beneficial microbes

Keywords

regulated cell death (RCD) / developmentally programmed cell death (dPCD) plant-microbe interaction / growth promotion / manipulation of dPCD / SMB / meristematic stem cells / BFN1 / aging cells

Introduction

Plant roots elongate by producing new cells in the root apical meristem at the root tip. As roots extend further into the soil environment, the newly formed root tissue is naturally exposed to microbial attack, challenging the successful establishment of root systems. However, microbial colonization at the meristematic zone is rarely detected (Deshmukh et al., 2006, Jacobs et al., 2011). The sensitive tissue of meristematic stem cells is surrounded by the root cap, a specialized root organ that orchestrates root architecture, directs root growth based on gravitropism and hydrotropism, and senses environmental stimuli. In addition, the root cap is presumed to have a protective function in soil exploration (Kumpf and Nowack, 2015, Moriwaki et al., 2013).

In *Arabidopsis thaliana* (hereafter Arabidopsis), the root cap consists of two distinct tissues: the centrally located columella root cap at the very root tip and the peripherally located lateral root cap (LRC), which flanks both the columella and the entire root meristem (Dolan et al., 1993). A ring of specific stem cells continuously generates both

new LRC cells and root epidermal cells (Dolan et al., 1993). However, despite the constant production of LRC cells, the root cap itself does not grow in size but matches the size of the meristem (Fendrych et al., 2014, Barlow, 2002, Kumpf and Nowack, 2015). To maintain size homeostasis, root cap development is a highly regulated process that varies in different plant species. In Arabidopsis, a combination of dPCD and shedding of old cells into the rhizosphere has been described (Bennett et al., 2010, Kumpf and Nowack, 2015). The centrally located columella root cap along with adjacent proximal LRC cells is shed as a cell package, followed by a PCD process (Shi et al., 2018, Feng et al., 2022, Huysmans et al., 2018). In contrast, LRC cells at the distal end of the root tip elongate and reach the edge of the meristematic zone where they undergo dPCD followed by corpse clearance on the root surface, orchestrated as part of a terminal differentiation program by the root cap-specific transcription factor ANAC33/SOMBRERO (SMB) (Fendrych et al., 2014, Willemsen et al., 2008, Bennett et al., 2010). SMB belongs to a plant-specific family of transcription factors carrying a NAC domain (NAM - no apical meristem; ATAF1 and -2, and CUC2 - cup-shaped cotyledon). SMB promotes the expression of genes associated with the initiation and execution of LRC cell death, including the senescence-associated bifunctional nuclease *BFN1* and the putative aspartic protease *PASPA3* (Fendrych et al., 2014, Huysmans et al., 2018). BFN1 localizes in the ER, but upon cell death the protein is released and its nuclease activity ensures rapid and irreversible degradation of RNA and DNA in the nucleus and cytoplasm as part of a rapid cell-autonomous corpse clearance at the root surface (Reza et al., 2018, Fendrych et al., 2014, Farage-Barhom et al., 2011). Accordingly, DNA and RNA fragmentation in *bfn1-1* loss-of-function mutants is delayed (Fendrych et al., 2014). Precise timing of cell death and elimination of LRC cells before they fully enter the elongation zone is essential for maintaining root cap size and optimal root growth (Fendrych et al., 2014). Loss of SMB activity results in a delayed cell death, causing LRC cells to enter the elongation zone where they eventually die without expression of dPCD executor genes in the root cap (Fendrych et al., 2014). Interestingly, the aberrant cell death of LRC cells in the elongation zone of *smb-3* mutants is not followed by corpse clearance, resulting in an accumulation of uncleared cell corpses along the entire root surface (Fendrych et al., 2014).

Despite its importance in root morphology and plant development, little is known about
the importance of dPCD and rapid cell corpse clearance on plant-microbe interactions.
To address this question, we tested two well-characterized loss-of-function T-DNA
insertion lines, *smb-3* and *bfn1-1*, during colonization with *Serendipita indica*, a beneficial
fungus of the order Sebacinales. As a root endophyte, *S. indica* colonizes the epidermal
and cortex layers of a broad range of different plant hosts, conferring various beneficial
effects, including plant growth promotion, protection against pathogenic microbes and
increased tolerance to abiotic stresses (Boorboori and Zhang, 2022, Mahdi et al., 2022,
Fesel and Zuccaro, 2016). The colonization strategy of *S. indica* comprises an initial
biotrophic interaction, followed by a growth phase associated with a restricted host cell
death that does not, however, diminish the beneficial effects on the plant host. The
induction of restricted cell death in the epidermal and cortex layers is a crucial component
of the colonization strategy of *S. indica* and is accompanied by an increased production
of fungal hydrolytic enzymes (Zuccaro et al., 2011, Deshmukh et al., 2006). The switch
between the biotrophic and the cell death-associated phase can vary depending on the
host system and environmental conditions, but has been postulated to occur
approximately 6 to 8 days post inoculation (dpi) in Arabidopsis (Zuccaro et al., 2011).
Although several effector proteins involved in fungal accommodation have been
described (Dunken et al., 2022, Weiss et al., 2016, Akum et al., 2015, Wawra et al., 2016,
Nostadt et al., 2020, Nizam et al., 2019), the exact mechanism by which *S. indica*
manipulates host cell death and the role of dPCD in fungal accommodation in the roots
are largely unclear.

Here, we show that the accumulation of uncleared LRC cell corpses on the roots of *smb-*
3 mutants triggers hypercolonization by *S. indica*, especially around the meristematic
zone, and delays *S. indica*-induced root growth promotion. We propose that a tight
regulation of host dPCD and rapid and complete clearance of root cap cell corpses play
important roles in restricting fungal colonization at the root apical meristem. Furthermore,
we show that *S. indica* downregulates *BFN1* in older and differentiated epidermal cells to
promote fungal accommodation. Our results emphasize that beneficial microbes have the
ability to modify plant dPCD processes to enhance host colonization.

Results

The SMB-mediated clearance of dead cells protects the root meristem and regulates symbiosis

dPCD and corpse clearance are the final steps in LRC differentiation, maintaining root cap organ size in Arabidopsis root tips. This process is orchestrated by the LRC-specific transcription factor SMB and executed by its direct and indirect downstream targets (**Fig. 1A**). To characterize the role of disrupted dPCD in Arabidopsis LRCs, we analyzed the phenotypic implications of the SMB loss-of-function allele *smb-3* (Willemsen et al., 2008). We employed two staining methods to visualize the extent of cell death and protein aggregation in the *smb-3* T-DNA insertion line. We first used Evans blue staining, a viability dye that penetrates damaged/dying cells (Vijayaraghavareddy et al., 2017). In *smb-3* mutants, this staining revealed uncleared LRC cell corpses along the surface of primary roots, starting at the distal border of the meristematic zone (**Fig. 1B, C and S1A**). We further characterized *smb-3* mutants with Proteostat staining, a fluorescent dye that binds to quaternary protein structures typically found in misfolded and/or aggregated and condensed proteins (hereafter referred to as protein aggregates) (Llamas et al., 2021). In *smb-3* mutants, Proteostat staining showed an accumulation of protein aggregates in uncleared dead LRC cells attached to the roots (**Fig. 1D – F**).

The transcription factor SMB promotes the expression of various dPCD executor genes, including proteases that break down and clear cellular debris and protein aggregates following cell death induction. In the LRCs of *smb-3* mutants, the absence of induction of these proteases potentially explains the accumulation of protein aggregates in uncleared dead LRC cells.

Filter trap analysis further confirmed elevated levels of protein aggregates in *smb-3* mutants compared to WT roots (**Fig. 1G**). Under physiological conditions in WT roots, we previously observed protein aggregate accumulation in sloughed columella cell packages, but not during dPCD of distal LRC clearance (Llamas et al., 2021). This aligns with the findings that dPCD of the columella is affected by the loss of autophagy, while dPCD of the LRC is not (Feng et al., 2022).

In *smb-3* mutants, neither Proteostat nor Evans blue staining was detected in LRC cells
covering the meristem or in epidermal cells beneath the uncleared LRC cell corpses along
the elongation and differentiation zone (**Fig. 1B, D and F**). This observation highlights
that the loss of SMB activity specifically affects the induction of dPCD in LRC cells at the
transition between meristematic and elongation zone. These data show that the delayed
cell death of LRC cells in *smb-3* mutants in the elongation zone is accompanied by
impaired protein homeostasis (proteostasis), resulting in the accumulation of misfolded
and aggregated proteins in uncleared LRC cell corpses.

To assess the effects of impaired dPCD processes in the LRC of *smb-3* mutants on plant-
microbe interactions, we measured colonization rates of *S. indica*. We quantified
extraradical colonization using the chitin-binding fluorescent marker Alexa Fluor 488
conjugated to Wheat Germ Agglutinin (WGA-AF 488) as a proxy for fungal biomass. We
compared staining intensities of WGA-AF 488 between *S. indica*-colonized WT and *smb-*
3 roots. *S. indica* showed a clear hypercolonization phenotype along the main root axis
of *smb-3* mutants (**Fig. 2A, B**). On WT roots, *S. indica* preferentially colonized the
differentiation zone, leaving the meristematic and elongation zone largely uncolonized,
whereas mycelial growth was clearly detectable at the root tips of *smb-3* plants (**Fig. 2A**
and Fig. S1B) (Deshmukh et al., 2006, Jacobs et al., 2011). Intraradical colonization by
S. indica was quantified by comparing fungal and plant single-copy housekeeping marker
genes using quantitative PCR (qPCR), after washing of roots to remove outer fungal
mycelium. The results showed a significant increase in intraradical fungal accommodation
in *smb-3* mutants compared to WT roots (**Fig. 2C**). To assess the biological implications
of hypercolonization, we measured *S. indica*-induced root growth in WT and *smb-3*
mutants. While *S. indica* consistently and significantly increased root length in WT plants
at 8, 10 and 14 dpi, increased length of *smb-3* mutant roots was only observable at later
stages of colonization, indicating a delayed growth promotion phenotype (**Fig. 2D**).
Detailed cytological analysis confirmed that *S. indica* grew extensively around the
meristematic zone of *smb-3* mutants but not of WT roots (**Fig. 3A, B**) and showed
increased colonization of *smb-3* mutants in the differentiation zone (**Fig. 3C**). We further
observed that *S. indica* was accommodated in cells that were subject to cell death and
protein aggregation in the *smb-3* background (**Fig. 3B, D, E**). Together, these findings

indicate that loss of SMB activity in the root cap results in an accumulation of uncleared
LRC cell corpses that promotes fungal colonization from the meristematic to the
differentiation zone. Therefore, we postulate that the continuous clearance of root cap
cells in WT roots is important to limit microbial colonization along the entire root axis and
prevent microbial colonization in the meristematic zone.

Interestingly, Evans blue cell death staining of *S. indica*-inoculated *smb-3* mutants
displayed a clearing of LRC cell corpses from the surface of *smb-3* mutant roots over
time, while mock-treated *smb-3* mutant roots remained littered with LRC cell corpses (**Fig.**
3F and Fig. S1C). This observation indicates that *S. indica* is able to degrade uncleared
cell corpses, which likely provide additional nutrients that fuel fungal hypercolonization in
the *smb-3* mutant background.

The senescence associated plant nuclease BFN1 is exploited by beneficial
microbes to facilitate root accommodation

To further explore the role of root dPCD during *S. indica* accommodation in Arabidopsis,
we performed transcriptome analysis, tracking developmental cell death-marker gene
expression during different colonization stages (Olvera-Carrillo et al., 2015). The major
regulator in LRCs, *SMB*, showed no significant changes in expression during fungal
colonization (**Fig**. **4A, C**). However, in Arabidopsis colonized by *S. indica*, there was a
significant decrease in *BFN1* expression observed after 6 dpi (**Fig. 4B, C**). To validate
the RNA-Seq analysis, we performed whole-root qPCR of WT mock- and colonized-roots,
confirming *BFN1* downregulation at the onset of cell death in *S. indica*-colonized plants
(**Fig. 4D**).

While *SMB* expression is restricted to the LRC, *BFN1* exhibits a broader expression
pattern across various cell types and tissues, such as root cap cells, cells adjacent to
emerging lateral root primordia, differentiating xylem tracheary elements, as well as
senescent leaves, and abscission zones of flowers and seeds (Farage-Barhom et al.,
2008, Escamez et al., 2020). This widespread expression establishes BFN1 as a key
player in the general regulation of dPCD and senescence processes in various tissues in
Arabidopsis. To visualize the extent of *BFN1* downregulation upon *S. indica* colonization
in different zones of the root, we used a transgenic *BFN1* promoter-reporter line

(*pBFN1::NLS-tdTOMATO*) (Huysmans et al., 2018). In agreement with the previously

described GUS reporter lines (Farage-Barhom et al., 2008), we detected activation of the

BFN1 promoter via accumulation of the fluorescent tdTOMATO signal in the nuclei of root

cap and xylem cells in mock-treated roots (**Fig. S2A**). Additionally, we observed promoter

activation in epidermal root cells of the differentiation zone (**Fig. 4E**). In the distal region

of the differentiation zone in young epidermal cells, the tdTOMATO signal was observed

in nuclei, while in the basal region of the differentiation zone in older epidermal cells, the

tdTOMATO signal appeared to be dispersed (**Fig. 4E, Fig S2C**). These findings indicate

ongoing nuclear envelope breakdown as a hallmark of cell death (Wang et al., 2024) in

the older part of the root, independent of fungal colonization and suggest activation of

BFN1 during root epidermal cell aging/senescence. Next, we inoculated the *pBFN1*

reporter lines with *S. indica* and observed a reduction in promoter activity in epidermal

cells that were in contact with the fungus compared with mock-treated roots (**Fig. 4E, F**).

BFN1 expression and nuclear localization in the root cap or xylem was not affected by *S.*

indica colonization (**Fig. S2B**). This indicates that the downregulation of *BFN1* by *S. indica*

occurs mainly in epidermal cells of the differentiation zone and is regulated independently

of SMB and its activity in the root cap.

To assess the phenotypic effects of *BFN1* downregulation, we analyzed Arabidopsis *bfn1-*

1 null mutants (Fendrych et al., 2014) using the cell death and protein aggregates

markers, Evans blue and Proteostat. Staining of *bfn1-1* mutants with Evans blue showed

an increase of cell remnants in the epidermal cell layer of the differentiation zone (**Fig.**

5A, B), consistent with a proposed delay of dPCD and cell corpse clearance by BFN1

activity. Furthermore, while WT roots were devoid of protein aggregates, *bfn1-1* mutants

exhibited aggregates along the primary root axis, starting at the transition between

elongation and differentiation zone. The meristematic zone remained free of protein

aggregates (**Fig. 5C, D**). These data suggest that the lack of BFN1 activity in the root cap,

xylem, and senescent epidermis creates a general cellular stress in the roots that affects

proteostasis in the differentiation zone (**Fig. S3**). Similar to *bfn1-1*, WT roots colonized by

S. indica showed increased Evans blue and Proteostat signal in the differentiation zone.

Aggregated proteins were detected in colonized and adjacent non-colonized cells along

the differentiation zone, suggesting a non-cell autonomous host response to the fungus
(**Fig. S4**).

To investigate the biological relevance of *BFN1* downregulation during *S. indica* root
colonization, we quantified extraradical fungal growth using the WGA-AF 488 stain. When
comparing staining intensities of *S. indica*-inoculated *bfn1-1* mutants and WT seedlings,
we observed a significantly stronger fluorescence signal at the roots of *bfn1-1* mutants,
indicating a higher extraradical fungal colonization along the differentiation zone (**Fig. 5E,**
F and Fig. S3D). However, similar to WT roots, *bfn1-1* mutants did not exhibit fungal
colonization around the meristematic zone as observed in *smb-3*-colonized roots (**Fig.**
5E). Quantification of intraradical colonization by qPCR, after removal of outer fungal
mycelium, showed a significant increase of *S. indica* biomass in *bfn1-1* mutants at later
stages of interaction (**Fig. 5G**). Together, these results emphasize that downregulation of
BFN1 during colonization is beneficial for intra- and extraradical fungal accommodation
in the differentiation zone.

Next, we investigated the impact of other beneficial microbes on dPCD by examining
transcriptional responses in Arabidopsis roots colonized by different organisms. These
included *Serendipita vermifera*, an orchid mycorrhizal fungus closely related to *S. indica*,
and two bacterial synthetic communities (SynComs) derived from either Arabidopsis roots
or the rhizosphere of *Hordeum vulgare* (Mahdi et al., 2022). In all three interactions, *BFN1*
expression was consistently decreased in Arabidopsis roots (**Fig. S5A**). Additionally,
RNA-Seq analysis of Arabidopsis dPCD marker genes during *S. vermifera* colonization
confirmed the downregulation of *BFN1* (**Fig. S5 B, C**). Our findings indicate that microbes
may benefit from delayed post-mortem corpse clearance after dPCD in host plants and
suggest that beneficial microbes may have evolved mechanisms to manipulate dPCD
pathways to increase colonization (**Fig. 6**).

Discussion

In this study, we investigated the functional link between dPCD and microbial
accommodation in roots. Impaired dPCD in the Arabidopsis *smb-3* and *bfn1-1* mutants
increased colonization by the beneficial endophyte *S. indica*. The *smb-3* mutants
displayed hypercolonization along the entire primary root, suggesting that the extra sheet

of cell corpses surrounding *smb-3* roots provide additional and easily accessible nutrients that fuel fungal colonization. In fact, we observed a clearing effect with progressive colonization stages. Furthermore, hypercolonization of Arabidopsis, caused by the loss of dPCD in the root cap, mitigated the beneficial effects of *S. indica* by delaying the induction of growth promotion. Most notably, we observed hypercolonization of the meristematic zone of *smb-3* mutants (**Fig. 6**). The root apical meristem embedded in this zone of the root tip is essential for root growth, as all primary root tissue originates from these continuously dividing stem cells (Hamamoto et al., 2006). This sensitive tissue is surrounded by the root cap, which protects it from external stresses (Kumar and Iyer-Pascuzzi, 2020). The phenotype of *smb-3* mutants shows resemblance to the human skin disease hyperkeratosis. In healthy human skin, a pool of stem cells produces layers of cells that divide, differentiate, die, and are shed. Such developmental programs form a physical and dynamic barrier against environmental factors. Microbes attempting to establish themselves are consistently removed by skin exfoliation (Dettmer, 2021). Patients with hyperkeratosis show an accumulation of dead cells on the outer skin layer, making them more susceptible to microbial infection (Cheng et al., 1992). Likewise, malfunctions within regulated cell death in mammal gut epithelial cells produce death-induced nutrient release (DINNR) that can fuel bacterial growth and infection and could cause a variety of disorders such as inflammatory diseases (Anderson et al., 2021). The importance of an intact root cap in plant-microbe interactions is further highlighter by the fact that the physical removal of root caps in maize plants leads to increased colonization of the root tip by the plant growth promoting rhizobacterium (PGPR) *Pseudomonas fluorescens* (Humphris et al., 2005) and to changes in the rhizosphere microbiome composition along the root axis (Ruger et al., 2023). Moreover, it has been suggested that border cells released from the root cap may distract root-feeding nematodes from attacking plant roots (Rodger et al., 2003). Our results provide strong evidence that root cap size maintenance in the form of constant root cap cell turnover in Arabidopsis acts as a dynamic barrier, analogous to epidermal cell turnover in animals. It thus represents a sophisticated physical mechanism to prevent or reduce intracellular microbial colonization near the root meristematic tissue and contributes to the maintenance of a beneficial interaction with root endophytes.

Mutation of *BFN1* displayed significantly increased colonization in the differentiation zone but not in the meristematic zone (**Fig. 6**). The differences in colonization patterns to the *smb-3* mutants likely reflect spatial expression and localization patterns of SMB and BFN1 activity throughout Arabidopsis roots. While the expression of the transcription factor *SMB* is restricted to the LRC, the senescence-associated nuclease *BFN1* is expressed in different tissues undergoing dPCD and senescence below and above ground (Farage-Barhom et al., 2008, Escamez et al., 2020). Here, we show that *BFN1* is additionally expressed in differentiated root epidermal cells that undergo nuclear degradation during root maturation. This suggests an age-dependent dPCD in the outer epidermal layer of Arabidopsis roots where expression of *BFN1* possibly pre-dates cortical and epidermal abscission during secondary root growth and contributes to clearance of cell corpses during periderm emergence (Wunderling et al., 2018). This process resembles root cortical senescence and cell death in grass species such as wheat, barley, and corn, which typically start in the epidermis and spread toward the endodermis (Drew et al., 2000).

Additionally, we observed distinct differences in the presence and distribution of protein aggregates in *smb-3* and *bfn1-1* mutants. In *smb-3* mutants protein aggregates are restricted to LRC cells adhering to the primary root. This localized distribution may be attributed to the absence of SMB-induced proteases in the LRC cells of these mutants. The absence of these proteases likely impairs protein turnover and degradation, leading to accumulation and aggregation specifically in these cells. In contrast, *bfn1-1* mutation results in a more extensive and diffuse pattern of protein aggregation in the epidermal cell layer of the differentiation zone, regardless of the occurrence of cell death. This widespread distribution suggests that BFN1 plays a broader role in maintaining cellular homeostasis and protein quality control throughout the root epidermis and differentiation zone. These distinct patterns of protein aggregation and cell death in *smb-3* and *bfn1-1* mutants underscore the importance of these genes in maintaining cellular integrity and protein homeostasis. The contrasting phenotypes highlight the distinct roles of these genes in cellular processes.

While we show that dPCD protects the meristem from microbial colonization, we propose that some adapted microbes manipulate host dPCD processes by affecting the transcriptional expression of *BFN1* to facilitate accommodation in the root. Whether active interference by fungal effector proteins, fungal-derived signaling molecules or a systemic response of Arabidopsis roots underlies *BFN1* downregulation by *S. indica* remains to be investigated. It was recently demonstrated that small active metabolites produced either by Toll/interleukin-1 receptor (TIR)-containing leucine-rich repeat (NLR) receptors (Yu et al., 2022) or by fungal-derived enzymes (Dunken et al., 2022) through hydrolysis of RNA/DNA can mediate host cell death. This raises an exciting possibility that RNAse and DNAse activities of BFN1 may be involved in producing small, active nucleotide-derived metabolites that affect cell death, cell corpse clearance, proteostasis and fungal accommodation in the differentiation zone. To explore this hypothesis further, metabolomic and proteomic approaches should be employed in future studies. Notably, we have shown that colonization by other beneficial microbes such as the closely related fungus *S. vermifera* and bacterial members of the Arabidopsis and *H. vulgare* microbiota (Mahdi et al., 2022) also led to the downregulation of *BFN1* in Arabidopsis. These findings emphasize the presence of a conserved pathway influenced by diverse beneficial microbes to downregulate *BFN1* expression in epidermal tissue to facilitate symbioses.

Transcriptomic analysis of both established and predicted key dPCD marker genes revealed diverse patterns of upregulation and downregulation during *S. indica* colonization. These findings provide a valuable foundation for future studies investigating the dynamics of dPCD processes during beneficial symbiotic interactions and the potential manipulation of these processes by symbiotic partners.

In conclusion, our data show that tight regulation of host dPCD in epidermal- and root cap-tissue plays an important role in restricting fungal colonization. From a microbial perspective, dPCD pathways represent a potential nexus for enhancing symbiotic interactions and potentially improving nutrient availability (**Fig. 6**). These results shed light on the complex relationship between PCD and microbial accommodation in plant roots, offering valuable insights into the development of plants that establish more efficient partnerships with beneficial microbes.

Materials and methods

Fungal strains and growth conditions

Fungal experiments were performed with *Serendipita indica* strain DSM11827 (German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany). *S. indica* was grown on complete medium (CM) containing 2% (w/v) glucose and 1.5% (w/v) agar (Hilbert et al., 2012). Fungal material was grown at 28°C in the dark for 4 weeks before spore preparation. For additional experiments, *S. vermifera* (MAFF305830) was used and grown on MYP medium (7 g/l malt extract (Sigma-Aldrich), 1 g/l peptone (Sigma-Aldrich), 0.5 g/l yeast extract (Carl Roth) containing 1.5% agar at 28°C in darkness for 3 weeks before mycelium preparation for root inoculation.

Plant material and growth conditions

Seeds of *Arabidopsis thaliana* wild-type (WT) ecotype Columbia 0 (Col-0) and T-DNA insertion mutants (*bfn1-1* [GK-197G12] and *smb-3* [SALK_143526C]) in Col-0 background were used for experiments. Seeds were surface sterilized in 70% ethanol for 15 min and 100% ethanol for 12 min, stratified at 4°C in the dark for 3 days and germinated and grown on ½ MS medium (Murashige-Skoog Medium, with vitamins, pH 5.7, Duchefa Biochemie) containing 1% (w/v) sucrose and 0.4% (w/v) Gelrite (Duchefa Biochemie) under short-day conditions (8 h light, 16 h dark) with 130 μmol $m^{-2\,s^{-1}}$ light and 22°C/18 °C.

Fungal inoculation

One-week-old seedlings were transferred to 1/10 PNM (Plant minimal Nutrition Medium, 0.5 mM KNO_3, 0.367 mM KH_2PO_4, 0.144 mM K_2HPO_4, 2 mM $MgSO_4$ x H_2O, 0.2 mM $Ca(NO_3)_2$, 0.25% (v/v) Fe-EDTA (0.56% w/v $FeSO_4$ x $7H_2O$ and 0.8% w/v Na_2EDTA x $2H_2O$), 0.428 mM NaCl; pH-adjusted to 5.7 and buffered with 10 mM MES. For solid media, 0.4% (w/v) Gelrite (Duchefa Biochemie) was added) plates without sucrose, using 15 to 20 seedlings per plate. Under sterile conditions, spores of *S. indica* were scraped from agar CM plates using 0.002% Tween water (Roth), washed two times with ddH_2O and pipetted in a volume of 2 ml on plant roots and surrounding area in a concentration of $5x10^5$ spores per plate. ddH_2O was used for inoculation of mock plants. For *S.*

vermifera inoculation, mycelium was scrapped from plates in ddH$_2$O, washed and added to Arabidopsis roots in a volume of 1 ml of a stock solution of 1 g / 50 ml.

In case of experiments using seeds inoculation with *S. indica*, Arabidopsis seeds were surface sterilized, incubated with fungal spore solution at 5x10^5 concentration for 1 hour and plated on ½ MS plates (without sucrose).

Evans blue staining

For the visualization of cell death in Arabidopsis roots a modified protocol by (Vijayaraghavareddy et al., 2017) was used. Roots were washed three times in ddH$_2$O to remove loose external fungal mycelium and then stained for 15 min in 2 mM Evans blue (Sigma-Aldrich) dissolved in 0.1 M CaCl$_2$ pH 5.6. Subsequently, roots were washed extensively with ddH$_2$O for 1 hour and a Leica M165 FC microscope was used for imaging. 4 pictures were taken along the main root axis of each plant and averaged together, for an overview of cell death in the differentiation zone of one root. To quantify Evans blue staining intensity, ImageJ was used to invert the pictures, draw out individual roots and measure and compare mean grey values.

Extraradical colonization assays

To quantify extraradical colonization of *S. indica* on Arabidopsis, seed-inoculated plants were grown for 10 days. Inoculated and mock-treated seedlings were stained directly on agar plate by pipetting 2 ml of 1X Phosphate-buffered saline (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$) solution containing Alexa Fluor 488 conjugated with Wheat Germ Agglutinin (5 µl/mL from 1mg/ml stock) (WGA-AF 488, Invitrogen). After 2 min of incubation, the roots were washed twice on the plate with 1X PBS solution. The stained seedlings were transferred to a fresh ½ MS square plate (Greiner Bio-One). In order to perform a correct and focused scan of the agar plate with the roots, it was checked that the solid MS medium was flat and even and had no unevenness. To scan the agar plate, we used an Odyssey M Imaging System (LI-COR Biosciences) and the LI-COR Acquisition Software 1.1 (LI-COR Biosciences). In the software, we selected custom assay and then membrane. We selected the area in the agar plate to be scanned and selected the channels 488 (for WGA-Alexa Flour 488) and

RGB trans (for bright field). We defined the focus offset between 3.5 mm and 4.0 mm (depending on the thickness of the MS medium). For resolution we selected 10 μm when scanning two plants or 100 μm when scanning several plants. Quantification of WGA-AF 488 fluorescence was performed using EmpiriaStudio Software (LI-COR Biosciences).

RNA extraction (intraradical colonization assay)

To measure intraradical colonization via RNA extraction and PCR, plants were harvested at three time points around 7, 10 and 14 dpi. The roots were extensively washed with ddH$_2$O and tissue paper was used to carefully wipe off mycelium on the surface of the roots. After cleaning, the roots were shock frozen in liquid nitrogen and RNA was extracted with TRIzol (Invitrogen, Thermo Fisher Scientific, Schwerte, Germany). After a DNase I (Thermo Fisher Scientific) treatment according to the manufacturer's instructions to remove DNA, RNA was used to generate cDNA through the utilization of the Fermentas First Strand cDNA Synthesis Kit (Thermo Fisher Scientific).

Quantitative RT-PCR analysis

The quantitative real time-PCR (qRT-PCR) was performed using a CFX connect real time system (BioRad) with the following program: 95 °C 3min, 95 °C 15s, 59 °C 20s, 72 °C 30 s, 40 cycles and melting curve analysis. Relative expression was calculated using the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). qRT-PCR primers can be found in Table S1.

Filter trap analysis

Filter trap assays were performed as previously described (Llamas et al., 2023, Llamas et al., 2021). Protein extracts were obtained using native lysis buffer (300 mM NaCl, 100 mM HEPES pH 7.4, 2 mM EDTA, 2% Triton X-100) supplemented with 1x plant protease inhibitor (Merck). Cell debris was removed by several centrifugation steps at 8,000 x g for 10 min at 4 °C. The supernatant was separated, and protein concentration determined using the Pierce BCA Protein Assay Kit (Thermo Fisher). A cellulose acetate membrane filter (GE Healthcare Life Sciences) and 3 filter papers (BioRad, 1620161) were immersed in 1x PBS and placed in a slot blot apparatus (Bio-Rad) connected to a vacuum system. The membrane was equilibrated by 3 washes with equilibration buffer (native buffer containing 0.5% SDS). 300, 200 and 100 μg of the protein extract were

mixed with SDS at a final concentration of 0.5% and filtered through the membrane. The membrane was then washed with 0.2% SDS and blocked with 3% BSA in Tris-Buffered Saline 0.1% Tween (TBST, 50mM Tris-CL, 150mM NaCl, pH 7.5) for 30 minutes, followed by 3 washes with TBST. Incubation was performed with anti-polyQ [1:1000] (Merck, MAB1574). The membrane was washed 3 times for 5 min and incubated with secondary antibodies in TBST 3% BSA for 30 min. The membrane was developed using the Odyssey M Imaging System (LI-COR Biosciences). Extracts were also analyzed by SDS-PAGE and western blotting against anti-Actin [1:5000] (Agrisera, AS132640) to determine loading controls.

Confocal laser scanning microscopy (CLSM) and Proteostat staining quantification

CLSM images were acquired using either the FV1000 confocal laser scanning microscope (Olympus) or a Meta 710 confocal microscope with laser ablation 266 nm (Zeiss). All images were acquired using the same parameters between experiments. Excitation of WGA-AF 488 was done with an argon laser at 488 nm and the emitted light was detected with a hybrid detector at 500-550 nm. Proteostat was excited at 561 nm and the signal was detected between 590-700 nm. Hoechst was excited with a diode laser at 405 nm and the emitted light was detected with a hybrid detector at 415-460 nm.

Proteostat staining

For the detection of aggregated proteins, we used the Proteostat Aggresome detection kit (Enzo Life Sciences). Seedlings were stained according to the manufacturer's instructions. Seedlings were incubated with permeabilizing solution (0.5% Triton X-100, 3 mM EDTA, pH 8.0) for 30 minutes at 4°C with gentle shaking. The seedlings were washed twice with 1X PBS. Then the plants were incubated in the dark with 1x PBS supplemented with 0.5 µl/ml Proteostat and 0.5 µl/ml Hoechst 33342 (nuclear stain) for 30 min at room temperature. Finally, the seedlings were washed twice with 1x PBS and mounted on a slide for CLSM analysis or in mounted in fresh MS phytoagar plates for LI-COR analysis. Quantification of Proteostat fluorescence was performed using Fiji software or EmpiriaStudio Software (LI-COR Biosciences).

Transcriptomic analysis

Arabidopsis Col-0 WT roots were inoculated with *S. indica*. Arabidopsis roots were harvested from mock-treated plants and inoculated plants at four different time points post inoculation: 1, 3, 6 and 10 dpi. Three biological replicates were used for each condition. The RNA-seq libraries were generated and sequenced at US Department of Energy Joint Genome Institute (JGI) under a project proposal (Proposal ID: 505829) (Zuccaro and Langen, 2020, Eichfeld et al., 2023). For each sample, stranded RNA-Seq libraries were generated and quantified by qRT-PCR. RNA-Seq libraries were sequenced with Illumina sequencer. Raw reads were filtered and trimmed using the JGI QC pipeline. Filtered reads from each library were aligned to the Arabidopsis genome (TAIR10) using HISAT2 (Kim et al., 2015) and the reads mapped to each gene were counted using featureCounts (Liao et al., 2014). Samples harvested at 1 dpi were omitted from the analysis because we decided to focus on the time points at which the interaction between Arabidopsis and *S. indica* is well established. Differential gene expression analysis was performed using the R package DESeq2 (Love et al., 2014). Genes with an FDR adjusted p-value < 0.05 were considered as differentially expressed genes (DEGs). The adjusted p-value refers to the transformation of the p-value obtained with the Wald test after considering multiple testing. To visualize gene expression, genes expression levels were normalized as Transcript Per kilobase million (TPM).

Statistical analyses

All statistical analyses, except for the transcriptomic analysis, were performed using Prism8. Individual figures state the applied statistical methods, as well as p and F values. p-*values* and corresponding asterisks are defined as following, $p < 0.05$ *, $p < 0.01$**, $p < 0.001$***.

Acknowledgements

We thank the imaging facilities of CECAD (A. Schauss and C. Jüngst) and CEPLAS (P.S. Tan) for their assistance with CLSM. We thank Lisa Mahdi for conducting the experiments and providing the samples used for the RNA-seq analysis. We further would like to thank Yu Zhang, Sravanthi Tejomurthula, Daniel Peterson, Vivian Ng & Igor Grigoriev and their

work performed in the work proposal (10.46936/10.25585/60001292) conducted by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231. AZ, NC and EL acknowledge support by the German Research Foundation (DFG) - Excellence Strategy of the Federal Republic of Germany - EXC-2048/1 - project ID 390686111. AZ and NC acknowledge support by the SFB-1403-414786233 and DV by the German Excellence Strategy-CECAD, EXC 2030-390661388. MKN gratefully acknowledges funding from the European Research Council (ERC) (StG PROCELLDEATH 639234 and CoG EXECUT.ER 864952) as well as the Research Foundation – Flanders (FWO) projects no. G002120N and G002620N.

Contributions

References

AKUM, F. N., STEINBRENNER, J., BIEDENKOPF, D., IMANI, J. & KOGEL, K.-H. 2015. The Piriformospora indica effector PIIN_08944 promotes the mutualistic Sebacinalean symbiosis. *Frontiers in Plant Science,* 6.

ANDERSON, C. J., MEDINA, C. B., BARRON, B. J., KARVELYTE, L., AAES, T. L., LAMBERTZ, I., PERRY, J. S. A., MEHROTRA, P., GONCALVES, A., LEMEIRE, K., BLANCKE, G., ANDRIES, V., GHAZAVI, F., MARTENS, A., VAN LOO, G., VEREECKE, L., VANDENABEELE, P. & RAVICHANDRAN, K. S. 2021. Microbes exploit death-induced nutrient release by gut epithelial cells. *Nature,* 596**,** 262-267.

BARLOW, P. W. 2002. The Root Cap: Cell Dynamics, Cell Differentiation and Cap Function. *Journal of Plant Growth Regulation,* 21**,** 261-286.

BENNETT, T., VAN DEN TOORN, A., SANCHEZ-PEREZ, G. F., CAMPILHO, A., WILLEMSEN, V., SNEL, B. & SCHERES, B. 2010. SOMBRERO, BEARSKIN1, and BEARSKIN2 regulate root cap maturation in Arabidopsis. *Plant Cell,* 22**,** 640-54.

BOORBOORI, M. R. & ZHANG, H. Y. 2022. The Role of Serendipita indica (Piriformospora indica) in Improving Plant Resistance to Drought and Salinity Stresses. *Biology (Basel),* 11.

CHENG, J., SYDER, A. J., YU, Q. C., LETAI, A., PALLER, A. S. & FUCHS, E. 1992. The genetic basis of epidermolytic hyperkeratosis: a disorder of differentiation-specific epidermal keratin genes. *Cell,* 70**,** 811-9.

DESHMUKH, S., HUCKELHOVEN, R., SCHAFER, P., IMANI, J., SHARMA, M., WEISS, M., WALLER, F. & KOGEL, K. H. 2006. The root endophytic fungus Piriformospora indica requires host cell death for proliferation during mutualistic symbiosis with barley. *Proc Natl Acad Sci U S A,* 103**,** 18450-7.

DETTMER, P. 2021. *Immune: The new book from Kurzgesagt - a gorgeously illustrated deep dive into the*
immune system, Hodder & Stoughton.
DOLAN, L., JANMAAT, K., WILLEMSEN, V., LINSTEAD, P., POETHIG, S., ROBERTS, K. & SCHERES, B. 1993.
Cellular organisation of the Arabidopsis thaliana root. *Development,* 119**,** 71-84.
DREW, M. C., HE, C. J. & MORGAN, P. W. 2000. Programmed cell death and aerenchyma formation in
roots. *Trends Plant Sci,* 5**,** 123-7.
DUNKEN, N., WIDMER, H., BALCKE, G. U., STRAUBE, H., LANGEN, G., CHARURA, N. M., SAAKE, P., LESON,
L., RÖVENICH, H., WAWRA, S., DJAMEI, A., TISSIER, A., WITTE, C. P. & ZUCCARO, A. 2022. A
fungal endophyte-generated nucleoside signal regulates host cell death and promotes root
colonization. *bioRxiv,* 2022.03.11.483938.
EICHFELD, R., MAHDI, L. K., QUATTRO, C. D., ARMBRUSTER, L., ENDESHAW, A. B., MIYAUCHI, S.,
HELLMANN, M. J., CORD-LANDWEHR, S., GRIGORIEV, I., PETERSON, D., SINGAN, V., LAIL, K.,
SAVAGE, E., NG, V., LANGEN, G., MOERSCHBACHER, B. M. & ZUCCARO, A. 2023. Time-resolved
transcriptomics reveal a mechanism of host niche defense: beneficial root endophytes deploy a
host-protective antimicrobial GH18-CBM5 chitinase. *bioRxiv,* 2023.12.29.572992.
ESCAMEZ, S., ANDRE, D., SZTOJKA, B., BOLLHONER, B., HALL, H., BERTHET, B., VOSS, U., LERS, A., MAIZEL,
A., ANDERSSON, M., BENNETT, M. & TUOMINEN, H. 2020. Cell Death in Cells Overlying Lateral
Root Primordia Facilitates Organ Growth in Arabidopsis. *Curr Biol,* 30**,** 455-464 e7.
FARAGE-BARHOM, S., BURD, S., SONEGO, L., METT, A., BELAUSOV, E., GIDONI, D. & LERS, A. 2011.
Localization of the Arabidopsis Senescence- and Cell Death-Associated BFN1 Nuclease: From the
ER to Fragmented Nuclei. *Molecular Plant,* 4**,** 1062-1073.
FARAGE-BARHOM, S., BURD, S., SONEGO, L., PERL-TREVES, R. & LERS, A. 2008. Expression analysis of the
BFN1 nuclease gene promoter during senescence, abscission, and programmed cell death-
related processes. *J Exp Bot,* 59**,** 3247-58.
FENDRYCH, M., VAN HAUTEGEM, T., VAN DURME, M., OLVERA-CARRILLO, Y., HUYSMANS, M., KARIMI,
M., LIPPENS, S., GUERIN, C. J., KREBS, M., SCHUMACHER, K. & NOWACK, M. K. 2014.
Programmed cell death controlled by ANAC033/SOMBRERO determines root cap organ size in
Arabidopsis. *Curr Biol,* 24**,** 931-40.
FENG, Q., DE RYCKE, R., DAGDAS, Y. & NOWACK, M. K. 2022. Autophagy promotes programmed cell
death and corpse clearance in specific cell types of the Arabidopsis root cap. *Curr Biol,* 32**,** 2110-
2119 e3.
FESEL, P. H. & ZUCCARO, A. 2016. Dissecting endophytic lifestyle along the parasitism/mutualism
continuum in Arabidopsis. *Curr Opin Microbiol,* 32**,** 103-112.
HAMAMOTO, L., HAVES, M. C. & ROST, T. L. 2006. The production and release of living root cap border
cells is a function of root apical meristem type in dicotyledonous angiosperm plants. *Ann Bot,*
97**,** 917-23.
HILBERT, M., VOLL, L. M., DING, Y., HOFMANN, J., SHARMA, M. & ZUCCARO, A. 2012. Indole derivative
production by the root endophyte Piriformospora indica is not required for growth promotion
but for biotrophic colonization of barley roots. *New Phytol,* 196**,** 520-534.
HUMPHRIS, S. N., BENGOUGH, A. G., GRIFFITHS, B. S., KILHAM, K., RODGER, S., STUBBS, V., VALENTINE,
T. A. & YOUNG, I. M. 2005. Root cap influences root colonisation by Pseudomonas fluorescens
SBW25 on maize. *FEMS Microbiol Ecol,* 54**,** 123-30.
HUYSMANS, M., BUONO, R. A., SKORZINSKI, N., RADIO, M. C., DE WINTER, F., PARIZOT, B., MERTENS, J.,
KARIMI, M., FENDRYCH, M. & NOWACK, M. K. 2018. NAC Transcription Factors ANAC087 and
ANAC046 Control Distinct Aspects of Programmed Cell Death in the Arabidopsis Columella and
Lateral Root Cap. *Plant Cell,* 30**,** 2197-2213.

JACOBS, S., ZECHMANN, B., MOLITOR, A., TRUJILLO, M., PETUTSCHNIG, E., LIPKA, V., KOGEL, K. H. &
SCHAFER, P. 2011. Broad-spectrum suppression of innate immunity is required for colonization
of Arabidopsis roots by the fungus Piriformospora indica. *Plant Physiol,* 156**,** 726-40.
KIM, D., LANGMEAD, B. & SALZBERG, S. L. 2015. HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods,* 12**,** 357-60.
KUMAR, N. & IYER-PASCUZZI, A. S. 2020. Shedding the Last Layer: Mechanisms of Root Cap Cell Release.
Plants (Basel), 9.
KUMPF, R. P. & NOWACK, M. K. 2015. The root cap: a short story of life and death. *J Exp Bot,* 66**,** 5651-
62.
LIAO, Y., SMYTH, G. K. & SHI, W. 2014. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics,* 30**,** 923-30.
LIVAK, K. J. & SCHMITTGEN, T. D. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods,* 25**,** 402-8.
LLAMAS, E., KOYUNCU, S., LEE, H. J., WEHRMANN, M., GUTIERREZ-GARCIA, R., DUNKEN, N., CHARURA,
N., TORRES-MONTILLA, S., SCHLIMGEN, E., MANDEL, A. M., THEILE, E. B., GROSSBACH, J.,
WAGLE, P., LACKMANN, J.-W., SCHERMER, B., BENZING, T., BEYER, A., PULIDO, P., RODRIGUEZ-
CONCEPCION, M., ZUCCARO, A. & VILCHEZ, D. 2023. In planta expression of human polyQ-
expanded huntingtin fragment reveals mechanisms to prevent disease-related protein
aggregation. *Nature Aging*.
LLAMAS, E., TORRES-MONTILLA, S., LEE, H. J., BARJA, M. V., SCHLIMGEN, E., DUNKEN, N., WAGLE, P.,
WERR, W., ZUCCARO, A., RODRIGUEZ-CONCEPCION, M. & VILCHEZ, D. 2021. The intrinsic
chaperone network of Arabidopsis stem cells confers protection against proteotoxic stress.
Aging Cell, 20**,** e13446.
LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
MAHDI, L. K., MIYAUCHI, S., UHLMANN, C., GARRIDO-OTER, R., LANGEN, G., WAWRA, S., NIU, Y., GUAN,
R., ROBERTSON-ALBERTYN, S., BULGARELLI, D., PARKER, J. E. & ZUCCARO, A. 2022. The fungal
root endophyte Serendipita vermifera displays inter-kingdom synergistic beneficial effects with
the microbiota in Arabidopsis thaliana and barley. *ISME J,* 16**,** 876-889.
MORIWAKI, T., MIYAZAWA, Y., KOBAYASHI, A. & TAKAHASHI, H. 2013. Molecular mechanisms of
hydrotropism in seedling roots of Arabidopsis thaliana (Brassicaceae). *Am J Bot,* 100**,** 25-34.
NIZAM, S., QIANG, X., WAWRA, S., NOSTADT, R., GETZKE, F., SCHWANKE, F., DREYER, I., LANGEN, G. &
ZUCCARO, A. 2019. Serendipita indica E5'NT modulates extracellular nucleotide levels in the
plant apoplast and affects fungal colonization. *EMBO reports,* 20**,** e47430.
NOSTADT, R., HILBERT, M., NIZAM, S., ROVENICH, H., WAWRA, S., MARTIN, J., KÜPPER, H.,
MIJOVILOVICH, A., URSINUS, A., LANGEN, G., HARTMANN, M. D., LUPAS, A. N. & ZUCCARO, A.
2020. A secreted fungal histidine- and alanine-rich protein regulates metal ion homeostasis and
oxidative stress. *New Phytologist,* 227**,** 1174-1188.
OLVERA-CARRILLO, Y., VAN BEL, M., VAN HAUTEGEM, T., FENDRYCH, M., HUYSMANS, M., SIMASKOVA,
M., VAN DURME, M., BUSCAILL, P., RIVAS, S., COLL, N. S., COPPENS, F., MAERE, S. & NOWACK,
M. K. 2015. A Conserved Core of Programmed Cell Death Indicator Genes Discriminates
Developmentally and Environmentally Induced Programmed Cell Death in Plants. *Plant Physiol,*
169**,** 2684-99.
REZA, S. H., DELHOMME, N., STREET, N. R., RAMACHANDRAN, P., DALMAN, K., NILSSON, O., MININA, E.
A. & BOZHKOV, P. V. 2018. Transcriptome analysis of embryonic domains in Norway spruce
reveals potential regulators of suspensor cell death. *PLoS One,* 13**,** e0192945.

RODGER, S., BENGOUGH, A. G., GRIFFITHS, B. S., STUBBS, V. & YOUNG, I. M. 2003. Does the Presence of
Detached Root Border Cells of Zea mays Alter the Activity of the Pathogenic Nematode
Meloidogyne incognita? *Phytopathology,* 93**,** 1111-4.
RUGER, L., GANTHER, M., FREUDENTHAL, J., JANSA, J., HEINTZ-BUSCHART, A., TARKKA, M. T. &
BONKOWSKI, M. 2023. Root cap is an important determinant of rhizosphere microbiome
assembly. *New Phytol,* 239**,** 1434-1448.
SHI, C. L., VON WANGENHEIM, D., HERRMANN, U., WILDHAGEN, M., KULIK, I., KOPF, A., ISHIDA, T.,
OLSSON, V., ANKER, M. K., ALBERT, M., BUTENKO, M. A., FELIX, G., SAWA, S., CLAASSEN, M.,
FRIML, J. & AALEN, R. B. 2018. The dynamics of root cap sloughing in Arabidopsis is regulated by
peptide signalling. *Nat Plants,* 4**,** 596-604.
VIJAYARAGHAVAREDDY, P., ADHINARAYANREDDY, V., VEMANNA, R. S., SREEMAN, S. & MAKARLA, U.
2017. Quantification of Membrane Damage/Cell Death Using Evan's Blue Staining Technique. *Bio
Protoc,* 7**,** e2519.
WANG, J., BOLLIER, N., BUONO, R. A., VAHLDICK, H., LIN, Z., FENG, Q., HUDECEK, R., JIANG, Q., MYLLE, E.,
VAN DAMME, D. & NOWACK, M. K. 2024. A developmentally controlled cellular
decompartmentalization process executes programmed cell death in the Arabidopsis root cap.
Plant Cell, 36**,** 941-962.
WAWRA, S., FESEL, P., WIDMER, H., TIMM, M., SEIBEL, J., LESON, L., KESSELER, L., NOSTADT, R., HILBERT,
M., LANGEN, G. & ZUCCARO, A. 2016. The fungal-specific β-glucan-binding lectin FGB1 alters
cell-wall composition and suppresses glucan-triggered immunity in plants. *Nature
Communications,* 7**,** 13188.
WEISS, M., WALLER, F., ZUCCARO, A. & SELOSSE, M. A. 2016. Sebacinales - one thousand and one
interactions with land plants. *New Phytol,* 211**,** 20-40.
WILLEMSEN, V., BAUCH, M., BENNETT, T., CAMPILHO, A., WOLKENFELT, H., XU, J., HASELOFF, J. &
SCHERES, B. 2008. The NAC domain transcription factors FEZ and SOMBRERO control the
orientation of cell division plane in Arabidopsis root stem cells. *Dev Cell,* 15**,** 913-22.
WUNDERLING, A., RIPPER, D., BARRA-JIMENEZ, A., MAHN, S., SAJAK, K., TARGEM, M. B. & RAGNI, L.
2018. A molecular framework to study periderm formation in Arabidopsis. *New Phytol,* 219**,**
216-229.
YU, D., SONG, W., TAN, E. Y. J., LIU, L., CAO, Y., JIRSCHITZKA, J., LI, E., LOGEMANN, E., XU, C., HUANG, S.,
JIA, A., CHANG, X., HAN, Z., WU, B., SCHULZE-LEFERT, P. & CHAI, J. 2022. TIR domains of plant
immune receptors are 2',3'-cAMP/cGMP synthetases mediating cell death. *Cell,* 185**,** 2370-2386
e18.
ZUCCARO, A., LAHRMANN, U., GULDENER, U., LANGEN, G., PFIFFI, S., BIEDENKOPF, D., WONG, P.,
SAMANS, B., GRIMM, C., BASIEWICZ, M., MURAT, C., MARTIN, F. & KOGEL, K. H. 2011.
Endophytic life strategies decoded by genome and transcriptome analyses of the mutualistic
root symbiont Piriformospora indica. *PLoS Pathog,* 7**,** e1002290.
ZUCCARO, A. & LANGEN, G. 2020. Host-specific regulation of effector gene expression in mutualistic root
endophytic fungi (Proposal ID: 505829). JGI Award
671

Figure 1. _smb-3_ mutant roots exhibit uncleared cell corpses loaded with misfolded / aggregated proteins. (A) Schematic representation of lateral root cap (LRC) development in WT and _smb-3_ mutant plants, impaired in dPCD. (B) Evans blue staining of 10-day-old WT and _smb-3_ mutant roots, showing an overview of the meristematic, elongation and differentiation zone. Evans blue highlights the accumulation of LRC cell corpses on _smb-3_ mutant roots starting at the transition into the elongation zone. Scale indicates 150 µm. (C) Relative quantification of Evans blue staining of the differentiation zone of 14-day-old WT and _smb-3_ mutant roots (refers to image shown in B). 5 plants per genotype were used. All data points were normalized to the mean of the WT control. Statistical relevance was determined by unpaired, two-tailed Student's t test (F [4, 4] = 3.596; p < 0.001). (D) Confocal Scanning Laser Microscopy (CLSM) images of 10-day-old WT and _smb-3_ mutant roots stained with Proteostat (red) showing the meristematic- and the beginnings of the elongation-zone. Scale indicates

100 µm. (E) Magnification of the differentiation zone of WT and *smb-3* mutant roots, both

stained with Proteostat (red) and Hoechst (blue). WT roots do not show any Proteostat signal,

while *smb-3* mutants display extensive protein aggregate accumulation in uncleared LRC cell

corpses. Scale indicates 50 µm. (F) Quantification of relative Proteostat fluorescence levels,

comparing the differentiation and meristematic zones of WT and *smb-3* mutants. Five 10-day-

old plants were used for each genotype. All data points were normalized to the mean of the

WT control, analyzing differentiation and meristematic zone separately. Statistical

significance was determined by one-way ANOVA and Tukey's post hoc test ($F_{[3, 16]} = 8,314$; p

< 0.01). (G) Filter trap and SDS-PAGE analysis with anti-poly-glutamine (polyQ) antibodies of

15-day-old WT and *smb-3* mutant roots. The images are representative of two independent

experiments.

Figure 2. *smb-3* mutants display extraradical hypercolonization and increased intraradical colonization by *S. indica*. (A) Representative images show extraradical colonization of 10-day-old WT and *smb-3* mutant seedlings (seed inoculated). *S. indica* was stained with WGA-AF 488. Roots were scanned and captured with a LI-COR Odyssey M imager using the bright field (BF) and Alexa Fluor 488 channel (green). White arrowheads indicate colonization of the root tip in the *smb-3* mutant background. Scale indicates 5 mm. (B) Relative quantification of WGA-AF 488 signal as proxy for extraradical colonization on *smb-3* mutants and WT roots (refers to image shown in A). The statistical comparison was made by two-tailed Student's t test for unpaired samples (F [9, 9] = 11.85; p < 0.001) using 10 plants. (C) Measurement of intraradical colonization in WT and *smb-3* mutant roots at 10dpi were performed by qPCR. Roots from 7

biological replicates were collected and washed to remove extra-radical hyphae, pooling
approximately 30 seedlings for each genotype per replicate. The graph is normalized to WT.
Statistical analysis was done via two-tailed Student's t test for unpaired samples (F [6, 6] =
8.905; p < 0.05). (D) Root length measurements of WT plants and *smb-3* mutant plants, during
S. indica colonization (seed inoculated) or mock treatment. 50 plants for each genotype and
treatment combination were observed and individually measured over a time period of two
weeks. WT roots show *S. indica*-induced growth promotion, while growth promotion of *smb-*
3 mutants was delayed and only observed at later stages of colonization. This experiment was
performed three times, with fresh fungal material, showing similar results. Statistical analysis
was performed via one-way ANOVA and Tukey's post hoc test (F [11, 1785] = 1149; p < 0.001).
For visual representation of statistical relevance each time point was additionally evaluated
via one-way ANOVA and Tukey's post hoc test at 8 dpi (F [3, 593] = 69.24; p < 0.001), 10dpi (F
[3, 596] = 47.59; p < 0.001) and 14dpi (F [3, 596] = 154.3; p < 0.001).

Figure 3. Cytological analyses of *S. indica*-colonized *smb-3* mutants and WT roots. For CLSM analyses, 7-day-old seedlings were inoculated with *S. indica* spores and roots were analyzed at 10 dpi. (A) Representative images of the meristematic zone of Arabidopsis WT and *smb-3* mutants during *S. indica* colonization. WGA-AF 488 stain (green) was used to visualize fungal structures. Transmitted light (TL) images are also shown. Scale indicates 100 μm (B) Magnification of a *smb-3* mutant root tip colonized with *S. indica*. Asterisks indicate penetration of hyphae into dead cells stained with propidium iodide (PI, Sigma-Aldrich) shown

in magenta). Scale indicates 100 µm. (C) Representative images of the differentiation zone of
WT and *smb-3* mutants colonized with *S. indica* and stained with WGA-AF 488 and PI. Scale
indicates 100 µm. (D) Representative images of the meristematic zone of WT and *smb-3*
mutant root tips inoculated with *S. indica*, stained with WGA-AF 488 and Proteostat (red).
Scale indicates 100 µm (E) Magnification of the root differentiation zone of *smb-3* mutants
showing *S. indica* colonization, stained with WGA-AF 488, Hoechst and Proteostat.
Penetration of fungal hyphae into uncleared cell corpses is marked with asterisks. Dotted
yellow line indicates lateral root cap (LRC) cell corpse. Scale indicates 50 µm. (F)
Representative images of the differentiation zone of *S. indica*-colonized WT and *smb-3* roots
at 10 dpi, stained with Evans blue. Scale indicates 100 µm.

Figure 4. *BFN1* is downregulated during interaction with *S. indica*. RNA-Seq expression profiles of (A) *SMB* and (B) *BFN1* in Arabidopsis roots mock-treated or inoculated with *S. indica* at 3, 6 and 10 dpi. The log2-transformed Transcript per Kilobase million (TPM) values are shown and the lines indicate average expression values among the 3 biological replicates. Asterisk indicates significantly different expression (adjusted p-value < 0.05) (C) The heat map shows the expression values (TPM) of Arabidopsis dPCD marker genes with at least an average of 1 TPM across Arabidopsis roots mock-treated or inoculated with *S. indica* at 3, 6 and 10 dpi. The TPM expression values are log2 transformed and row-scaled. Genes are clustered using spearman correlation as distance measure. Each treatment displays the average of three biological replicates. The dPCD gene markers were previously defined (Olvera-Carrillo et al., 2015). (D) *BFN1* expression in WT Arabidopsis roots during *S. indica* colonization at 8 and 11 dpi. RNA was isolated from 3 biological replicates, pooling 30 plants per conditions for qPCR analysis, comparing *BFN1* expression with an Arabidopsis ubiquitin marker gene. Statistical significance was determined by one-way ANOVA and Tukey's post hoc test (F [3, 8]

= 7,263; p < 0.05). (E) Representative CLSM images of the differentiation zone of mock- and *S.*
indica- colonized *pBFN1::NLS-tdTOMATO* reporter roots at 7 dpi. The tdTOMATO signal
(magenta) represents BFN1 expression and *S. indica* was stained with WGA-AF 488 (green).
Scale indicates 100µm. (F) Whole seedling scans of mock- and *S. indica*-treated *pBFN1::NLS-*
tdTOMATO plants taken with a LI-COR Odyssey M imager at 7 dpi. Images show *BFN1*
expression via tdTOMATO signal in mock conditions and BFN1 expression in presence of *S.*
indica (stained with WGA-AF 488). Scale indicates 5mm.

Figure 5. *BFN1* downregulation promotes fungal accommodation. (A) Microscopy images of the differentiation zone of 14-day-old WT and *bfn1-1* mutant roots, stained with Evans blue. Scale indicates 100 µm. (B) Relative quantification of Evans blue staining (refers to image shown in A), comparing 14-day-old WT and *bfn1-1* mutants. 10 plants were used for each genotype. Data were normalized to the WT control. Statistical significance was determined using an unpaired, two-tailed Student's t test (F [9, 9] = 2.033; p < 0.001). (C) Proteostat staining of 10-day-old WT and *bfn1-1* mutant root tips. Scale indicates 100 µm. (D) Quantification of Proteostat staining (refers to image shown in C) using 4 to 5 10-day-old WT and *bfn1-1* mutants. Statistical analysis was performed via one-way ANOVA and Tukey's post hoc test (F [3, 14] = 33,55; p < 0.001). (E) Extraradical colonization of 10-day-old WT and *bfn1-1* mutant plants, seed-inoculated with *S. indica* and stained with WGA-AF 488 (green). Roots were scanned with a LI-COR Odyssey M imager. Arrowheads indicate the position of the uncolonized root tips. Scale indicates 5 mm. (F) Relative quantification of extraradical colonization of *bfn1-1* mutant and WT roots, using WGA-AF 488 signal as a proxy for fungal biomass (refers to image shown in E). Data were normalized to the WT control. Statistical comparisons were made by unpaired, two-tailed Student's t test for unpaired samples (F [5, 5] = 3.597; p < 0.01). (G) Intraradical colonization of WT and *bfn1-1* mutants was measured via qPCR. Roots from 7 biological replicates were collected and washed to remove outer extraradical mycelium, pooling approximately 30 plants per time point and replicate for each genotype. Each time point was normalized to WT for relative quantification of colonization. Statistical analysis was performed via one-way ANOVA and Tukey's post hoc test (F [5, 33] = 5.358; p < 0.01).

Figure 6. dPCD and its proposed effects on plant-microbe interactions. The root cap protects
and covers the stem cells of the root apical meristem. Its size in Arabidopsis is maintained by
a high cellular turnover of root cap cells. While the columella root cap is shed from the root
tip, a dPCD machinery marks the final step of LRC differentiation and prevents LRC cells from
entering into the elongation zone. Induction of cell death by the transcription factor SMB is
followed by irreversible DNA fragmentation and cell corpse clearance, mediated by the
nuclease BFN1, a downstream executor of dPCD (Fendrych et al., 2014). The absence of dPCD
induction in the *smb-3* knockout mutant leads to a delay in LRC differentiation and allows LRC
cells to enter the elongation zone, where they die uncontrolled, resulting in an accumulation
of LRC cell corpses along the differentiation zone. In a WT background, the fungal endophyte
S. indica colonizes the differentiation zone of Arabidopsis roots and can also be found in shed
columella cell packages. The impaired dPCD of the s*mb-3* mutant phenotype results in a
hypercolonization of Arabidopsis roots, along the differentiation zone as well as the
meristematic zone, highlighting that the continuous clearance of root cap cells is necessary
for restricting microbial accommodation at the meristematic zone. Loss of the downstream
dPCD executor *BFN1* does not affect fungal colonization in the meristematic zone but
increases accommodation by *S. indica* in the differentiation zone, where BFN1 appears to be
involved in dPCD of senescent epidermal cells and undergoes downregulation during *S. indica*
colonization.