Uncertainty in crop phenology simulations is driven primarily by parameter variability

Wallach[1], Daniel; Palosuo[2,*], Taru; Mielenz[3], Henrike; Buis[4], Samuel; Thorburn[5], Peter; Asseng[6], Senthold; Dumont[7], Benjamin; Ferrise[8], Roberto; Gayler[9], Sebastian; Ghahramani[10], Afshin; Harrison[11], Matthew Tom; Hochman[5], Zvi; Hoogenboom[12], Gerrit; Huang[13], Mingxia; Jing[14], Qi; Justes[15], Eric; Kersebaum[16,17,18], Kurt Christian; Launay[19], Marie; Lewan[20], Elisabet; Liu[11], Ke; Luo[21], Qunying; Mequanint[9], Fasil; Nendel[16,17,22], Claas; Padovan[8], Gloria; Olesen[23], Jørgen Eivind; Pullens[23], Johannes Wilhelmus Maria; Qian[14], Budong; Seserman[16], Diana-Maria; Shelia[12], Vakhtang; Souissi[24], Amir; Specka[16], Xenia; Wang[13], Jing; Weber[25], Tobias, K.D.; Weihermüller[26], Lutz, Seidel, Sabine J.[1]

[1]Institute of Crop Science and Resource Conservation, University of Bonn, Germany

[2]Natural Resources Institute Finland (Luke), Helsinki, Finland

[3]Julius Kühn Institute (JKI) – Federal Research Centre for Cultivated Plants, Institute for Crop and Soil Science , Braunschweig, Germany

[4]INRAE, UMR 1114 EMMAH, Avignon, France

[5]CSIRO Agriculture and Food, Brisbane, Queensland, Australia

[6]Technical University of Munich, Department of Life Science Engineering, Digital Agriculture, 85354 Freising, Germany

[7]Plant Sciences & TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium

[8]Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, Italy

[9]Institute of Soil Science and Land Evaluation, Biogeophysics, University of Hohenheim, Stuttgart, Germany

[10]Centre for Sustainable Agricultural Systems, Institute for Life Sciences and the Environment, University

of Southern Queensland, Toowoomba, Queensland, Australia

[11]Tasmanian Institute of Agriculture, University of Tasmania, Launceston, Tasmania, Australia, 7248

[12]Agricultural and Biological Engineering Department, University of Florida, Gainesville, Florida, USA &

Global Food Systems Institute, University of Florida, Gainesville, Florida, USA

[13]College of Resources and Environmental Sciences, China Agricultural University, Beijing, China

[14]Ottawa Research and Development Centre, Agriculture and Agri-Food Canada, Ottawa, Canada

[15]CIRAD, Univ Montpellier, Persyst Department, Montpellier, France

[16]Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

[17]Global Change Research Institute CAS, Brno, Czech Republic

[18]Institute for Tropical Plant Production and Agricultural Systems Modelling, Georg-August University,

Göttingen, Germany

[19]INRAE, US 1116 AgroClim, Avignon, France

[20]Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Uppsala,

Sweden

[21]Hillridge Technology Pty Ltd, Sydney, Australia

[22]Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[23]Department of Agroecology, Aarhus University, Tjele, Denmark

[24]Swift Current Research and Development Centre, Agriculture and Agri-Food Canada, Swift Current,

Saskatchewan, Canada

[25] Soil Science Section, Faculty of Organic Agricultural Sciences, University of Kassel, Germany (before [9])

[26]Institute of Bio- and Geosciences - IBG-3, Agrosphere, Forschungszentrum Jülich GmbH, Jülich,

Germany

*Corresponding author. Email: taru.palosuo@luke.fi

Author contribution:

Daniel Wallach: Conceptualization, methodology, project administration, writing - original draft,

validation. Taru Palosuo, Henrike Mielenz, Peter Thorburn, Samuel Buis, Sabine Seidel:

Conceptualization, methodology, project administration, writing, review &editing. All other

authors: expertise on models and simulation results, writing, review & editing.

Abstract

Crop phenology has a major influence on crop yield and is a major aspect of crop response to global warming. Process-based models of phenology are often used to predict the effect of weather on the development rate of crops through their growth phases, but such models are associated with large uncertainties, as demonstrated by the large variability between simulation results of different models. The purpose of this study is to estimate the relative importance of model structure uncertainty (due to uncertainty in the model equations) and model parameter uncertainty. Previous studies have assumed that the choice of model parameters to be fitted to data is fixed, and have evaluated the effect of uncertainty in those parameters. This underestimates parameter variance, as it ignores uncertainty in many aspects of the calibration approach, in particular the choice of parameters to estimate, as well as uncertainty in the parameters not fitted by calibration. Here, we propose and apply two approaches for estimating parameter variance that take into account uncertainty in all aspects of calibration, and, for one approach, uncertainty in parameters not fitted by calibration. Both approaches are based on previously reported large multi-model studies. The first approach uses components of variance analysis, which is possible because these studies included different modeling groups using the same model structure. The second approach is based on a study where multiple modeling groups each applied two different calibration procedures. The variance calculated from the two calibration approaches gives an estimate of parameter variance. Both methods give an overall result of 69% of total variance coming from parameter variance, for those simulated variables for which there are observed data. We conclude that in order to reduce uncertainty in crop phenology predictions, improving and sharing calibration procedures is as or more important than improving model structure.

Keywords: Crop modeling, Model calibration, Parameter uncertainty, Phenology, Structure uncertainty, Wheat

1. Introduction

Crop phenology has a major effect on crop yield and is a major aspect of crop response to global warming (Cleland et al., 2007; Liu et al., 2022; Parmesan, 2006). Process-based models of phenology, usually embedded in crop or agroecosystem models, are often used to predict the effect of weather on phenology (Bindi et al., 2015; Muleke et al., 2022). It has become common to study crop performance in general and crop phenology, in particular, using multi-model ensembles, where multiple models are provided with identical inputs and the range of simulated outputs is examined. These studies have shown that the variability in simulated values is substantial (Asseng et al., 2013; Durand et al., 2018; Sándor et al., 2020; Wallach et al., 2021a, 2021b). In several studies where multiple crop models were combined with multiple global circulation models (GCMs), it was been found that the uncertainty contribution from crop models is as large or larger than that from the GCMs (Asseng et al., 2013; Müller et al., 2021; Wang et al., 2020). A reduction in uncertainty from crop models is necessary in order to increase confidence in the results of these models, especially for the simulation of climate change impact assessment and options for adaptation (Maiorano et al., 2017; Rötter et al., 2011).

For given inputs (i.e. weather, crop management, soil characteristics and initial conditions), the uncertainty in crop model simulations has two origins, the uncertainty in crop model structure, which encompasses both the processes included in the model and the specific forms of the equations used to model those processes, and the uncertainty in the parameter values. Only relatively few studies have targeted the relative importance of both structure and parameter uncertainty, and those studies mostly indicate that the contribution of parameter uncertainty is rather small compared to the contribution from structure uncertainty (Tao et al., 2018; Wallach et al., 2017; Xiong et al., 2020; Zhang et al., 2017). Here, we argue that those studies do not take into account the full extent of parameter uncertainty, and we explain why it is difficult to do so. Therefore, we present here a new analysis, based on multi-model

ensemble simulation studies of crop phenology, that show that parameter uncertainty is the major

contributor to overall simulation uncertainty.

Model structure uncertainty refers to the range of plausible model structures, where a

"plausible" structure is one that would be deemed acceptable by researchers knowledgeable in the field

of crop modeling. Often an explicit or implicit requirement for participating in multi-model studies is that

the model structure has been published in the scientific literature, and has thus undergone peer review.

This can be considered to satisfy the plausibility condition. We also assume that in a multi-model

ensemble study, the represented model structures are a random sample from the population of

plausible structures, even though this assumption may be hard to fully justify. The problem is that crop

multi-model ensembles, like those in other fields, are generally "ensembles of opportunity" (Knutti,

2010; Wallach et al., 2016). That is, whichever modeling groups wish to participate are accepted. In fact,

this openness is a principle of the Agricultural Modeling Intercomparison and Improvement Project

(AgMIP, Rosenzweig et al., 2013). As a result, the ensemble may well have the same model structure

represented more than once, or there may be complex relationships among models, where some models

are derived from others or share similarities in structure (Fronzek et al., 2018; Holzworth et al., 2014; van

Ittersum et al., 2003) so that the structures are not independent. There have been attempts to account

for model interrelatedness in climate modeling (Sanderson et al., 2015) , but this has not been done for

crop models so far.

Parameter uncertainty for a given model structure and data set refers to the range of plausible

parameter vectors for that structure and data. Here "plausible" means that research teams working with

that structure would consider those parameters, or the way in which the parameters are obtained, i.e.

model calibration, reasonable. In almost all multi-model ensemble studies, models are calibrated using

some data set (which is the same for all models). Thus, the uncertainty in parameterization is due to

variability in the calibration method and in the choice of default parameter values (the values of those

parameters that are not affected by calibration).

To study the consequences of uncertainty in model structure and/or in model parameters, one

needs a sample drawn from the range of plausible structures and/or from the range of plausible

parameters for one or more structures. As discussed, the models represented in multi-model ensemble

studies are generally accepted as a sample from plausible structures. How to sample from plausible

parameter vectors on the other hand is not clear.

Wallach et al. (2022) identified several steps in crop model calibration, including the choice

of default parameter values, the choice of observed data to use in the objective function for frequentist

calibration, the choice of parameters to be estimated from the data and finally the choice of algorithm

and software for doing the calibration. They showed that different modeling groups made different

choices, and that even different modeling groups using the same model structure made different

choices. We suggest that sampling from plausible parameter vectors thus involves sampling from the

plausible choices for the default parameter values and from the plausible choices at each step of the

calibration procedure.

There has been a range of previous studies that have sampled model parameters Zhang et al.

(2017) used five rice phenology models and generated 50 randomly perturbed sets of parameter values

for each to obtain a sample of parameter values. Wallach et al. (2017) estimated parameters for two rice

phenology models using weighted least squares, then generated samples of parameters using the

estimated variance-covariance matrix of the estimated parameters. Tao et al. (2018) using seven barley

models identified a reasonable range for each parameter and generated a super ensemble, where each

model was associated with multiple parameter values drawn from the range of that parameter. Xiong et

al. (2020) compared three different crop models and four parametrization strategies that differed in the

amount of data available for calibration. Other studies have specifically considered the way in which the

model parameters are estimated. Kawakita et al. (2019), in a study of time to heading in wheat, assessed

three models and three different optimization algorithms for calibration, leading to three different

parameter vectors for each model. Jha et al. (2022) compared three calibration approaches for CERES

Maize, namely grid search, GLUE, and a genetic algorithm. Gao et al. (2020) compared the predictions

and uncertainty estimates for the CERES Rice model using a weighted least squares calibration method,

the GLUE method, and a Bayesian approach. Harrison et al. (2019) compared search techniques for

calibrating the APSIM model. In these studies, the choice of parameters to be treated as uncertain, as

well as the choices for most of the calibration steps listed above, are taken as given. Also, no uncertainty

in the default parameter values is taken into account. Thus those studies ignore several causes of

uncertainty in parameters that exist in multi-model ensemble studies (Albanito et al., 2022; Wallach et

al., 2021c) and could be important, and thus, those studies are likely to underestimate the importance of

parameter uncertainty.

A straightforward approach to estimating parameter and model structure uncertainty would be

to specifically create a multi-model ensemble in which each model structure is used by more than one

modeling group, each of which independently estimates the model parameters. In practice, this is

difficult to achieve. Confalonieri et al. (2016) organized a study with five crop models and 20 users,

leading to different model parameterizations. However, in that case, the groups responsible for

calibration were students and not model experts, so the parameter vectors might not be considered

"plausible" by model experts.

The goal of this study was to obtain estimates of parameter uncertainty, and of the relative

importance of parameter and structure uncertainty, which are more realistic than previous estimates.

The specific application is the estimation of wheat phenology in multiple environments. Two different

approaches are employed. The first is based on two multi-model ensemble studies which were not

specifically designed to study parameter uncertainty, but where it turned out that some of the model

structures were used by more than one modeling team. Those teams independently chose default

parameter values and the calibration procedure. The differences between modeling groups using the

same structure give information about parameter uncertainty. The second approach is based on a study

where multiple modeling teams each used two different calibration procedures. The differences

between the two procedures are used to estimate parameter uncertainty.

178   2.   Materials and Methods

179   2.1 Data

This study revisits three simulation experiments on wheat phenology that have been previously

reported (Wallach et al., 2021a, 2021b, 2022). In the previous studies, the emphasis was on prediction

accuracy, whereas in the present study the simulation results are reanalyzed with the objective of

estimating the contribution of parameter uncertainty to the overall uncertainty of the phenology

prediction.

The simulations concern three data sets (**Table 1**). Two of the data sets are from winter wheat

variety trials in France. These data sets have identical structures (same sites and sowing dates) but

concern two different varieties, "Apache" and "Bermude". Both varieties are of intermediate precocity,

but Bermude has a longer cycle. The observed data from each environment consist of daes of stem

elongation (BBCH30 on the BBCH scale, Meier, 1997) and middle of heading (BBCH55). The third data set

is from a multi-site sowing date experiment on spring wheat variety "Janz" in Australia. The original data

were observations of growth stage at weekly intervals. The data provided to modeling groups were the

dates of each BBCH stage, from the first to the last observed stage, derived from interpolation of the

original data. All data sets were split into two subsets, one for calibration and the other for evaluation.

The calibration and evaluation data sets had neither site nor year in common. Thus evaluation was a true

test of how well models could predict for out-of-sample conditions.

Table 1

Data set	Number of calibration environments	Number of evaluation environments	Observed development stages*	Simulated development stages used for analysis*
France Variety Apache	14	8	BBCH30, BBCH55	BBCH10, BBCH30, BBCH55
France Variety Bermude	14	8	BBCH30, BBCH55	BBCH10, BBCH30, BBCH55
Australia Variety Janz	24	18	BBCH stage at weekly intervals	BBCH10, BBCH30, BBCH65, BBCH90

197   *The development stages, using the BBCH scale (Meier, 1997) are emergence (BBCH10), start of

198   stem elongation (BBCH30), middle of heading (BBCH55), full flowering (BBCH65), and maturity (BBCH90).

199   **Table 1.** The three data sets used in the multi-model ensemble studies analyzed here.

200   **2.2 Participating modeling groups and model structures**

201   The call for participants in the first study was published using the mailing lists of several models

and of AgMIP. All modeling groups that volunteered (a modeling group is one or more individuals that

work together to run a model) were accepted. All groups used model structures that have been

published. A total of 30 modeling groups using 23 different structures (SI) participated in one or more of

the simulation experiments. In all cases, the phenology models were embedded in general crop models,

so model structure here refers to the overall crop model. Most but not all of the groups participated in

all the studies.

No specific attempt was made to encourage or discourage participation by different groups that

used the same structure. In the event, there were two or three (depending on the simulation study)

model structures were used by more than one group. In particular, in the first two studies three

structures were used by 2-4 different modeling groups (Table 2).

In all studies, it was decided to identify the different modeling groups were identified by a code

(M1, M2 etc.) without further identification of the group or giving the name of the model structure used.

The same codes are used here. In addition, we assign here codes to the different model structures (S1,

S2, etc.). The same model structure can be associated with more than one modeling group.

Table 2

Simulation study	Data set (country, cultivar)	Calibration procedure	Number of modeling groups	Repeated structures*
1	France, Apache	Usual	27	S2 (4), S4 (3), S18 (2)
1	France, Bermude	Usual	27	S2 (4), S4 (3), S18 (2)
2	Australia, Janz	Usual	28	S2 (3), S4 (3), S18 (2)
3	France, Apache	Protocol	19	S2 (3), S4 (2)
3	France, Bermude	Protocol	19	S2 (3), S4 (2)
3	Australia, Janz	Protocol	19	S2 (3), S4 (2)

217   *The model structures used by more than one modeling group, and in parentheses, the number of

218   groups that used that structure.

219   **Table 2.** The three different simulation studies analyzed here.

2.3 Simulation studies

In the first simulation study (Wallach et al., 2021a), based on the French data sets, each participating modeling group implemented their usual calibration method to calibrate their model using the calibration data subsets and then simulated days from sowing to phenological stages BBCH10, BBCH30, and BBCH55 for both varieties and for both the calibration and evaluation environments (Table 2). In the second study (Wallach et al., 2021b), each participating modeling group again used their usual calibration method and then simulated days from sowing to phenological stages BBCH10, BBCH30, BBCH65, and BBCH90 for the Australian environments. Neither the French nor the Australian data sets had observations of BBCH10, which was included in the variables to simulate in order to have an example of a simulated stage without calibration data.

The detailed descriptions of the calibration procedures used by each team in the first two studies were used to identify the range of decisions for each calibration step. Based on an analysis of the different choices, and following discussions with the participating teams, a new detailed calibration procedure (hereafter the calibration "protocol") was proposed and applied to all three data sets by all participating teams. Any group that participated in the third study but that had not participated in the first two studies first did the calibration using their usual method, before applying the protocol.

The proposed calibration protocol is described in detail in Wallach et al. (2022). The protocol covers all steps in phenology model calibration and includes templates for documenting the different steps. A major innovation, compared to the usual calibration procedures, is in the choice of parameters to estimate. In the protocol, this is done in two stages. First, parameters that play an almost additive role, i.e.affect all environments to a similar extent are identified. At most, there can be as many almost additive parameters as the number of different variables used for calibration (for example, two almost additive parameters if BBCH30 and BBCH55 are used for calibration). In most cases, the almost additive parameters are degree days to development stages at or near the observed stages. Those parameters

are estimated based on the calibration data. This should remove bias in the simulations, or nearly do so.

Then a series of candidate parameters, thought likely to explain part of the remaining unexplained

variability between environments, is identified, and those parameters are tested one by one. If the

estimation of a candidate parameter, together with all previously chosen parameters, sufficiently

improves the fit to the data, as measured by the Bayesian Information Criterion (BIC), it is included in the

final list of estimated parameters. If not, it is kept at its default value. A comparison between usual and

protocol calibration for the participants in the third study showed major differences between the two.

For example, the number of estimated parameters differed between the two procedures for the great

majority of participants (Wallach et al., 2021c).

2.4 Estimation of structure and parameter variance

Variance components approach

For a given environment, with given inputs, the variability in the simulated values comes from

the variability in structures and parameters. We assume that for each simulation experiment, the model

structures are a random sample from a plausible structure. It is further assumed that the parameter

vector used by each modeling group in simulation studies 1 and 2 is an independent random draw from

plausible parameter vectors for the corresponding model structure, consisting of a random draw from

plausible default parameter values and a random draw from plausible calibration approaches.

Presumably all the modeling teams were well-versed in using their model, which fulfills the plausibility

requirement for the parameters.

The statistical model for a simulated value is then

264
$$y_{sp} = \mu + \alpha_s + \beta_{sp} \tag{1}$$

where y_{sp} is the simulated value for model structure *s* and parameter vector *p*, μ is the overall mean ,a_s

is the effect of model structure and β_{sp} is the effect of parameter vector nested within the model

structure. There is no residual term because these are simulated values with no measurement error. We

treat a_s and β_{sp} as random effects, since we are interested in the variability in the population not in the

specific model structures or parameter vectors in the sample. The variances are noted

$\mathrm{var}(\alpha_s) = \sigma^2_{structure}$ for all *s* and $\mathrm{var}(\beta_{sp}) = \sigma^2_{parameters}$ for all *sp*. The random terms are assumed

independent so that the total variance for one environment and one simulated variable is the sum of

model structure variance and of parameter variance (Scheffé, 1959):

$$\sigma^2_{total} = \sigma^2_{structure} + \sigma^2_{parameters} \tag{2}$$

To estimate parameter and model structure variance simultaneously , we use the lmer function

of the lme4 package (Bates et al., 2015) in R (R Core Team, 2017), which is designed to analyze linear

mixed effect models (SI). This components of variance approach to estimating parameter variance is not

used for simulation study 3, since there were too few model structures used by more than one modeling

group. An illustration of the data used in this approach is given in Figure 1. As that figure shows, three of

the model structures (S2, S4, and S18) were used by more than one modeling group. It is the variability

between the different groups using the same model structure that allows us to estimate parameter

variance. The results for all groups contribute to the estimation of structure variance. Note, that the

simplified approach to estimating parameter uncertainty using these same data in Wallach et al. (2021a,

2021b) did not fully take into account the contribution of parameter uncertainty.

Figure 1

Figure 2. Example of data used for variance components approach to estimating parameter and structure

uncertainty. Simulation of days from sowing to mid-heading (stage BBCH55) for variety Apache in one

environment from simulation study 1. All modeling groups used the same inputs. The x-axis shows the

model structure. In most cases, each structure was used by a single modeling team, but structures S2, S4,

and S18 were used respectively by 4, 3, and 2 teams (data shown as stars). It is the variability between

simulations from different groups using those model structures that is the basis for estimating parameter

uncertainty. The horizontal line is the average of all simulated values.

Comparison of usual and protocol calibration

The second estimation of parameter uncertainty is based on all three simulation studies (see

Table 2). In studies 1 and 2 Every group used their usual calibration approach and in this study used the

newly proposed protocol calibration approach to obtain parameters for each data set. The estimate of

parameter uncertainty here is based on the difference between these two calibration methods. The two

simulation results for every modeling group for one environment are shown in Figure 2.

Each group's usual calibration approach can be considered to be a random draw from plausible

calibration approaches for their model structure. The protocol calibration approach, which was

developed in collaboration with all participating modeling teams, is also by construction a plausible

calibration approach. It is essentially independent of the usual approach since the contribution of any

single group to the protocol was limited. Furthermore, the protocol did not mimic closely any single

usual calibration approach. Therefore, for each modeling group we treat the usual and the protocol

calibration as two random draws from the population of plausible calibration approaches for the model

structure used by that group. For each environment, we calculate the variance of the two calibration

approaches, noted var_i for modeling group i (eq. 3), and then average over modeling groups to obtain an

estimated parameter variance for that environment (eq. 4).

309
$$\mathrm{var}_i = \left(y^{usual} - \bar{y} \right)^2 + \left(y^{protocol} - \bar{y} \right)^2 \tag{3}$$

310
$$\hat{\sigma}^2_{parameters} = 1/n \sum_{i=1}^{R} \mathrm{var}_i \tag{4}$$

where $\bar{y} = (y^{usual} + y^{protocol})/2$.

We do not have a rigorous method of estimating structure variance from these data. The

difficulty is that the assumptions underlying the random effects model described above are not satisfied;

the parameter vectors obtained by protocol calibration are not independent between modeling groups,

since all use the same protocol. We therefore use, for each environment, the estimate of structure

variance from the variance components approach above.

Figure 2

Figure 2. Example of data used for usual compared to protocol approach to estimating parameter

uncertainty Simulated days from sowing to stem elongation (stage BBCH30) for variety Bermude in one

environment from simulation study 3, which compared usual and protocol calibration for each modeling

group. Each code (M1, etc.) refers to a specific modeling group (x-axis). For each modeling group, there

are two simulated values, one using parameters based on usual calibration and the other using parameters

based on protocol calibration. The differences between the two values are used to estimate parameter

variance. The horizontal line shows the mean of simulations using usual calibration.

3.   Results

First, all calculations were done for the calibration and evaluation data subsets separately,

however we found that there were no systematic differences between the two (Fig. S1 and Fig. S2). All

results here are therefore based on the combined calibration and evaluation data.

Figure 3 shows parameter variance as a fraction of total variance for each environment of the

two French data sets, using variance components or usual compared to protocol calibration. Figure 4

shows the same information for the Australian data set. Table 3 and Table 4 show parameter variance,

and parameter variance as a fraction of total variance, averaged over all environments for the French

and Australian data sets, respectively.

Table 4 shows results averaged over all environments of both the French and Australian data

sets. Separate values are given for BBCH10 (for which there was no observed data) and for all other

stages.

Two aspects of the results in Table 3 and Table 4 are worth emphasizing. First, the estimated

contribution of parameter variance to total variance is always much less for BBCH10 than for the other

simulated development stages. This is the case for both approaches, but the difference is particularly

important for the approach based on comparison of usual and protocol calibration.

Secondly, for development stages other than BBCH10, the estimated contribution of parameter

uncertainty is large. It ranges from 24 to 93%, and is larger than 50% for 11 out of the 14 combinations of

data set, development stage and approach for estimating the contribution of parameter variance.

Figure 3

Figure 3. Fraction of total variance due to parameter variance, for the French data sets, based on two

different approaches: based on variance components (top) or based on the comparison of usual and

protocol calibration (bottom). A10, A30, and A55 (B10, B30, B55) refer respectively to simulations of

days from sowing to emergence, to start of stem elongation and to middle of heading for the variety

Apache (Bermude). Boxes indicate the lower and upper quartiles. The solid line within the box is the

median. Whiskers indicate the most extreme data point which is no more than 1.5 times the interquartile

range from the box, and the outlier dots are those observations that are beyond that range.

     Figure 4

Figure 4. Fraction of total variance due to parameter variance, for the Australia data set, based on two

different approaches: based on variance components (top) or based on comparison of usual and protocol

calibration (bottom). Simulations are of days from sowing to emergence, start pf stem elongation, full

flowering and maturity. Boxes indicate the lower and upper quartiles. The solid line within the box is the

median. Whiskers indicate the most extreme data point which is no more than 1.5 times the interquartile

range from the box, and the outlier dots are those observations that are beyond that range.

Table 3

		Apache			Bermude		
	Development stage	BBCH10	BBCH30	BBCH55	BBCH10	BBCH30	BBCH55
Method of estimating parameter variance as fraction of total							
Variance components	Average number of modeling groups	23	25	27	23	25	27
	Number of model structures	17	19	21	17	19	21
	Parameter variance	11.74	52.94	26.77	12.05	71.25	36.41
	Structure variance	5.05	4.29	9.65	4.99	10.86	12.73
	Parameter variance as fraction of total	0.55	0.93	0.81	0.56	0.92	0.8
Usual compared to protocol calibration	Average number of modeling groups	14	16	17	14	16	16.9
	Parameter variance	0.75	16.62	10.35	2.57	28.52	10.01
	Parameter variance as fraction of total	0.28	0.88	0.71	0.39	0.89	0.65

365 **Table 3.** Parameter and structure variance averaged over environments of the French data sets. The variance components estimate is

366 based on ensemble studies where some model structures were used by multiple modeling groups. The usual compared to protocol

367 estimate is based on comparing usual and protocol calibration results for each modeling group.

368   **Table 4**

Method for estimating parameter variance as fraction of total	Development stage	BBCH10	BBCH30	BBCH65	BBCH90
Variance components	Average number of modeling groups	22.95	23.93	27.64	26.05
	Average number of model structures	17.95	19.93	22.64	21.05
	Parameter variance	40.69	102.88	47.1	30.3
	Structure variance	276.33	45.74	48.4	90.71
	Parameter variance as fraction of total	0.12	0.66	0.49	0.24
Usual compared to protocol calibration	Average Number of modeling groups	13.9	16.9	18.9	17.3
	Parameter variance	8.41	62.54	67.12	80.76
	Parameter variance as fraction of total	0.026	0.65	0.61	0.46

370   **Table 4.** Parameter and structure variance averaged over environments of the Australian data set. The variance components estimate is

371   based on ensemble studies where some model structures were used by multiple modeling groups. The usual compared to protocol

372   estimate is based on comparing usual and protocol calibration results for each modeling group.

373   **Table 5**

Method for estimating parameter variance as fraction of total	BBCH10	All other stages
Variance components	0.41	0.69
Usual compared to protocol calibration	0.23	0.69

375   **Table 5.** Fraction of total variance due to parameter variance based on two estimation procedures:

376   variance components and comparison of usual and protocol calibration. Averages are over all

377   environments of both the French and Australian data sets, either for the development stage BBCH10 or for

378   all other development stages.

380   4.   Discussion

There are relatively few studies that have estimated separately the contributions of parameter

uncertainty and model structure uncertainty to total simulation uncertainty in ensemble studies with

crop models or crop phenology models. Those studies focused on parameter variability when the list of

parameters to be estimated is fixed. This includes studies where the parameter distribution is based on

expert opinion (Tao et al., 2018; Zhang et al., 2017) where parameters differ due to the use of different

levels of gridded information (Xiong et al., 2020) or where the variance-covariance matrix of the

parameter vector is estimated during least squares parameter estimation (Wallach et al., 2017). These

studies have systematically found that structural uncertainty is substantially more important than

parameter uncertainty.

390  Those studies sampled from the distribution of values for some fixed subset of model

391 parameters. However, this does not really take into account the full parameter uncertainty in crop or

392 phenology models, which arises from all aspects of the calibration approach, not just from uncertainty in

393 a fixed subset of parameters (Albanito et al., 2022; Wallach et al., 2021c). In addition, there is

394 uncertainty in the values of the fixed parameters. The two approaches we propose here take into

395 account more realistically the full uncertainty in model parameterization.

396  Our first approach is based on variability in simulated values between different modeling groups

397 using the same model structure. The underlying hypothesis is that the most realistic way to sample true

398 parameter uncertainty in crop or phenology models is to sample established crop modeling groups using

399 the same model structure. If they are provided with the same input data and the same observational

400 data for calibration, the parameter vectors they obtain is by construction a sample from plausible

401 parameter vectors for that structure and those data. It was found in the studies that were analyzed here,

402 that different groups took different decisions at all steps in the calibration procedure, including the

403 choice of which parameters to estimate, the form of the objective function, and the algorithm used for

404 searching for the best parameters (Wallach et al., 2021c). This confirms that uncertainty in calibrated

405 parameters arises from uncertainty in all aspects of the calibration procedure. Our approach

406 automatically takes all those differences in calibration procedure into account, as well as differences in

407 the choice of default parameter values. A limitation of the present study is that only three model

408 structures were used by more than one group, and at most four groups used the same model structure.

409  Our second estimate of parameter uncertainty is based on a comparison, for multiple modeling

410 groups, between two different calibration procedures, their usual procedure, and a new protocol

411 procedure. In general, there were major differences between the two calibration procedures for each

412 group (Wallach et al., 2022). Both are plausible in the sense that they were developed and used by

413 modeling teams well-versed in the use of their model structure. It seems reasonable then to assume that

for each modeling group, the two are a random sample of size two of plausible procedures for the

corresponding model structure. One limitation of this second estimate is that the same modeling group

implemented both calibration procedures. It seems likely that the default parameter values were mostly

unchanged between the two calibration approaches. Since the variability in the fixed parameter values is

not sampled, this approach underestimates the contribution of parameter uncertainty to the total

uncertainty.

In the estimates obtained here, parameter variance as a fraction of total variance is

systematically smaller for emergence (stage BBCH10) than for the other stages. The fraction is

particularly small for the estimates based on usual compared to protocol calibration. This is likely due to

the fact that there were no data for days to emergence, and so simulated days to emergence were

usually determined by the default parameter values. Probably, most modeling groups used the same

default values for usual and protocol calibration, leading to no difference in simulated days to

emergence between usual and protocol calibration. The contribution of parameter uncertainty to total

uncertainty for stages other than BBCH10, as estimated here, is similar for the two approaches to

estimation of parameter uncertainty. This suggests that when there are data related to a development

stage, differences between groups are mostly due to differences in calibration procedure rather than to

differences in default values.

The contribution of parameter uncertainty to total uncertainty, averaged over development

stages and data sets, is 69%. This implies that in a multi-model ensemble study simulating phenology,

over two thirds of the variability is due to parameter uncertainty and only 31%, or about one third is due

to structure uncertainty, i.e. to the variability among plausible model structures. The estimates obtained

here come from three different data sets, with 22 or 42 different environments. At least 19 modeling

groups participated in each study used here. There is substantial variability between environments and

simulated variables, but the range of situations analyzed gives some confidence in the overall conclusion,

that parameter uncertainty is the major contribution to overall model uncertainty

It is important to obtain information on the contribution of parameter uncertainty in other

conditions than those considered here. The most realistic estimate of parameter uncertainty is based on

independent modeling groups using the same model structure. One possibility would be to proactively

create multi-model ensembles where the same structure is represented by multiple modeling groups.

Another approach would be to take advantage of those multi-model ensemble studies where, by chance,

the same structure is represented by more than one modeling group. The variance components

calculation applied here could be applied quite generally to those studies. In some cases, the results of

past multi-model ensemble studies have been documented in data journals (e.g. Martre et al., 2018).

Those studies could be revisited in order to estimate parameter and structure uncertainty.

The major role of parameter uncertainty emphasizes the importance of documenting all

parameter values, both fixed and calibrated, for each model in each study, since these may have as much

or more effect on simulated values as the model equations. Without that information, the results of a

simulation study cannot be reproduced.

Much, though not all, of the parameter uncertainty, comes from the variability in calibration

approaches between modeling groups. The obvious path to the reduction of this source of uncertainty is

to have a single calibration approach for each model structure. Of course, this is only worthwhile if the

common calibration approach leads to a parameterized model that predicts well. There is no consensus

on the best calibration approach for crop models. However, there have been recent attempts to define a

calibration protocol that could be applied generally to phenology models, and that has been found both

to reduce uncertainty and to reduce mean squared error of prediction (Wallach et al., 2022).

The results here focus on simulating phenology, one component of crop models. Full crop

models simulate many other processes and not just development, which may lead to larger structure

uncertainty. However, full crop models have many more parameters than phenology models and

parameter estimation of an entire crop model is more complex, which may increase parameter

uncertainty. It is thus not clear how the results here scale up to more complex models. In any case,

however, it is important to take into account the full extent of parameter uncertainty in analyzing

structural versus parameter uncertainty for full crop models.

466   5.   Conclusions

The large uncertainties associated with crop models in general and crop phenology models in

particular clearly limit the utility and acceptance of these models for impact assessment and other

practical applications. Previous studies that have aimed at quantifying the separate contributions of

model structure and model parameter uncertainty have concluded that model parameter uncertainty is

relatively less important, leading to the implicit conclusion that research should be focused on the

processes and equations in crop models. We find here that when one takes into account the full

uncertainty in calibration procedure and also uncertainty in the default parameter values, parameter

uncertainty is the major cause of uncertainty in simulated values of crop phenology models The

development and application of improved, shared calibration procedures as well as increased attention

to the choice of default parameter values should thus be a major goal of the crop modeling community.

477   6.   Acknowledgements

This work was in part supported by the Collaborative Research Center 1253 CAMPOS (Project 7:

Stochastic Modelling Framework), funded by the German Research Foundation (DFG, Grant Agreement

SFB 1253/1 2017), the Academy of Finland through projects AICropPro (316172) and DivCSA (316215)

and Natural Resources Institute Finland (Luke) through a strategic project BoostIA, the BonaRes projects

''Soil3'' (BOMA 03037514) and "I4S" (031B053I) of the Federal Ministry of Education and Research

(BMBF), Germany, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under

Germany's Excellence Strategy - EXC 2070 – 390732324 EXC (PhenoRob), the project BiomassWeb of the

GlobeE programme (Grant number: FKZ031A258B) funded by the Federal Ministry of Education and

Research (BMBF, Germany), the INRA ACCAF meta-programme, the German Federal Ministry of

Education and Research (BMBF) in the framework of the funding measure "Soil as a Sustainable

Resource for the Bioeconomy –- BonaRes", project "BonaRes (Module B): BonaRes Centre for Soil

Research, subproject B" (grant 031B0511B), the National Key Research and Development Program of

China (2017YFD0300205), the National Science Foundation for Distinguished Young Scholars (31725020),

the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), the 111

project (B16026), and China Scholarship Council, the Agriculture and Agri-Food Canada's Project J-

002303 under the Interdepartmental Research Initiative in Agriculture, the DFG Research Unit FOR 1695

'Agricultural Landscapes under Global Climate Change – Processes and Feedbacks on a Regional Scale,

the U.S. Department of Agriculture National Institute of Food and Agriculture (award no. 2015-68007-

23133) and USDA/NIFA HATCH grant N. MCL02368, the National Key Research and Development

Program of China (2016YFD0300105), The Broadacre Agriculture Initiative, a research partnership

between University of Southern Queensland and the Queensland Department of Agriculture and

Fisheries, the Academy of Finland through project AI-CropPro (315896), the JPI FACCE MACSUR2 project,

funded by the Italian Ministry for Agricultural, Food, and Forestry Policies (D.M. 24064/7303/15 of

6/Nov/2015), the SYSTEMIC funded by JPI HDHL, JPI-OCEANS and FACCE-JPI under ERA-NET (n.696295),

the SustEs project funded by the Ministry of Education, Youth and Sports of the Czech Republic

(CZ.02.1.01/0.0/0.0/16_019/000797). The time of MTH and KL was supported by the Grains Research

and Development Corporation (Project UOT1906-002RTX). The observed Australian phenology data were

jointly funded by CSIRO, the Grains Research and Development Corporation (GRDC) under the "Adding

Value to GRDC's National Variety Trial Network" project (CSA00027). The order in which the donors are

listed is arbitrary.

7.   References

Albanito, F., McBey, D., Harrison, M., Smith, P., Ehrhardt, F., Bhatia, A., Bellocchi, G., Brilli, L., Carozzi, M.,

   Christie, K., Doltra, J., Dorich, C., Doro, L., Grace, P., Grant, B., Léonard, J., Liebig, M., Ludemann, C.,

   Martin, R., Meier, E., Meyer, R., De Antoni Migliorati, M., Myrgiotis, V., Recous, S., Sándor, R.,

   Snow, V., Soussana, J.-F., Smith, W.N., Fitton, N., 2022. How Modelers Model: the Overlooked

   Social and Human Dimensions in Model Intercomparison Studies. Environ. Sci. Technol. 56, 13485–

   13498. https://doi.org/10.1021/acs.est.2c02023

Asseng, S., Ewert, F., Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P.J.,

   Rötter, R.P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P.K., Angulo, C., Bertuzzi,

   P., Biernath, C., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J.,

   Hunt, L.A., Ingwersen, J., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Naresh Kumar, S., Nendel, C.,

   O'Leary, G., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A.,

   Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M.,

   Waha, K., Wallach, D., White, J.W., Williams, J.R., Wolf, J., 2013. Uncertainty in simulating wheat

   yields under climate change. Nat. Clim. Chang. 3, 827–832. https://doi.org/10.1038/nclimate1916

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J.

   Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bindi, M., Palosuo, T., Trnka, M., Semenov, M., 2015. Modelling climate change impacts on crop

production for food security. Clim. Res. 65, 3–5. https://doi.org/10.3354/cr01342

Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A., Schwartz, M.D., 2007. Shifting plant phenology in

response to global change. Trends Ecol. Evol. https://doi.org/10.1016/j.tree.2007.04.003

Confalonieri, R., Orlando, F., Paleari, L., Stella, T., Gilardelli, C., Movedi, E., Pagani, V., Cappelli, G.,

Vertemara, A., Alberti, L., Alberti, P., Atanassiu, S., Bonaiti, M., Cappelletti, G., Ceruti, M.,

Confalonieri, A., Corgatelli, G., Corti, P., Dell'Oro, M., Ghidoni, A., Lamarta, A., Maghini, A.,

Mambretti, M., Manchia, A., Massoni, G., Mutti, P., Pariani, S., Pasini, D., Pesenti, A., Pizzamiglio, G.,

Ravasio, A., Rea, A., Santorsola, D., Serafini, G., Slavazza, M., Acutis, M., 2016. Uncertainty in crop

model predictions: What is the role of users? Environ. Model. Softw. 81, 165–173.

https://doi.org/10.1016/j.envsoft.2016.04.009

Durand, J.-L., Delusca, K., Boote, K., Lizaso, J., Manderscheid, R., Weigel, H.J., Ruane, A.C., Rosenzweig, C.,

Jones, J., Ahuja, L., Anapalli, S., Basso, B., Baron, C., Bertuzzi, P., Biernath, C., Deryng, D., Ewert, F.,

Gaiser, T., Gayler, S., Heinlein, F., Kersebaum, K.C., Kim, S.-H., Müller, C., Nendel, C., Olioso, A.,

Priesack, E., Villegas, J.R., Ripoche, D., Rötter, R.P., Seidel, S.I., Srivastava, A., Tao, F., Timlin, D.,

Twine, T., Wang, E., Webber, H., Zhao, Z., 2018. How accurately do maize crop models simulate the

interactions of atmospheric CO_2 concentration levels with limited water supply on water use and

yield? Eur. J. Agron. 100, 67–75. https://doi.org/10.1016/J.EJA.2017.01.002

Fronzek, S., Pirttioja, N., Carter, T.R., Bindi, M., Hoffmann, H., Palosuo, T., Ruiz-Ramos, M., Tao, F., Trnka,

M., Acutis, M., Asseng, S., Baranowski, P., Basso, B., Bodin, P., Buis, S., Cammarano, D., Deligios, P.,

Destain, M.-F., Dumont, B., Ewert, F., Ferrise, R., François, L., Gaiser, T., Hlavinka, P., Jacquemin, I.,

Kersebaum, K.C., Kollas, C., Krzyszczak, J., Lorite, I.J., Minet, J., Minguez, M.I., Montesino, M.,

Moriondo, M., Müller, C., Nendel, C., Öztürk, I., Perego, A., Rodríguez, A., Ruane, A.C., Ruget, F.,

Sanna, M., Semenov, M.A., Slawinski, C., Stratonovitch, P., Supit, I., Waha, K., Wang, E., Wu, L.,

Zhao, Z., Rötter, R.P., 2018. Classifying multi-model wheat yield impact response surfaces showing

sensitivity to temperature and precipitation change. Agric. Syst. 159, 209–224.

https://doi.org/10.1016/J.AGSY.2017.08.004

Gao, Y., Wallach, D., Liu, B., Dingkuhn, M., Boote, K.J., Singh, U., Asseng, S., Kahveci, T., He, J., Zhang, R.,

Confalonieri, R., Hoogenboom, G., 2020. Comparison of three calibration methods for modeling rice

phenology. Agric. For. Meteorol. 280, 107785. https://doi.org/10.1016/J.AGRFORMET.2019.107785

Harrison, M.T., Roggero, P.P., Zavattaro, L., 2019. Simple, efficient and robust techniques for automatic

multi-objective function parameterisation: Case studies of local and global optimisation using

APSIM. Environ. Model. Softw.

Holzworth, D.P., Huth, N.I., deVoil, P.G., Zurcher, E.J., Herrmann, N.I., McLean, G., Chenu, K., van

Oosterom, E.J., Snow, V., Murphy, C., Moore, A.D., Brown, H., Whish, J.P.M., Verrall, S., Fainges, J.,

Bell, L.W., Peake, A.S., Poulton, P.L., Hochman, Z., Thorburn, P.J., Gaydon, D.S., Dalgliesh, N.P.,

Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li,

F.Y., Wang, E., Hammer, G.L., Robertson, M.J., Dimes, J.P., Whitbread, A.M., Hunt, J., van Rees, H.,

McClelland, T., Carberry, P.S., Hargreaves, J.N.G., MacLeod, N., McDonald, C., Harsdorf, J.,

Wedgwood, S., Keating, B.A., 2014. APSIM – Evolution towards a new generation of agricultural

systems simulation. Environ. Model. Softw. 62, 327–350.

https://doi.org/10.1016/j.envsoft.2014.07.009

Jha, P.K., Ines, A.V.M., Han, E., Cruz, R., Vara Prasad, P.V., 2022. A comparison of multiple calibration and

ensembling methods for estimating genetic coefficients of CERES-Rice to simulate phenology and

yields. F. Crop. Res. 284, 108560. https://doi.org/10.1016/J.FCR.2022.108560

Kawakita, S., Inaba, S., Takahashi, T., Kamada, E., Ishikawa, N., Takahashi, H., Okuno, R., 2019. Evaluation

of non-linear wheat development models and optimization methods for their parameter

determination. shiJournal Agric. Meteorol. 75, 120–128. https://doi.org/10.2480/agrmet.D-18-

00034

Knutti, R., 2010. The end of model democracy? Clim. Change 102, 395–404.

https://doi.org/10.1007/s10584-010-9800-2

Liu, Y., Bachofen, C., Wittwer, R., Silva Duarte, G., Sun, Q., Klaus, V.H., Buchmann, N., 2022. Using

PhenoCams to track crop phenology and explain the effects of different cropping systems on yield.

Agric. Syst. 195, 103306. https://doi.org/10.1016/J.AGSY.2021.103306

Maiorano, A., Martre, P., Asseng, S., Ewert, F., Müller, C., Rötter, R.P., Ruane, A.C., Semenov, M.A.,

Wallach, D., Wang, E., Alderman, P.D., Kassie, B.T., Biernath, C., Basso, B., Cammarano, D.,

Challinor, A.J., Doltra, J., Dumont, B., Rezaei, E.E., Gayler, S., Kersebaum, K.C., Kimball, B.A., Koehler,

A.-K., Liu, B., O'Leary, G.J., Olesen, J.E., Ottman, M.J., Priesack, E., Reynolds, M., Stratonovitch, P.,

Streck, T., Thorburn, P.J., Waha, K., Wall, G.W., White, J.W., Zhao, Z., Zhu, Y., 2017. Crop model

improvement reduces the uncertainty of the response to temperature of multi-model ensembles.

F. Crop. Res. 202. https://doi.org/10.1016/j.fcr.2016.05.001

Martre, P., Kimball, B.A., Ottman, M.J., Wall, G.W., White, J.W., Asseng, S., Ewert, F., Cammarano, D.,

Maiorano, A., Aggarwal, P.K., Anothai, J., Basso, B., Biernath, C., Challinor, A.J., De Sanctis, G.,

Doltra, J., Dumont, B., Fereres, E., Garcia-Vila, M., Gayler, S., Hoogenboom, G., Hunt, L.A.,

Izaurralde, R.C., Jabloun, M., Jones, C.D., Kassie, B.T., Kersebaum, K.C., Koehler, A.-K., Müller, C.,

Kumar, S.N., Liu, B., Lobell, D.B., Nendel, C., O'Leary, G., Olesen, J.E., Palosuo, T., Priesack, E.,

Rezaei, E.E., Ripoche, D., Rötter, R.P., Semenov, M.A., Stöckle, C., Stratonovitch, P., Streck, T., Supit,

I., Tao, F., Thorburn, P., Waha, K., Wang, E., Wolf, J., Zhao, Z., Zhu, Y., 2018. The Hot Serial Cereal

Experiment for modeling wheat response to temperature: field experiments and AgMIP-Wheat

multi-model simulations. Open Data J. Agric. Res. 4, 28–34.

https://doi.org/10.18174/odjar.v4i0.15830

Meier, U., 1997. Growth stages of mono- and dicotyledonous plants : BBCH-Monograph [WWW

Document]. URL

https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwjBo-

Ln4_r4AhVM3RoKHc-

TA5sQFnoECAcQAQ&url=https%3A%2F%2Fwww.politicheagricole.it%2Fflex%2FAppData%2FWebLi

ve%2FAgrometeo%2FMIEPFY800%2FBBCHengl2001.pdf&usg=AOvVaw1G_eUh_-qnVt91_qbj44d

(accessed 1.9.19).

Muleke, A., Harrison, M.T., de Voil, P., Hunt, I., Liu, K., Yanotti, M., Eisner, R., 2022. Earlier crop flowering

caused by global warming alleviated by irrigation. Environ. Res. Lett. 17, 044032.

https://doi.org/10.1088/1748-9326/ac5a66

Müller, C., Franke, J., Jägermeyr, J., Ruane, A.C., Elliott, J., Moyer, E., Heinke, J., Falloon, P.D., Folberth, C.,

Francois, L., Hank, T., Izaurralde, R.C., Jacquemin, I., Liu, W., Olin, S., Pugh, T.A.M., Williams, K.,

Zabel, F., 2021. Exploring uncertainties in global crop yield projections in a large ensemble of crop

models and CMIP5 and CMIP6 climate scenarios. Environ. Res. Lett. 16, 034040.

https://doi.org/10.1088/1748-9326/abd8fc

Parmesan, C., 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annu. Rev. Ecol.

Evol. Syst. 37, 637–669. https://doi.org/10.1146/annurev.ecolsys.37.091305.110100

R Core Team, 2017. A language and Environment for Statistical Computing.

Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P., Antle, J.M., Nelson, G.C.,

Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorria, G., Winter, J.M., 2013.

The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot

618  studies. Agric. For. Meteorol. 170, 166–182. https://doi.org/10.1016/j.agrformet.2012.09.011

619 Rötter, R.P., Carter, T.R., Olesen, J.E., Porter, J.R., 2011. Crop–climate models need an overhaul. Nat.

620  Clim. Chang. 1, 175–177. https://doi.org/10.1038/nclimate1152

621 Sanderson, B.M., Knutti, R., Caldwell, P., Sanderson, B.M., Knutti, R., Caldwell, P., 2015. Addressing

622  Interdependency in a Multimodel Ensemble by Interpolation of Model Properties. J. Clim. 28, 5150–

623  5170. https://doi.org/10.1175/JCLI-D-14-00361.1

624 Sándor, R., Ehrhardt, F., Grace, P., Recous, S., Smith, P., Snow, V., Soussana, J.-F., Basso, B., Bhatia, A.,

625  Brilli, L., Doltra, J., Dorich, C.D., Doro, L., Fitton, N., Grant, B., Harrison, M.T., Kirschbaum, M.U.F.,

626  Klumpp, K., Laville, P., Léonard, J., Martin, R., Massad, R.-S., Moore, A., Myrgiotis, V., Pattey, E.,

627  Rolinski, S., Sharp, J., Skiba, U., Smith, W., Wu, L., Zhang, Q., Bellocchi, G., 2020. Ensemble

628  modelling of carbon fluxes in grasslands and croplands. F. Crop. Res. 252, 107791.

629  https://doi.org/10.1016/j.fcr.2020.107791

630 Scheffé, H., 1959. The analysis of variance. John Wiley & Sons, New York.

631 Tao, F., Rötter, R.P., Palosuo, T., Gregorio Hernández Díaz-Ambrona, C., Mínguez, M.I., Semenov, M.A.,

632  Kersebaum, K.C., Nendel, C., Specka, X., Hoffmann, H., Ewert, F., Dambreville, A., Martre, P.,

633  Rodríguez, L., Ruiz-Ramos, M., Gaiser, T., Höhn, J.G., Salo, T., Ferrise, R., Bindi, M., Cammarano, D.,

634  Schulman, A.H., 2018. Contribution of crop model structure, parameters and climate projections to

635  uncertainty in climate change impact assessments. Glob. Chang. Biol. 24, 1291–1307.

636  https://doi.org/10.1111/gcb.14019

637 van Ittersum, M.., Leffelaar, P.., van Keulen, H., Kropff, M.., Bastiaans, L., Goudriaan, J., 2003. On

638  approaches and applications of the Wageningen crop models. Eur. J. Agron. 18, 201–234.

639  https://doi.org/10.1016/S1161-0301(02)00106-5

Wallach, D., Mearns, L.O., Ruane, A.C., Rötter, R.P., Asseng, S., 2016. Lessons from climate modeling on

the design and use of ensembles for crop modeling. Clim. Change 139, 551–564.

https://doi.org/10.1007/s10584-016-1803-1

Wallach, D., Nissanka, S.P., Karunaratne, A.S., Weerakoon, W.M.W., Thorburn, P.J., Boote, K.J., Jones,

J.W., 2017. Accounting for both parameter and model structure uncertainty in crop model

predictions of phenology: A case study on rice. Eur. J. Agron. 88.

https://doi.org/10.1016/j.eja.2016.05.013

Wallach, D., Palosuo, T., Thorburn, P., Gourdain, E., Asseng, S., Basso, B., Buis, S., Crout, N., Dibari, C.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani, A., Hochman, Z., Hoek, S.,

Hoogenboom, G., Horan, H., Huang, M., Jabloun, M., Jing, Q., Justes, E., Kersebaum, K.C.,

Klosterhalfen, A., Launay, M., Luo, Q., Maestrini, B., Mielenz, H., Moriondo, M., Nariman Zadeh, H.,

Olesen, J.E., Poyda, A., Priesack, E., Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A.,

Specka, X., Srivastava, A.K., Stella, T., Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D.,

Weihermüller, L., de Wit, A., Wöhling, T., Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021a. How well do

crop modeling groups predict wheat phenology, given calibration data from the target population?

Eur. J. Agron. 124, 126195. https://doi.org/https://doi.org/10.1016/j.eja.2020.126195

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Andrianasolo, F., Asseng, S., Basso, B., Buis, S., Crout,

N., Dumont, B., Ferrise, R., Gaiser, T., Gayler, S., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G.,

Huang, M., Jabloun, M., Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.C., Launay, M., Lewan, E.,

Luo, Q., Maestrini, B., Moriondo, M., Olesen, J.E., Padovan, G., Poyda, A., Priesack, E., Pullens,

J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Kumar Srivastava, A., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021b. Multi-model evaluation of phenology prediction for

wheat in Australia. Agric. For. Meteorol. 298–299, 108289.

https://doi.org/10.1016/J.AGRFORMET.2020.108289

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Gourdain, E., Andrianasolo, F., Asseng, S., Basso, B.,

Buis, S., Crout, N., Dibari, C., Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani,

A., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G., Huang, M., Jabloun, M., Jansson, P.-E., Jing,

Q., Justes, E., Kersebaum, K.C., Klosterhalfen, A., Launay, M., Lewan, E., Luo, Q., Maestrini, B.,

Mielenz, H., Moriondo, M., Nariman Zadeh, H., Padovan, G., Olesen, J.E., Poyda, A., Priesack, E.,

Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Srivastava, A.K., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021c. The chaos in calibrating crop models: Lessons learned

from a multi-model calibration exercise. Environ. Model. Softw. 145, 105206.

https://doi.org/10.1016/J.ENVSOFT.2021.105206

Wallach, D., Palosuo, T., Thorburn, P., Mielenz, H., Buis, S., Hochman, Z., Gourdain, E., Andrianasolo, F.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Hiremath, S., Horan, H., Hoogenboom, G.,

Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.-C., Launay, M., Lewan, E., Mequanint, F.,

Moriondo, M., Nendel, C., Padovan, G., Qian, B., Schütze, N., Seserman, D.-M., Shelia, V., Souissi, A.,

Specka, X., Srivastava, A.K., Trombi, G., Weber, T.K.D., Weihermüller, L., Wöhling, T., Seidel, S.J.,

2022. Calibration of crop phenology models: Going beyond recommendations. bioRxiv

2022.06.08.495355. https://doi.org/10.1101/2022.06.08.495355

Wang, B., Feng, P., Liu, D.L., O'Leary, G.J., Macadam, I., Waters, C., Asseng, S., Cowie, A., Jiang, T., Xiao,

D., Ruan, H., He, J., Yu, Q., 2020. Sources of uncertainty for wheat yield projections under future

climate are site-specific. Nat. Food 1, 720–728. https://doi.org/10.1038/s43016-020-00181-w

Xiong, W., Asseng, S., Hoogenboom, G., Hernandez-Ochoa, I., Robertson, R., Sonder, K., Pequeno, D.,

Reynolds, M., Gerard, B., 2020. Different uncertainty distribution between high and low latitudes in

modelling warming impacts on wheat. Nat. Food 1, 63–69. https://doi.org/10.1038/s43016-019-

0004-2

Zhang, S., Tao, F., Zhang, Z., 2017. Uncertainty from model structure is larger than that from model

parameters in simulating rice phenology in China. Eur. J. Agron. 87, 30–39.

https://doi.org/10.1016/j.eja.2017.04.004