Why is there so much variability in crop multi-model studies?

Wallach[a], Daniel; Palosuo[b,*], Taru; Mielenz[c], Henrike; Buis[d], Samuel; Thorburn[e], Peter; Asseng[f],

Senthold; Dumont[g], Benjamin; Ferrise[h], Roberto; Gayler[i], Sebastian; Ghahramani[j,1], Afshin; Harrison[k],

Matthew Tom; Hochman[e], Zvi; Hoogenboom[l], Gerrit; Huang[m], Mingxia; Jing[n], Qi; Justes[o], Eric;

Kersebaum[p,q,r], Kurt Christian; Launay[s], Marie; Lewan[t], Elisabet; Liu[k], Ke; Luo[u], Qunying; Mequanint[i], Fasil;

Nendel[p,q,v], Claas; Padovan[h], Gloria; Olesen[x], Jørgen Eivind; Pullens[x], Johannes Wilhelmus Maria; Qian[n],

Budong; Seserman[p], Diana-Maria; Shelia[l], Vakhtang; Souissi[y], Amir; Specka[p], Xenia; Wang[m], Jing; Weber[z],

Tobias, K.D.; Weihermüller[aa]; Lutz, Seidel, Sabine J.[a,ab]

[a]Institute of Crop Science and Resource Conservation, University of Bonn, Germany

[b]Natural Resources Institute Finland (Luke), Helsinki, Finland

[c]Julius Kühn Institute (JKI) – Federal Research Centre for Cultivated Plants, Institute for Crop and Soil

Science , Braunschweig, Germany

[d]INRAE, UMR 1114 EMMAH, Avignon, France

[e]CSIRO Agriculture and Food, Brisbane, Queensland, Australia

[f]Technical University of Munich, Department of Life Science Engineering, Digital Agriculture, HEF World

Agricultural Systems Center, Freising, Germany

[g]Plant Sciences & TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liege,

Gembloux, Belgium

[h]Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, Italy

---

[1] Present address: Department of Environment and Science and Innovation, Queensland Government, Toowoomba, QLD 4350, Australia.

[i]Institute of Soil Science and Land Evaluation, Biogeophysics, University of Hohenheim, Stuttgart,

Germany

[j]Centre for Sustainable Agricultural Systems, Institute for Life Sciences and the Environment, University

of Southern Queensland, Toowoomba, Queensland, Australia

[k]Tasmanian Institute of Agriculture, University of Tasmania, Launceston, Tasmania, Australia, 7248

[l]Agricultural and Biological Engineering Department, University of Florida, Gainesville, Florida, USA &

Global Food Systems Institute, University of Florida, Gainesville, Florida, USA

[m]College of Resources and Environmental Sciences, China Agricultural University, Beijing, China

[n]Ottawa Research and Development Centre, Agriculture and Agri-Food Canada, Ottawa, Canada

[o]CIRAD, Univ Montpellier, Persyst Department, Montpellier, France

[p]Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

[q]Global Change Research Institute CAS, Brno, Czech Republic

[r]Institute for Tropical Plant Production and Agricultural Systems Modelling, Georg-August University,

Göttingen, Germany

[s]INRAE, US 1116 AgroClim, Avignon, France

[t]Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Uppsala,

Sweden

[u]Hillridge Technology Pty Ltd, Sydney, Australia

[v]Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[x]Department of Agroecology, Aarhus University, Tjele, Denmark

[y]Swift Current Research and Development Centre, Agriculture and Agri-Food Canada, Swift Current,

Saskatchewan, Canada

[z]Soil Science Section, Faculty of Organic Agricultural Sciences, University of Kassel, Germany (before [i])

[aa]Institute of Bio- and Geosciences - IBG-3, Agrosphere, Forschungszentrum Jülich GmbH, Jülich,

Germany

[ab]Institute of Organic Farming, Department of Agricultural Sciences, University of Natural Resources and

Life Sciences, Vienna 1180, Austria

[*]Corresponding author. Email: taru.palosuo@luke.fi

Author contribution:

51   Daniel Wallach: Conceptualization, methodology, project administration, writing - original draft,

52   validation. Taru Palosuo, Henrike Mielenz, Peter Thorburn, Samuel Buis, Sabine Seidel:

53   Conceptualization, methodology, project administration, writing, review &editing. All other

54   authors: expertise on models and simulation results, writing, review & editing.

Abstract

It has become common to compare crop model results in multi-model simulation experiments. In general, one observes a large variability in such studies, which reduces the confidence one can have in such models. It is important to understand the causes of this variability as a first step toward reducing it. For a given data set, the variability in a multi-model study can arise from uncertainty in structure or in parameter values for a given structure. Previous studies have made assumptions about the origin of parameter uncertainty, and then quantified its contribution, generally finding that parameter uncertainty is less important than structure uncertainty. However, those studies do not take account of the full parameter variability in multi-model studies. Here we propose estimating parameter uncertainty based on open-call multi-model ensembles where the same structure is used by more than one modeling group. The variability in such a case is due to the full variability of parameters among modeling groups. Then structure and parameter contributions can be estimated sing random effects analysis of variance. Based on three multi-model studies for simulating wheat phenology, it is found that the contribution of parameter uncertainty to total uncertainty is, on average, more than twice as large as the uncertainty from structure. A second estimate, based on a comparison of two different calibration approaches for multiple models leads to a very similar result. We conclude that improvement of crop models requires as much attention to parameters as to model structure.

Keywords

Crop model, structure uncertainty, parameter uncertainty, multi-model studies

1. Introduction

Process-based crop models are widely used tools in agronomy, to help analyze past results and to explore scenarios related to alternative management options, effect of climate change, impact of new varieties etc. (Boote et al., 2010). Since the inception of the Agricultural Modeling Inter-comparison and Improvement Project (AgMIP, Rosenzweig et al., 2013), it has become common to organize multi-model simulation studies, where multiple modeling groups are provided with the same input data (i.e. weather, crop management, soil characteristics and initial conditions) and the same observational data for model calibration, and simulate the same response variables. Such studies have shown that the variability in simulated values is very substantial (Asseng et al., 2013; Durand et al., 2018; Salo et al., 2016; Sándor et al., 2020; Wallach et al., 2021a, 2021b). In several studies where multiple crop models were combined with multiple global circulation models (GCMs), it was found that the uncertainty contribution from crop models is as large or larger than that from the GCMs (Asseng et al., 2013; Müller et al., 2021; Wang et al., 2024, 2020). The variability in multi-model studies is probably the most pertinent measure of overall crop model uncertainty, since it reflects the current variability in simulated results among modeling groups. The large uncertainty clearly limits the confidence that one can have in the results of crop models, since it implies that results can vary widely depending on the specific model and modeling group providing the results. A reduction in uncertainty from crop models is necessary in order to increase confidence in the results of these models (Maiorano et al., 2017; Rötter et al., 2011).

In multi-model studies, all modeling groups are given the same input and observation data. Thus the variability between groups has two possible origins, model structure and model parameters. Crop model structure encompasses both the processes taken into account in the models and the specific forms of the equations used to model those processes. The second source of variability is uncertainty in the parameter values. Different groups using the same model structure can have different values for the

parameters (Albanito et al., 2022; Confalonieri et al., 2016), again leading to differences in simulated

values. It is important to quantify the contributions of structure and parameter uncertainty to overall

uncertainty, as a first step in reducing the overall uncertainty. In general, uncertainty is summarized as a

variance value, so the question is the relative contributions of structure and parameter variance to the

overall variance of simulated values in multi-model studies.

To proceed, it is helpful to have a clear definition of the parameter and structure uncertainties of

interest. We propose that an appropriate definition of model structure uncertainty is the range of plausible

crop model structures, where by "plausible" we mean a structure that would be deemed acceptable by

researchers knowledgeable in the field of crop modeling. The structures represented in open-call multi-

model studies can be considered a random sample of plausible structures; random because all modeling

groups were invited to participate, plausible because the structures are used by researchers in the field.

Our definition of parameter uncertainty for a given model structure and data set is the range of plausible

parameter vectors for that structure and data. Here "plausible" means that research groups working

with that structure would consider those parameters, or the way in which the parameters are obtained,

reasonable. Note that parameter uncertainty is nested within structure uncertainty. That is, the

uncertainty in question is different for each model structure, since each structure has its own

parameterization. Overall parameter uncertainty is an average over the uncertainties for different

structures. Parameter uncertainty covers both the plausible range of parameter default values and the

plausible range of calibration procedures, including the choice of which parameters to fit to the data and

the procedures of doing the fitting. Each parameter vector represented in open-call multi-model studies

is plausible, since it is derived by a practicing research group. For a given model structure, each

parameter vector can be considered a random draw from among plausible vectors for that structure,

that is, all plausible parameter vectors have an equal chance of being represented in the multi-model

study.

A major source of parameter uncertainty is variability in calibration approach. Calibration of crop

models and more generally process based models in other fields is difficult because of the large number

of parameters and the multiple types of observed variables. As a result, there is no general consensus on

the best calibration approach for crop modes (Ahuja and Ma, 2011). One question is how to choose which

parameters to estimate and which to treat as fixed. Various studies have proposed sensitivity analysis

approaches for choosing the parameters to estimate (e.g. Zadeh et al., 2022 for water quality modeling;

Zhang et al., 2014 for rice phenology). Other studies have compared different algorithms for searching for

the best parameter values (e.g. Gao et al., 2020; Harrison et al., 2019). The choice of objective function is

also variable, and several choices are possible (e.g. Houska et al., 2015 for ecological models). Wallach et

al. (2021c) specifically studied the range of calibration approaches used by different modeling groups in

three multi-model studies. They found that some groups used a frequentist approach, with various

objective functions, and others a Bayesian approach, with various assumptions about the likelihood

function. There were also differences in the way to choose the parameters to estimate, with some groups

basing the choice on sensitivity analysis, others on expert knowledge about the most important model

parameters, sometimes combined with trial and error to choose the parameter set that gives the best fit

to the data. Even for modeling groups using the same model structure, there were multiple differences in

calibration approach, including different choices of which parameters to fit to the data. . Albanito et al.

(2022) also emphasized that different modeling groups make different decisions about calibration

practices. In order to quantify the variability in plausible parameter values, we need to take into account

all these differences in calibration approach.

Several previous studies have specifically quantified the contributions of structure and parameter

uncertainty to total uncertainty for crop models or other process-based models. In these studies, a single

modeling group generates multiple parameter vectors for each of several models. Zhang et al. (2017)

considered five different models used to predict rice phenology. For each model, 50 parameter sets were

generated, based on an assumed uncertainty range for each parameter. Analysis of variance was used to

estimate the separate contributions of structure and parameter uncertainty. Tao et al. (2018) used a

similar approach for an ensemble of seven models for simulating barley growth and yield. Parameter

ranges were based on expert opinion. Both studies identify a fixed subset of parameters that are

considered uncertain. These approaches are appropriate for estimating *a priori* uncertainty in a fixed

subset of parameter , when no data is available for calibration. However, in multi-model studies, and in

our definition of plausible parameter values, models are calibrated using provided data. Calibration

modifies the uncertainty in parameter values compared to their *a priori* range. Furthermore, these studies

ignore uncertainty in the choice of parameters to estimate. In practice, different modeling groups using

the same model structure choose to estimate different parameters, but this variability is not taken into

account in these studies. ed. Thus, this approach does not provide a sample of plausible parameter values

according to our definition, and is therefore not appropriate for estimating the contribution of parameter

uncertainty to total uncertainty in multi-model studies.

A different approach assumes that parameter uncertainty derives from error in observations. The

variance of observation error is estimated and then used to estimate the distribution of the parameter

estimators. Thus Wallach et al (2017) estimated parameters of two models using the standard regression

technique of generalized least squares, which also provides estimates of the variance-covariance matrix

of the parameter estimators (Seber and Wild, 1989). Then samples of parameter vectors were generated

for each model, and analysis of variance was used to estimate structure and parameter variance. Another

approach explores the parameter space around the optimal parameter values, and accepts parameter

values that satisfy a goodness of fit test, which is a way of relating parameter uncertainty to observation

uncertainty. Migliavacca et al. (2012), looking at 12 budburst models, used this approach to generate 1000

parameter vectors that give fits to the data that are statistically equivalent to the fit using the best

parameter values for each model. However, while the effect of observation error on parameter

uncertainty is important, it is not relevant to uncertainty in multi-model studies, which use a fixed data

set. The variability in parameter values between different modeling groups in multi-model studies is not

due to observation error, since all groups use exactly the same data. The variability between groups using

the same model structure is due to differences in the choice of default parameter values and in calibration

approach, which is not captured in these studies.

In a somewhat different context, Smallman et al. (2021) studied structure and parameter

uncertainty for five terrestrial ecosystem models. Bayesian calibration was used to determine parameter

distributions for each model. This approach evaluates parameter uncertainty as the result of *a priori*

uncertainty and observation error. Since as noted above observation error is not considered in open-call

multi-model ensembles, this approach again does not recreate the variability of plausible parameter

values, according to our definition.

Xiong et al. (2020) considered structure and parameter contributions to uncertainty in wheat

model simulations applied to two Chinese regions. Three models were considered. Parameter variability

was due to different levels of spatial detail (same parameters at all locations, different parameters for each

major agro-ecological zone, parameters fit to phenology data or parameters fit to both phenology and

yield data). In this case parameter uncertainty measures the variability between different specific choices

of the way to determine the parameter values. The approach is not aimed at reproducing the variability in

multi-model studies.

These previous approaches have sampled from specific types of parameter uncertainty, namely a

priori uncertainty, uncertainty due to uncertainty in the data, or uncertainty related to different choices

of the way to treat spatial data. None of these approaches is designed to recreate the parameter variability

between different groups using the same structure in multi-model studies. That variability results from

differences in default parameter values and from multiple differences in calibration approach. To date,

there do not seem to have been any studies which realistically take into account the full uncertainty in

plausible parameter values.

The purpose of this study is to propose, for the first time, estimates of structure and parameter

uncertainty in multi-model studies that fully account for parameter uncertainty. Our case study concerns

prediction of wheat phenology. Crop phenology is important since it has a major effect on crop yield and

is a major aspect of crop response to global warming (Ahmad et al., 2019; Fatima et al., 2020; Liu et al.,

2022). Process-based models of phenology, usually embedded in crop or agroecosystem models, are often

used to predict the effect of weather on phenology (Bindi et al., 2015; Muleke et al., 2022). Multi-model

studies specifically of crop phenology have shown wide variability in simulated results (Wallach et al.,

2021b, 2021a). Thus, the uncertainty in prediction of crop phenology is of interest in its own right, as well

as serving as an example for crop models more generally.

We estimate parameter uncertainty based on the differences between different modeling groups

using the same model structure, in multi-model studies. This automatically takes into account all the

differences in choice of default parameters and in calibration approach between different

groups.Specifically, we revisit three published open-call multi-model studies on predicting wheat

phenology (Wallach et al., 2021b, 2021a), where some model structures were used by more than one

modeling group. Given the results from multiple different model structures, and from multiple groups

using the same structure, we estimate structure and parameter variances using random effects analysis of

variance. Our approach, therefore, provides more realistic estimates for the relative contributions of

structure and parameter uncertainty than have previously been available The methodology employed

here, namely using random effects analysis of variance on data from multi-model studies, is applicable in

general to process-based models.

217   In the particular cases treated here, we can obtain a second estimate of the variance in simulated

218 values due to parameter uncertainty. In the multi-model studies revisited here, each participating group

219 used their "usual" calibration approach. Subsequently an original calibration protocol was developed, and

220 most of the groups redid the calibration using the new protocol. Thus for multiple groups we have results

221 for two different calibration procedures, "usual" and "protocol", both of which may be considered

222 plausible The comparison between them can be used to estimate potential variability in parameters due

223 to calibration approach. These results can be compared with the results based on the original multi-model

224 studies.

2. Materials and Methods

2.1 Observational data

227   The multi-model studies use three different data sets (**Table 1**~~Table 1~~). Two of the data sets are

228 from winter wheat variety trials in France. These two data sets have identical structures (same sites and

229 sowing dates) but concern two different wheat varieties, "Apache" and "Bermude". Both varieties are of

230 intermediate precocity, but Bermude has a longer cycle. The observed data from each environment consist

231 of dates of stem elongation (BBCH30 on the BBCH scale, Meier, 1997) and of the middle of heading

232 (BBCH55). The third data set is from a multi-site sowing date experiment on spring wheat variety Janz in

233 Australia. The original data were observations of growth stage at weekly intervals. The data provided to

234 modeling groups were the dates of each BBCH stage, from the first to the last observed stage, derived

235 from interpolation of the original data. All data sets were split into two subsets, one for calibration and

236 the other for evaluation. The calibration and evaluation data sets had neither site nor year in common.

237 Thus evaluation was a true test of how well models could predict out-of-sample conditions.

Data set	Number of calibration environments	Number of evaluation environments	Observed development stages*	Simulated development stages used for analysis*
France Variety Apache	14	8	BBCH30, BBCH55	BBCH10, BBCH30, BBCH55
France Variety Bermude	14	8	BBCH30, BBCH55	BBCH10, BBCH30, BBCH55
Australia Variety Janz	24	18	BBCH stage at weekly intervals	BBCH10, BBCH30, BBCH65, BBCH90

*The development stages, using the BBCH scale (Meier, 1997) are emergence (BBCH10), start of

stem elongation (BBCH30), middle of heading (BBCH55), full flowering (BBCH65), and maturity (BBCH90).

Table 1. Data sets

The three data sets used in the multi-model ensemble studies analyzed here. The

French data sets are described in (Wallach et al., 2021a)**. The Australian data set is described**

in (Wallach et al., 2021b)**.**

2.2 Participating modeling groups and model structures

The call for participants in these studies was published using the mailing lists of AgMIP and of

several crop models. All modeling groups that volunteered (a modeling group is one or more individuals

who work together to run a model) were accepted. No specific attempt was made to encourage or

discourage participation by different groups that used the same structure. In the event, there were two or

three (depending on the simulation study) model structures used by more than one group (Table 2). These

are thus what we have termed "open-call" multi-model studies.

A total of 30 modeling groups using 23 different structures participated in one or more of the

simulation experiments. For the purposes of this study, each model with a unique name was considered

as a different model structure. A list of structures represented here, together with references that describe

the model equations, is given in Supplementary Table 1. In all cases, the phenology models were

embedded in general crop models, so the model structure here refers to the overall crop model. Most but

not all of the groups participated in all the studies.

In the original reports of these studies, the different modeling groups were identified by a code

(M1, M2 etc.) without identifying the group or the model structure used (Wallach et al., 2021b, 2021a).

The same codes are used here. In addition, we assign here codes to the different model structures (S1, S2,

etc.). The same model structure can be associated with more than one modeling group.

2.3 Simulation studies

The multi-model simulation studies, based on the three data sets, are summarized in Table 2. In

the first two studies, based on the two French data sets, each participating modeling group implemented

their usual calibration method to calibrate their model using the calibration data subsets and then

simulated days from sowing to development stages BBCH10, BBCH30, and BBCH55 for both the calibration

and evaluation environments (Wallach et al., 2021a) (Table 2). In the study based on the Australian data

set (Wallach et al., 2021b), each participating modeling group again used their usual calibration method

and then simulated days from sowing to development stages BBCH10, BBCH30, BBCH65, and BBCH90 for

the Australian environments. Neither the French nor the Australian data sets had observations of BBCH10

(emergence), which was included in the variables to simulate in order to have an example of a simulated

stage without calibration data. Detailed information about the variability in calibration procedures among

the different modeling groups is presented in Wallach et al. (2021c).

Simulation study	Data set (country, cultivar)	Calibration procedure	Number of modeling groups	Repeated structures*
multi-model study 1	France, Apache	Usual	27	S2 (4), S4 (3), S18 (2)
multi-model study 2	France, Bermude	Usual	27	S2 (4), S4 (3), S18 (2)
multi-model study 3	Australia, Janz	Usual	28	S2 (3), S4 (3), S18 (2)
multi-model study 4	All 3 data sets above	Protocol	19	not relevent

274   *The model structures used by more than one modeling group, and in parentheses, the number of groups

275   that used that structure.

276   **Table 2**

The three open-call multi-model simulation studies analyzed here (Wallach et al., 2021a,

2021b).

Following the three multi-model studies, a standardized calibration procedure (hereafter the

calibration "protocol") was proposed (Wallach et al., 2023). In a fourth simulation experiment, each of 19

participating groups used the protocol calibration to simulate for the same three data sets used in the first

studies. Any group that participated in the fourth study but that had not participated in the first three

studies first did the calibration using their usual method, before applying the protocol. Thus for each data

set and each participant, there were two parameter vectors, based on usual or protocol calibration.

2.4 Estimation of structure and parameter variance

Variance components approach

We assume that for each of the open-call simulation experiments, the model structures are a

random sample from plausible structures. We further assume that each parameter vector is a random

draw from plausible parameter vectors for the corresponding model structure. The statistical model for a

simulated value is

291
$$y_{sp} = \mu + \alpha_s + \beta_{sp} \tag{1}$$

where y_{sp} is the simulation result for the output y (for example days to BBCH30 in a particular

environment) obtained using the s^{th} draw of model structure s and the p^{th} draw of parameter vector for

that model structure. μ is the overall mean of the simulation results, α_s is the effect of model structure

and β_{sp} is the effect of parameter vector nested within the model structure. The parameter effect is nested

within structure, because different structures have different parameterizations. There is no residual term

because these are simulated values with no measurement error.

In a fixed effects analysis of variance, we are mainly interested in the effects α and β of the

specific structures and parameters of the sample. Here, on the other hand, we are mainly interested in

the variability between model structures and between parameter vectors for each given structure. The

appropriate analysis tool is a random effects analysis of variance.

In random effects analysis of variance, α_s and β_{sp} are treated as random effects. α_s is random

because the structure chosen at the s^{th} draw could be any plausible structure. All the α_s have the same

distribution, because every draw is from the same population of model structures. We are interested in

the variance of the α_s, noted $\text{var}(\alpha_s) = \sigma^2_{structure}$ for all s. Similarly, each β_{sp} has the same distribution. We

note $\mathrm{var}(\beta_{sp}) = \sigma^2_{parameters}$ for all *sp*. The random terms are assumed independent so that the total

variance for one environment and one simulated variable is the sum of model structure variance and

parameter variance (Scheffé, 1959):

309
$$\sigma^2_{total} = \sigma^2_{structure} + \sigma^2_{parameters} \quad (1)$$

The fraction of total variance due to parameter variance is then

$\sigma^2_{parameters} / \sigma^2_{total} = \sigma^2_{parameters} / (\sigma^2_{structure} + \sigma^2_{parameters})$.

The variances $\sigma^2_{structure}$ and $\sigma^2_{parameters}$ can be estimated using random effects analysis of variance.

In the case of a balanced sample (same number of parameter vectors for each model structure), there are

analytical expressions for the maximum likelihood (ML) estimators of the variances. In the general,

unbalanced case, the estimators are calculated iteratively. We use the lmer function of the lme4 package

(Bates et al., 2015) in R (R Core Team, 2017), which is designed to analyze linear mixed effect models. We

use the REML option, which corrects for the fact that the ML estimators are biased. Commented R script

for calling the lmer function and for extracting the variances of interest are shown in Supplementary.

An illustration of the data used in this approach is given in Fig. 1. As that figure shows, three of the

model structures (S2, S4, and S18) were used by more than one modeling group. It is the variability

between the different groups using the same model structure that allows us to estimate parameter

variance. The results for all groups contribute to the estimation of structure variance. Note that the

simplified approach to estimating parameter uncertainty using these same data in Wallach et al. (2021a,

2021b) underestimates the contribution of parameter uncertainty.

Figure 1

Example of data used for estimating parameter and structure uncertainty in the "variance components" approach (location Foreste, sown 10/11/2011). Each circle or star is a simulated value of days from sowing to mid-heading (stage BBCH55) for variety Apache in

one environment from simulation study 1. All modeling groups used the same inputs. The x-axis shows the model structure. In most cases, each structure was used by a single modeling group (open circles), but structures S2, S4, and S18 were used respectively by 4, 3, and 2 groups (data shown as stars). It is the variability between simulations from different groups using those model structures that is the basis for estimating parameter uncertainty. The horizontal line is the average of all simulated values.

Comparison of usual and protocol calibration

A second estimation of parameter uncertainty is based on comparing the results between usual and protocol calibration, for each group. The major difference between the calibration approaches was in the choice of which parameters to estimate from the data. For "usual" calibration, there was a wide diversity of methods of choosing parameters to estimate (Wallach et al., 2021c). None of the calibration methods was explicitly designed to avoid over-fitting. The new protocol based the choice of parameters to estimate on standard statistical model selection methods. For each development stage in the calibration data, one parameter that had a similar effect in all environments was automatically chosen to be calibrated. This was usually the number of degree days to the stage. Then additional "candidate" parameters were identified. These were tested in a procedure like forward regression. A candidate was chosen to be estimated if it led to a reduction in the corrected Akaike Information Criterion (AICc). This criterion has the form of a penalized likelihood; it has a term which decreases if the error in fitting the data decreases, and a term which increases as the number of estimated parameters increases. It is designed to avoid-over-fitting.

For each environment, we calculate the variance of the difference between the two calibration approaches (usual and protocol), noted var_i for modeling group i (eq. 1), and then average over modeling groups to obtain an estimated parameter variance for that environment (eq. 2).

354
$$\mathrm{var}_i = \left(y^{usual} - \bar{y}\right)^2 + \left(y^{protocol} - \bar{y}\right)^2 \ (2)$$

355
$$\hat{\sigma}^2_{parameters} = 1/R\sum_{i=1}^{R}\mathrm{var}_i \ (3)$$

356 where $\bar{y} = (y^{usual} + y^{protocol})/2$ and R is the number of modeling groups.

357   We do not have a rigorous method of estimating structure variance from these data. The

358 difficulty is that the assumptions underlying the random effects model described above are not satisfied;

359 the parameter vectors obtained by protocol calibration are not independent between modeling groups,

360 since all use the same protocol. Therefore, or each environment we use the estimate of structure

361 variance from the variance components approach above.

 Figure 2

364 **Example of simulated values where each group used two calibration approaches.**

365 **Simulations are of days from sowing to stem elongation (stage BBCH30) for variety**

366 **Bermude , location Boigneville, sown 21/10.2013. Each code (M1, etc.) refers to a specific**

367 **modeling group (x-axis). For each modeling group, there are two simulated values, one**

368 **using parameters based on usual calibration and the other using parameters based on**

369 **protocol calibration. The differences between the two values are used to estimate parameter**

370 **variance. The horizontal line shows the mean of simulations using usual calibration.**

371

372 3. Results

373 Initially, all calculations were done for the calibration and evaluation data subsets separately.

374 However, there were no systematic differences between the two (Fig. S1 and Fig. S2). Since both

375 calibration and evaluation data are sampled from the same target population, this is perhaps not

376 surprising. All results here are therefore based on the combined calibration and evaluation data. Table

377 3~~Table 3~~ and Table 4~~Table 4~~ show parameter variance, and parameter variance as a fraction of total

378 variance, averaged over all environments for the French and Australian data sets, respectively. Note that

379 the estimated contribution of parameter variance to total variance is always much less for BBCH10, for

380 which there were no observed values, than for the other simulated development stages.

381 Table 5~~Table 5~~ shows results averaged over all environments of both the French and Australian

382 data sets, for both the variance-components approach and the approach based on comparing usual and

383 protocol calibration. Separate values are given for BBCH10 and for all other stages. For development stages

384 other than BBCH10, the estimated contribution of parameter uncertainty is large. It ranges from 24 to

385 93%, and is larger than 50% for 11 out of the 14 combinations of data set, development stage and approach

386 for estimating the contribution of parameter variance.

387

	Apache			Bermude		
Development stage	BBCH10	BBCH30	BBCH55	BBCH10	BBCH30	BBCH55

Ave number of modeling groups	23	25	27	23	25	27
Number of model structures	17	19	21	17	19	21
Parameter variance (days²)	11.74	52.94	26.77	12.05	71.25	36.41
Structure variance (days²)	5.05	4.29	9.65	4.99	10.86	12.73
Parameter variance/total	0.55	0.93	0.81	0.56	0.92	0.8

Table 3

Parameter and structure variance for French data sets estimated using the variance

components calculation. The variance components estimate is based on ensemble studies

where some model structures were used by multiple modeling groups. Results are averages

over environments. The ratio of parameter variance to total (last row) is an average over the

ratios for each environment.

Development stage	BBCH10	BBCH30	BBCH65	BBCH90
Average number of modeling groups	23	24	28	26
Average number of model structures	17.95	19.93	22.64	21.05
Parameter variance (days²)	40.69	102.88	47.1	30.3
Structure variance (days²)	276.33	45.74	48.4	90.71
Parameter variance / total	0.12	0.66	0.49	0.24

396 **Table 4**

397   **Parameter and structure variance for the Australian data set estimated using the**

398   **variance components calculation. The variance components estimate is based on ensemble**

399   **studies where some model structures were used by multiple modeling groups. Results are**

400   **averages over environments. The ratio of parameter variance to total (last row) is an average**

401   **over the ratios for each environment.**

Method	Fraction of total variance due to parameter variance	
	BBCH10	All other stages
Variance components	0.41	0.69
Usual compared to protocol calibration	0.23	0.69

402

403

404

405   **Table 5**

406   **Fraction of total variance due to parameter variance based on two estimation procedures:**

407   **variance components and comparison of usual and protocol calibration. Averages are over**

408   **all environments of both the French and Australian data sets, either for the development**

409   **stage BBCH10 or for all other development stages.**

410

4. Discussion

. The variability in parameters between different groups using the same model structure results from differences in fixed parameter values and in calibration approach. None of the data sets here included time to emergence (BBCH10), so variability in simulating time to emergence likely depends only on variability in default values. On the other hand, there are data for subsequent development stages, so simulation of those stages will depend on the values of calibrated parameters. The parameter uncertainty contribution to variability in days to emergence is always smaller (41% on average) than the contribution to variability in days to other stages (69% on average). This shows that the importance of parameter uncertainty depends on both the specific variable simulated and on the data available for calibration. It seems that variability in simulated variables affected by calibration is larger than variability in simulated variables which depend just on default parameter values.

When averaging over data sets and simulated variables, for variables other than time to emergence, the resuts show that parameter uncertainty represents 69% of total uncertainty. The same value is obtained using variance components or the comparison of usual and protocol calibration. Structure uncertainty contributes 31%, so the contribution of parameter uncertainty is more than twice as large as the contribution of structure uncertainty. Previous studies that have estimated structure and parameter uncertainty contributions to overall uncertainty of plant or ecosystem models have generally found that structure uncertainty is larger than parameter uncertainty (Tao et al., 2018; Xiong et al., 2020), including studies that have specifically considered simulation of phenology (Migliavacca et al., 2012; Wallach et al., 2017; Zhang et al., 2017). This is in fact not too surprising. Previous studies have quantified the effect of some specific source of parameter uncertainty, such as observation error, or *a priori* uncertainty, or choice of data to use. The study here on the other hand takes into account the full

variability in default values and calibration approach between different modeling groups who could

participate in open-call multi-model studies.

Several studies have emphasized the uncertainty in parameter values and the necessity of better

evaluating parameter uncertainty and improving calibration practices (Iizumi et al., 2009; Ramirez-Villegas

et al., 2017; Seidel et al., 2018; Yang et al., 2024). The conclusion from this study, which remains to be

confirmed more generally, is that parameter uncertainty, in particular due to uncertainty in calibration

approach, is as or more important than structure uncertainty as a cause of variability in multi-model

studies. This implies that improving calibration approach, and sharing improved practices widely, should

be a major priority of the crop modeling community.

In multi-model study number 4 (Wallach et al., 2023) It was found that total variability between

simulated values was reduced by 22% when all groups used the new protocol, compared to the case where

each group used its usual calibration approach. Furthermore, prediction error was reduced by 11% by use

of the new protocol. This is a clear indication that improved, shared calibration approaches can

substantially reduce variability in multi-model studies and thereby enhance confidence in crop model

simulations. Recently, building on the calibration protocol for phenology data, a crop model calibration

approach has been proposed that is generic (i.e. can be applied to essentially all crop models and data

sets). It is directly based on statistical principles, which increases confidence that the approach uses

calibration data effectively (Wallach et al., 2025, 2024). This calibration protocol is being tested in several

studies, and would be a candidate for a standard calibration approach for crop models.

The present study has several limitations, which must be taken into account before generalizing

the results. First, this is a case study, so the specific results refer to the specific cases studied here. Also,

there are quite large differences for different environments and development stages (Supplementary Figs.

S1, S2. However, the studies use three different data sets, with 22 or 42 different environments each. The

French and Australian data sets not only have quite different environmental conditions and plant material

(winter wheat in France and spring wheat in Australia), but also quite different measured variables (two

development stages for the French data set, weekly development stage for the Australian data set). There

were also substantial differences within the French and Australian data sets. For variety Apache,

considering just the calibration data, time from sowing to stages BBCH30 and BBCH55 ranged from 126 to

192 days and from 177 to 234 days respectively. For the Australian data, time from sowing to stages

BBCH30, BBCH65 and BBCH90 ranged from 49 to 114 days, from 90 to 177 days and from 130 to 205 days

respectively. There is substantial variability in results for different environments and different simulated

variables, but the range of situations analyzed gives some confidence in the overall conclusion, that

parameter uncertainty is a major contribution to overall model uncertainty when simulating phenology. A

second limitation is that the study concerns only simulation of phenology. It remains to be seen whether

parameter uncertainty is also dominant when simulating crop growth and yield. Calibration in that case is

more complex, which may lead to larger calibration uncertainties, but also the simulations involve more

equations, which may lead to more structure uncertainty.. Thus it is not clear if the same conclusions as

here, that parameter uncertainty contributes more to overall uncertainty than structure uncertainty, also

applies in general to crop models In any case, however, it seems probable that taking into account the full

variability in plausible parameters will lead to a larger contribution of parameter uncertainty to total

uncertainty than previously thought, for all types of process-based models. A third limitation is that the

results rely on the variability between different groups that use the same model structure, but only three

model structures were used by more than one group, and at most four groups used the same model

structure. The results here thus represent only the parameter variability between a small number of

groups, using only three models. This is somewhat mitigated by the fact that that we have averaged over

a fairly large number of environments, in three different data sets. Most importantly, we also have a

second estimate of the contribution of parameter uncertainty, based on the comparison of usual and

protocol calibration. This second estimate is based on a sample of two different calibration approaches for

all models. The average contribution of parameter to total uncertainty using this second approach, again

ignoring simulations of time to emergence, was 69%, essentially the same as the value obtained using the

variance components analysis. This increases confidence in the variance component results.

. It should also be recognized that the assumption of a random sample of structures and

parameters underlying the random effects analysis of variance is probably not exactly satisfied. The

modeling groups in this study were not proactively chosen to represent a random sample, but rather

were simply the modeling groups that volunteered to participate. Such ensembles have been called

"ensembles of opportunity" (Tebaldi and Knutti, 2007). This could for example lead to underestimation

of structure uncertainty, if the full range of plausible structures was not sampled. It is not clear how to

quantify or correct this.   Perhaps the best one can do is to ensure that the invitation to participate is

widely disseminated.

The question of the relative importance of different sources of uncertainty in process-based

models, and in particular of structure and parameter uncertainty, arises in many fields, such as hydrology

(Butts et al., 2004) or earth system models (Ricciuto et al., 2021). Many of these fields also employ multi-

model studies (Thébault et al., 2024). The analysis of variance done here can be easily applied to these

other fields. , in order to estimate the contributions of structure and parameter uncertainty to the overall

variability. The requirement is to have the results of an open-call multi-model study where some model

structures are used by more than a single modeling group. The analysis proposed here does not require

any additional simulations, and the calculations can be easily done using existing software, for example

the R software package used here.

5. Conclusions

The large variability in crop multi-model simulations reduces the confidence one can have in such simulations, so it is important to understand the causes of this variability. A realistic estimate of parameter uncertainty must take into account the actual variability in parameters between different modeling groups. We illustrate a simple method of doing so, based on results of multi-model studies associated with random effects analysis of variance. This methodology is easily applicable to any case where the necessary multi-model results exist.

The parameter uncertainty taken into account here is different than in previous studies, where specific sources of parameter uncertainty were examined. Unlike those previous studies, we find in the case study here that parameter uncertainty makes the largest contribution to overall uncertainty, substantially larger than structure uncertainty. The larger parameter uncertainty found in the present study is unsurprising, since multiple sources of parameter uncertainty are included here, which suggests that it may be generally true that parameter uncertainty makes a larger contribution to total uncertainty than previously thought.

The importance of parameter uncertainty emphasizes the need for giving more attention to parameter values and in particular to Improved calibration approaches and procedures. Improving parameter values should be a major goal of the crop modeling community, on a par with work on improving model equations.

6. Acknowledgements

This work was in part supported by the Collaborative Research Center 1253 CAMPOS (Project 7: Stochastic Modelling Framework), funded by the German Research Foundation (DFG, Grant Agreement SFB 1253/1 2017), the Academy of Finland through projects AICropPro (316172) and DivCSA (316215), the

BonaRes projects ''Soil3'' (BOMA 03037514) and "I4S" (031B053I) of the Federal Ministry of Education and

Research (BMBF), Germany, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)

under Germany's Excellence Strategy - EXC 2070 – 390732324 EXC (PhenoRob), the project BiomassWeb

of the GlobeE programme (Grant number: FKZ031A258B) funded by the Federal Ministry of Education and

Research (BMBF, Germany), the INRA ACCAF meta-programme, the German Federal Ministry of Education

and Research (BMBF) in the framework of the funding measure "Soil as a Sustainable Resource for the

Bioeconomy –- BonaRes", project "BonaRes (Module B): BonaRes Centre for Soil Research, subproject B"

(grant 031B0511B), the National Key Research and Development Program of China (2017YFD0300205),

the National Science Foundation for Distinguished Young Scholars (31725020), the Priority Academic

Program Development of Jiangsu Higher Education Institutions (PAPD), the 111 project (B16026), and

China Scholarship Council, the Agriculture and Agri-Food Canada's Project J-002303 under the

Interdepartmental Research Initiative in Agriculture, the DFG Research Unit FOR 1695 'Agricultural

Landscapes under Global Climate Change – Processes and Feedbacks on a Regional Scale, the U.S.

Department of Agriculture National Institute of Food and Agriculture (award no. 2015-68007-23133) and

USDA/NIFA HATCH grant N. MCL02368, the National Key Research and Development Program of China

(2016YFD0300105), The Broadacre Agriculture Initiative, a research partnership between University of

Southern Queensland and the Queensland Department of Agriculture and Fisheries, the Academy of

Finland through project AI-CropPro (315896), the JPI FACCE MACSUR2 project, funded by the Italian

Ministry for Agricultural, Food, and Forestry Policies (D.M. 24064/7303/15 of 6/Nov/2015), the SYSTEMIC

funded by JPI HDHL, JPI-OCEANS and FACCE-JPI under ERA-NET (n.696295), the SustEs project funded by

the Ministry of Education, Youth and Sports of the Czech Republic (CZ.02.1.01/0.0/0.0/16_019/000797).

The time of MTH and KL was supported by the Grains Research and Development Corporation (Project

UOT1906-002RTX). The observed Australian phenology data were jointly funded by CSIRO, the Grains

7. References

Ahmad, S., Abbas, G., Ahmed, M., Fatima, Z., Anjum, M.A., Rasul, G., Khan, M.A., Hoogenboom, G., 2019.

Climate warming and management impact on the change of phenology of the rice-wheat cropping

system in Punjab, Pakistan. F. Crop. Res. 230, 46–61. https://doi.org/10.1016/j.fcr.2018.10.008

Ahuja, L.R., Ma, L., 2011. A Synthesis of Current Parameterization Approaches and Needs for Further

Improvements, in: Methods of Introducing System Models into Agricultural Research, Advances in

Agricultural Systems Modeling. pp. 427–440.

https://doi.org/https://doi.org/10.2134/advagricsystmodel2.c15

Albanito, F., McBey, D., Harrison, M., Smith, P., Ehrhardt, F., Bhatia, A., Bellocchi, G., Brilli, L., Carozzi, M.,

Christie, K., Doltra, J., Dorich, C., Doro, L., Grace, P., Grant, B., Léonard, J., Liebig, M., Ludemann, C.,

Martin, R., Meier, E., Meyer, R., De Antoni Migliorati, M., Myrgiotis, V., Recous, S., Sándor, R.,

Snow, V., Soussana, J.-F., Smith, W.N., Fitton, N., 2022. How Modelers Model: the Overlooked

Social and Human Dimensions in Model Intercomparison Studies. Environ. Sci. Technol. 56, 13485–

13498. https://doi.org/10.1021/acs.est.2c02023

Asseng, S., Ewert, F., Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P.J.,

Rötter, R.P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P.K., Angulo, C., Bertuzzi,

P., Biernath, C., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J.,

Hunt, L.A., Ingwersen, J., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Naresh Kumar, S., Nendel, C.,

O'Leary, G., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A.,

Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M.,

Waha, K., Wallach, D., White, J.W., Williams, J.R., Wolf, J., 2013. Uncertainty in simulating wheat

yields under climate change. Nat. Clim. Chang. 3, 827–832. https://doi.org/10.1038/nclimate1916

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J.

Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bindi, M., Palosuo, T., Trnka, M., Semenov, M., 2015. Modelling climate change impacts on crop

production for food security. Clim. Res. 65, 3–5. https://doi.org/10.3354/cr01342

Boote, K.J., Jones, J.W., Hoogenboom, G., White, J.W., 2010. The Role of Crop Systems Simulation in

Agriculture and Environment. Int. J. Agric. Environ. Inf. Syst. 1, 41–54.

Butts, M.B., Payne, J.T., Kristensen, M., Madsen, H., 2004. An evaluation of the impact of model structure

on hydrological modelling uncertainty for streamflow simulation. J. Hydrol. 298, 242–266.

https://doi.org/10.1016/J.JHYDROL.2004.03.042

Confalonieri, R., Orlando, F., Paleari, L., Stella, T., Gilardelli, C., Movedi, E., Pagani, V., Cappelli, G.,

Vertemara, A., Alberti, L., Alberti, P., Atanassiu, S., Bonaiti, M., Cappelletti, G., Ceruti, M.,

Confalonieri, A., Corgatelli, G., Corti, P., Dell'Oro, M., Ghidoni, A., Lamarta, A., Maghini, A.,

Mambretti, M., Manchia, A., Massoni, G., Mutti, P., Pariani, S., Pasini, D., Pesenti, A., Pizzamiglio, G.,

Ravasio, A., Rea, A., Santorsola, D., Serafini, G., Slavazza, M., Acutis, M., 2016. Uncertainty in crop

model predictions: What is the role of users? Environ. Model. Softw. 81, 165–173.

https://doi.org/10.1016/j.envsoft.2016.04.009

Durand, J.-L., Delusca, K., Boote, K., Lizaso, J., Manderscheid, R., Weigel, H.J., Ruane, A.C., Rosenzweig, C.,

Jones, J., Ahuja, L., Anapalli, S., Basso, B., Baron, C., Bertuzzi, P., Biernath, C., Deryng, D., Ewert, F.,

Gaiser, T., Gayler, S., Heinlein, F., Kersebaum, K.C., Kim, S.-H., Müller, C., Nendel, C., Olioso, A.,

Priesack, E., Villegas, J.R., Ripoche, D., Rötter, R.P., Seidel, S.I., Srivastava, A., Tao, F., Timlin, D.,

Twine, T., Wang, E., Webber, H., Zhao, Z., 2018. How accurately do maize crop models simulate the

interactions of atmospheric CO_2 concentration levels with limited water supply on water use and

yield? Eur. J. Agron. 100, 67–75. https://doi.org/10.1016/J.EJA.2017.01.002

Fatima, Z., Ahmed, M., Hussain, M., Abbas, G., Ul-Allah, S., Ahmad, S., Ahmed, N., Ali, M.A., Sarwar, G.,

Haque, E. ul, Iqbal, P., Hussain, S., 2020. The fingerprints of climate warming on cereal crops

phenology and adaptation options. Sci. Rep. 10, 18013. https://doi.org/10.1038/s41598-020-

74740-3

Gao, Y., Wallach, D., Liu, B., Dingkuhn, M., Boote, K.J., Singh, U., Asseng, S., Kahveci, T., He, J., Zhang, R.,

Confalonieri, R., Hoogenboom, G., 2020. Comparison of three calibration methods for modeling rice

phenology. Agric. For. Meteorol. 280, 107785. https://doi.org/10.1016/J.AGRFORMET.2019.107785

Harrison, M.T., Roggero, P.P., Zavattaro, L., 2019. Simple, efficient and robust techniques for automatic

multi-objective function parameterisation: Case studies of local and global optimisation using

APSIM. Environ. Model. Softw. 117, 109–133. https://doi.org/10.1016/j.envsoft.2019.03.010

Houska, T., Kraft, P., Chamorro-Chavez, A., Breuer, L., 2015. SPOTting Model Parameters Using a Ready-

Made Python Package. PLoS One 10, e0145180. https://doi.org/10.1371/journal.pone.0145180

Iizumi, T., Yokozawa, M., Nishimori, M., 2009. Parameter estimation and uncertainty analysis of a large-

scale crop model for paddy rice: Application of a Bayesian approach. Agric. For. Meteorol. 149,

333–348.

Liu, Y., Bachofen, C., Wittwer, R., Silva Duarte, G., Sun, Q., Klaus, V.H., Buchmann, N., 2022. Using

PhenoCams to track crop phenology and explain the effects of different cropping systems on yield.

Agric. Syst. 195, 103306. https://doi.org/10.1016/J.AGSY.2021.103306

Maiorano, A., Martre, P., Asseng, S., Ewert, F., Müller, C., Rötter, R.P., Ruane, A.C., Semenov, M.A.,

Wallach, D., Wang, E., Alderman, P.D., Kassie, B.T., Biernath, C., Basso, B., Cammarano, D.,

Challinor, A.J., Doltra, J., Dumont, B., Rezaei, E.E., Gayler, S., Kersebaum, K.C., Kimball, B.A., Koehler,

A.-K., Liu, B., O'Leary, G.J., Olesen, J.E., Ottman, M.J., Priesack, E., Reynolds, M., Stratonovitch, P.,

Streck, T., Thorburn, P.J., Waha, K., Wall, G.W., White, J.W., Zhao, Z., Zhu, Y., 2017. Crop model

improvement reduces the uncertainty of the response to temperature of multi-model ensembles.

F. Crop. Res. 202. https://doi.org/10.1016/j.fcr.2016.05.001

Meier, U., 1997. Growth stages of mono- and dicotyledonous plants : BBCH-Monograph [WWW

Document]. URL

https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwjBo-

Ln4_r4AhVM3RoKHc-

TA5sQFnoECAcQAQ&url=https%3A%2F%2Fwww.politicheagricole.it%2Fflex%2FAppData%2FWebLi

ve%2FAgrometeo%2FMIEPFY800%2FBBCHengl2001.pdf&usg=AOvVaw1G_eUh_-qnVt91_qbj44d

(accessed 1.9.19).

Migliavacca, M., Sonnentag, O., Keenan, T.F., Cescatti, A., O'Keefe, J., Richardson, A.D., 2012. On the

uncertainty of phenological responses to climate change, and implications for a terrestrial

biosphere model. Biogeosciences 9, 2063–2083. https://doi.org/10.5194/bg-9-2063-2012

Muleke, A., Harrison, M.T., de Voil, P., Hunt, I., Liu, K., Yanotti, M., Eisner, R., 2022. Earlier crop flowering

caused by global warming alleviated by irrigation. Environ. Res. Lett. 17, 044032.

https://doi.org/10.1088/1748-9326/ac5a66

Müller, C., Franke, J., Jägermeyr, J., Ruane, A.C., Elliott, J., Moyer, E., Heinke, J., Falloon, P.D., Folberth, C.,

Francois, L., Hank, T., Izaurralde, R.C., Jacquemin, I., Liu, W., Olin, S., Pugh, T.A.M., Williams, K.,

Zabel, F., 2021. Exploring uncertainties in global crop yield projections in a large ensemble of crop

models and CMIP5 and CMIP6 climate scenarios. Environ. Res. Lett. 16, 034040.

https://doi.org/10.1088/1748-9326/abd8fc

R Core Team, 2017. A language and Environment for Statistical Computing.

Ramirez-Villegas, J., Koehler, A.-K., Challinor, A.J., 2017. Assessing uncertainty and complexity in regional-

scale crop model simulations. Eur. J. Agron. 88, 84–95. https://doi.org/10.1016/J.EJA.2015.11.021

Ricciuto, D.M., Yang, X., Wang, D., Thornton, P.E., 2021. The impacts of model structure, parameter

uncertainty and experimental design on Earth system model simulations of litter bag

decomposition experiments. https://doi.org/10.5194/bg-2021-163

Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P., Antle, J.M., Nelson, G.C.,

Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorria, G., Winter, J.M., 2013.

The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot

studies. Agric. For. Meteorol. 170. https://doi.org/10.1016/j.agrformet.2012.09.011

Rötter, R.P., Carter, T.R., Olesen, J.E., Porter, J.R., 2011. Crop–climate models need an overhaul. Nat.

Clim. Chang. 1, 175–177. https://doi.org/10.1038/nclimate1152

Salo, T.J., Palosuo, T., Kersebaum, K.C., Nendel, C., Angulo, C., Ewert, F., Bindi, M., Calanca, P., Klein, T.,

Moriondo, M., Ferrise, R., Olesen, J.E., Patil, R.H., Ruget, F., Takac, J., Hlavinka, P., Trnka, M., Rotter,

R.P., 2016. Comparing the performance of 11 crop simulation models in predicting yield response

to nitrogen fertilization. J. Agric. Sci. 154, 1218–1240. https://doi.org/10.1017/S0021859615001124

Sándor, R., Ehrhardt, F., Grace, P., Recous, S., Smith, P., Snow, V., Soussana, J.-F., Basso, B., Bhatia, A.,

Brilli, L., Doltra, J., Dorich, C.D., Doro, L., Fitton, N., Grant, B., Harrison, M.T., Kirschbaum, M.U.F.,

Klumpp, K., Laville, P., Léonard, J., Martin, R., Massad, R.-S., Moore, A., Myrgiotis, V., Pattey, E.,

Rolinski, S., Sharp, J., Skiba, U., Smith, W., Wu, L., Zhang, Q., Bellocchi, G., 2020. Ensemble

modelling of carbon fluxes in grasslands and croplands. F. Crop. Res. 252, 107791.

https://doi.org/10.1016/j.fcr.2020.107791

Scheffé, H., 1959. The analysis of variance. John Wiley & Sons, New York.

Seber, G.A.F., Wild, C.J., 1989. Nonlinear regression. Wiley , New York.

Seidel, S.J., Palosuo, T., Thorburn, P., Wallach, D., 2018. Towards improved calibration of crop models –

Where are we now and where should we go? Eur. J. Agron. 94, 25–35.

https://doi.org/10.1016/J.EJA.2018.01.006

Smallman, T.L., Milodowski, D.T., Neto, E.S., Koren, G., Ometto, J., Williams, M., 2021. Parameter

uncertainty dominates C-cycle forecast errors over most of Brazil for the 21st century. Earth Syst.

Dynam. 12, 1191–1237. https://doi.org/10.5194/esd-12-1191-2021

Tao, F., Rötter, R.P., Palosuo, T., Gregorio Hernández Díaz-Ambrona, C., Mínguez, M.I., Semenov, M.A.,

Kersebaum, K.C., Nendel, C., Specka, X., Hoffmann, H., Ewert, F., Dambreville, A., Martre, P.,

Rodríguez, L., Ruiz-Ramos, M., Gaiser, T., Höhn, J.G., Salo, T., Ferrise, R., Bindi, M., Cammarano, D.,

Schulman, A.H., 2018. Contribution of crop model structure, parameters and climate projections to

uncertainty in climate change impact assessments. Glob. Chang. Biol. 24, 1291–1307.

https://doi.org/10.1111/gcb.14019

Tebaldi, C., Knutti, R., 2007. The use of the multi-model ensemble in probabilistic climate projections.

Philos. Trans. A. Math. Phys. Eng. Sci. 365, 2053–75. https://doi.org/10.1098/rsta.2007.2076

Thébault, C., Perrin, C., Andréassian, V., Thirel, G., Legrand, S., Delaigue, O., 2024. Multi-model approach

in a variable spatial framework for streamflow simulation. Hydrol. Earth Syst. Sci. 28, 1539–1566.

https://doi.org/10.5194/hess-28-1539-2024

Wallach, D., Buis, S., Seserman, D.-M., Palosuo, T., Thorburn, P.J., Mielenz, H., Justes, E., Kersebaum, K.-

C., Dumont, B., Launay, M., Seidel, S.J., 2024. A calibration protocol for soil-crop models. Environ.

Model. Softw. 180, 106147. https://doi.org/10.1016/J.ENVSOFT.2024.106147

Wallach, D., Kim, K.S., Hyun, S., Buis, S., Thorburn, P., Mielenz, H., Seidel, S.J., Alderman, P.D., Dumont,

B., Fallah, M.H., Hoogenboom, G., Justes, E., Kersebaum, K.-C., Launay, M., Leolini, L., Mehmood,

M.Z., Moriondo, M., Jing, Q., Qian, B., Susanne, S., Seserman, D.-M., Shelia, V., Weihermüller, L.,

Palosuo, T., 2025. Evaluating the AgMIP calibration protocol for crop models; case study and new

diagnostic tests. Eur. J. Agron. 168, 127659. https://doi.org/10.1016/J.EJA.2025.127659

Wallach, D., Nissanka, S.P., Karunaratne, A.S., Weerakoon, W.M.W., Thorburn, P.J., Boote, K.J., Jones,

J.W., 2017. Accounting for both parameter and model structure uncertainty in crop model

predictions of phenology: A case study on rice. Eur. J. Agron. 88.

https://doi.org/10.1016/j.eja.2016.05.013

Wallach, D., Palosuo, T., Thorburn, P., Gourdain, E., Asseng, S., Basso, B., Buis, S., Crout, N., Dibari, C.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani, A., Hochman, Z., Hoek, S.,

Hoogenboom, G., Horan, H., Huang, M., Jabloun, M., Jing, Q., Justes, E., Kersebaum, K.C.,

Klosterhalfen, A., Launay, M., Luo, Q., Maestrini, B., Mielenz, H., Moriondo, M., Nariman Zadeh, H.,

Olesen, J.E., Poyda, A., Priesack, E., Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A.,

Specka, X., Srivastava, A.K., Stella, T., Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D.,

Weihermüller, L., de Wit, A., Wöhling, T., Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021a. How well do

crop modeling groups predict wheat phenology, given calibration data from the target population?

Eur. J. Agron. 124, 126195. https://doi.org/https://doi.org/10.1016/j.eja.2020.126195

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Andrianasolo, F., Asseng, S., Basso, B., Buis, S., Crout,

N., Dumont, B., Ferrise, R., Gaiser, T., Gayler, S., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G.,

Huang, M., Jabloun, M., Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.C., Launay, M., Lewan, E.,

Luo, Q., Maestrini, B., Moriondo, M., Olesen, J.E., Padovan, G., Poyda, A., Priesack, E., Pullens,

J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Kumar Srivastava, A., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021b. Multi-model evaluation of phenology prediction for

wheat in Australia. Agric. For. Meteorol. 298–299, 108289.

https://doi.org/10.1016/J.AGRFORMET.2020.108289

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Gourdain, E., Andrianasolo, F., Asseng, S., Basso, B.,

Buis, S., Crout, N., Dibari, C., Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani,

A., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G., Huang, M., Jabloun, M., Jansson, P.-E., Jing,

Q., Justes, E., Kersebaum, K.C., Klosterhalfen, A., Launay, M., Lewan, E., Luo, Q., Maestrini, B.,

Mielenz, H., Moriondo, M., Nariman Zadeh, H., Padovan, G., Olesen, J.E., Poyda, A., Priesack, E.,

Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Srivastava, A.K., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021c. The chaos in calibrating crop models: Lessons learned

from a multi-model calibration exercise. Environ. Model. Softw. 145, 105206.

https://doi.org/10.1016/J.ENVSOFT.2021.105206

Wallach, D., Palosuo, T., Thorburn, P., Mielenz, H., Buis, S., Hochman, Z., Gourdain, E., Andrianasolo, F.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Harrison, M., Hiremath, S., Horan, H.,

Hoogenboom, G., Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.-C., Launay, M., Lewan, E., Liu, K.,

Mequanint, F., Moriondo, M., Nendel, C., Padovan, G., Qian, B., Schütze, N., Seserman, D.-M.,

Shelia, V., Souissi, A., Specka, X., Srivastava, A.K., Trombi, G., Weber, T.K.D., Weihermüller, L.,

Wöhling, T., Seidel, S.J., 2023. Proposal and extensive test of a calibration protocol for crop

phenology models. Agron. Sustain. Dev. 43, 46. https://doi.org/10.1007/s13593-023-00900-0

Wang, B., Feng, P., Liu, D.L., O'Leary, G.J., Macadam, I., Waters, C., Asseng, S., Cowie, A., Jiang, T., Xiao,

D., Ruan, H., He, J., Yu, Q., 2020. Sources of uncertainty for wheat yield projections under future

climate are site-specific. Nat. Food 1, 720–728. https://doi.org/10.1038/s43016-020-00181-w

Wang, B., Jägermeyr, J., O'Leary, G.J., Wallach, D., Ruane, A.C., Feng, P., Li, L., Liu, D.L., Waters, C., Yu, Q.,

Asseng, S., Rosenzweig, C., 2024. Pathways to identify and reduce uncertainties in agricultural

climate impact assessments. Nat. Food 1–7. https://doi.org/10.1038/s43016-024-01014-w

Xiong, W., Asseng, S., Hoogenboom, G., Hernandez-Ochoa, I., Robertson, R., Sonder, K., Pequeno, D.,

Reynolds, M., Gerard, B., 2020. Different uncertainty distribution between high and low latitudes in

modelling warming impacts on wheat. Nat. Food 1, 63–69. https://doi.org/10.1038/s43016-019-

0004-2

Yang, C., Lei, N., Menz, C., Ceglar, A., Torres-Matallana, J.A., Li, Siqi, Jiang, Y., Tan, X., Tao, L., He, F., Li,

Shigui, Liu, B., Yang, F., Fraga, H., Santos, J.A., 2024. Regional uncertainty analysis between crop

phenology model structures and optimal parameters. Agric. For. Meteorol. 355, 110137.

https://doi.org/10.1016/J.AGRFORMET.2024.110137

Zadeh, F.K., Nossent, J., W., B., Bauwens, W., Van Griensven, A., 2022. A fast and effective

parameterization of water quality models. Environ. Model. Softw. 149, 105331.

https://doi.org/https://doi.org/10.1016/j.envsoft.2022.105331

Zhang, S., Tao, F., Shi, R., 2014. Modeling the rice phenology and production in China with SIMRIW:

sensitivity analysis and parameter estimation. Front. Earth Sci. 8, 505–511.

https://doi.org/10.1007/s11707-014-0468-1

Zhang, S., Tao, F., Zhang, Z., 2017. Uncertainty from model structure is larger than that from model

parameters in simulating rice phenology in China. Eur. J. Agron. 87, 30–39.

https://doi.org/10.1016/j.eja.2017.04.004